



UNION OF SOUTH AFRICA

DEPARTMENT OF AGRICULTURE

Dipping and Tick-destroying Agents

PART III

BY

LT.-COL. H. WATKINS-PITCHFORD, F.R.C.V.S., F.R.S.E.

Government Bacteriologist, Natal

PRETORIA

THE GOVERNMENT PRINTING AND STATIONERY OFFICE

1911

PRINTED BY THE GOVERNMENT PRINTER, PRETORIA

5311—25/7/11--5000.

INDEX TO SCHEDULES

OF

REPORTS I, II, and III on "Dipping and Tick-Destroying Agents".

REPORT I.

I.—Estimation of efficacy of Proprietary Dips.	I.—Cooper's Tixol.
Schedule	II.—McDougall's Dip.
"	III.—Quibell's Dip.
"	IV.—Cooper's Powder Dip.
"	V.—Demuth Dip.
"	VI.—Izo-Izal Sheep Dip.
"	VII.—Thomas's Dip.
"	VIII.—Holmes' Dip.
"	IX.—Ialine Sheep Dip.
"	X.—Hypochlorite, or Electrolysed Sea Water.
"	XI.—Arsenite of Soda.
"	XII.—Erkenbrach's Cattle Dip.
"	XIII.—Alderson's Cattle Dip.
"	XIV to XVIII.—"Laboratory" Spraying Fluid.
"	XIX to XXI.—"Laboratory" Dipping Fluid.
Appendix	A.—Methods of Preparation.
"	B.—Composition of "Queensland" Dip.
"	C.—Author's Spraying Device.


REPORT II.

Schedule	A.—The ability of an animal to kill ticks placed upon it increases with the number of dippings such animal receives.
"	B.—The length of time elapsing since the last dipping determines the ability of an habituated animal to kill ticks which may become attached to it.
"	C.—Observations to show that in habituated animals the effect on ticks of a given dipping is more rapid than in non-habituated animals.
"	D.—Observations to show the retardation of the infected process in the habituated animal.
"	E.—Estimation of the amount of arsenic retained by the hair and hide of animals for varying periods after dipping.
"	F.—To show the elimination of arsenic of the kidneys at varying periods after dipping.

REPORT III.

Schedule	I.—Shortest periods for engorgement of larva and nymph of the Brown tick.
"	II.—Details of adjustment of "Three-Day" Dip.
"	III.—Lethal effect of "Three-Day" and "Laboratory" Dips on distended female Brown ticks.
"	IV.—Effect of "Three-Day" Dip at intervals of 72 hours upon adult tick-life generally.
"	V.—Lethal effect of dips half and five-twelfth strengths at intervals of 48 hours, showing the increase of efficiency resulting from the addition of soap and paraffin.
"	VI.—The "revulsive effect" of dipping on immature tick-life.
"	VII.—Details for preparation of "Three-Day" dipping fluid.
"	VIII.—Details of preparation of "Three-Day" ear dressing.
"	IX.—General map of experimental paddocks.
"	X.—Diagram illustrating rate of tick-travel, etc.
"	XI.—Map illustrating rate at which tick-infested paddocks are capable of being cleared.
"	XII.—Relative tick activity during day and night.
"	XIII.—Residual effect of "Three-Day" Dip in horses and cattle.
"	XIV.—The agency of the tail-tuft in the collection of ticks.
"	XV.—The suitability of "Three-Day" Dip in the treatment of scab in sheep.
"	XVI.—Effect of weekly dipping process (with "Laboratory" Dip) upon secretion of milk.
"	XVII.—Illustration of "Isometer" apparatus for estimation of arsenic.

Dipping and Tick-Destroying Agents.



GROUP OF FREQUENTLY DIPPED CATTLE.

These experimental cattle (about 30 in number) have been regularly dipped for more than six months every 72 hours. The above photograph serves to show that, in spite of 68 such dippings, the condition of the herd has been maintained.