

Biological activities of extracts and isolated compounds from *Bauhinia galpinii* (Fabaceae) and *Combretum vendae* (Combretaceae) as potential antidiarrhoeal agents

Ahmed Aroke Shahid

BSc (Chemistry) (ABU), MSc (Chemistry) (UNILAG)

A dissertation submitted in fulfilment of the requirements for the degree of Doctor of

Philosophy (PhD)

in the

Phytomedicine Programme, Department of Paraclinical Sciences,

Faculty of Veterinary Science



UNIVERSITEIT VAN PRETORIA
UNIVERSITY OF PRETORIA
YUNIBESITHI YA PRETORIA

Promoter: Prof. Jacobus N. Eloff

Co-promoters: Dr. Nivan Moodley (CSIR)

Prof. Vinny Naidoo

Dr. Lyndy McGaw

January 2012

Declaration

The research work described in the thesis was conducted in the Phytomedicine Programme in the Department of Paraclinical Sciences, Faculty of Veterinary Science, University of Pretoria under the supervision of Professor JN. Eloff, Dr. N. Moodley, Prof. V. Naidoo and Dr. LJ. McGaw

The results presented herewith were generated from my own experiments, except where the work of others are quoted and referenced. There is no part of this work that has been submitted to any other University.

Aroke Shahid, Ahmed

Dedication

This work is dedicated to the memory of the following: My Father (Late Mr. Ahmed Aninya Aroke), my brothers (Late Salihu Aroke and Late Ibrahim Onimisi Ahmed), Late Olukemi Ore Udom (A friend and colleague who started her PhD, but could not finish the programme before death) and my dear sister (Late Mrs. Husseinatu Ohunene Abubakar).

Acknowledgements

This Ph.D. study would not have been possible without the support and encouragement of many people. I will thank my mum Mrs Omeneke Aminatu Ahmed who has been the pillar of my life. The love of my wife Mrs Rabiatu Isoyiza Ahmed and my children Enehu Mazidah, Onize Nusrah, Ometere Shamsiyah, Eneze Azeemah and Ava'ami AbdulAleem have been essential to my success. I thank you all for your understanding and endurance during my absence. I also wish to thank my siblings Mr. Aroke Haruna, Mr. Umar Omeiza Aroke and his wife Rukkayat, Mrs. Zulaiha Mubarak, Dr. Halidu Aroke Ahmed, Mrs. Khaltum Khamilu and Aisha Ahmed for their enormous support. I am also deeply indebted to my friends who stuck by me through the years: Dr. Muhammed Awwalu Usman, Engr. Yakubu Adajah, Mr. Yahaya Ohida Yusuf, Mr. Tijani Muhammed Isah, Abdulmumuni Enesi Umar, Dr. Muhammed Onujagbe Onoda, Mrs. Ukachi Ezenwa Igbo, Dr. Caroline Anyakorah, Dr. Oluwatoyin Taiwo and Dr. Chima Cartney Igwe among others. Financial support was a crucial element of my ability to devote so much time to this research. Major support that keeps me and my family afloat during the study was provided by a study leave with pay provide by Federal Institute of Industrial Research Oshodi (FIIRO), Lagos, Nigeria. The University of Pretoria has provided support through University bursary, National Research Foundation (NRF) of South Africa provide fund for the research, Faculty of Veterinary Science also provide research fund through the research committee (RESCOM), and Department of Paraclinical Science also provide research fund.

I am extremely grateful to my supervisor, Prof. J. N. Eloff (overall supervisor) and my co-supervisors, Dr. Nivan Moodley (characterization of isolated compounds), Prof. Vinny Naidoo (isolated organ studies), and Dr. Lyndy J. McGaw (cellular toxicology) for allowing me to tap from the wealth of their knowledge and also giving me much of their time and energy through the duration of this study. I have a great respect and appreciated the instruction and assistance I received from each and every one of you. I sincere thank the Secretary to Phytochemical Programme, Tharien de Winnaar for all her assistance in coordinating the purchase of materials and other important aspects of this project.

I feel honoured to have opportunity to do some of the work at Bioscience, CSIR South Africa and lucky to meet with Dr. Vinesh Maharaj, Dr. Jacqueline Ndlebe, and Ms. Teresa Faleschini who are friends as well as crucial resources of information, techniques and instructions. I express my gratitude to CSIR for allowing me to use her NMR spectroscopic facilities.

At the UPBRC where I got my isolated organ part, the assistance of Dr. Tamsyn Pulleer, Mrs. Stephanie Keulder and Mrs. Ilse Janse van Rensburg were highly appreciated. The ability to collect plant material was essential to this project, and for that I want to recognize the help and outstanding collaboration of Ms Magda Nel of the Manie van der Schyff Botanical Garden and Ms Elsa van Wyk (Curator of the HGW Schweickert Herbarium of the University of Pretoria) for assistance in collection, identification and authentication of the plant samples. My appreciation also goes to Ms Lita Pauw of the Phytomedicine Programme for allowing me access her stored

plant samples. I also appreciated working in harmony with all other students of the Phytomedicine Programme, most especially Mr. Thanyani Ramandwa, Ms Bellonah Sakong, Ms Salaelo Raphatlelo, Ms Imelda Ledwaba and Mrs Edwina Muleya. I thank you all for your co-operation and immeasurable assistance.

Table of Contents

	Page number
Abstract	xiv
List of abbreviations	xvi
List of figures	xix
List of tables	xxi
List of appendix	xxii
 CHAPTER ONE	
Gastrointestinal disorders in diarrhoea diseases mechanisms and medicinal plants potentiality as therapeutic agents	
1.0. Introduction	1
1.1. Plant metabolites as potential therapeutic agent	2
1.2. Aims	3
1.3. Specific objectives	4
1.4. Hypothesis	4
 CHAPTER TWO	
2.0 Literature review	
2.1. Diarrhoea as a disease	5
2.2. Pathophysiology of Diarrhoea	6
2.3. Detailed pathophysiology of diarrhoea	8
2.3.1. Inflammation in diarrhoea	8
2.3.2. Oxidative damage in diarrhoea	11
2.3.3. Enteric nervous system in diarrhoea	15
2.3.4. Cystic fibrosis transmembrane conductance regulator (CFTR) regulation	16
2.4. Specific Agents of Diarrhoea	16
2.4.1. Bacterial causes of diarrhoea	16
2.4.1.1. <i>Escherichia coli</i>	16
2.4.1.2. <i>Staphylococcus aureus</i>	17
2.4.1.3. <i>Campylobacter jejuni</i>	18
2.4.1.4. <i>Shigella</i> spp	18
2.4.1.5. <i>Vibrio cholerae</i>	18
2.4.1.6. <i>Bacillus cereus</i>	19
2.4.1.7. <i>Yersinia enterocolitica</i>	19
2.4.1.8. <i>Listeria monocytogenes</i>	20

2.4.1.9. <i>Clostridium</i> spp	20
2.4.1.10. <i>Salmonella typhimurium</i>	20
2.4.1.11. <i>Enterococcus faecalis</i>	21
2.5. Fungal induced diarrhoea symptoms	21
2.5.1. <i>Candida albicans</i>	21
2.6. Viral induced diarrhoea	21
2.6.1. Rotavirus	21
2.6.2. Norovirus	22
2.6.3. Hepatitis A virus	22
2.6.4. Human immunodeficiency virus (HIV)	22
2.7. Protozoa induced diarrhoea	22
2.7.1. <i>Giardia intestinalis</i>	22
2.7.2. <i>Entamoeba histolytica</i>	23
2.7.3. <i>Cryptosporidium parvum</i>	23
2.7.4. <i>Cyclospora cayetanensis</i>	23
2.8. Parasitic induced diarrhoea	23
2.8.1. <i>Trichinella spiralis</i>	23
2.9. Immune disordered induced diarrhoea	24
2.9.1. Compromised immune system	24
2.9.2. Hyperactive immune system	24
2.10. Antibiotic therapy induced diarrhoea	24
2.10.1. Antibiotic toxicity	24
2.10.2. Alteration of digestive functionality	25
2.10.3. Overgrowth of pathogenic microorganisms	25
2.11. Diabetic complications induced diarrhoea	25
2.12. Food allergy induced diarrhoea	26
2.13. Potential mechanisms in the control of diarrhoea	26
2.13.1. Oxidative damage and antioxidants in diarrhoeal management	26
2.13.2. Inflammation and anti-inflammatory agents in diarrhoea management	26
2.13.3. Enteric nervous system in diarrhoea symptoms and treatment	26
2.14. Plants as potential source of therapeutic agents in alleviating diarrhoeal symptoms	28
2.14.1. Anti-infectious mechanisms of plant secondary metabolites against diarrhoeal pathogens	28
2.14.2. Antioxidative mechanisms of plant phytochemical as potential antidiarrhoeal agents	29
2.14.3. Anti-inflammatory mechanisms of plant phytochemical in diarrhoea management	29
2.14.4. Antidiarrhoeal mechanisms of plant phytochemical	30
2.15. Classification of phytochemicals with antidiarrhoea potential	30
2.15.1. Terpenoids	30

2.15.2. Alkaloids	33
2.15.3. Phenolic	35
2.16. Ethnobotany and scientific investigation of plant species used traditionally in treating diarrhoea in South Africa	38
2.17. Conclusion	38
CHAPTER THREE	
Plant selection, collection, extraction and analysis of selected species	
3.1. Introduction	39
3.2. Solid-liquid extraction	40
3.3. Liquid-liquid fractionation	41
3.4. Thin layer chromatography (TLC)	41
3.4.1. Phytochemical fingerprints	41
3.5. Materials and Methods	42
3.5.1. Selection of South Africa medicinal plants for antidiarrhoeal screening	42
3.5.2. Collection of plant materials	42
3.5.3. Preparation of plant material and optimization of phenolic-enriched extraction process	42
3.5.4. Phytochemical profiling	44
3.6. Quantification of the phenolic constituents of the extracts	45
3.6.1. Determination of total phenolic constituents	45
3.6.2. Determination of total tannin	45
3.6.3. Determination of proanthocyanidin	45
3.6.4. Determination of condensed tannin	46
3.6.5. Determination of hydrolysable tannin (gallotannin)	46
3.6.6. Determination of total flavonoids and flavonol	46
3.6.7. Determination of anthocyanin	47
3.7. Results	47
3.7.1. Yield of extractions and fractionations processes	47
3.7.2. Phytochemical screening (fingerprints)	48
3.7.3. Phenolic composition of the crude extracts	52
3.8. Discussion	57
3.8.1. Yield	57
3.8.2. Thin layer chromatogram	57
3.8.3. Phenolic constituents of the crude extract	58
3.9. Conclusion	60
CHAPTER FOUR	
Antimicrobial activities of the plant extracts against potential diarrhoeal pathogens	
4.0. Introduction	61

4.1. Qualitative antimicrobial (Bioautography) assay	62
4.2. Quantitative antimicrobial activity (Minimum inhibitory concentration (MIC)) assay	63
4.3. Selection of microorganisms used in the study	63
4.4. Material and Methods	64
4.4.1. Microorganism strains	64
4.4.2. Culturing of the Bacteria	64
4.4.3. Bioautography against some pathogenic microorganisms	64
4.4.4. Determination of Minimum Inhibitory Concentration (MIC) against the bacteria pathogens	64
4.4.5. Determination of Minimum Inhibitory Concentration (MIC) against the fungal pathogens	65
4.5. Results	65
4.5.1. Microbial bioautography	65
4.5.2. Minimum inhibitory concentration against bacteria	70
4.5.3. Minimum inhibitory concentration (MIC)	73
4.6. Discussion	75
4.6.1. Antimicrobial bioautography	75
4.6.2. Minimum inhibitory concentration (MIC)	75
4.7. Conclusion	77
CHAPTER FIVE	
Free radical scavenging and antioxidant activities of the extracts and fractions as antidiarrhoeal mechanism	
5.1. Introduction	79
5.1.1. Superoxide ion	81
5.1.2. Hydrogen peroxide	81
5.1.3. Hydroxyl radical	81
5.1.4. Peroxyl radical	82
5.1.5. Hypochlorous acid	82
5.1.6. Nitric oxide	82
5.2. Antioxidant assays	83
5.2.1. Antioxidant bioautography	83
5.2.2. The chemistry of some common antioxidant assays	83
5.2.2.1. Hydroxyl radical	83
5.2.2.2. Hydrogen peroxide scavenging	84
5.2.2.3. Superoxide scavenging capacity	84
5.2.2.4. DPPH	84
5.2.2.5. ABTS	85
5.2.2.6. Ferric reducing antioxidant power (FRAP)	85
5.3. Materials and Methods	85

5.3.1. Antioxidative profile of the crude extracts and fractions using DPPH radical solution	86
5.3.2. Antioxidative assays	86
5.3.2.1. DPPH free radical-scavenging method	86
5.3.2.2. ABTS free radical-scavenging method	86
5.3.2.3. Ferric reducing antioxidant power (FRAP)	87
5.3.2.4. Hydroxyl radical scavenging assay	87
5.3.2.5. Lipid peroxidation inhibition assay	87
5.4. Result	87
5.4.1. TLC-DPPH analyses	87
5.4.2. DPPH effective concentration (EC ₅₀)	90
5.4.3. ABTS effective concentration (EC ₅₀)	92
5.4.4. FRAP gradient	93
5.4.5. Hydroxyl radical effective concentration (EC ₅₀)	94
5.4.6. Lipid peroxidation inhibition effective concentration (EC ₅₀)	95
5.5. Discussion	96
5.5.1. Qualitative antioxidant analyses (DPPH-TLC bioautography)	96
5.6. Conclusion	98
CHAPTER SIX	
Anti-inflammatory activities of the crude extracts as antidiarrhoeal mechanisms	
6.0. Introduction	100
6.1. Effect of cyclooxygenases (COX) on GIT	101
6.2. Effects of lipoyxygenase (LOX) on GIT	101
6.3. Effects of cytokines on GIT	102
6.4. Oxidative species as inflammatory mediator	102
6.5. Allopathic anti-inflammatory therapies and adverse effects on GIT	103
6.6. Plant phytochemicals as anti-inflammatory agents	105
6.7. Mechanisms of anti-inflammatory assay models	105
6.8. Materials and Methods	106
6.8.1. COX assay	106
6.8.2. LOX assay	106
6.9. Results	107
6.9.1. COX	107
6.9.2. LOX	108
6.10. Discussion	109
6.10.1. COX	109
6.10.2. LOX	109
6.11. Conclusion	110

CHAPTER SEVEN

Cytotoxicity evaluation of the crude extracts against Vero African green monkey kidney cell lines

7.0. Introduction	111
7.1. Materials and Methods	112
7.1.1. Preparation of plant extracts	112
7.1.2. Cytotoxicity assay against Vero cell	112
7.2. Results	113
7.3. Discussion	114
7.4. Conclusion	115

CHAPTER EIGHT

Motility modulation potential of *Bauhinia galpinii* and *Combretum vendae* phenolic-enriched leaf extracts on isolated rat ileum

8.0. Introduction	116
8.1. Drugs and reagents	117
8.2. Animal care	117
8.2.1. Isolated ileum preparation	118
8.3. Contractility test	118
8.3.1. Spasmogen assay	118
8.3.2. Spasmolytic assays	118
8.3.2.1. Effects on acetylcholine-induced contractility	118
8.3.2.2. Effects on serotonin-induced contractility	118
8.3.2.3. Effects on KCl-induced contractility	119
8.4. Data analysis	119
8.5. Results	119
8.5.1. Effect of <i>B. galpinii</i> crude extract on isolated rat ileum	119
8.5.2. Effect of <i>C. vendae</i> crude extract on isolated rat ileum	122
8.6. Discussion	123
8.7. Conclusion	126

CHAPTER NINE

Isolation and characterization of antimicrobial and antioxidant compounds from *Bauhinia galpinii* and *Combretum vendae*

9.0. Introduction	127
9.1.1. Column chromatography	128
9.1.2. Mass spectrometry	128
9.2. Materials and Methods	128
9.2.1. Preparation of plant extracts	128
9.2.2. Bioautography	128

9.2.3. Isolation of bioactive triterpenoids from <i>C. vendae</i>	129
9.2.4. Isolation of phenolic compounds from <i>C. vendae</i>	130
9.3. Isolation of compounds from <i>B. galpinii</i>	130
9.3.1. Isolation of bioactive triterpenoids from <i>B. galpinii</i>	130
9.3.2. Isolation of phenolic compounds from <i>B. galpinii</i>	130
9.4. Characterization of the isolated compounds	131
9.4.1. NMR spectroscopy	132
9.4.2. Mass spectrometry	132
9.4.3. Ultra-violet spectroscopy	132
9.5. Results	132
9.5.1. Identification of the chemical structures of isolated compounds from <i>C. vendae</i>	132
9.5.2. Antimicrobial activity of isolated compounds from <i>C. vendae</i>	135
9.5.3. Identification of the chemical structures of isolated compounds from <i>B. galpinii</i>	136
9.5.4. Antimicrobial activity of isolated compounds from <i>B. galpinii</i>	140
9.6. Discussion	140
9.6.1. Bioactive compounds from <i>C. vendae</i>	140
9.6.2. Bioactive compounds from <i>B. galpinii</i>	141
9.7. Conclusion	143
CHAPTER 10	
General conclusion and future prospects	
10. Introduction	144
10.1. Identification of diarrhoeal pathogenesis and medicinal plants used as therapeutic Agents	145
10.2. Antimicrobial evaluation of the extracts against infectious pathogens	145
10.3. Antioxidant evaluation of the extracts	145
10.4. Anti-inflammatory potential of the extracts	146
10.5. Toxicity risk of the extracts	146
10.6. Motility modulatory effects of <i>Bauhinia galpinii</i> and <i>Combretum vendae</i>	147
10.7. Isolation and characterisation of bioactive compounds	147
CHAPTER 11	
References	149

Abstract

Diarrhoea is one of the killer diseases resulting from the dehydration and loss of electrolytes through profuse and excessive excretion of loose stool. The pathoaetiologies include infections, intestinal inflammation, imbalanced intestinal oxidative homeostasis and altered motility. Treatment with oral rehydration therapy (ORT) is a key intervention especially in secretory diarrhoea as supportive therapy. Symptomatic and non-symptomatic therapies directed at treating the intestinal tissues are available. However, these conventional treatments are still not sufficient in curing diarrhoea due to their associated hazards such as the development and spread of drug-resistant pathogens, changes in normal intestinal bacteria flora and potential chronic toxicity. Therapies targeted at intestinal tissue include antimotility and antisecretory agents have adverse effects such as addictiveness, constipation and fatal ischaemic colitis. Many ethnopharmacological and ethnobotanical therapies for treating diarrhoea exist among different cultures. The aims of this study were to evaluate the biological activities of plant extracts against some diarrhoeal pathophysiology.

A literature search in English of published articles and books that discussed ethnobotanical uses of medicinal plants in southern Africa was conducted. A list of 230 medicinal plants used in South African traditional medicines for treating diarrhoea and associated complications was created. The list included family, genus, species, biological activities and bioactive isolates as well as the remedies for diarrhoea. Twenty seven species were selected to evaluate for antimicrobial, antioxidant and anti-inflammatory activities. Safety of the plants was determined by determining the cytotoxicity of the crude extracts against Vero African green monkey kidney cell lines using a standard method. Motility effects of *Bauhinia galpinii* (BGE) and *Combretum vendae* (CVE) were determined by modulation of the contractility process of the isolated rat ileum induced by spasmogens.

Phenolic compositions of the crude extract were determined using various standard methods and finally bioactivity guided isolation of antimicrobial and antioxidant compounds from BGE and CVE were carried out using open column chromatography. Identification and characterization of the isolated compounds was achieved by NMR, EI-MS and UV spectroscopy.

The non-polar fractions had good antimicrobial activities with MIC ranged between 19 – 1250 µg/ml while the polar fraction had moderate antimicrobial activities with MIC ranged between 39 - >2500 µg/ml. In general the non-polar fractions had a higher antimicrobial activity.

The crude extracts contained wide range phenolic compounds with a total phenolic (74.91±1.26 to 467.04±15.82 mg GAE/g plant material), and total flavonoids (11.27±3.37 to 176±5.96 mg EQ/g plant material). The antioxidant activities were concentrated and potentiated in the polar fractions. The non-polar fractions had poor antioxidant activities with EC₅₀ values ranging from 0.21±0.03 to 303.65±3.84 µg/ml for DPPH radical scavenging and 0.43±0.03 to 1709±91.44 µg/ml for ABTS radical scavenging.

The crude extracts had selective COX-1 inhibitory activities ranging between 41.70 to 84.61% and had no COX-2 inhibitory activity. All the extracts tested had 15-LOX inhibitory capacity with LC₅₀ values ranging between 0.86±0.27 and 111.44±37.28 µg/ml. The cytotoxicity results indicated a wide variation in toxic potential of the crude extracts with LC₅₀ values ranging from 3.51 to 741.90µg/ml.

The BGE extracts had dual activities as spasmolytic by stimulating the spontaneous contractility and also agonised contractions induced by spasmogens but it inhibited K^+ induced contraction. CVE had spasmodic activities through a multiple mechanisms inhibiting contractions induced by spasmogens and K^+ in a dose-dependent manner.

Several bioactive compounds were isolated from the *Combretum vendae* leaves, There were triterpenoids (ursol-12-en-28-oic acid, mixtures of corosolic acid and maslinic acid, and asiatic acid and arjunolic acid) as well as bibenzyls combretastatin B5-O-2'- β -D-glucopyranoside, combretastatin B1-O-2'- β -D-glucopyranoside and a flavonoid (apigenin)..

From *Bauhinia galpinii* the following bioactive compounds were isolated and characterized: β -3 ethoxy sitosterol, one new flavone (5, 7, 4', 5' tetrahydroxy-2'-methoxyflavone (isoetin 2'-methyl ether) or 5, 7, 2', 5' tetrahydroxy-4'-methoxyflavone (isoetin 4'-methyl ether)), 3, 5, 7, 3', 4'-pentahydroxyflavone and 3, 5, 7, 3', 4', 5'-hexahydroxyflavone, quercetin-3-O- β -galactopyranoside and myricetin-3-O- β -galactopyranoside

The extraction protocol used in this work potentiated the antimicrobial activities in the non-polar fractions while antioxidant activities were potentiated in the polar fractions. This indicated that using polar solvents as extractant for treating infectious diarrhoea may not be quite effective unless some other antidiarrhoeal mechanisms are involved. Therefore, mixture of organic solvent (ethanol) and water can be recommended for broad-based activity.

Bauhinia galpinii extracts had a dual- mechanism of action (prokinetic and relaxant) on gastro-intestinal motility, depending on the prevalent patho-physiological condition and *Combretum vendae* mediated spasmolytic effects on isolated rat ileum through multiple inhibitions of a wide range of contractile stimuli. Hence, the presence of multiple acting spasmolytic activities in the plant extract might be contributing towards its effectiveness in treating diarrhoea and abdominal spasm. The uses of these plants in traditional medicine need to be monitored closely because of the selective inhibition of COX-1 and its associated GIT injury, and the high toxicity potential of some of the extracts.

Further work evaluating the antidiarrhoea mechanisms, identification and isolation of bioactive compounds, sub-acute and acute toxicity of the plant extracts is recommended.

Key words: Antimicrobial, antioxidant, anti-inflammatory, diarrhoeal, antispasmodic, enteric nervous system, cytotoxicity.

List of Abbreviations

A

ABTS=2,2'-azino-bis-(3-ethylbenzothiazoline-6-sulfonic acid)

AMP=Antimicrobial peptides

B

BAB= *Bauhinia bowkeri*

BAG= *Bauhinia galpinii*

BAP= *Bauhinia petersiana*

BAV= *Bauhinia variegata*

BGE= *Bauhinia galpinii* extract

C

Ca²⁺= Calcium ion

Cl⁻= chloride ions

CNF-1= Cytotoxic necrotising factor 1

CNS= Central nervous system

COB= *Combretum bracteosum*

COP= *Combretum padoides*

COV= *Combretum vendae*

COX= Cyclooxygenase

COW= *Combretum woodii*

CVE= *Combretum vendae* extract

D

DAEC= diffusively adherent *Escherichia coli*

DNA

DPPH=2, 2-diphenyl-1-picrylhydrazyl

E

EAEC= Enteroaggregative *Escherichia coli*

EHEC= Enterohaemorrhagic *Escherichia coli*

EIEC= Enteroinvasive *Escherichia coli*

ENS= Enteric nervous system

EPEC= Enteropathogenic *Escherichia coli*

ETEC= Enterotoxigenic *Escherichia coli*

EUC=*Euclea crispa*

EUN= *Euclea natalensis*



F

FIC= *Ficus cratostoma*

FIG=*Ficus glumosa*

FRAP= Ferric reducing antioxidant capacity

G

GIT= Gastrointestinal tract

H

HIV/AIDS= Human immune deficiency virus/Acquired immune deficiency syndrome

HOCl= hypochlorite

HUB= Haemolytic uremic syndrome

I

IBS= Irritable bowel syndrome

IL= Interleukin

INC= *Indigofera cylindrica*

iNOS= inducible nitric oxide synthase

INT= p-iodonitrotetrazolium

L

LT= Heat labile enterotoxin

LTB= Leukotriene B

M

MDA= Malondialdehyde

MCP-1= Monocyte chemoattractant protein

MIC= Minimum inhibitory concentration

MPD= *Maytenus peduncularis*

MPR= *Maytenus procumbens*

MSE= *Maytenus senegalensis*

MUN= *Maytenus undata*

N

Na⁺= sodium ions

NAME= nitro

NH₂Cl= Ammonium chloride

NO= Nitric oxide

O

OH⁻ = Hydroxyl radical

ORT=Oral rehydration therapy

OZM= *Ozoroa mucronata*

OZP= *Ozoroa paniculosa*

P

PG= Prostaglandin

R

ROS= Reactive oxygen species

RNS= Reactive nitrogen species

S

SCB=*Schotia brachypetala*

SLE= *Searsia leptodictya*

SPD= *Searsia pendulina*

SPT= *Searsia penteri*

ST= Heat stable enterotoxins

SYP= *Syzygium paniculatum*

T

TLC=Thin layer chromatography

TNF- α = Tumour necrosis factor- α

Trolox= 6-hydroxy-2, 5, 7, 8-tetrahydroxyl-chroman-2-carboxylic acid

U

UNICEF=United Nation Children Fund

W

WHO= World Health Organization

List of Figures

Chapter 2

Fig. 2.1. Classification of the diarrhoea and the stimulants	6
Fig. 2.2. Cytokines production network in the tissues	8
Fig. 2.3. Biosynthetic pathways for the eicosanoids	9
Fig. 2.4. Intestinal epithelial TJs as a physical barrier	10
Fig.2.5. The integrative pathophysiology and mechanism of diarrhoeal disease	13
Fig. 2.6. Lipid peroxidation chain reactions	14
Fig. 2.7. Chemical structures of the lipid peroxidation intermediates	14
Fig. 2.8. Mechanisms of antibiotic-induced diarrhoea	25
Fig. 2.9. Chemical structures of bioactive terpenoids against diarrhoeal mechanisms	31
Fig. 2.10. Chemical structures of bioactive alkaloids against diarrhoeal mechanisms	33
Fig. 2.11. Sub-classes of biologically important phenolic compounds	34
Fig. 2.12. Chemical structures of bioactive phenolics against diarrhoeal mechanisms	36

Chapter 3

Fig.3.1. Flow chart for the extraction, phytochemical analysis and fractionation of the crude extracts	43
Fig.3.2. TLC phytochemical profile of the crude extracts	48
Fig.3.3. TLC phytochemical profile of the hexane fractions	49
Fig.3.4. TLC phytochemical profile of the dichloromethane fraction	50
Fig.3.5. TLC phytochemical profile of the ethyl acetate fraction	51
Fig.3.6. Total phenolic and non-tannin constituents of the crude extract	52
Fig.3.7. Total tannin and condensed tannin constituents of the crude extracts	53
Fig.3.8. Proanthocyanidin and gallotannin constituents of the crude extract	54
Fig.3.9. Total flavonoid and flavonol constituents of the crude extract	56

Chapter 4

Fig.4.1. The classification of microbiological methods for biological detection	63
Fig.4.2. Bioautography of the hexane fractions against <i>S. aureus</i>	65
Fig.4.3. Bioautography of the dichloromethane fractions against <i>S. aureus</i>	66
Fig.4.4. Bioautography of hexane fractions of different plant species against <i>E. faecalis</i>	66
Fig.4.5. Bioautography of dichloromethane fractions of different plant species against <i>E. coli</i>	67
Fig.4.6. Bioautography of dichloromethane fractions of different plant species against <i>E. faecalis</i>	67
Fig.4.7. Bioautography of hexane of different plant species against <i>C. neoformans</i>	68
Fig.4.8. Bioautography of dichloromethane fractions against <i>C. neoformans</i>	68
Fig.4.9. Bioautography of hexane fractions against <i>A. fumigatus</i>	69
Fig.4.10. Bioautography of dichloromethane fractions against <i>A. fumigatus</i>	69
Fig.4.11. Bioautography of hexane fractions against <i>C. albicans</i>	70

Fig.4.12. Bioautography of dichloromethane fractions against <i>C. albicans</i>	70
Chapter 5	
Fig.5.1. Some deleterious reactions from the production of reactive free radicals in biological systems	80
Fig.5.2. TLC-DPPH profiles of the crude extracts of extracts of different plants	88
Fig.5.3. TLC-DPPH profile of the hexane fractions of different plants	89
Fig.5.4. TLC-DPPH profiles of the dichloromethane fractions of different plants	89
Fig.5.5. TLC-DPPH profiles of the ethyl acetate fractions of different plants	90
Chapter 6	
Fig.6.1: Roles of COX in the pathogenesis mechanism of NSAID-induced intestinal damage	103
Fig.6.2: Factors involved in the pathogenesis of indomethacin-induced small intestinal lesions	104
Fig.6.3: COX-1 inhibitory activity of some selected phenolic-enriched crude extracts	106
Chapter 8	
Fig.8.1.Schematic presentation of the contractility assay using isolated rat ileum	118
Fig.8.2.Stimulatory effects <i>B. galpinii</i> on spontaneous contraction of isolated rat ileum	119
Fig.8.3.Effects of 70% acetone crude leaf extract of <i>B. galpinii</i> on the acetylcholine cumulative concentration dependent-induced contraction in the absence and presence of atropine	119
Fig.8.4.Agonized effects of <i>B. galpinii</i> on serotonin-induced contraction of the isolated rat ileum	120
Fig.8.5.Relaxant effects of <i>B. galpinii</i> on KCl-induced contraction of the isolated rat ileum	120
Fig.8.6.Spasmolytic effects of 70% acetone crude leaf extract of <i>C. vendae</i> on acetylcholine-induced contraction of the isolated rat ileum	121
Fig.8.7.Spasmolytic effects of 70% acetone crude leaf extract of <i>C. vendae</i> on serotonin-induced contraction of the isolated rat ileum	121
Fig.8.8.Spasmolytic effect of the <i>C. vendae</i> on the depolarised KCl-induced isolated rat ileum contractions	122
Chapter 9	
Fig.9.1. Extraction, fractionation and isolation of bioactive compounds from the leaf extract of <i>Combretum vendae</i>	128
Fig.9.2. Extraction, fractionation and isolation of bioactive compounds from the leaf extract of <i>Bauhinia galpinii</i>	130
Fig. 9.3 Chemical structures isolated bioactive compounds from the leaf extract of <i>C. vendae</i>	134
Fig. 9.4 Chemical structures isolated bioactive compounds from the leaf extract of <i>B. galpinii</i>	138

List of Tables

Chapter 2

Table 2.1. The mechanism of action and symptoms of enteric pathogenic <i>E.coli</i>	17
Table 2.2. Neurotransmitters of ENS causing intestinal secretion in diarrhoea	27

Chapter 3

Table 3.1. Medicinal plants selected for the antidiarrhoeal investigations in this study	42
Table 3.2. The percentage yield of the crude extract and fractions (g/g dried plant material)	47

Chapter 4

Table 4.1. The minimum inhibitory concentration (MIC) of the crude extracts and fractions against bacterial strains tested	72
Table 4.2. The minimum inhibitory concentration (MIC) of the crude extracts and fractions against fungal strains tested	74

Chapter 5

Table 5.1. DPPH radical scavenging potential of the crude extracts and fractions expressed as EC ₅₀ (µg/ml)	91
Table 5.2. ABTS radical scavenging potential of the crude extracts and fractions expressed as EC ₅₀ (µg/ml)	92
Table 5.3. FRAP	94
Table 5.4. Hydroxyl radical scavenging potential of the crude extracts and fractions expressed as EC ₅₀ (µg/ml)	94
Table 5.5. Linoleic acid peroxidation inhibition expressed as LC ₅₀ (µg/ml)	95

Chapter 6

Table 6.1. Lipoxygenase inhibitory activity of the crude extracts	107
---	-----

Chapter 7

Table 7.1. The LD ₅₀ of the cytotoxicity assay of some medicinal plants used in South African traditional medicine to treat diarrhoea and related ailments	112
---	-----

Chapter 9

Table 9.1: NMR experiments commonly applied for natural product structural elucidation	126
Table 9.2: Minimum inhibitory concentration (µg/ml) of the isolated compounds from the leaf extract of <i>C. vendae</i>	135
Table 9.2: Minimum inhibitory concentration (µg/ml) of the isolated compounds from the leaf extract of <i>B. galpinii</i>	139

List of Appendix

	Page number
Appendix 2.1. Ethnobotanical and literature information of medicinal plant species used traditionally for treating diarrhoea in South Africa	172
Appendix 9.1. 1D and 2D NMR spectra data of Ursolic acid	197
Appendix 9.2. 1D and 2D NMR spectra data of mixture of corosolic acid and maslinic acid	198
Appendix 9.3. 1D and 2D NMR spectra data of mixture of asiatic acid and arjunolic acid	199
Appendix 9.4. 1D and 2D NMR spectra data of combretastatin B5-2'-O- glucopyranoside	200
Appendix 9.5. 1D and 2D NMR spectra data of combretastatin B1-2'-O- glucopyranoside	200
Appendix 9.6. 1D and 2D NMR spectra data of 3 β -ethoxy sitosterol	201
Appendix 9.7. 1D and 2D NMR spectra data of quercetin	201
Appendix 9.8. 1D and 2D NMR spectra data of myricetin	202
Appendix 9.9. 1D and 2D NMR spectra data of isoetin 2' methyl ether/ isoetin 4' methyl ether	202
Appendix 9.10. 1D and 2D NMR spectra data of quercetin-3-O- β -galactopyranoside	203
Appendix 9.11. 1D and 2D NMR spectra data of myricetin-3-O- β -galactopyranoside	203

CHAPTER ONE

Gastrointestinal disorders in diarrhoea diseases mechanisms and medicinal plants potentiality as therapeutic agents.

1. Introduction

The gastrointestinal tract (GIT) is dedicated to processing and absorbing nutrients and fluids essential for the maintenance of good health (Martinez-Augustin *et al.*, 2009). For the GIT to function optimally, a balance is maintained between intestinal motility and intestinal fluid volume. The latter process is finely regulated through the control of fluid absorption via intestinal villous epithelial cells and secretion across the intestine via intestinal crypt cells (Martinez-Augustin *et al.*, 2009). Net fluid absorption driven by osmotic gradients controlling the movement of electrolytes (sodium ions [Na⁺] and chloride ions [Cl⁻]), sugars and amino acids across the epithelial lining of the lumen, predominate in these opposing processes (Pash *et al.*, 2009). In contrast, motility is controlled by the activation of enteric nervous system (ENS) by either neurotransmitters, inflammatory mediators or epithelium membrane lipid peroxidation by-products (Wood, 2004). Any upset of this delicate intestinal fluid balance (decrease fluid absorption and increase fluid secretion), and/or changes in GIT motility usually causes intestinal disorders clinically evident as diarrhoea (Vitali *et al.*, 2006).

Diarrhoea is loosely defined as an alteration in the normal bowel movement characterized by an increase in the volume, frequency and water content of stool (Baldi *et al.*, 2009). The pathophysiology of diarrhoea include microbial and parasitic infections (Hodges and Gill, 2010), stress (oxidative or physical) (Soderholm and Perdue, 2006), dysfunctional immunity (Schulzke *et al.*, 2009), disrupt GIT integrity and neurohumoral mechanisms (Vitali *et al.*, 2006; Spiller, 2004). Diarrhoea can also be a symptom of other diseases such as cholera, irritable bowel syndrome (IBS), gastroenteritis (intestinal inflammation and ulcerative colitis) (Schiller, L. R., 1999; Baldi *et al.*, 2009), malaria (Gale *et al.*, 2007) and diabetes mellitus (Forgacs and Patel, 2011).

The mechanism causing diarrhoea can be secretory (resulting from osmotic load within the intestine), hyper motility (resulting from rapid intestinal transitions) or hypo motility (resulting in decreased intestinal fluid re-absorption) or combination of these mechanisms (Vitali *et al.*, 2006). The symptoms are either caused by an increase in fluid and electrolyte secretion predominantly in the small intestine or a decrease in absorption which can involve both small and large intestine (Pash *et al.*, 2009; Spiller and Garsed, 2009). Physiologically, diarrhoea is considered beneficial to the GIT as it provides an important mechanism of flushing away harmful luminal substances (Valeur *et al.*, 2009). However, diarrhoea becomes pathological when the loss of fluids and electrolytes exceeds the body's ability to replace the losses.

As a disease, diarrhoea is considered one of the most dangerous GIT disorders as death can result in severe cases due to dehydration and loss of electrolytes (WHO and UNICEF, 2004). According to the World Health Organization (WHO)/United Nation Children Fund (UNICEF) report, more than 1 billion diarrhoeal episodes occurred in human across the world yearly, with about 5 million deaths especially in infants (Thapar and

[Sanderson, 2004](#)). In addition to causing acute disease and mortality, diarrhoea associated malnutrition could result in stunted growth, non-optimal immune functionality and increase susceptibility to infections. Diarrhoea therefore poses a major health challenge to human, as it could lead to premature mortality, disability and/or increase health-care costs ([Guerrant et al., 2005](#)).

In animal production, diarrhoea is presumed to impose heavy productivity losses on affected farms, although true effects in monetary terms cannot be easily appreciated. The apparent on-farm losses are reduction in productivity (milk, wool, egg, meat and meat quality), increased mortality and morbidity, weight loss and abortion ([Chi et al., 2002](#)). Episodes of diarrhoeal diseases can also affect the export market and hurt consumer's confidence in the products ([Yarnell, 2007](#)).

The most common modern method of managing diarrhoea is the replacement of lost fluids and electrolytes with either oral or intravenous electrolyte preparation ([Thapar and Sanderson, 2004](#)). While fluid replacement is usually effective, severe fluid losses requires additional pharmacological treatment to mitigate the on-going fluid loss. For this, drugs with antispasmodic, antimotility, antioxidative, anti-secretory/pro-absorptive and/or anti-inflammatory properties (depending on the causative agents) may be used to treat diarrhoeal ([Wynn and Fougere, 2007](#)). The issue of antimicrobial therapy for self-limiting and non-infectious diarrhoea is usually not encouraged to avoid development of drug resistance microbes. However, in cases of established infectious diarrhoea with known pathogenic agents, specific therapeutic intervention using antimicrobial drugs targeting the causative microbes may be applied. At present, the current standard therapeutic options are insufficient because of limited available modalities with broad based activities against the large number of diarrhoeal disease mechanisms and apparent side effects. The problems associated with some of the standard therapies include antimicrobial resistance, drug toxicity, constipation and addiction.

As a result there is an urgent need for new therapeutic drugs with lower cost, high efficacy, little or no side effects and wider availability especially in rural areas where diarrhoea causes large scale infant mortality. Plants, which serves as dietary source to animals and people, may also provide a good source of new therapeutic drugs.

1.2. Plant metabolites as potential therapeutic agent

Plant serves as dietary source to animals and humans providing sufficient nutrients to meet metabolic requirements for their well being, growth and productivity. However, it can also contribute to achieving optimal health and development as well as serving an essential role in reducing the risk or delaying the onset of diseases and disorders ([Kosar et al, 2006](#); [Halliwell, 1997](#)). Medicinal plants have therapeutic properties due to biosynthesis of various complex phytochemical substances grouped broadly as phenolics, alkaloids and terpenoids. Synergistic interaction among the multiple phytochemicals may be responsible for the overall bioactivity of a given medicinal plant. Pharmacological and clinical studies of phytochemical in plants have shown that they exhibit various medicinal uses and serve as the major backbone of traditional medicine ([Van Wyk and Wink, 2004](#)). Medicinal plants have played some key roles in the health care needs of rural and urban

settlements for human, livestock and animals. Plant extracts, formulations, or pure natural compounds are used in controlling diverse diseases ranging from coughs, inflammation, and diarrhoea to parasitic infection in human and veterinary medicine. A large number of these medicinal plants have been screened and validated for their ethnopharmacological use as antidiarrhoeal agents of varied mechanisms (Gutierrez *et al.*, 2007). However, the literatures available on the pharmacological evaluation of medicinal plants used traditionally in treating diarrhoea in South Africa are mainly on antimicrobial screening models. Little literature information is available on other antidiarrhoeal mechanisms and *in vivo* study

For this study, 27 South African medicinal plants used as diarrhoeal remedies with ethnopharmacological background identified as requiring further biological evaluation. Thereafter, *Bauhinia galpinii* and *Combretum vendae* were choosing for further investigation based on the results from preliminary screening. *Bauhinia galpinii* was previously investigated for its antioxidant activities and three compounds (two active and one inactive) were isolated from the acetone leave extract (Aderogba *et al.*, 2007). Methanol and dichloromethane leaf extracts of *B. galpinii* are reported to have antimutagenic property (Reid *et al.*, 2006). The acetone root extract of *B. galpinii* has also been found to be highly cytotoxic (LD₅₀ 2.70 µg/ml) against Vero cell lines (Samie *et al.*, 2009). Antimicrobial activity of *Combretum vendae* against four bacterial pathogens (Ahmed *et al.*, 2009) and apigenin has been isolated from the acetone leaf extract (Eloff *et al.*, 2008).

In many previous studies relatively non-polar extractants were used despite the fact that traditionally aqueous extracts are used. This is probably due to difficulties in analyzing complex molecules extracted by polar extractants, because phenolics may play an important role in managing diarrhoea the focus of this study will be on more polar extractant.

1.3. Aims

To investigate the biological activities of the phenolic-enriched extracts and fractions of 27 medicinal plants against some diarrhoea pathoaetiologies and evaluating the antidiarrhoeal mechanisms of *Bauhinia galpinii* and *Combretum vendae* extracts using *in vitro* isolated organ methods, as means of validating their ethnopharmacological used in South African traditional medicine to treat diarrhoea.

1.4. Specific objectives

- ❖ To evaluate the effect of the extracts, fractions and isolated compound(s) against pathogenic microbes that are known to induce diarrhoea.
- ❖ To determine the antioxidative properties of the extracts, fractions and isolated compound(s) using the DPPH radical scavenging, the ABTS radical scavenging, the hydroxyl radical scavenging, the linoleic acid peroxidation inhibition and the ferric reducing antioxidant power (FRAP).
- ❖ To determine the effects of the most promising extracts on the contractility process of the isolated rat ileum induced by spasmogens, receptor agonists, antagonists and ion channels activators.

- ❖ To fractionate the extracts and elucidate the component(s) that exhibit antimicrobial and antioxidant properties.
- ❖ To evaluate the safety, efficacy and toxicity of the crude extracts and the pure active component(s).

1.5. Hypothesis

The phytochemical constituents of medicinal plants used in traditional medicine have antioxidant, anti-inflammatory, antimicrobial and /or anti-spasmodic activities that could help in alleviating diarrhoeal diseases in human and animals.

CHAPTER TWO

2.0. Literature review

2.1. Diarrhoea as a disease

Diarrhoea is a common clinical sign following on altered bowel movement, decreased intestinal absorption of fluids and increased intestinal electrolyte secretion resulting in loose and watery stool (Baldi *et al.*, 2009). The mechanisms of diarrhoea diseases can be secretory due to impaired electrolyte absorption and osmotic load within the intestine, hyper motility resulting from rapid intestinal transitions of material or hypo motility resulting in decreased intestinal fluid re-absorption or combination of these mechanisms (Vitali *et al.*, 2006). The symptoms are either caused by an increase in fluid and electrolyte secretion predominantly in the small intestine or a decrease in absorption which can involve both small and large intestine (Pash *et al.*, 2009; Spiller and Garsed, 2009).

Diarrhoeal disease can be either infectious or non-infectious in nature with infection pathogenesis responsible for the major total episode worldwide. In infectious diarrhoea, the potential causative pathogens include bacterial agents (Mathabe *et al.*, 2006), rarely fungal (Robert *et al.*, 2001), viral and parasite pathogens (Brijesh *et al.*, 2006). Non-infectious diarrhoea can be caused by adverse reactions to drugs, toxins, allergy to food, poisons and acute inflammation which promote the release of secretagogues and some enteric nervous system (ENS) receptors (prostaglandin, serotonin, substance P, vasoactive intestinal peptides, and hormone) in the GIT (Wynn and Fougere, 2007). Diarrhoea is usually classified according to the duration of the symptoms:

- Acute diarrhoea: mostly caused by enteric pathogenic infections, intoxicants or food allergy. This type of diarrhoea is self-limiting without pharmacological intervention and usually resolves within two week from onset or,
- Persistent diarrhoea: mostly result from a secondary cause such as enteric infections or malnutrition, and usually last for more than 14 days, or
- Chronic diarrhoea: mostly result from congenital defects of digestion and absorption. This usually last for more than 30 days (Thapar and Sanderson, 2004; Baldi *et al.*, 2009).

Other methods of classifying diarrhoea include stool characteristics or pathological mechanisms such as watery, osmotic, altered motility or inflammatory diarrhoea (Ravikumara, 2008) as shown in Fig. 2.1.

- Watery diarrhoea typically referred to as secretory diarrhoea results from increased chlorine secretion, decreased sodium absorption and increased mucosal permeability.
- Osmotic diarrhoea, also a watery form of diarrhoea, is caused by the ingestion of non-absorbable indigestible material (Baldi *et al.*, 2009) or absence of brush border enzymes required for the digestion of dietary carbohydrates (Podewils *et al.*, 2004).

- Inflammatory diarrhoea is characterized by the presence of mucus, blood, and leukocytes in the stool, and is usually induced by an infectious process, allergic colitis or inflammatory bowel disease (IBD) (Ravikumara, 2008).




Fig.2.1: Classification of diarrhoea and the stimulants (modified from Ebert, 2005)

PGE₂= prostaglandin E₂, PGDF_a= prostaglandin D_{Fa}, NO=nitric oxide, HOCl=hydrogen chlorate, H₂O₂=hydrogen peroxide, OH=hydroxyl radical, SBBO = small-bowel bacterial overgrowth

2.2. Pathophysiology of Diarrhoea

A “healthy gastrointestinal tract (GIT)” can be defined as one where a balance is reached between the bacteria colonising the environment and the immune system. Any disturbance in this homeostasis will result in GIT disorders like diarrhoea. The general causes of diarrhoea are: microbial infection (bacteria, viruses and parasites), intestinal inflammation, altered GIT motility as a result of damage to enteric nervous system (ENS) and immune dysfunctions. The mechanisms of infectious diarrhoea include:

- Microbial attachment and localized effacement of the intestinal epithelium: Some enteric pathogens have the ability to attach and alter the surface of the invaded cell characterized as attaching and effacing lesion. Attachments of the infectious pathogens to the apical surface of the enterocyte-like cells create favourable conditions for bacterial growth and multiplication (Ramaroa and Lereclus, 2006). The mechanism of effacing lesion involves the localized destruction of the adjacent epithelial microvilli and

the formation of a pedestal-like structure from the accumulation of cytoskeletal proteins, such as actin, beneath the site of attachment (Thapar and Sanderson, 2004; Guerrant *et al*, 1999).

- Production of enterotoxins that subvert mucosal transport systems: In the case of intoxication, the causative organisms may or may not be present in the transmitting medium, but act through preformed enterotoxins (examples of such organisms are *S. aureus* (α -haemolysin) and emetic type *Bacillus cereus* (cereulide) (Granum, 2006). The toxins may be cytotoxic or haemolytic, thus causing damage to intestinal epithelial cells. The mechanism of actions used by the enterotoxins to cause diarrhoea include (Laohachai *et al*, 2003):
 - (1) Decrease in intestinal surface area and, hence a decreased fluid absorption rate.
 - (2) Changes in mucosal osmotic permeability, resulting from mucosal destruction.
 - (3) Changes in fluid and electrolytes homeostasis through the toxin's action on ion channels.
- Direct epithelial cell invasion: Epithelial cells are the first major cell type encountered by infectious pathogens in the intestinal mucosa and the main site of host-pathogens interaction (Ramaroa and Lereclus, 2006). Intestinal mucosal epithelial cells are essential for initiation of the immune response to different microorganisms (Hodges and Gill, 2010). In addition to forming a physical barrier that protects the host's internal organs from the external environment, epithelial cells produce a variety of cytokines and chemokines in response to microbial infection (MacNaughton, 2006). The survival strategy of some pathogens is the invasion of epithelial mucosa cells through the activation or inhibition of different signal transduction pathways and induces cytoskeletal rearrangement within the host cell.
- Production of cytotoxins: Microbial cytotoxins degrades of the epithelial cell surface membrane and consequently results in loss of host epithelial cell layer (Ramaroa and Lereclus, 2006; Thapar and Sanderson, 2004). Diarrhoea occurs through the destruction of the epithelial cells due to loss of absorptive surface area and impaired secretion mechanisms.
- Immune activation: Some pathogens may induce diarrhoea indirectly via excessively stimulation of the immune system. The inflammatory mediators such as cytokines (interferon- γ , tumour necrosis factor (TNF- α), interleukin-6 (IL-6), and IL-1 β) (Johnson *et al.*, 2010), reactive oxygen species (ROS), reactive nitrogen species (RNS) (Sprague and Khalil, 2009) can all interfere with the epithelial tight junctions (TJs), thereby resulting in diarrhoea.
- Immune inhibition: With the direct involvement of the immune system in the protection of the body against both pathogenic and enteric flora, any descent in functionality in the immune could allow the opportunity for pathogenic and/or enteric microbes to establish themselves in abundance with resultant decrease in GIT function and diarrhoea.

2.3. Detailed pathophysiology of diarrhoea

2.3.1. Inflammation in diarrhoea

Inflammation is the body's first line of defence against infection and hazardous stimuli in people and animals (Iwalewa *et al.*, 2007) with injury or infection in the GIT, resulting in the activation of neutrophils and macrophages. Once activated, the immune cell (e.g. macrophage) assist with the killing of pathogenic microorganisms and/ or the removal of harmful and cell debris (Stables and Gilroy, 2010). This task is achieved through the release of numerous pro-inflammatory cytokines (tumour necrosis factor [TNF]- α and interleukin [IL]-1 β , IL-3, IL-6); chemokines and chemoattractants (IL-8 and monocyte chemoattractant protein [MCP]-1) (Conforti *et al.*, 2008) (Fig 2.2).





Fig. 2.2. Cytokines production network in the tissues (modified from Hopkins, 2003)

(IL=interleukin, TNF=tumour necrosis factor, CSF=colony stimulating factor, PGs=prostaglandins, IFN=interferon).

Other inflammatory mediators include ROS and RNS, eicosanoids such as cyclooxygenase products (prostaglandin E₂ (PGE₂) or lipoxygenase products (leukotrienes (LTB₄) (Nardi *et al.*, 2007) (Fig 2.3), pain provoking mediators (histamine and bradykinin) (Matu and van Staden, 2003), and/ or cationic antimicrobial peptides (CAMP). Another antimicrobial properties of inflammation is disruption of the epithelial lining which limit microbial survival and colonization of the GIT in inflamed intestine due to loss of replication niche






Fig.2.3. Biosynthetic pathways for the eicosanoids (modified from Haeggstrom *et al.*, 2010)

(COX=cyclooxygenase, LOX=lipoxygenase, S=synthase, PG=prostaglandin, LT= leukotriene, TXA=thromboxane, PGI=prostacyclin, cPLA₂=phospholipase A₂)

While inflammation process is beneficial to the body as it removes the insulting cause (Lee *et al.*, 2007a; Pharaoh *et al.*, 2006) the large recruitment and activation of neutrophil and macrophages can induced changes in gut motility, neuronal functionality, and hydroelectrolyte movement with resultant diarrhoea (Gelberg, 2007). Some infectious enteric pathogens elicit inflammatory cascade and mediators to manifest diarrhoea (Guttman and Finlay, 2009).The mechanisms involved in the inflammatory modulated diarrhoea may include several factors listed below and shown in Fig 2.1.

-  Epithelial barrier disruption: Gastrointestinal epithelium barrier provide a physical defence against hostile environment within the intestine lumen (Blikslager, 2010). The intestinal barrier is determined by interactions among several barrier components including the adhesive mucous gel layer, the mucosal immune system and the tight junctions (TJs) (Schenk and Mueller, 2008). The intercellular TJs are the most essential component of the intestinal physical barrier. TJs are multiple protein complexes located around the apical end of the lateral membrane of the epithelial cells. It performs dual functions as a selective/semipermeable paracellular barriers allowing movement of ion, solutes and water through the intestinal epithelium while also preventing the translocation of luminal antigens, microorganisms and their toxins into the mucosa (Groschwitz and Hogan, 2009; Guttman and Finlay, 2009). Disruption of the intestinal TJ barrier by inflammatory cytokines, reactive oxygen species and pathogens (Guttman *et al.*, 2006) impair intestinal TJ function cause an increase in intestinal permeability resulting diarrhoea (Schenk and Mueller, 2008) as shown in Fig. 2.4.




Fig. 2.4. The mechanisms of intestinal epithelial tight junctions as a physical barrier to movement of selected solute materials across the GIT. The intestinal TJs tightly regulate intestinal paracellular permeability. The barrier impairment induced by extracellular stimuli, such as inflammatory cytokines and reactive oxygens, allows the lumina bacterial products and dietary antigens to cross the epithelium and enter circulation (Suzuki and Hara, 2010).

- ✚ **Reduced absorption capacity:** Nutrient-coupled absorption of electrolytes takes place in the brush border microvilli (Dudeja and Ramaswamy, 2006). In an inflamed or infected intestinal tract, the total absorptive surface area is decreased due to brush border shortening resulting in malabsorption (Cotton *et al.*, 2011). Small intestinal malabsorption occurs due to impaired absorption of water, glucose and electrolytes creating an osmotic gradient that draws water into the small intestinal lumen resulting small intestinal distension and rapid peristalsis, consequently diarrhoea (Schulke *et al.*, 2009; Gelberg, 2007; Beavis and Weymouth, 1996).
- ✚ **Chloride ion hypersecretion:** Diarrhoeal agents such as inflammatory mediators, microbial toxins, neurotransmitters and endogenous hormones can activate inappropriate chloride ion (Cl^-) secretion from the colonic crypt epithelial cells. Excessive secretion of chloride ion (Cl^-) from the intestinal crypt is the driving force for many diarrhoea aetiologies. The underlying mechanism is the increase in intracellular levels of cyclic nucleotides (cAMP and cGMP) and/or cytosolic calcium. This process, in turn, drives the secretion of fluid and electrolytes into the intestinal lumen, which may overwhelm the intestinal absorptive mechanism, thereby resulting in secretory diarrhoea with potential effect of severe dehydration (Petri Jr. *et al.*, 2008).
- ✚ **Interference with ability to digest:** Inflammatory response in the intestine may negatively affect the ability of the enterocyte to digest nutritional material. The process causes maldigestion occurs due to a deficiency in various brush border digestive enzymes, especially for carbohydrate and lipids (Schulke *et*

al., 2009). The high level of undigested carbohydrate and lipids are conversion to short chain fatty acids by the colonic microbiota and the amount may exceed colonic capacity for their absorption. Excess short chain fatty acids induced osmotic gradient pulling water and secondarily, ions into the intestinal lumen resulting in osmotic diarrhoea of colonic origin (Field, 2003). Maldigestion of ingested food coupled with osmotic diarrhoea ultimately lead to long-term malnutrition in affected host (Ogoina and Onyemelukwe, 2009).

- ✚ Inflammatory mediators as secretagogue: The release of cytokines such as interleukin-8 (IL-8) and eicosanoids into the gastrointestinal tract activated the macrophage of the immune system. The activated macrophages release inflammatory mediators such as PGE₂, LTE₄, platelet activating factor (PAF), histamine, serotonin, adenosine resulting in cell damage mediated by T lymphocytes or proteases and oxidants generated (H₂O₂, O₂⁻, OH[·], NO) by mast cells. Some of the inflammatory mediators (PGE₂, LTB₄, histamine) also serve as secretagogue causing secretory diarrhoea (Field, 2003).
- ✚ Stimulate enteric nervous system (ENS): Inflammation causes structural changes to the ENS that ranges from axonal damage to neuronal death (Stanzel *et al.*, 2008). The changes include altered neurotransmitters synthesis, storage and release, therefore contributing to the altered intestinal motility during the onset and progression of many GIT disorder (Stanzel *et al.*, 2008) (See section 2.3.3 for more detailed).

2.3.2. Oxidative damage in diarrhoea

Excessive generation of reactive oxygen species (ROS) or reactive nitrogen species (RNS) by the intestinal immunological system as a result of intestinal infection, irritation, inflammation, and depleted endogenous antioxidant defence causes oxidative stress (Granot and Kohen, 2003). This condition has been implicated as one of the causes of diarrhoea (Peluso *et al.*, 2002; Granot and Kohen, 2003).

The pathophysiology of oxidative stress (production) is complex and results from the normal immune response in conditions of disease (infectious and non-infectious), and is initiated by activated mitochondrial of the leukocyte. The free radicals produced are unstable and highly reactive charged function to destroy invading organism (Dwyer *et al.*, 2009). The mechanisms of ROS and RNS production involved an incomplete reduction of oxygen and nitrogen in the electron transfer chain of respiratory process in the mitochondria. In addition, immune reactions during infection or autoimmune responses through inflammation activation of a variety of inflammatory cells, which in turn activate the oxidant-generating enzymes including NADPH oxidase, inducible nitric oxide synthase (iNOS), myeloperoxidase, and eosinophil peroxidase. The ROS generated in the body are superoxide anions, hydroxyl radical, singlet oxygen and hydrogen peroxide (from leukocyte respiratory burst). The RNS included nitric oxide (NO) (produced by inducible nitric oxide synthase (iNOS). Other miscellaneous reactive species are reactive halogen and pseudohalogen species (produced by myeloperoxidase, eosinophil peroxidase,


lactoperoxidase). It is well-established in vitro that free radicals may also be generated via transition metal-mediated oxidation, the so-called Fenton type chemistry, but due to the limited availability of unbound transition metals, these reactions are probably unlikely to play a major role as a source of oxidants in vivo (Chen et al., 2000).

However, since their effect is usually non-specific and aimed at the lipid membrane, the chain reaction initiated by the immune system will destroy the body's macromolecules unless scavenged (terminated). At normal physiological conditions a balance is maintained between amounts of free radicals generate and endogenous antioxidant defence system that scavenged/quenched the radicals preventing their harmful effects. Cellular antioxidant endogenous defence mechanisms are divided into three parts depending on their function:

- Quenching antioxidants: The tissue have inherent antioxidant network capable of donating electrons to oxidants, thus quenching their reactivity under controlled conditions and the derivatives become harmless to cellular macromolecules. The antioxidants however become radicals themselves, but far more stable incapable of inducing cellular damage. The oxidised antioxidants are subsequently recycled to their active reduced state by a number of efficient cellular processes fuelled by energy from NADPH. This recycling process is the main key to the efficiency of the antioxidant network.
- Repairing/removing oxidative damage: This level of antioxidant defence involves the ability to detect and repair or remove oxidised and damaged molecules before it become a threat to normal body.physiological process.
- Encapsulating non-repairable damage: Finally, the body is also equipped with controlled cell suicide or apoptosis, if the extent of the oxidative damage exceeds the capacity of repair and removal.

However, a shift in favour of the radical generation, increase the burden in the body (oxidative stress) which causes tissue injury and subsequently diseases. The proposed mechanisms through which these products induced diarrhoea are presented in Fig 2.5 and discussed below:

- ❖ Lipid peroxidation are primary mechanisms for intestinal cellular malfunction, and can destroy the capacity of membranes to maintain ionic gradients resulting in an aberration in ion transport, particularly affecting potassium efflux and sodium/calcium influx (Dudeja and Ramaswamy, 2006). The production of arachidonic acid metabolites in the lipid peroxidation process can also contribute to intestinal dysfunction including diarrhoea. The ROS and RNS-induced lipid peroxidation process involves three major stages (Catala, 2009): the initiation stage, where the oxidant abstracts hydrogen from polyunsaturated fatty acids of the cell membrane, forming a radical lipid. The propagation stage may involve the rearrangement of the lipid radical to form conjugated dienes and can interact with oxygen to form lipid peroxide radicals.







Fig.2.5 The integrative pathophysiology and mechanism of diarrhoeal disease

Bold (pathogenesis and areas of possible pharmacological intervention in diarrhoea), italic (mechanisms through which diarrhoea manifest)

The peroxide radicals can in turn abstract hydrogen from lipids to produce lipid hydroperoxide and a new radical. Lipid hydroperoxides can be oxidized, via reaction with reduced iron (Fe^{2+}) to lipid alkoxy radical and lipid peroxide, thus continuing the chain reaction of lipid peroxidation. In the final stage, the lipid peroxide radicals in the presence of reduced metals can be degraded to form highly reactive and potent toxic aldehydes such as malondialdehyde (MDA) (Fig 2.6 and 2.7). The chain reaction can be terminated by endogenous antioxidant enzymes and exogenous antioxidant molecules by forming non-reactive substances (Catala, 2009).






Figure 2.6: Lipid peroxidation chain reaction (Valko *et al.*, 2007).

(Equation 1 is generation of superoxide by enzymes such as NAD(P)H oxidase, xanthine oxidase and mitochondria, 2) Superoxide radical is dismutated by the superoxide dismutase (SOD) to hydrogen peroxide, 3 and 4) hydroxyl radical and hydroxyl ion hydrogen peroxide in the presence of transitional metal, 5 and 6 chain reaction to generate more radicals, 7 lipid peroxidation of phospholipids, 8 cyclization and scission of the lipid peroxide radical to generate cytotoxic malonydialdehyde (MDA), hydroxyacrolein (HAC), 4-Hydroxy nonenal (4-HNE) (Fig.2.7).




Fig.2.7. Chemical structures of lipid peroxidation intermediates outlined in Fig 2.6

- ❖ Some of the reactive species such as HOCl and NH₂Cl can also act as secretagogues on their own or can evoke the release of acetylcholine or other neurotransmitters, thus stimulating the enteric nervous system (ENS) to cause increased contractility or motility of intestinal tract (Gaginella *et al.*, 1992). The reactive species also induce gene expression by stimulating signal transduction such as Ca²⁺-signalling and protein phosphorylation.
- ❖ Increased production of inflammatory mediators: The onset of lipid peroxidation process leads to changes in the physiological integrity of the cell membrane. The body responds to the process by the release of pro-inflammatory eicosanoids such as (prostaglandins, prostacyclins and leukotrienes) and pro-inflammatory cytokines (Nardi *et al.*, 2007) such as interleukins (IL-1B, IL-3,IL-6), interferons (IFN), tumor nuclear factor (TNF-α) and platelet-activating factor (PAF) (Conforti *et al.*, 2008; Kunkel *et al.*, 1996).

2.3.3. Enteric nervous system in diarrhoea

The enteric neural network is responsible for the control of propulsive transport and segmental peristalsis in the GIT, as well as secretion and absorption across the intestinal lumen (Wood, 2004; Bohn and Raehal, 2006). While enteric nervous system (ENS) functions independently of the central nervous system (CNS), it is modulated by the parasympathetic and sympathetic autonomic nervous system (Farthing, 2003). As a unit, the ENS is a complicated physiological with autoregulation being mediated by a number of neurotransmitters such as acetylcholine, serotonin, substance P, histamine and endorphin (Farthing, 2002). Diarrhoea can result from the alteration of these systems:

- ❖ Smooth muscle contractility: Many agonists and/or antagonists elicit contractility in GIT smooth muscle (longitudinal or circular) through activation of various receptors located within the muscle (Holzer, 2004). In some cases the activation of the smooth muscle receptors by neurotransmitters and inflammatory mediators include reactive oxygen species causes relaxation (spasmolytic). While in other cases, the process lead to increase in spontaneous or induced contraction (spasmogenic). Ionic channel (Ca^{2+} and Cl^-) are also known to play important roles in smooth muscle contraction (Giorgio *et al.*, 2007). Anion and fluid secretion into the intestine lumen are stimulated through activation of the receptors on enteric secretomotor pathways and epithelial cells, consequently causing secretory diarrhoea.
- ❖ Motility: Intestinal motility dysfunctions include situations in which movement of material along the GIT is repetitive and rapid (diarrhoea) and/or too slow (pseudo-obstruction, slow transit constipation) (Talley, 2006; Giorgio *et al.*, 2007) are controlled by activities of neurotransmitters on the ENS. Pathogenic bacterial overgrowth is common as a result of intestinal hypomotility or low transit time which may lead to mucosal inflammation, increased accumulation and absorption of toxins which are known pathophysiology of diarrhoea. The mechanisms may include impaired digestion as in the deconjugation of bile salts with subsequent fat malabsorption, leading to fatty acid diarrhoea or osmotic effects of malabsorption of sugars resulting in osmotic diarrhoea. Diarrhoea also results from an increase in the gut motility (hypermotility) inducing an accelerated transit of food intake. The net fluid absorption from the food intake is reduced due to less adequate contact time with the GIT epithelial lining for the absorption of fluids before excretion.

2.3.4. Cystic fibrosis transmembrane conductance regulator (CFTR) regulation

Cystic fibrosis transmembrane conductance regulator (CFTR) is a cyclic adenylyate monophosphate (cAMP)-activated Cl^- channel expressed in epithelial cells in the intestine and other fluid-transporting tissues (Thiagarajah and Verkman, 2003). Diarrhoeal pathogens and their toxins can induce secretory diarrhoea by simultaneously stimulation of the active secretion of Cl^- and inhibition of Na^+ absorption across the apical membrane of enterocyte with resulting massive fluid and electrolyte loss into GIT (Schuier *et al.*, 2005). The cellular signalling mechanisms include an increase in cellular cAMP and cyclic guanylate monophosphate (cGMP), which may result in activation of the CFTR Cl^- channel. Pharmacological blocking of CFTR with drugs such as glibenclamide and CFTR_{inh}-172 inhibits salt and water loss in diarrhoea (Schuier *et al.*, 2005).

2.4. Specific Agents of Diarrhoea

2.4.1. Bacterial causes of diarrhoea

2.4.1.1. *Escherichia coli*

E. coli is a gram-negative rod shaped bacteria that shares a symbiotic relationship with animal host as part of normal digestive intestinal flora. Under certain define conditions these organisms or pathogenic strains of these organisms are known to induce diarrhoea (Clarke, 2001; Le Bouguenec, 2005). There are six main types of

pathogenic *E. coli* associated with diarrhoea, namely enterotoxigenic *E. coli* (ETEC), enteroinvasive *E. coli* (EIEC), enteropathogenic *E. coli* (EPEC), enterohaemorrhagic *E. coli* (EHEC), enteroaggregative *E. coli* (EAEC) and diffusively adherent *E. coli* (DAEC) (Clarke, 2001). While the exact process by which each type of these *E. coli* induces diarrhoea symptoms varies significantly, the basic pathophysiology involves their inherent ability to adhere to epithelial cells and colonize the host tissues (Le Bouguenec, 2005). The characteristics and mode of actions of each type of the pathological strains in diarrhoea diseases are listed in Table 2.1. Infections from some of the strains of *E. coli* are self-limiting and can resolve without pharmacological intervention. However, symptomatic, supportive and antibiotic, or a combination of the therapies may be beneficial in the chemotherapeutic management of some cases involving ETEC, EIEC and EPEC infection (Elsinghorst, 2002). The use of antibiotics recommended, antimicrobial chemotherapeutic agents such as tetracycline, doxycycline, and ciprofloxacin may be used (Casburn-Jones and Farthing, 2004; Elsinghorst, 2002) in infectious diarrhoea.

Table 2.1: The mechanism of actions and symptoms of enteric pathogenic *E. coli* (Thapar and Sanderson, 2004; Clarke, 2001).

Strain	Mechanism of action	Symptoms	Treatments
Enterotoxigenic <i>E. coli</i> (ETEC)	Colonization of the small bowel mucosa, followed by elaboration of heat-labile (LT) and heat stable (ST) enterotoxins. The ST enterotoxins are classified as STa and STb. The binding of STa to guanylate cyclase C receptor results in increased intracellular cyclic guanylate monophosphate (cGMP) level. The resultant effect is the stimulation of chloride secretion or inhibition of sodium chloride absorption causing intestinal fluid secretion. LT enterotoxins consist of two serotypes (LT-I and LT-II). LT activate adenylate cyclase causing intracellular increase in cyclic adenosine monophosphate (cAMP) levels resulting in decrease sodium absorption by villous cells and subsequent active chloride secretion by crypt cells thus leading to osmotic diarrhoea.	Watery diarrhoea ranging from mild, self-limiting disease to severe purging.	Supportive therapy with antibiotic in cases of severe infection.
Enteroinvasive <i>E. coli</i> (EIEC)	Invasion of the epithelium and mucosal destruction eliciting inflammatory response accompanied by necrosis and ulceration of the large bowel with resultant release of blood and mucosa into the stool.	Bloody, mucoid diarrhoea and dysentery	Antibiotic in cases of bloody diarrhoea
Enterohaemorrhagic <i>E. coli</i> (EHEC)	Adhesion followed by liberation of a potent toxin which is cytotoxic to Vero cells (referred to as shiga-like cytotoxin I and II). Other mechanism attributed to the EHEC virulence includes adhesin	Bloody diarrhoea, fever, vomiting, haemorrhagic colitis, haemolytic uremic syndrome (HUS), acute renal failure, haemolytic anaemia,	Symptomatic therapy
Enteropathogenic <i>E. coli</i> (EPEC)	Adherence of the bacterium to the gut epithelium causing attachment and effacement lesion on intestinal epithelial cells, alteration of intracellular calcium and cytoskeleton.	Self-limiting watery diarrhoea with fever and vomiting	Antibiotic in severe cases
Enterotoxigenic <i>E. coli</i> (EAEC)	Aggregating pattern of adherence to intestinal mucosa produces enterotoxigenic heat-stable (EAEC) enterotoxins causing cellular damage and function similar to, but distinct from ST enterotoxins.	Watery mucoid diarrhoea	Antibiotic in severe cases
Diffusively adherent <i>E. coli</i> (DAEC)	Elaboration of α -haemolysin and cytotoxic necrotising factor 1 (CNF-1).	Watery diarrhoea	Supportive therapy

2.4.1.2. *Staphylococcus aureus*

S. aureus is a gram-positive coccus present in normal intestinal and skin flora of human and homeothermic animal. Under define conditions, the pathogenic strains produces heat stable staphylococcal enterotoxins (SETs) and toxic shock syndrome toxins (TSST-1) (de Oliveira, 2010) both of which are known to induce diarrhoea. Toxicity from SET results from the consumption of the preformed heat-stable enterotoxins (α -haemolysin) in contaminated food. Upon ingestion of the food contaminated with the SETs, the toxins results in signs of nausea, vomiting, fluid accumulation in ileal loops, and diarrhoea associated with fever (Rosengren *et al*, 2010; Perez-

Bosque and Moreto, 2009). The main sources of *S. aureus* toxin contaminants are raw material and food processing unit such as human handling, water and environment (Linscott, 2011).. Serotonin receptor antagonists have been reported to ameliorate the vomiting, diarrhoea and prostration induced by SETs . The mechanism behind toxicity results from the activation of autonomous nervous system with resultant hyperperistalsis as well as activation of central pathways which control vomiting and diarrhoea (Podewils *et al*, 2004).

In contrast, TSST-1 is characterized by sudden onset of fever, vomiting, diarrhoea, erythematous rash with skin peels, hypotensive shock, impairment of renal and hepatic functions, and sometime death. Toxicity results via the production of pro-inflammatory cytokines and chemokines. Toxicity is usually exacerbated by further interaction between the activated immune system and inflammatory mediators (Krakauer *et al.*, 2001).

2.4.1.3. *Campylobacter jejuni*

C. jejuni is an invasive Gram-negative, spiral-shaped rod bacterium present in the GIT of mammals, birds and primates (Lengsfeld *et al.*, 2007). The major source of *Campylobacter* infection in mammals is from poultry and poultry products (Podewils *et al*, 2004). The clinical signs of *campylobacter* infections include pyrexia, abdominal pains, watery diarrhoea and dysentery (Podewils *et al*, 2004). The characteristic mechanisms *Campylobacter* infection involves invasion and translocation of the epithelium with a concomitant induction of inflammation (Hu *et al.*, 2008).

2.4.1.4. *Shigella* spp

Shigella (*Shigella flexneri*, *Shigella dysenteriae*, *Shigella sonnei* and *Shigella boydii*) is a Gram-negative rod, non motile and facultative anaerobic bacterium that invades the colon with resulting inflammation and diarrhoea (Podewils *et al*, 2004). *Shigella flexneri* is responsible for dysenteric symptoms and persistent illness while *Shigella dysenteriae* type-1 produces Shiga-toxin like EHEC causes bloody diarrhoea (Podewils *et al*, 2004). *Shigella sonnei* causes bacterial gastroenteritis and bacillary dysentery and *Shigella boydii* causes fever, chills, abdominal pain and diarrhoea.

2.4.1.5. *Vibrio cholerae*

V. cholerae is a motile, facultative anaerobic Gram-negative rod associated with potentially fatal diarrhoea (Granum, 2006) that results from the ingestion of the cholera enterotoxins (CT) from contaminated water and seafood (Podewils *et al*, 2004). Watery, colourless mucous- flecked stool and vomiting are the main clinical signs associated with cholera which in severe cases can result in a life-threatening fluid and electrolyte imbalance (Podewils *et al*, 2004). Pathophysiologically, toxicity results from the CT induction of intestinal hypersecretion through the activation of the mucosal epithelium cAMP-adenylate cyclase system in the mucosal epithelium (Casburn-Jones and Farthing, 2004).

In addition, it has been speculated that ROS/RNS production in *V. cholerae* infection could also contribute to intestinal damage through lipid peroxidation of the cellular and mitochondrial membrane thereby further increasing membrane permeability and fluid loss (Gorowara *et al.*, 1998). Other species of *Vibrio* such as *V. parahaemolyticus* and *V. vulnificus* also caused watery diarrhoea, abdominal cramps, nausea, vomiting. These organisms infect host from raw or undercooked seafood or cooked seafood contaminated with seawater (Linscott, 2011).

2.4.1.6. *Bacillus cereus*

B. cereus is a sporulating bacterium that causes both food poisoning and non-gastrointestinal infection (Al-khatib *et al.*, 2007; Ramarao and Lereclus, 2006). In food poisoning, two main types of diseases namely diarrhoeal and emetic food poisoning are common. The diarrhoeal type of *B. cereus* food poisoning is caused by enterotoxins such as haemolysin BL (HBL), non-haemolytic enterotoxin (NHE) and cytotoxin K (CytK) (Lund *et al.*, 2000).with clinical signs of abdominal pain with diarrhea. Causes of the diarrhoeal forms always occurs from accidental contamination of food like meat, vegetables, pasta, deserts cakes, sauces and milk (Linscott, 2011).

In constrast the emetic form is induced by a small preformed heat and acid stable cyclic peptide (cereulide) (Agata *et al.*, 1995; Ehling-Schulz *et al.*, 2004) with clinical symptoms of sudden onset of nausea and vomiting, with or without diarrhoea (Linscott, 2011). The major sources include cooked foods, like meat or fried rice that have not been properly refrigerated. While the other species of *Bacillus* such as *B. subtilis*, *B. licheniformis*, *B. pumilus* and *B. megaterium* are usually considered relatively safe, but they can also produce enterotoxins and emetic toxins involved in foodborne illness (From *et al*, 2007)

2.4.1.7. *Yersinia* species

Yersinia species are Gram-negative facultative anaerobic nonsporing rods or coccobacilli bacteria belonging to the Enterobacteriaceae family. Three human pathogenic species namely: *Y. pestis*, *Y. enterocolitica*, and *Y. pseudotuberculosis* are recognized (Fallman and Gustavsson, 2005). *Y. pestis* is the causative agent of bubonic plague characterized with the onset of fever, chills, headache, and weakness, followed by swelling and tenderness of lymph nodes while *Y. enterocolitica* and *Y. pseudotuberculosis* cause an enteric infection in humans called yersiniosis with clinical signs such as diarrhoea, vomiting, fever and abdominal pain (may mimic appendicitis) following ingestion from undercooked pork, unpasteurized milk, tofu, contaminated water, chitterlings (Linscott, 2011, Damme *et al.*, 2010).

2.4.1.8. *Listeria monocytogenes*

L. monocytogenes is a Gram-positive bacterium which causes life-threatening invasive diseases referred to listeriosis in human and animals (Chaturongakul *et al.*, 2008; Todd and Notermans, 2011). Upon ingestion of the bacteria from contaminated foods such as unpasteurized milk, soft, cheese made with unpasteurized milk

Linscott, 2011), the organism may, colonize the intestinal tract with resultant diarrhoea (Chaturongakul *et al.*, 2008).

2.4.1.9. *Clostridium* spp

C. difficile is an anaerobic, spore-forming, Gram-positive bacillus widely distributed in the environment and present in the colon flora of less than 3% of healthy adults (Beaugerie *et al.*, 2003). *C. difficile* causes a spectrum of diseases ranging from benign diarrhoea to fatal colitis and most often as a consequent of antibiotics treatment. Most antibiotics predispose *C. difficile* overgrowth leading to the production and accumulation of and diarrhoea are Toxins A (enterotoxin) and B (cytotoxin) in the intestine. Both toxins A and B inactivate intracellular Rho-proteins by glycosylation, leading to desorption of the cytoskeleton, production of inflammatory cytokines and damage to tight junctions. The most commonly antibiotics associated with *C. difficile* overgrowth include cephalosporins, clindamycin and broad-spectrum penicillins (Wistrom *et al.*, 2001).

In contrast, *C. perfringens* is an important food poisoning bacterium with clinical sign as diarrhoea, abdominal cramping and nausea. The main sources of infection include contaminated meat, poultry, gravy and inadequately reheated food (Linscott, 2011). *C. botulinum* may also play a role in diarrhoeal diseases when the preformed botulinum toxin is consumed from improperly canned foods, herb-infused oils, baked and potatoes in aluminium foil. Symptoms of infection include abdominal cramping, nausea, vomiting, diarrhea, double vision, long term nerve damage and possible even death from paralysis (Linscott, 2011).

2.4.1.10. *Salmonella typhimurium*

S. typhimurium is a bacterium that may be associated with mild gastroenteritis to enteric (typhoid) fever, bacteraemia and septicaemia commonly referred to as salmonellosis (Mastroeni and Maskell, 2006). The clinical signs of salmonellosis include diarrhoea, fever and abdominal cramps. In people with typhoid fever, *Salmonella* spreads systemically from the gut to blood stream and other parts of the body resulting in mortality if not treated adequately with antibiotics (Castillo *et al.*, 2011). The virulence of the *Salmonella* bacterium differs among the different animal species depending on *Salmonella* serovar involved, strain, infective dose, host animal species, age and immune status of the host (Castillo *et al.*, 2011). The pathogenesis of *Salmonella* involves adhesion/invasion to specific intestinal epithelial cells, mainly in the ileum (Guttman and Finlay, 2009).

2.4.1.11. *Enterococcus faecalis*

E. faecalis is a gram-positive bacterium that survives symbiotically in the human or animal's intestinal tract. However, under conditions such as the disruption delicate host-commensal relationship following antibiotic use, abdominal surgery or changes in host immunity, the enterococci becomes invaders of the intestinal wall (Butler, 2006) through the production of adhesin, aggregating and binding substances (Butler, 2006). *E. faecalis* is known to produce superoxide (O_2^-) that can results in hydroxyl radical formation which contributes to oxidative

stress in the intestine and membrane lipid peroxidation (Huycke and Moore, 2002; Sun *et al*, 2010; Peluso *et al.*, 2002; Granot and Kohen, 2003) (Fig 2.5).

2.5. Fungal induced diarrhoea symptoms

2.5.1. *Candida albicans*

C. albicans is a yeast fungus and exist as a member of normal flora in the GIT and mucocutaneous membrane (Forbes *et al.*, 2001). However, following the use of antibiotic therapy that result in sterilization of the GIT flora, *C. albicans* can overgrowth to take the place of removed organisms with end result of diarrhoeal symptoms(Henry-Stanley *et al.*, 2003). Other predisposing factors include altered intestinal permeability and diminished host immunity response. It has been postulated that this organism produces virulence factors which increases fungal adherence to host cells, fungal secretion of proteolytic enzymes and fungal morphological switching (ability to change and grow in several distinct morphological forms: yeast, hyphae, and pseudohyphae, according to environmental conditions) (Henry-Stanley *et al.*, 2003). Clinical signs associated with enteric candidiasis are abdominal pain, cramping, rectal irritation and absence of nausea, vomiting, bloody and mucus stool, and pyrexia (Levine *et al.*, 1995).

2.6. Viral induced diarrhoea

2.6.1. Rotavirus

Rotavirus is a major cause of severe diarrhoea and account for 30% and 80% of all cases of acute gastroenteritis (Savi *et al*, 2010). The diarrhoeal mechanism of the organism includes the production of enterotoxin NSP4 which induced Na^+ -glucose dependent malabsorption and destruction of enterocytes (cytotoxicity). The toxin also has a direct effect on the intestinal barriers by blocking TJs formation with resultant diarrhoea through a 'leak flux' mechanism in which water is secreted into the lumen of the intestine (Dickman *et al.*, 2000).

2.6.2. Norovirus

Norovirus is considered one of the major global causes of gastroenteritis (Mattison, 2010) The diseases is opportunistic and is usually transmitted through faecal contamination of food, water or contact with an infected host following poor hygiene. The organism has the ability to infect small intestine and induce intestinal TJ dysfunction, malabsorption through villus surface area reduction and an increased number of cytotoxic intraepithelial lymphocytes (Troeger *et al.*, 2009). Clinical signs associated with infection are nausea, vomiting, diarrhoea and abdominal pains (Koopmans, 2008).

2.6.3. Hepatitis A virus

Hepatitis A is a small, non-enveloped spherical with cubic symmetry, thermostable and acid resistant virus. While the primary target organ of infection is the liver, the resultant hepatitis (Koff, 1998) result in clinical signs dark

urine, jaundice, malaise, weakness, fever, anorexia, nausea and vomiting, abdominal pains, and diarrhoea (Koff, 1992). Sources of infection usually result from contaminated water on raw produce, food, and shellfish or exposure to the water itself (Linscott, 2011).

2.6.4. Human immunodeficiency virus (HIV)

Chronic diarrhoea is one of the complications associated with HIV infection and acquired immune deficiency syndrome (AIDS) due to multiple enteric opportunistic microbes (DuPont and Marshall, 1995). While HIV is important in secondary enteric diseases as a result of immune suppression (CD4+ T-lymphocytes destruction), the virus can result in diarrhoea directly by altering the mucosa structural arrangement viz referred to as HIV-enteropathy (Epple *et al.*, 2009). The diarrhoea resulting from HIV appears to be caused by the releases of cytokines from the infected immune cells (Schmitz *et al.*, 2002).

2.7. Protozoa induced diarrhoea

2.7.1. *Giardia intestinalis*

G. intestinalis (syn. *Giardia doudevalis*, *Giardia lamblia*) is a flagellate protozoa parasite of the upper small intestine that exists as a motile trophozoite (Cotton *et al.*, 2011). The organism colonizes the small intestinal lumen and induces non-inflammatory and malabsorptive diarrhoea (Schulzke *et al.*, 2009). The pathophysiology of Giardiasis involves Na⁺-dependent D-glucose absorption impairment, active electrogenic anion secretion activation, mucosal inflammation and leak flux (Buret, 2007; Troeger *et al.*, 2007). Clinical signs of *Giardia* infection include bloating, steatorrhea and nausea. Chronic infection cause weight loss, growth retardation and development in young children

2.7.2. *Entamoeba histolytica*

E. histolytica is a protozoa parasite which infects the large intestine with resultant intestinal dysfunction characterized by invasive illness and severe dehydration commonly referred to as amoebiasis (Ralston and Petri, 2011). The pathophysiology of amoebiasis involves villus structural destruction, increased epithelial permeability and alteration of TJs (Lauwaet *et al.*, 2004). The organism also causes invasive disease such as colitis and abscess through massive host tissue destruction. The clinical signs are usually similar to *S.dysenteriae* or enteroinvasive *E. coli* with blood and pus contaminated stool. Other related infectious species include *E. dispar* and *E. moshkovskii* (Ralston and Petri, 2011).

2.7.3. *Cryptosporidium parvum*

C. parvum is an intracellular protozoa parasite that infects epithelia causing cryptosporidiosis (Kenny and Kelly, 2009) which manifest clinically as profuse watery diarrhea, containing mucus, but rarely blood or leukocytes

(O'Hara and Chen, 2011). Some other clinical signs of cryptosporidiosis include nausea, vomiting, cramp-like abdominal pain and mild fever (Linscott, 2011). The period and severity of clinical symptoms of intestinal cryptosporidiosis depends on the immune status of the infected individual (Linscott, 2011). Cryptosporidiosis in the healthy individuals is usually a self-limiting illness with approximate duration of 9-15 days while in immunocompromised patient the infection is severe and life-threatening (O'Hara and Chen, 2011).

2.7.4. *Cyclospora cayetanensis*

C. cayetanensis is a protozoan parasite which invades the epithelial cells of the small intestine upon ingestion of sporulated oocysts in contaminated food or water (Chacin-Bonilla, 2010; Manfield and Gajadhar, 2004). Pathomechanisms of *C. cayetanensis* infection are intestinal inflammation associated with pathological lesions such as villus blunting, and malabsorption. The clinical signs of the infection include watery diarrhoea, loss of appetite, weight loss, abdominal bloating and cramping, increased flatulence, nausea, fatigue, and low-grade fever (Linscott, 2011).

2.8. Parasitic induced diarrhoea

2.8.1. *Trichinella spiralis*

T. spiralis is a food-borne zoonotic parasite responsible an enteral phase and a muscular phase (Cui *et al.*, 2011). The adult worms live in the duodenal and jejunal mucosa of flesh-eating animals and humans, while the larvae live in skeletal muscles of the same hosts (Cui *et al.*, 2011). The source of infection is raw or undercooked contaminated meat (pork, bear, walrus and moose), cross-contaminated ground beef and lamb (Linscott, 2011). The clinical intestinal symptoms manifest one or 2 days after ingesting the contaminated meat due to the matured adults penetrating the intestinal mucosa, resulting in nausea, abdominal pain, vomiting, and diarrhoea (Linscott, 2011). *T. Spiralis* induced changes in intestinal function by hypersensitivity mechanism resulting in an increased intestinal chloride and fluid secretion (Cui *et al.*, 2011).

2.9. Immune disordered induced diarrhoea

2.9.1. Compromised immune system

The main function of the immune system is to protect against disease through the recognition and removal of pathogens or their sequelae (Gertsch *et al.*, 2010). To fulfil this role, the body make use of an innate immune system that defends it in a non-specific manner via molecular interaction and inflammation (Gertsch *et al.*, 2010) and an adaptive system comprised of specialized effector cells (T and B cells) which not only recognize antigens but play a role in their active removal (Gertsch *et al.*, 2010). In the GIT, the important protective system which prevent the normal flora from becoming pathogenic are gut-associated lymphoids tissue (GALT), epithelial-derived antimicrobial peptides (AMPs) (such as defensins, cathelicidin and lysozyme present in the secretion which constantly washes the mucosal surfaces), the mesenteric lymph nodes, the liver's Kupfer cells, mast cells, within the intestinal walls and the reticuloendothelial system of the intestinal walls. In diseases characterized by immune suppression such as HIV, the immune system is destroyed which makes a person more susceptible to

infectious diseases (Gertsch *et al.*, 2010). One of the clinical signs that may result is diarrhoea as the above-mentioned micro-organisms colonize the compromised GIT.

2.9.2. Hyperactive immune system

The normal response of the immune system during conditions of antigen stimulation is generally an inflammatory response with the release of numerous inflammatory mediators (interferon- γ , tumour necrosis factors (TNF- α), interleukin-6 (IL-6), and IL-1 β) which in conjunction plays a role in the removal of the causative agents (Sprague and Khalil, 2009). On the removal of inciting cause, the inflammatory response is usually terminated. However, on occasion when the body failed to terminate the inflammatory response, the inflammatory agents can be equally as destructive to the host's own tissues. The latter usually result from the damage of the epithelial mucosa cells, from the generated ROS/RNS and subsequent lipid peroxidation. With the destruction of the epithelial cells, the body loses absorptive capacity with resultant diarrhoea. The produced cytokines also has the ability to directly increase intestinal mucosa permeability and fluid loss. Inflammatory bowel disease is one of the diseases caused by up-regulated immune system (Gertsch *et al.*, 2010).

2.10. Antibiotic therapy induced diarrhoea

Diarrhoea develops in some patient following antibiotic chemotherapy with one of the following mechanisms:

- Antibiotic toxicity: Some antibiotic (levofloxacin, azithromycin and amoxicillin-clavulanate) may have a direct negative effect on the GIT with resultant poor absorption or enteropathy characterized by infiltration of the lamina propria by macrophage (Dobbins, 1968, Hartmut, 2010). In addition, antibiotic such as erythromycin has prokinetic action on the GIT, mediated through motilin receptor stimulating potential (Catnach and Fairclough, 1992, Annese *et al.*, 1992).
- Alteration of digestive function: The removal of some commensal organisms by antibiotic (Hofmann, 1977) could result in decreased carbohydrate digestion (Saunders and Wiggins, 1981) which leads to accumulation of these osmotically active substances in the intestinal lumen. The net result is the accumulation of electrolytes and water from osmotic pull in the intestinal lumen which is evidence as diarrhoea (Beaugerie and Pettit, 2004).
- Overgrowth of pathogenic microorganisms: The use of antibiotic may result in the removal of beneficial GIT flora. As a result of the disruption of this balance ecosystem, various pathogenic organisms (as listed above) can overgrow and colonise the GIT (Beaugerie and Pettit, 2004). The predisposing antibiotic for various diarrhoeal pathogens include cephalosporins, clindamycin and broad-spectrum penicillins for *C. difficile* (Wistrom, 2001; Stoddart and Wilcox, 2002); β -lactams or pristinamycin for *Klebsiella oxytoca* (Wu *et al.*, 1999) or tetracycline for *S. aureus*. *Salmonella* and *Candida* overgrowth can be a consequence of wide range of antibiotic with broad base actions (Danna, 1991; Gupta and Ehrinpreis, 1990).




Fig. 2.8. Mechanism of antibiotic induced diarrhoea

2.11. Diabetic complications induced diarrhoea

Gastrointestinal disorders manifesting as diarrhoea (watery stool) or constipation (dry and hard stool) is a common symptom in the diabetic patient (Gould and Sellin, 2009) with a prevalence of approximately 12.5 to 32.4% and 60% respectively. In addition the oral hypoglycaemia medications used for the management of diabetes viz metformin, acarbose, miglitol (Forgacs and Patel, 2011), exenatide and orlistat (Gould and Sellin, 2009) may also induced diarrhoea as side effect while the recommended dietary material such as the non-digestible sweeteners (sorbitol, mannitol and D-xylose) induce an osmotic diarrhoea (Forgacs and Patel, 2011). Bacterial overgrowth may also result in diabetic patient from decreased functioning of the immune system as described above (Forgacs and Patel, 2011).

2.12. Food allergy induced diarrhoea

Diarrhoea is one of the clinical manifestations of food allergy (Wang *et al.*, 2010). The mechanisms of action include active ion secretion, altered epithelial barrier function (Groschwitz and Hogan, 2009), and mucosal damage resulting in malabsorption and osmotic diarrhoea (Wang *et al.*, 2010). The initiation of food induced intestinal allergy is regulated by numerous inflammatory cells and mediators, including mast cells and T_H2 cytokines (IL-4, IL-5, and IL-13) (Wang *et al.*, 2010). The release of neurotransmitters (serotonin, histamine) and inflammatory mediator (COX-2 and LOX) by mast cell induced ion secretion in the presence of allergen (Schenk and Mueller, 2008). These neurotransmitters and inflammatory mediators also stimulate intestinal contractions (altered intestinal motility) which act synergistically with ion secretion to cause diarrhoea.

2.13. Potential mechanisms in the control of diarrhoea

2.13.1. Oxidative damage and antioxidants in diarrhoeal management

Several endogenous strategies are available in human and animal body to combat oxidative damage. These provide ways for normal oxidative metabolism to occur in the body without damaging the cells and also allow for normal ROS/RNS-mediated cellular response such as phagocytosis and intracellular signalling (Valko *et al.*, 2007). Therefore, the possibility exists that returning the animal to a more neutral oxidative balance, may promote repair of damaged membranes (Nose, 2000). As a result antioxidants and/or radical scavengers may be

beneficial in the attenuation of diarrhoea. The best known antioxidants as treatment are selenium, vitamin E, vitamin C and the proanthocyanidins in red wine and resverastrol in commercial grape seed extract.

2.13.2. Inflammation and anti-inflammatory agents in diarrhoea management

As a result of the negative impact the inflammatory cascade can have on the functionality of the GIT, modulation of these processes through the use of drugs may be of benefit. Possible mechanisms include attenuation of inflammatory process through the use of anti-inflammatory, antioxidative and radical scavenging mechanisms. Potential target include drugs with cyclooxygenase and lipoxygenase enzyme inhibitory activity. The drugs that are used commonly for this are non-steroidal anti-inflammatory drugs (NSAIDs) like indomethacin, aspirin, ibuprofen, diclofenac and coxibs.

2.13.3. Enteric nervous system in diarrhoea symptoms and treatment

The ENS is an important target for pharmacological intervention in diarrhoea through the use of agonists and antagonists that target these ENS endogenous receptors. Numerous pharmaceutical agents are currently available for alleviating many of the clinical signs of diarrhoea. The effects and possible sites of pharmacological intervention against the activities of neurotransmitters in diarrhoeal symptoms are presented in Table 2.2.

Table 2.2: Neurotransmitters and modulators of ENS causing intestinal secretion in diarrhoea

Neurotransmitters	Effects on GIT	Receptors	Potential pharmacological intervention	modulator
Acetylcholine	Main endogenous excitatory neurotransmitter in the GIT.	Nicotinic and muscarinic receptor subtypes M ₁ and M ₃ .	Non selective nicotinic acetylcholine receptor antagonist or specific muscarinic acetylcholine receptor antagonist	atropine
Serotonin	Modulate muscular contraction and relaxation, intestinal fluid secretion and enhanced colonic transit.	5-HT ₃ , 5-HT ₄	5-HT ₃ receptor antagonist (diarrhoea) and 5-HT ₄ receptor agonist (constipation)	Metoclopramide, granisetron, ondansetron, tropisetron, palonsetron, alosetron, cisapride
Substance P	Transmitter of enteric neurones and extrinsic afferent fibre, control of GI motility, secretion, vascular permeability, immune function and pain sensitivity	NK ₁ , NK ₂ and NK ₃	NK ₁ and NK ₂ receptor antagonist	
Histamine	Modulation of GIT motility, enhancement of gastric acid secretion, increases mucosal Cl ⁻ ion secretion, and	H ₁ , H ₂ , H ₃	H ₁ , H ₂ antagonist	Cimetidine and ranitidine

	modulator of immune functions.			
Opioid peptide	decrease motility, increase transit time, increase fluid absorption from the intestine	Mu (μ), delta (δ) and kappa (κ)	μ , δ agonist	loperamide racecadotril, diphenoxylate
Nitric oxide	ENS neurotransmitter, pro-absorptive and secretory agent		iNOS inhibitor, NO modulator	N ^G -nitro-L-arginine methyl ester (L-NAME)
Dopamine	Prokinetic and antiemetic, enhance antral contractility and inhibit fundus receptive relaxation	D ₂	antagonist	Domperidone
Motilin	induced antral contractility			Erythromycin, motilides, motilactides

2.14. Plants as potential source of therapeutic agents in alleviating diarrhoeal symptoms

Due to the widespread occurrence of diarrhoea as a disease together with the prevalence coinciding with human development, plants and fungi have featured widely in the management of the disease. Their use has become so common in human and veterinary medicines that a number of compounds considered to be allopathic are of natural origin. A non exhaustive list includes:

- Antioxidant: The natural vitamins and red pigments present in plant.
- Anti-inflammatory: Salicylic acid from willow bark.
- ENS modifiers: Atropine from *Atropa belladonna*, tincture of opium from *Papaver somniferum*.
- Antibiotic: By definition all antibiotics are natural product produced by fungi. Almost all the available classes of antimicrobials are of fungal origin.

Alongside these naturally derived allopathic medicines, medicinal plant have been widely use in alleviating diarrhoeal symptoms in humans and animals (Brijesh *et al.*, 2006; Gutierrez *et al.*, 2007). Numerous species of these plants have been screened and validated for their use in treatment of diarrhoea (Gutierrez *et al.*, 2007).

2.14.1. Anti-infectious mechanisms of plant secondary metabolites against diarrhoeal pathogens

- Antimicrobial: Many plant metabolites are known to exhibit some level of toxicity toward microorganisms. Numerous mechanisms of actions have been hypothesized to explain their antimicrobial activity such as microbial enzymes inhibition, deprivation of essential growth substrates, cell membrane disruption (Cowan, 1999) or direct interference with metabolic pathways.
- Antiadhesion: Adhesion of some enteric pathogen to the mucosa epithelium of the host cells is the first important step in intestinal infections that may lead to the development of diseases (Ofek and Sharon, 1990). Application of antiadhesives chemotherapy can be effective only against microorganisms that

depend on the surface contact with host cells as prerequisite for survival, multiplication and virulence (Lengsfeld *et al.*, 2007).

- Antitoxin: Since enteric pathogens may induce diarrhoea through the production of toxin (endotoxin or cytotoxin) the neutralization with plant antidiarrhoeal compounds may be beneficial in the management of diarrhoea. Activated charcoals processed from plants are also used as toxin binders. Other binder includes pectin obtained from apples.
- Immunomodulatory: With immune suppression being a pre-disposing, drugs or medicinal plant preparations with immune stimulating activities may help in attenuating many infectious diseases.

2.14.2. Antioxidative mechanisms of plant phytochemical as potential antidiarrhoeal agents:

- Free radical scavenging: Many plant preparations and phytochemicals have strong antiradical activities which can antagonize the deleterious action of free radical. The mechanism may be electron transfer or hydrogen donating to stabilize the free radical molecules generating radicals that are relatively stable due to delocalization resulting from resonance and unavailability of site for further attack by molecular oxygen (Mello *et al.*, 2005).
- Complexation of catalytic metallic ion: Metallic ions such as ferrous ion (Fe^{2+}), cuprous ion (Cu^{2+}), Manganese ion (Mg^{2+}) and Zinc ion (Zn^{2+}) can also generate free radicals (Kane, 1996). Many plant molecules moderate oxidation activity by complexing with the free transition metal thereby inactivating their capacity to catalyze oxidative process.
- Pro-oxidation enzymatic pathways: With the generation of oxidant being enzyme driven, the antioxidant activities of plant phytochemical may be able to inhibit these enzymatic pathways or inactivation of the enzyme.
- Lipid peroxidation inhibition: Scavenging of free radicals is one of the major antioxidation mechanisms to inhibit the chain reaction of lipid peroxidation and reduction of the deleterious effect of the cytotoxic products.
- Inhibition of nitric oxide (NO): NO generated by inducible NOS (iNOS) can act synergistically with other inflammatory mediators in the development of diarrhoea. The inhibition or down-regulation of iNOS expression may be beneficial to reduce the inflammatory response.
- Immune system optimization: Over expression of immune system may cause damage to surrounding tissues and consequently results in a host of diseases and illness including diarrhoea. Many medicinal plants and phytochemical compounds protect against oxidative stress due to immunomodulatory activity (Wang *et al.*, 2002).

2.14.3. Anti-inflammatory mechanisms of plant phytochemical in diarrhoea management:

- Cyclooxygenase inhibition: Compounds with COX enzyme inhibitory potential could be use as anti-inflammatory agents. Some plant secondary metabolites such as alkaloids, phenols, , terpenoids and their derivatives have potential to inhibit the formation of pro-inflammatory signalling molecules such as prostaglandin (Polya, 2003)
- Lipoxygenase inhibition: Lipoxygenase metabolites are critical mediators of inflammation and thus important in the pathogenesis of abdominal distress and diarrhoea associated with intestinal inflammation. Plant phytochemicals with lipoxygenase inhibitory potential are candidate for anti-inflammation and the resulting diarrhoea.
- Modulation of cytokines activity: A number of diarrhoea pathogenesis causes severe intestinal inflammatory with hypersecretion of pro-inflammatory cytokines (MacNaughton, 2006). Inhibition of the pro-inflammatory cytokine mediator's can remove their negative activities associated with gastrointestinal disorders including diarrhoea.

2.14.4. Antidiarrhoeal mechanisms of plant phytochemicals

- ✓ Antispasmodic: Spasmolytic agents are used in the treatment hypermotility of the GIT while prokinetic agents are used in treatment of hypomotility (Gilani, 2005). Many phytochemicals demonstrated various range of spasmolytic or antispasmodic activities against spontaneous or agonist induced contraction on isolated parts of the GIT.
- ✓ Antisecretory: Microbial enterotoxins cause diarrhoea by disturbing the balance between intestinal absorption and secretion in favour of the later. Therefore, inhibition of the intestinal secretion is one therapeutic model for treating diarrhoea (Velazquez *et al.*, 2009).

2.15.0. Classification of phytochemicals with antidiarrhoea potential

2.15.1. Terpenoids

Terpenoids are the most structurally diverse groups of natural products formed by fusion of isoprene monomers in plants. This class of plant secondary metabolites are grouped according to the number isoprene units or numbers of carbon in their skeletal structure (Zwenger and Basu, 2008). The group include monoterpenes which contain two units of isoprene with C₁₀ and are present as essential or volatile components of herbs, spices and flowers. Sesquiterpenes are derivatives of three isoprene units containing 15 carbon atoms in their structures and are present in essential oil. This group of compounds acts as phytoalexins, antimicrobial and antifeedant in plants. Diterpenes contains 20 carbon atoms derived from four units of isoprene and pharmacological activities such as taxol (anticancer) and forskolin (for treating glaucoma). Triterpenes are contains 30 carbon atoms skeleton formed by the head-to-head joining of two C₁₅ chains, each of which contains three isoprene units joined head-to-tail. Tetraterpenes such as carotenoids are 40 carbon atoms made of 8 isoprene units.

Several terpenoids have been identified so far to have good activity in one or more of the antidiarrhoeal mechanisms described above. Some of the compounds include:

- ✚ Niloticane from *Acacia nilotica* (L.) Willd ex Del. subsp. *kraussiana* (Benth) Brenan (Fabaceae) had inhibitory effect on (*Bacillus subtilis*, *Staphylococcus aureus*, and *Escherichia coli* at 4.0, 8.0 and 33.0 µg/ml respectively) (Eldeen *et al.*, 2010).
- ✚ 1,3-Dihydroxy-12-oleanen-29-oic acid; 3,30-Dihydroxyl-12-oleanen-22-one; 1,3,24-Trihydroxyl-12-olean-29-oic acid; 1,23-Dihydroxy-12-oleanen-29-oic acid-3-O-2,4-di-acetyl-1-rhamnopyranoside isolated from *Combretum imberbe* have been reported to have MIC of 16.0 µg/ml against *E. coli* (Angeh *et al.*, 2007).
- ✚ Oleanolic acid, ursolic acid, and betulinic acid from *Chaenomeles speciosa* have the potential of blocking the binding of virulence heat labile unit B (LTuB) of *E. coli* enterotoxin to ganglioside receptor (GM 1). The IC₅₀ values enterotoxin binding activity were 202.8±47.8, 493.6±100.0, and 480.5±56.9 µM respectively (Chen *et al.*, 2007).
- ✚ Glycyrrhizin from *Glycyrrhiza uralensis* have been reported to have LTuB-binding inhibitory activity with IC₅₀ of 3.26±0.17 mM, therefore, can suppress LT-induced intestinal fluid accumulation (Chen *et al.*, 2009).
- ✚ Oleanolic acid and echinocystic acid both isolated from *Luffa cylindrica* (cucurbitaceae) were reported to increase phagocyte index, stimulate macrophage, increase humoral and cell-mediated immune responses (Khajuria *et al.*, 2007).




Fig. 2.9. Chemical structures of bioactive terpenoids against diarrhoeal mechanisms

-  Sugiol (a diterpenoid) isolated from *Amentotaxus formosana* reported to exhibit good xanthine oxidase inhibitory activity with IC_{50} of $6.8 \pm 0.4 \mu M$ compared to standard allopurinol with IC_{50} of $2.0 \pm 0.7 \mu M$ (Lin *et al.*, 2010).


- ✚ *Epibetulinic acid* and *betulonic acid* isolated from *Maytenus cuzcoina* Loesener demonstrated nitric oxide inhibition of $89.1 \pm 4.4\%$ (IC_{50} of $0.7 \mu M$) and $69.2 \pm 5.1\%$ (IC_{50} of $0.3 \mu M$) respectively in vitro (Reyes *et al.*, 2006).
- ✚ *Betulonic acid* isolated *Maytenus cuzcoina* Loesener to have demonstrated PGE inhibition activity of $58.4 \pm 3.9\%$ (IC_{50} of $2.7 \mu M$) (Reyes *et al.*, 2006) in vitro.
- ✚ *Maslinic acid*, *oleanolic acid*, *erythrodiol* and *uvaol* isolated from olive pomace oil was shown to have concentration dependent IL-6, TNF- α modulatory effects in a human mononuclear cells culture assay (Marquez-Martin *et al.*, 2006).
- ✚ *Ganoderic acids C and D* isolated from *Ganoderma lucidum* have been reported to have anti-allergic properties through histamine release inhibitory mechanisms (Rios, 2008). *Friedoolean-type triterpenoid* (*bryonolic acid*) (Rios, 2005) and *cucurbitacin* structure, including *dihydrocucurbitacin B* (Escandell *et al.*, 2007) and *cucurbitacin R* (Escandell *et al.*, 2010) from *Cayaponia tayuya* (Cucurbitaceae) were reported to exhibit anti-allergic activities through inflammatory responses modulation.

2.15.2. Alkaloids


Alkaloid refers to a group of heterocyclic nitrogen compounds and many exhibit remarkable physiological and pharmacological activities (Samy and Gopalakrishnakone, 2008). Most alkaloids are derived from amino acid precursor and are classified based on their structure as pyridine, tropane or pyrrolizidine alkaloids. Though, alkaloids have many pharmacological mechanisms such as microbiocidal effect on diarrhoeagenic pathogens, the main antidiarrhoea effect is probably that of delayed intestinal transition of bowel materials (Cowan, 1999).

Some of the pharmacological important antidiarrhoeal alkaloids include:


- ✚ *Kurryam*, *koenimbine* and *koenine* isolated from *Murraya koenigii* Spreng (Rutaceae) seed were reported to be active against various diarrhoeal mechanisms such as castor-induced diarrhoea, GIT motility, PGE₂ induced enterpooling (Mandal *et al.*, 2010).
- ✚ *8-acetyldihydronitidine* and *8-acetyldihydroavicine* both isolated from *Zanthoxylum tetraspermum* had strong antistaphylococcal activities with MICs of 1.56 and 3.12 $\mu g/ml$ respectively. However, the two compounds were reported to have no inhibitory activity against *E. coli* (Nissanka *et al.*, 2001).
- ✚ *Boldine* isolated from *Peumus boldus* Molina and *Corydalis ternate* Nakai had good antioxidant properties, indicating the effectiveness of the compound in preventing various oxidative-stress related illnesses like inflammatory cascades, immune dysfunctions. However, at high concentration, it causes cellular damage and potentiates lipoperoxidation, which is a pro-oxidant property.




Kurryam




Koenimbine




Koenine




(S)-boldine




Laurotetanine




Magnocurarine



(S)-reticuline



8-acetyldihydronitidine



8-acetyldihydroavicine

Fig. 2.10. Chemical structures of bioactive alkaloids against diarrhoeal mechanisms

2.15.3. Phenolic

Phytophenolic compounds are widely distributed as secondary metabolites of medicinal plants, as well as in some edible plants (Nacz and Shahidi, 2004). The consumption of diet rich in phenolic compounds has been hypothesized to be important in health promotion and disease prevention in humans and animals (Ramful *et al*, 2010). Phenolic compounds are characterized as aromatic metabolites that have one or more acidic hydroxyl groups attached to the phenyl ring. The sub-class of phenolics compounds are presented in Fig. 2.8. This group of compounds exhibits numerous biological activities directly or indirectly on intestinal epithelium which contribute to alleviation of diarrhoea symptoms.





Fig. 2.11 Sub-classes of biologically important group of phenolic compounds.

Phenolic compounds with anti-diarrhoeal activities against some of the mechanisms described above include:

- ✚ 2(S)-5'-(-1'',1'''-dimethylallyl)-8-(3'',3'''-dimethylallyl)-2',4',5,7-tetrahydroxyflavanone; 2(S)-5'- (1'',1'''-dimethylallyl)-8-(3'',3'''-dimethylallyl)-2'-methoxy-4',5,7-trihydroxyflavanone and 5'- (1'',1'''-dimethylallyl)-8-(3'',3'''-dimethylallyl)-2',4',5,7-tetrahydroxyflavone] from *Dalea scandens* var. *paucifolia* with MIC of 1.56 µg/ml against standard and Methicillin-resistant *S. aureus* (MRSA) (Nanayakkara *et al.*, 2002).
- ✚ Moracin T isolated from *Morus mesozygia* was reported to exhibit antimicrobial activities with MIC of 5.0 µg/ml against *E. coli*, *S. dysenterica*, *P. aeruginosa*, *S. typhi*, and 10 µg/ml against *S. aureus*, *C. albicans* (Kuethe *et al.*, 2009).
- ✚ Isoquercitrin, catechin and epicatechin isolated from *Chiranthodendron pentadactylon* flowers have antisecretory effect on *Vibrio cholerae* toxin induced intestinal fluid accumulation with ID₅₀ of 19.2, 51.7 and 8.3 µg/ml against loperamide (positive control) with ID₅₀ of 6.1 µg/ml.

- Davidigenin isolated from *Mascarenhasia arborescens* inhibits histamine induced contractile response of rat ileum and Ach induced contractile response on rat duodenum. The compound was reported to have non-competitive concentration dependent inhibitory activity (Desire *et al.*, 2010).
- Vitexin isolated from *Aloysia citriodora* has also been reported to have antispasmodic activities (Ragone *et al.*, 2007).
- Quercetin has 90.7±0.3% and 79.6±2.3% inhibition of Ach and the depolarized KCl induced contractions on guinea pigs isolated ileum at IC₅₀ <0.1 µg/ml respectively (Cimanga *et al.*, 2010).
- Quercitrin was reported to have spasmolytic activities of 82.3±2.3% and 72.1±0.6% against Ach and the depolarized KCl induced contractions on guinea pigs isolated ileum at IC₅₀ <0.01 µg/ml respectively (Cimanga *et al.*, 2010).
- Spasmolytic activities of rutin were also reported as 93.4±1.6% and 86.3±2.1% against Ach and the depolarized KCl induced contractions on guinea pigs isolated ileum at IC₅₀ <0.01 µg/ml respectively (Cimanga *et al.*, 2010).
- Luteolin isolated from *Pogonatherum crinitum* has iNOS inhibitory activity with E_{max} equals 100.00±0.00% and IC₅₀ of 10.41 µM while Kaempferol isolated from the same plant has iNOS inhibitory activity with E_{max} of 95.12±1.15% and the IC₅₀ equals 10.61±0.44 µM (Wang *et al.*, 2008)
- Stilbenoids such as r-2-viniferin, trans-amurensin and cis-amurensin isolated from *Vitis amurensis* have LOX inhibitory activity with IC₅₀ of 6.39±0.08 µM, 12.1±0.32 µM and 16.3±0.52 µM respectively (Ha *et al.*, 2009).
- Laurentixanthone A isolated from *Vismia laurentii* has good activities with MIC of 4.88 µg/ml against *S. dysenterica*, *S. flexneri*, *B. subtilis* (Nguemeving *et al.*, 2006).
- 5,7-dihydroxy-2-[14-methoxy-15-propyl phenyl]-4H-chromen-4-one isolated from *Leuca aspera* has superoxide radical scavenging activity of 75.4±0.31% and lipid peroxidation inhibition of 68.7±0.41% at a concentration of 40 ppm (Meghashri *et al.*, 2010).
- Lipid peroxidation inhibitory activity of foeniculose X isolated from *Foeniculum vulgare* at concentration of 10⁻⁵M was reported to be 6.30 nm TBARS/mg LDL (Marino *et al.*, 2007).
- An *in vivo* lipid peroxidation inhibitory activity of arzanol isolated from *Helichrysum italicum* has also been reported (Rosa *et al.*, 2011).



A survey on traditional practice in South Africa indicates that diarrhoea is one of the most prominent ailments being treated with medicinal plants (Dambisya and Tindimwebwa, 2003). At this stage, the scientific validation of their therapeutic potential, standardization, safety and mechanisms of actions of most of the plants is still lacking (Havagiray *et al.*, 2004). Ethnopharmacological properties and phytochemistry of the medicinal plants used for the treatment of diarrhoea in South Africa is reviewed and presented in Appendix 1.

2.17. Conclusion

Many medicinal plants are used in various traditional cultures of South Africa to treat diarrhoea and the associated complications. The traditional recipes include infusions, decoctions, and tinctures of the leaves, stems, roots, seeds and bark of medicinal plants. Several scientific methods have been used to evaluate and validate the traditional use of some of the plants as antidiarrhoeal agents. Such properties investigated as the potential antidiarrhoeal mechanism are the antimotility, antipropulsive, antioxidant, anti-inflammation, antispasmodic, antimicrobial, antiprotozoal, and immunomodulatory activity of the medicinal plant preparations. Several of these aspects will be examined for some selected medicinal plants in this study.

CHAPTER THREE

Plant selection, collection, extraction and analysis of selected species

3.1. Introduction

Renewed interest in the therapeutic potential of medicinal plants means that researchers are concerned not only with validating ethnopharmacological usage of plant, but also with identifying, isolating and characterizing the active components (Fennell *et al.*, 2004). However, the presence of numerous inactive components makes the screening and isolation of the target component(s) extremely cumbersome (Sticher, 2008). In choosing medicinal plants for scientific evaluation of their biological activities and validation of ethnopharmacological usage, some criteria such as

- Evidence of ethnopharmacological usage by the native population.
- The ailment(s) which the plant(s) is used to cure.
- The availability of the plant in its natural habitat.
- The sustainable use of the part(s) of the plant (root, leaves, stem, bark or whole plant) (Baker *et al.*, 1995; Van der Watt and Pretorius, 2001).
- Mode of preparation and administration by traditional healers must also be considered.

Plant quality and pre-treatment are also important determinant of the phytochemical constituents and invariably the biological activities of an extract. These factors depend on plant parts used, genetic variation, geographical location, climatic conditions, collection period, drying methods, and storage conditions. Due to these possible variations, plant material from recognized botanical gardens or herbaria is usually recommended because they are protected, correctly identified and serve as reliable sources for subsequent collections. Preparation of voucher specimens is also an important aspect of medicinal plant research. Standard procedures for pre-treatment of plant materials have been developed (Eloff *et al.*, 2008). The basic steps include pre-washing if necessary, air drying under shade at room temperature, grinding into powder and storage in an air tight container at appropriate temperature (room or refrigerated).

In view of limited resources and the large number of potential medicinal plants to be studied, efficient systems of evaluation need to be developed for rapid phytochemical and biological screening. The first step is the use of appropriate extraction process to remove the phytochemical from the plant cellular matrix (Sticher, 2008). Extraction processes need to be exhaustive, efficient, simple, rapid and inexpensive in extracting targeted compound(s). A number of extraction methods such as soxhlet, percolation, maceration, digestion, reflux, and steam distillation have been developed over the years (Sticher, 2008). However, solid-liquid extraction with a suitable range of solvents remains the most viable, convenient and effective procedure widely in use. The plant extracts are usually qualitatively analyzed for chemical composition (phytochemical fingerprint) and biological activities (for example bioautography for antimicrobial assay, antioxidative profiling with DPPH radical solution, acetyl cholinesterase inhibition) on thin layer chromatography (TLC). These plant pre-treatment methods,

extraction and analyses were employed in this chapter to determine qualitatively the phytochemical constituents and biological activities of selected plant extracts. The plants studied were selected based on literature documentation of their use in South African traditional medicine (SATM) as antidiarrhoeal agents and results from preliminary antimicrobial studies on some of the species in the phytomedicine tree project.

3.2. Solid-liquid extraction

Extraction is first pre-purification step in the isolation and characterization of active compound(s) of a medicinal plant (Sticher, 2008). Selective removal of interfering components from solid plant material involves a five-unit operation:

- Mixing of plant material and extractant.
- Solubilisation of the solute with the aid of a shaker or sonicator.
- Filtration of the mixture to remove solutes and extractant from the plant residue.
- Drying of sample using technique such as freeze drying, evaporation under vacuum (rota-evaporation) or air drying.
- Recovery of the solute extract.

The type of extractant may range from non-polar to polar solvent depending on the targeted class of bioactive component(s). Though the method is relatively simple, some of the drawbacks include: long extraction time, labour intensivity, high solvent consumption and inadequate reproducibility. In traditional medicine practice, ethanol and water are the most widely used extractants. The bioactive components of medicinal plants are usually unknown, and the nature of the extractant used affects the composition of the crude extract. Therefore, solvents such as hexane, dichloromethane, ethyl acetate, acetone, methanol, propanol, water or a combination of solvents are used in laboratory settings. Acetone has been adjudged to be the best extractant of plant extract for bioassay because it extracts a broad spectrum of components (polar and non-polar), is miscible with all other solvents, is highly volatile, and exhibit low toxicity to biological organisms in various assays (Eloff, 1998).

Temperature is also an important factor in extraction, drying and storage of plant extracts because of varying compound stability due to chemical degradation, losses by volatilization and oxidation. Milder extracting and drying temperatures are required to avoid loss of activity by plant extracts possibly due to thermal decomposition. Storage of plant extracts, fractions or isolated pure compounds should be done at 4°C in the dark to avoid any negative influence of temperature and light.

3.3. Liquid-liquid fractionation

Solvent partitioning of extracts allows a finer separation of the plant constituents into fractions of different polarity. Bioactivity-guided fractionation, where the fractions are tested following separation to quickly identify and isolate the agents responsible for bioactivity is a desirable step in medicinal plant research. The solvent partition process involves the use of two immiscible solvents of different polarities. Various solvents are used

starting with non-polar (hexane, dichloromethane, diethyl ether) to medium polar (chloroform, ethyl acetate), and finally more polar solvent (acetone, methanol, butanol and water).

3.4. Thin layer chromatography (TLC)

3.4.1. Phytochemical fingerprints

TLC is widely used in natural product extract analysis, stability tests of extracts and finished products, and in sample quality control (Cimpoi, 2006). TLC fingerprints of medicinal plants and extracts can be used for identification and quality control of medicinal preparations. The identification of separated components can be achieved on the basis of retention factor (R_f) values and colour spots. In relation to other chromatographic methods, TLC offers the simplest and cheapest means of detecting natural product constituents, requiring little sample clean-up and equipment (Nyiredy and Glowniak, 2001). Characteristic features of TLC include: analysis of many samples and comparison of their phytochemical profiles on the same plate; results can be stored and communicated as images (picture, video or scanned) and flexibility in the choice of mobile and stationary phase (Cimpoi, 2006). Identification of compounds can be done using three different mobile phases on the same stationary phase or three different stationary phases with one mobile phase to develop the fingerprint of the extracts and standards. If the difference in R_f values is less than 0.03, then the compounds is identified without further isolation (Nyiredy and Glowniak, 2001). However, position isomeric compounds such as ursane and oleanane derivatives can have superimpose or close R_f values, making them inseparable.

Visualization of separated compounds is achieved by natural colour in daylight or by fluorescent quenching on 254 nm (for conjugated double bonds or extended π electron systems) or 366 nm UV light. Some commercial plate absorbents contain fluorescent dye that lights when placed under UV light and compounds are indicated with blue, green, brown, red or purple areas against a fluorescent background. Visualization of chromatogram under UV light at 366 nm shows orange-yellow bands for flavonoids and blue fluorescent bands for phenolic acids (Males and Medic-Saric, 2001). Many chromogenic spray reagents are also available for specific classes of compounds or serve as indicators for broad classes of compounds. Examples are vanillin/sulphuric acid solution, anisaldehyde and ferric chloride-potassium ferricyanide given intense blue bands for phenolic compounds (Wettasinghe *et al.*, 2001).

3.5. Materials and Methods

3.5.1. Selection of South Africa medicinal plants for antidiarrhoeal screening

For this project, 27 plant species from nine families (Table 3.1) were selected for preliminary screening based on the following criteria:

1. Ethnopharmacological use of the plant in the management of diarrhoea locally,
2. Phylogenetic relationship to other plants used in treatment diarrhoea due to the possibility of their producing related chemical compounds (chemotaxonomy),

3. Medicinal plants reportedly used in countries other than South Africa but naturalized or endogenous in South African flora,
4. Preliminary pharmacological evaluation of the medicinal plant from the phytomedical laboratory of the Department of Paraclinical Sciences (University of Pretoria),
5. Absence of published literature describing antidiarrhoea and biological studies, and
6. Their availability for evaluation.

A literature review on the selected plants for antidiarrhoea and other biological studies yielded little or no previous research work.

3.5.2. Collection of plant materials

The leaves of the 27 plants were collected from the Marie van der Schijff Botanical Garden University of Pretoria Main Campus at Hatfield, Pretoria or from Phytomedicine Programme tree project stored samples. The plants were identified and authenticated by Ms. Lorraine Middleton and Magda Nel at the University of Pretoria Botanical Garden. Voucher specimens were maintained at the HGWJ Schweikert Herbarium of the Department of Plant Science, University of Pretoria, Hatfield Campus, Pretoria, South Africa.

3.5.3. Preparation of plant material and optimization of phenolic-enriched extraction process

Plant leaves collected were pre-treated according to Phytomedicine programme (University of Pretoria) standard protocol. In brief, the leaves were sorted from the stem, packed in a well perforated bag and air dried under shade at room temperature for 2 week. The dried leaves were ground, powdered and kept in an air tight polyethylene bag until needed for the extraction process. Simultaneous extraction and fractionation of the leaves using a mixture of 70% acetone acidified with 0.1% HCl and hexane. The chlorophyll, fat and wax-enriched hexane fraction was decanted from the phenolic-enriched 70% acetone fraction.


Table 3.1: Medicinal plants selected for antidiarrhoeal investigation in this study

Genera	Family/Species	syn	Voucher specimen information	Reasons for selection
Anacardiaceae	<i>Ozoroa mucronata</i> (Bernh.ex C.Krauss) R.fern & A. Fern	Ozm	PRU 068928	2, 4, 5, 6
	<i>Ozoroa paniculosa</i> (Sond.) R.fern & A. Fern	Ozp	PRU 66851	1, 2, 4, 5, 6
	<i>Searsia leptodictya</i> Diels	Sle	PRU 70151	2, 4, 5, 6
	<i>Searsia pendulina</i> Jacq.	Spd	PRU 84141	2, 4, 5, 6
	<i>Searsia pentheri</i> Zahlbr.	Spt	PRU 709769	2, 4, 5, 6
Apocynaceae	<i>Carissa macrocarpa</i> (Eckl.) A.DC	Cam	PRU 37819	2, 4, 5, 6

Burseraceae	<i>Commiphora harveyi</i> (Engl.) Engl.	Com	PRU 49952	2, 4, 5, 6
Celastraceae	<i>Maytenus peduncularis</i> (Sond.) Loes.	Mpd	PRU 76382	2, 4, 5, 6
	<i>Maytenus probumbens</i> (L.f.) Loes.	Mpr	PRU 77119	2, 4, 5, 6
	<i>Maytenus senegalensis</i> (Lam.) Exell	Mse		1, 2, 3, 4, 5
	<i>Maytenus undata</i> (Thunb.) Blakelock	Mun	PRU 18576	1, 2, 3, 4, 5, 6
Combretaceae	<i>Combretum bracteosum</i> (Hochst.) Brandis ex Engl.	Cob	PRU 117443	1, 2, 4, 5, 6
	<i>Combretum padoides</i> Engl. & Diels	Cop	PRU 115416	1, 2, 4, 5, 6
	<i>Combretum vendae</i> A.E. van Wyk	Cov	PRU 50800	1, 2, 4, 5, 6
	<i>Combretum woodii</i> Dummer	Cow	PRU 20544	1, 2, 4, 5, 6
Ebenaceae	<i>Euclea crispa</i> (Thunb.) Gurke	Euc	PRU 76444	2, 4, 5, 6
	<i>Euclea natalensis</i> A.DC.	Eun	PRU 66327	1, 2, 4, 5, 6
Fabaceae	<i>Bauhinia bowkeri</i> Harv	Bab	PRU 44967	2, 4, 5, 6
	<i>Bauhinia galpinii</i> N. E. Br	Bag	PRU 28944	1, 2, 4, 5, 6
	<i>Bauhinia petersiana</i> Bolle	Bap	PRU 66874	2, 4,5
	<i>Bauhinia variegata</i> L.	Bav	PRU 38533	1, 2, 3, 4, 5, 6
	<i>Erythrina latissima</i> E. Mey	Erl	PRU 16349	2, 4, 5, 6
	<i>Indigofera cylindrical sensu</i> E. Mey	Inj		2, 3, 4, 5, 6
	<i>Schotia brachypetala</i> Sond.	Scb	PRU 55333	1, 2, 4, 5, 6
Moraceae	<i>Ficus craterostoma</i> Warb.ex Mildbr. & Burret	Fic	PRU 38554	2, 4, 5, 6
	<i>Ficus glumosa</i> Delile	Fig	PRU 48293	1,2, 4, 5
Myrtaceae	<i>Syzygium paniculatum</i> Gaertner	Syp	PRU 115417	2, 3, 5, 6

(1) Ethnopharmacological use of the plant in the management of diarrhoea locally, (2) phylogenetic relationship to other plants used in treatment diarrhoea due to the possibility of their producing related chemical compounds (chemotaxonomy), (3) medicinal plants reportedly used in countries other than South Africa but naturalized or endogenous in South African flora, (4) preliminary pharmacological evaluation of the medicinal plant from the Phytomedicine Programme of the Department of Paraclinical Sciences (University of Pretoria), (5) absence of published literature describing anti-diarrhoea and biological studies, and (6) their availability for evaluation.

The acetone residue was removed by evaporation under vacuum using a rotary evaporator at 40°C. The residual water fractions were divided into two portions (A and B). Portion A was freeze dried and served as the crude extract while portion B was fractionated using solvents of increasing polarities as presented in Figure 3.1. The crude extracts and fractions were reconstituted in various suitable solvents for the biological assays.






Fig. 3.1. Flow chart for the extraction, phytochemical analysis and fractionation of plant material

3.5.4. Phytochemical profiling

The phytochemical profiles of the crude extracts and fractions were determined using thin layer chromatography (TLC) by spotting 10 μ l of solution at a concentration of 10 mg/ml. The plates were developed with various combinations of hexane (H), ethyl acetate (E), formic acid (F), acetic acid (A), chloroform (C), methanol (M), water (W), benzene (B) and ammonia (Am) at different ratio to create eluting solvent of varied polarities. The combination with ratios in parenthesis that were used:

- (1) E: F: A: W (70:5:5:10)
- (2) E: F: A: W (70:5:15:10)
- (3) E: M: Am (90:20:15)
- (4) H: E: F (90:10:2),
- (5) H: E: F (70:30:2)
- (6) H: E: F (50:50:2)
- (7) (B: E: Am (90:10:1)
- (8) C: E: F (50:40:10)
- (9) E: M: W: F (50:6.5:5:2)
- (10) H: E: F (20:80:2)

The developed TLC plates were sprayed with vanillin/ H_2SO_4 solution and heated at 100°C to allow colour development (FAO/IAEA, 2000). Other reagents such as ferric chloride-potassium ferricyanide and p-anisaldehyde/ H_2SO_4 (acetic acid, 5 ml; conc. H_2SO_4 , 25 ml; ethanol, 425 ml; water, 25 ml) (Kubata *et al.*, 2005) were also used.

3.6. Quantification of the phenolic constituents of the extracts

3.6.1. Determination of total phenolic constituents

The total phenolic constituents of the extracts were determined using Folin-Ciocalteu method as described by Makkar 2003, with some modifications. The crude extracts at concentration of 1:1 (mg/ml) plant material: extracting solvent (50 μ l) was dispensed into a test tube and made up to 500 μ l with distilled water. Folin-Ciocalteu reagent (250 μ l) diluted with distilled water (1:1) and 1250 μ l of 20% sodium carbonate solution were

added to the extract. The mixture was vortexed and absorbance recorded at 725 nm after 40min incubation at room temperature. The amount of polyphenols (expressed as mg Gallic /g dry weight) was calculated from a prepared standard curve for gallic acid (0.0019-0.25 mg/ml gallic acid). The standard curve equation is $y = 4.9022x + c$, where y is absorbance, x is mg Gallic acid, $c=0$, $R^2=0.9804$)

3.6.2. Determination of total tannin

The total tannin content of the extracts was determined using polyvinylpyrrolidone (PVPP) binding method (Makkar, 2003). The bound mixtures were prepared by mixing 100 mg of PVPP, 1.0 ml of distilled water and 1.0 ml of tannin-containing extracts in a centrifuge tube. The mixtures were mixed thoroughly and kept at 4°C for 15 min and then filtered. The filtrate (100 µl) was transferred into a test tube and the phenolic content was evaluated as described in section 3.7.1 above. Non-tannin phenolic constituents were determined from the standard curve of catechin expressed as catechin equivalent in mg/g dry material. The standard curve equation is $y = 4.9022x + c$, where y is absorbance, x is mg Gallic acid, $c=0$, $R^2=0.9804$). The tannin content was calculated as the difference between the total phenolic and non-phenolic content of the extracts because the tannin was bound and precipitated by PVPP.

3.6.3. Determination of proanthocyanidin

The proanthocyanidin content of the extracts was determined using the butanol-HCl assay as described by Makkar, 2003. The extract (500 µl) was dispensed into a test tube and diluted to 10 ml with 70% acetone. To this 3 ml of butanol/HCl (95/5%) and 100 µl of 2% ferric ammonium sulphate in 2N HCl were added. The test tubes were loosely covered and heated in a boiling water bath for 50min. The absorbance was recorded at 550 nm after the tubes were allowed to cool to room temperature. Absorbance of the unheated mixture was used as blank.

3.6.4. Determination of condensed tannin

The condensed tannin content of the extracts was determined using vanillin/HCl assay as described by Heimler *et al*, 2006. To 0.5 ml of the extract measured into a test tube, 3 ml of vanillin reagent containing 4% concentrated HCl and 0.5% of vanillin in methanol was added. The mixture was allowed to stand for 15 min. The absorbance was recorded at 500 nm against methanol as blank. The amount of condensed tannin in the extracts was expressed as catechin equivalent (CE)/g dry plant material. The standard curve ranged from 0.0019 to 0.25 mg/ml (Absorbance= 0.1791 mg catechin + 0.0504, $R^2=0.944$).

3.6.5. Determination of hydrolysable tannin (gallotannin)

The gallotannin content of the extracts was determined using the potassium iodate assay (Vermeris and Nicholson, 2006). To 3 ml of the extract, 1 ml of saturated solution of potassium iodate was added and allowed to stand at room temperature for 40 min. The absorbance was read at 550 nm. A standard curve was prepared

using gallic acid under the same conditions as the extracts and results were expressed as gallic acid equivalent (GAE)/g dry plant material (Absorbance= 0.8264mg catechin + 0.0392, R²=0.9155).

3.6.6. Determination of total flavonoids and flavonol

The total flavonoids content of the extracts was determined by aluminium chloride method as described by [Abdel-Hamed et al \(2009\)](#) with some modification. Briefly 100 µl of the extract was mixed with 100 µl of 20% AlCl₃ and two drops of glacial acetic acid. The mixture was diluted with methanol to 3000 µl. Absorbance was read at 415 nm after 40 mins. Blank samples were prepared with the extract without AlCl₃. Standard curve was prepared using quercetin (3.9-500 µg/ml) in methanol under the same condition. The amount of flavonoids was expressed as mg quercetin equivalent/g of dry plant material (Absorbance= 4.9747 mg quercetin, R²=0.9846).

The flavonol content of the extracts was determined by aluminium chloride method as described by [Abdel-Hamed et al \(2009\)](#) with some modification. One ml of the extract was mixed with 1 ml of 20 mg/ml of AlCl₃ and 3 ml of 50 mg/ml of CH₃COONa. Standard curve was prepared using quercetin (0.0019 - 0.0312 mg/ml) in methanol under the same condition. Absorbance was read at 440 nm after 2.5 hr. The amount of flavonol was expressed as mg quercetin equivalent/g of dry plant material (Absorbance= 34.046mg quercetin, R²=0.9853).

3.6.7. Determination of anthocyanin

Total anthocyanin content of the extracts was determined by a pH differential method with 96 well microplate ([Lee et al, 2008](#), [Lee et al., 2005](#)) using spectrophotometer. Absorbance was measured at 520 nm and 700 nm in buffers at pH 1.0 and 4.5 using a molar coefficient of 29,600. Results were expressed as mg cyanidin-3-glucoside equivalent/g dry plant material using equation 1 and 2 ([Lee et al., 2005](#)).

$$\text{Equation 1} \quad A = (A_{520} - A_{700})_{\text{pH } 1.0} - (A_{520} - A_{700})_{\text{pH } 4.5}$$

$$\text{Equation 2} \quad \text{Anthocyanin (cyanidin-3-glucose equivalent mg/L)} = A \times \text{MW} \times \text{DF} \times 10^3 / \square \times l$$

Where MW (molecular weight) = 449.2 g/mol for cyanidin-3-glucose (cyn-3-glu); DF = dilution factor; l = pathlength in cm; \square = 26900 molar extinction coefficient in L × mol⁻¹ × cm⁻¹ for cyn-3-glu; 10³ factor for conversion from g to mg.

3.7. Results

3.7.1. Yield of extractions and fractionations processes

The yield of the phenolic-enriched crude extracts and the fractions of various polarities using hexane, dichloromethane, ethyl acetate, butanol and residual water are presented in Table. 3.2. The 70% acetone was an extremely efficient extractant with an average of 34.61±5.84% extracted. The maximum yield was obtained for the crude extracts of *S. leptodicya* (48.50±12.47% g/g dried plant material) followed by *O paniculosa*

($43.87 \pm 6.60\%$ g/g dried plant material) while *S. pentheri* (21.13 ± 2.67 g/g dried plant material) yielded the least. There was a surprisingly high standard deviation between the three repetitions with a single extraction with new plant material. This may have been caused by a difference in the particle size of samples. The extraction process efficiently removed the chlorophyll from the bulk 70% acetone extractant into hexane fraction. In most cases there was a difference between the percentage extracted and the total percentage of all the fractions. This loss may be ascribed to solubility difficulties encountered with the dried residual water fraction which could not be reconstituted due to the formation of insoluble complexes between the polyphenolics and other high molecular weight components such as polysaccharides and possibly alkaloids. Unfortunately at that stage a freeze drying was not available. This problem may have been partially resolved if the water fraction was freeze dried. To evaluate the degree to which the different plant species contain compounds of different polarity the percentage of quantity present in the crude extract into the different fractions was calculated (Table 3.2).

Table 3.2. The percentage yield of the crude extracts and various fractions (g/g dried plant material)

Plant spp	Crude	Hexane	DCM	ETOAc	Butanol	Residual Water	Insoluble ppt
Bab	33.25±0.83	3.44±0.15	1.39±0.16	2.45±0.34	9.91±1.09	7.47±0.41	8.27±1.87
Bag	38.83±6.18	2.38±0.35	1.31±0.27	2.70±0.21	15.05±1.16	8.93±0.18	10.90±2.94
Bap	32.23±2.84	1.67±0.17	1.71±0.17	3.30±0.78	11.82±0.48	8.71±1.23	7.69±2.53
Bav	31.62±5.46	1.83±0.07	1.90±0.23	2.98±0.17	10.65±1.68	8.07±0.13	
Erl	22.12±0.32	1.35±0.55	0.34±0.05	0.26±0.06	6.63±2.20	10.44±0.96	2.77±1.87
Inc	36.15±1.62	1.58±0.17	0.78±0.08	1.71±0.32	12.10±0.66	9.31±1.51	7.84±2.47
Scb	30.15±3.47	1.53±0.49	1.14±0.40	1.58±0.06	11.54±1.00	7.39±1.93	6.21±1.30
Cob	34.24±3.08	1.27±0.21	3.39±1.13	3.28±0.44	8.15±0.30	8.81±1.40	
Cop	39.96±0.78	2.33±0.51	3.31±0.68	3.56±0.18	17.42±0.79	7.43±1.06	1.84±0.96
Cov	38.77±0.48	1.33±0.47	2.86±0.24	3.13±0.51	14.82±2.53	12.08±0.16	2.74±1.06
Cow	36.88±3.39	3.95±1.75	2.07±0.38	2.67±0.41	12.35±2.99	8.39±0.52	7.41±3.29
Ozm	30.65±2.44	2.0±0.20	0.86±0.11	1.03±0.01	7.28±1.53	14.25±2.08	2.66±1.02
Ozp	43.87±6.60	6.57±0.55	1.55±0.38	4.30±0.82	14.54±0.96	8.81±2.02	8.62±1.77
Sle	48.50±12.47	5.85±0.61	1.49±0.30	4.25±0.52	10.30±1.82	9.29±0.89	13.28±3.34
Spd	33.76±0.28	5.05±0.69	0.98±0.28	3.03±0.35	12.21±0.81	11.40±3.31	1.96±0.53
Spt	21.13±2.67	2.96±0.30	1.50±0.25	1.04±0.28	6.34±0.17	8.80±1.50	1.62±0.89
Mpd	33.12±1.07	3.80±0.04	1.22±0.14	1.32±0.08	8.89±0.92	13.39±1.92	4.24±1.98
Mpr	35.10±4.77	3.05±0.28	1.18±0.31	0.90±0.28	8.50±1.31	12.41±0.47	4.78±1.35
Mse	37.89±3.05	3.75±0.40	1.20±0.14	1.30±0.08	8.77±0.91	13.21±1.89	10.08±2.71
Mun	36.89±4.67	1.42±0.18	1.63±0.48	1.08±0.46	10.88±0.46	12.08±1.77	3.13±1.57
Euc	34.97±1.90	2.76±0.56	1.49±0.32	2.05±0.16	13.44±0.86	10.81±0.30	2.84±0.73
Eun	32.83±3.19	2.05±0.82	1.73±0.21	2.35±0.45	10.89±1.39	12.76±1.96	2.77±1.87
Fic	25.68±3.22	1.50±0.06	0.63±0.07	0.91±0.07	7.59±1.93	9.39±0.30	2.69±1.36
Fig	35.22±4.04	1.82±0.11	1.13±0.16	1.44±0.17	12.84±1.43	8.35±0.58	10.54±2.55
Cam	40.80±1.57	2.21±0.13	0.90±0.08	2.73±0.42	10.77±2.67	11.04±1.92	9.77±3.54
Com	33.09±1.19	1.01±0.04	1.06±0.28	1.64±0.08	10.39±4.81	7.80±0.59	5.37±1.85
Syp	36.80±8.10	1.44±0.20	0.78±0.06	1.10±0.17	8.34±1.89	10.60±0.40	5.95±2.64

3.7.2. Phytochemical screening (fingerprints)

The TLC phytochemical profiles of the crude extracts and fractions of the 27 plant species investigated are presented in Figs 3.1–3.4. Figures 3.1, 3.2, 3.3, and 3.4 are the TLC fingerprints of the crude, hexane fraction, dichloromethane fraction, ethyl acetate fraction developed with three mobile phases of different polarities for each fraction.

In each chromatogram the order from left to right was **Bab** (*Bauhinia bowkeri*), **Bag** (*Bauhinia galpinii*), **Bap** (*Bauhinia petersiana*), **Bav** (*Bauhinia variegata*), **Erl** (*Erythrina latissima*), **Inc** (*Indigofera cylindrica*), **Scb**

(*Schotia brachypetala*), **Cob** (*Combretum bracteosum*), **Cop** (*Combretum padoides*), **Cov** (*Combretum vendae*), **Cow** (*Combretum woodii*), **Ozm** (*Ozoroa mucronata*), **Ozp** (*Ozoroa paniculosa*), **Sle** (*Searsia leptodictya*), **Spd** (*Searsia pendulina*), **Spt** (*Searsia pentheri*), **Mpd** (*Maytenus peduncularis*), **Mpr** (*Maytenus procumbens*), **Mse** (*Maytenus senegalensis*), **Mun** (*Maytenus undata*), **Cam** (*Carissa macrocarpa*), **Com** (*Commiphora harveyi*), **Syp** (*Syzygium paniculatum*).




Fig.3.2. Chromatograms of 100 µg of crude extracts of different plant species developed with ethyl acetate: acetic acid: formic acid: water (75:5:5:10) (top), ethyl acetate: acetic acid: formic acid: water (70:5:15:10) (middle) and ethyl acetate: methanol: ammonia (90:20:15) (bottom) and visualized with vanillin sulphuric acid. For identity of plant species see under section 3.7.2 or under abbreviations used.

The chromatograms revealed complex mixture of compounds which exhibited different coloured reactions with the vanillin/H₂SO₄ spray reagent. The classes of compounds in the extracts include terpenoids (purple or bluish purple) (Taganna *et al.*, 2011) and phenolics such as flavonoids (yellow, pinkish or orange), stilbenes (bright red to dark pink colour), and proanthocyanidins (pink colour). The phenolic components were confirmed by blue-black spots with ferric chloride-potassium ferric cyanide reagents (Wettasinghe *et al.*, 2001) while the flavonoids were confirmed by yellow spot (Rijke *et al.*, 2006) with aluminium chloride/acetic acid spray reagent (AlCl₃/CH₃COOH).




Fig. 3.3: Chromatograms of the hexane fractions of different plant species developed with hexane: ethyl acetate: formic acid (90:10:2) (top), hexane: ethyl acetate: formic acid (70:30:2) (middle), and benzene: ethyl acetate: ammonia (90:10:1) (bottom) and visualized with vanillin sulphuric acid. For identity of plant species see under section 3.7.2 or under abbreviations used.

Characterization of the phytochemical profiles of the extracts indicated that the extraction method and extractants used resulted in splitting the complex mixtures into polar components concentrated in the 70% acetone component (crude extracts) and non-polar compounds concentrated in the hexane component. From the chromatogram, the crude extracts contained phenolics (especially flavonoids and proanthocyanidin) and terpenoids. The hexane and dichloromethane fractions contained prominent spots for terpenoids while the ethyl acetate fractions had prominent spots typical of flavonoid and other phenolic compounds.




Fig 3.4: Chromatograms of the dichloromethane fractions of different plant species developed with hexane: ethyl acetate: formic acid (70:30:2) top, hexane: ethyl acetate: formic acid (50:50:2) (middle) and chloroform: ethyl acetate: formic acid (50:40:10) bottom and visualized with vanillin sulphuric acid. For identity of plant species see under section 3.7.2 or under abbreviations used.




Fig 3.5 Chromatograms of the ethyl acetate fractions of different plant species developed with chloroform:ethyl acetate:formic acid (50:40:10) (top), hexane: ethyl acetate:formic acid (10:90:10) (middle) and ethyl acetate:methanol:water:formic acid (100:13:10:2) (bottom) and visualized with vanillin sulphuric acid. For identity of plant species see under section 3.8.2 or under abbreviations used.

3.7.3. Phenolic composition of the crude extracts

In this study, the total phenolic, total tannin, condensed tannin, proanthocyanidin, hydrolysable tannin as gallotannin, flavonoids and flavonol constituents of the phenolic-enriched crude extracts were evaluated using various standard protocols. The total polyphenolic and non-tannin phenolic constituent of each crude extract was evaluated using the Folin-Ciocalteu reagent. All the 27 extracts contain significant amount of polyphenols and non-tannin compounds; however, the quantity varied widely between the species ($74.91 \pm 1.26 - 467.0 \pm 15.8$ mg GAE/g plant material) (Fig.3.6).

Among the different extracts tested, the highest content of polyphenols was *Combretum padoides* (467.0 ± 15.8 mg GAE/g plant material) which did not differ significantly ($P < 0.05$) to *Combretum vendae* with (444.20 ± 15.4 mg GAE/g plant material). These two plant species were followed by *Carissa macrocarpa* (354.15 ± 3.01 mg GAE/g plant material), *Commiphora harveyi* (362.60 ± 2.10 mg GAE/g plant material), *Euclea natalensis* (204.98 ± 1.89 mg GAE/g plant material), *Ozoroa paniculosa* (370.89 ± 4.80 mg GAE/g plant material) and *Searsia pendulina* (339.80 ± 5.10 mg GAE/g plant material) all of which are significantly similar. The lowest content of polyphenols was *Ozoroa mucronata* with (74.91 ± 1.26 mg GAE/g plant material) which is not significant different from that of

the *Erythrina lattisima* with (76.08±2.59 mg GAE/g plant material) followed by *Maytenus procumbens* (112.71±1.51 mg GAE/g plant material) significantly similar to *Maytenus undata* (123.82±1.45 mg GAE/g plant material). The non-tannin phenolic constituent of the crude extracts ranges from 31.45±1.16 to 174.72±0.39 mg GAE/g plant material (Fig 3.6). The plant with the highest non-tannin phenolics was *C. macrocarpa* (174.72±0.39 mg GAE/g plant material) followed by *C. vendae* (155.80±6.40 mg GAE/g plant material) which was not significantly different to *O. paniculosa* (139.93±5.93 mg GAE/g plant material) $p < 0.05$.




Fig. 3.6. Total phenolic and non tannin constituent of the crude plant extracts

The plant with lowest content of non-tannin phenolic was *M. procumbens* (31.46±1.16 mg GAE/g plant material) which was not significantly different to *M. undata* (35.64±2.12 mg GAE/g plant material) $p < 0.05$.

The total tannin content of the extracts ranged from 25.55±0.81 to 359.40±8.30 mg GAE/g plant material (Fig.3.6). The highest tannin constituent was *C. padoides* (359.40±8.30 mg GAE/g plant material) and was

mainly hydrolysable gallotannin (305.80 ± 19.09 mg GAE/g plant material) (Fig 3.7). This was followed by *C. vendae* (288.40 ± 8.30 mg GAE/g plant material) which also contained high hydrolysable gallotannin (197.60 ± 12.79 mg GAE/g plant material)




Fig. 3.7. Total tannin and condensed tannin of the crude extracts

The hydrolysable gallotannin constituents of *C. vendae* were not significantly different ($p < 0.05$) to that for *Euclaea crispa* and *Indigofera cylindrical* at 199.36 ± 17.61 and 185.21 ± 11.50 mg GAE/g plant material respectively. *E. latissima* had the lowest tannin content at 25.55 ± 0.81 mg GAE/g plant material followed by *O. mucronata* at 27.17 ± 0.18 mg GAE/g plant material. For both these plants the tannin content was mainly proanthocyanidin at 33.42 ± 3.76 and 19.88 ± 2.51 mg LE/g plant material respectively (Fig 3.8).




Fig. 3.8. Proanthocyanidin and gallotannin constituent of the crude extracts

The condensed tannin content ranged from 6.99 ± 0.32 to 183.53 ± 10.10 mg CE/g plant material. *Bauhinia galpinii* had the highest condensed tannin at 183.53 ± 10.10 mg CE/g plant material. This was followed by *C. macrocarpa*, *Bauhinia bowkeri*, and *Combretum bracteosum* at 125.0 ± 2.72 , 120.02 ± 8.37 and 121.90 ± 5.50 mg CE/g plant material respectively, which are not significantly different ($p < 0.05$) from each other. *O. mucronata* had the lowest condensed tannin at 6.99 ± 0.32 mg CE/g plant material followed by *M. pendulina* at 7.32 ± 1.20 mg CE/g plant material and *E. latissima* at 11.90 ± 0.8 mg CE/g plant material.

The highest proanthocyanidin content was found in *C. macrocarpa* at 213.10 ± 7.00 mg LE/g plant material followed by *S. leptodictya*, *B. galpinii*, and *Searsia pendulina* at 126.54 ± 6.46 , 121.08 ± 2.20 and 117.83 ± 2.24 mg LE/g plant material respectively. Statistically, *S. leptodictya*, *B. galpinii*, and *Searsia pendulina* were not

significantly different ($p < 0.05$) for their proanthocyanidin content. *M. procumbens* and *S. pentheri* had the lowest proanthocyanidin content at 10.46 ± 1.76 and 10.08 ± 2.24 mg LE/g plant material respectively.




Fig. 3.9. Total flavonoids and the flavonol constituent of the crude extracts

The total flavonoid ranged from 11.27 ± 3.37 to 176.87 ± 5.96 mg QE/g plant material. The highest flavonoid content was present in *Schotia brachypetala* at 176.87 ± 5.96 mg QE/g plant material followed by *O. paniculosa* 168.27 ± 5.96 mg QE/g plant material. No significant difference ($p < 0.05$) was present between the two plant (Fig 3.9). The lowest flavonoid content was present in *M. senegalensis* at 11.27 ± 3.37 mg QE/g plant material. The highest flavonol content was present for *M. procumbens* at 17.85 ± 0.20 mg QE/g plant material followed by *C. padoides* at 8.81 ± 0.13 mg QE/g plant material. The lowest flavonol content was *C. bracteosum* at 0.13 ± 0.07 mg QE/g plant material.

3.8. Discussion

3.8.1. Yield

The extraction of phenolic constituents from plant matrix is complex and is influenced by their chemical nature, extraction method, sample particle size, solvent as well as presence of interfering substances. Phenolics can also complex with carbohydrates, proteins, and other plant components like alkaloids. High molecular weight phenolics and their complexes are usually insoluble and solubility is also a function of the solvent polarity (Naczek and Shahidi, 2004). Consequently, phenolic extracts composed of varied classes of phenolics present in different proportion with the degree of solubility in the solvent system as the primary determinant. In this experiment, simultaneous extraction and fractionation using acidified 70% acetone and n-hexane was adopted. Two immiscible phases of phenolic-enriched acetone solution (low phase) and chlorophyll-enriched terpenoids containing hexane (upper phase) were obtained.

The extraction process is an important factor for assessing the biological activity of medicinal plant extracts (Berlin and Berlin, 2005) as it influence yield of the extracts, extractive capacity of an extractant, and quality parameter of the herbal preparations (Albuquerque and Hanazaki, 2006). Low polar solvent extractants such as hexane, petroleum ether and dichloromethane extract non polar compounds mainly of terpenoids or highly methoxylated phenolics. In contrast, medium and high polar solvents such as ethyl acetate, acetone, methanol, ethanol water or mixture of these solvents extract the polar compounds ranging from simple phenolics to complex polymeric phenolics (tannins).

3.8.2. Thin layer chromatogram

TLC fingerprints of the plant leaf crude extracts and fractions showed complex mixtures of non-polar to polar compounds. TLC was used as qualitative method to characterize and document the phytochemical profiles of the extracts as a fingerprint. The phytochemical constituents of plants depend on several factors including seasonal changes, biotic (genetic) and abiotic (climatic stress, infection and soil fertility) factors (Moure *et al.*, 2001). TLC analyses help in monitoring composition of the extracts and fractions to ensure that no component(s) are lost during processing. It also provides a means of comparing phytochemical composition of different plant extracts developed side by side.

When comparing TLC fingerprints of the hexane fractions and the crude extracts from the extraction process, hexane fractions were enriched with non-polar components while the crude extracts were enriched with polar components, mostly of phenolic compounds. Solvents (2) E: F: A: W (70:5:15:10), (5) H: E: F (70:30:2), (8) C: E: F (50:40:10) and (9) E: M: W: F (50:6.5:5:2) were the best mobile phase obtainable for preparing TLC fingerprint of the crude extract, hexane fractions, dichloromethane fractions and ethyl acetate fractions respectively in this work.

Polyphenolic compounds are important bioactive component of medicinal plant extract exhibiting various pharmacological properties (Vundac *et al.*, 2007). Phenolic-enriched extracts have been reported to correlate with a wide range of physiological and health benefits which include antiallergenic, antiviral, antibacterial, antifungal (Pietta, 2000), antisecretory, antispasmodic, antimotility (Yue *et al.*, 2004), anti-inflammatory,

immunomodulatory and parasitic activities. In traditional medicine preparation of plant extract recipe, water or ethanolic solutions are the main extractants.

3.8.3. Phenolic constituents of the crude extracts

Polyphenolic compounds are important bioactive component of medicinal plant extract exhibiting various pharmacological properties (Vundac *et al.*, 2007). Phenolics form one of the main classes of secondary metabolites and several thousand (among them over 8,150 flavonoids) different compounds have been identified with a large range of structures: monomeric, dimeric and polymeric phenolics (Lattanzio *et al.*, 2006). Several classes of phenolics have been categorized on the basis of their basic skeleton. These groups of phytochemicals are primarily natural antioxidants which act as reducing agent, metal chelators and single oxygen quenchers. Phenolic-enriched extracts have been reported to correlate with a wide range of physiological and health benefits other than antioxidative activity.

Polyphenolic compounds have antidiarrhoea properties exhibiting one or more activities against diarrhoea pathogenesis. These may include antiallergenic, antiviral, antibacterial, antifungal (Pietta, 2000), antisecretory, antispasmodic, antimotility (Yue *et al.*, 2004), anti-inflammatory, immunomodulatory and parasitic activities. In traditional medicine preparation of plant extract recipe, water or ethanolic solutions are the main extractants. These extractants extract more or less polar compounds made majorly of phenolic compounds. Specific types of phenolic compounds present in the crude extracts are therefore evaluated.

Flavonoids are C₆-C₃-C₆ polyphenolic compounds present in food, beverage and medicinal plants. They have been reported to have useful pharmacological properties including anti-inflammatory activity, enzyme inhibitors, antiallergic, anti-inflammatory, antiviral, antispasmodic, pro-secretory (Yue *et al.*, 2004) and antimicrobial activity. Flavonoids are known to act on the inflammatory response via many routes and block molecules like COX, iNOS, cytokines, nuclear factor- κ B and matrix metalloproteinases. In addition, flavonoids have good antioxidant, free radical scavengers that donate hydrogen, inhibit lipid peroxidation (Rauha, 2001; Havsteen, 2002) and metal ion chelators. However, the antioxidant power of flavonoids depends on some important structural prerequisites such as the number and the arrangement of hydroxyl groups, the extent of structural conjugation and the presence of electron-donating and electron-accepting substituents on the ring structure (Miliauskas *et al.*, 2005). These groups of phytochemicals are known to play some beneficial roles in the prevention of many oxidative and inflammatory diseases (Arts and Hollman, 2005) inhibiting oxidative and inflammatory enzymes (Middleton *et al.*, 2000).

Gallotannins are complex sugar esters of gallic acid and together with the related sugar esters of ellagic acid (ellagitannins) made up the hydrolysable tannins. Gallotannins exhibit biological activities including antimicrobial, antiviral, anti-inflammatory to anticancer and antiviral properties (Erde'lyi *et al.*, 2005). The mechanisms underlying the anti-inflammatory effect of tannins include the scavenging of radicals, and inhibition of the

expression of inflammatory mediators, such as some cytokines, inducible nitric-oxide synthase, and cyclooxygenase-2 (Polya, 2003; Erde'lyi *et al.*, 2005).

Condensed tannins also referred to as proanthocyanidins are oligomers or polymers essentially derived from flavan-3-ol and their derivatives via carbon to carbon (C–C) or rarely C–O–C links. They differ structurally according to the number of hydroxyl groups on both aromatic rings (ring A and B) and the stereochemistry of the asymmetric carbons of the heterocyclic ring (ring C). Condensed tannins are classified according to their hydroxylation pattern into several subgroups including procyanidins (3,5,7,3',4'-OH), prodelphinidins, (3,5,7,3',4',5'-OH), propelargonidins (3,5,7,4'-OH), profisetinidins (3,7,3',4'-OH), prorobinetinidins (3,7,3',4',5'-OH), proguibourtinidins (3,7,4'-OH), proteracacinidins (3,7,8,4'-OH), and promelacacinidins (3,7,8, 3',4'-OH) (Cos *et al.*, 2003). As with other polyphenols, tannin structures are suitable for free radical scavenging activities serving as an excellent hydrogen or electron donors to form radicals that are relatively stable due to delocalization resulting from resonance and unavailability of site for attack by molecular oxygen (Mello *et al.*, 2005). Tannins can also bind to some free radical producing enzymes forming an insoluble tannin-protein complex (astringent characteristic), complex with catalytic metallic ions making it unavailable to initiate oxidation reaction, and inhibiting lipid peroxidation process (Russo *et al.*, 2005; Mello *et al.*, 2005). These compounds are antagonists of hormone receptors or inhibitors of enzymes such as cyclooxygenase enzymes (Polya, 2003).

Tannins have the ability to protect renal cells against ischemia reperfusion injury (Yokozawa *et al.*, 1997) characterized by an overproduction of $O_2^{\cdot-}$ due to both an electron leak in the mitochondrial respiration chain and the conversion of xanthine dehydrogenase to xanthine oxidase (Wernes and Lucchesi, 1990). The protective action of tannins in this process is related to direct inhibition of enzymatic function of xanthine oxidase activity (Russo *et al.*, 2005).

Production of reactive species (H_2O_2 , $O_2^{\cdot-}$, and OH^{\cdot}) and per-oxy-nitrite occurs at the site of inflammation and contributes to the exacerbation of inflammatory disease and tissue damage. In acute inflammation or chronic inflammations, the production of $O_2^{\cdot-}$ is increased at a rate that overwhelms the capacity of the endogenous SOD enzyme defense to dissipate. Reduction in the $O_2^{\cdot-}$ generation can decrease side-effects of the radical in inflammatory conditions. Tannins have been demonstrated to exhibit anti-inflammatory activity by exerting anti-oxidative properties in reducing $O_2^{\cdot-}$ and malondialdehyde (MDA) production, plasma extravasations and cell migration mainly of leukocytes and potentiates the activity of SOD in radical scavenging (Nardi *et al.*, 2007). It shows that reactive species are most important mediators that provoke or sustain inflammatory processes and consequently, their annihilation by antioxidants and radical scavenger can alleviate inflammation (Delaporte *et al.*, 2002; Geronikaki and Garalas, 2006).

3.9. Conclusion

The extraction methods used optimally extract the phytochemical constituent from the powdered leaves. The extraction process adopted in this work separated the phytochemicals into non-polar hexane portion and polar

water soluble portion in the first step. In addition to taxonomic identification and authentication of medicinal plant, Chemical characterization is also an important and useful means of quality control as it directly correlate with pharmacological functions. The TLC fingerprints revealed the complexity of plant extracts and fractions with chemical compositions of a wide range of polarities. For the optimization of the TLC fingerprinting more than one mobile phase were used to obtain a representative chromatogram of the extracts. In this study, combination of fingerprint with multicomponent quantification of the phenolic compositions was adopted as a good method for chemical profiling of the plants.

There was a strong similarity in the chromatograms of *Erythrina latissima*, *Combretum vendae* and *Combretum woodii*. *Erythrina* and *Combretum* are not closely related and the similarity may be an example of convergent evolution. *Combretum vendae* and *C. woodii* are however, closely related as part of the subgenus *Combretum* and the results indicate the potential use of chemical markers in taxonomy.

CHAPTER FOUR

Antimicrobial activities of the plant extracts against potential diarrhoeal pathogens

4.0. Introduction

Infectious disease defined as an illness caused by a specific pathogen or its toxins that result from transmission of the causative agent or its virulence effectors from an infected person, animal or reservoir to a vulnerable host. The susceptibility of host to infectious pathogens, disease development, progression and severity depends on the age, gender, genetic, immune and nutritional status. Infectious diseases represent a leading cause of morbidity and mortality worldwide despite the advancement in orthodox medicine accounting for more than 26% of all death with developing countries carrying the major burden (Becker *et al.*, 2006). Infections are also considered to be a major contributing factor associated with reduced performance in food animals during growth. Particularly, persistent infections account for slow growth, suboptimal feeding efficiency and economic loss in the livestock industry (Borghetti *et al.*, 2009). The infective pathogens include bacteria, fungi, viruses, protozoa and parasites which manifest their virulence through different mechanisms (see section 2.2 for detailed discussion).

The discovery of antibiotics in 1928 and subsequent development in 1940 as medical treatment provides effective and efficient therapeutic agents for controlling almost all infectious diseases including many feared and contagious infections. Antibiotics are effective in curing many infectious diseases, but they also enhance selection of resistant microbes as some pathogens rapidly became resistant to many of the originally susceptible drugs (Barbour *et al.*, 2004).

At present, the pharmaceutical drugs available to control antibiotic-resistant bacteria are becoming limited. The indiscriminate use and abuse of antibiotics has led to the development of antimicrobial resistance strains and toxicity of some drugs to human and animals (Barton, 2000; Parekh and Chanda, 2007). As a result of these problems, European Union (EU) with EU-directive 1831/2003 imposed ban on the use of antibiotics as growth factor in animal production with effect from 2006 to avoid cross resistance problem with human pathogens and chemical residues in foods (Makkar *et al.*, 2007).

Drug resistance of human and animal pathogenic microbes and parasites has created a serious problem worldwide as previously treatable ailments such as diarrhoea (including dysentery and cholera), and tuberculosis are now more difficult and expensive to treat. The mechanisms of microbial resistance to antibiotic include (Dwyer *et al.*, 2009):

- Genetic alterations which involved the physical exchange of genetic material with another organism (via plasmid conjugation, phage-based transduction, or horizontal transformation), the activation of latent mobile genetic elements (transposons or cryptic genes), and the mutagenesis of its own DNA.

- Chromosomal mutagenesis arises directly from interaction between the chromosome and the antibacterial agent or antibiotic-induced oxidative stress, or indirectly from the bacterium's error prone DNA polymerases during the repair of a broad spectrum of DNA lesions.

The situations have complicated by the treatment of infectious diseases in immunocompromised patients. These negative health trends necessitate for a new prevention and treatment of infectious diseases including diarrhoea.

Medicinal plants have also featured as therapeutic agents used by the world population for basic health care needs and to combat many kinds of infectious diseases worldwide (Voravuthikunchai and Limsuwan, 2006). Medicinal plants have curative properties due to the presence of complex mixture of phytochemicals acting individually or synergistically to exert the associated therapeutic effects. Some of the plant compounds may be novel bioactive substances that can be effective as therapeutic agents for treating ailments such as infectious diarrhoea. These phytochemicals exhibit their antidiarrhoeal effects through various mechanisms such as antimicrobial (Lutherodt *et al.*, 1999), increasing colonic water and electrolytic re-absorption, inhibition of intestinal motility (Oben *et al.*, 2006) and anti secretory effects (Rao *et al.*, 1997). There is considerable research in the screening of natural products from extracts of edible and medicinal plants for the development of alternative drugs to prevent and curtail the emergence of drug-resistance pathogens or other forms of ailments.

4.1. Qualitative antimicrobial (Bioautography) assay

This refers to the direct bioactivity test on developed TLC plates as a means of localizing the biological activity such as microbial growth inhibition, enzymatic inhibition or antioxidative properties of extracts to the particular active compound(s). This helps in focusing attention on the relevant components of an extract (Saxena *et al.*, 1995). Fractionation of medicinal plant extracts in combination with bioautography provides an efficient and relative cheap method for bioactivity-guided isolation of target compound(s) (Hostettman *et al.*, 1997). Practical application of bioautography in activity guided isolation includes enzyme inhibition assay such as the Ellman method for cholinesterase inhibitors (Ellman *et al.*, 1961). In this method, the developed TLC plate is sprayed with a substrate, enzyme and indicator to determine the inhibition by colour variation (white zone against yellow background) (Rhee *et al.*, 2001). In the antimicrobial bioassay, two bioautography methods are available. Firstly, the agar diffusion method involves pouring a layer of inoculated agar solution of the microbes on the developed TLC plate and allowed to set, and the bioactive zone(s) are transferred to the agar gel by diffusion where they can inhibit the growth of the microorganism (Fig 4.1). Secondly, the direct method involves spraying of microorganism broth inocula onto the TLC plate (Homans and Fuchs, 1970) and incubating in humid conditions

to facilitate the growth of the organism.




Fig 4.1. The classification of microbiological methods for biological detection (Adopted from Choma and Grzelak, 2011)

Microbial growth inhibitions are recognized based on the ability of the living microorganism to transform tetrazolium salts to a coloured formazan product. White spots against an intense purple coloured background indicate the compound(s) that kill the tested microorganism (Hostettman and Martson, 2002).

4.2. Quantitative antimicrobial activity (Minimum inhibitory concentration (MIC)) assay

Among the quantitative antimicrobial methods used in evaluating plant extract activity, agar diffusion assay (Greenwood, 1989) and two-fold serial micro-dilution assay (Eloff, 1998) are the most common in phytomedicine research. Sensitivity of the two protocols and their mechanisms varied widely. The mechanism of agar diffusion is the movement of bioactive compounds through the solid agar medium to kill or inhibit the growth of organism it may come in contact with. However, agar diffusion assays may sometimes lead to a false negative result, due to influence of the agar type, salt concentration, incubation temperature molecular size of the antimicrobial components (Greenwood, 1989), and limited diffusion of bioactive component in agar medium. The two-fold serial micro-dilution assay depend on direct contact between the test sample and organism is adjudged to be 30 times more sensitive than the other methods used to screen plant extracts for antimicrobial activity (Eloff, 1998). Although, the effective solubility and miscibility of the bioactive component in the test medium such as the non-polar compounds like terpenes, alkaloid and highly methoxylated phenolics is a limiting factor.

4.3. Selection of microorganisms used in the study

The selection of the microorganisms for antibacterial evaluation in this study was based on their known pathogenic effects in both human and animals with emphasis on diarrhoeal pathogens. Pathogenic *E. coli* has been implicated in diseases such as diarrhoea, hemorrhagic colitis, haemolytic uremic syndrome and thrombocytopenic purpura (Voravuthikunchai and Limsuwan, 2006). *Enterococcus faecalis* has been implicated in causing enteric infection with diarrhoeal effects (Butler, 2006). *Pseudomonas aeruginosa* strains cause diseases such as mastitis, abortions and upper respiratory complications (Masika and Afolayan, 2002). *Staphylococcus aureus* is one of the prominent microbes causing skin infection such as boils, abscesses, carbuncles and sepsis of wounds and it also produces toxins causing diarrhoea and vomiting (Maregesi *et al*,

2008). *Candida albicans* is a typical opportunistic pathogen causing diarrhoea (Gambhir *et al*, 2006), oral and vaginal candidiasis (Shai *et al*, 2008) especially in immunocompromise individuals due to unexpected opportunity by a failure of host defence. *Cryptococcus neoformans* has been implicated in causing life-threatening meningoencephalitis (Xue *et al*, 2007) and pneumonia in immunocompromise individuals (Hamza *et al*, 2006).

4.4. Material and Methods

4.4.1. Microorganism strains

Two standard strains of Gram-positive bacteria (*Staphylococcus aureus* ATCC 29213, *Pseudomonas aeruginosa* ATCC 25922) and two standard strains of Gram-negative bacteria (*Escherichia coli* ATCC 27853, *Enterococcus faecalis* ATCC 29212) were used for antibacterial assay. Three clinical pathogenic fungi namely yeasts (*Candida albicans*, *Cryptococcus neoformans*) and mould (*Aspergillus fumigatus*) (All fungal strains obtained from the Department of Veterinary Tropical Diseases, Faculty of Veterinary Sciences, University of Pretoria) were used.

4.4.2. Culturing of the Bacteria

The bacterial strains were maintained in Mueller Hinton agar (MHA) (Fluka, Spain) while the fungi were maintained in Sabouraud dextrose agar (Merck, Germany) at 4°C under anaerobic conditions. All the organisms were subcultured every 2 weeks. Before testing, the bacterial inoculums were prepared and cultivated in Mueller Hinton broth for 12 h at incubation temperature of 37°C. The fungi inoculums were prepared in Sabouraud dextrose broth (SDB.). The microbial cultures were serially diluted (10 fold increments) in sterile broth to obtain the cell suspension of 1.0×10^5 CFU/ml.

4.4.3. Bioautography against some pathogenic microorganisms

Bioautography was undertaken to ascertain the number of active compound(s) present in crude extracts and fractions. TLC plates were developed as described in section 3.5.4 (Pp 43 - 44), and sprayed with overnight cultures of *E. coli*, *S. aureus*, *P. aeruginosa* or *E. faecalis* and incubated at 37°C for 12-16 h prior to being spray with tetrazolium violet (INT). The inhibitory activity of any components was evident as clear white zones against the purple/red background.

4.4.4. Determination of Minimum Inhibitory Concentration (MIC) against the bacteria pathogens

The minimum inhibitory concentration (MIC) for the crude extract and fractions against bacteria were evaluated using the twofold serial dilution assay with tetrazolium violet added as growth indicator (Eloff, 1998). The extracts (100 µl) at an initial concentration of 1.0×10^4 µg/ml was serially diluted with distilled water up to 50% in 96-well microtitre plate to prepare solution range between 5000 µg/ml first well and 40 µg/ml last well. The bacterial (100 µl) inoculants from 12 h broth cultures (section 4.4.2) diluted to 1:100 were added to each well to obtain final extract concentration range of 2500 µg/ml first well and 20 µg/ml last well. Gentamicin (25 µg/ml first well and 0.18 µg/ml last well) was used as positive control and the solvent used in dissolving the extract was used as

negative control. Final volume in each well was 200 μ l. The plates were incubated for 24 h at 37°C and 100% relative humidity. The inhibition of the bacteria were visualised by adding 40 μ l of aqueous p-iodonitrotetrazolium violet (INT) (Sigma) to each well (concentration 200 μ g/ml). The plates were incubated for another 1 h and MIC was determined as the lowest concentrations of test sample before purple formazan colour were observed.

4.4.5. Determination of Minimum Inhibitory Concentration (MIC) against the fungal pathogens

Minimum inhibitory concentrations (MIC) against three pathogenic fungi were determined using twofold serial dilution assay as described above by Eloff, 1998 with the following modification of Masoko *et al*, 2005. The fungal inoculants (100 μ l) were in fresh Sabouraud dextrose broth and positive controls was amphotericin B (50 μ g/ml first well and 0.4 μ g/ml last well) and negative controls was 70% acetone, final visualization of inhibitory activity was obtained after an incubation for 24 h at 37°C, and 100% relative humidity.

4.5. Results

4.5.1. Microbial bioautography

The TLC bioautography of the crude extracts and fractions of the 27 plant species tested against standard strain bacteria and clinical fungal isolates are presented in Fig 4.2-4.12.




Fig. 4.2. Bioautography of hexane (upper) fraction of different plant species against *S. aureus* (Bab (*Bauhinia bowkeri*), Bag (*Bauhinia galpinii*), Bap (*Bauhinia petersiana*), Cam (*Carissa macrocarpa*), Cop (*Combretum padoides*), Cov (*Combretum vendae*), Cow (*Combretum woodii*), Com (*Commiphora harveyi*), Euc (*Euclea crista*), Eun (*Euclea natalensis*), Erl (*Erythrina latissima*), Fic (*Ficus craterestoma*), Fig (*Ficus glumosa*) developed with hexane: ethyl acetate: formic acid (70:30)

The antimicrobial activities of the extracts were concentrated on the non-polar-enriched hexane fraction while the polar enriched components no sign of microbial inhibition.



Fig. 4.3. Bioautography of dichloromethane fractions of different plant species against *S. aureus* (Bab (*Bauhinia bowkeri*), Bag (*Bauhinia galpinii*), Bap (*Bauhinia petersiana*), Bav (*Bauhinia variegata*), Cam (*Carissa macrocarpa*), Cop (*Combretum padoides*), Cov (*Combretum vendae*), Cow (*Combretum woodii*), Com (*Commiphora harveyi*), Euc (*Euclea crispa*), Eun (*Euclea natalensis*), Erl (*Erythrina latissima*), Fic (*Ficus craterestoma*), Fig (*Ficus glumosa*) developed with chloroform: ethylacetate: formic acid (100:13:10)).




Fig. 4.4. Bioautography of hexane fractions of different plant species against *E. faecalis* (Bab (*Bauhinia bowkeri*), Bag (*Bauhinia galpinii*), Bap (*Bauhinia petersiana*), Cam (*Carissa macrocarpa*), Cop (*Combretum padoides*), Cov (*Combretum vendae*), Cow (*Combretum woodii*), Com (*Commiphora harveyi*), Euc (*Euclea crispa*), Eun (*Euclea natalensis*), Erl (*Erythrina latissima*), Fic (*Ficus craterestoma*), Fig (*Ficus glumosa*) developed with hexane: ethyl acetate: formic acid (70:30)).




Fig. 4.5. Bioautography of dichloromethane fractions of different plant species against *E. coli* (Bab (*Bauhinia bowkeri*), Bag (*Bauhinia galpinii*), Bap (*Bauhinia petersiana*), Bav (*Bauhinia variegata*), Cam (*Carissa macrocarpa*), Cop (*Combretum padoides*), Cov (*Combretum vendae*), Cow (*Combretum woodii*), Com (*Commiphora harveyi*), Euc (*Euclea crispa*), Eun (*Euclea natalensis*), Erl (*Erythrina latissima*), Fic (*Ficus craterestoma*) developed with chloroform: ethylacetate: formic acid (100:13:10)).




Fig. 4.6. Bioautography of dichloromethane fractions of different plant species against *E. faecalis* (Bab (*Bauhinia bowkeri*), Bag (*Bauhinia galpinii*), Bap (*Bauhinia petersiana*), Bav (*Bauhinia variegata*), Cam (*Carissa macrocarpa*), Cop (*Combretum padoides*), Cov (*Combretum vendae*), Cow (*Combretum woodii*), Com (*Commiphora harveyi*), Euc (*Euclea crispa*), Eun (*Euclea natalensis*), Erl (*Erythrina latissima*), Fic (*Ficus craterestoma*), Fig (*Ficus glumosa*) developed with chloroform: ethylacetate: formic acid (100:13:10)).




Fig. 4.7. Bioautography of hexane of different plant species against *C. neoformans* (Ozm (*Ozoroa mucronata*), Ozp (*Ozoroa paniculosa*), Sle (*Searsia leptodictya*), Spd (*Searsia pendulina*), Spt (*Searsia penterii*), Mpd (*Maytenus peduncularis*), Mpr (*Maytenus procumbens*), Mse (*Maytenus senegalensis*), Mun (*Maytenus undata*), Cam (*Carissa macrocarpa*), Com (*Commiphora harveyi*), Syp (*Syzygium paniculatum*)).



Fig. 4.8. Bioautography of dichloromethane fractions against *C. neoformans* (Ozm (*Ozoroa mucronata*), Ozp (*Ozoroa paniculosa*), Sle (*Searsia leptodictya*), Spd (*Searsia pendulina*), Spt (*Searsia penterii*), Mpd (*Maytenus peduncularis*), Mpr (*Maytenus procumbens*), Mse (*Maytenus senegalensis*), Mun (*Maytenus undata*), Cam (*Carissa macrocarpa*), Com (*Commiphora harveyi*), Syp (*Syzygium paniculatum*)).




Fig. 4.9. Bioautography of hexane fractions against *A. fumigatus* (Ozm (*Ozoroa mucronata*), Ozp (*Ozoroa paniculosa*), Sle (*Searsia leptodictya*), Spd (*Searsia pendulina*), Spt (*Searsia penterii*), Mpd (*Maytenus peduncularis*), Mpr (*Maytenus procumbens*), Mse (*Maytenus senegalensis*), Mun (*Maytenus undata*), Cam (*Carissa macrocarpa*), Com (*Commiphora harveyi*), Syp (*Syzygium paniculatum*))




Fig. 4.10. Bioautography of dichloromethane fractions against *A. fumigatus* (Ozm (*Ozoroa mucronata*), Ozp (*Ozoroa paniculosa*), Sle (*Searsia leptodictya*), Spd (*Searsia pendulina*), Spt (*Searsia penterii*), Mpd (*Maytenus peduncularis*), Mpr (*Maytenus procumbens*), Mse (*Maytenus senegalensis*), Mun (*Maytenus undata*), Cam (*Carissa macrocarpa*), Com (*Commiphora harveyi*), Syp (*Syzygium paniculatum*))




Fig. 4.11. Bioautography of hexane fractions against *C. albicans* (Ozm (*Ozoroa mucronata*), Ozp (*Ozoroa paniculosa*), Sle (*Searsia leptodictya*), Spd (*Searsia pendulina*), Spt (*Searsia pentheri*), Mpd (*Maytenus peduncularis*), Mpr (*Maytenus procumbens*), Mse (*Maytenus senegalensis*), Mun (*Maytenus undata*), Cam (*Carissa macrocarpa*), Com (*Commiphora harveyi*), Syp (*Syzygium paniculatum*)).




Fig. 4.12. Bioautography of dichloromethane fractions against *C. albicans* (Ozm (*Ozoroa mucronata*), Ozp (*Ozoroa paniculosa*), Sle (*Searsia leptodictya*), Spd (*Searsia pendulina*), Spt (*Searsia pentheri*), Mpd (*Maytenus peduncularis*), Mpr (*Maytenus procumbens*), Mse (*Maytenus senegalensis*), Mun (*Maytenus undata*), Cam (*Carissa macrocarpa*), Com (*Commiphora harveyi*), Syp (*Syzygium paniculatum*)).

4.5.2. Minimum inhibitory concentration (MIC)

The antibacterial activities of the phenolic-enriched extracts, fractions and sub-fractions of different polarities of 27 plant species evaluated against 4 microorganisms (two gram positive and two gram negative bacteria) are presented in Table 4.1. The results are presented as minimum inhibitory concentrations ($\mu\text{g/ml}$) against tested bacteria and the fungi. The extracts and fractions of the plant species tested exhibited average to good degree of

inhibition against the growth of all the tested bacteria and fungi strains. The extracts and fractions exhibited high potency or growth inhibition at concentration between 19 and >2500 µg/ml for the various different organisms. In this investigation, hexane and dichloromethane fractions had significant and broad-spectrum antimicrobial activities against all tested microbial strains.

The most susceptible bacterium to the crude extract is *E. faecalis* with MIC ranged from 78 to 1250 µg/ml and the least susceptible bacterium to the crude phenolic-enriched extract is *E. coli* with MIC ranging from 312 to 2500 µg/ml. However, all the microorganisms tested are highly susceptible to hexane and dichloromethane fractions with MICs ranging from 19 to 1250 µg/ml. Some of the interesting results include the hexane and dichloromethane fractions of *C. padoides*, *C. vendae*, *C. woodii*, *B. galpinii*, *M. pendularis*, *M. procumbens*, *S. leptodictya* and *S. pendulina* with MICs of 19 - 39 µg/ml against *E. coli*. The hexane and dichloromethane fractions of *C. padoides*, *C. vendae*, *C. woodii*, *M. pendularis*, *M. procumbens*, *M. senegalensis*, *O. mucronata*, *O. paniculosa*, *S. leptodictya* and *S. pentheri* also exhibited good microbial growth inhibitory activity against *E. faecalis* with MICs between 19 - 78 µg/ml.

Growth inhibition activities against *S. aureus* of interest include hexane, dichloromethane, ethyl acetate, butanol fractions of *B. galpinii* and *C. vendae* (MIC 39 - 156 µg/ml); hexane and dichloromethane fractions of *C. padoides*, *O. mucronata* and *S. pentheri* (19 - 78 µg/ml), and hexane, dichloromethane, ethyl acetate fractions of *O. paniculosa*, *S. leptodictya* and *S. pendulina* (19 - 39 µg/ml).

Pseudomonas aeruginosa is also susceptible to growth inhibition by hydrogen and dichloromethane of *C. padoides*, *E. latissima*, *M. pendularis*, *M. procumbens*, and *M. senegalensis* with MICs of 19 - 78 µg/ml; hydrogen, dichloromethane, ethyl acetate *M. undata*, *O. paniculosa*, *S. leptodictya*, *S. pendulina* and *S. pentheri* with MICs of 19 - 78 µg/ml. The reference antibiotic (gentamicin) exhibited good antibacterial activity against the four tested bacterial strains with MIC ranged between 0.18 and 1.56 µg/ml.

Table 4.1. Minimum inhibitory concentration (MIC) of the crude extracts and fraction against *E. coli* and *E. faecalis*

Plant spp	<i>E. coli</i>						<i>E. faecalis</i>					
	CRE	Hf	DCMf	ETOAc	Butanol	Water	CRE	Hf	DCMf	ETO Ac	Butanol	Water
Bab	625	39	156	312	156	>2500	78	312	312	39	78	312
Bag	312	39	156	312	156	312	156	156	39	156	78	312
Bap	312	78	156	312	625	312	78	156	312	78	39	312
Bav	625	78	156	156	156	312	312	312	625	39	39	312
Cam	312	156	156	156	312		312	312	312	78	156	
Cob	312	39	156	312	156	>2500	156	312	312	39	78	312
Cop	156	19	19	312	39	>2500	78	19	19	156	78	312
Cov	312	39	39	156	156	312	156	39	78	78	78	39
Cow	625	39	39	312	156	>2500	156	39	78	78	78	312
Cmh	625	312	39	78	312		156	156	156	78	156	
Erl	1250	156	19	156	156	1250	156	78	19	78	78	1250
Euc	312	625	78	78	156		156	312	156	156	78	
Eun	312	312	312	78	156		625	312	312	156	156	
Fic	2500	312	78	312	312		2500	78	312	312	312	
Fig	1250			1250	312		312			625	156	
Inc	1250				312		1250				312	
Mpd	312	39	39	78	625			39	39	156	1250	
Mpr	1250	39	39	156	1250		>2500	39	39	156	625	
Mse	2500	78	78	156	1250		2500	39	39	312	625	
Mun	1250	312	156	156	>2500		312	156	156	312	2500	
Ozm	1250	78	39	156	312		625	39	19	625	1250	
Ozp	1250	78	39	39	625		1250	39	19	156	78	
Sle	1250	39	39	78	2500		625	39	19	156	625	
Spd	1250	39	39	78	1250		625	156	19	312	625	
Spt	2500	39	39	156	2500		1250	39	19	156	1250	
Scb	625	312	156	39	156	1250	1250	78	312	39	78	2500
Syp	625	312	312	39	156	312	1250	312	156	39	156	1250

Table 4.1. Cont.....Minimum inhibitory concentration (MIC) of the crude extracts and fraction against *S. aureus* and *P. aeruginosa*

Plant spp	<i>S. aureus</i>						<i>P. aeruginosa</i>					
	CRE	Hf	DCMf	ETOAc	Butanol	Water	CRE	Hf	DCMf	ETOAc	Butanol	Water
Bab	625	78	312	625	78	312	312	78	312	156	312	312
Bag	625	39	78	39	78	312	312	312	78	312	156	312
Bap	312	156	312	78	78	78	78	312	625	625	312	625
Bav	625	78	1250	-	-	-	156	78	625	625	312	625
Cam	312	312	156	156	78		312	156	625	312	78	
Cob	625	78	312	156	78	312	312	78	312	625	156	312
Cop	156	19	39	156	156	312	156	78	39	312	19	312
Cov	156	39	78	78	39	39	156	156	156	156	39	312
Cow	312	39	78	156	156	312	625	312	156	312	312	312
Cmh	312	78	156	78	156	2500	1250	312	312	156	312	
Erl	1250	78	312	78	312		2500	78	78	156	78	1250
Euc	156	312	312	78	312		156	312	625	156	39	
Eun	312	312	625	312	312		625	625	625	312	78	
Fic	2500	156	312	1250	1250		2500	156	312	625	156	
Fig	625			312	625		1250			312	78	
Inc	625				156		625				312	
Mpd	>2500	78	39	312	78		312	19	19	156	312	
Mpr	1250	39	78	312	625		312	78	39	156	625	
Mse	625	78	39	156	625		625	39	39	156	1250	
Mun	625	156	156	156	312		312	78	78	78	2500	
Ozm	2500	19	19	625	1250		312	156	39	312	2500	
Ozp	156	19	19	78	156		156	39	39	78	156	
Sle	312	19	19	78	312		625	39	39	39	625	
Spd	312	78	78	39	625		625	78	78	19	2500	
Spt	312	19	19	156	312		156	19	19	78	2500	
Scb	156	1250	156	156	156	1250	156	312	156	156	156	1250
Syp	156	39	156	312	312	2500	156	156	156	78	312	1250

4.5.3. Minimum inhibitory concentration (MIC)

The phenolic-enriched crude extracts and fraction exhibited good to moderate growth inhibitory activities against the three fungal strains of different morphology with MICs ranging from 19 to 2500 µg/ml (Table 4.2). *Candida albicans* demonstrated resistance to all the crude extracts and fractions with the exception of dichloromethane and butanol fractions which had MICs of 19 - 78 µg/ml. In contrast, *Cryptococcus neoformans* was sensitive to majority of many crude extracts and fractions at the concentration ranging from 19 - 78 µg/ml. The fungi were susceptible to amphotericin B with the MIC ranges from 0.78 - 6.25 µg/ml.

Table 4.2. Minimum inhibitory concentration (MIC) of the crude extracts and fraction against *C. albicans*, *C. neoformans* and *A. fumigatus* Values below 100 µg/ml in different colour

Plant spp	<i>C. albicans</i>						<i>C. neoformans</i>						<i>A. fumigatus</i>					
	CRE	H	DCM	ETOAc	But	Water	CRE	H	DCM	ETOAc	But	Water	CRE	H	DCM	ETOAc	But	Water
Bab	625	625	78	156	39	39	156	312	78	78	39	39	2500	156	312	312	156	625
Bag	312	625	78	156	78	625	78	315	78	39	39	312	625	156	156	156	78	156
Bap	1250	156	78	156	156	625	78	625	39	39	156	78	625	312	78	78	156	156
Bav	312	-	-	156	78	625	78	-	-	78	78	78	625			312	156	156
Cam	625	156	625	312	19	156	78	78	156	19	19	1250	625	19	1250	625	312	156
Cob	625	625	312	156	156	1250	78	625	625	39	19	39	312	312	156	156	312	625
Cop	312	312	19	312	156	625	39	156	625	19	312	312	156	156	78	78	156	625
Cov	1250	625	312	78	312	312	78	19	156	78	156	312	156	156	156	156	156	625
Cow	1250	625	19	156	78	625	78	312	312	39	156	312	156	156	78	78	78	625
Cmh	625	156	39	156	39	625	156	39	19	39	19	1250	625	39	19	1250		312
Erl	1250	156	156			625	312	78	19			625	1250	156	156	312		625
Euc	625	156	156	156	78	156	78	156	78	39	39	156	312	625	156	625	156	156
Eun	625	156	78	78	78	625	78	78	39	156	78	625	312	625	625	312	78	312
Fic	2500	78	39	156	39	312	312	78	19	156	39	625	1250	312	156	156	312	156
Fig	625	78	39	78	39	312	78	78	19	78	39	1250	625	78	78	78	625	312
Inc	1250	78	19	156			78	78	39	39			1250	78	156			
Mpd	312	312	78	312	156	312	78	312	78	78	1250	2500	312	78	156	156	312	625
Mpr	1250	156	39		312	625	625	156	39		156	2500	1250	78	78		156	625
Mse	1250	78	39	156	156	625	156	156	39	78	78	1250	625	156	78	156	1250	312
Mun	1250	625	78	312	625	1250	156	1250	78	78	312		1250	625	625	156	625	625
Ozm	625	312	156	156	625	625	156	312	156	39	156	2500	625	39	156	39	156	1250
Ozp	312	156	78	156	312	625	78	312	39	39	156	1250	312	78	78	312	156	625
Sle	312	39	39		312	625	156	156	39		312	1250	625	78	39		156	312
Spd	625	39	78	156	625	625	312	156	39	39	156	625	625	78	39	156	78	312
Spt	2500	625	78	312	156	1250	312	1250	19	39	312	1250	2500	78	19	625	156	625
Scb	625	156	312			625	78	312	78			2500	312	78	19			625
Syp	312	156	156	625	312	625	156	78	39	78	39	1250	312	156	19	1250	2500	312

4.6. Discussion

4.6.1. Antimicrobial bioautography

The crude extracts and various fractions were screened qualitatively for growth inhibitory activity against 4 bacteria and 3 fungi representing different morphologies as yeasts and moulds. Many compounds present in the non-polar enriched hexane fractions inhibited the growth of the organisms tested with several zones of inhibitions.

However, the polar fractions exhibited poor individual inhibitory activities, and this may be due to the high solubility of polar compounds in water, like flavonoids resulting in washing and spread of the compounds on the TLC plate surface. Therefore, reducing the threshold inhibitory concentrations of the bioactive compounds on the spot of the chromatogram against the organisms tested. Other factors may include the disruption of synergistic effects of the individual compounds separated on TLC plates or the concentration of the bioactive components is not sufficient to inhibit microbial growth.

A number of methods have been developed for effective and quick screening of microbial growth inhibitory properties of compounds like the disc or agar diffusion assay adapted as agar-overlay methods ([Rasoanaivo and Ratsimamanga-Urverg, 1993](#)). However, the differential diffusion of the bioactive compounds from the TLC plate to the agar layer make the method unsuitable for certain class of compounds, especially the water-insoluble types like terpenoids and non-polar compounds ([Eloff, 1998](#)). The direct bioautography method allows the localization of a number of components with significant individual inhibitory activities against the tested organisms. The characteristic features of this method are its quickness, efficiency, simplicity, high sample throughput, small test sample size and no sophisticated equipment required. The method is adaptable and applicable to all extracts that can be separated on TLC, against any organism capable of growing directly on TLC plate surfaces.

4.6.2. Minimum inhibitory concentration (MIC)

In this investigation, *in vitro* antimicrobial efficacy of the crude 70% acetone leaf extracts and fractions derived from 27 plants (13 genera across 9 families) used in South African traditional medicine for treating diarrhoea and related ailments was quantitatively assessed on the basis of minimum inhibitory concentration (MIC). All the plants evaluated exhibited varying degree of inhibitory effect against the standard strain of human and animal pathogenic bacteria (Gram-positive as well as Gram-negative) and clinical isolate of pathogenic fungi. There have been no specific cut-off values as a reference or standard for categorizing antimicrobial activity of plant extracts and fractions. In this study, crude extracts and fractions with an MIC value less than 500 µg/ml were considered to have good activity and MIC value less than 100 µg/ml were considered to have significant antimicrobial activity of pharmacological interest according to the criterion by [Rios and Recio \(2005\)](#). A lower MIC values indicated high effectiveness of the compound as antimicrobial agent as little quantity which may be below toxicity level of the extracts can be applied without being harmful to the host.

Crude extracts of 4 out of 27 had an MIC less than a 100 µg/ml (*Bauhinia bowkeri*, *Bauhinia galpinii*, and *Combretum padoides*) against *E. faecalis* and *Bauhinia petersiana* against *P. aeruginosa* (78 µg/ml). However, the antimicrobial activities were potentiated in the fractions with the non-polar fractions of hexane (MIC ranges from 19 to 312 µg/ml) and dichloromethane (MIC ranges from 19 to 625 µg/ml) enriched with terpenoids exhibiting more broad-based potency compared with the polar fractions of ethyl acetate (MIC ranges from 39 to 1250 µg/ml) and butanol (MIC ranges from 39 to <2500 µg/ml). The MIC value less than 100 µg/ml obtained for some fractions were significant although much higher than that of the control antibiotic (gentamicin with an MIC ranged from 0.18 to 1.56 µg/ml against bacteria and amphotericin B with an MIC ranged from 0.78 to 6.25 µg/ml against fungi).

The water fractions have relatively low antimicrobial activities (MIC ranges from 312 to <2500 µg/ml) except for the *C. vendae* with an MIC value 39 µg/ml against *E. faecalis* and *S. aureus*. In traditional medicine plant preparation, water is used as the major extractant. The poor antimicrobial activity of water fractions of most of the plants indicated that decoction or infusion may be less effective in infectious diseases. These observations are consistent with most of the findings in other studies.

From the phytochemical evaluation, the crude extracts contain high level of polyphenolic compounds. The activity of the extracts and polar fractions (ethyl acetate, butanol, and residual), though not exclusive to polyphenolic compounds only would be expected to correlate to the respective constituents, the structural configuration, functional groups and possible synergistic effects among the constituents. Members of this class of compounds are known to have either bacteriostatic or bactericidal properties against most microorganisms depending on the structure and concentration used. The mechanism of their antimicrobial activity may be related to their fundamental properties of having the ability to form complex with protein and polysaccharides, thus the capacity to inactivate microbial adhesions, enzymes, and cell envelope transport protein. The presence and position of hydroxyl group in the phenolic structure determine and influence the antimicrobial activities of this class of compounds (Taguri *et al*, 2004). Phenolic compounds including tannins and flavonoids were found to have high antimicrobial activity (Majhenic *et al*, 2007; Vaquero *et al*, 2007). Some mechanisms of antimicrobial activity of phenolic compounds includes their ability to denature microbial proteins as surface-active agents (Sousa *et al*, 2006), ability to react with cellular membrane component which impairs both function and integrity of cells (Raccach, 1984), and the reducing property of phenolics can influence the redox potential (E_h) of microbial growth causing growth inhibition (Jay, 1996).

However, two methods are widely used in quantitative evaluation antimicrobial activities of plants extracts: Agar disc diffusion method (NCCLS, 2002) and serial dilution method (Eloff, 1998). Both methods depend on the effective solubility of the extracts in the test medium in order to obtain a maximum efficacy against the organisms. However, some phenolic compounds form complex with proteins and other macromolecules present in the test medium, therefore, get precipitated. While some extract components especially the non-polar are not readily soluble in test medium which is more than 90% water in most cases. These factors may at times cause

reduction in the effectiveness of the plant extracts to inhibit microbial growth. The antimicrobial profiles indicated that the extracts and fractions there from were active against Gram-positive and Gram negative bacteria, yeast and mould fungi. The susceptibility of both bacteria and fungi to the extracts may be indicative of the presence of broad-based bioactive compounds or general metabolic toxins.

Pathogenic enteric microorganisms present in contaminated food and water produces enterotoxins or irritants that cause intestinal disorder such as diarrhoea. *In vitro* antimicrobial assays against standard strains of the intestinal pathogens using the polyphenolic-rich crude extracts and fractions have demonstrated various degree of microbial growth inhibition. The plant extracts and fractions investigated have moderate to good activities against diarrhoeal standard strains such as *E. coli*, *S. aureus* and *C. albicans* and *P. aeruginosa*, thus validating their use in traditional medicine for treatment of diarrhoea symptoms. The mechanisms involved in diarrhoea symptoms are multifaceted and interwoven. It is, therefore, possible that extracts and fractions with moderate antimicrobial activities could still have good antidiarrhoeal effects by elaborating other biological activities such as antioxidant, anti-inflammatory, antisecretory, binding of toxins, and antimotility effects on the gastrointestinal tract.

4.7. Conclusion

In infectious diarrhoea many bacteria, protozoa, virus and parasites have been implicated as causative agents. These agents include *Vibrio cholera*, *Escherichia coli*, *Shigella dysenteriae*, *Bacillus cereus*, *Staphylococcus aureus*, *Clostridium difficile*, *Entamoeba histolytica*, *Salmonella typhi* and *Giardia lamblia*. Some viruses such as Rotavirus and adenovirus have also been implicated as causative agent of diarrheal diseases. The infectious mechanisms of the pathogenic strains of the enteric microbes include microbial adhesion and attachment, localized effacement of the epithelial mucosa lining, production and elaboration of secretory enterotoxins, production of cell-destroying cytotoxins, and direct epithelial cell invasion.

In this study, the emphases were on *E. coli*, *S. aureus*, *E. faecalis*, *C. albicans* as diarrhoeal pathogens. Seven virulence groups of diarrheagenic *E. coli*, namely enteropathogenic *E. coli* (EPEC), enterohaemorrhagic *E. coli* (EHEC), enteroinvasive *E. coli* (EIEC), enterotoxigenic *E. coli* (ETEC), enteroaggregative *E. coli* (EAaggEC), diarrhoea-associated haemolytic *E. coli* (DHEC) and cytolethal distending toxin (CDT)-producing *E. coli* have been classified (Clarke, 2001). On a global scale EPEC, EHEC, ETEC and EIEC are the most important diarrhoeal agents accounting for 4-8%, 0-1%, 12-20% and 0-2% respectively in terms of total episodes (Bhan, 2000). The virulence mechanisms of ETEC, EHEC, *S. aureus*, *E. faecalis* and some strains of *V. cholerae* include production of endotoxin, cytotoxins and reactive species. The use of antimicrobial therapy with microbicidal or microbistatic mechanisms may not be effective in the diarrhoea cases involving these organisms because the toxins if already present in contaminated food or water does not need the pathogens to exert activity. Therefore, non-antimicrobial therapy may be required in such cases but antitoxins which can antagonize

toxin and receptor interactions. More work is needed in evaluating the antitoxin and antiadhesion of medicinal plant extracts as other forms anti-infectious mechanisms.

Many of the plant extracts and fractions used have good activities especially the non-polar fractions of hexane and dichloromethane against the pathogens tested, and this may explain the traditional use of these medicinal plants.


Considering importance of oxidative burst such as ROS/RNS in the immune mechanisms and possible consequences of cellular damages, if the resultant oxidative stress is not resolved by the endogenous antioxidant system, the antioxidant potentials of the plants will be evaluated in the next chapter.

CHAPTER FIVE

Free radical scavenging and antioxidant activities of the extracts and fractions as anti-diarrhoeal mechanism.

5.1. Introduction

The intestinal mucosa lining is constantly exposed to the oxidants and toxins from the diet, as well as to endogenous free radicals and other highly reactive species commonly referred to as reactive oxygen species (ROS) or reactive nitrogen species (RNS). These reactive species are generated endogenously in many basic biochemical processes of the body from the respiration (Stojiljkovic *et al.*, 2009) and some cell-mediated immune functions (activated neutrophils and macrophages) in response to microbial infection (Neish, 2009). The enzymatic sources of ROS/RNS include NAD(P)H oxidase, xanthine oxidase, uncoupled endothelial nitric oxide (NO) synthase (eNOS), arachidonic acid metabolizing enzymes such as cytochrome P-450 enzymes, lipoxygenase and cyclooxygenase, and the mitochondrial respiratory chain (Griendling, 2005; Mueller *et al.*, 2005). Exogenous hazards such as exposure to ionizing radiation, smoke and toxins can also generate free radicals (Masoko and Eloff, 2007; Li and Trush, 1994). Free radicals such as trichloromethyl ($\text{CCl}_3\cdot$), superoxide ($\text{O}_2^{\cdot-}$), hydroxyl ($\cdot\text{OH}$), peroxy ($\text{ROO}\cdot$), and nitric oxide ($\text{NO}\cdot$) are produced metabolically in living organisms. In addition, some non-radical derivatives of oxygen molecules (hydrogen peroxide (H_2O_2), hypochlorous acid (HOCl)), are also generated in biological systems. The formation of ROS/RNS have been implicated in the pathogenesis of several human and animal diseases such as atherosclerosis, diabetes mellitus, chronic inflammation, neurodegenerative disorders, gastrointestinal disorders and certain types of cancer (Catalá, 2006). The mechanisms involved in diseases initiation by free radical or oxidative species are outlined in Fig 5.1. In physiological conditions, the epithelial mucosa cell integrity and homeostasis are protected from deleterious effects of ROS by antioxidant defence system consisting of nonenzymatic antioxidants (glutathione (GSH), vitamins A, C, E, carotenoids) and antioxidant enzymes superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPx) and glutathione reductase (GR) (Krishnaiah *et al.*, 2011). However, in pathological conditions excessive oxidation (oxidative stress) in the intestinal tract results in lipid peroxidation of the membrane phospholipids. The peroxidation of membrane phospholipids is basically damaging because the formation of lipid peroxidation products leads to the spread of free radical reactions and cytotoxic aldehydes by-products. The general process of lipid peroxidation consists of three stages: initiation, propagation, and termination (Catalá, 2006). The initiation phase of lipid peroxidation includes hydrogen atom abstraction. Several species can abstract the first hydrogen atom and include the radicals: hydroxyl ($\cdot\text{OH}$), alkoxy ($\text{RO}\cdot$), peroxy ($\text{ROO}\cdot$), and possibly $\text{HO}_2\cdot$ but not H_2O_2 or $\text{O}_2^{\cdot-}$ (Gutteridge, 1988).






Fig. 5.1. Deleterious reactions from the production of reactive free radicals in biological system (amended from Slater *et al.*, 1987)

The initial reaction of $\cdot\text{OH}$ with polyunsaturated fatty acids produces a lipid radical ($\text{L}\cdot$), which in turn reacts with molecular oxygen to form a lipid peroxy radical ($\text{LOO}\cdot$). The $\text{LOO}\cdot$ can abstract hydrogen from an adjacent fatty acid to produce a lipid hydroperoxide (LOOH) and a second lipid radical (Catalá, 2006). The LOOH formed can suffer reductive cleavage in the presence reduced metals, such as Fe^{2+} , producing lipid alkoxy radical ($\text{LO}\cdot$). Both alkoxy and peroxy radicals stimulate the chain reaction of lipid peroxidation by abstracting additional hydrogen atoms (Buettnner, 1993) (See Fig. 2.6 for detailed reaction mechanisms). Peroxidation of lipids can disturb the assembly of the membrane, causing changes in fluidity and permeability, alterations of ion transport and inhibition of metabolic processes (Nigam and Schewe, 2000). Injury to mitochondria induced by lipid peroxidation causes further ROS generation (Green and Reed, 1998). In addition, LOOH break down, frequently in the presence of reduced metals or ascorbate, to reactive aldehyde products, including malondialdehyde (MDA), 4-hydroxy-2-nonenal (HNE), 4-hydroxy-2-hexenal (4-HHE) and acrolein (Esterbauer *et al.*, 1991; Parola *et al.*, 1999; Uchida, 1999; Kehrer and Biswal, 2000; Lee *et al.*, 2001).

Lipid peroxidation and the metabolites are the main oxidative biochemical processes contributing to the disruption of detoxifying pathways in intestine and to dysfunction of enterocytes, which may cause various disorders of digestive tract including diarrhoea. GSH redox cycle plays the main role in lipid peroxidation scavenging in the intestine. Enzyme GPx reduces LOOH using GSH as a reducing factor, while GR regenerates GSH from its oxidized form glutathione disulfide (GSSG), with simultaneous oxidation of nicotinamide adenine dinucleotide phosphate (NADPH). GSH acts not only as an enzyme cofactor, but can react directly with free radicals and is involved in recycling other cellular antioxidants. Excessive generation of ROS/RNS and depleted endogenous antioxidant defences have been implicated in the pathogenesis and perpetuation of intestinal damage which can clinically manifest as diarrhoea. Under disease conditions, more ROS/RNS is generated by the body enhancing the oxidative stress. ROS are also effective in activating redox-responsive pro-inflammatory transcription factors, nuclear factor (NF)- β and activator protein (AP)-1 (Rahman and Adcock, 2006).

Supplementary therapy with antioxidant compounds provides an additional relief against deleterious effect of ROS/RNS.

5.1.1. Superoxide ion

Superoxide radical anion ($O_2^{\cdot-}$) generated from an electron leakage in the mitochondrial respiration chain and the conversion of xanthine dehydrogenase to xanthine oxidase (Wernes and Lucchesi, 1990) as a result of electron donation to oxygen molecule is regarded as the primary ROS in biological system. Although $O_2^{\cdot-}$ is not very active but the radical interact with other molecules to produce highly potent secondary ROS either directly or indirectly through enzyme and/or metal catalyzed mechanisms (Valko *et al.*, 2005). In acute inflammation or chronic inflammations, the production of $O_2^{\cdot-}$ is increased at a rate that overwhelms the capacity of the endogenous SOD enzyme defence to dissipate.

5.1.2. Hydrogen peroxide

The generation of hydrogen peroxide (H_2O_2) by activated phagocytes plays an important part in the killing of several bacterial and fungal strains (Sanchez-Moreno, 2002). Additionally, H_2O_2 is generated *in vivo* under physiological conditions by peroxisomes and several oxidative enzymes including glucose oxidase, d-amino acid oxidase, and dismutation of superoxide radical, catalysed by superoxide dismutase. There is increasing evidence that H_2O_2 , either directly or indirectly via its reduction product (hydroxyl ion ($OH\cdot$)), acts as a messenger molecule in the synthesis and activation of inflammatory mediators (Auroma *et al.*, 1989).

5.1.3. Hydroxyl radical

Hydroxyl radical ($\cdot OH$) is the neutral form of hydroxyl ion and the most reactive free radical in biological systems generated from free metal ions (copper or iron) catalyzed breakdown of H_2O_2 (Fenton reaction) or superoxide ion reaction with H_2O_2 (Haber-Weiss reaction, Fig 2.6). Hydroxyl radicals have short half of 10^{-9} s with the highest 1-electron reduction potential of 2310 mV, and is primarily responsible for the cytotoxic effect in aerobic organism. The radical reacts with every cell components in living organisms at the second-order rate constants of 10^9 – 10^{10} mol/s (Siddhuraju and Becker, 2007) such as lipid, polypeptides, proteins, and DNA, especially thiamine and guanosine. Unlike $O_2^{\cdot-}$ and H_2O_2 , which can be enzymatically eradicated by the activity of superoxide dismutases ($2O_2^{\cdot-} + 2H^+ \rightarrow H_2O_2 + O_2$) and catalases/peroxidises ($2H_2O_2 \rightarrow 2H_2O + O_2$), respectively, there exists no known enzyme that catalyzes the cellular detoxification of $\cdot OH$ (Dwyer *et al.*, 2009).

5.1.4. Peroxyl radical

Peroxyl radicals are important reactive species in living systems formed by a direct reaction of oxygen with alkyl radicals or the protonation of the superoxide ions. Peroxyl radicals are potent oxidants with standard reduction potential of more than 1000 mV (Decker, 1998). The radicals abstract hydrogen from other molecules with lower standard reduction potential to perpetuate chain reaction such as propagation stage of lipid peroxidation. Cell

membranes including intestinal epithelial mucosa are phospholipid bilayers with extrinsic proteins and are the primary target of lipid peroxidation (Girotti, 1998) causing cell dysfunction and tissue injury. Lipid peroxidation cytotoxic by-products such as malonaldehyde can react with free amino group of proteins, phospholipid, and nucleic acids leading to structural modification, which induce dysfunction of immune systems.


5.1.5. Hypochlorous acid

Hypochlorous acid (HOCl) is also a strong oxidant generated *in vivo* by neutrophil myeloperoxidase (MPO) catalyzed oxidation of chloride ions and H₂O₂. The cytotoxicity of this reaction contributes to the phagocytosis of infectious microorganisms in the host defence system. However, HOCl generated by MPO also inactivate some enzymes such as α -antiproteinase contributing to proteolytic damage of healthy human tissues in inflammatory disease (Halliwell and Gutteridge, 1990; Hippeli and Elstner, 1999). The oxidant has also been implicated as a secretagogue.

5.1.6. Nitric oxide

Biological tissues generate nitric oxide (NO[•]) by specific nitric oxide synthases (eNOS, iNOS) metabolization of arginine to citrulline via a five electron oxidative mechanisms. In normal physiological processes, nitric oxide (NO[•]) acts as an important oxidative biological signalling molecule in neurotransmission, blood pressure regulation, defence mechanisms, smooth muscle relaxation and immune regulation (Bergendi *et al.*, 1999). Nitric oxide (NO[•]) has greater stability in an environment with a lower oxygen concentration compared to the hydroxyl radical with half life >15 s.

Cells of the immune system produce both the superoxide anion and nitric oxide in the oxidative burst inducing inflammatory processes. Under these conditions, nitric oxide and the superoxide anion may react together to produce significant amounts highly reactive oxidative molecule (peroxynitrite anion (ONOO⁻)). This potent oxidising agent that can cause DNA fragmentation and initiate lipid peroxidation (Carr *et al.*, 2000):



The NO[•] toxicity is predominantly linked to its ability to combine with superoxide anions with the rate constants known for reactions of NO[•], 7.0×10⁹M⁻¹ s⁻¹.

5.2. Antioxidant assays

Several standardized methods have been proposed to analyze the antioxidant potential of a substrate including plant extract and isolated compounds from it. Criteria for the standard methods include (i) measurement of the chemical process actually occurring in potential applications; (ii) utilization of biological relevant molecules; (iii) technically simple; (iv) with a defined endpoint and chemical mechanism; (v) readily available instrumentation; (vi) good repeatability and reproducibility; (vii) adaptable for assay of both hydrophilic and lipophilic antioxidants; (viii) and adaptable to high-throughput analysis (Prior, *et al.*, 2005). The assays are based on scavenging capacity against specific biological ROS/RNS and/or against stable, non-biological radicals and evaluation of

total reduction capacity such as 1,1-diphenyl-2-picrylhydrazine (DPPH) radical scavenging assay (Brand-Williams *et al.*, 1995); 2,2'-azinobis(3-ethylbenzthiazoline-6-sulphonic acid) (ABTS) radical scavenging method (Re *et al.*, 1999); β -carotene linoleic acid bleaching assay (Siddhuraju and Becker, 2003); inhibition of linoleic acid peroxidation (Osawa and Namiki, 1981); ferric reducing antioxidant power (FRAP) (Benzie and Szeto, 1999); total radical trapping antioxidant potential (TRAP) assay (Leontowicz *et al.*, 2002); hydroxyl radical scavenging activity (Jodynis-Liebert *et al.*, 1999); hydrogen peroxide scavenging activity (Ruch *et al.*, 1989); nitro blue tetrazolium (NBT) reduction assay or superoxide anion scavenging activity (Kirby and Schmidt, 1997) and oxygen radical absorbance capacity (ORAC) assay (Silva *et al.*, 2007). Each method has its own merit and demerit in evaluating antioxidant capacity of plant extracts and their components. Based on the criteria enumerated above, the most common and reliable methods are the ABTS and DPPH methods.

5.2.1. Antioxidant bioautography

For the qualitative detection of free radical scavengers and the number of antioxidant compounds present, DPPH or β -carotene is usually the spraying reagents (Martson, 2011) of TLC chromatograms. DPPH is a purple-coloured free radical that turns yellow on reduction by an antioxidative component of an extract. Yellow spots on TLC plates sprayed with DPPH solution against the purple background indicate the presence of an active compound(s). In the β -carotene assay, the TLC plate is sprayed with a solution of β -carotene, dried and exposed to 254 nm UV light to bleach the β -carotene. Areas where antioxidants inhibit degradation of β -carotene appear as orange zones on a pale background.

5.2.2. The chemistry of some common antioxidant assays

5.2.2.1. Hydroxyl radical

Hydroxyl radical is the most reactive species and source of many other secondary free radicals in biological systems; thus, it is important to evaluate hydroxyl radical scavenging capability. Hydroxyl radical (HO^\bullet) scavenging is usually evaluated using the "deoxyribose assay": a mixture of ferric chloride (FeCl_3) and ethylenediamine tetraacetic acid (EDTA) in the presence of ascorbate reacts to form iron(II)-EDTA plus oxidized ascorbate, H_2O_2 then reacts with iron(II)-EDTA to generate iron(III)-EDTA plus HO^\bullet from the Fenton reaction ($\text{Fe}_2 + \text{H}_2\text{O}_2 \rightarrow \text{Fe}_3 + \text{HO}^\bullet + \text{HO}^-$). The radicals not scavenged by other components of the reaction mixture attack the sugar deoxyribose, and degrade it into a series of fragments, some or all of which react on heating with thiobarbituric acid at low pH to give a pink chromogen. Thus the scavenging activity towards HO^\bullet of a substance added to the reaction mixture is measured on the basis of the inhibition of the degradation of deoxyribose.

Another spectrophotometric method developed for assessment of hydroxyl radical scavenging capacity of antioxidants includes Fenton reaction as the hydroxyl radical generation system and salicylate as a spectrophotometric indicator. Attack by OH^\bullet radicals on salicylate produce 2, 3-dihydroxybenzoate, 2, 4-dihydroxybenzoate, and 2, 5-dihydroxybenzoate as major products. The hydroxylated products can be identified and quantified by Beer's law testing the additivity of absorbances of the hydroxybenzoates. This method is able

to measure the hydroxyl radical scavenging capability of individual antioxidants with a wide dynamic activity range, i.e., 635-637 nm. These spectrophotometric methods may not be the most sensitive, but they are simple, reproducible, and cost effective method valuable in antioxidant studies.

5.2.2.2. Hydrogen peroxide scavenging

Hydrogen peroxide-scavenging activity is measured by using a peroxidase-based assay system. The most common used peroxidase is horseradish, which uses H_2O_2 to oxidize scopoletin into a nonfluorescent product. In the presence of a putative scavenger, the oxidation of scopoletin is inhibited and the H_2O_2 scavenging can be monitored by decay in H_2O_2 concentration spectrophotometrically from absorption at 230 nm using the molar optical density of $81 M^{-1}cm^{-1}$.

5.2.2.3. Superoxide scavenging capacity

The scavenging activity towards superoxide by antioxidants is measured by the inhibition of generation of superoxide with the hypoxanthine-xanthine oxidase (HX-XO) system. The superoxide generating from HX-XO reduces nitro-blue tetrazolium (NBT) to formazan at pH 7.4 and room temperature which can be followed spectrophotometrically at 560 nm. Any added molecule capable of reacting with superoxide inhibits the production of formazan and the reduction of the absorbance is estimated as superoxide scavenging activity compared to the value obtained with no test added sample.

5.2.2.4. 2, 2-diphenyl-1-picrylhydrazyl radical (DPPH[•] assay)

The DPPH radical scavenging assay is hydrogen atom transfer processes widely used evaluate the antioxidant activity of reductants (plant extracts, phytochemical or pharmaceutical drugs) (Kaviarasan et al., 2007). Although DPPH assay has no direct biological relevance, the process is related to the inhibition of lipid peroxidation (Rekka and Kourounakis, 1991). The DPPH free radical is reduced to the corresponding hydrazine when it reacts with hydrogen donors (antioxidant) using decolouration mechanisms (purple to yellow), which are monitored by the decrease in absorbance at 515–528 nm. From the methodological point of view the assay conducted in ethanol or methanol solution of DPPH is considered a valid, easy and accurate assay to evaluate radical scavenging activity of antioxidants, since the radical compound is stable and does not have to be generated as in other radical scavenging assays. The results are highly reproducible and comparable to other free radical scavenging methods such as ABTS.

5.2.2.5. 2, 2'-azinobis-(3-ethylbenzothiazoline-6-sulphonate) radical cation (ABTS^{•+}) free radical-scavenging method


The ABTS radical scavenging method is based on the reduction of blue/green ABTS^{•+} chromophore generated from the reaction between ABTS and potassium persulphate ($K_2S_2O_7$) by an electron-donating antioxidant. The

decolourization of the ABTS^{•+} chromophore is measured spectrophotometrically at 734 nm in both lipophilic and hydrophilic medium.



5.2.2.6. Ferric reducing antioxidant power (FRAP)

The FRAP method is based on the reduction at low pH, of a colourless ferric complex to an intense blue-coloured ferrous complex by an electron donating antioxidant. The reduction of ferric complex is monitored by measuring the increase in absorbance at 750 nm. The chemistry of FRAP assay can be summarized with equation (1) with oxidant accepting an electron from antioxidant to be reduced an intense coloured molecule.



The FRAP assay is a robust and potentially useful test using inexpensive reagents and equipment and a speedy reaction applicable over a wide concentration range.

5.3. Materials and Methods

5.3.1. Qualitative antioxidant assay using TLC-DPPH method

This was done to determine the number of active compound(s) present in the crude extracts and fraction(s) with free radical scavenging capacity against DPPH radical. Chromatograms were developed as described in section 3.6.4 and sprayed with 0.2% methanolic DPPH solution. The presence of yellow spots against a purple background indicated component(s) with antioxidant activities.

5.3.2.1. DPPH[•] radical-scavenging assay

The antioxidant activities of the samples were measured in term of radical scavenging ability using the stable radical (DPPH[•]) of [Brand-Williams *et al.*, \(1994\)](#) with some modifications. Methanol solutions (40 µl) of the samples at various concentrations (19–2000 µg/ml), and positive control (trolox and ascorbic acid) at concentration (19 to 250 µg/ml) were added to 160 µl of DPPH in methanol (25 µg/ml) in a 96 well-microtitre plate. The change in absorbance (516 nm) measured after at 1, 10, 20 and 30 min (A_t) with a microtitre plate reader (Versamax). The sample concentrations were corrected for the dilutions.

5.3.2.2. ABTS^{•+} radical-scavenging assay

The free radical-scavenging activity as a measure of hydrogen donating capacity was determined by using ABTS cation decolourization method of [Re *et al.*, 1999](#) with some modifications. ABTS radical solution (7 µM) was prepared by dissolving 1.32×10^4 µg of ABTS in 10 ml of 50% methanolic solution and 7.68×10^4 µg of potassium persulphate ($\text{K}_2\text{S}_2\text{O}_8$) in 10 ml of distilled water. The two solutions were mixed together and made up to 200 ml with 50% methanolic solution, and kept in the dark at room temperature for 12 h. Prior to running the assay, the ABTS radical solution was diluted with a 50% methanolic solution to an absorbance (A_{62}) between

0.7- 0.8 at 734 nm. The extracts were serially diluted (40 μ l) (19 to 2000 μ g/ml) in 96 well-microtitre plate and 160 μ l of ABTS radical solution added to each well. The absorbance were taken exactly after 6 min of reaction (A_{t2}) and blank absorbance (A_{b2}) were prepared using the respective extracts without ABTS radical.

5.3.2.3. FRAP assay

The FRAP of the samples was determined by direct reduction of potassium ferric cyanide ($K_3Fe_3(CN)_6$) to potassium ferrocyanide ($K_3Fe_2(CN)_6$) (electron transfer process from the antioxidant). The increase in absorbance from the formation of Pearl's Prussian blue complex following the addition of excess ferric ion was measured as described by [Berker *et al.*, \(2007\)](#) with some modification. The reaction medium (200 μ l) containing 40 μ l of the test samples or positive controls (trolox and ascorbic acid) (concentration range between 19 -2000 μ g/ml), 100 μ l of 1.0M hydrochloric acid, 20 μ l of 1% (w/v) of SDS, 30 μ l of 1% (w/v) of potassium ferric cyanide was incubated for 20 min at 50°C, cooled to room temperature and finally 20 μ l of 0.1% (w/v) of ferric chloride was added. The absorbance at 750 nm was read and blank absorbance was taken by preparing the reaction medium the same way except the addition of ferric chloride. The reducing capacities were taken as slope obtained from the line of best fit of the absorbance against concentration using the linear regression curve.

5.3.2.4. Hydroxyl radical scavenging assay

The hydroxyl radical scavenging activities of the test samples were measured by the salicylic acid method. The hydroxyl radical scavenging activity of the extracts was determined according to method of [Smirnoff and Cumbes, \(1996\)](#) with some modifications. The hydroxyl radical was generated by using Fenton reaction which contains 50 ml of $FeCl_3$ (8.0 mM), 80 ml of H_2O_2 and 50 ml of distilled water was allowed to stand for 1h. The mixture was filtered to remove the debris. Hydroxyl radical was determined by mixing 120 μ l of the hydroxyl radical solution with 66 μ l of the extracts followed by 14 μ l of salicylic acid (20 mM). The mixture was incubated for 30min at 37°C and absorbance taken at 510 nm.

5.3.2.5. Lipid peroxidation inhibition assay

Lipid peroxidation of linoleic acid was determined as described by [Kishida *et al.*, 1993](#). Each reaction mixture contained 4.1 ml of 2.5% linoleic acid in ethanol and 10 ml of 0.2M phosphate buffers (pH 7.4), 1.0 ml of 0.0025 mg/ml $FeSO_4$ was added as catalyst. Different concentrations of the samples (10-500 μ g/ml) were added to the reaction mixture in a centrifuge tube. The reaction mixture was incubated at 40°C for 2 h. The reaction was terminated by adding 1.0 ml of 25% trichloroacetic acid and 1.0 ml of 0.67% thiobarbituric acid. The reaction mixture was heated at 95°C for 30 min and cooled for 15 min. The mixture was extracted with butanol and absorbance measured at 500 nm.

5.4. Results

5.4.1. TLC-DPPH analyses (Antioxidant bioautography)

The qualitative antioxidant screening of spraying DPPH on TLC plate indicated the presence of a number of antioxidant compounds in the crude extracts and fractions (hexane, dichloromethane ethyl acetate, and butanol). Antioxidant compounds were visualized as yellow spot against the purple background of DPPH as shown in Fig. 5.2 - 5.5. The numbers of active compounds identifiable on a plate depend on the mobile phase used in the development of the plate. For the crude extracts, solvents 1 and 2 separated the antioxidative compounds but solvent 3 revealed close similarity between *Combretum vendae* and *Combretum woodii*, both exhibiting activities at three different spots of same with R_f values of 0.94, 0.53 and 0.48 respectively.




Fig.5.2. TLC-DPPH profiles of the crude extracts of extracts of different plants (left to right Bab (*Bauhinia bowkeri*), Bag (*Bauhinia galpinii*), Bap (*Bauhinia petersiana*), Bav (*Bauhinia variegata*), Erl (*Erythrina latissima*), Inc (*Indigofera cylindrica*), Scb (*Schotia brachypetala*), Cob (*Combretum bracteosum*), Cop (*Combretum padoides*), Cov (*Combretum vendae*), Cow (*Combretum woodii*), Ozm (*Ozoroa mucronata*), Ozp (*Ozoroa paniculosa*), Sle (*Searsia leptodictya*), Spd (*Searsia pendulina*), Spt (*Searsia pentheri*), Mpd (*Maytenus peduncularis*), Mpr (*Maytenus procumbens*), Mse (*Maytenus senegalensis*), Mun (*Maytenus undata*), Cam (*Carissa macrocarpa*), Com (*Commiphora harveyi*), Syp (*Syzygium paniculatum*)) developed with ethyl acetate: acetic acid: formic acid: water (75:5:5:10) (top), ethyl acetate: acetic acid: formic acid: water (70:5:15:10) (middle) and ethyl acetate: methanol: ammonia (90:20:15) (bottom)).

For the hexane fraction using eluent 5, the antioxidant compound was present only in *Erythrina latissima* sample with R_f value of 0.26. The dichloromethane fraction of the extract exhibited antioxidant activity at many spots with some being minor while *E. latissima*, *C. vendae*, and *C. woodii* revealed major antioxidant spots at R_f values of 0.6, 0.46 and 0.33 using solvent 5. Spots at R_f values of 0.53 and 0.40 were peculiar to *C. vendae*, and *C. woodii* respectively. Eluent 6 separated more antioxidant components in all the plants tested with *Bauhinia bowkeri*, *E.*

latissima, *C. vendae*, and *C. woodii* exhibiting activity at 0.8, *E. latissima*, *C. vendae*, and *C. woodii* at 0.66, *B. bowkeri*, *B. galpinii*, *B. petersiana*, *B. variegata*, *E. latissima*, *C. vendae*, and *C. woodii* at 0.51, *C. vendae* and *C. woodii* at 0.17 and *C. vendae* at 0.1.




Fig. 5.3. TLC-DPPH profile of the hexane fractions of different plants (Bab (*Bauhinia bowkeri*), Bag (*Bauhinia galpinii*), Bap (*Bauhinia petersiana*), Bav (*Bauhinia variegata*), Erl (*Erythrina latissima*), Inc (*Indigofera cylindrica*), Scb (*Schotia brachypetala*), Cob (*Combretum bracteosum*), Cop (*Combretum padoides*), Cov (*Combretum vendae*), and Cow (*Combretum woodii*) developed with hexane: ethyl acetate: formic acid (70:30:2))




Fig.5.4 TLC-DPPH profiles of the dichloromethane fractions of different plants (left to right Bab (*Bauhinia bowkeri*), Bag (*Bauhinia galpinii*), Bap (*Bauhinia petersiana*), Bav (*Bauhinia variegata*), Erl (*Erythrina latissima*), Inc (*Indigofera cylindrica*), Scb (*Schotia brachypetala*), Cob (*Combretum bracteosum*), Cop (*Combretum padoides*), Cov (*Combretum vendae*), Cow (*Combretum woodii*), Ozm (*Ozoroa mucronata*), Ozp (*Ozoroa paniculosa*), Sle (*Searsia leptodictya*), Spd (*Searsia pendulina*), Spt (*Searsia pentheri*), Mpd (*Maytenus peduncularis*), Mpr (*Maytenus procumbens*), Mse (*Maytenus senegalensis*), Mun (*Maytenus undata*), Cam (*Carissa macrocarpa*), Com (*Commiphora harveyi*), Syp (*Syzygium paniculatum*) developed with hexane:ethyl acetate: formic acid (70:30:2) top, hexane:ethyl acetate: formic acid (50:50:2) (middle) and chloroform:ethyl acetate:formic acid (50:40:10) bottom.




Fig. 5.5. TLC-DPPH profiles of the ethyl acetate fractions of different plants left to right (left to right Bab (*Bauhinia bowkeri*), Bag (*Bauhinia galpinii*), Bap (*Bauhinia petersiana*), Bav (*Bauhinia variegata*), Erl (*Erythrina latissima*), Inc (*Indigofera cylindrica*), Scb (*Schotia brachypetala*), Cob (*Combretum bracteosum*), Cop (*Combretum padoides*), Cov (*Combretum vendae*), Cow (*Combretum woodii*), Ozm (*Ozoroa mucronata*), Ozp (*Ozoroa paniculosa*), Sle (*Searsia leptodictya*), Spd (*Searsia pendulina*), Spt (*Searsia pentheri*), Mpd (*Maytenus peduncularis*), Mpr (*Maytenus procumbens*), Mse (*Maytenus senegalensis*), Mun (*Maytenus undata*), Cam (*Carissa macrocarpa*), Com (*Commiphora harveyi*), Syp (*Syzygium paniculatum*) developed with chloroform:ethyl acetate:formic acid (50:40:10) (top), hexane: ethyl acetate:formic acid (10:90:10) (middle) and ethyl acetate:methanol:water:formic acid (100:13:10:2) (bottom).

5.4.2. Effective concentration required to reduce DPPH radical (oxidant) by half (EC₅₀)

The phenolic-enriched crude extracts and fractions exhibited strong radical scavenging activity against DPPH radicals in a dose dependent manner (Table 5.1). There are significant variations in the capacity of the test samples to scavenge the DPPH radical with EC₅₀ ranging from 0.21±0.03 to 303.65±3.84 µg/ml. Butanol fractions of *Combretum padoides* had the highest anti DPPH radical activities compared to crude extracts and other fractions with EC₅₀ 0.21±0.03 µg/ml followed by butanol fractions of *Combretum vendae* and *Combretum woodii* with EC₅₀ 0.25±0.06 and 0.33±0.01 µg/ml respectively. The EC₅₀ of these fractions are lower than the EC₅₀ of the positive controls (trolox 1.18±0.06 – 1.31±0.07 µg/ml and ascorbic acid 1.50±0.06 – 1.68±0.07 µg/ml). As expected the non-polar compounds present in the hexane and dichloromethane fractions had poor radical scavenging activities compared to the controls.

Table 5.1. DPPH radical scavenging potential of the crude extract and fractions expressed as EC₅₀ (µg/ml)

Plant species	Crude	Hexane	DCM	ETOAc	Butanol	Residual Water
Bab	19.53±4.83 ^a	11.14±3.59 ^b	5.21±1.04 ^{b,c}	1.25±0.23 ^c	0.64±0.05 ^c	4.99±0.56 ^c
Bag	14.39±0.48 ^a	79.58±13.14 ^b	9.92±1.16 ^{a,c}	2.82±0.44 ^{a,c}	1.02±0.06 ^c	2.86±0.42 ^{a,c}
Bap	43.29±5.05 ^a	47.45±2.91 ^a	8.18±1.11 ^b	3.21±1.01 ^{b,c}	1.51±0.07 ^c	15.20±1.66 ^d
Bav	123.60±11.05 ^a	97.02±30.03 ^a	8.40±0.62 ^b	1.88±0.10 ^b	0.89±0.05 ^b	23.07±3.83 ^c
Erl	2.54±1.40	76.71±20.25	6.02±2.0	5.61±0.37	2.24±0.05	57.98±13.94
Inc						
Scb						
Cob	5.72±1.21	85.04±10.56	20.53±0.40	7.76±0.68	4.97±0.19	35.65±4.78
Cop	4.44±0.35	12.65±1.3	3.33±0.30	0.44±0.06	0.21±0.03	0.84±0.15
Cov	1.65±0.20	16.88±2.66		1.02±0.14	0.25±0.06	0.96±0.12
Cow	3.88±1.78	9.41±3.51	2.16±0.284	1.24±0.13	0.33±0.01	1.10±0.07
Ozm	15.82±4.02	132.13±5.0	63.48±4.00	7.92±1.63	-	-
Ozp	1.29±0.07	31.95±5.6	9.77±0.71	1.22±0.44	11.79±1.12	-
Sle	1.81±0.09	54.88±2.53	10.14±1.51	0.91±0.04	38.93±0.28	-
Spd	1.19±0.15	138.5±9.50	16.51±0.77	1.26±0.03	41.8±3.37	-
Spt	4.26±0.40	139.63±10.62	4.91±0.69	2.09±0.32	-	-
Mpd	3.81±0.03	113.4±12.60	28.20±4.14	2.33±0.21	29.95±4.01	76.79±10.30
Mpr	7.39±0.32	111.2±10.69	30.80±4.56	2.52±0.30	20.71±0.90	189.50±7.56
Mse	13.46±0.52	253.0±29.69	121.46±11.0	4.73±0.06	24.01±0.13	81.17±11.39
Mun	6.99±0.14	160.4±31.4	42.88±6.16	2.23±0.15	7.86±15	303.67±3.84
Euc	3.00±0.37	134.46±10.8	4.70±0.72	0.84±0.00	0.91±0.19	2.04±0.24
Eun		84.88±9.74	7.23±4.40	1.66±0.22	1.34±0.00	2.62±1.19
Fic						
Fig						
Cam	5.85±0.55	138.46±35.73	11.03±0.77	1.85±0.27	2.55±0.23	10.4±2.36
Com	19.31±1.40	103.22±30.37	17.08±0.27	0.90±0.00	1.13±0.22	310.53±12.0
Syp						

Bab (*Bauhinia bowkeri*), Bag (*Bauhinia galpinii*), Bap (*Bauhinia petersiana*), Bav (*Bauhinia variegata*), Cam (*Carissa macrocarpa*), Cob (*Combretum bracteosum*), Cop (*Combretum padoides*), Cov (*Combretum vendae*), Cow (*Combretum woodii*), Com (*Commiphora harveyi*), Erl (*Erythrina latissima*), Euc, (*Euclea crispa*), Eun (*Euclea natalensis*), Fic (*Ficus craterostoma*), Fig (*Ficus glumosa*), Inc (*Indigofera cylindrica*), Mpd (*Maytenus peduncularis*), Mpr (*Maytenus procumbens*), Mse (*Maytenus senegalensis*), Mun (*Maytenus undata*), Ozm (*Ozoroa mucronata*), Ozp (*Ozoroa paniculosa*), Sle (*Searsia leptodictya*), Spd (*Searsia pendulina*), Spt (*Searsia pentheri*), Scb (*Schotia brachypetala*), Syp (*Syzygium paniculatum*)

Surprisingly the residual water fractions of many of the plant species also had a low antioxidant activity this may be due to the insolubility of the dried water fraction. If polyphenolics reacted with e.g. sugars to form insoluble complexes it would explain the results. The least active samples are the residual water fractions of *C. harveyi* (310.53±12.00 µg/ml) and *Maytenus undata* (303.61±3.84 µg/ml).

Other fractions with notable antioxidant activity were the butanol fractions of *Bauhinia bowkeri* (0.64±0.05 µg/ml), *Bauhinia galpinii* (1.02±0.06 µg/ml), *Bauhinia variegata* (1.51±0.07 µg/ml) and *Commiphora harveyi* (1.13±0.22

$\mu\text{g/ml}$); the ethyl acetate fractions of *C. padoides* ($0.44\pm 0.06 \mu\text{g/ml}$), *C. vendae* ($1.02\pm 0.14 \mu\text{g/ml}$), *Ozoroa paniculosa* ($1.22\pm 0.44 \mu\text{g/ml}$), *Searsia leptodictya* ($0.91\pm 0.04 \mu\text{g/ml}$), *C. harveyi* ($0.90\pm 0.00 \mu\text{g/ml}$); and residual water fractions *C. padoides* ($0.84\pm 0.15 \mu\text{g/ml}$), *C. vendae* ($0.96\pm 0.12 \mu\text{g/ml}$) and *C. woodii* ($1.10\pm 0.07 \mu\text{g/ml}$). The ethyl acetate fractions of all the 27 plant samples exhibited good antiradical activities against DPPH radical with EC_{50} ranging between 0.44 ± 0.06 (*C. padoides*) – $7.92\pm 1.63 \mu\text{g/ml}$ (*Ozoroa mucronata*). From the estimated EC_{50} values, the order of potency is butanol fraction > ethyl acetate fraction > crude extract > dichloromethane fraction > residual water fraction > hexane fraction.

5.4.3. Effective concentration required to reduce ABTS radical (oxidant) by half (EC_{50})

The ABTS^{•+} radical scavenging capacity of the crude extract and the fractions expressed as EC_{50} are presented in Table 5.2 with lower EC_{50} indicate higher antiradical activity. A wide variation in the ABTS^{•+} radical scavenging capacity of the crude extracts and the fractions which range from 0.43 ± 0.03 to $1709 \pm 91.44 \mu\text{g/ml}$.

The same trend of DPPH radical scavenging activity is also noticeable with the ABTS radical scavenging assay though the EC_{50} are slightly higher. Butanol fractions of *C. padoides* had the highest anti DPPH radical activities compared with crude extracts and other fractions with EC_{50} $0.21\pm 0.03 \mu\text{g/ml}$ followed by butanol fractions of *C. vendae* and *C. woodii* with EC_{50} 0.25 ± 0.06 and $0.33\pm 0.01 \mu\text{g/ml}$ respectively. The EC_{50} of these fractions are notably lower than the EC_{50} of the positive controls (Trolox 1.18 ± 0.06 – $1.31\pm 0.07 \mu\text{g/ml}$ and ascorbic acid 1.50 ± 0.06 – $1.68\pm 0.07 \mu\text{g/ml}$). Other fractions which have notable antioxidant activity were the butanol fractions of *B. bowkeri* ($0.88\pm 0.18 \mu\text{g/ml}$), *B. galpinii* ($0.89\pm 0.04 \mu\text{g/ml}$), *B. variegata* ($1.05\pm 0.11 \mu\text{g/ml}$), *C. vendae* ($0.60\pm 0.03 \mu\text{g/ml}$), *C. woodii* ($0.89\pm 0.06 \mu\text{g/ml}$) and *E. crispa* ($1.45\pm 0.08 \mu\text{g/ml}$). The ethyl acetate fractions of *C. padoides* ($0.79\pm 0.01 \mu\text{g/ml}$), *C. vendae* ($1.20\pm 0.30 \mu\text{g/ml}$), *C. woodii* ($1.30\pm 0.13 \mu\text{g/ml}$), *S. pentheri* ($1.25\pm 0.08 \mu\text{g/ml}$), *E. crispa* ($1.34\pm 0.03 \mu\text{g/ml}$) including the crude extract of *O. paniculosa* ($0.99\pm 0.05 \mu\text{g/ml}$) also have EC_{50} lower or comparable to the control. The hexane fractions exhibited poor antiradical activity with the EC_{50} being 6.78 ± 0.29 and $1709.0\pm 91.44 \mu\text{g/ml}$ for *C. woodii* and *C. bracteosum* respectively. From the estimated EC_{50} values, the order of potency is ethyl acetate fraction > butanol fraction > crude extract > dichloromethane fraction > residual water fraction > hexane fraction.

Table 5.2. ABTS radical scavenging potential of the crude extract and fractions expressed as EC_{50} ($\mu\text{g/ml}$)

Plant spp	Crude	Hexane	DCM	ETOAc	Butanol	Residual Water
Bab	14.50 ± 0.50	50.63 ± 6.37^b	5.54 ± 0.48^c	1.81 ± 0.13^c	0.88 ± 0.18^c	17.99 ± 2.34^a
Bag	55.01 ± 0.25	102.25 ± 5.04^b	11.81 ± 0.90^c	3.21 ± 0.22^d	0.89 ± 0.04^d	6.21 ± 1.03^c
Bap	17.19 ± 0.14	116.27 ± 0.57^b	9.76 ± 0.20^a	5.01 ± 0.07^c	7.31 ± 0.85^c	40.44 ± 7.09^c
Bav	9.24 ± 1.30^a	85.84 ± 2.55^b	8.58 ± 0.43^a	2.40 ± 0.22^a	1.05 ± 0.11^a	73.91 ± 3.68^b
Erl	246.37 ± 17.73	50.89 ± 6.08	8.76 ± 0.22	6.52 ± 0.19	18.52 ± 1.10	125.00 ± 4.22

Inc	44.95±5.60	1017.73±65.41	49.89±10.43	3.19±1.16	ND	ND
Scb	4.12±0.53	276.76±54.53	ND	5.3.3±0.60	2.09±0.45	61.64±15.23
Cob	11.34±1.5	1709±91.44	59.12±2.33	7.38±0.71	22.57±1.03	190.64±16.3
Cop	4.17±0.02	22.72±1.46	4.03±0.20	0.71±0.01	0.43±0.03	1.47±0.82
Cov	6.01±0.07	23.93±0.95	2.82±0.44	1.20±0.03	0.60±0.03	4.26±0.12
Cow	9.78±0.08	6.78±0.29	1.71±0.03	1.30±0.13	0.83±0.06	5.73±1.50
Ozm	15.93±2.10	43.48±4.20	17.55±2.79	10.74±0.45	68.85±23.76	288.17±7.05
Ozp	0.99±0.05	191.47±10.94	161.60±4.16	1.60±0.61	7.74±4.57	172.67±29.37
Sle	5.43±0.07	129.17±20.20	113.80±11.07	4.63±0.56	23.43±7.72	278.77±16.25
Spd	1.94±0.20	213.73±18.31	153.00±15.48	2.19±0.24	20.39±0.24	87.85±8.60
Spt	4.70±0.24	142.80±6.62	169.33±4.38	1.25±0.08	ND	ND
Mpd	8.64±0.13	114.64±25.93	33.54±1.29	6.33±0.18	52.79±14.43	74.89±2.80
Mpr	4.03±0.18	277.80±16.13	22.26±1.33	1.71±0.13	8.99±2.86	130.70±15.05
Mse	5.34±0.39	312.73±43.83	139.90±13.65	3.59±0.06	7.78±3.13	62.86±3.90
Mun	7.89±0.30	286.30±7.78	55.30±5.09	6.66±1.53	5.74±1.37	220.27±30.15
Euc	4.18±0.86	83.39±15.89	2.50±0.39	1.34±0.03	1.45±0.08	4.51±0.13
Eun	3.53±0.55	281.77±42.97	7.41±0.33	2.10±0.23	3.25±0.16	6.54±0.40
Fic		548.43±191.60	273.36±31.81	4.63±0.92	42.09±8.04	ND
Fig		285.43±12.83	187.53±4.54	9.06±1.51	2.80±0.50	13.93±1.02
Cam	7.87±0.93	293.97±77.70	11.68±3.66	2.66±0.14	2.45±0.21	17.92±1.37
Com	19.13±0.69	51.11±2.36	6.14±0.49	2.07±0.10	3.35±	270.03±20.94
Syp						

Bab (*Bauhinia bowkeri*), Bag (*Bauhinia galpinii*), Bap (*Bauhinia petersiana*), Bav (*Bauhinia variegata*), Cam (*Carissa macrocarpa*), Cob (*Combretum bracteosum*), Cop (*Combretum padoides*), Cov (*Combretum vendae*), Cow (*Combretum woodii*), Com (*Commiphora harveyi*), Erl (*Erythrina latissima*), Euc, (*Euclea crispa*), Eun (*Euclea natalensis*), Fic (*Ficus craterestoma*), Fig (*Ficus glumosa*), Inc (*Indigofera cylindrica*), Mpd (*Maytenus peduncularis*), Mpr (*Maytenus procumbens*), Mse (*Maytenus senegalensis*), Mun (*Maytenus undata*), Ozm (*Ozoroa mucronata*), Ozp (*Ozoroa paniculosa*), Sle (*Searsia leptodictya*), Spd (*Searsia pendulina*), Spt (*Searsia pentheri*), Scb (*Schotia brachypetala*), Syp (*Syzygium paniculatum*)

5.4.4. Ferric reducing antioxidant power (FRAP) gradient

The FRAP results are presented in Table 5.3 as the slope of the best fit linear regression analysis. Some of the ethyl acetate and butanol fractions had moderate to good dose-dependent ferric ion reducing capability comparable to the controls (trolox and ascorbic acid). The ethyl acetate fractions *Carissa macrocarpa* (45.0±3.7), *Combretumpadoides* (54.15±4.87), *Combretum vendae* (49.87±2.91), *Combretum woodii* (45.89 ± 3.87), *Commiphora harveyi* (50±3.5), *Euclea crispa* (48.0±4.8) and *Euclea natalensis* (42.0±3.9) have reducing power gradients compared to the trolox and ascorbic acid. The butanol fractions of *Bauhinia bowkeri* (40.92±2.14), *C. padoides* (44.11±4.06), *C. vendae* (42.59± 3.81), *C. woodii* (41.19 ± 1.28) and *E. crispa* (45.0±4.2) also exhibited good reducing power comparable with the control. The orders of reducing capacity is ethyl acetate > butanol > DCM > water > hexane > crude extract.

Table 5.3. Ferric reducing antioxidant power (FRAP) of the crude extracts and fractions expressed as the gradient of the linear curve

Plant species	Crude extract	Hexane fraction	DCM fraction	ETOAc fraction	Butanol fraction	Water fraction
Bab	3.30±0.17	8.15±0.76	18.75±1.18	31.84±3.36	40.92±2.14	10.26±1.21
Bag	3.63±0.19	8.36±0.41	15.20±0.59	21.48±1.55	32.70±2.10	31.75±2.08
Bap	1.51±0.05	7.57±0.34	16.42±0.83	15.50±1.69	24.19±1.22	24.63±1.14
Bav	2.63±0.27	11.74±0.62	15.22±0.75	27.75±2.58	32.06±1.11	0.47±0.05
Cam	1.70±0.10	6.1±0.57	17.0±0.87	45.0±3.7	29.1±1.7	5.2±0.59
Cob	9.4±0.32	0.27±0.037	1.6±0.10	2.7±0.40	5.2±0.25	11.0±0.37
Cop	6.36±0.26	16.96±0.91	22.68±1.04	54.15±4.87	44.11±4.06	9.59±0.58
Cov	3.310 ± 0.25	14.28±0.72	27.84±1.66	49.87±2.91	42.59±3.81	3.59±0.37
Cow	4.98±0.21	24.02±1.79	25.91±2.98	45.89±3.87	41.19±1.28	3.43±0.67
Com	1.1±0.15	14.0±0.57	26±1.7	50±3.5	36±3.0	0.26±0.08
Erl	0.29±0.07	16.0±0.60	23.0±1.3	33.0±2.9	18.0±0.84	1.8±0.13
Euc	5.5±0.17	11.0±1.2	25.0±2.1	48.0±4.8	45.0±4.2	20.0±0.91
Eun	5.3±0.27	6.1±0.18	21.0±1.2	42.0±3.9	34.0±2.2	20.0±1.9

Bab (*Bauhinia bowkeri*), Bag (*Bauhinia galpinii*), Bap (*Bauhinia petersiana*), Bav (*Bauhinia variegata*), Cam (*Carissa macrocarpa*), Cob (*Combretum bracteosum*), Cop (*Combretum padoides*), Cov (*Combretum vendae*), Cow (*Combretum woodii*), Com (*Commiphora harveyi*), Erl (*Erythrina latissima*), Euc, (*Euclea crispa*), Eun (*Euclea natalensis*)

5.4.5. Effective concentration required to reduce hydroxyl radical (oxidant) by half (EC₅₀)

The scavenging ability of the crude extracts and fractions expressed as EC₅₀ are presented in Table 5.4. The EC₅₀ ranged from 11.03±2.80 µg/ml (dichloromethane fraction of *O. paniculosa*) to 356.80±2.39 µg/ml (dichloromethane fraction of *M. senegalensis*). The order of hydroxyl radical inhibition is butanol > ethyl acetate > crude extract > dichloromethane > hexane.

5.4. Hydroxyl radical scavenging potential of the crude extract and fraction expressed as EC₅₀ (µg/ml)

Plant species	Crude	Hexane	DCM	ETOAc	Butanol	Residual Water
Mpd	23.92±2.28	110.54±17.91	122.07±20.50	70.86±18.09	49.55±5.70	
Mpr	107.69±12.32	179.70±41.17	223.96±42.04	76.70±11.56	48.79±12.42	
Mse	146.30±21.60	187.40±55.56	356.80±2.39	42.06±12.90	30.81±1.78	
Mun	80.68±2.90	284.36±27.04	311.90±150.33	30.81±1.78	51.19±5.30	
Ozm	44.29±4.20	175.56±6.88	45.77±0.98		82.24±0.97	
Ozp	33.02±6.46	35.90±3.20	11.03±2.80	33.07±0.85	17.17±3.39	
Sle	43.88±8.57	128.69±8.96	27.76±0.90	41.17±7.90	64.23±9.55	
Spd	83.46±10.45	130.45±5.70		39.02±0.90	74.51±15.46	
Spt	74.69±2.87	73.93±4.93	19.02±2.70	22.59±6.5	38.16±	

Mpd (*Maytenus peduncularis*), Mpr (*Maytenus procumbens*), Mse (*Maytenus senegalensis*), Mun (*Maytenus undata*), Ozm (*Ozoroa mucronata*), Ozp (*Ozoroa paniculosa*), Sle (*Searsia leptodictya*), Spd (*Searsia pendulina*), Spt (*Searsia pentheri*)

5.4.6. Lipid peroxidation inhibition effective concentration (EC₅₀)

The inhibitory effect on the lipid peroxidation expressed as EC₅₀ values are presented in Table of 5.5. The most active are *O. mucronata* and *C. woodii* with EC₅₀ of 13.95±2.25 and 13.24±1.17 µg/ml respectively followed by *C. bracteosum* with 17.89±1.72 µg/ml. The least active extracts were *S. leptodictya* and *M. peduncularis* with EC₅₀ of 40.45±13.38 and 39.84±5.52 µg/ml respectively.

Table 5.5: Linoleic acid peroxidation inhibition expressed as EC₅₀ (µg/ml)

Plant species	EC ₅₀ (µg/ml)
<i>Combretum bracteosum</i>	17.89±1.72
<i>Combretum padoides</i>	35.62±4.37
<i>Combretum vendae</i>	30.91±2.53
<i>Combretum woodii</i>	13.24±1.17
<i>Maytenus peduncularis</i>	39.84±5.52
<i>Maytenus procumbens</i>	34.21±1.63
<i>Maytenus senegalensis</i>	27.21±2.30
<i>Maytenus undata</i>	33.70±0.85
<i>Ozoroa mucronata</i>	13.95±2.25
<i>Ozoroa paniculosa</i>	25.20±8.10
<i>Searsia leptodictya</i>	40.45±13.38
<i>Searsia pendulina</i>	30.21±5.49
<i>Searsia pentheri</i>	25.53±6.20

5.5. Discussion

5.5.1. Qualitative antioxidant analyses (DPPH-TLC bioautography)

The antioxidant assay using DPPH on TLC plates to screen plant extracts is a quick method used to confirm the potential of the extracts for further evaluations. The intensity of the yellow spot depends on the amount and chemical characteristics of the compound present. The reaction kinetics between the DPPH radical and the active compounds varies as some compounds react slowly while others react fast. Some antioxidant spots were not readily visible immediately after sprayed with DPPH but appeared after incubation at room temperature for 2-12h. Also the mechanism of the reaction may differ as some of the compounds act as hydrogen donors and others may act as electron donors. In the DPPH radical scavenging process hydrogen donation is the predominant mechanism (Rekka and Kourounakis, 1991). This antioxidant assay is fast, simple and the image can be stored for future reference. All the extracts and fractions of the 27 plants used had antioxidant properties with varying number of yellow spots as free radical scavenging potential against the purple background of the DPPH radical on the plate. The three different mobile phases (5, 6 and 8) used to develop the chromatogram for

TLC-DPPH analyses demonstrated close relationships between the compounds present in the dichloromethane fraction of *E. latissima*, *C. vendae*, and *C. woodii*. These results indicate the danger of considering only chemical markers in taxonomy because *Combretum* and *Erythrina* are not closely related based on classical taxonomic parameters.

The antioxidant activities of the crude extracts and their fractions of varying polarities were quantified several different antioxidant assays such as DPPH[•] and ABTS^{•+} synthetic free radicals, the hydroxyl radical and their ferric reducing capacities using the FRAP assay and lipid peroxidation inhibition. More than one type of antioxidant capacity measurement usually performed to take into account the various modes of antioxidant mechanism. These methods were not specific to any particular antioxidant component rather to the overall capacity of the extract. Of these methods only hydroxyl radical scavenging, FRAP and LPO assays have direct physiological importance as a measure of plant extracts protective performance against free radical chain reactions in cellular membranes.

For the DPPH and ABTS assay which involves hydrogen atom transfer and electron transfer processes respectively, all the extracts and fractions have a dose-dependent radical activity with butanol fraction being the most active with EC₅₀ ranges from 0.21 ± 0.03 µg/ml for *Combretum padoides* to 41.8 ± 3.37 µg/ml for *Searsia pendulina* (DPPH) and EC₅₀ ranges from 0.43 ± 0.03 µg/ml for *C. padoides* to 68.85 ± 23.76 µg/ml for *Ozoroa mucronata* (ABTS). These results suggested that the strong DPPH radical scavenging ability of ethyl acetate fraction was closely related to the high levels of phenolic compounds and due to the scavenging of the radical by hydrogen donation. Ethyl acetate fractions with EC₅₀ ranges from 0.44 ± 0.06 µg/ml for *C. padoides* to 7.92 ± 1.63 µg/ml for *Ozoroa mucronata* (DPPH) and EC₅₀ ranges from 0.71 ± 0.01 µg/ml for *C. padoides* to 10.74 ± 0.45 µg/ml for *O. mucronata* (ABTS) and the crude extracts with EC₅₀ ranges from 1.91 ± 0.15 µg/ml for *S. pendulina* to 123.60 ± 11.05 µg/ml *Bauhinia variegata* (DPPH) and EC₅₀ ranges from 0.99 ± 0.05 µg/ml for *Ozoroa paniculosa* to 246.37 ± 17.73 µg/ml for *Erythrina latissima* (ABTS).

In the results presented in Table 5.3, the higher the slope value the stronger the total antioxidant capacity (reduction power) of the tested extracts or fractions. The ferric reducing antioxidant power (FRAP) serve as a significant indicator of antioxidant potential of medicinal plant preparations and the activity are potentiated in the butanol and ethyl acetate fractions as observed in the DPPH and ABTS assays results. Antioxidant capacity of plant extracts and fractions depend on factors such as the compositions, chemical structures of the constituents and conditions of the test used. These results indicates that the phytochemical present in the plants performed as good electron or hydrogen donors and therefore should be able to terminate radical chain reaction by converting free radicals and reactive oxygen species to more stable products.

Ferric ions generate hydroxyl radical in vivo through Haber-Weiss and Fenton reaction mechanisms to hydroxyl. Hydroxyl radicals are highly strong reactive oxygen species, and there is no specific enzyme to defend against them in living organisms (Liu *et al.*, 2005). Hydroxyl radicals and other reactive species are also produced by

activated neutrophils, eosinophils monocytes and macrophages during inflammatory responses of the immune process. Hydroxyl radicals-mediated and propagated lipid peroxidation of the gastrointestinal tract mucosa phospholipids are considered to play a crucial role in the pathophysiology of numerous chronic diseases. The major toxic products of LPO are 4-hydroxyl-2-nonenal and malondialdehyde (MDA) which can react with intestinal epithelium mucosa resulting in altered transport process (fluid and ions). These mechanisms are involved in varieties of diarrhoea aetiology such as infection, toxin, and inflammations.

ROS are important mediators that initiate and propagate inflammatory responses by inducing the formation of pro-inflammatory cytokines such as interleukin-1 β (IL-1 β) and tumour necrosis factors (TNF- α). ROS/RNS are generated directly by COX at the site of inflammation and have regulatory role in the expression of COX and subsequent synthesis of PGE, therefore amplifying the acute phase of the inflammatory responses. These inflammatory mediators contribute to diarrhoea aetiology as direct secretagogue (pro-secretory), reduced fluid absorption capacity due to damaged to mucosa epithelial tissue and/or modulation of the intestinal contractility through enteric nervous system (ENS).

Oxidative damage to cellular components such as cell membrane by free radicals is believed to be associated with immune system decline and hyperactivation. Immune activation of PMNs and monocytes result in formation of potent hypochlorous acid (HOCl) from myeloperoxidase (MPO)-catalyzed oxidation of Cl⁻ by H₂O₂. In addition, the HOCl react with primary amine groups (RNH₂) to produce N-chloramines (RNHCl). Both HOCl and RNHCl are cytotoxic (Pavlick *et al.*, 2002). Some of the ROS/RNS and their products enhances intestinal and colonic Cl⁻ acts directly or indirectly to initiate diarrhoea as secretagogue (Gaginella *et al.*, 1995).

Considerable interest is focused in finding natural antioxidants which can help on the management of numerous diseases with oxidative stress aetiology and maintenance of good health. Oxidative stress and the associated diseases resulting from an imbalance between the endogenous antioxidant defence mechanisms and pro-oxidative forces in favour can be alleviate by increased expression of antioxidant (Pavlick *et al.*, 2002). Antioxidants have the capacity of stopping the chain reaction of oxidative species and the deleterious health hazard to the body. The crude extract of these medicinal plants inhibit Fenton-generated hydroxyl radical-mediated peroxidation of a heterogeneous phospholipid-aqueous phosphate buffered system and scavenged hydroxyl radical which are important characteristic of phenolic compounds. Free radical scavenging and antioxidant activity of these medicinal plants contribute their therapeutic effect against diarrhoea diseases and other GIT disorders for which they are being use ethnopharmacological in South African traditional medicine to treat.

5.6. Conclusion

In this study, the extracts and their fractions were found to have various forms of antioxidant activities that could possibly be attributed to the phenolic constituents. The extraction and fractionation protocols potentiate the

antioxidant components in the polar fractions while the non-polar hexane fractions demonstrated little or no antioxidant activity except one prominent spot from the *Erythrina latissima* extract in TLC-DPPH analyses.

Linoleic acid and arachidonic acid are indigenous compounds of the cell membrane with a task to protect the cell. The two membrane lipids are prone to attack during induce inflammatory and oxidative stress. However, the increases of intracellular ROS level, due to increased production or impaired removal, can also cause cell damage ranging from cytoplasmic swelling to cell death. In view of the involvement of the many oxidative mechanisms in the pathogenesis of various diseases, free radicals scavenging and removal of excessive ROS are important for restoring normal conditions, which might be the possible reasons of the correlation between antioxidant activity and other therapeutic activities. This investigation provided data clarifying the potentials of some of the plants as promising sources of natural antioxidants.

Further work on *in vivo* verification of the antioxidant therapeutic effectiveness, bioavailability, absorption and metabolism of the active component is needed. Finally identification, isolation, characterization and absence of possible toxicity of the bioactive compounds also required further investigation. Free radicals and oxidative species play some critical roles in diseases with inflammatory aetiologies including the GIT disorders and immunosuppression mechanism. The crude extracts and the polar fractions (ethyl acetate and butanol) of many of the plant species have strong antioxidant activities, consequently may reduce inflammation or stimulate the immune system of host. This could be one of the anti-diarrhoeal mechanisms and therefore explaining the traditional use of these medicinal plants. The anti-inflammatory potentials of the crude extracts will be investigated in the next chapter

CHAPTER SIX

Anti-inflammatory activities of the crude extracts as antidiarrhoeal mechanisms

6.0. Introduction

Inflammation is an important component of immune response to pathogens and damaged cell characterized by heat, redness, pains, swelling and sometimes loss of tissue functionality in chronic situation (MacNaughton, 2006). Although inflammatory response provides an important defence mechanism to injurious agents, injury to some healthy cells at the inflammatory site could also occur (Sprague and Khalil, 2009).

The cellular immune systems including that of the gastrointestinal tract (GIT) act as defence mechanisms by mobilizing white blood cells (leukocytes and other chemicals) to fight infections and harmful stimuli. The body's reaction to this phenomenon may trigger inflammatory responses through the release of pro-inflammatory eicosanoids such as prostaglandins, prostacyclins and leukotrienes, and pro-inflammatory cytokines (Nardi *et al*, 2007) such as interleukins (IL-1B, IL-3,IL-6), interferons (IFN), tumor nuclear factor (TNF- α) and platelet-activating factor (PAF) (Conforti *et al*, 2008; Kunkel *et al*, 1996).

Cyclooxygenase (COX) and lipoxygenase (LOX) oxidation of polyunsaturated fatty acid (PUFA) such arachidonic acid or linoleic acid forming bioactive eicosanoids are the major features of inflammatory response (Haeggstrom *et al*, 2010) (Fig 6.1). The generation and release of reactive species (ROS/RNS) by inflammatory cells in response to pathogens and stimuli is considered the major microbicidal mechanism in the body. However, excessive generation of ROS/RNS exacerbate inflammatory responses that may lead to development of disease state.

In the GIT, inflammation affects epithelial cells as well as the more specialised mucus secreting and enteroendocrine cells of the gut mucosa (Spiller, 2004). Inflamed intestinal epithelial mucosa usually results in increased permeability, increased bowel movement or contractility, inadequate digestion of food materials and impaired absorption of essential food components (Spiller, 2004). Some inflammatory mediators are ion or fluid secretagogue and prokinetic of enteric nervous system (ENS) causing diarrhoea, and malnutrition (See section 2.10.2 for detailed discussion).

Intestinal inflammation causes damage to mucosal barrier function comprised of physical diffusion barriers, physiologic and enzymatic barriers, and immunologic barriers (Soderholm and perdue, 2006). The continuous layer of epithelial cells interconnected by tight junction, restricts both transcellular and paracellular permeation, therefore, constituting the major part of the mucosal barrier. Active secretion of fluid and mucus containing secretory immunoglobulin isotype A (IgA) also serve to bind, dilute, and cleaning mechanisms of the intestine. The intestinal propulsive movement is also an important protective process against noxious substances (Soderholm and perdue, 2006).

6.1. Effect of cyclooxygenases (COX) on GIT

Cyclooxygenases (COXs) are oxidizing enzymes which metabolize polyunsaturated fatty acid (PUFA) such as arachidonic acid liberated from membrane phospholipid by phospholipases A to various eicosanoids such as prostaglandin D (PGD), prostaglandin E (PGE), prostacyclin (PC), thromboxane (TXA₂) (Xu *et al*, 2007; Simon *et al*, 2004). The physiological activities of these inflammatory mediators are mediated by G-protein-coupled prostanoid receptors such as DP, EP₁₋₄, FP, IP and TP which preferentially respond to PGD, PGE, PGF_{2α}, PGI and TXA respectively. There are two isoforms of COX namely COX-1 and 2. COX-1 is constitutively expressed to produce PG series which are involve in the regulation of physiological housekeeping such as platelet aggregation, homeostasis of the GIT and the kidney. The COX-2 is an inducible enzyme expressed to produce PG series which are responsible for pro-inflammatory stimuli such as cytokines, growth factors, tumor promoting agents and bacterial endotoxins.

Prostaglandins are widely distributed along the GIT and are involved in a number of physiological and pathological processes including motility, blood flow, water and electrolyte absorption, and mucus secretion. PGE₂ is cytoprotective to the intestinal epithelium by decreasing gastric acid secretion, thus prevents ulceration. However, in pathological situation, it also increases intestinal motility and intestinal secretion causing secretory diarrhoea.

6.2. Effects of Lipoxygenase in GIT

Lipoxygenases (LOXs) comprise a family of non-heme iron-containing dioxygenases, representing the key enzymes in the biosynthesis of leukotrienes from PUFA. Leukotrienes have been postulated to play essential role in the pathophysiology of several inflammatory and allergic diseases. The LOXs are classified with respect to their positional specificity of arachidonic acid oxygenation as 5-LOX, 9-LOX, 12- LOX, 15-LOX. The products of LOXs catalysed oxygenation include leukotrienes, lipoxins, hydroperoxyeicosatetraenoic acids (HPETE), and hydroxyeicosatetraenoic acids (HETE).

LTB₄ is synthesized by 5-LOX from arachidonic acid. It is a potent chemotactic agent for inflammatory cells such as neutrophils and macrophages. It elicits leukocytes migration towards inflammatory sites and activates neutrophils, causing their degranulation associated with enzyme release as well as superoxide radicals. It also plays an important role in immune systems by enhancing the release of pro-inflammatory cytokines by macrophages and lymphocytes.

6.3. Effects of cytokines on GIT

Secretion of cytokines by the intestinal immune system is one of the main factors in maintaining the gut integrity in quiescent homeostasis. Cytokines are classified as pro-inflammatory (TNF-α, IL-1, -6, -12, -15, -18, and -32, as well as the anti-inflammatory cytokines IL-10 and TGF-β produced predominantly by activated macrophages, involved in the up-regulation of inflammatory reactions and IFN-γ and IL-4 from T-cells (MacNaughton, 2006; Sprague and Khalil, 2009). Anti inflammatory cytokines such as IL-4, IL-10, IL-13, IFN-α, and TGF-β are involved

in the down regulation of inflammatory reactions (Sprague and Khalil, 2009). Of major importance is the balance between pro-inflammatory cytokines such as TNF- α , IL-1 and IFN- γ and regulatory cytokines like IL-10 and transforming growth factor- β . The features ultimately determine the capacity of an immune response as either detrimental or innocuous to the gut. TNF- α is a critical cytokine that elaborates inflammatory responses by activating a number of inflammatory cells including neutrophils, macrophages and NK cells which induces the production of inflammatory cytokines such as IL-1 β , IL-6 and IL-8 and upregulation of adhesion molecules on cell surface. In addition TNF- α also directly potentiates the immune response of other pro-inflammatory cytokines such as IL-1, IL-6, IL-12 and IFN- γ consequently enhancing the anti-inflammatory and anti-apoptotic effect. IFN- γ can stimulate the production of IL-1 β , platelet-activating factor, H₂O₂, NO and downregulate IL-8. As a pro-inflammatory cytokine, IFN- γ sensitizes intestinal epithelial cells to physiological and therapeutic inducers of apoptosis.

6.4. Oxidative species as inflammatory mediator

Production of reactive species (H₂O₂, O₂⁻, and OH[·]), nitric oxide (NO) and per-oxy-nitrite occurs at the site of inflammation and contributes to the exacerbation of inflammatory disease and tissue damage. Oxidative species stimulates the release of pro-inflammatory cytokines such as interleukin-1 β (IL-1 β) and tumor necrosis factor (TNF- α). In addition, ROS induced by activated neutrophils, eosinophils, monocytes and macrophages during the inflammation process leads to tissue injury by damaging macromolecules and effecting the lipid peroxidation of membranes. In acute or chronic inflammations, the production of O₂⁻ is increased at a rate that overwhelms the capacity of the endogenous SOD enzyme defence to dissipate. Reduction in the O₂⁻ generation can decrease side-effects of the radical in inflammatory conditions.

Nitric oxide (NO) is a free radical gas synthesized by nitric oxide synthase (NOS) from L-arginine and initiates diverse physiological and pathological processes (Lee *et al.*, 2007b). Three iso-forms of NOS had been identified; they are neuronal NOS (nNOS), endothelial NOS (eNOS) and inducible NOS (iNOS). The first two iso-forms (nNOS and eNOS) are constitutive NOS (cNOS) while iNOS is produced only by specific stimulants in some cells. The iNOS stimulants include cytokines or bacterial lipopolysaccharides or endotoxin. Inflammatory responses are associated with the production of large quantity of NO (Cuzzocrea *et al.*, 2001). The deleterious effects of NO include mitochondrial enzymes inhibition (Nathan, 1992) and activation of COXs to produce inflammatory PGs (Salvemini and Masferrer, 1996) and interaction with superoxide to generate cytotoxic peroxynitrite. NO is an important mediator in the inflammatory process and is produced at inflamed sites by iNOS. High levels of NO have been linked to a number of pathological processes including various forms of inflammation, circulatory shock, and carcinogenesis. Therefore, an inhibitor of NOS might be effective as a therapeutic agent for inflammatory diseases.

6.5. Allopathic anti-inflammatory therapies and adverse effects on GIT

In view of the importance of PGs in inflammatory response, the rate-limiting enzymes for PGs synthesis are the therapeutic targets in controlling inflammation. Non-steroidal anti-inflammatory drugs (NSAIDs) exert their action by inhibiting the activity of COX enzymes, thereby reducing the production of pro-inflammatory prostaglandins. NSAIDs are structurally diverse, including compounds in the salicylic acid, arylalkanoic acid, propionic acid (profens), N-arylanthranilic acid (fenamic acid), pyroolidine derivatives, oxicam and sulphonamide families. Classical NSAIDs exhibited non-selective inhibition of both COX-1 and COX-2 while some other NSAIDs however show preferential inhibitory activity toward one isoform or the other.

Although NSAIDs provides good therapeutic relief against inflammation, some of these drugs currently in use have various side effects, particularly in the gastrointestinal tract ulceration and kidney (Charlier and Michaux, 2003). Prolonged use of nonselective NSAIDs has adverse effects such as nausea, dyspepsia, gastritis, abdominal pain, peptic ulceration, gastrointestinal bleeding, and/or perforation of gastroduodenal ulcers. In addition, NSAIDs are postulated to shift the metabolite profile from COX derivatives to lipoxygenase (LOX) derivatives resulting in the accumulation of substrate for the LOX-derived metabolites. LOX products stimulate neutrophil migration, increase adhesion of leukocytes to endothelial cells, cause smooth muscle contraction, increase vascular permeability, and increase ion and mucus secretion. Inhibition of leukotriene biosynthesis decreases inflammation and accelerates gastrointestinal healing. However, there are controversies in the recent findings that non selective NSAID (indomethacin) causes GIT damage and neither selective COX-1 inhibitors nor selective COX-2 inhibitor causes any intestinal damage. The combine use of selective COX-1 and selective COX-2 inhibitors produces intestinal haemorrhage (Takeuchi *et al.*, 2010) (Fig 6.2). Aspirin a known non selective NSAID as its metabolite such as salicylic acid causes no intestinal damage but instead provided protection against ulcerogenic response induced by other classical NSAID. This also indicates that some complex mechanisms are responsible for the intestinal damage by a number of non selective NSAIDs other than COX-1 inhibition only (Takeuchi *et al.*, 2010). Some of the factors involved in the pathogenesis of NSAIDs toxicity include

- Bile acids secretion: NSAIDs increase the secretion of bile acids in the GIT causing complications such as colonic mucosa damage and diarrhoea. Bile acids induce the liberation of arachidonic acid from epithelial membrane, and the generation of COX and LOX metabolites along with the secondary active oxygen radicals.
- Intestinal motility: NSAIDs such as COX-1inhibitors causes marked enhancement of intestinal motility with regard to both the amplitude and frequency of contractility. Intestinal hypermotility caused mucosal hypoxia and microvascular injury due to smooth muscle contraction (Takeuchi *et al.*, 2002).
- Neutrophil infiltration: NSAIDs cause severe damage to the GIT resulting in loss of surface epithelium, mucosal necrosis and massive neutrophil infiltration.

- Bacterial flora: Non selective and COX-1 selective NSAIDs increased number of enterobacteria in the intestinal mucosa homogenates and luminal bacterial adherence to the mucosa induced severe intestinal injury (Takeuchi *et al.*, 2011).
- Nitric oxide (NO): Non selective and COX-1 selective NSAIDs cause an up-regulation of inducible NO synthases (iNOS) in the GIT. This is due to bacterial endotoxin increased intestinal permeability which induces the expression of iNOS and enhanced the generation of NO in the mucosa (Takeuchi *et al.*, 2011).
- Prostaglandins (PGs) deficiency: COX-1 isoform is expressed in most tissues, producing prostaglandins that play an important protective role in the gut by stimulating the synthesis and secretion of mucus and bicarbonate, increasing mucosal blood flow and promoting epithelial proliferation. The inhibitions of this enzyme by NSAIDs create PGs deficiency. In addition the inhibition of the COX-1 blocks platelet production of thromboxane, which increases bleeding when an active GI bleeding site is present. COX-2 isoform is induced in most tissues in response to inflammatory stimuli. Prostaglandins derived from COX-2 can be generated at the ulcer margin and appear to play an important role in ulcer healing through triggering the cell proliferation, promotion of angiogenesis and restoration of mucosal integrity (Takeuchi *et al.*, 2011). Effects of NSAIDs on GIT are presented in Fig 6.1 and Fig 6.2.




Fig. 6.1. Roles of COX-1 and COX-2 in the pathogenic mechanism of NSAID-induced intestinal damage (Takeuchi *et al.*, 2010)

- Effect COX-2 inhibition: The NSAIDs selective inhibition of COX-2 has adverse effect on cardiovascular function (Grosser *et al.*, 2006) due to suppression of PGI (anti-thrombotic) promoting hypertension and blood coagulation while the synthesis of TXA₂ (pro-thrombotic) by COX-1 remain unchanged (Fitzgerald, 2004).




Fig. 6.2. Various factors involved in the pathogenesis of indomethacin-induced small intestinal lesions (Takeuchi *et al*, 2010)

6.6. Plant phytochemicals as anti-inflammatory agents

Plant extracts are suspected to contain potential bioactive component that can strongly inhibit the expression of LOX and COX. Therefore, there is continuous need to search for new drugs from natural products with anti-inflammatory properties and minimum side effects. Modulation of the activities of the enzymes implies that the inflammation process can be modified. Anti-inflammatory agent may also have an antioxidant and radical-scavenging mechanisms as part of it activity.

Phenolics, alkaloids and triterpenoids have been demonstrated to exhibit anti-inflammatory activity by exerting anti-oxidative properties in reducing $O_2^{\cdot-}$ and malondialdehyde (MDA) production, plasma extravasations and cell migration mainly of leukocytes and potentiates the activity of SOD in radical scavenging (Nardi *et al*, 2007). Reactive species are one of the most important mediators that provokes or sustain inflammatory processes and consequently, their annihilation by antioxidants and radical scavenger such as phenolic compounds can alleviate inflammation (Delaporte *et al*, 2002; Geronikaki and Garalas, 2006).

6.7. Mechanisms of anti-inflammatory assay models

Anti-inflammatory potential were determined by in vitro assays based on the inhibitory effect on the biosynthesis of 12(S)-hydroxy-(5Z, 8E, 10E)-heptadecatrienoic acid (12-HHT), and 12(S)-hydroxy-(5Z, 8Z, 10E, 14Z)-eicosatetraenoic acid (12-HETE). 12-HHT and 12-HETE are inflammation mediators derived from arachidonic acid metabolism, which is catalysed by enzymes of inflammatory response, cyclooxygenase (COX-1) and lipoxygenase (12-LOX), respectively. The advantage of this type of experiment is avoidance of the undesirable in vivo tests on experimental animals, since the tests commonly used to detect the anti-inflammatory activity is carrageenan induced paw edema in rats.

6.8. Materials and Methods

6.8.1. Lipoxygenase inhibition assay

Lipoxygenase activity was determined spectrophotometrically according to [Taraporewala and Kauffman, \(1990\)](#); [Lyckander and Malterud, \(1992\)](#) which is based on the enzymatic oxidation of linoleic acid to the corresponding hydroperoxide. To determine hydroperoxide, soy lipoxygenase-1 (200 U) was incubated with linoleic acid (50 μM) in sodium borate buffer (200 mM, pH 9.0) for 4 min at 25°C. The absorbance at 234 nm was measured on a Helios β (Thermo Electron Corporation) spectrophotometer using a quartz cuvette. The inhibitory assays were performed in presence of extracts in different concentrations ranging from 0.15 - 25 $\mu\text{g/ml}$. The anti-inflammatory effect was evaluated by calculating percentage inhibition of hydroperoxide production from the changes in optical density values at 234 nm for 5 min. The test compound concentration causing 50% inhibition of hydroperoxide-release (IC_{50}) was calculated from the concentration–inhibition response curve by best fit non-linear regression analysis. The extinction coefficient of 25 $\text{mM}^{-1} \text{cm}^{-1}$ was used for quantification of lipid hydroperoxides. DMSO was used as negative control.

6.8.2. Cyclooxygenase enzymes inhibition (COX 1 and 2) assay

The experiments were performed using an assay originally described by [Noreen *et al.*, \(1998\)](#), with some modification by [du Toit *et al.*, \(2005\)](#). For COX-1 assay, commercial COX-1 (from ram seminal vesicles, Sigma-Aldrich) (10 $\mu\text{l/sample}$) enzymes and Hematin (co-factor) (50 $\mu\text{l/sample}$) was pre-incubated for 5 min on ice. The mixture (enzyme and cofactor) was added to the test sample (2.5 μl of test sample and 17.5 μl of water) to make a concentration of 0.25 $\mu\text{g}/\mu\text{l}$ in the final assay volume and pre-incubated for 5 min at room temperature. 1- ^{14}C -arachidonic acid (20 μl) was added to the enzyme-test sample mixture and incubated for 10 min in a water bath at 37°C. The reaction was terminated by adding 10 μl of 2M HCl. The amount of [^{14}C]-labelled PG synthesized was measured using a Packard scintillation counter after removing the unmetabolized [^{14}C]-arachidonic acid substrate by column chromatography using Pasteur pipette as column. Unmetabolized [^{14}C]-arachidonic acid substrate was eluted with n-hexane-dioxane-glacial acetic acid (70:30:1) while [^{14}C]-labelled PG synthesized in the reaction was eluted with ethyl acetate: methanol (85:15).

The same procedure was adopted for COX-2 assay using three units of COX-2 enzymes (human recombinant, Sigma-Aldrich). Indomethacin at 12.5 μM and 200 μM used as positive controls for the COX-1 assay and COX-2 assays respectively. Two background controls in which the enzymes were inactivated with HCl before the addition of [^{14}C]-arachidonic acid and two solvent blanks were prepared for experiment. The results were expressed as percentage inhibition (% I) using equation described by [Lin *et al.*, 1999](#).

$$\% \text{ I} = [(1 - \text{DPM of sample}) - (\text{DPM of background} / \text{DPM of blank} - \text{DPM of background})] \times 100$$

Where DPM is disintegration min^{-1}

6.9. Results

6.9.1. Cyclooxygenase inhibition assay

Results from cyclooxygenase assay against COX-1 are presented in Fig. 6.3. All the extracts exhibited moderate to good activity with the inhibitory effects ranging between 41.70 to 84.61%. The most active extract against COX-1 was *Carissa macrocarpa* with $82.98 \pm 1.62\%$ inhibition at 250 $\mu\text{g/ml}$ and $69.72 \pm 1.91\%$ at 62.5 $\mu\text{g/ml}$. The inhibition of COX-1 enzyme was concentration dependent as *Bauhinia petersiana* inhibited 50% of the enzyme at a concentration of 167 $\mu\text{g/ml}$ ($R^2=0.989$). *Bauhinia bowkeri* and *Bauhinia galpinii* inhibited 50% of the enzyme at the concentrations of 241 $\mu\text{g/ml}$ ($R^2=0.9645$) and 377.66 $\mu\text{g/ml}$ ($R^2=0.9216$) respectively. *Commiphora harveyi* was only active at concentration of 250 $\mu\text{g/ml}$ by inhibiting $45.45 \pm 2.96\%$ of COX-1 enzyme.




Fig. 6.3. COX-1 inhibitory activity of some selected phenolic-enriched crude extracts

The phenolic-rich crude extracts of these plants had no inhibitory activity against COX-2 at the maximum concentration tested (250 $\mu\text{g/ml}$) indicating that the extracts were COX-1 selective inhibitor. The phenolic rich crude extracts of *C. padoides*, *C. vendae* and *C. woodii* exhibit no cyclooxygenase inhibitory activity against COX-1 and 2 enzymes. Indomethacin used as reference compound had IC_{50} of 3.30 ± 0.006 and 122.5 μM against COX-1 and COX-2 respectively.

6.9.2 Lipoxygenase inhibitory assay

The lipoxygenase inhibitory capacity expressed as LC_{50} and percentage inhibition are presented in Table 6.1. The activity of the extract varied widely ranging between 0.86 ± 0.27 and 111.44 ± 37.28 $\mu\text{g/ml}$. The most active extract was obtained from *Syzygium paniculatum* with LC_{50} of 0.86 ± 0.27 $\mu\text{g/ml}$ and percentage inhibition of 66.74 ± 2.07 followed by *Euclea crispa* with LC_{50} of 2.55 ± 0.13 $\mu\text{g/ml}$ and percentage inhibition of 63.06 ± 1.75 . The least active extract was from *Commiphora harveyi* with LC_{50} of 111.44 ± 37.28 $\mu\text{g/ml}$ and percentage inhibition of 39.15 ± 1.92 (extract concentration equals 25.6 $\mu\text{g/ml}$).

Table 6.1: Lipoxygenase inhibitory activity of the crude extracts

Plant species	LC_{50} ($\mu\text{g/ml}$)	% inhibition (25 $\mu\text{g/ml}$)

Bag	4.10±0.62	56.31±4.36
Bap	10.18±2.25	52.15±2.1
Bav	5.07±1.11	56.00±0.97
Cam	30.22±0.83	46.29±1.69
Cop	25.12±2.05	44.97±0.53
Cov	33.48±6.01	45.30±0.70
Cow	15.70±5.57	49.21±1.25
Com	111.44±37.28	39.15±1.92
Euc	2.55±0.13	63.06±1.75
Eun	17.23±0.13	50.46±1.49
Erl	7.25±1.84	55.09±1.85
Fic	8.48±4.35	53.87±2.25
Fig	5.02±1.46	66.66±3.64
Inc	7.90±1.87	55.37±4.25
Mpd	4.08±0.51	61.54±4.19
Mpr	11.08±3.50	55.80±3.61
Mse	10.88±1.92	53.18±2.31
Mun	4.68±2.44	56.50±1.81
Ozm	2.88±1.10	57.90±1.62
Ozp	27.33±9.16	46.80±2.20
Sle	11.60±2.61	54.80±2.21
Spd		
Spt	9.16±2.07	54.76±2.79
Scb	4.09±2.37	60.19
Syp	0.86±0.27	66.74±2.07

6.9. Discussion

6.9.1. Cyclooxygenase assay

COX enzymes are the rate-determining enzymes in the prostaglandin biosynthetic pathways. The modulation of the enzymes can help in anti-inflammatory treatments due to the key role of PG especially PGE₂ in the inflammatory response (Gale *et al.*, 2007). ROS have been reported to have a regulatory role in the expression of COX, particularly COX-2 and subsequent synthesis of PGE₂ which is responsible for inflammation. Classification of inflammatory activity of extract based on extractants as 59% (minimum inhibition) by aqueous extracts tested at a final concentration of 250 µg per test solution and for organic extracts is 70%, when tested at a final concentration of 250 µg per test solution (Fennell *et al.*, 2004). The polyphenolic-rich extracts of the plant species tested exhibited selective inhibition of COX-1. The results confirmed the postulation that most phenolic compounds like flavonoids exhibit COX-1 selective inhibitory activity and have no effect on COX-2 isoform (Kim *et al.*, 2004). COX-1 is also reported to be involved in the inflammatory response and compensatory mechanisms

between COX-1 and COX-2 have been demonstrated (Gale *et al.*, 2007). COX-1 is the predominant isoenzyme in the normal gastrointestinal tract (Radi and Khan, 2006) and modulates neurogenic contraction (Smid and Svensson, 2009), while COX-2 expression is up-regulated during inflammation, where it modulates cholinergic contraction and small bowel motility. COX-2 mediated PGs from inflamed gastrointestinal mucosa may play a role in the chloride and fluid flux that helps flush GI bacteria.

However, the inhibition of COX-2 in the inflamed GI mucosa has been hypothesized to delay the resolution of GI injury. Since the polyphenolic-rich crude extracts of *C. macrocarpa*, *B. bowkeri*, *B. galpinii* and *B. petersiana* exhibited selective inhibition of COX-1, the use of these plants in traditional medicine as anti-diarrhoea agents need to be monitored critically especially in term of dosage. The phenolic-enriched extracts of *C. padoides*, *C. vendae*, *C. woodii* and *Syzygium paniculatum* did not exhibit activities against COX-1 and COX-2. However, the anti-inflammatory activities of these extracts cannot be ruled out as it may involve in other inflammatory mediators. In diarrhoea disease, ROS and RNS are known to activate many pro-inflammatory cytokines (interleukins and TNF α), cell adhesions and COX enzymes. Oxidative damage exacerbates intestinal inflammatory response and causes a virulent cycle of oxidative stress, inflammation and increased mucosal permeability (Chen *et al.*, 2007). Though, the phenolic-enriched extracts of the three *Combretum* species may not have a direct effect on COX enzymes, the significant free radical, ferric reducing properties and inhibition of lipid peroxidation may probably influence inflammation process.

6.9.2. Lipoygenase assay

The anti-inflammatory activities of phenolic-enriched crude extracts on inflammatory mediators were measured against soybean lipoygenase enzyme (Table 6.1). All medicinal extracts inhibited the lipoygenase enzyme and these inhibitory effects are concentration dependent. Lipoygenases inhibition correlate to antioxidants because lipid hydroperoxide formations are usually inhibited as a result of the scavenging of lipid-oxy- or lipid-peroxy-radicals formed in the course of enzymatic peroxidation. Consequently, limiting the availability of lipid hydroperoxide substrates required for the catalytic cycle of lipoygenase oxidative process (Cuello *et al.*, 2011)

6.10. Conclusion

Some of the crude extract exhibited selective COX-1 and LOX inhibitory activities in the *in vitro* enzymatic assays conducted in this study. The release of arachidonic acid is closely related to the cyclooxygenase (COX) and 5-lipoygenase (LOX) enzyme systems. The ability of plant extracts, fractions and isolated pure compounds to inhibit both COX and LOX pathways of the arachidonate metabolism have been suggested to contribute to anti-inflammatory action (Middleton *et al.*, 2000). The inhibition of COX enzymes result in the shifting of arachidonic acid to the LOX pathway, which promotes gastrointestinal damage by recruiting leukocytes to the mucosal and stimulating gastric acid secretion. It is proposed that drugs that are capable of block both COX and LOX metabolic pathways (dual inhibitors) are best option in terms of NSAIDs. The dual inhibition of COX and LOX enhances their individual anti-inflammatory effects and reduce the undesirable side effects associated with

NSAIDs, especially of the gastrointestinal tract (Fiorucci *et al.*, 2001). Further work on *in vivo* anti-inflammatory evaluation of the extracts in an animal model is needed to confirm the therapeutic potentials of these plant extracts. The crude *B. galpinii* had COX-1 and LOX inhibitory activity above 50% at concentration of 250 and 25 µg/ml respectively. Considering the GIT injury potential of some of the plants due to selective COX-1 inhibition, cellular toxicity will be evaluated in the next chapter.

Chapter Seven

Cytotoxicity evaluation of the crude extracts against Vero African green monkey kidney cell lines

7.0. Introduction

Medicinal plants are assumed to be non-toxic and regarded safe due to their natural origin and long use in traditional medicine to treat various forms of diseases (Chen *et al.*, 2011; Fennell *et al.*, 2004). Medicinal plant preparations are administered with the hope of promoting health and treating various diseases such as infections, colds, inflammation, GIT disorders, insomnia, depression, heart diseases, diabetes, cancer, acquired immunodeficiency syndrome, and liver diseases has increased in recent times (Chen *et al.*, 2011). However, scientific studies on efficacy and safety of some medicinal plants indicated that there are many phytochemicals that have cytotoxic, genotoxic, and carcinogenic effects when used chronically (Ernst, 2004; Rietjens *et al.*, 2005). It should also be kept in mind that if a different extractant is used, the safety ascribed to traditional use based mainly on aqueous extracts may not be relevant at all.

The adverse effects of medicinal plant use arise due to organ toxicity, adulteration, contamination, contents of heavy metals, herb–drug interactions, poor quality control and inherent poisonous phytochemical (Jordan *et al.*, 2010). Some medicinal plant phytochemicals are associated with toxicities of the heart, liver, blood, kidney, central nervous system, gastrointestinal disorder such as diarrhoea, and less frequently carcinogenesis (Jordan *et al.*, 2010). In the formal herbal industry the toxicity problems of medicinal plants are attributable to insufficient quality assurance and non compliance to the standards of Good Manufacturing Practise (Palombo, 2006). Furthermore, the problem is complicated by adulteration of herbal remedies by surreptitious addition of synthetic drugs and other potentially toxic compounds such as other botanicals, microorganisms, toxins, pesticides, and fumigants (Palombo, 2006).

More importantly, if herbal medicines are used with prescription drugs especially those with narrow therapeutic indices it can result in potential harmful herb–drug interactions that cause altered drug response and toxicity (Chen *et al.*, 2011). The fact that herbal medicines contain many compounds (active and non active), the large number of pharmacologically active compounds also increases the chance of herb–drug interaction (Palombo, 2006). Like synthetic drugs, herbal bioactive compounds can also undergo Phase I and Phase II enzymatic transformations to form nontoxic metabolites which are excreted through the faeces and urine. However, the production of reactive and potentially toxic metabolites is feasible with associated toxicity implications (Chen *et al.*, 2011).

With the current emphasis on research and development of medicinal plant worldwide, it is important to have some information regarding the toxicity potential and efficacy of plants utilized ethnobotanically to treat ailments. As part of ethnopharmacological studies of medicinal plant available literature should be searched for known toxic properties of plants of interest before embarking on biological activity studies. However, where toxic effects are unavailable, the inclusion of cytotoxicity and other toxicity protocols in the study are useful in detecting

potential toxicity. This strategy is applicable when screening plant extracts or isolated natural products for some other biological activities such as anti-infectious, anti-inflammatory, antioxidant, anti-diarrhoea and anti-parasitic property. The aim of this work was to determine the potential risk of the crude phenolic-enriched extracts by evaluating the cytotoxicity using Vero cell lines.

7.1. Materials and Methods

7.1.1. Preparation of plant extract

The plant extracts were prepared as described in section 3.6.3. The dried sample were reconstituted in 70% acetone at the concentration of 1.0 mg/ml (3 ml) and from it a serial dilution of the concentration range of 1.0 to 0.001 mg/ml were made on the 96 well tissue culture plate.

7.1.2. Cytotoxicity assay against Vero cell

Cytotoxicity of the extract was determined by MTT [3-(4, 5-dimethylthiazol-2-yl)-2, 5 diphenyltetrazolium bromide] assay (Mosmann, 1983) using Vero African green monkey kidney cell lines. The cells were cultured in Minimal Essential Medium (MEM) Earle's Base, supplemented with 20 mM L-glutamine, 16.5 mM NaHCO₃ supplemented with 0.1% gentamicin and 5% foetal calf serum. Confluent monolayer culture suspensions of the cells were seeded into 96-well tissue culture microtitre plate at a density of 0.5×10^3 cells per well and incubated for 24 h. at 37°C in a 5.0% CO₂ incubator. The cells were washed with cultured media and extract (1.0, 0.1, 0.001 mg/ml), positive control (berberine at concentrations of 100, 10, 1.0, 0.1 µg/ml) were added and incubated for 5 days. The cells were observed using inverted microscope to check for cytopathic effect from the extract. The cells proliferation and viability was examined by addition of 30 µl of a 5 mg/ml solution of MTT in PBS to each well and incubated for another 4 h. at 37°C. The medium was carefully removed from the wells without disturbing the MTT concentrate and washed twice with PBS. The liquid was aspirated from the cells and 50 µl of DMSO was added to each well to dissolve the crystallized MTT formazan. The amount of reduced MTT was measured as absorbance at 570 nm using microtitre plate reader. The result expressed as a percentage of the control cells and IC₅₀ was calculated

Dose response curves were obtained by plotting the percentage growth of cells versus log concentration of the compound. The LC₅₀ (50% inhibitory concentration) values were calculated from a non-linear regression model (best fit curve) of sigmoidal dose-response curve (variable) and computed using GraphPad Prism 5.04 (Graphpad, USA).

7.2. Results

The cytotoxicity of phenolic-enriched crude leaf extracts of the 19 medicinal plants used ethnopharmacologically in treating diarrhoea and other GIT disorders is presented in Table 7.1. The results indicate that the extracts had varying degrees of toxicity to Vero cell lines with LC₅₀ ranging from 3.51 ± 2.03 to 741.90 ± 44.22 µg/ml. The most cytotoxic extract was *Combretum woodii* (3.51 ± 2.03 µg/ml) followed by *Combretum vendae* (5.70 ± 1.25 µg/ml)

while the least cytotoxic extract was *O. mucronata* (741.90±44.22 µg/ml) followed by *Maytenus procumbens* (187.71±19.92 µg/ml).

Table 7.1 The LD₅₀ of the cytotoxicity assay of some medicinal plants used in South African traditional medicine to treat diarrhoea and related ailments

Plant species	LC ₅₀ (µg/ml)
Bab	17.90±2.56
Bag	35.68±2.15
Bap	40.68±18.13
Bav	76.37±7.50
Cam	ND
Cob	48.81±6.15
Cop	9.03±0.20
Cop	5.70±1.25
Cow	3.51±2.03
Com	-ND
Euc	31.61±4.04
Eun	26.99±4.48
Erl	ND
Fic	ND
Fig	ND
Inc	ND
Mpd	89.41±16.37
Mpr	187.71±19.92
Mse	87.62±3.03
Mun	99.17±11.88
Ozm	741.90±44.22
Ozp	16.58±1.85
Sle	25.09±2.40
Spd	22.30±2.42
Spt	50.62±4.30
Scb	ND
Syp	ND

ND = not determined

7.3. Discussion

For medicinal plant extracts to be useful in clinical application, the preparation must be selectively toxic to the targeted organism or interfere directly with specific reaction pathway without a major effect on the host cell or interference with normal physiological pathways. In categorization of crude extract safety, IC₅₀ value of 20 µg/ml and below were considered to be cytotoxic in an in vitro assay according to US National Cancer Institute (NCI) plant screening program (Kuete *et al.*, 2011) following incubation for more than 48 h. Some of the phenolic-rich crude leaf extract of the medicinal plants tested in this study are relatively toxic compared to the positive berberine control.

The cellular toxicity effects of the crude extracts were evaluated by MTT-formazan viability assay. Cellular viability and proliferation are considered to be an important functional characteristic of healthy growing cells. Increase in cell viability indicate cell proliferation, while decrease in cell viability indicate cell death as a result of either toxic effects of the test extracts or sub optimal culture conditions. With the cell viability of the negative control (DMSO) at the highest concentration of 1000 µg/ml under the same experimental condition, the latter postulate is eliminated. Therefore, all the phenolic-enriched extracts of the medicinal plants tested may be suggested to be safe for use in treating diarrhoea if the dosage is below the cytotoxic level. Although, *Ozoroa paniculosa* (16.58±1.85 µg/ml), *Searsia pendulina* (22.30±2.42 µg/ml), *Searsia leptodictya* (25.09±2.40 µg/ml) and *Euclea natalensis* (26.99±4.48 µg/ml) are within the defined cytotoxicity range, therefore the use of these extracts in traditional medicine need to be monitored carefully. It is also important to note that no report of toxicity has been recorded for the traditional use of these plant extracts. One should however remember that cellular toxicity does not necessarily equate to whole animal toxicity due to possibly interactions in the gut and bioavailability issues.

C. woodii acetone extracts have however been reported to be toxic in an *in vivo* test as anticoccidial in poultry at concentration of 160 mg/kg (Naidoo *et al.*, 2008). Furthermore, several cytotoxic and anti-tumour derivative of stilbenoids such as Combretastin A and Combretastatin B5 (IC₅₀ value of 10 µg/ml) have been isolated from the genus *Combretum*.

Toxicity is usually encountered due to irrational use causing accumulation of potentially toxic constituents or interactions between herbal medicinal products and conventional therapies. Indicative observations of toxicity is alterations of one of the clinical signs such as diarrhoea, weight loss, agitation, hispid hair, convulsions, tremors, dyspnoea among other) and mortality (Caparroz-Assef *et al.*, 2005).

Some medicinal plant metabolites can cause GIT toxicity. The mechanism of action can be primarily irritative or cytotoxic in nature resulting in an initial release of mucus from goblet cells, hypersecretion from crypt cells, and maladsorption causing diarrhoea and emesis. Administration of high dose of some phytochemicals can cause effects such as necrosis, haemorrhage, and even ulceration on the GIT. Medicinal plant toxins can have additional toxicity or more directly life-threatening effects on other organ system.

7.4. Conclusion

These results are important because they show that there are risks of toxicity with an inappropriate use of some of these extracts as therapeutics for any ailments. *In vivo* acute toxicity studies may be necessary to establish the safety level of the extracts as *in vitro* assay results not necessarily translate to *in vivo* activity. Long term effect of the use of the extracts such as mutagenicity and genotoxicity also need to be determined.

In vivo animal studies are frequently very expensive and requires much work to establish changes in enzyme concentrations or histological evaluation of toxicity. It is also possible to do *ex vivo* studies using isolated organs.

In the next chapter some *ex vivo* studies will be described to investigate the possible mechanism of activity of two selected species.

CHAPTER EIGHT

Motility modulation potential of *Bauhinia galpinii* and *Combretum vendae* phenolic-enriched leaf extracts on isolated rat ileum

8.0. Introduction

Gastrointestinal tract (GIT) used the smooth muscle of the mucosal lining enriched with an enteric neural network to regulate propulsive transport and mixing of food material directionally through the digestive systems (Wood, 2004). The neural network initiates and coordinates secretion and absorption across the intestinal lumen as well (Bohn and Raehal, 2006). The enteric neurons function independent of the central nervous system (CNS), therefore referred to as enteric nervous system (ENS). Enteric nervous system controls the motility and contractility of the GIT as its rate and intensity of contraction regulates the absorption of fluid, and expulsion of solid material. Therefore ENS exhibit significant role in GIT disorders such as diarrhoea and constipation through these means. Neurotransmitters such as acetylcholine (ACh), serotonin (5-hydroxytryptamine (5-HT)), substance P, histamine and opioids are the important chemical mediators in contractile regulatory actions of ENS (Farthing, 2002). The activities of the neurotransmitters in the intestine are coordinated by a large number of receptors and sub-receptors. Some of the receptors have been proved to play essential roles in GIT disorders such as peristaltic colonic motility, diarrhoeal and constipation diseases.

Some of the diarrhoea aetiologies such as infectious pathogens or their toxins, inflammatory mediators and oxidation by-products targets to control the peristaltic colonic movement by manipulating the ENS, and also control fluid and electrolyte movement across the intestinal mucosa (Guttman and Finlay, 2008). The modulations in the quantity of the neurotransmitters or the activity of the receptors can have enormous effects on intestinal motility and contraction. The process may help in regulating absorption or secretion of fluid and electrolyte by the intestine; hence provide relief against GIT disorders including diarrhoea and constipation diseases (Sikander *et al.*, 2009).

Enteric nervous system presents an attractive potential target for pharmacological intervention in diarrhoea. The use of agonists and antagonist that target these ENS hormone receptors are routinely used clinically to modulate intestinal motility, absorption and secretion. Antispasmodic or antimotility (atropine, clonidine and deodorized tincture opium), and antisecretory agents (racecadotril, octreotide) are used to treat or prevent smooth muscle contraction and control intestinal secretion, thus alleviating many symptoms of GIT disorder including diarrhoea. However, prolonged uses of these drugs are often associated with some side effects such as dry mouth and urinary retention for antimuscarinic drugs, headache, nausea, vomiting and constipation for calcium blockers. Several medicinal plants are used by different traditional cultures across the world in alleviating GIT disorders clinically manifesting as diarrhoea without reported cases of adverse effects. These provide the rationale in continuous search for safer and efficient drugs from plant phytochemicals that might target a specific receptor.

In South Africa and other developing countries treatment of gastrointestinal disorders such as diarrhoea with medicinal plants are particularly common in rural areas. The antidiarrhoea activities of medicinal plant extracts can be exhibited through spasmolytic effects (intestinal smooth muscle relaxation), delay gastrointestinal transit, suppress gut motility, stimulate water absorption or reduce electrolyte secretion. In contrast, the mechanism of actions of medicinal plants used in constipation include spasmogenic effects (intestinal smooth muscle contraction), rapid gastrointestinal transit, activated gut motility, suppressed water absorption or increase electrolyte secretion (Gilani *et al*, 2005a). All these effects are related to the regulation of ENS motility and contractility. However, scientific evaluations of the therapeutic claims as well as mechanisms of action are still unreported for many of the antidiarrhoeal plants used in traditional medicine. The aim of this study therefore is to evaluate motility regulatory potentials and determine possible mechanism of action of phenolic-enriched leaf of *Bauhinia galpinii* and *Combretum vendae* as antidiarrhoea medicinal plants on isolated rat ileum.

8.1. Drugs and reagents

Acetylcholine hydrochloride (ACh), serotonin (5-HT), nicotine, Histamine, Prostaglandin E₂ (PGE₂), Prostaglandin F_{2α} (PGF_{2α}), N^G-nitro-L-arginine methyl ester (L-NAME), Carbachol, Pilocarpine, Cyclopiazonic acid, Dimethylsulphoxide (DMSO), Sodium chloride (NaCl), Potassium chloride (KCl), Calcium chloride (CaCl₂), Sodium bicarbonate (NaHCO₃), Magnesium sulphate (MgSO₄), Potassium hydrogen phosphate (KH₂PO₄), Glucose, and carbogen

8.2. Animals

Male Wistar rats (250-300 g) obtained from University of Pretoria Biological Research Centre (UPBRC), Faculty of Veterinary Science, Onderstepoort, Pretoria were used. All animals were housed under standard environmental conditions and provided with food and water ad libitum. All the procedures were in accordance with the guidelines for use of experimental animals established by the Animal Use and Care Committee (AUCC), University of Pretoria based on specification in the South African National Standard (SAN 10386-2008). The approval of ethical committee at Faculty of Veterinary Science, University of Pretoria was obtained before the start of the work. The project was also approved by Faculty of Veterinary Science, University of Pretoria research committee (UP-RESCOM) with approval number of V027-10.

8.2.1. Isolated ileum preparation

The animal was humanely sacrificed with inhalation of isoflurane and dissected immediately. The ileum was removed and placed in carbogenated (95% O₂ and 5% CO₂) Krebs solution with the following composition (g/l): NaCl, 6.94; KCl, 0.354; KH₂PO₄, 0.163; NaHCO₃, 2.1; MgSO₄, 0.370; CaCl₂, 0.367; glucose, 2.07 and pH 7.4.

The intestinal content was removed by washing with Krebs's solution and the mesenteric constituents were eliminated. Longitudinal segments (1.5–2.0 cm) obtained from the distal ileum were placed in a 50 mL thermostatically controlled (37°C) organ bath containing Krebs solution gassed with carbogen. The preparations were connected to an isotonic transducer (load 0.5 g) in such a way as to record contractions mainly from the longitudinal axis and allowed to equilibrate for 60 min before the start of experiment: contractions were recorded using Bioscience transducers.

8.3. Contractility test

8.3.1. Spasmogenic assays

The crude extracts was prepared in stock solution of 20 mg/ml in DMSO and cumulatively added to the organ bath from concentration of 10, 25, 50, 100, 250, 500, 750 and 1000 µg/ml. The effective concentration of DMSO in the waterbath was less than 5% in all the experiments. Effect of the extracts on spontaneous motility of the ileal preparations were monitored at 20 min contact time for each concentration and cumulative dose-dependent curves for the extracts were determined to measure stimulatory effects.

8.3.2. Spasmolytic assays

8.3.2.1. Effects on acetylcholine-induced contraction

Acetylcholine hydrogen chloride was added to the organ bath cumulatively in the absence of test extracts at concentration ranging between 0.01-1.00 µg/ml in water. The process was repeated with addition of ACH (0.01-1.00 µg/ml) after 20 min pre-incubation of the isolated ileum with the extracts (10, 25, 50, 100, 250, 500, 750 and 1000 µg/ml).

8.3.2.2. Effects on Serotonin-induced contraction

Serotonin was added to the organ bath cumulatively in the absence of test extracts at concentration ranging between 0.001 - 0.1 µg/ml. The process was repeated with addition of 5-HT after 20 min pre-incubation of the isolated ileum with the extracts (10, 25, 50, 100, 250, 500, 750 and 1000 µg/ml).

8.3.2.3. Effects on K-induced contraction

The isolated ileum preparation was washed with K⁺ free Krebs's solution (composition (g/l): NaCl, 6.94; KCl, 0.354; KH₂PO₄, 0.163; NaHCO₃, 2.1; MgSO₄, 0.370; CaCl₂, 0.367; glucose, 2.07 and pH 7.4) for 20 min after equilibration and incubated with the extracts for 20 min. Thereafter, KCl solution (100 µl) was added cumulatively.




Fig. 8.1. Schematic presentation of the contractility assay

8.4 Data analysis

The inhibition of ileum contraction by test sample was normalized and expressed as a percentage of mean \pm SEM from 3-4 experiments of the references responses induced by acetylcholine (10 μ g/ml), other spasmogens, receptor agonists and antagonists using the following formula:

$$\% \text{ Inhibition} = [A_C - A_T/A_C] \times 100$$

Where A_C is the amplitude (cm) of the ileum contraction induced by the agonists and antagonists in the absence of the test sample; A_T is the amplitude (cm) of the ileum contraction by the agonists and antagonists in the presence of the test sample. The changes in EC_{50} will be used to compare the effect of the extracts using an ANOVA.

8.5. Results.

8.5.1. Effect of *B. galpinii* crude extract on isolated rat ileum

The 70% acetone extract which should have high concentration of phenolics of *B. galpinii* (10 - 1000 μ g/ml) stimulate spontaneous contraction of the rat ileum as shown in Fig 8.2 with EC_{50} value of 27.85 μ g/ml. Maximum contraction (E_{max}) of 44 mm was obtained at 200 μ g/ml and additional doses causes suboptimal response but increase duration of response caused an irreversible spasm at the maximum dosage of 1000 μ g/ml. Repeated administration of the extract at maximum dosage (1000 μ g/ml) caused exhaustion of the ileum.

Effects of the extract on acetylcholine, serotonin, K^+ induced contractions and acetylcholine in the presence of atropine (acetylcholine non-specific muscarinic receptors antagonist) indicated dual mechanisms of being an agonist (prokinetic) and an antagonist (relaxant) agent. The extract also exhibited additive contractility activity with acetylcholine and agonistic tendency to serotonin-induced contraction of the isolated rat ileum (Fig 8.3 and 8.4).




Fig.8.2: Stimulatory effect of 70% acetone leaf extract of *B. galpinii* on spontaneous contractility of isolated rat ileum and the antagonised effect of atropine.




Fig. 8.3. Effect of 70% acetone leaf extract of *B. galpinii* (200 µg/ml) on the acetylcholine cumulative concentration-effect curves in the presence and absence of atropine

From the concentration-response curve (CRC) for acetylcholine-induced contraction, the EC_{50} value in the absence of *B. galpinii* was 0.033 µg/ml and the EC_{50} in the presence of *B. galpinii* was 0.049 µg/ml. The stimulation of spontaneous contraction and agonistic effects on acetylcholine-induced contraction were partially abolished by atropine (Fig 8.3). In the CRC for serotonin-induced contraction, the EC_{50} value in the absence of *B. galpinii* was 0.0025 µg/ml and the EC_{50} in the presence of *B. galpinii* was 0.0014 µg/ml. In contrast, the *B. galpinii* extract resulted in a concentration-dependent spasmolytic effect (antagonist) on K^+ -induced contraction of the isolated rat ileum (Fig. 8.5) with maximum effect (E_{max}) of 40.66 ± 5.13 mm at concentration of 200 µg/ml.




Fig. 8.4. Agonised effect of 70% acetone leaf extract of *B. galpinii* (200 $\mu\text{g/ml}$) on serotonin induced-contraction on rat isolated ileum.




Fig. 8.5. Relaxant effect of 70% acetone leaf extract of *B. galpinii* on KCl induced contractility of isolated rat ileum

8.5.2. Effect of *C. vendae* crude extract on isolated rat ileum

The phenolic-enriched extract leaf extract of *C. vendae* do not stimulate spontaneous contraction (spasmogenic) of the isolated rat ileum, we therefore conclude that the extract have spasmolytic potential. The crude extract of *C. vendae* exhibited concentration-dependent spasmolytic effect on acetylcholine-induced contraction with EC_{50} values of 0.037, 0.027, 0.117, 0.365, and 0.396 $\mu\text{g/ml}$ at the concentration of 0, 100, 200, 400, and 600 $\mu\text{g/ml}$ of *C. vendae* in the organ bath (Fig 8.6) and concentration-dependent spasmolytic effect on serotonin-induced contraction of isolated rat ileum with EC_{50} value of 0.0017, 0.0044 and 0.012 $\mu\text{g/ml}$ at the concentration of 0, 100, 200 $\mu\text{g/ml}$ of *C. vendae* in the organ bath respectively (Fig 8.7). Equivalent volume of the solvent (5% DMSO) used in dissolving the extract had no effect on the spontaneous contraction or on 5-HT-induced contraction.




Fig.8.6. Spasmolytic effect of 70% acetone leaf extract of *C. vendae* on Ach-induced contractility of isolated rat ileum




Fig.8.7. Relaxant effect of 70% acetone leaf extract of *C. vendae* on 5-HT-induced contractility of isolated rat ileum

Addition of depolarised KCl solution (80mM) caused sustained contractions which were inhibited by *C. vendae* phenolic enriched leaf extracts in concentration-dependent response (Fig. 8.8). Therefore, agent that inhibits contraction induced by depolarised KCl solution is considered to be a calcium channel blocker (Godfraind et al., 1986). The spasmolytic effects were reversible and the spontaneous contraction returned to normal after washing three times with kerbs' solution.




Fig. 8.8. Spasmolytic effect of the *C. vendae* on the depolarised KCl-induced isolated rat ileum contractions

8.6. Discussion

Gastrointestinal motor tone is modulated through multiple physiological mediators which include neurotransmitters, inflammatory mediators and oxidative metabolites (Hoogerwerf and Pasricha, 2006). The release of these chemical modulators in GIT causes stimulatory effect mediated through an ultimate increase in cytosolic Ca^{2+} (Burks, 1987). Drug substances with ability to block or alter any of the above pathways or with non-receptor specific inhibitory action such as Ca^{2+} antagonists could be considered to be effective as therapeutic agent in hyperactive or hypoactive GIT disorders. These are important in control or alleviating diseases such as diarrhoea, constipation, emesis and dyspepsia. To study the pharmacology and possible mechanism of smooth muscle excitatory or inhibitory effect of drugs and medicinal plant extracts isolated tissue preparations of laboratory animal are usually used for *in vitro* assays.

Acetylcholine (ACh) is a neurotransmitter released by the parasympathetic nervous system mediating its action in the GIT by stimulation of nicotinic acetylcholine receptors (nAChR) and muscarinic acetylcholine receptors (mAChR). In the GIT, five subtypes of the muscarinic receptors, namely M_1 , M_2 , M_3 , M_4 and M_5 have been identified (Tobin *et al.* 2009). However, M_2 and M_3 receptors play some essential roles in the smooth muscle contraction/relaxation of GIT (Matsui *et al.*, 2002; Takeuchi *et al.*, 2005; Unno *et al.*, 2005). Through this mechanism, acetylcholine plays a critical physiological role in regulating the peristaltic movements of the GIT (Brown and Taylor, 1996). The possible mechanisms responsible for contractility mediating action of drugs including medicinal plant extracts may include one or combinations of:

- Stimulation/inhibition of ACh release from the cholinergic nerve endings.
- Stimulation/inhibition of acetylcholinesterase (AChE) enzyme at the neuro-effector junction.

- Direct activation/inactivation of the muscarinic receptors of all smooth muscles, including those of GIT.

The effects of serotonin in the ENS are complex and diverse including modulation of smooth muscle function (promoting both contraction and relaxation), potent intestinal secretagogue (predominantly pro-secretory) and responses to visceral pain. Serotonin (5-HT) receptors found within the ENS and motor neurones of the GIT include 5-HT₁, 5-HT₂, 5-HT₃, 5-HT₄, 5-HT₅, 5-HT₆ and 5-HT₇. However, only 5-HT₁, 5-HT₂, 5-HT₃, 5-HT₄ and 5-HT₇ receptor subtypes are known to affect GIT motor functionality.

The 5-HT₃ and 5-HT₄ receptors are the most studied subtypes with the regards to physiological function and histological distribution in GIT (Chetty *et al.*, 2006; Cellek *et al.*, 2006). The 5-HT₃ receptor induces a rapid depolarization of the myenteric neuron through enhancing ACh release (Kim, 2009), while 5-HT₄ receptor expressed in the nerve terminal facilitates the releases of neurotransmitters including ACh, substance P and vasoactive intestinal peptides (Kim, 2009; Wouters *et al.* 2007). These cellular events of 5-HT lead to an up-stream regulation enhancing the excitatory activity of GIT smooth muscles through mediating the ACh release. Serotonin is involved in cholera toxin-and bile salt-induced fluid and electrolyte secretion by activating the ENS.

Contractions of all smooth muscles, including those of GIT depend on the presence of Ca²⁺. Increase and decrease in intracellular free Ca²⁺ are the principal mechanisms that initiate contraction and relaxation respectively in smooth muscle (Sanders, 2001). Agonists-induced contractions are related to the release of intracellular Ca²⁺ from sarcoplasmic stores and extracellular Ca²⁺ influx through L-type channels (Makhlouf, 1994). Therefore smooth muscle relaxations can be effected by antispasmodic drugs through the inhibition of Ca²⁺ entry or release into the cells. Exposure of smooth muscle cells to high concentration of K⁺ (>30 mM) stimulate contractions through opening of voltage-dependent L-type Ca²⁺ channels and influx of extracellular Ca²⁺ (Bolton, 1979; Godfraind *et al.*, 1986).

The results obtained in this work indicated that phenolic-enriched crude leaf extract of *B. galpinii* contracted the rat ileum dose-dependently and its initial contractile phase was partially blocked by atropine, a naturally occurring alkaloid and a well known non-selective muscarinic receptor antagonists. Atropine competes with ACh and other muscarinic agonists for a common binding site on the muscarinic receptor. This result shows the involvement of cholinergic muscarinic receptors alongside with other stimulatory receptors exhibiting initial contraction by *B. galpinii* on isolated rat ileum.

The phenolic-enriched crude leaf extract of *B. galpinii* also exhibited dose-dependent stimulating activity on serotonin-induced contraction of isolated rat ileum. The spasmogenic effects of the extract on ileum longitudinal muscle may be direct serotogenic activation of 5-HT receptor pathways or through the enhanced release of other neurotransmitters without serotogenic potential.

Addition of KCl (80 mM) caused sustained contractions which were inhibited by *B. galpinii*. Therefore the inhibitory effect of the crude extract of *B. galpinii* against K⁺-induced contractions can be as result of the blockade

of Ca^{2+} channels. Thus it can be concluded that *B. galpinii* has a dual-mechanism of action (prokinetic and relaxant) on gastro-intestinal motility, depending on the prevalent patho-physiological condition. The *B. galpinii* 70% acetone leaf extract can therefore be clinically relevant as therapeutic agent in diarrhoea and constipation which are both diseases with aetiology based on motility disturbances to a large extent.

Fumaria indica crude extract also has dual-spasmogenic and spasmolytic effects on isolated organs (Gilani *et al.*, 2005a). The aqueous-ethanolic extract (80% ethanol) of the aerial parts of *Hibiscus rosasinensis* Linn (Malvaceae) contains spasmogenic and spasmolytic constituents mediating their effect through cholinergic receptors activation and blockade of Ca^{2+} influx, respectively (Gilani *et al.*, 2005b). Crude aqueous leaf extracts of *Morinda morindoides* (Baker) Milne-Redh (Rubiaceae) agonise spontaneous contractility of isolated rat ileum (Cimanga *et al.*, 2010). The petroleum ether soluble fraction and the crude saponin constituents of the extract are responsible for the spasmogenic activities. The spasmogenic and spasmolytic effect of a particular medicinal plant extract on the isolated ileum depends on predominant phytochemical constituents. Phenolic compounds exhibit spasmolytic activity while saponins are responsible for the spasmogenic activities of many plant extract preparation. From the phytochemical analysis of the extract of *B. galpinii*, the extract contains high content of phenolics. However, the result obtained in this study indicated that the crude extract of *B. galpinii* also contains other active ingredients with spasmodic effect higher than the anticholinergic effect of the phenolics.

C. vendae extract did not stimulate spontaneous contractility of the rat ileum. Further investigation of its effects on ACh-induced contraction led to a concentration-dependent inhibitory activity against ACh contraction of the rat ileum. Anti-tractility effects of *C. vendae* against ACh-induced contraction are similar to atropine indicating that the extract may be acting via nAChR or mAChR.

Addition of KCl (80 mM) caused sustained contractions which were inhibited by both *B. galpinii* and *C. vendae* phenolic enriched leaf extracts in concentration-dependent response. Agents that inhibit contraction induced by KCl are considered to be a calcium channel blocker (Godfraind *et al.*, 1986). The spasmolytic effects were reversible and the spontaneous contraction returned to normal after washing three times with Ca^{2+} free-krebs solution.

The results indicate that *C. vendae* extract is capable of mediating spasmolytic effects on isolated rat ileum through multiple inhibitions of a wide range of contractile stimuli, such as neurotransmitters (acetylcholine and serotonin) and high potassium (depolarizing stimulus). This suggests that the ileum relaxant effects of the extract are not specific to a type of receptor but rather due to either general receptor inactivation or membrane depolarization. Muscarinic receptor antagonists, 5-HT receptor antagonist and Calcium channel blockers of the L-type are known to be effective as antispasmodic, anti-motility and antidiarrhoeal agents (Lee *et al.*, 1997; Brown and Taylor, 2006; Pasricha, 2006). Hence, the presence of multiple acting spasmolytic activities in the plant extract might be contributing towards its effectiveness in diarrhoea and abdominal spasm. The isolated triterpenoids such as ursolic acid, maslinic acid, corosolic acid, asiatic acid and arjunolic acid from the plant also

have good antimicrobial activity and the stilbenoid glycosides such as combretastatin B5-O-2'- β -D-glucopyranoside and combretastatin B1-O-2'- β -D-glucopyranoside has good antioxidant activity. Such activities of the plant could account for additional benefits providing a wider cover for its use in diarrhoea of different aetiologies. This is also in accordance with the general understanding that plants contain multiple active constituents with effect enhancing activities ([Gilani and Rahman, 2005](#)).

8.7. Conclusion

The result indicated the *B. galpinii* have dual activities with the capacity of acting as prokinetic and spasmolytic agent while *C. vendae* acts as spasmolytes against the three spasmogens used to induce contraction of the ileum. Further studies aiming to identify the targeted receptor subtype and the type of interaction with muscarinic receptors as well as the identification of the main active principle are needed.

The results indicate that there is a scientific rationale for using extracts of these plant species to treat diarrhoea in humans or animals. In the next section some of the antimicrobial and anti-oxidant compounds present in these extracts will be isolated and characterized.

CHAPTER NINE

Isolation and characterization of antimicrobial and antioxidant compounds from *Bauhinia galpinii* and *Combretum vendae*

9.0. Introduction

One of the cardinal objectives in medicinal plant research and development is identification, isolation and characterization of the bioactive components present in an extract. Medicinal plant extracts are inherently complex mixture of diverse chemical components. Separation of the active components of plant phytochemicals from the inactive components are categorized into three parts: extraction, purification and chromatography. Extraction and purification involved sample preparations schematically represented in Fig. 9.1 and Fig. 9.2. Various chromatographic methods are available for qualitative and quantitative (TLC fingerprint, high performance liquid chromatography (HPLC) fingerprints) as well as for isolations (open column chromatography (OCC), vacuum liquid chromatography (VLC), HPLC, high-speed counter-current chromatography (HSCCC), gas-liquid chromatography (GLC) and/or gel permeation chromatography (GPC)). The principles of separation are based on molecular size, adsorption to the stationary phase, polarity and solubility in the mobile phase.

Structural information on isolated compounds are usually obtained from different spectroscopic techniques namely: nuclear magnetic resonance spectroscopy (NMR), mass spectroscopy (MS), and to a lesser extent infrared spectroscopy (IR) and ultraviolet-visible spectroscopy (UV-visible). The characteristic features of each NMR experiments are summarized in Table 9.1.

Table 9.1 NMR experiments commonly applied for natural product structural elucidation (Simpson *et al.*, 2011)

NMR experiment	Information/interpretation
Proton NMR (1D ^1H NMR)	Quantitative overview of the distribution of protons in a sample.
Carbon-13 NMR (1D ^{13}C NMR)	Can provide a quantitative overview of the carbon distribution.
Distortionless enhancement through polarization transfer (DEPT) (1D ^{13}C NMR)	Separate the carbon of a compound into primary (CH_3), secondary (CH_2), Tertiary (CH) and quaternary (C) spectra.
^1H - ^1H Correlation spectroscopy (COSY) (2D ^1H NMR)	Connectivity information of protons on adjacent carbons. Cross-peaks connect the chemical shifts of protons that are coupled. Symmetrical cross peaks appear around a central diagonal.
^1H - ^{13}C Heteronuclear single quantum correlation (HSQC) and heteronuclear multiple quantum correlation (HMQC) (2D ^1H - ^{13}C NMR)	^1H - ^{13}C 1 bond correlation. Cross peaks represent carbon chemical shifts in one dimension and proton chemical shifts in the other dimension.
^1H - ^{13}C Heteronuclear bond multiple correlation (HMBC) (2D ^1H - ^{13}C NMR)	^1H - ^{13}C 2–4 bond correlations. Quaternary carbons are observed. Connectivity information is read as vertical lines.

9.1.1. Column chromatography

Isolation of such compounds is usually carried out by open column chromatography under gravitation force using silica gel, sephadex, polyamides or reverse phase (RP) mode on C8 or C18-bonded silica gel stationary phase. The separation of individual compounds from the complex extract mixture is based on the compound characteristic ability for the stationary phase in the column relative to the polarity of the mobile phase. Changing the polarity (gradient elution) of the mobile phase will allowed all target compounds to elute in a sequential manner. The chromatographic process should be rapid, do not lead to decomposition of compounds, material loss, or formation of artefacts.

Open column chromatography is simple, cheap and universally practiced despite some obvious disadvantage of method being slow and often produces irreversible adsorption of sample onto the stationary phase. The method is also encumbered with large sample and solvent requirement. The bioactive compounds of interests in this project are non volatile.

9.1.2. Mass spectrometry

Mass spectrometry (MS) is an important physico-chemical tool applied for structural elucidation of compounds from natural products including medicinal plants. The fundamental principle of MS is the use of different physical means for sample ionization and separation of the ions generated based on their mass (m) to charge (z) ratio (m/z) (Rijke *et al.*, 2006). The ionization techniques available include electrospray ionization (ESI), atmospheric pressure chemical ionization (APCI), electron ionization (EI), chemical ionization (CI), fast atom bombardment (FAB), and matrix-assisted laser desorption ionization (MALDI) (Rijke *et al.*, 2006). Mass spectrometry has high sensitivity with detection limit of femtogram compared to NMR with sensitivity limit of nanogram range and above (Simpson *et al.*, 2011). The high sensitivity and the flexibility for hyphenation with other chromatographic technique made MS a versatile analytical instrument.

9.2. Materials and Methods

9.2.1. Preparation of plant extracts

The extraction and fractionation protocol was followed as described in Chapter 3. The schematic diagrams of the extraction, fractionation and isolation processes for *Combretum vendae* and *Bauhinia galpinii* are presented in Fig 9.1 and 9.2.

9.2.2. Bioautography

The bioautography against bacteria (*E. coli* and *S. aureus*) protocol were followed as described in chapter 4 while the TLC-DPPH antioxidant assay were carried out as described in chapter 5.

9.2.3. Isolation of bioactive terpenoids from *Combretum vendae*

The n-Hexane and ethyl acetate fractions showed one and two clear zone(s) of microbial growth inhibition respectively. The two fractions were subjected to gravitational column chromatography on silica gel

(2.5cm×73cm using 150 g silica, particle size 0.063–0.200 nm, Merck 70–230 mesh ASTM) using the solvent mixture of hexane: ethyl acetate starting with 100% n-Hexane, 99: 1, 98:2, 97:3, 96:4, 95:5, 94:6, 93:7 and finally 90:10 as mobile phase. Schematic representation of the isolation procedure is presented in Fig 9.1.




Fig. 9.1. Extraction, fractionation and isolation of bioactive compounds from the leaf extract of *Combretum vendae*

The dried fractions were reconstituted (10 mg/ml) and 10 µl of the aliquot spotted on TLC. Three mixtures of n-Hexane: ethyl acetate (90:10; 95:5 and 98:2) was used to develop the plates. The fractions with R_f corresponding to the R_f values of bioautography assay were combined. The purification of the compounds was achieved by repeated column chromatography until single spot was obtained for each compound using three different mobile phases to develop the TLC.

9.2.4. Isolation of bioactive phenolics from *Combretum vendae*

The ethyl acetate, n-Butanol and residual water fractions exhibited good antimicrobial and antioxidant activities. In TLC-DPPH assay four clear zones of antioxidant components were observed in ethyl acetate fraction. The fraction was subjected to open column chromatography under gravitational force (2.5cm×73cm using 150 g silica, particle size 0.063–0.200 nm, Merck 70–230 mesh ASTM) using the solvent mixture of dichloromethane: methanol starting with 100% dichloromethane, 98:2, 96:4, 94:6, 92:8, 90:10, 85:15, 80:20, 70:30, 60:40, 50:50 as mobile phase. The eluents were monitored with DPPH-TLC antioxidant assay and vanillin/H₂SO₄ spray reagent. The fractions with the same compounds were combined and subjected to further cleaning by re-chromatography until single spots were obtained on TLC chromatogram using three different mobile phases. Schematic representation of the isolation procedure is presented in Fig 9.1.


9.3. Isolation of compounds from *B. galpinii*

9.3.1. Isolation of bioactive terpenoid from *Bauhinia galpinii*

The n-Hexane fractions showed one clear zone of microbial growth inhibition and was subjected to open column chromatography under gravity on silica gel (2.5cm×73cm using 150 g silica, particle size 0.063–0.200 nm, Merck 70–230 mesh ASTM) using the solvent mixture of hexane: ethyl acetate starting with 100% n-Hexane, 98:2, 96:4, 94:6, 92:8 and finally 90:10 as mobile phase. The eluent were monitored using TLC and vanillin/H₂SO₄ spray. The fractions containing the target compound was combined and the chromatography process was repeated until a single spot using three mixtures of n-Hexane: ethyl acetate (90:10; 95:5 and 98:2) as mobile phases for TLC chromatogram was obtained. Schematic representation of the isolation procedure is presented in Fig 9.2.

9.3.2. Isolation of bioactive phenolics from *Bauhinia galpinii*

The ethyl acetate, n-Butanol and residual water fractions exhibited good antimicrobial and antioxidant activities. In TLC-DPPH assay four clear zones of antioxidant components were observed in ethyl acetate and butanol fractions. The fractions were subjected to open column chromatography under gravitational force (2.5cm×73cm using 150 g silica, particle size 0.063–0.200 nm, Merck 70–230 mesh ASTM) using the acidified solvent mixture of dichloromethane: methanol starting with 100% dichloromethane, 98:2, 96:4, 94:6, 92:8, 90:10, 85:15, 80:20, 70:30, 60:40, 50:50 as mobile phase. The eluents were monitored with DPPH-TLC antioxidant assay and vanillin/H₂SO₄ spray reagent. The fractions with the same compounds were combined and subjected to further cleaning by re-chromatography using silica gel or Sephadex L20 until single spots were obtained on TLC chromatogram using three different mobile phases. Schematic representation of the isolation procedure is presented in Fig 9.2.






Fig. 9.2. Extraction, fractionation and isolation of bioactive compounds from the leaf extract of *Bauhinia galpinii*

9.4 Characterization of the compounds

9.4.1 NMR spectroscopy

One dimensional (1D) (^1H and ^{13}C) and two dimensional (2D) NMR spectra (^1H - ^1H COSY, HMQC and HMBC) NMR spectra were recorded on a Varian-NMR-vnmrs 600 spectrometer with tetramethylsilane (TSM) as internal standard. Standard pulse sequences were used for homo- and heteronuclear correlation experiments. ^1H NMR spectra were measured at 599 MHz whereas ^{13}C NMR spectra were run at 150 MHz. Multiplicities of ^{13}C NMR resonances were determined by DEPT experiments. All NMR experiments were performed at constant temperature (27 °C) using software supplied by the manufacturer, employing deuteriochloroform, deuteriomethanol, or deuteriodimethylsulphoxide as solvent on the basis of solubility of the sample and literature data.

9.4.2 Mass Spectroscopy:

Electrospray ionization mass spectrometric analyses (negative and positive mode) were carried out to obtain the molecular weight and fragmentations patterns of the isolated compound(s) using TOF mass spectrometer (WATERS HPLC).

9.4.3 UV spectroscopy:

The UV-spectrum of the isolated bioactive compound(s) was recorded using Agilent 1200 UV-Visible spectrophotometer.

9.5 Results

9.5.1 Identification of the chemical structures of isolated compounds from *Combretum vendae*

The n-Hexane and ethyl acetate fractions from the acidified 70% acetone leaf extract of *Combretum vendae* A. E. van Wyk through bioassay guided fractionation were repeatedly subjected to gravity column chromatography to yield one pure, mixture of four position isomer antimicrobial triterpenoids and two stilbene glucopyranoside. The structures of the compounds were determined by extensive NMR techniques and chemical methods mainly by 1D NMR (^1H , ^{13}C and DEPT) and 2D NMR (HSQC, HMBC and COSY), ESIMS, UV-visible spectra and by comparison with the literature data.

Compound 1 was obtained as an amorphous white powder. Detailed analyses of the 1D and 2D NMR spectra indicated the presence of 30 carbons which revealed 7 methyl, 9 methylene, 6 methane, 6 quaternary, 1 carboxylic acid at δ_{C} 178.9, an olefinic broad triplet proton at δ_{H} 5.25 (H-12) coupled to a carbon at δ_{C} 125.4, a quaternary carbon at δ 138.8.0 (C-13) and a β -18 proton at δ_{H} 2.05 characteristic of signal of an ursol-12-ene skeleton (Appendix 9.1). The Comparison of these NMR data with the literature confirms the compound as **ursol-12-en-28-oic acid (ursolic acid)** (Mahato and Kundu, 1994).

Compound 2 and 3 were obtained as position isomeric mixture; the TLC fingerprint indicated unresolved single spot with three different mobile phases. However, the 1D and 2D NMR spectra exhibited chemical shift characteristic of both olean-12-ene and ursol-12-ene. ^{13}C NMR (DMSO): δ 179.7(C-28, olean), 178.0 (C-28, urs), 145.08 (C-13, olean), 122.72 (C-12, olean), 139.42 (C-13, urs), 125.55 (C-12, urs), 41 (C-18, olean), 53 (C-18, urs), 27.97 (C-19, olean), 39.07 (C-19, urs) and 46.13 (C-20, olean), 39.02 (C-20, urs) (Appendix 9.2). Comparing the data with the literature values, the two compounds in the mixture were identified to be **corosolic acid and maslinic acid** (Mahato and Kundu, 1994)

Compound 4 and 5 were also obtained as position isomeric mixture. The spectra 1D and 2D were similar to the spectra of compound 2 and 3 except the presence of additional hydroxyl at C-24) (appendix 9.3). From the correlation data and literature values, compound 4 and 5 were identified to be **asiaitic acid and arjunolic acid** respectively (Mahato and Kundu, 1994).

Compound 6 was obtained as an amorphous powder from the ethyl acetate fraction. The proton NMR spectrum shows signal for apigenin: δ_{H} 6.19 (d, $J=2.0$ Hz, H-6), 6.38 (d, $J=2.0$ Hz, H-8), 7.2 (d, $J=2.1$ Hz, H-3), 6.50 (d, $J=8.9$ Hz, H-3' and H-5'), and 7.30 (d, $J=2.1$ Hz, 8.9, 2H' and H-6').

Compound 7 was obtained as creamy glass-like solid mass and deduced to have molecular formula $\text{C}_{23}\text{H}_{29}\text{O}_{11}$ with molecular weight 481 by EIMS. The ^{13}C -NMR spectrum indicated 17 carbon signals due to aglycone and six carbon signals of a glycoside group. The signal consist twelve aromatic signals, three methoxy and two aliphatic (Appendix 5). The ^1H -NMR spectra revealed the presence of two aromatic benzylic rings. The ^1H detected heteronuclear multiple-bond connectivity (HMBC) spectrum indicated long-range correlations from ^1H -1a' (δ_{H} 2.9, 3.0) to C-1' (δ_{C} 128.55), C-2' (δ_{C} 144.35), C-6' (δ_{C} 118.95), C-1a (δ_{C} 36.96) and C-1 (δ_{C} 132.72). Long range correlation observed between ^1H -1a (δ_{H} 2.7) and C-1 (δ_{C} 132.72, C-2 (δ_{C} 106.16), C-6 (δ_{C} 106.16), C-1a' (δ_{C} 31.73) and C-1' (δ_{C} 128.55). Additional long-range correlation were observed between ^1H -2, 6 (δ_{H} 6.5) and C-1 (δ_{C} 132.72), C-3, C-5 (δ_{C} 148.16), C-4 (δ_{C} 133.79), C-1a (δ_{C} 36.96); ^1H -5' (δ_{H} 6.7) to C-1' (δ_{C} 128.55), C-3' (δ_{C} 139.71), C-6' (δ_{C} 118.95), C-4' (δ_{C} 147.26) ; and ^1H -6' (δ_{H} 6.6) to C-1a' (δ_{C} 31.73), C-3' (δ_{C} 139.71), C-4' (δ_{C} 147.26). Long-range correlation also observed between methoxyl signal (δ_{H} 3.7) and the C-3, C-5 (δ_{C} 148.16), C-4' (δ_{C} 147.26) (appendix 9.4). Based on these data, the aglycone of compound 6 was determined to be stilbenes derivative. The signals for anomeric proton and carbon (δ_{H} 4.5 and δ_{C} 106.26) indicated the presence of a sugar moiety. The long-range correlation in the HMBC experiment between the anomeric proton signal (δ_{H} 4.5) of the β -D-glucopyranosyl group and the C-2' signal (δ_{C} 144.35) confirmed the position of the attachment of the glycopyranosyl moiety on the phenolic ring. The structure of compound 6 was determined to be **combretastatin B5-O-2'- β -D-glucopyranoside**. The EIMS, 1D (^1H and ^{13}C , DEPT), and 2D (HSQC, HMBC, COSY) data correlate with literature information on the compound (Pettit *et al.*, 1985).

Compound 8 had molecular weight of 496 and the molecular formula was deduced to be $\text{C}_{24}\text{H}_{32}\text{O}_{11}$ by EIMS. The compound differs from compound 6 by the presence of extra methyl group. This was confirmed by 1D and 2D NMR spectra. From HSQC one additional methoxyl signal (δ_{H} 3.6 and δ_{C} 59.69) having long range correlation C-4 (δ_{C} 135) (appendix 9.5). The long range correlation from HMBC experiment between ^1H -4 proton of the glycopyranosyl moiety (δ_{H} 3.7) and the C-2' signal (δ_{C} 143.78) confirmed the position of the structure of compound 7 to be **combretastatin B1-O-2'- β -D-glucopyranoside**. The EIMS, 1D (^1H and ^{13}C , DEPT), and 2D (HSQC, HMBC, COSY) data correlate with literature information on the compound (Schwickard *et al.*, 2000).




Fig 9.3 Chemical structures of isolated bioactive compounds from 70% acetone leaf extract of *Combretum vendae*

9.5.2 Antimicrobial assay of isolated compound from *C. vendae*

Antimicrobial activities of the isolated compounds against standard and clinical isolate pathogens expressed as MIC are presented in Table 9.1. Some of the compounds exhibited good microbial growth inhibitory potential worthy of pharmacological considerations with MIC ranging from 3.9-31 µg/ml.

Table 9.2: Minimum inhibitory concentration (µg/ml) of the isolated compounds from the leaf extract of *Combretum vendae*

Microorganisms	Compound 1	Compounds 2 and 3	Compounds 4 and 5	Compound 7	Compound 8
<i>E. coli</i>	62	250	250	250	250
<i>E. faecalis</i>	31	31	31	31	31
<i>S. aureus</i>	62	125	125	125	62
<i>P. aeruginosa</i>	125	125	62	250	250
<i>C. albicans</i> (M0825)	15	7.8	7.8	31	31
<i>C. albicans</i> (M0824)	62	62	31	125	125
<i>C. albicans</i> (1051604)	3.9	3.9	3.9	3.9	3.9
<i>C. albicans</i> (1051608)	7.8	7.8	7.8	15	15
<i>C. albicans</i> (ATCC 10231)	7.8	15	7.8	31	31
<i>C. neoformans</i>	15	15	15	125	125
<i>A. fumigatus</i>	31	31	31	250	250

1: Ursolic acid, 2 and 3: mixture of maslinic and corosolic acid, 4 and 5: mixture of asiatic and arjulongic acid, 6: combretastatin B5-O-2'- β -D-glucopyranoside. 7: combretastatin B1-O-2'- β -D-glucopyranoside. *C. albicans* (M0824), *C. albicans* (M0825), *C. albicans* (1051604), *C. albicans* (1051608) are clinical isolate obtained from National Health Laboratory Service, Pretoria, South Africa.

9.5.3 Identification of the isolated bioactive compounds from *Bauhinia galpinii*

The combine TLC fingerprint, TLC-DPPH assay and bioactivity guided fractionation of acidified 70% acetone leaf extract of *B. galpinii* detected four major flavonoids from ethyl acetate fraction with antioxidant activity. Bioautography against fungal and bacterial pathogens revealed two microbial inhibitory growth spot in Hexane fraction and one in DCM fraction respectively. The bioactive compounds were isolated from each fraction using open column chromatography with silica gel as stationary phase. The phenolics compounds were further purified using Sephadex L-20 as stationary phase and acetone/methanol (50:50) as mobile phase at a rate of 2ml/5min. The chemical structures of the compounds were determined by detailed nuclear magnetic resonance (NMR) techniques including the one dimensional (1D) NMR (proton (^1H), carbon-13 (^{13}C) and distortion enhancement DEPT) and two dimensional (2D) NMR (HSQC, HMBC and COSY). Mass spectrometry and the fragmentation patterns of the compounds were extensively used for the structural elucidation.

Compound 9 obtained as white amorphous powder from hexane fraction was characterized by ^1H NMR spectra (in CDCl_3), ^{13}C -NMR spectra (in CDCl_3), HSQC, HMBC, DEPT and COSY. The ^1H NMR spectrum of 9 showed a one-proton doublet at δ_{H} 5.33 (J^{H} 5.5 Hz) assigned to a vinylic H-6 proton. A one-proton broad multiplet at δ_{H} 3.46 with half-width of 18.5 Hz was attributed to carbinol H-3 proton. Two three-proton broad signals at δ_{H} 0.67 and 1.01 were attributed to a tertiary C-18 and C-19 methyl protons. A six-proton broad signal at δ_{H} 0.83 was associated with C-29 and C-20 methyl protons. Two three proton doublets at δ_{H} 0.91 (J^{H} 6.2 Hz) and 0.85 (J^{H} 6.3 Hz) were due to secondary C-26 and C-27 methyl protons. The remaining methylene and methine protons appeared between δ_{H} 2.50 and 1.27. The presence of all the methyl signals in the range δ_{H} 1.01–0.67 suggested that all these functionalities were located on the saturated carbons. The ^{13}C NMR spectrum of 9 exhibited signals for vinylic carbons at δ_{C} 140.97 (C-5) and 121.96 (C-6). The two carbinol signals appeared at δ_{C} 72.04 (C-3) and


76.86 (CH₃ CH₂O-) respectively. The carbon signals in the upfield region at δ_C 12.20, 21.30, 19.25, 20.04, 19.62, 12.08 and 19.00 were associated with the methyl functionalities. The remaining methylene and methine carbon resonated between δ_C 56.99 and 23.29. The ¹H–¹H COSY spectrum of 9 showed correlations of H-6 with H2-7 and H-8; H-3 with H2-2, H2-4 (appendix 9.6). The ¹H and ¹³C NMR spectral data of steroidal nucleus of 9 were compared with related steroidal constituents (Alam *et al.*, 1994). On the basis of the foregoing discussion, the structure of 9 was elucidated as 3 β -ethoxy stigmast-5-en-ol (3 β -ethoxy sitosterol).

Compound 10 and 11 was isolated from the ethyl acetate fraction as a yellow powder by repeated gravitational column chromatography on silica gel and Sephadex L-20 stationary phases. The ¹H NMR spectrum in deuterated methanol of 10 showed signals for **quercetin**: δ_H 6.19 (d, J=2.0 Hz, H-6), 6.38 (d, J=2.0 Hz, H-8), 7.57 (d, J=2.1 Hz, H-2', δ_C), 6.86 (d, J=8.9 Hz, H-5', δ_C), and 7.56 (d, J=2.1 Hz, 8.9, H-6', δ_C). Carbon 13 NMR (100 MHz, in ppm, methanol-d₄) shows 15 signals and the data (appendix 6) correlated well with literature for 3, 5, 7, 3', 4'-pentahydroxyflavone (Quercetin) (Said *et al.*, 2009)


The ¹H NMR of compound 11 in deuterated methanol at 599.74MHz revealed the presence of myricetin aglycone: δ_H 7.383 (s, H-2',6', δ_C 107.09) for the B phenyl ring, and 6.369 (d, H-8, δ_C 92.95), 6.338 (d, H-8, δ_C 92.95), 6.172 (d, H-6, δ_C 97.80) and 6.170 (d, H-6, δ_C 97.80) for the meta substituted A phenyl ring. The structure was confirmed by ¹³C-NMR which showed 15 signals without methoxy or glycoside substituent and 2D correlation (HSQC and HMBC) (appendix 9.7). The data correlated well with literature data for **3, 5, 7, 3', 4', 5'-hexahydroxyflavone (myricetin)** (Said *et al.*, 2009)

The NMR spectrum of compound 12 was typical for a flavone with one meta-substituted and para-substituted phenolic moiety. The UV spectrum showed λ_{max} 261.27 and 369.27 nm which compared favourably with 263 and 367 nm reported by Abdurrahman and Moon, 2007 for isoetin moiety. The compound was assigned molecular formula C₁₆H₁₀O₇ with the aid of a peak observed in the ESI-MS experiment at 315.054. The carbon chemical shifts were assigned by the combination ¹H-¹³C HSQC and long range couplings in the ¹H-¹³C HMBC experiments. The ¹H-NMR spectrum was indicative of five aromatic protons, one methoxyl group. The ¹H and ¹³C NMR spectra data correlated with those of isoetin flavone moiety (Abdurrahman and Moon, 2007; Pauli and Junior, 1995; Voirin *et al.*, 1975) (Table 1). The ¹H-¹³C HSQC correlations were used to assign signal at δ_H 7.19, 6.27 and 6.40 to the protons at C-3 (s, δ_C 108.85), C-6 (d, J = δ_C 99.97) and C-8 (d, J = δ_C 94.86) positions of A ring, and the signal at δ_H 6.65 and 7.38 to the proton at C-3' (s, δ_C 101.36) and C-6' (s, δ_C 114.45) positions of B ring of the isoetin moiety respectively. The long range coupling in the HMBC presented in appendix 9.10 also supported the isoetin flavone moiety. The methoxy protons signal at δ_H 3.8 was correlated with the quaternary carbons (C-2') at δ_C 153.28 indicating the attachment of methoxy group at the carbon (appendix 9.8). From all the correlations, compound 13 was determined to be a new flavone named as **5, 7, 4' 5' tetrahydroxy-2'-methoxyflavone (isoetin 2'-methyl ether)** or 5, 7, 2' 5' tetrahydroxy-4'-methoxyflavone (isoetin 4'-methyl ether)


Compound 13 and 14 obtained as yellow powder respectively. UV spectra in MeOH showed λ_{\max} of 203.27, 255.27, 355.27 nm for compound 14, and 207.27, 257, 354.27 nm for compound 13 respectively. The ESI-MS peaks of compound 13 and compound 14 in negative mode, were observed at m/z 463 [M-H]⁻ and m/z 479 [M-H]⁻ respectively. The molecular ion of compound 13 was 16 mass units smaller than that of compound 14, which corresponds to the difference in the number of hydroxyl groups on the B-ring of the flavonol aglycone. The ¹H NMR spectrum in deuterated methanol of 13 showed signals for quercetin: δ_{H} 6.19 (d, $J=2.0$ Hz, H-6, δ_{C} 99.17), 6.38 (d, $J=2.0$ Hz, H-8, δ_{C} 94.09), 7.57 (d, $J=2.1$ Hz, H-2', δ_{C}), 6.86 (d, $J=8.9$ Hz, H-5', δ_{C}), and 7.56 (d, $J=2.1$ Hz, 8.9, H-6', δ_{C}) and anomeric protons at δ_{H} 5.192 (s, δ_{C} 104.14) and 5.22 (s, δ_{C} 104.14) characteristic of galactopyranose. The ¹H NMR of 14 in deuterated methanol at 599.74MHz revealed the presence of myricetin aglycone with five aromatic proton: δ_{H} 6.19 (s, H-6, δ_{C} 98.48), 6.38 (s, H-8, δ_{C} 93.25), 7.37 (s, H-2', H-6', δ_{C} 108.52) and similar anomeric protons to those in 13 (appendix 9.9). The ¹³C NMR and HMBC, HSQC spectra in deuterated methanol at 150MHz of 13 and 14 were very similar, except for the signals corresponding to the flavonol aglycone (appendix 9.8). The spectral data and correlation with literature information revealed the two compounds to be [quercetin-3-O- \$\beta\$ -galactopyranoside](#) (Rayyan *et al.*, 2004; Yan *et al.*, 2002) and [myricetin-3-O- \$\beta\$ -galactopyranoside](#) (Yan *et al.*, 2002) respectively.




3 beta-O-ethyl sitosterol (9)




5, 7, 4', 5' tetrahydroxy-2-methoxyflavone
(Isoetin 2'-methyl ether) (12)




3, 5, 7, 3' 4' pentahydroxyflavone
(Quercetin) (10)



3, 5, 7, 3' 4', 5' Hexahydroxyflavone
(Myricetin) (11)



Quercetin-3-O-galactopyranose (13)



Myricetin-3-O-galactopyranose (14)

Fig. 9.4. Chemical structure of bioactive compound isolated from the leaf extract of *bauhinia galpinii*

9.5.4. Antimicrobial assay of isolated compounds from *B. galpinii*

Antimicrobial potential of the isolated compounds against some diarrhoeal pathogens and organisms of other important infectious diseases are presented in Table 9.2 as minimum inhibitory concentration (MIC) ($\mu\text{g/ml}$).

Table 9.3. Minimum inhibitory concentration ($\mu\text{g/ml}$) of the isolated compounds from the leaf extract of *Bauhinia galpinii* and positive control

Microorganisms	9	10	11	13	14	Gentamicin	Amphoteric B

E.coli	125	31	31	31	62	15.5	-
<i>E. faecalis</i>	31	31	7.8	62	62	7.75	-
<i>S. aureus</i>	62	62	15	31	31	3.87	-
<i>P. aeruginosa</i>	125	62	15	15	15	3.87	-
<i>C. albicans</i> (M0825)	15	3.9	3.9	31	31	-	3.9
<i>C. albicans</i> (M0824)	31	31	31	31	31	-	3.9
<i>C. albicans</i> (1051604)	3.9	3.9	3.9	3.9	3.9	-	0.8
<i>C. albicans</i> (1051608)	62	3.9	3.9	15	15	-	1.93
<i>C. albicans</i> (ATCC 10231)	62	3.9	3.9	31	31	-	1.93
<i>C. neoformans</i>	125	31	31	125	125	-	7.75
<i>A. fumigatus</i>	125	31	31	125	125	-	7.75

1: 3 β -ethoxyl sitosterol; 2 Quercetin; 3 Myricetin, 4 Quercetin-3-O- β -galactopyranose; 5 Myricetin- 3-O- β -galactopyranose. *C. albicans* (M0824), *C. albicans* (M0825), *C. albicans* (1051604), *C. albicans* (1051608) are clinical isolate obtained from National Health Laboratory Service, Pretoria, South Africa.

9.6. Discussion

9.6.1. Bioactive compounds from *Combretum vendae*

Bioactivity guided investigation of *Combretum vendae* afforded ursolic acid ((3 β)-3-hydroxyurs-12-en-28-oic acid), mixture of asiatic and arjunolic acid, mixture of maslinic and corosolic acid, apigenin, Combretastatin B5-2'-O- β -glucopyranoside and Combretastatin B1-2'-O- β -glucopyranoside all exhibiting broad based microbial growth inhibitory potentials.

Ursolic acid and its derivatives biosynthetically derived from the cyclization of squalene (and other triterpene acids) have been extensively studied as pharmacological active molecules in many *in vitro* and *in vivo* studies. Some of the biological activities include antioxidant, hepatoprotective, anti-inflammatory, anticancer, anti-HIV, vasorelaxant (Aguirre-Crespo *et al.*, 2006) and antidiabetic activities. Several mechanisms have been proposed to explain its anti-inflammatory activity, including inhibition of secretory PLA₂ enzymes, IL-1 β secretion, iNOS and COX-2. Ursolic acid have no antispasmodic effect (Estrada-Soto *et al.*, 2007), however the broad base antimicrobial activities (3.9-125 μ g/ml) obtained in this work and other reports of antioxidant, anti-inflammatory and antidiabetic provide pharmacological bases for further investigation of the compound as anti-diarrhoeal agent.

Biological activities such as α -glucosidase inhibition of arjunolic acid (18.63 \pm 0.32 μ g/ml), asiatic acid (30.03 \pm 0.41 μ g/ml), maslinic acid (5.52 \pm 0.19 μ g/ml) and corosolic acid (3.53 \pm 0.27 μ g/ml) isolated from *Lagerstroemia speciosa* were reported. The α -glucosidase inhibition of these compounds shows their antidiabetic and antiadhesion potential against microbial pathogens both of which are also important in anti-diarrhoea therapy. However the compounds have no α -amylase inhibitory activities (Hou *et al.*, 2009). Mixture of arjunolic acid and asiatic acid isolated from *Combretum nicholsonii* have antifungal activities with MIC of 0.2-1.5 μ g/ml. However, antimicrobial activities obtained in the work is slightly higher for the mixture (3.9-250 μ g/ml), the difference in result may be due to experimental variable such as concentrations of the culture media, composition of the

mixture, incubation time and strains. These results indicate that individual compound or the mixtures have pharmacological potentiality against infectious pathogens.

Apigenin have been isolated previously from the acetone leaf extract of *C. vendae*. The antibacterial activity of the compound was evaluated (Eloff *et al.*, 2008).

Combretastatin B5-2'-O- β -glucopyranoside have been isolated previously from the seed of *Combretum kraussii*. The aglycone moiety of this compound was isolated from the acetone leaf extract of *Combretum woodii* reported to have antibacterial activity. This is the first report on the isolation of the compound from the leaf extract and its antimicrobial activities.

Combretastatin B1-2'-O- β -glucopyranoside have been previously isolated from the seeds of *Combretum kraussii* (Pettit *et al.*, 1987) and wood bark of *Combretum erythrophylum* (Schwikkard *et al.*, 2000). The stilbenes have been reported to have cytotoxic activity with effect on tubulin polymerization, the primary protein component of microtubules in cancer hence the potential of the compound as anticancer drug is being explored. The compound has been evaluated for selective inhibitory activity against the DNA-damaging repair-deficient strain of *Saccharomyces cerevisiae* deficient in the RAD52 recombination repair gene and exhibited no activity while the derivative with unsaturated bond at 1aC- 1a'C (combretastatin A1-2'-O- β -glucopyranoside) was active (Schwikkard *et al.*, 2000). This indicated the importance of the unsaturated bond in the structure-activity relationship for cytotoxicity effects. However, there is no literature report on other biological activity potential such as antimicrobial, antioxidant, anti-inflammatory and antidiarrhoea. This is the first report on the isolation of the compound from the leaf extract and its antimicrobial activities. The non cytotoxic effect of the compound against the cancer cells lines (Schwikkard *et al.*, 2000) indicates that the compound can be exploited for other biological activities.

9.6.2. Bioactive compounds from *Bauhinia galpinii*

Isoetin (5, 7, 2' 4' 5'-pentahydroxyl flavone) and its various derivatives are rare compounds formed by insertion of a 2' hydroxyl group into luteolin to give characteristic yellow pigments in some plant part (Lattanzio *et al.*, 2006). 5, 7, 4' 5' tetrahydroxy-2'-methoxyflavone (isoetin 2'-methyl ether) is a new compound (unfortunately the quantity obtained was not enough for bioassays). A related compound 5, 7, 2' 4' tetrahydroxy-5'-methoxyflavone (isoetin 5'-methyl ether) isolate from *Trihosanthes kirilowii* (Cucurbitaceae) was reported to be cytotoxic against human lung cell line A549 (IC₅₀ 0.92 μ g/ml), human melanoma Sk-Mel-2 (IC₅₀ 8.0 μ g/ml), and mouse melanoma B16F1 cell lines (IC₅₀ 7.23 μ g/ml). High cytotoxicity (IC₅₀ 2.5 μ g/ml) of the acetone root extract of *B. galpinii* against Vero cell lines has been reported (Samie *et al.*, 2009). Isolation of more 5, 7, 4' 5' tetrahydroxy-2'-methoxyflavone (isoetin 2'-methyl ether) from *B. galpinii* for further studies on cytotoxicity effect important.

Quercetin and myricetin are the common flavonol present in dietary and constitute the active component of medicinal plant with characteristic hydroxyl substitutions at the 3, 5, 7, 3', 4' (Quercetin), and 3, 5, 7, 3', 4' and 5'

(myricetin) positions of flavone ring. The compounds occur in nature mostly as glycoside with D-glucose, L-rhamnose, D-galactose or arabinose rather than free aglycone. Biological effects of phenolic compounds depend on their bioavailability which is determined by the lipophilicity of each molecule. The mechanisms involved in digestion and absorption of phenolic compound is complex but passive transports through the membrane have been proposed. The glycosides and methylated phenolics are not readily absorbed in native form but need to be hydrolysed by intestinal enzymes or colonic microflora to aglycone before absorption.

Quercetin and myricetin alongside their galactopyranoside derivatives were isolated from the acidified 70% acetone leaf extract of *Bauhinia galpinii*. Quercetin galactopyranose and myricetin galactopyranose were previously isolated from this plant and their antioxidant activity and cytotoxicity was evaluated (Aderogba *et al.*, 2007).

A wide range of biological activities related to diarrhoeal pathogenesis including antimicrobial (Naz *et al.*, 2007), anti-inflammatory and spasmolytic due to their antioxidant and/or free radical scavenging (Aderogba *et al.*, 2007) as well as ability to interfere with several enzymatic pathways have been reported for the compounds. Quercetin and myricetin are active against microorganisms of the genera *Bacillus*, *Corynebacterium*, *Salmonella*, *Shigella*, *Staphylococcus*, *Streptococcus*, and against *Escherichia coli* and *Vibrio cholerae* (Naz *et al.*, 2007).

Both compounds also have protective and promotive effects on intestinal TJ barrier function through interaction with intracellular signaling molecules, tyrosine kinases and protein kinase C δ (PKC δ) (Suzuki and Hara, 2010). The intercellular TJs are the major determinant of the intestinal physical barrier regulating the paracellular movement of ions, solute, and water through the intestinal epithelium. Impaired intestinal TJs functions are involved in several intestinal and metabolic diseases, such as diarrhoea, inflammatory bowel disease and food allergy (Suzuki and Hara, 2010). Myricetin inhibits the generation of MDA a cytotoxic by-product of lipid peroxidation of arachidonic acid liberated from membrane phospholipids (Robak *et al.*, 1986).

Myricetin has potential as an antiviral agent by its ability to inhibit the reverse transcriptase from Moloney murine leukaemia virus, Rauscher murine leukaemia virus and human immunodeficiency virus (Ono *et al.*, 1990). Myricetin also has antidiabetic activity with ability to stimulate lipogenesis and enhanced glucose uptake into adipocytes. The mechanisms postulated include changes in lipid-protein interaction or increase membrane fluidity. Myricetin inhibits the intraluminal accumulation of fluid and prevent diarrhoea induced by castor oil.

9.7. Conclusion

All the isolated compounds from the two plant species have biological activities with relevance against one or more diarrhoeal pathophysiology. The antimicrobial of some of the compounds are worthy of pharmacological consideration. In order to exploit the full potentials of these compounds some *in vitro* and *in vivo* studies are required to determine the mechanism of action.

CHAPTER TEN

General conclusion and future prospects

10. Introduction

Diarrhoea is one of the major health challenges facing the world and especially developing countries. The problem is aggravated due to the increasing number of immunocompromised people infected by HIV, with associated opportunistic infections and other health complications manifesting as diarrhoeal symptoms. The emergence of more virulent strains resulting from drug resistant pathogens and the apparent side effects of some conventional drugs currently in use is also serious concerns in diarrhoeal control and management. In animal production, diarrhoeal outbreaks usually cause serious economic losses due to reduced productivity, cost of treatment, lower level of reproduction and increased mortality.

However, the success of oral rehydration therapy in reducing mortality and lack of commercial interest in drugs for developing countries has slowed the progress in the development of novel agents for treating diarrhoeal diseases. Therefore, there is an urgent need for new therapeutic drugs or herbal products with lower cost, high efficacy and little or no side effects. Plants and plant preparations have been used ethnopharmacologically in treating diarrhoea successfully, although their efficacies, mechanisms of action and safety have generally not been proven scientifically. Thus, the overall aim of this project was to systematically determine the efficacy, mode of action and safety of some plants used traditionally in South African traditional medicine as diarrhoea therapy.

The following objectives were identified to attain this aim:

- ❖ To conduct comprehensive literature works on diarrhoeal aetiologies and mechanisms and, medicinal plants use for treating diarrhoea symptoms in Southern Africa.
- ❖ To determine the phenolic compositions of the crude extract.
- ❖ To evaluate the effects of some selected medicinal plant species against pathogenic microbes known to induce diarrhoea.
- ❖ To determine the antioxidative properties of the selected plants using various standard protocols.
- ❖ To determine the anti-inflammatory potentials of the selected plants using various standard protocols.
- ❖ To evaluate the toxicity risk of the crude extracts.
- ❖ To determine the intestinal motility modulatory effects of the most promising extracts on the contractility process of the isolated rat ileum induced by spasmogens and ion channels activators.
- ❖ To isolate and characterize the component(s) that exhibit antimicrobial and antioxidant properties from the most promising extracts.

The achieved objectives of the study are outlined as follows:

10.1. Identification of diarrhoeal pathogenesis and medicinal plants used as therapeutic agents

The data generated from the literature work indicate that diarrhoea has a number of pathogenesis such as microbial infection, chronic inflammation, oxidative injury to intestinal mucosal lining, and deranged intestinal motility. The mechanisms involved include one or combination of ionic and water secretion into the lumen and reduced absorption of fluid from the intestine. The compendium of the medicinal plants used as antidiarrhoeal agents (254 species) in Southern Africa also revealed the diarrhoeal challenges and the wide acceptability of medicinal for cure.

10.2. Antimicrobial evaluation of the extracts against infectious pathogens

The results obtained by the antimicrobial screening indicate the presence of many compounds with potent antibacterial and antifungal activity against the standard strains of microbes responsible for infectious diarrhoea and other important infectious diseases in humans and animals. The significant inhibitory activity exhibited by the water fraction of the *C. vendae* against *S. aureus* can be considered as important for the traditional use of this plant, where water is the main extractant available. Generally, the results revealed that the antimicrobial potential of the extracts are potentiated in the hexane and dichloromethane fractions. Future investigation of the potent extracts and fractions against resistant and virulent pathogens might indicate new mechanisms for the growth inhibition of the microorganisms.

10.3. Antioxidant evaluation of the extracts

The crude extracts and the polar fractions of ethyl acetate and butanol had significant radical scavenging activity, through hydrogen and proton donation mechanisms. These activities are ascribed to the presence of large quantity of phenolics. In view of the oxidative stress in the pathogenesis of diarrhoea through tissue injury by ROS/RNS involvement in lipid peroxidation, exacerbation of inflammatory processes and some of the reactive species serving as secretagogues, the strong antioxidant activity could indicate the presence of compounds with potentially important mechanisms of pharmacological relevance in reducing the deleterious effects of the oxidative species in diarrhoea. *In vitro* results however, cannot be literally translated into *in vivo* situation due the problem of bioavailability, absorption and possible metabolic transformations of the bioactive compounds in the intestine. Further research are needed to verify using other models with different mechanisms against substrates which are generated in human or animal cells as well as *in vivo* studies to evaluate their efficacy and safety.

10.4. Anti-inflammatory potential of the extracts

Inflammation is regarded as the hallmark of many diseases aetiology and the significant mediators are eicosanoids (prostaglandins, prostacyclin) from cyclooxygenase (COX) and leukotrienes from lipoxygenase (LOX) pathways. These two enzymes are the target for modulating the inflammatory process. The result obtained indicated that the polar extracts of *Bauhinia*, *Carissa*, and *Syzygium* species used for COX inhibitory assay were

active against COX-1 with no activity against COX-2 while the *Combretum* species were inactive against both enzymes. COX-1 selective inhibitors are considered to cause GIT injury while selective COX-2 inhibitors are more beneficial against inflammatory processes, therefore these plant polar extracts should be used with caution because possible intestinal injury. Most of the plant extracts however, had good LOX inhibitory activity. Current research on anti-inflammatory agent focus on dual activity as COX and LOX inhibitors since both pathways uses the same substrate. If one pathway is closed down, more substrate will be available to the other unperturbed pathway, thus increasing its products and consequently promoting some other inflammatory mechanisms. Additional work is required to determine the fraction(s) in which the active component is present and the probable mechanisms of action. Since the polar extract are not active against COX-2, the non-polar extracts or fractions still have to be tested. The plant extracts also have to be tested against other inflammatory biomarkers and mediators including an *in vivo* studies using laboratory animal model.

10.5. Toxicity risk of the extracts

The toxicity risk assessment using MTT assay (Mosmann, 1983) using Vero African green monkey kidney cell lines indicated that the extracts of *Combretum* species except *C. bracteosum* were highly toxic. The other extracts have varying degree of toxicity with *Ozoroa mucronata* being the least toxic. These results are important because they show that there are risks of toxicity with an inappropriate use of some of these extracts as therapeutics for any ailments except perhaps cancer. The toxicity of medicinal plants depends on many factors such as the plant part used, and the solvents used as extractant which are determinant of the compositional characteristics and biological activity of extracts. Most of the highly toxic extracts also contain a high quantity of hydrolysable tannin. Poor handling of raw or processed materials may produce exogenous toxic contaminants not inherent as plant phytochemicals. It should be kept in mind that the results of cytotoxicity testing may vary considerably depending on the cell type used, the initial cell density to which the extracts are exposed, and the duration of exposure. Vero cells were selected as these are readily available and are commonly used in cytotoxicity tests. In this study, a low cell density was used and the cells were exposed for a long time, 5 days (McGaw et al., 2007). Hence, relatively low LC₅₀ values were obtained and differences in cytotoxicity between the extracts were maximised. Further work is needed to test the extracts, fractions and subfractions against other cell lines and, also to conduct acute and chronic toxicity assays with a view of determining the toxic constituents present in the plants.

It is a pity that one of the two plant species selected for in depth work *C. vendae* had a high cellular toxicity. In future studies toxicity of extracts should be investigated at an early stage.

10.6 Motility modulatory effects of *Bauhinia galpinii* and *Combretum vendae*

Considering the wide ethnopharmacological use of *B. galpinii* and *C. vendae* against GIT disorders and their excellent activity in some of the preliminary screening, the two plants were chosen for motility modulatory assays despite the toxicity potential of *C. vendae*. This was with the view that the toxic component(s) will be determined

and separated from the other active components. The data generated by the study indicate that *B. galpinii* has a dual-mechanism of action (prokinetic and relaxant) on gastro-intestinal motility while *C. vendae* extracts exhibited spasmolytic (relaxant) effects on isolated rat ileum through multiple mechanisms. These results were important as they indicate that *B. galpinii* extract can clinically be relevant as therapeutic agent in diarrhoea and constipation which are both diseases with aetiology based on motility disturbances to a large extent while the presence of multiple acting spasmolytic activities in the *C. vendae* extract might be contributing towards its effectiveness in diarrhoea and abdominal spasm therapy. Further work are needed for the identification of the specific ENS receptors through which these extract acts as well as the phytochemical compounds responsible for their activities.

10.7. Isolation and characterisation of antimicrobial compounds

Bioassay-guided protocols for antibacterial and antioxidant activity were adopted for the identification and isolation of 14 compounds (8 from *C. vendae* and 6 from *B. galpinii*) using open column chromatography with silica gel and Sephadex LH 20 as stationary phases. However, some of the compounds are mixtures of position isomers which are extremely difficult to separate. The compounds were characterised as ursol-12-en-28-oic acid (ursolic acid), a mixture of corosolic acid and maslinic acid, and a mixture of asiatic acid and arjunolic acid, two stilbenoid glycosides (combretastatin B5-O-2'- β -D-glucopyranoside and combretastatin B1-O-2'- β -D-glucopyranoside) and one flavone (apigenin) from the *Combretum vendae*.

One phytosterol (β -3 ethoxy sitosterol), one new flavone (5, 7, 4', 5' tetrahydroxy-2'-methoxyflavone (isoetin 2'-methyl ether) or 5, 7, 2', 5' tetrahydroxy-4'-methoxyflavone (isoetin 4'-methyl ether)), two known flavonols (3, 5, 7, 3', 4'-pentahydroxyflavone (Quercetin) and 3, 5, 7, 3', 4', 5'-hexahydroxyflavone (myricetin)) and their galactoside derivatives (quercetin-3-O- β -galactopyranoside and myricetin-3-O- β -galactopyranoside) were isolated from *Bauhinia galpinii*.

The results from this study indicate that medicinal plants used in ethnopharmacology are reservoirs of bioactive compounds. Some of the medicinal plants may serve as potential sources of novel active compounds or lead molecule for synthesis of more potent drugs. There is also a distinct possibility of developing plants extracts that could be used by poor rural people or sophisticated herbal medicines from some of the species investigated in this study. The information gained from this work provides a baseline study for other scientist to explore other medicinal plant species in depth with possible commercial application.

References

- Abbas, A. K., Lichtman, A. H., 2003. Cellular and Molecular Immunology. Elsevier, Amsterdam.
- Abdel-Hameed, E. S. S., 2009. Total phenolic contents and free radical scavenging activity of certain Egyptian *Ficus* species leaf samples. *Food Chemistry* 114 (4): 1271-1277.
- Abdur Rahman, M. A., Moon, S-S., 2007. Isoetin 5'-Methyl Ether, A Cytotoxic Flavone from *Trichosanthes kirilowii*. Bulletin of Korean Chemical Society 28 (8), 1261-1264.
- Abo K. A. Ashidi J. S., 1999. Antimicrobial screening of *Bridelia micrantha*, *Alchornea cordifolia* and *Boerhavia diffusa*. African Journal of Medicine & Medical Sciences 28(3-4), 167-169.
- Abroscia, B. D., Fiorentino, A., Monaco, P., Oriano, P. Pacifico, S., 2006. Annurcoic acid: A new antioxidant ursane triterpene from fruits of cv. Annurca apple. *Food Chemistry* 98, 285–290.
- Acharya, S., Datta, A., Bag, P. K., 2009. Evaluation of the antimicrobial activity of some medicinal plants against enteric bacteria with particular reference to Multi-Drug Resistant *Vibrio cholerae*. *Tropical Journal of Pharmaceutical Research* 8, 231-237.
- Adedapo, A. A., Sofidiya, M. O., Masika, P. J., Afolayan, A. J., 2008. Anti-Inflammatory and Analgesic Activities of the Aqueous Extract of *Acacia karroo* Stem Bark in Experimental Animals. *Basic & Clinical Pharmacology & Toxicology* 103, 397–400.
- Adedapo, A. A., Sofidiya, M. O., Masika, P. J., Afolayan, A. J., 2008. Safety evaluations of the aqueous extract of *Acacia karroo* stem bark in rats and mice. *Scientific Commons*
- Aderibigbe, A.O., Emudianughe, T.S., Lowal, B.A., 2001. Evaluation of antidiabetic action of *Mangifera indica* in mice. *Phytotherapy Research* 15, 456–458.
- Aderogba, M. A., McGaw, L. J., Ogundaini, A. O., Eloff, J. N., 2007. Antioxidant activity and cytotoxicity study of the flavonol glycosides from *Bauhinia galpinii*. *Natural Product Research* 21, 591-599.
- Adewusi, E. A., Afolayan, A. J., 2009a. Antibacterial, antifungal and antioxidant activity of the root and leaves of *Pelargonium reniforme* Curtis (Geraniaceae). *African Journal of Biotechnology*, 8(22), 6425-6433.
- Adewusi, E. A., Afolayan, A. J., 2009b. Safety evaluation of the extract from the roots of *Pelargonium reniforme* Curtis in male Wistar rats. *African Journal of Pharmacy and Pharmacology* 3(8), 368-373.
- Adzu, B., Abubakar, M. S., Izebe, K. S., Akumka, D. D., Gamaniel, K. S., 2005. Effect of *Annona senegalensis* rootbark extracts on *Naja nigricollis nigricollis* venom in rats. *Journal of Ethnopharmacology* 96, 507–513.
- Afolayan, A. J., Yakubu, M. T., 2009. Effect of *Bulbine natalensis* Baker stem extract on the functional indices and histology of the liver and kidney of male Wistar rats. *Journal of Medicinal Food*. 12(4): (in press), DOI:10.1089/jmf.2008.0221.
- Agnihotri, V. K., Elsohly, H. N., Smillie, T. J., Khan, I. A., Walker, L. A., 2009. Constituents of *Leonitis leonurus* flowering tops. *Phytochemistry letters* 2:103-105.
- Aguirre-Crespo, F., Vergara-Galicia, J., Villalobos-Molina, R., L'opez-Guerrero, J. J., Navarrete-V'azquez, G., Estrada, S., 2006. Ursolic acid mediates the vasorelaxant activity of *Lepechinia caulescens* via NO release in isolated rat thoracic aorta. *Life Science* 79, 1062–1068.
- Ahmed, A. S., Igwe, C. C., Eloff, J. N., 2009. Preliminary studies of the antibacterial activities of *Combretum vendae* leave extract. *African Journal of Traditional Medicine* 366-367.
- Akah, P. A., 1996. Antidiarrhoeal activity of *Kigalia africana* in Experimental animal. *Journal of Herbs and Spices Medicinal plants* 4, 31-38.

- Akah, P. A., Nwafor, S. V., 1999. Studies on anti-ulcer properties of *Cissampelos mucronata* leaf extract. *Indian Journal of Experimental Biology* 37 (9), 936–938.
- Akula, U. S., Odhav, B., 2008. In Vitro 5-lipoxygenase inhibition of polyphenolic antioxidant from undomesticated plants of South Africa. *Journal of Medical plant Research* 2(9), 207-212.
- Alam, M. S., Chopra, N., Ali, M., Niwa, M., Sakae, T., 1994. Ursane and sterol derivatives from *Pluchea lanceolata*. *Phytochemistry*, 37, 521–524.
- Albuquerque, U. P., Hanazaki, N., 2006. As pesquisas etnorigidas na descoberta de novos farmacos de interesse medico e farmaceutico: fragilidades e perspectivas. *Brazilian Journal of Pharmacognosy* 16, 678–689.
- Al-Khatib, M. S., Khyami-Horani, H., Badran, E., Shehabi, A. A., 2007. Incidence and characterization of diarrheal enterotoxins of fecal *Bacillus cereus* isolates associated with diarrhea. *Diagnostic Microbiology and Infectious Disease* 59, 383–387.
- Almeida, C. E., Karnikovski, M. G., Foletto, R., Baldisserotto, B., 1995. Analysis of antidiarrheic effect of plants used in popular medicine. *Rev. Saude Publica* 29,428-433.
- Amabeoku, G. J., 2009. Antidiarrhoeal activity of *Geranium incanum* Burm. f. (Geraniaceae) leaf aqueous extract in mice. *Journal of Ethnopharmacology* 123,190–193.
- Amabeoku, G. J., Bamuamba, K., 2010. Evaluation of the effects of *Olea europaea L.subsp. africana* (Mill.) P. S. Green (Oleaceae) leaf methanol extract against Castor oil-induced diarrhoea in mice. *Journal of Pharmacy and Pharmacology* 62(3), 368-373.
- Amara, A. A., El-Masry, M. H., Bogdady, H. H., 2008. Plant crude extracts could be the solution: Extracts showing in vivo antitumorigenic activity. *Pakistan Journal of Pharmaceutical Sciences*, 21(2), 159-171.
- Amusan, O. O. G., Sukati, N. A., Dlamini, P. S., and Sibandz, F. G., 2007. Some Swazi phytomedicines and their constituents. *African Journal of Biotechnology* 6(3), 267-272.
- Angeh, J. E., Huang, X., Sattler, I., Swan, G. E., Dahse, H., Hartl, A., Eloff, J. N., 2007. Antimicrobial and anti-inflammatory activity of four known and one new triterpenoid from *Combretum imberbe* (Combretaceae). *Journal of Ethnopharmacology* 110, 56–60.
- Angeh, J. E., Huang, X., Swan, G. E., Mollman, U., Sattler, I., Eloff, J. N., 2007. Novel antibacterial triterpenoid from *Combretum padoides* (Combretaceae). *ARKIVOC* (ix) 113-120.
- Annese, V., Janssens, J., Vantrappen, G. *et al.*, 1992. Erythromycin accelerates gastric emptying by inducing antral contractions and improved gastroduodenal coordination. *Gastroenterology* 102, 823–828.
- Appidi, J. R., Grierson, D. S., Afolayan, A. J., 2008. Ethnobotanical study of plants used for the treatment of diarrhoea in the Eastern Cape, South Africa. *Pakistan Journal of Biological Sciences* 11, 1961–1963.
- Arnold, H-J., Gulumian, M., 1984. Pharmacopoeia of traditional medicine in Venda. *Journal of Ethnopharmacology* 12, 35–74.
- Arts, I. C. W., Hollman, P. C. H., 2005. Polyphenols and disease risk in epidemiologic studies. *American Journal of Clinical Nutrition*, 81(1), 317S–325S.
- Aruoma, O. I., Colognato, R., Fontana, I., Gartlon, J., Migliore, L., Koike, K., Coecke, S., Lamy, E., Mersch-Sundermann, V., Laurenza, I., Benzi, L., Yoshino, F., Kobayashi, K., Lee, M. C., 2006. Molecular effects of fermented papaya preparation on oxidative damage, MAP kinase activation and modulation of the benzo[a]pyrene mediated genotoxicity. *Biofactors* 26, 147–159.

- Asase, A. A., Oteng-Yeboah, G. T., Odamtten Simmonds, M. S., 2005. Ethnobotanical study of some Ghanaian anti-malarial plants. *Journal of Ethnopharmacology* 99, 1221-1229
- Asres, K., Bucar, F., Knauder, E., Yardley, V., Hendrick, H., Croft, S. L., 2001. *In vitro* antiprotozoal activity of extract and compound from the stem bark of *Combretum molle*. *Phytotherapy Research* 15(7), 613-617.
- Athanasiadou, S., Houdijk, J., Kyriazakis, I., 2008. Exploiting synergisms and interactions in the nutritional approaches to parasite control in sheep production systems. *Small Ruminant Research* 76, 2-11.
- Atta, A. H., Mouneir, S. M., 2005. Evaluation of some medicinal plant extracts for anti-diarrhoeal activity. *Phytotherapy Research* 19(6), 481-485.
- Auroma, O. I., Halliwell, B., Loughton, M. J., Quinlan, G. J., Gutteridge, J. M. C., 1989. The mechanism of initiation of lipid peroxidation. Evidence against a requirement for an iron(II)-iron(III) complex. *Biochemistry Journal* 258:617- 620.
- Aworet-Samseny, R. R., Souza, A., Kpahé, F., Konaté, K., Datté, J. Y., 2011. *Dichrostachys cinerea* (L) Wight et Arn. (Mimosaceae) hydro-alcoholic extract action on the contractility of tracheal smooth muscle isolated from guinea-pig. *BMC Complementary and Alternative Medicine* 11, 23.
- Ayinde, B. A., Owolabi, O. J., 2009. Effect of the aqueous extract of *Ficus capensis* Thunb (Moraceae) leaf on gastrointestinal motility. *Journal of Pharmacognosy and Phytotherapy* 1(3), 031-035.
- Babajide, O. J., Mabusela, W. T., Green, I. R., Ameer, F., Weitz, F., Iwouha, E. I., 2010. Phytochemical screening and biological activity studies of five South African indigenous medicinal plants. *Journal of Medicinal Plant Research* 2(18), 1924-1932.
- Baker, J. T., Borris, R. P., Carte, B., Cardell, G. A., Soejarto, D. D., Gragg, G. M., Gupta, M. P., Iwu, M. M., Madilid, D. R., and Tyler, V. E., 1995: . Natural product drug discovery and development: New perspectives on international collaboration. *Journal of Natural products* 58(9), 1325-1357.
- Baldi, F., Bianco, M. A., Nardone, G., Pilotto, A., Zamparo, E., 2009. Focus on acute diarrhoeal disease. *World Journal of Gastroenterology* 15(27), 3341-3348.
- Bamuamba, K., Gammon, D. W., Meyers, P., Dijoux-Franca, M.-G., Scott, G., 2008. Anti-mycobacterial activity of five plant species used as traditional medicines in the Western Cape Province (South Africa). *Journal of Ethnopharmacology* 117, 385-390.
- Bandaranayake, W. M., 2006. Quality control, Screening Toxicity and Regulation of Herbal Drugs. In: *Modern phytomedicine. Turning medicinal plants into drugs* (Edited by Ahmed F Aqil and Owais M) WILEY-VCH Verlag GmbH and Co KGaA, Weinheim Pp 25-53.
- Barbosa, E., Calzada, F., Campos, R., 2007. In vivo anti-giardial activity of three flavonoids isolated of some medicinal plants used in Mexican traditional medicine for the treatment of diarrhea. *Journal of Ethnopharmacology* 109, 552-554.
- Barbour, E. K., Sharif, M. A., Sagherian, V. K., Habre, A. N., Talhouk, R. S., Talhouk, S. N., 2004. Screening of selected indigenous plants of Lebanon for antimicrobial activity. *Journal of Ethnopharmacology* 93, 1-7.
- Barton, M. D., 2000. Antibiotic use in animal feed and its impact on human health. *Nutritional Research Review* 13, 279-299.
- Batawila, K., Kokou, K., Koumaglo, K., Gbeassor, M., de Foucault, B., Bouchet, Ph. Akpagana, K., 2005. Antifungal activities of five Combretaceae used in Togolese traditional medicine. *Fitoterapia* 76(2), 264-268.
- Beaugerie L, Flahault A, Barbut F et al. Antibiotic-associated diarrhoea and *Clostridium difficile* in the community. *Alimentary Pharmacology and Therapeutics* 2003; 17: 905-912
- Beaugerie, L., Pettit, J-C., 2004. Antibiotic-associated diarrhoea. *Best Practice & Research Clinical Gastroenterology* 18, 337-352.

- Beavis, K. G., Weymouth, L. A., 1996. Cellular and Molecular pathology of Infectious diseases. In: Cellular and Molecular Pathogenesis. Editor, Alphonse E. Sirica: Lippincott-Raven Publishers, 277 East Washington Square, Philadelphia, Pennsylvania 19106-3780 Pp 219-244.
- Berker, K. I., Guclu, K., Tor, I., Apak, R., 2007. Comparative evaluation of Fe(III) reducing power-based antioxidant capacity assays in the presence of phenanthroline, batho-phenanthroline, tripyridyltriazine (FRAP) and ferricyanide reagent. *Talanta* 72, 1157-1165.
- Berlin, E.A., Berlin, B., 2005. Some field methods in medical ethnobiology. *Field Methods* 17, 235–268.
- Bessong, P. O., Obi, C. L., Andreola, M. L., Rojas, L. B., Pouysegu, L., Igumbor, E., Marion Meyer, J. J., Quideau, S., Litvak, S., 2005. Evaluation of selected South African medicinal plants for inhibitory properties against human immunodeficiency virus type 1 reverse transcriptase and integrase. *Journal of Ethnopharmacology* 99, 83–91.
- Bhan, M. K., 2000. Current and Future: Management of childhood diarrhoea. *International Journal of Antimicrobial Agents* 14, 71-73.
- Bhargav, B., Rupal, A. V., Reddy, A. S., Narasimhacharya, A. V. R. L., 2009. Antihyperglycaemic and hypolipidemic effect of *Adansonia digitata* L. On Alloxan induced diabetic rats. *Journal of Cells and Tissue Research* 9(2), 1879-1882.
- Bhat, R. B., Jacobs, T. V., 1995. Traditional herbal medicine in Transkei. *Journal of Ethnopharmacology*, 48, 7-12.
- Biennu, E., Amabeoku, G. J., Eagles, P. K., Scott, G., Springfield, E. P., 2002. Anticonvulsant activity of aqueous extract of *Leonotis leonurus*. *Phytomedicine* 9, 217–223.
- Bisi-Johnson, M. A., Obi, C. L., Kambizi, L., Nkomo, M., 2010. A survey of indigenous herbal diarrhoeal remedies of O.R. Tambo district, Eastern Cape Province, South Africa. *African Journal of Biotechnology* 9(8), 1245-1254.
- Bizimenyera, E. S., Githiori, J. B., Eloff, J. N., Swan, G. E., 2006. In vitro activity of *Peltophorum africanum* (Fabaceae) extracts on the egg hatching and larval development of the parasitic nematode *Trichostrongylus colubriformis*. *Veterinary Parasitology* 142, 336-343.
- Blikslager, A. T., 2010. Mucosal epithelial barrier repair to maintain pig health. *Livestock Science* 133, 194–199.
- Bohn, L. M., Raehal, K. M., 2006. Opioid receptors signalling: relevance for gastrointestinal therapy. *Current Opinion in Pharmacology* 6, 559-563.
- Borghetti, P., Saleri, R., Mocchegiani, E., Corradi, A., Martelli, P., 2009. Infection, Immunity and the Neuroendocrine Response. *Veterinary Immunology and Immunopathology* 130, 141–162.
- Borrelli, F., Capasso, F., Capasso, R., Ascione, V., Aviello, G., Longo, R., Izzo, A. A., 2006. Effects of *Boswellia serrata* on intestinal motility in rodents: inhibition of diarrhoea without constipation. *British Journal of Pharmacology* 148, 553-560
- Botes, L., Van der Westhuizen, F. H., Loot du, T., 2008. Phytochemical and antioxidant capacity of two *Aloe greatheadii* var. *davyana* extract. *Molecules* 13(9), 2169-2180.
- Brand-Williams, W., Cuvelier, M.E., Berset, C., 1995. Use of a free radical method to evaluate antioxidant activity. *Lebensmittel Wissenschaftund Technologie* 28, 25-30.
- Brendler, T., Van Wyk, B. E., 2008. A historical, scientific and commercial perspective on the medicinal use of *Pelargonium sidioides* (Geraniaceae). Review. *Journal of Ethnopharmacology* 119, 420–433.
- Breytenbach, J. C., Malan, S. F., 1989. Pharmacological properties of *Combretum zeyheri*. *South African Journal of Sciences* 85(6), 372-374.

- Brijesh, S., Daswani, P. G., Tetali, P., Rojatkhar, S. R., Antia, N. H., Birdi, T. J., 2006. Studies on *Pongania pinnata* (L) Pierre leaves: Understanding the mechanism(s) of action in infectious diarrhea. *Journal of Zhejiang University SCIENCE B* 7(8), 665-674.
- Brown, J.H., Taylor, P., 1996. Muscarinic receptor agonists and antagonists. In: Gilman, A.G., Hardman, J.G., Limbird, L.E., Molinoff, P.B., Ruddon, R.W. (Eds.), *Goodman & Gillman's: The Pharmacological Basis of Therapeutics*. McGraw-Hill, New York, pp. 141–159.
- Buettner, G.R., 1993. The pecking order of free radicals and antioxidants lipid peroxidation alpha-tocopherol and ascorbate, *Archive Biochemistry Biophysics* 300, 535–543.
- Buret, A. G. 2007. Mechanisms of epithelial dysfunction in giardiasis. *GUT* 56(3), 328–335.
- Butler, K. M., 2008. Enterococcal Infection in Children. *Seminars in Pediatric Infectious diseases* 17, 128-139
- Buwa, L. V., Van Staden, J., 2006. Antibacterial and Antifungal activity of traditional medicinal plants used against venereal diseases in South Africa. *Journal of Ethnopharmacology* 103(1), 139-142.
- Cai, Y. Z., Sun, M., and Corke, H., 2003. Antioxidant activity of betalains from plants of the Amaranthaceae. *Journal of Agricultural and Food chemistry* 51(8), 2288-2294.
- Calis, I., Yürüker, A., Tas, demir, D., Wright, A.D., Sticher, O., Luo, Y.D., Pezzuto, J. M., 1997. Cycloartane triterpene glycosides from the roots of *Astragalus melanophrurius*. *Planta Medica* 63, 183–186.
- Calzada, F., Yépez Mulia, L., Aguilar, A., 2006. In vitro susceptibility of *Entamoeba histolytica* and *Giardia lamblia* to plants used in Mexican traditional medicine for the treatment of gastrointestinal disorders. *Journal of Ethnopharmacology* 108, 367–370.
- Caparroz-Assef, S. M., Grespan, R., Batista, R. C. F., Bersani-Amado, F. A., Baroni, S., Dantas, J. A., Cuman, R. K. N., Bersani-Amado, C.A., 2005. Toxicity studies of *Cordia salicifolia* extract. *Acta Scientiarum Health Science* 27, 41–44.
- Casburn-Jones, A. C., and Farthing, M. J. D., 2004. Management of infectious diarrhoea. *GUT* 53, 296-305.
- Castillo, N. A., de LeBlanc, A. M., Galdeano, C. M., Perdigón, G., 2011. Probiotics: An alternative strategy for combating salmonellosis Immune mechanisms involved. *Food Research International* doi:10.1016/j.foodres.2011.04.031
- Catalá, A., 2006. An overview of lipid peroxidation with emphasis in outer segments of photoreceptors and the chemiluminescence assay. *Int. J. Biochem. Cell Biol.* 38, 1482–1495, Review.
- Catala, A., 2009. Lipid peroxidation of membrane phospholipids generates hydroxy-alkenals and oxidized phospholipids active in physiological and/or pathological conditions. *Chemistry and Physics of Lipid* 157: 1-11.
- Catnach, S. M., Fairclough, P. D., 1992. Erythromycin and the gut. *Gut* 33, 397–401.
- Cellek, S., John, A. K., Thangiah, R., Dass, N. B., Bassil, A. K., Jarvie, E.M., Lalude, O., Vivekanandan, S., Sanger, G. J., 2006. 5-HT₄ receptor agonists enhance both cholinergic and nitrgergic activities in human isolated colon circular muscle. *Neurogastroenterology and Motility* 18, 853–861.
- Ceylan, E., Fung, D. Y. C., 2004. Antimicrobial activity of spices. *Journal of RapidMethods and Automation in Microbiology*, 12(1), 1–55.
- Chacin-Bonilla, L., 2010. Epidemiology of *Cyclospora cayetanensis*: A review focusing in endemic areas. *Acta Tropica* 115, 181–193.
- Chang, S-L., Chiang, Y-M., Chang, C. L-T., Yeh, H-H., Shyur, L-F., Kou, Y-H., Wu, T-K., Yang, W-C., 2007. Flavonoid, Centaurein and Centaurenidin, from *Bidens pilosa* stimulate IFN- γ expression. *Journal of Ethnopharmacology*, 112, 232-236.

- Charlier, C., Michaux, C., 2003. Dual inhibition of cyclooxygenase-2 (COX-2) and 5-lipoxygenase (5-LOX) as a new strategy to provide safer non-steroidal anti-inflammatory drugs. *European Journal of Medicinal Chemistry* 38, 645-659.
- Chaturongakul, S., Raengpradub, S., Wiedman, M., Boor, K. J., 2008. Modulation of stress and virulence in *Listeria monocytogenes*. *Trend in Microbiology* 16, 388-396.
- Chen, J.-C., Chang, Y.-S., Wu, S.-L., Chao, B.-C., Chang, C.-S., Li, C.-C., Ho, T.-Y., Hsiang, C.-Y., 2007. Inhibition of *Escherichia coli* heat labile enterotoxins induced diarrhoea by *Chaenomeles speciosa*. *Journal of Ethnopharmacology* 113, 233-239.
- Chen, J.-C., Ho, T.-Y., Chang, Y.-S., Wu, S.-L., Li, C.-C., Hsiang, C.-Y., 2009. Identification of *Escherichia coli* enterotoxin inhibitors from traditional medicinal herbs by *in silico*, *in vitro*, and *in vivo* analyses. *Journal of Ethnopharmacology* 121, 372–378.
- Chen, K., Suh, J., Carr, A. C., Morrow, J. D., Zeind, J., Frei, B., 2000. Vitamin C suppresses oxidative lipid damage *in vivo*, even in the presence of iron overload. *American Journal of Physiology, Endocrinology and Metabolism* 279, E1406–E1412.
- Chen, X.-W., Serag, E. S., Sneed, K. B., Zhou, S.-F., 2011. Herbal bioactivation, molecular targets and the toxicity relevance. *Chemico-Biological Interactions* 192, 161–176.
- Chetty, N., Irving, H. R., Coupar, I. M., 2006. Activation of 5-HT₃ receptors in the rat and mouse intestinal tract: a comparative study. *British Journal of Pharmacology* 148, 1012–1021.
- Chi, J., VanLeeuwen, J. A., Weersink, A., Keefe, G. P., 2002. Direct production losses and treatment costs from bovine viral diarrhoea virus, bovine leukosis virus, *Mycobacterium avium* subspecies *paratuberculosis*, and *neosporea caninum*. *Preventive veterinary medicine* 55, 137-153.
- Chiang, Y.-M., Chang, C. L.-T., Yang, S.-L., Shyur, L.-F., 2007. Cytopyloine, a novel polyacetylenic glycoside from *Biden pilosa*, function as a T-helper cell modulator. *Journal of Ethnopharmacology* 110(3), 532-538.
- Chinsamy, M., Finnie, J. F., Van Staden, J., 2010. The ethnobotany of South African medicinal orchids. *South African Journal of Botany*
- Choma, I. M., Grzelak, E. M., 2011. Bioautography detection in thin-layer chromatography. *Journal of Chromatography A* 1218, 2684–2691.
- Christophersen, O. A., Haug, A., 2005. Possible roles of oxidative stress, local circulatory failure and nutrition factors in the pathogenesis of hypervirulent influenza: Implications for therapy and global emergency preparedness. *Microbial Ecology in Health and Disease* 17, 189-199.
- Cimanga, R. K., Kambu, K., Tona, L., Hermans, N., Apers, S., Totte, J., Pieters, L., Vlietinck, A. J., 2006. Cytotoxicity and *in vitro* susceptibility of *Entamoeba histolytica* to *Morinda Morindoides* leaf extract and its isolated constituents. *Journal of Ethnopharmacology* 107, 83-90.
- Cimanga, R. K., Mukenyi, P. N. K., Kambu, O. K., Tona, G. L., Apers, S., Totté, J., Pieters, L., Vlietinck, A. J., 2010. The spasmolytic activity of extracts and some isolated compounds from the leaves of *Morinda morindoides* (Baker) Milne-Redh. (Rubiaceae). *Journal of Ethnopharmacology* 127, 215–220
- Cimpoi, C., 2006. Analysis of some natural antioxidants by Thin-Layer Chromatography and High performance Thin-layer Chromatography. *Journal of liquid chromatography and related Technologies*, 29: , 1125-1142.
- Clarke, S. C., 2001. Diarrhoeagenic *Escherichia coli*-an emerging problem? *Diagnostic Microbiology and infectious disease* 41, 93-98.
- Coates-Palgrave, M., 2002. Keith Coates-Palgrave Trees of Southern Africa, 3rd edition, imp. Struik Publishers. Cape Town.

- Conforti, F., Sosa, S., Marrelli, M., Menichini, F., Statti, G. A., Uzunov, B., Tubaro, A., Menichini, F., Loggia, R. D., 2008. *In vivo* anti-inflammatory and *in vitro* antioxidant activities of Mediterranean dietary plants. *Journal of Ethnopharmacology* 116, 144-151.
- Cooposamy, R. M., 2011. Traditional information and antibacterial activity of Four *Bulbine* species (Wolf). *African Journal of Biotechnology* 10(2), 220-224.
- Cos, P., Vanden Berghe, D.; De Bruyne, T.; Vlietinck, A.J. *Curr. Org. Chem.*, **2003**, 7, 1163.
- Cotton, J. A., Beatty, J. F., Buret, A. G., 2011. Host parasite interactions and pathophysiology in *Giardia* infections. *International Journal for Parasitology* xxx , xxx-xxx
- Coutino, R. R., Hernandez, C. P., Giles, R. H., 2001. Lectins in fruits having gastrointestinal activity: their participation in the hemagglutinating property of *Escherichia coli* O157:H7. *Archives of Medical Research* 32, 251-257.
- Cowan, M. M., 1999. Plant products as antimicrobial agents. *Clinical Microbiology Review* 12, 564-582.
- Cuello, S., Alberto, M. R., Zampini, I. C., Ordonez, R. M., Isla, M. I., 2011. Comparative study of antioxidant and anti-inflammatory activities and genotoxicity of alcoholic and aqueous extracts of four *Fabiana* species that grow in mountainous area of Argentina. *Journal of Ethnopharmacology* xxx , xxx- xxx
- Cui, J., Wang, Z. Q., Xu, B. L., 2011. Review: The epidemiology of human trichinellosis in China during 2004-2009. *Acta tropica* 118, 1-5.
- Cuzzocrea, S., Mazzon, E., Serraino, I., Dugo, L., Centorrino, T., Ciccolo, A., Sautebin, L., Caputi, A.P., 2001. Celecoxib, a selective cyclo-oxygenase-2 inhibitor reduces the severity of experimental colitis induced by dinitrobenzene sulfonic acid in rats. *European Journal of Pharmacology*. 431, 91-102.
- da Silva, G., Tamia, M., Rocha, J., Serrona, R., Gomes, E. T., Sepodea, B., Silva, O., 2010. *In vivo* Anti-inflammatory effect and toxicological screening of *Maytenus heterophylla* and *Maytenus senegalensis* extracts. *Human and Experimental Toxicology* 000(00) 1-8 DOI: 10: 1177/0960327110379242
- Dambisya, Y. M., Tindimwebwa, G., 2003. Traditional remedies in children around Eastern Cape, South Africa. *East African Medical Journal* 80, 401-405.
- Danna, P. L., Urban, C., Bellin, E *et al.*, 1991. Role of candida in pathogenesis of antibiotic-associated diarrhoea in elderly patients. *Lancet* 337, 511-514.
- de Boer, H. J., Kool, A., Broberg, A., Mziray, W. R., Hedberg, I., Levenfors, J. J., 2005. Antifungal and antibacterial activity of some herbal remedies from Tanzania. *Journal of Ethnopharmacology* 96(3), 461-469.
- de Oliveira, C. E. V., Stamford, T. L. M., Neto, N. J. G., de Souza, E. L., 2010. Inhibition of *Staphylococcus aureus* in broth and meat broth using synergies of phenolics and organic acids. *International Journal of Food Microbiology* 137, 312-316.
- De Villiers, B. J., Van Vuuren, S. F., Van Zyl, Z. L., Van Wyk, B-E., 2010. Antimicrobial and antimalarial activity of *Cussonia* species (Araliaceae). *Journal of Ethnopharmacology* 129: 189-196.
- de Wet, H., Nkwanyana, M. N., van Vuuren, S. F., 2010. Medicinal plants used for the treatment of diarrhoea in northern Maputaland, KwaZulu-Natal Province, South Africa. *Journal of Ethnopharmacology* 130. 284-289.
- Deeni, Y. Y., Hussain, H. S. N., 1994. Screening of *Vernonia kotschyana* for antimicrobial activity and alkaloids. *International Journal of Pharmacognosy* 32, 388-395.
- Dekker, T. G., Fourie, T. G., Matthee, E., Snycker, F. O., Ammann, W., 1987. Studies of South African medicinal plants. Part 4: Jaherin, a new daphnane diterpene with antimicrobial properties from *Jatropha zeyheri*. *South African Journal of Chemistry*, 40, 74-76

- Delaporte, R. H., Sanchez, G. M., Cuellar, A. C., Giuliani, A., Palazzo de Mello, J. C., 2002. Anti-inflammatory activity and lipid peroxidation inhibition of iridoid lamiide isolated from *Bouchea fluminensis* (vell.) Mold. (Verbenaceae). *Journal of Ethnopharmacology* 82: 127-130.
- Delves, P., Martin, S., Burton, D., Roitt, I., 2006. *Roitt's Essential Immunology*, 11th ed. Wiley-Blackwell, Hoboken, NJ, USA.
- Desire, O., Riviere, C., Razafindrazaka, R., Goosens, L., Moreau, S., Guillon, J., Uverg-Ratsimamanga, S., Andriamadio, P., Moore, N., Randriantsoa, A., Raharisolalao, A., 2010. Antispasmodic and antioxidant activities of fraction sand bioactive constituent davidigenin isolated from *Mascarenhasia arborescens*. *Journal of Ethnopharmacology* 130, 320-328.
- Deutschlander, M. S., Van de Venter, M., Louw, J., Lall, N., 2009. Hypoglycaemic activity of four plant extracts traditionally used in South Africa for diabetes. *Journal of Ethnopharmacology*, 124(3), 619-624.
- Dickman, K. G., Hempson, S. J., Anderson, J., Lippe, S., Zhao, L., Burakoff, R., et al. 2000. Rotavirus alters paracellular permeability and energy metabolism in Caco-2 cells. *American Journal of Physiology, Gastrointestinal and Liver Physiology*, 279(4), G757–G766.
- Dobbins, W. O., Herrero, B. A., Mansbach, C. M., 1968. Morphologic alterations associated with neomycin induced malabsorption. *The American Journal of Medical Sciences* 255, 63–77.
- Dold, A. P., Cocks, M. L., 2001. Traditional veterinary medicine in the Alice district of the Eastern Cape Province, South Africa. *South African Journal of Science* 97, 375-379.
- Du Toit, K., Elgorashi, E. E., Sarel, F. M., Drewes, S. E., Van Staden, J., Crouch, N. R., Mulholland, D. A., 2005. Anti-inflammatory activity and QSAR studies of compounds isolated from Hyacinthaceae and *Tachyadenus longiflorus* Griseh (Gentianaceae). *Bioorganic and medicinal chemistry* 13, 2561-2568.
- Dudeja, P. K., and Ramaswamy, K., 2006. *Physiology of Gastrointestinal Tract, Fourth edition*, Edited by Leonard R. Johnson. Academic Press, Chicago USA.
- Dupont, H. L., Marshall, G. D., 1995. HIV-associated diarrhoea and wasting. *Lancet* 346, 352-356.
- Dwyer, D. J., Kohanski, M. A., Collins, J. J. 2009. Role of reactive oxygen species in antibiotic action and resistance. *Current Opinion in Microbiology* 12, 482–489.
- Ebert, E. C., 2005. Gastrointestinal Complications of Diabetes Mellitus. *Dis Mon* 51, 620-663.
- Ehling-Schulz, M., Fricker, M., Scherer, S., 2004. *Bacillus cereus*, the causative agent of an emetic type of food-borne illness. *Molecular Nutrition and Food Research* 48, 479–487.
- Eldeen, I. M. S., Eldeen, I. M., Van Heerden, F. R., Van Staden, J., 2008. Isolation and Biological activities of Termilignan B and arjunic acid from *Terminalia sericea*. *Planta Medica* 74(4), 411-413.
- Eldeen, I. M. S., Elgorashi, E. E., Mulholland, B. A., Van Staden, J., 2006. Anolignan B: A bioactive compound from the roots of *Terminalia sericea*. *Journal of Ethnopharmacology* 103(1), 135-138.
- Eldeen, I. M. S., Elgorashi, E. E., Van Staden, J., 2005. Antibacterial, antiinflammatory, anti-cholinesterase and mutagenic effects of extrats obtained from some trees used in South African traditional medicine. *Journal of Ethnopharmacology* 102, 457-464
- Eldeen, I. M. S., Van Heerden, F. R., Van Staden, J., 2007. Biological activities of cycloart-2, 3-ene-3, 25-diol isolated from the leaves of *Trichilia drageana* South African Journal of botany 73(3), 366-371.
- Eldeen, I. M. S., Van Heerden, F. R., Van Staden, J., 2010. In vitro biological activities of niloticane, a new bioactive cassane diterpene from the bark of *Acacia nilotica* subsp. *Kraussiana*. *Journal of Ethnopharmacology* 128, 555–560.

- Elgayyar, M., Droughon, F. A., Golden, D. A., Mount, J. R., 2001. Antimicrobial activity of essential oils from plants against selected pathogenic and saprophytic microorganisms. *Journal of Food Protection* 64, 1019-1024.
- Elgorashi, E. E., Taylor, J. L. S., Maes, A., van Van Staden, J., De Kimpe, N., Verschaeve, L., 2003. Screening of medicinal plants used in South African traditional medicine for genotoxic effects. *Toxicology Letters* 143, 195–207.
- Ellman, G. L., Coutney, D., Andies, V., Featherstone, R. M., 1961. A new and rapid colourimetric determination of acetylcholinesterase activity. *Biochemical Pharmacology* 7, 88–95.
- Eloff, J. N. 2001. Antibacterial activity of Marula (*Sclerocarya birrea* (A. Rich.) Hochst. subsp. *caffra* (Sond.) Kokwaro) (Anacardiaceae) bark and leaves. *Journal of Ethnopharmacology* 76, 305-308.
- Eloff, J. N., 1998. A sensitive and quick micro-plate method to determine the minimal inhibitory concentration of plant extracts for bacteria. *Planta Medica*, 64: 711-713.
- Eloff, J. N., Famakin, J. O., Katerere, D. R., 2005. Isolation of antibacterial stilbenes from *Combretum woodii* (Combretaceae) leaves. *African Journal of Biotechnology* 4, 1166–1171.
- Eloff, J. N., Katerere, D. R., McGaw, L. J., 2008. The Biological Activity and Chemistry of the Southern African Combretaceae. *Journal of Ethnopharmacology* 119, 686–699.
- Elsinghorst, E. A., 2002. Enterotoxigenic *Escherichia coli*, In *Escherichia coli: Virulence Mechanisms of a Versatile Pathogen*. Elsevier Science (USA) Pp 155-187.
- Epple, H. J., Schneider, T., Troeger, H., Kunkel, D., Allers, K., Moos, V., *et al.*, 2009. Impairment of the intestinal barrier is evident in untreated but absent in suppressively treated HIV-infected patients. *Gut* 58, 220–227.
- Erde'lyi, K., Kiss, A., Bakondi, E., Bai, P., Szabó, C., Gergely, P., Erdó di, F., Virá g, L., 2005. Gallotannin inhibits the expression of chemokines and inflammatory cytokines in A549 Cells. *Molecular Pharmacology* 68, 895–904.
- Ernst, E., 2004. Risks of herbal medicinal products. *Pharmacoepidemiology and Drug Safety* 13, 767–771.
- Escandell, J. M., Recio, M. C., Giner, R. M., Mánez, S., Cerdá-Nicolás, M., Merfort, I., Ríos, J. L., 2010. Inhibition of delayed-type hypersensitivity by cucurbitacin R through the curving of lymphocyte proliferation and cytokine expression by means of NF-AT translocation to the nucleus. *Journal of Pharmacology and Experimental Therapeutics* 332, 352–363.
- Escandell, J. M., Recio, M. C., Mánez, S., Giner, R. M., Cerdá-Nicolás, M., Gil-Benso, R., Ríos, J. L., 2007. Dihydrocucurbitacin B inhibits delayed-type hypersensitivity reaction by suppressing lymphocyte proliferation. *Journal of Pharmacology and Experimental Therapeutics* 322, 1261–1268.
- Essop, A. B., van Zyl, R. L., van Vureen, S. F., Mulholland, D and Viljoen, A. M. 2008. The *in vitro* pharmacological activities of South African *Hermania* species. *Journal of Ethnopharmacology* 119, 615-619.
- Esterbauer, H., Schaur, R.J., Zollner, H., 1991. Chemistry and biochemistry of 4- hydroxynonenal, malonaldehyde and related aldehydes. *Free Radical Biology and Medicine* 11, 81–128.
- Estrada-soto, S., Rodriguez-Aviles, A., Castaneda-Avila, C., Castillo-Espana, P., Navarrete-Vazquez, G., Hernandez, L., and Aguirre-Crespo, F., 2007. Spasmolytic action of *Lepechinia caulescens* is through calcium blockage and NO release. *Journal of Ethnopharmacology* 114:364-370.
- Fabry, W., Okemo, P., Ansorg, R., 1996. Activity of East African medicinal plants against *Helicobacter pylori*. *Chemotherapy* 42, 315–317.
- Fallman, M., Gustavsson, A., 2005. Cellular Mechanisms of Bacterial Internalization Counteracted by *Yersinia*. *International Review of Cytology* 246, 135-188.
- Farthing, M. J. G., 2002. Novel target for the control of secretory diarrhoea. *Gut*, 50(Suppl III):iii15–iii18

- Farthing, M. J. G., Casburn-Jones, A., Banks, M. R., 2003. Getting control of intestinal secretion: thoughts for 2003. *Digestive and Liver Disease* 35, 378–385.
- Fawole, O. A., Finnie, J. F., Van Staden, J., 2009. Antimicrobial activity and mutagenic effects of twelve traditional medicinal plants used to treat ailments related to the gastro-intestinal tract in South Africa. *South African Journal of Botany* 75, 356–362.
- Fennell, A. W., Lindsey, K. L., McGaw, L. J., Sparg, S. G., Stafford, G. I., Elgorashi, E. E., Grace, O. M., van Staden, J., 2004. Assessing African medicinal plants for efficacy and safety: Pharmacological screening and toxicology. *Journal of Ethnopharmacology* 95, 205-217.
- Field, M., 2003. Intestinal ion transport and the pathophysiology of diarrhea. *Journal of Clinical Investigation* 111, 931–943.
- Fiorucci, S., Meli, R., Bucci, M., Cirino, G., 2001. Dual inhibitors of cyclooxygenase and 5-lipoxygenase. A new avenue in anti-inflammatory therapy? *Biochemical Pharmacology* 62, 1433–1438.
- Fitzgerald, G. A., 2004. Coxibs and cardiovascular disease. *New England Journal of Medicine* 351, 1709–1711.
- Forbes, V.S. (Ed.), 1986. *Carl Peter Thunberg Travels at the Cape of Good Hope 1772–1775*. Van Riebeeck Society, Cape Town, ISBN: 0620109815.
- Forgacs, I., Patel, V., 2011. Diabetes and gastrointestinal tract. *Medicine* 39 (5), 288-292.
- Fouche, G., Cragg, G.M., Pillay, P., Kolesnikova, N., Maharaja, V.J., Senabe, J., 2008. *In vitro* anticancer screening of South African plants. *Journal of Ethnopharmacology* 119, 455–461.
- From, C., Hormazabal, V., Granum, P. E., 2007. Food poisoning associated with pumilacidin-producing *Bacillus pumilus* in rice. *International Journal of Food Microbiology* 115, 319–324.
- Gadewar, S., Fasano, A., 2005. Current concepts in the evaluation, diagnosis and management of acute infectious diarrhea. *Current Opinion in Pharmacology* 5, 559–565.
- Gaginella, S. T., Kachur, J. F., Tamai, H., Kershavarzian, A., 1995. Reactive oxygen and Nitrogen metabolites as mediators of Secretory Diarrhea. *Gastroenterology* 109, 2019-2028.
- Gale, G. A., Kirtikara, K., Pittayakhajounwut, P., Sivichai, S., Thebtaranonth, Y., Thongpanchang, C., Vichai, V., 2007. In search of cyclooxygenase inhibitors, anti-*Mycobacterium tuberculosis* and anti-malarial drugs from Thai flora and microbes. *Pharmacology and Therapeutics* 115, 307-351.
- Galvez, J., Zarzuelo, A., Crespo, M. E., Lorente, M. D., Ocete, M. A., Jimenez, J., 1993. Antidiarrhoeic activity of *Euphorbia hirta* extract and isolation of an active flavonoid constituent. *Planta Medica* 59(4), 333-336.
- Gambhir, I. S., Nath, G., Jaiswal, J., Gopal, A., 2006. *Candida* Associated Acute Diarrhoea in Elderly. *Journal of the Indian Academy of Geriatrics* 2, 57-60.
- Gangoue-Pieboji, J., Baurin, S., Frere, J. M., Ngassam, P., Ngameni, B., Azebaze, A., Pegnyemb, D. E., Watchueng, J., Goffin, C., Galleni, M., 2007. Screening of some medicinal plants from cameroonCameroon for beta-lactamase inhibitory activity *Phytotherapy Research* 21(3), 284-287.
- Gautam, R., Saklani, A., Jachak, S., 2007. Indian medicinal plants as a source of antimycobacterial agents. *Journal of Ethnopharmacology* 110, 200–234.
- Gelberg, B. H., 2007. Pathology of organ systems: Alimentary system. In: *Pathologic Basis of Veterinary Diseases*. (Edited by McGavin MD and Zachary JF) Mosby Elsevier 11830 West line Industrial Drive, St Louis, Mussolini 63146 Pp 301-391.
- Geronikaki, A. A., Gavalas, A. M., 2006. Antioxidants and anti-inflammatory Diseases: synthetic and natural antioxidants with anti-inflammatory activity. *Combinatorial chemistry and High Throughput Screening* 9, 425-442.

- Gertsch, J., Viverose-Paredes, J. M., Taylor, P., 2010. Plant immunostimulants—Scientific paradigm or myth?. *Journal of Ethnopharmacology* xxx, xxx–xxx.
- Giess, W., Snyman, J. W., 1986. The naming and utilization of plant life by the Zu'hoasi Bushmen of the Kau-Kauveld. In: Vossen, R., Keuthmann, K. (Eds.), *Contemporary Studies on Khoisan I and II. In honour of Oswin Köhler on the Occasion of His 75th Birthday*. Buske, Hamburg, pp. 237–346.
- Gilani, A. H., Bashir, S., Janbaz, K. H. and Khan, A., 2005a. Pharmacological basis for the use of *Fumaria indica* in constipation and diarrhoea. *Journal of Ethnopharmacology* 96, 585–589.
- Gilani, A. H., Bashir, S., Jambaz, K. H., Shah, A. J., 2005b. Presence of cholinergic and calcium channel blocking activities explains the traditional use of *Hibiscus rosasinensis* in constipation and diarrhoea. *Journal of Ethnopharmacology* 102, 289–294.
- Gilani, A. H., Bashir, S., Jambaz, K. H., Shah, A. J., 2005. Presence of cholinergic and calcium channel blocking activities explains the traditional use of *Hibiscus rosasinensis* in constipation and diarrhoea. *Journal of Ethnopharmacology* 102, 289–294.
- Gilani, A. H., Rahman, A., 2005. Trends in Ethnopharmacology. *Journal of Ethnopharmacology*, 100, 43–49.
- Godfraind, T., Miller, R., Wibo, M., 1986. Calcium antagonism and calcium entry blockade. *Pharmacological Reviews* 38, 321–416.
- Goncalves, J. L., Lopes, R. C., Oliviera, D. B., Costa, S. S., Miranda, M. M., Romanos, M. T., Santos, N. S., Wigg, M. D., 2005. *In vitro* anti-rotavirus activity of some medicinal plants used in Brazil against diarrhea. *Journal of Ethnopharmacology* 99, 403–407.
- Gonzalez, A. G., Bazzocchi, I. L., Moujir, L., Jimenez, I. A., 2000. Ethnobotanical uses of Celastraceae: Bioactive metabolites studies. *Natural Products Chemistry* 23, 649-738.
- Gorowara, S., Sapru, S., Ganguly, N. K., 1998. Role of intracellular second messengers and ROS in the pathophysiology of *V. Cholerae* 0139 treated rabbit ileum. *Biochimica Biophysica Acta* 1407, 21–30.
- Gould, M., Sellin, J. H., 2009. Diabetic diarrhea. *Current gastroenterology reports* 11950, 354-359.
- Granot, E., Kohen, R., 2004. Oxidative stress in childhood-in health and disease states. *Clinical Nutrition* 23, 3–11.
- Granum, E. P., 2006. Bacterial toxins as food poisons: Clinical, immunological aspects and applications of Bacterial protein toxins: The comprehensive sourcebook of Bacterial protein toxins (Ed. Alouf and Popoff) Pp. 949-958.
- Greca, M. D., Manaco, P., Previtiera, L., 1990. Stagmasterol from *Typha latifolia*. *Journal of Natural Products*, 53, 1430–1435.
- Green, D. R., Reed, J. C., 1998. Mitochondria and apoptosis. *Science* 281, 1309– 1312.
- Green, E., Samie, A., Obi, C. L., Bessong, P. O., Ndip, R. N., 2010. Inhibitory properties of selected South Africa medicinal plants against *Mycobacterium tuberculosis*. *Journal of Ethnopharmacology* 130, 151-157.
- Greenwood, D. (ed.) 1989. Antibiotic sensitivity testing. In: *Antimicrobial Chemotherapy*. Oxford University, Oxford, p.91-100.
- Griendling, K. K., 2005. ATVB in focus: redox mechanisms in blood vessels. *Arterios Thrombotic and Vascular Biol.* 25, 272–273.
- Grierson, D. S., Afolayan, A. J., 1999. An ethnobotanical study of plants used for the treatment of wounds in the Eastern Cape, South Africa. *Journal of Ethnopharmacology* 67, 327–332.
- Groschwitz, K. R., Hogan, S. P., 2009. Intestinal barrier function: Molecular regulation and disease pathogenesis. *Journal of Allergy and Clinical Immunology* 124, 3–20 quiz 21–22.

- Grosser, T., Fries, S., FitzGerald, G. A., 2006. Biological basis for the cardiovascular consequences of COX-2 inhibition: Therapeutic challenges and opportunities. *Journal of Clinical Investigation* 116, 4-15.
- Guerrant, R. L., Oria, R., Bushen, O. Y., Patrick, P. D., Houpt, E., Lima, A. A. M., 2005. Global impact of diarrhoeal diseases that are sampled by Traveller: The rest of the Hippopotamus. *Clinical Infectious Diseases* 4, 5524-5530.
- Guerrant, R. L., Steiner, T. S., Lima, A. A. M., Bobak, D. A., 1999. How Intestinal Bacteria Cause Disease. *The Journal of Infectious Diseases* 179(2), S331-S337.
- Gupta, S., Ali, M., Alam, M. S., Niwa, M., Sakai, T., 1992. 2, 4 β -Ethylcholest-4-en-3 β -ol from the roots of *Lawsonia inermis*. *Phytochemistry* 31, 2558–2560.
- Gupta, T. P., Ehrinpreis, M. N., 1990. Candida-associated diarrhea in hospitalized patients. *Gastroenterology* 98, 780–785.
- Gutierrez, R. M. P., Mitchell, S., Solis, R. V., 2008. *Psidium guajava*: A review of its traditional uses, phytochemistry and pharmacology. *Journal of Ethnopharmacology* 117, 1–27
- Gutierrez, S. P., Sanchez, M. A. Z., Gonzalez, C. P., Garcia, L. A., 2007. Antidiarrhoeal activities of different plants used in traditional medicine. *African Journal of Biotechnology* 6(25), 2988-2994.
- Gutteridge, J.M.C., 1988. Lipid peroxidation: some problems and concepts, in: B. Halliwell, (Ed.), *Oxygen Radicals and Tissue Injury*. FASEB, Bethesda, MD, pp. 9–19.
- Guttman, J. A., Finlay, B. B., 2008. Subcellular alterations that lead to diarrhea during bacterial pathogenesis. *Trends in Microbiology* 16(11), 535-542.
- Guttman, J. A., Samji, F.N., Li, Y., Vogl, A. W., Finlay, B. B., 2006. Evidence that tight junctions are disrupted due to intimate bacterial contact and not inflammation during attaching and effacing pathogen infection in vivo. *Infections and Immunology* 74, 6075–6084.
- Ha, D. T., Kim, H., Thuong, P. T., Ngoc, T. M., Lee, I., Hung, N. D., Bae, K., 2009. Antioxidant and lipoxygenase inhibitory activity of oligostilbenes from the leaf and stem of *Vitis amurensis*. *Journal of Ethnopharmacology* 125, 304–309.
- Ha, T. J., Kubo, I., 2005. Lipoxygenase Inhibitory Activity of Anacardic Acids. *Journal of Agricultural and Food Chemistry* 53(11), 4350-4354.
- Haeggström, J. Z., Rinaldo-Matthis, A., Wheelock, C. E., Wetterholm, A., 2010. Advances in eicosanoid research, novel therapeutic implications. *Biochemical and Biophysical Research Communications* 396, 135–139.
- Halliwell, B., 1997. Antioxidants and human disease: a general introduction. *Nutrition Reviews* 55, 544-552.
- Halliwell, B., 2008. Are polyphenols antioxidants or pro-oxidants? What do we learn from cell culture and in vivo studies? *Archives of Biochemistry and Biophysics* 476, 107–112.
- Halliwell, B., Gutteridge, J. M., 1990. Role of free radicals and catalytic metal ions in human disease: an overview. *Methods in Enzymology*. 186, 1–85.
- Hamza, O. J. M., van den Bout-van den Beukel, C. J. P., Mates, M. I. N., Moshi, M. J., Mikx, F. H. M., Selemani, H. O., Mbwambo, Z. H., Van der Van, A. J. A. M., Verweij, P. E., 2006. Antifungal activity of some Tanzania plants used traditionally for the treatment of fungal infection. *Journal of Ethnopharmacology* 108, 124-132.
- Han, T., Li, H-L., Zhang, Q-Y., Han, P., Zheng, H-C., Rahman, K., Qin, L-P., 2007. Bioactivity guided fraction for antiinflammatory and analgesic properties and constituent of *Xynthium strumarium* .L. *Phytomedicine*, 14(12), 825-829.
- Havagiray, R., Ramesh, C., Sadhna, K., 2004. Study of antidiarrhoeal activity of *Calotropis gigantea* r.b.r. in experimental animals. *Journal of Pharmacology and Pharmaceutical Science* 7, 70–75.

- Havsteen B. H., 2002. The biochemistry and medical significance of the flavonoids. *Pharmacology and Therapeutics* 96, 67–202 *Sci.* 57, 6–15.
- Heimler, D., Vignolini, P., Dini, M. D., Vincieri, F. F., Romani, A., 2006. Antiradical activity and polyphenol composition of local Brassicaceae edible varieties. *Food Chemistry*, 99, 464-469.
- Henry-Stanley, M. J., Hess, D. J., Erickson, E. A., Garni, R. M., Wells, R. M., 2003. Effect of Lipopolysaccharide on Virulence of Intestinal *Candida albicans*. *Journal of Surgical Research* 113, 42–49.
- Hermon-Taylor, J., 2006. Gut pathogens: Invaders and turncoats in a complex cosmos. *Gut pathogens* 1:3 [hppt://www.gutpathogen.com/content/1/1/3](http://www.gutpathogen.com/content/1/1/3)
- Hodges, K., Gill, R., 2010. Infectious diarrhea Cellular and molecular mechanisms. *Gut Microbes* 1, 4-21.
- Hofmann, A. F., 1977. Bile acids, diarrhea, and antibiotics: data, speculation, and a unifying hypothesis. *Journal Infectious Diseases* 135(supplement), S126–S132.
- Holzzapfel, C., Van Wyk, B., Castro, A., Marias, W., Herbst, M., 1995. A Chemotaxonomic Survey of Kaurene Derivatives in the Genus *Alepidea* (Apiaceae). *Biochemikal .Qstematics and Ecology*, Vol. 23, No. 718, pp. 799-803.
- Holzer, P. 2004. Gastrointestinal afferents as targets of novel drugs for the treatment of functional bowel disorders and visceral pain. *European Journal of Pharmacology* 429, 177–193.
- Homans, A. L., Fuchs A., 1970. Direct bioautography on thin layer chromatograms as a method for detecting fungitoxic substances. *Journal of Chromatography* 51, 327–329.
- Hoogerwerf, W. A., 2006. Prokineticin 1 inhibits spontaneous giant contractions in the murine proximal colon through nitric oxide release. *Neurogastroenterology and Motility* 18, 455–463.
- Hopkins, M. J., Macfarlane, G. T., 2003. Nondigestible oligosaccharides enhance bacterial colonization resistance against *Clostridium difficile* in vitro. *Applied and Environmental Microbiology* 69, 1920–1927.
- Hostettmann, K., Marston, A., 2002. Twenty years of research into medicinal plants: Results and perspectives. *Phytochemistry Review*1, 275-285.
- Hostettmann, K., Terreaux, C., Marston, A., Potterat, O., 1997. The role of planar chromatography in the rapid screening and isolation of bioactive compounds from medicinal plants. *Journal of Planar Chromatography-Mod. TLC*, 10, 251–257.
- Hostettmann-Kaldas, M., Nakanishi, K., 1979. Moronic acid, a simple triterpenoids keto acid with antimicrobial activity isolated from *Ozoroa mucronata*. *Planta Medica* 37(4), 358-360.
- Hou, W., Li, Y., Zhan, Q., Wei, X., Peng, A., Chen, L., Wei, Y., 2009. Triterpene Acids Isolated from *Lagerstroemia speciosa* Leaves as α -Glucosidase Inhibitors. *Phytother. Res.* 23, 614–618
- Hu, L., Tall, B. D., Curtis, S. K., Kopecko, D. J., 2008. Enhanced microscopic definition of *Campylobacter jejuni* 81-176 adherence to, invasion of, translocation across, and exocytosis from polarized human intestinal Caco-2 cells. *Infection and Immunity* 76(11), 5294–5304.
- Hutchings, A., Scott, A. H., Lewis, G., 1996. *Zulu Medicinal Plants: An Inventory*. University of Natal Press, Durban, p. 266–267.
- Huycke, M. M., Moore, D. R., 2002. *In vivo* production of hydroxyl radical by *enterococcus faecalis* colonizing the intestinal tract using aromatic hydroxylation. *Free Radical Biology & Medicine*, 33(6), 818–826.
- Imanishi, K., 1993. Aloctin A, an active substance of *Aloe arborescens* Miller as an immunomodulator. *Phytotherapy Research*, 7(7), S20-S22.

- Iwalewa, E. O., McGaw, L. J., Naidoo, V., Eloff, J. N., 2007. Inflammation: the foundation of diseases and disorders. A review of phytomedicines of South African origin used to treat pain and inflammatory conditions. *African Journal of Biotechnology* 6 (25), 2868-2885.
- Iwalokun, B. A., Gbenle, G. O., Adewole, T. A., Akinsinde, K. A., 2001. Shigellocidal properties of three Nigerian medicinal plants: *Ocimum gratissimum*, *Terminalia avicennoides*, and *Momordica balsamina*. *Journal of Health, Population and Nutrition* 19(4), 331-335.
- Jager, A. K., Van Staden, J., 2005. Cyclooxygenase inhibitory activity of South African plants used against inflammation. *Phytochemistry Review* 4, 39-46.
- Jay, J. M., 1996. *Modern Food Microbiology* 5th edn New York, NY: Chapman and Hall.
- Jeller, A. H., Silva, D. H. S., Liao, L. M., Bolzani, V. S., Furlan, M., 2004. *Phytochemistry* 65, 1977.
- Johns, T., Faubert, G. M., Kokwaro, J. O., Muhunnah, R. L. A., Kimanam, 1995. Anti-giardial activity of gastrointestinal remedies of the Luo of East African. *Journal of Ethnopharmacology* 46, 17-23.
- Johnson, A. M., Kaushik, R. S., Hardwidge, P. R., 2010. Disruption of transepithelial resistance by enterotoxigenic *Escherichia coli*. *Veterinary Microbiology* 141, 115-119.
- Jordan, S. A., Cunningham, D. G., Marles, R. J., 2010. Assessment of herbal medicinal products: challenges, and opportunities to increase the knowledge base for safety assessment, *Toxicol. Ogy and Appl. ied Pharmacol. ogy* 243 (2010) 198–216.
- Joseph, C. C., Moshi, M. J., Innocent, E., Nkunya, M. H. H., 2007. Isolation of a Stilbenes Glucoside and Other constituent of *Terminalia sericea*. *African Journal of Traditional and Complementary, Alternative Medicine*, 4(4), 383-386.
- Kaikabo, A. A., Suleiman, M. M., Samuel, B. B., Eloff, J. N., 2008. Antibacterial activity of eleven South African plants use in treatment of diarrhoea in folkloric medicine. *African Journal of Traditional, Complementary and Alternative Medicine* 5, 315–316.
- Kamatou, G. P. P., Viljoen, A. M., Steenkamp, P., 2010. Antioxidant, anti-inflammatory activities and HPLC analysis of South African *Salvia* species. *Food Chemistry* 119, 684–688.
- Kane, A. B., 1996. Mechanisms of Cell and Tissue Injury. In: *Cellula and Molecular Pathogenesis*. Editor, Alphonse E. Sirica: Lippincott-Raven Publishers, 277 East Washington Square, Philadelphia, Pennsylvania 19106-3780 Pp 1-22.
- Katerere, D. R., Gray, A. I., Nash, R. J., Waigh, R. D., 2003. Antimicrobial activity of pentacyclic triterpenes isolated from African Combretaceae. *Phytochemistry* 63, 81-88.
- Katsoulis, L. C., Veale, D. J. H., Havlik, I., 2000. The pharmacological action of *Rhoicissus tridentata* on isolated rat uterus and ileum. *Phytotherapy* 14(6), 460-462.
- Kaviarasan, S., Naik, G. H., Gangabhagirathi, R., Anuradha, C. V., Priyadarsini, K. I., 2007. In vitro studies on antiradical and antioxidant activities of fenugreek (*Trigonella foenum graecum*) seeds. *Food Chemistry* 103, 31–37.
- Kayser, O., Kolodziej. H., 1997. Antibacterial activity of extracts and constituent of *Pelargonium sidoides* and *Pelargonium reniforme* Curtis
- Kayser, O., Kolodziej. H., Kiderlen, A. F., 2001. Immunomodulatory principle of *Pelargonium sidoides*. *Phytotherapy Research* 15(2), 122-126.
- Kehrer, J. P., Biswal, S. S., 2000. The molecular effects of acrolein. *Toxicological Sciences* 57, 6–15.
- Kenny, J. M., Kelly, P., 2009. Protozoal gastrointestinal infections. *Medicine* 39, 599-602.

- Kermanshai, R., McCamy, B. E., Rosenfeld, J., Summers, P. S., Weretilnyk, E. A., Sorger, G. J., 2001. Benzyl isothiocyanate is the chief or sole anthelmintic in Papaya seed extracts. *Phytochemistry* 57(3), 427-435.
- Khajuria, A., Gupta, A., Garai, S., Wakhloo, B. P., 2007. Immunomodulatory effects of two saponins 1 and 2 isolated from *Luffa cylindrica* in Balb/C mice. *Bioorganic and Medicinal Chemistry Letters* 17, 1608–1612.
- Khaled, M. A., 1994. Oxidative stress in childhood malnutrition and diarrheal disease. *Journal of Diarrheal Disease Research* 12, 165–72.
- Kim, H. S., 2009. 5-Hydroxytryptamine 4 receptor agonists and colonic motility. *Journal of Smooth Muscle Research* 45, 25–29.
- Kinghorn, A. B., Chai, H-B., Sung, C-K., Keller, W-J., 2011. The classical drug discovery approach to defining bioactive constituents of botanicals. *Fitoterapia* 82, 71-79.
- Kishida, E., Tokumaru, S., and Ishitani, Y., 1993. Comparison of the formation of malondialdehyde and thiobarbituric acid reactive substances from autoxidized fatty acids based on oxygen consumption. *Journal of Agricultural and Food Chemistry*, 41, 1598-1600.
- Koduru, S., Grierson, D. S., Afolayan, A. J., 2006. Antimicrobial activity of *Solanum aculeastrum*. *Pharmaceutical Biology* 44(4), 283-286.
- Koduru, S., Grierson, D. S., Van der Venter, M., Afolayan, A. J., 2007: Anticancer activity of steroid alkaloids isolated from *Solanum aculeastrum*. *Pharmaceutical Biolog* 45(8), 613-618.
- Koff, R. S., 1992. Clinical manifestation and diagnosis of hepatitis A virus infection. *Vaccine* 10, S15-S17.
- Koff, R. S., 1998. Hepatitis A. *Lancet* 341, 1643–1649.
- Komane, B. M., Olivier, E., Alvaro, I. Viljoen, M., 2011. *Trichilia emetica* (Meliaceae) – A review of traditional uses, biological activities and Phytochemistry. *Phytochemistry Letters* 4, 1–9.
- Koopmans, M., 2008. Progress in understanding norovirus epidemiology. *Current Opinion on Infectious Diseases* 21, 544–552.
- Kosar, M., Bozan, B., Temelli, F., Baser, K. H. C., 2007. Antioxidant activity and phenolic composition of sumac (*Rhus coriaria* L.) extracts. *Food Chemistry* 103, 952-959.
- Kougan, G. B., Miyamoto, T., Mirjolet, J-F., Duchamp, O., Sondengam, B. L., Lacaille-Dubios, M-A., 2009. Arboreasides A-E, Triterpene Saponins from the Bark of *Cussonia arborea*. *Journal of Natural Products* 72, 1081–1086.
- Krakauer, T., Li, B. Q., Young, H. A., 2001. The flavonoids baicalin inhibits superantigens-induced inflammatory cytokines and chemokines. *Federation of European Biochemical Societies Letter* 500, 52-55.
- Krishnaiah, D., Sarbatly, R., Nithyanandam, R., 2011. A review of the antioxidant potential of medicinal plant species. *Food and Bioprocess Technology* 89, 217–233.
- Kubo, I., Kim, M., Naya, K., Komatsu, S., Yamagiwa, Y., Ohashi, K., Sakamoto, Y., Hirakawa, S., Kamikawa, T., 1987. Prostaglandin synthetase inhibitors from the African medicinal plant " *Ozoroa mucronata*. *Chemistry letters*, 1101-1104.
- Kuete, V. Ngameni, B., Fotso Simo, C. C., Kengap Tankeu, R., Tchaleu Ngadjui, B., Meyer d, N., Lall, J. J. M., Kuate, J. R., 2008. Antimicrobial activity of the crude extracts and compounds from *Ficus chlamydocarpa* and *Ficus cordata* (Moraceae). *Journal of Ethnopharmacology* 120, 17–24.
- Kuete, V., Krusche, B., Youns, M., Voukenga, I., Fankam, A. G., Tankeo, S., Lacmata, S., Efferth, T., 2011. Cytotoxicity of some Cameroonian spices and selected medicinal plant extracts. *Journal of Ethnopharmacology* 134, 803–812.

- Kuete, V., Nana, F., Ngameni, B., Mbaveng, A. T., Keumedjio, F., Chaleu Ngadjui, B. T., 2009. Antimicrobial activity of the crude extract, fractions and compounds from stem bark of *Ficus ovata* (Moraceae). *Journal of Ethnopharmacology* 124, 556–561.
- Kunkel, S. L., Lukacs, N., Strieter, R. M., 1996. Cytokines and Inflammatory disease. In: *Cellular and Molecular Pathogenesis*. Editor, Alphonse E. Sirica: Lippincott-Raven Publishers, 277 East Washington Square, Philadelphia, Pennsylvania 19106-3780 Pp 23-36.
- Kunle, O. O., Shittu, A., Nasipuri, R. N., Kunle, O. F., Wambembe, C., Akah, P. A., 1999. Gastrointestinal activity of *Ficus sur*. *Fitoterapia* 70(6), 542-547.
- Lajubutu, B. A., Pinney, R. L., Roberts, M. F., Odelola, H. A., Oso, B. A., 1995. Antibacterial activities of diosquinone and plumbagin from the root of *Diospyros mespiliformis* (Hostch) (Ebenaceae). *Phytotherapy Research*, 9(5), 346-350.
- Lall, N., Meyer, J. J. M., 2001. Inhibition of drug-sensitive and drug-resistant strains of *Mycobacterium tuberculosis* by Diospyrin, isolated from *Euclea natalensis*. *Journal of Ethnopharmacology* 78(2-3), 213-216.
- Laohachai, K. N., Bahadi, R., Hardo, M. B., Hardo, P. G., Kourie, J. I., 2003. The role of bacterial and non-bacterial toxins in the induction of changes in membrane transport: implications for diarrhea. *Review: Toxicon* 42, 687–707.
- Lansky, E. P., Newman, R. A., 2007. Review: *Punica granatum* (pomegranate) and its potential for prevention and treatment of inflammation and cancer. *Journal of Ethnopharmacology* 109, 177–206.
- Lattanzio, V., Lattanzio, V. M. T., Cardinali, A., 2006. Role of phenolics in the resistance mechanisms of plants against fungal pathogens and insects. *Phytochemistry: Advances in Research* 23-67.
- Lauwaet, T., Oliveira, M. J., Callewaert, B., De Bruyne, G., Mareel, M., Leroy, A., 2004. Proteinase inhibitors TPCCK and TLCK prevent *Entamoeba histolytica* induced disturbance of tight junctions and microvilli in enteric cell layers in vitro. *International Journal for Parasitology* 34(7), 785–794.
- Le Bouguenec, C., 2005. Adhesins and Invasins of pathogenic *Escherichia coli*. *International journal of medical microbiology* 295, 471-478.
- Lee, C. W., Sarma, S. K., Singaram, C. Casper, M. A., 1997. Ca²⁺ channel blockage by verapamil inhibits GMCs and diarrhea during small intestinal inflammation. *American Journal of Physiology* 273, 785-794.
- Lee, G. Y., Byeon, S. E., Kim, J. Y., Lee, J. Y., Rhee, M. H., Hong, S., Wu, C. J., Lee, H. S., Kim, M. J., Cho, D. H., Cho, J. Y., 2007a. Immunomodulating effect of *Hibiscus cannabinus* extract on macrophage functions. *Journal of Ethnopharmacology* 113, 62-71.
- Lee, H. M., Lee, J. M., Jun, S. H., Lee, H. S., Kim, W. N., Lee, J. H., Ko, N. Y., Mun, S. H., Kim, B. K., Lim, B. O., Choi, D. K., Choi, W. S., 2007b. The anti-inflammatory effects of *Pyrolae herba* extract through the inhibition of the expression inducible nitric oxide synthase (iNOS) and NO production. *Journal of Ethnopharmacology* 112, 49-54.
- Lee, J., Durst, R. W., Wrolstad, R. E., 2005. Determination of Total Monomeric Anthocyanin pigment content of fruit juices, Beverages, Natural colourants, and Wines by the pH differential method: Collaborative study. *Journal of AOAC International* 88, 1269-1278.
- Lee, J., Rennaker, C., Wrolstad, R. E., 2008. Correlation of two anthocyanin quantification methods: HPLC and spectrophotometric method. *Food Chemistry* 110, 782-786.
- Lee, S.H., Oe, T., Blair, I.A., 2001. Vitamin C-induced decomposition of lipid hydroperoxides to endogenous genotoxins. *Science* 292, 2083–2086.
- Lengsfeld, C., Faller, G., Hansel, A., 2007. Okra polysaccharides inhibit adhesion of *Campylobacter jejuni* to mucosa isolated from poultry *in vitro* but not *in vivo*. *Animal Feed Science and Technology* 135, 113–125.

- Levine, J., Dykoski, R. K., Janoff, E. N., 1995. Candida-Associated Diarrhea: A Syndrome in Search of Credibility. *Clinical Infectious Diseases* 21 (4), 881-886.
- Lewu, F. B., Afolayan, A. J., 2009. Ethnomedicine in South Africa: The role of weedy species. *African Journal of Biotechnology* 8 (6), 929-934.
- Li, Y., Trush, M. A., 1994. Reactive oxygen dependent DNA damage resulting from the oxidation of phenolic compounds by a copper redox cycle. *Cancer Research* 54, 1895s-1898s.
- Lin, C-N., Haung, A-M., Lin, K-W., Hour, T-C., Ko, H-H., Yang, S-C., Pu, Y-S., 2010. Xanthine oxidase inhibitory terpenoids of *Amentotaxus formosana* protect cisplatin-induced cell death by reducing reactive oxygen species (ROS) in normal human urothelial and bladder cancer cells. *Phytochemistry* 71, 2140–2146.
- Lin, J., Opoku, A. R., Geheeb-Keller, M., Hutchings, A. D., Terblanche, S. E., Jager, A. K., van Staden, J., 1999. Preliminary screening of some traditional Zulu medicinal plants for anti-inflammatory and anti-microbial activities. *Journal of Ethnopharmacology* 68, 267–274.
- Lin, J., Puckree, T., Mvelase, T. P., 2002. Anti-diarrhoeal evaluation of some medicinal plants used by Zulu traditional healers. *Journal of Ethnopharmacology* 79(1), 53-6.
- Lindsey, K. L., Motsei, M. L., Jager, A. K., 2002. Screening of South African food plants for anti oxidant activity. *Journal of Food Science* 64, 2129–2931.
- Linscott, A. J., 2011. Food-borne illnesses. *Clinical Microbiology Newsletter* 33, 41-45.
- Lis-Balchin, M., Hart, S., Simpson, E., 2001. Buchu (*Agathosma betulina*, *Agathosma crenulata* (Rutaceae)) essential oils: their pharmacological activity on guinea pig ileum and antimicrobial activity on microorganism. *Journal of Pharmacy and Pharmacology* 53(4), 579-582.
- Liu, Y., Abreu, P., 2006. Tirucallane triterpenes from the roots of *Ozoroa insignis*. *Phytochemistry* 67, 1309–1315.
- Lode, H., 2010. Safety and Tolerability of Commonly Prescribed Oral Antibiotics for the Treatment of Respiratory Tract Infections. *The American Journal of Medicine* 123, S26-S38.
- Lorrot, M., Vasseur, M., 2007. How do the rotavirus NSP4 and bacterial enterotoxins lead differently to diarrhea? *Virology Journal* 4, 31
- Lourens, A. C. U., Van Vuuren, S. F., Viljoen, A. M., Van Herdeen, F. R., 2011. Antimicrobial activity and in vitro cytotoxicity of selected South African *Helichrysum* species. *South African Journal of Botany*, 77(1), 229-235.
- Lourens, A. C. U., Viljoen, A. M., van Heerden, F. R., 2008. South African *Helichrysum* species: A review of the traditional uses, biological activity and phytochemistry. *Journal of Ethnopharmacology* 119, 630–652.
- Lund, T., DeBuyser, M. L., Granum, P. E., 2000. A new cytotoxin from *Bacillus cereus* that may cause necrotic enteritis. *Molecular Microbiology* 38, 254–261.
- Luseba, D., Elgorashi, E. E., Ntloedibe, B. T., Van Staden, J., 2007. Antibacterial, anti-inflammatory and mutagenic effects of some medicinal plants used in South Africa for the treatment of wounds and retained placenta in Livestock. *South African Journal of Botany* 73, 378-383.
- Luseba, D., Van der Merwe, D., 2006. Ethnoveterinary medicine practices among Tsonga speaking people of South Africa. *Onderstepoort Journal of Veterinary Research* 73, 115–122.
- Lutterodt, G. D., Ismail, A., Basheer, R. H., Baharuddin, M. H., 1999. Antimicrobial effects of *Psidium guajava* extracts as one mechanism of its antidiarrhea action. *Malaysian Journal of medicinal Sciences*. Makkar HPS, Francis G, Becker K. 2007: Bioactivity of phytochemicals in some lesser-known plants and their effects and potential applications in livestock and aquaculture production system. *Animal* 1(9), 1371- 1391, 17-20.

- Lutterodt, G.D., 1992. Inhibition of Microlax-induced experimental diarrhoea with narcotic-like extracts of *Psidium guajava* leaf in rats. *Journal of Ethnopharmacology* 37, 51–157
- Lyckander, I. M., Malterud, K. E., 1992. Lipophilic flavonoids from *Orthosiphon spicatus* as inhibitors of 15-lipoxygenase. *Acta Pharmaceutica Nordica* 4, 159–166.
- Mabogo, D. E. N., 1990. The ethnobotany of the VhaVenda. M. Sc. dissertation, University of Pretoria.
- MacNaughton, W. K., 2006. Mechanisms and Consequences of Intestinal Inflammation. *Physiology of Gastrointestinal Tract*, Fourth Edition, edited by Leonard R. Johnson. Academic Press.
- Mahato, S. B., Kundu, A. P., 1994. ¹³C-NMR spectra of pentacyclic triterpenoids. A compilation and some salient features. *Phytochemistry* 37, 1517–1575.
- Majhenic, L., Skerget, M., and Knez, Z., 2007. Antioxidant and antimicrobial activity of guarana seed extracts. *Food Chemistry* 104, 1258-1268.
- Makhlouf, M. G., 1994. Neuromuscular function of the small intestine, In: Johnson, Leonard R. (Ed.), *Physiology of the Gastrointestinal Tract*, Third ed. Raven Press, New York, pp. 977–990.
- Makkar, H. P. S., 2003. Quantification of tannins in tree and foliage- a laboratory manual. Klumer Academic Press Dordrecht, The Netherland.
- Makkar, H. P. S., Francis G, Becker K. 2007. Bioactivity of phytochemicals in some lesser-known plants and their effects and potential applications in livestock and aquaculture production system. *Animal* 1(9), 1371- 1391.
- Males, Z., Medic-Saric, M., 2001. *Journal of Pharmacology and Biomedical Analysis* 24 (2001) 353.
- Mamphiswara, N. D., Mashela, P. W., Mdee, L. K., 2010. Distribution of total phenolics and antioxidant in fruit, leaf, stem and root of *Monsonia burkeana*. *African Journal of Agricultural Research*, 5(18), 2570-2575.
- Mandal, S., Nayak, A., Kar, M., Banerjee, S. K., Das, A., Upadhyay, S. N., Singh, R. K., Banerji, A., Banerji, J., 2010. Antidiarrhoeal activity of carbazole alkaloids from *Murraya koenigii* Spreng (Rutaceae) seeds. *Fitoterapia* 81, 72-74.
- Mansfield, L. S., Gajadhar, A. A., 2004. *Cyclospora cayetanensis* a food-and water-borne mcoccidian parasite. *Veterinary Parasitology* 126, 73–90.
- Maregesi, S. M., Pieters, L., Ngassapa, O. D., Apers, S., Vingerhoets, R., Cos Paul, Berghe, Van den Dirk, A., Vlietinck, A. J., 2008. Screening of some Tanzanian medicinal plants from Bunda district for antibacterial, antifungal and antiviral activities. *Journal of Ethnopharmacology* 119, 58-66.
- Marino, S. D., Gala, F., Borbone, N., Zollo, F., Vitali, S., Visioli, F., Lorizzi, M., 2007. Phenolic glycoside from *Foeniculum vulgare* fruit and evaluation of antioxidative activity. *Phytochemistry* 68(13), 1805-1812.
- Marquez-Martin, A., Puerta, R. D. L., Fernandez-Arche, A., Ruiz-Gutierrez, V., Yaqoob, P., 2006. Modulation of cytokine secretion by pentacyclic triterpenes from olive pomace oils in human mononuclear cells. *Cytokine*, 36(5-6), 211-217.
- Marston, A., 2011. Review: Thin-layer chromatography with biological detection in *Phytochemistry*. *Journal of Chromatography A*, 1218, 2676–2683.
- Martinez-Augustin, O., Romero-Calvo, I., Suarez, M. D., Zarzuelo, A., and Sanohez de Medina, F., 2009. Molecular bases of impaired water and ion movement in inflammatory bowel diseases. *Inflammatory Bowel Diseases* 15, 114-127.
- Masika, P. J., Afolayan, A. J., 2002. Antimicrobial activity of some plants used for the treatment of livestock disease in the Eastern Cape, South Africa. *Journal of Ethnopharmacology* 83, 129-134.
- Masika, P. J., Sultana, N., Afolayan, A. J., 2004. Antibacterial activity of two flavonoids isolated from *Schotia latifolia*. *Pharmaceutical Biology* 42(2), 105-108.

- Masoko, P., Picard, J., Eloff, J. N., 2005. Antifungal activities of six South African *Terminalia* species (Combretaceae). *Journal of Ethnopharmacology* 99, 301-308.
- Masoko, P., Picard, J., Eloff, J. N., 2007. The antifungal activities of twenty-four Southern southern Africa *Combretum* species (Combretaceae). *South African Journal of Botany* 73, 173-183.
- Mastroeni, P., Maskell, D., 2006. *Salmonella* infections: Clinical, immunological, and molecular aspects. Cambridge University Press.
- Mathabe, M. C., Nikolova, R. V., Lall, N., Nyazema, N. Z., 2006. Antibacterial activities of medicinal plants used in the treatment of diarrhea disease in Limpopo Province, South Africa. *Journal of Ethnopharmacology* 105, 286-293.
- Mativandela, S. P. N., Lall, N., Meyer, J, J, M., 2006. Antibacteria, antifungal, antitubercular activityactivities of the root of *Pelargonium reniforme* Curtis and *Pelargonium sidoides* (DC) (Geraniaceae) root extract. *South African Journal of Botany* 72(2), 233-237.
- Mativandela, S. P. N., Meyer, J, J, M., Hussein, A. A., Houghton, P. J., Hamilton, C. J., Lall, N., 2008. Activity against *Mycobacterium smegmatis* and *Mycobacterium tuberculosis* by South African Medicinal plants. *Phytotherapy Research* 22(6), 841-845.
- Matsui, M., Motomura, D., Fujikawa, T., Jiang, J., Takahashi, S., Manabe, T., Taketo, M.M., 2002. Mice lacking M2 and M3 muscarinic acetylcholine receptors are devoid of cholinergic smooth muscle contractions but still viable. *The Journal of Neuroscience* 22, 10627–10632.
- Mattison, K., 2010. Norovirus as a food-borne disease hazard. *Advances in Food and Nutrition Research* 62, 1-39.
- Matu, E. N., van Staden, J., 2003. Antibacterial and anti-inflammatory activity of some plants used for medical purposes in Kenya. *Journal of Ethnopharmacology* 87, 35-41.
- Mbagwu, H. O. C., Adeyemi O. O., 2008. Anti-diarrhoeal activity of the aqueous extracts of *Mezoneuron benthamianum* Baill (Caesalpiniaceae) *Journal of Ethnopharmacology* 116, 16-20.
- Mbatchi, S. F., Mbatchi, B., Banzouzi, J. T., Basmba, T., Nsonde Ntandou, G. F., Ouamba, J–M., Berry, A., Benoit-Vical, F., 2000. *In vitro* antiplasmodial activity of 18 plants used in Congo Brazzaville traditional medicine. *Journal of Ethnopharmacology* 104, 168-174.
- McGaw, L. J., Eloff, J. N., 2008. Ethnoveterinary use of Southern African plants and scientific evaluation of their medicinal properties. *Journal of Ethnopharmacology* 119: 559–574.
- McGaw, L. J., Jäger, A. K., van Staden, J., 2000. Antibacterial, anthelmintic and anti-amoebic activity in South African medicinal plants. *Journal of Ethnopharmacology* 72, 247–263.
- McGaw, L. J., Jager, A. K., van Staden, J., 2002. Isolation of antibacterial fatty acids from *Schotia brachypetala*. *Fitoterapia* 73(5), 431-433.
- McGaw, L. J., Rabe, T., Sparg, S. G., Jager, A. K., Eloff, J. N., van Staden, J., 2001. An investigation on the biological activity of *Combretum* species. *Journal of Ethnopharmacology* 75: 45–50.
- McGaw, L. J., Van der Merwe, D., Eloff, J. N., 2007. In vitro anthelmintic, antibacterial and cytotoxic effects of extracts from plant used in South African ethnoveterinary medicine. *The veterinary Journal* 173, 366-372.
- Meghashri, S., Vijar Kumar, S., Gopal, S., 2010. Antioxidant properties of aof novel flavonoids from leaves of *Leucas aspera*. *Food Chemistry* 122(1), 105-110.
- Mello, D. L., ALves, A. A., Macedo, D. V., Kkota, L. T., 2005. Peroxidase-based biosensor as a tool for a fast evaluation of antioxidant capacity of tea. *Food Chemistry* 92, 515-519.

- Michelangeli, F., Ruiz, M. C., 2003. Physiology and pathology of the gut in relation to viral diarrhoea. *Viral gastroenteritis*: U. Desselberger and J. Gray (Editors). Elsevier Science BV. Pp 23-50.
- Middleton, E., Jr., Kandaswami, C., and Theoharides, T. C., 2000. The effects of plant flavonoids on mammalian cells: Implications for inflammation heart disease and cancer. *Pharmacology Reviews*, 52(4), 673–751.
- Miliauskas, G., van Beek, T. A., de Waard, P., Venskutonis, R. P., and Sudholter, E. J. R. 2005. Identification of radical scavenging compounds in *Rhaponticum carthamoides* by means of LC-DADSPE- NMR. *Journal of Natural Products*, 68(2), 168–172.
- Mishra, A., Kumar, A., 2000. Medicinally important trees of Rajasthan. *International Journal of Mendel* 1–8, 37–38.
- Mlambo, N. P., 2008. The screening of medicinal plants traditionally used to treat diarrhoea, in Ongoye area, Kwazulu Natal. A dissertation submitted to the Faculty of Science at the University of Zululand in fulfillment of the requirement for the degree of Master of Science.
- Mølgaard, P., Nielsen, S. B., Rasmussen, D. E., Drummond, R. B., Makaza, N., Andreassen, J., 2001. Anthelmintic screening of Zimbabwean plants traditionally used against schistosomiasis. *Journal of Ethnopharmacology* 74, 257–264.
- Molla, A., Viljoen, A. M., 2008. Buchu- *Agathosma betulina* and *Agathosma crenulata* (Rutaceae): A Review. *Journal of Ethnopharmacology* 119: 413–419.
- More, G., Tshikalange, T. E., Lall, N., Botha, F., Meyer, J. J. M., 2008: . Antimicrobial activity of medicinal plants against oral microorganisms. *Journal of Ethnopharmacology*, 119, 473–477.
- Moshi, M. J., Mbwambo, Z. H., 2005. Some pharmacological properties of extract of *Terminalia sericea* roots. *Journal of Ethnopharmacology* 97(1), 43-47.
- Mosmann, T., 1983. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *Journal of Immunological Methods* 65, 55–63.
- Motlanka, D. M. T., 2008. Free radical scavenging activity of selected medicinal plants of Eastern Botswana. *Pakistan Journal of Biological Sciences* 11(5), 805-808.
- Moundipa, P. F., Flore, K. G. M., Bilong Bilong, C. F., Bruchhau, I., 2005. In vitro amoebicidal activity of some medicinal plants of the Bamu region (Cameroon). *African Journal of Traditional and Complimentary Medicine* 2(2), 113-121.
- Moure, A., Cruz, J. M., Franco, D., Dominquez, J. M., Sineiro, J., Dominquez, H., Nunez, M. J., and Parajo, J. C., 2001. Natural antioxidant from residual sources. *Food Chemistry* 72, 145-171.
- Mueller, C.F., Laude, K., McNally, J.S., Harrison, D.G., 2005. ATVB in focus: redox mechanisms in blood vessels. *Arterioscler. Thromb. Vasc. Biol.* 25, 274–278.
- Muhammed, I., El-Sayed, K. A., Mossa, J. S., Al-Said, M. S., El-Ferally, F. S., Clark, A. M., Hufford, C. D., Oh, S., Mayer, A. M. S., 2000. Bioactive 12-Olaenene triterpene and secotriterpene acids from *Maytenus undata*. *Journal of Natural Products* 63, 605-610.
- Mulaudzi, R. B., Ndhala, A. R., Kulkarni, M. G., Finnie, J. F., Van Staden, J., 2011. Antimicrobial properties and phenolic contents of medicinal plants used by Venda people for conditions related to venereal diseases. *Journal of Ethnopharmacology*, DOI: 10: 1016/J.jep2011.03.022.
- Mulaudzi, R. B., Ndhala, A. R., Finnie, J. F., Van Staden, J., 2009. Antimicrobial, anti-inflammatory and genotoxicity activity of *Alepidea amatymbica* and *Alepidea natalensis* (Apiaceae). *South African Journal of Botany* 75, 584–587.
- Murata, T., Miyase, T., Muregi, F. W., Naoshima-Ishibashi, Y., Umshara, K., Warashina, T., Kanou, S., Mkoji, G. M., Terada, M., Ishih, A., 2008. Antiplasmodial triterpenoids from *Ekebergia capensis*, *Journal of Natural Product* 71, 167-174.

- Murayama, T., Eizuru, Y., Yamada, R., Sadanari, H., Matsubara, K., Rukung, G., Tolo, F. M., Mungai, G. M., Kofi-Tsekpo, M., 2007. Anticytomegalovirus activity of pristimerin, a triterpenoids quinine methide isolated from *Maytenus heterophylla* (Eckl. & Zeyh.). *Antiviral Chemistry and Chemotherapy* 18(3), 133-139.
- Mutanyatta, J., Bezabih, M., Abegaz, B. M., Dreyer, M., Brun, R., Kocher, N., Bringmann, G., 2005. The first 6'-O-sulfated phenylanthraquinones: Isolation from *Bulbine frutescens*, structural elucidation enantiomeric purity and partial synthesis. *Tetrahedron* 61, 8475-8484.
- Muthaura, C. N., Rukungu, G. M., Chhabra, S. C., Omar, S. A., Guantai, A. N., Gathirwa, J. W., Tolo, F. M., Mwitari, P. G., Keter, L. K., Kirira, P. G., Kimani, C. W., Mungai, G. M., Njagi, E. N. M., 2007. Antimalarial activity of some plants traditionally used in treatment of malaria in Kwale district of Kenya. *Journal of Ethnopharmacology* 112, 545-551.
- Naczki, M., Shahidi, F., 2004. Extraction and analysis of phenolics in food. *Journal of Chromatography A*, 1054, 95-111.
- Naidoo, L. A. C., Drewes, S. E., Van Staden, J., Hutching, J., 1992. Exocarpic acid and other compounds from tubers and inflorescences of *Sarcophyte sanguine*. *Phytochemistry* 31(11), 3929-3931.
- Naidoo, V., McGaw, L. J., Bisschop, S. P. R., Duncan, N., Eloff, J. N., 2008. The value of plant extracts with antioxidant activity in attenuating coccidiosis in broiler chickens. *Veterinary Parasitology* 153, 215-219.
- Naidoo, V., Zweygarth, E., Eloff, J. N., Swan, G. E., 2005. Identification of anti-babesial activity for four ethnoveterinary plants in vitro *Veterinary Parasitology* 130(1-2), 9-13, 2005.
- Naidoo, V., Zweygarth, E., Swan, G. E., 2006. Determination and quantification of the in vitro activity of *Aloe marlothii* (A. Berger) subsp. *marlothii* and *Elephantorrhiza elephantina* (Burch.) skeels acetone extracts against *Ehrlichia ruminantium*. *Onderstepoort Journal of Veterinary Research* 73, 175-178.
- Nanayakkara, N. P., Burandt, C. L. Jr., Jacob, M. R., 2002. Flavonoid with activity against methicillin-resistant *Staphylococcus aureus* from *Dalea scandens* var. *paucifolia*. *Planta medica* 68, 519-522.
- Nardi, G. M., Siqueira Junior, J. M., Monache Belle, F., Pizzolatti, M. G., Ckless, K., Ribeiro-do-Valle, R. M., 2007. Antioxidant and anti-inflammatory effects of products from *Croton celtidifolius* Bailon on carrageenan-induced pleurisy in rats. *Phytomedicine* 14, 115-122.
- Nathan, C., 1992. Nitric Oxide as a secretory product of mammalian cells. *FASEB J.* 6: 3051-3064.
- National Committee for Clinical Laboratory Standards (NCCLS), 2002. *Performance Standards for Antimicrobial Susceptibility testing*. Twelfth Information Supplement M100 - S12 NCCLS, Wayne.
- Naz, S., Siddiqi, R., Ahmad, S., Rasool, S. A., Sayeed, S. A., 2007. Antibacterial activity directed isolation of compounds from *Punica granatum*. *Journal of Food Science* 72, M341-M345.
- Ndamba, J., Nyazema, N., Makaza, N., Anderson, C., Kaondera, K. C., 1994. Traditional herbal remedies used for the treatment of urinary schistosomiasis in Zimbabwe. [Comparative Study] *Journal of Ethnopharmacology* 42(2), 125-32
- Ndawonde, B. G., Zobolo, A. M., Dlamini, E. T., Siebert, S. J., 2007. A survey of plants sold by traders at Zululand muthi markets, with a view to selecting popular plant species for propagation in communal gardens. *African Journal of range and Forage Science* 42(2), 103-107(5).
- Ndhlala, A. R., Finnie, F. R., Van staden, J., 2010. Plant composition, Pharmacological properties and mutagenic evaluation of commercial Zulu herbal mixture: *Imbiza ephuzwato*. *Journal of Ethnopharmacology* 133(2), 663-674.
- Neish, A. S., 2009. Microbes in Gastrointestinal Health and Disease. *Gastroenterology* 136, 65-80.

- Nergard, C. S., Diallo, D., Michealsen, T. E., Malteru, K. E., Kiyohara, H., Matsumoto, T., Yamada, H., Paulsen, B. S., 2004. Isolation, partial characterization and immunomodulating activity of polysaccharides from *Vernonia kotschyana* Sch. Bip. Ex. Walp., *Journal of Ethnopharmacology* 91(1), 141-152.
- Neuwinger, H. D., 1996. *African ethnobotany: poisons and drugs: chemistry, pharmacology, toxicology*. Chapman and Hall, Germany.
- Nguemeving, J. R., Azebaze, A. G. B., Kuete, V., Carly, N. N. E., Beng, V. P., Meyer, M., Blond, A., Bodo, B., Nkengfack, A. E., 2006. Laurentixanthonones A and B, antimicrobial xanthonones from *Vismia laurentii*. *Phytochemistry* 67, 1341–1346.
- Ngueyem, T. A., Brusotti, G., Caccialanza, G., Vita Finzi, P., 2009. The genus *Bridelia*: A phytochemical and ethnopharmacological review. *Journal of Ethnopharmacology* 124, 339–349.
- Nhiem, N. X., Tai, B. H., Quang, T. H., Kiem, P. V., Minh, C. V., Nam, N. H., Kim, J-H., Im, L-R., Lee, Y-M., Kim, Y. H., 2011. A new ursane-type triterpenoid glycoside from *Centella asiatica* leaves modulates the production of nitric oxide and secretion of TNF- α in activated RAW 264.7 cells. *Bioorganic & Medicinal Chemistry Letters* 21, 1777–1781.
- Nigam, S., Schewe, T., 2000. Phospholipase A2 and lipid peroxidation. *Biochimie and Biophysics Acta* 1488, 167–181.
- Nissanka, A. P. K., Karunaratne, V., Bandara, B. M. R., Kumar, V., Nakanishi, T., Nishi, M., Inada, A., Tillekeratne, L. M. V., Wijesundara, D. S. A., Gunatilaka, A. A. L., 2000. Antimicrobial alkaloids from *Zanthophyllum tetraspermum* and *caudatum*. *Phytochemistry* 56, 857-861.
- Noreen, Y., Ringbom, T., Perera, P., Danielson, H., Bohlin, L., 1998. Development of a radiochemical Cyclooxygenase-1 and -2 in vitro assay for identification of natural products as inhibitors of prostaglandin biosynthesis. *Journal of Natural Products* 61, 2-7
- Nose, K., 2000. Role of reactive oxygen species in regulation of physiological functions. *Biological and Pharmaceutical Bulletin* 22, 897-903.
- Nyiredy, S. Z.; Glowniak, K., 2001. Planar chromatography in medicinal plant research. In *Planar Chromatography*; Nyiredy, Sz. Ed.; Springer Scientific Publisher: Budapest, Hungary 550.
- O'Hara, S. P., Chen, X-M., 2011. The cell biology of *cryptosporidium* *Cryptosporidium* infection. *Microbes and Infection* xx, 1-10.
- Oben, J. E., Assi, S. E., Agbor, G. A., Musoro, D. F., .2006. Effect of *Eremomastax speciosa* on experimental diarrhea. *African Journal of Traditional Complimentary Alternative Med.* 3(1), 95-100.
- Obi, C. L., Potgieter, N., Bessong, P. O., Masebe, T., Mathebula, H., Molobela, P., 2003. In vitro antibacterial activity of Venda medicinal plants. *South African Journal of Botany* 69(2), 1-5.
- Odhav, B., Kanasamy, T., Khumalo, N., Baijnath, H., 2010. Screening of African traditional vegetables for their alpha-amylase inhibitory effect. *Journal of Medicinal Plant Research*, 4(14), 1502-1507.
- Ofek, I., Sharon, N., 1990. Adhesins as lectins: Specificity and role in infection. In: K, Jann B (eds) *Bacterial capsules and adhesins: Fact and Principles*. Current topic in Microbiology and Immunology. Vol 151, Berlin: Springer, 91-113.
- Ogoina, D., Onyemelukwe, G. C., 2009. The role of infections in the emergence of non-communicable diseases (NCDs): Compelling needs for novel strategies in the developing world. *Journal of Infection and Public Health* 2, 14-29.
- Ojewole, J. A. O., 2004. Evaluation of the Antidiabetic, Anti inflammatory and Antidiabetic properties of *Sclerocarya birrea* (A. Rich) Hochst. Stem bark aqueous Extract in Mice and Rats. *Phytotherapy Research* 18, 601-608.
- Ojewole, J. A. O., 2005. Antinociceptive, antiinflammatory and antidiabetic effects of *Leonotis leonurus* (L) R. BR. [Lamiaceae] leaf aqueous extract in mice and rats. *Methods and Findings in Experimental and Clinical Pharmacology* 27, 257–264

- Ojewole, J. A. O., 2006. Antinociceptive, anti-inflammatory and antidiabetic properties of *Hypoxis hemerocallidea* Fisch. and C. A. Mey. (Hypoxidaceae) Corm ('African potato') in mice and rats. *Journal of Ethnopharmacology* 103(1), 126-134.
- Ojewole, J. A. O., Awe, E. O., Nyinawumuntu, A., 2009. Antidiarrhoeal Activity of *Hypoxis hemerocallidea* Fisch. and C. A. Mey. (Hypoxidaceae) Corm ('African potato') Aqueous Extract in Rodents. *Phytotherapy Research*, 23, 965–971.
- Ojewole, J. A. O., Mawoza, T., Chiwororo, W. D. H., Owira, P. M. O., 2010. *Sclerocarya birrea* (A. Rich.) Hochst. ['Marula'] (Anacardiaceae): a review of its phytochemistry, pharmacology and toxicology and its ethnomedicinal uses. *Phytotherapy Research* 24, 633–639.
- Okokon, J. E., Ita, B. N., Udokpoh, A. E., 2006. The in vivo antimalarial activities of *Uvaria chamae* and *Hippocratea africana*. *Annals of Tropical Medicine and Parasitology* 100, 585–590.
- Orabi, K. Y., Al-Qasoumi, S., El-Olemy, M. M., Mosaa, J. S., Muhammed, I., 2001. Dihydragarofuran alkaloid and triterpenes from *Maytenus hetrophylla* and *Maytenus arbutifolia*. *Phytochemistry* 58(3), 475-480.
- Oskay, M., Oskay, D., Kalyoncu, F., 2009. Activity of some plants extracts against multi drug resistant human pathogens. *Iranian Journal of Pharmaceutical Research* 8, 293-300.
- Oskay, M., Sari, D., 2007. Antimicrobial screening of some Turkish medicinal plants. *Pharmaceutical Biology* 45(3), 176-181.
- Ovenden, P. B., Yu, J., Bernays, J., Wanss Christophidis, L. J., Sberna, G., Tait, R. A., Wildman, H. G., Labeller, D., Lowther, J., Walsh, N. G., Meurer-Grimes, B. M., 2004. Physaloside A, an Acylated sucrose ester from *Physalis viscosa*.
- Owolabi, O. J., Omogbai, E. K. I., 2007. Analgesic and anti-inflammatory activities of the ethanolic stem bark extract of *Kigelia africana* (Bignoniaceae). *African Journal of Biotechnology*, 6(5), 582-585.
- Owolabi, O. J., Omogbai, E. K. I., Obasuyi, O., 2007. Antifungal and antibacterial activities of the ethanolic and aqueous extracts of *Kigelia africana* (Bignoniaceae). *African Journal of Biotechnology* 6(14), 1677-1680.
- Owolabi, O. J., Nworgu, Z. A., Odushu, K., 2010. Antidiarrhoeal evaluation of the ethanolic extract of *Nauclea latifolia* root bark. *Methods and Findings in Experimental and Clinical Pharmacology* 32(8), 551-555.
- Ozyurek, M., Bektasoglu, B., Guclu, K., Apak, R., 2008. Hydroxyl radical scavenging assay of phenolics and flavonoids with a modified cupric reducing antioxidant capacity (CUPRAC) method using catalase for hydrogen peroxide degradation. *Analytica Chimica Acta* 616, 196–206.
- Pallant, C. A., and Steenkamp, V., 2008. In-vitro bioactivity of Venda medicinal plants used in the treatment of respiratory conditions. *Human & Experimental Toxicology* 27, 859–866.
- Palombo, E. A., 2006. Phytochemicals from traditional medicinal plants used in the treatment of diarrhoea: modes of action and effects on intestinal function. *Phytotherapy Research* 20, 717–724.
- Palombo, E. A., 2006. Traditional plants and Herbal Remedies used in the treatment of Diarrhea Disease: Mode of action, Quality, Efficacy and Safety Consideration. In: *Modern phytomedicine, turning medicinal plants into drugs.* (Edited by Ahmed F Aqil and Owais M) WILEY-VCH Verlag GmbH and Co KGaA, Weinheim. Pp 248-269.
- Panda, D., Patra, C. R., Nandi, S., Swarup, D., 2009. Oxidative stress indices in gastroenteritis in dogs with canine parvoviral infection. *Research in veterinary science* 86, 36-42.
- Parekh, J., Chandra, S., 2007. Antibacterial and phytochemical studies on twelve species of Indian medicinal plants. *African Journal of Biomedical Research* 10, 175-181.
- Parola, M., Bellomo, G., Robino, G., Barrera, G., Dianzani, M. U., 1999. 4-Hydroxynonenal as a biological signal: molecular basis and pathophysiological implications. *Antioxidant and Redox Signalling* 1, 255–284.

- Pash, Y., Banks, M., Farthing, M., 2009. Causes and recommended management of acute diarrhoea. *Prescriber*, January, 16-34.
- Pauli, G. F., Junior, P., 1995. Phenolic glucosides from *Adonis Aleppica*. *Phytochemistry* 38, 1245-1250.
- Pavlick, K. P., Laroux, F. S., Fuseler, J., et al., 2002. Role of reactive metabolites of oxygen and nitrogen in inflammatory bowel disease. *Free Radical Biol. ogy and Medicine*. 33, 311–322.
- Peluso, I., Campolongo, P., Valeri, L. R., Palmery, M., 2002. Intestinal motility disorder induced by free radicals: a new model mimicking oxidative stress in gut. *Pharmacological Research* 46 (6), 533-538.
- Pendota, S. C., Yakubu, M. T., Grierson, D. S., Afolayan, A. J., 2010. Effect of administration of aqueous extracts of *Hippobromus pauciflorus* leaves in male Wistar rats. *African journal of Traditional, Complimentary and Alternative Medicine* 7(1): 40-46.
- Pendota, S. C., Yakubu, M. T., Grierson, D. S., Afolayan, A. J., 2009. Antiinflammatory, analgesic, antipyretic activities of the aqueous extract of *Pippobromus puaciflorus* (L. F.) Radk leaves in Male Wistar rats. *African Journal of Biotechnology*, 8(10), 2036-2041.
- Perez-Bosque, A. and Moreto, M., 2010. A rat model of mild intestinal inflammation induced by *Staphylococcus aureus* enterotoxins B. The 3rd International Immunonutrition Workshop held in Spain on 21-24 Oct., 2009. *Proceedings of the Nutrition Society*, 69, 447-453.
- Petri Jr., W. A., Miller, M., Binder, H. J., Levine, M. M., Dillingham, R., Guerrant, R. L., 2008. Enteric infections, diarrhea, and their impact on function and development. *Journal of Clinical. Investigation* 118:1277–1290.
- Pharoah, D. S., Varsan, H., Tathan, R. W., 2006. Expression of the inflammatory chemokines CCL5, CCL3 and CXCL10 in juvenile idiopathic arthritis and demonstration of CCL5 production by a typical subset of CD8⁺T cells. *Arthritis Research and Therapy* 8, R50.
- Picerno, P., Autore, G., Marzocco, S., Meloni, M., Sanago, R., Aquino, R. P., 2005. Anti-inflammatory activity of cutaneous irritation of cell cullins and reconstituted Human epidermis. *Journal of Natural Products* 68(11): 1610-1614.
- Pietta, P. G., 2000. Flavonoids as antioxidants. *Journal of Natural Products*, 63, 1035–1042.
- Pillai, N. R., 1992. Antidiarrhoeal activity of *Punica granatum* in experimental animals. *Pharmaceutical Biology* 30(3), 201-204.
- Pillay, P., Maharaj, V. J., Smith, P. J., 2008. Investigating South African plants as a source of new antimalarial drugs. *Journal of Ethnopharmacology* 119, 438-454.
- Podewils, L. J., Mintz, E. D., Nataro, J. P., Parashar, U. D., 2004. Acute, Infectious Diarrhoea Among Children in Developing Countries. *Journal of seminars in Pediatric Infectious Diseases* 155-168.
- Polya, G. M., 2003. *Biochemical Targets of Plant Bioactive Compounds. A Pharmacological Reference Guide to Sites of Action and Biological Effects*. CRC Press, Florida.
- Ponou, B. K., Barboni, L., Teponno, R. B., Mbiantcha, M., Nguelefack, T. B., Park, H-J., Lee, K-T., Taponjou, L. A., 2008. Polyhydrooleanane-type triterpenoids from *Combretum molle* and their anti-inflammatory activity. *Phytochemistry letters* 1(4), 183-187.
- Pooley, E., 1998. *A Field Guide to Wildflowers KwaZulu-Natal and the Eastern Region*. Natal Flora Publications Trust, Durban.
- Prior, R. L., Wu, X., Schaich, K., 2005. Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements. *Journal of Agriculture and Food Chemistry* 53 (10), 4290–4302.

- Prozesky, E. A., Meyer, J. J. M., Louw, A. L., 2001. In vitro antiplasmodial activity and cytotoxicity of ethnobotanically selected South African plants. *Journal of Ethnopharmacology* 76 (3), 81-308.
- Pujol J., 1990. *Natur Africa: The Herbalist Handbook*. Natural Healers Foundation, Durban.
- Puyvelde, V. L., De Kimpe, N., Costa, J., Munyjabo, V., Nyirankuliza, S., Hakizamungu, E., Schamp, N., 1989. Isolation of flavonoids and chalcone from *Helichrysum odoratissimum* and synthesis of Helichrysetin. *Journal of Natural Product* 52(3), 629-633.
- Quais, E. Y., Elokda, A. S., Ghalyun, Y. Y. A., Abdulla, F. A., 2007. Antidiarrhoeal activity of the aqueous extract of *Punica granatum* (Pomegranate) peels. *Pharmaceutical Biology* 45(9), 715-720.
- Rabe, T., Van Staden, J., 1997. Antibacterial activity of South African plants used for medicinal purposes. *Journal of Ethnopharmacology* 56, 81–87.
- Raccach, M., 1984. The antimicrobial activity of phenolic antioxidant in foods: a review. *Journal of Food Safety* 6, 141-170.
- Radi, Z. A., Khan, N. K., 2006. Effects of cyclooxygenase inhibition on the gastrointestinal tract. *Experimental and Toxicological pathology* 58, 163-173.
- Ragone, M. I., Sella, M., Cornforti, P., Volonte, M. G., Consolini, A. E., 2007. The spasmolytic effect of *Aloysia citriodora*, Palau (South American cedron) is partially due to its vitexin but not isovitexin on rat duodenum. *Journal of Ethnopharmacology* 113, 258–266.
- Rahman, I., Adcock, I. M., 2006. Oxidative stress and redox regulation of lung inflammation in COPD. *European Respiratory Journal* 28, 219-242.
- Ralston, K. S., Petri Jr., W. A., 2011. Tissue destruction and invasion by *Entamoeba histolytica*. *Trends in Parasitology*, 27, 253-262.
- Ramahivhana, J. N., Moyo, S. R., Obi, C. I., 2010. The possible role of medicinal plants in tackling resistant microbial pathogens in Limpopo Province, South Africa. *Journal of Medicinal plant Research*, 4(11), 999-1002.
- Ramaroa, N., Lereclus, D., 2006. Adhesion and cytotoxicity of *Bacillus cereus* and *Bacillus thuringiensis* to epithelial cells are FlhA and PlcR dependent, respectively. *Microbes and Infection* 8, 1483-1491.
- Ramful, D., Bahorun, T., Bourdon, E., Tarnus, E., Aruoma, O. I., 2010. Bioactive phenolics and antioxidant propensity of flavedo extracts of Mauritian citrus fruits: Potential prophylactic ingredients for functional foods application. *Toxicology xxx* (2010) xxx–xxx.
- Rao, V. S. N., Santos, F. A., Sobreika, T. T., Souza, M. F., Melo, L. L., Silveira, E. R., 1997. Investigations on the gastroprotective and antidiarrhea properties of ternatin, a tetramethoxyflavone from *Egletes viscosa*. *Planta Medica* 63(2), 146-149.
- Rasoanaivo, O., Ramanitrahambola, D., Rafatro, H., Rakotondramanana, D., Robijaona, B., Rakotozafy, A., Ratsimamanga-Urverg, S., Labaied, M., Grellier P., Allorge, L., Mambu, L., Frappier, F., 2004. Screening of extracts of Madagascan plants in search of antiplasmodial compounds, *Phytotherapy Research*, 18, 743-747.
- Rasoanaivo, P., Ratsimamanga-Urverg, S., 1993. Biological Evaluation of Plants with Reference to the Malagasy Flora. Monograph for the IFS-NAPRECA Workshop on Bioassays, Antananarivo, Madagascar, pp 9–43, 72–83.
- Rauha, J. P., 2001. The research for biological activity in Finnish plant extracts containing phenolic compounds, PhD Thesis, Pharmacognosy Department, Pharmacy Faculty of Science, University of Helsinki, pp. 31–45
- Ravikumara, M., 2008. Investigation of chronic diarrhoea. *Paediatrics and Child health* 18(10), 441-447.
- Re, R., Pellegrini, N., Proteggente, A., Pannala, A., Yang, M., Rice-Evans, C., 1999. Antioxidant activity applying improved ABTS radical cation decolourization assay. *Free Radical Biology and Medicine* 26, 1231-1237.

- Rea Al. Schmidt JM. Setzer WN. Sibanda S. Taylor C. Gwebu ET., 2003. Cytotoxic activity of *Ozoroa insignis* from Zimbabwe *Fitoterapia* 74(7-8), 732-735.
- Reid, K. A., Maes, J., Maes, A., van Staden, J., De Kimpe, N., Mulholland, D. A. Verschaeve, L., 2006. Evaluation of the mutagenic and antimutagenic effects of South African plants. *Journal of Ethnopharmacology* 106, 44–50.
- Rekka, E., Kourounakis, P. N., 1991. Effect of hydroxyethyl rutenosides and related compounds on lipid peroxidation and free radical scavenging activity-some structural aspects. *Journal of Pharmacy and Pharmacology* 43, 486–491.
- Reyes, C. P. Nunez, M. J. Jimenez, I. A. Busserolles, J., Alcaraz, M. J., Bazzocchi, I. L., 2006. Activity of lupane triterpenoids from *Maytenus* species as inhibitors of nitric oxide and prostaglandin E2. *Bioorganic & Medicinal Chemistry* 14, 1573–1579.
- Rhee, I. K., Van De Meent, M., Ingkaninan, K., Verpoorte, R., 2001: . A screening for acetylcholinesterase inhibitors from Amaryllidaceae using silica gel thin-layer chromatography in combination with bioactivity staining. *Journal of Chromatography A* 915, 217–223.
- Rietjens, I. M. C. M., Boersma, M. G., Van der Woude, H., Jeurissen, S. M. F., Schutte, M. E., Alink, G. M., 2005. Flavonoids and alkenylbenzenes: mechanisms of mutagenic action and carcinogenic risk. *Mutation Research* 574, 124–138.
- Rijke, E., Out, P., Niessen, W. M. A., Ariese, F., Gooijer C., Brinkman, U. A., 2006. Analytical separation and detection methods for flavonoids. *Journal of Chromatography A* 1112, 31–63.
- Ríos, J. L., 2008. *Ganoderma lucidum*, unhongos con potenciales propiedades inmunoestimulantes. *Revista de Fitoterapia* 8, 135–146.
- Ríos, J. L., Bas, E., Recio, M. C., 2005. Effects of natural products on contact dermatitis. *Current Medicinal Chemistry. Anti-inflammatory and Anti-allergic Agents* 4, 65–80.
- Rios, J. L., Recio, M. C., 2005. Medicinal plants and antimicrobial activity. *Journal of Ethnopharmacology* 100, 80-84.
- Risa, J., Risa, A., Adersen, A., Gauguin, B., Stafford, G. I., Van Staden, J., Jager, A. K., Screening of plants used in South Africa for epilepsy and convulsions in the GABA_A-benzodiazepine receptor assay. *Journal of Ethnopharmacology* 93(2-3), 177-182.
- Robak, J., Duniec, Z., Rzakowska Bodalska, H., Olechnowicz Stepien, W., Cisowski, W., 1986. The effect of some flavonoids on non-enzymatic lipid oxidation and enzymatic oxidation of arachidonic acid. *Pol. ish Journal of Pharmacognosy and Pharmaceutical* 38, 483-491.
- Robert, K., Egon, S., Daneila, B., Florian, D., Christoph, W., Gunter, J. K., Emil, C. R., 2001. Role of *Candida* in antibiotic-associated diarrhea. *Journal of infectious diseases* 184, 1065-1069.
- Roberts, M., 1990. *Indigenous Healing Herbs*. Southern Book Publishers, South Africa, pp. 1–285.
- Rocha Martin, L. R., Brenzan, M. A., Nakamura, C. V., Dias Filho, B. P., Nakamura, T. U., Ranei Cortez, L. E., Garcia Cortez, D. A., 2011. In vitro antiviral activity from *acanthospermum australei* on *herpesvirus* and *poliovirus*. *Pharmaceutical Biology* 49 (1), 26-31.
- Rocha-Martins, L. R. R., Benzan, M. A., Nakamura, C. V., Filho, B. P. D., Nakamura, T. U., Cortez, E. R., Cortez, D. A. G., 2010. In vitro antiviral activity of *Acanthospermum australe* on herpevirus and poliovirus. *Pharmaceutical biology* 49(1), 26-31.
- Rosa, A., Pollastro, F., Afzeri, A., Appendino, G., Melis, M. P., Dieana, M., Incani, A., Loru, D., Dessi, M. A., 2011. Protective role of arzanol against lipid peroxidation in biological systems. *Chemistry and Physics of lipids* 164, 24-32.

- Rosengren, Å., Fabricius, A., Guss, B., Sylvén S., Lindqvist, R., 2010. Occurrence of foodborne pathogens and characterization of *Staphylococcus aureus* in cheese produced on farm-dairies. International journal of Food microbiology
- Russo, A., Cardile, V., Lombardo, L., Vanela, L., Vanela, A., Garbarino, J. A., 2005. Antioxidant activity and antiproliferative action of methanolic extract of *Geum quellyon* sweet roots in human tumor cells lines. Journal of Ethnopharmacology 100, 323-332.
- Saini, N. K., Singhal, M., Srivastava, B., Sachdeva, K., Singh, C. C., 2011. Antimicrobial activity of *Tecomaria capensis* leaves extracts. International Journal of Pharmaceutical Sciences Review and Research 7(1), 121-124.
- Sairam, K., Hemalatha, S., Kumar, A., Srinivasan, T., Ganesh, J., Shankar, M., Venkataraman, S., 2003. Evaluation of anti-diarrhoeal activity in seed extracts of *Magnifera indica*. J. Ethnopharmacol. 84: 11-15.
- Salvemini, D., Masferrer, J. L., 1996. Interactions of nitric oxide with cyclooxygenase in vitro, ex vivo and in vivo studies. Methods of Enzymology 269:12-25.
- Samie, A., Obi, C. L., Bessong, P. O., Namrita, L., 2005. Activity profiles of fourteen selected medicinal plants from rural Venda communities in South Africa against fifteen clinical bacterial species. African Journal of Biotechnology, 4, 1443–1451.
- Samie, A., Obi, C. L., Lall, N., Meyer, J. J. M., 2009. In-vitro cytotoxicity and antimicrobial activities, against clinical isolates of *Campylobacter* species and *Entamoeba histolytica*, of local medicinal plants from the Venda region, in South Africa. Annals of Tropical Medicine & Parasitology 103 (2), 159–170.
- Samie, A., Tambani, J., Harshfield, E., Green, E., Ramalivhana, J. N., Bessong, P. O., 2010. Antifungal activities of selected Venda medicinal plants against *Candida albicans*, *Candida krusei* and *Cryptococcus neoformans* isolated from South African AIDS patients. African Journal of Biotechnology 9(20), 2965-2976.
- Samy, R. P., Gopalakrishnakone, P., 2008. Therapeutic Potential of Plants as Anti-microbials for Drug Discovery: Review eCAM 1-12.
- Sanders, K. M., 2001. Invited review: mechanisms of calcium handling in smooth muscles. Journal of Applied Physiology 91, 1438–1449.
- Saunders, D. R., Wiggins, H. S., 1981. Conservation of mannitol, lactulose, and raffinose by the human colon. American Journal of Physiology 1241, G397–E402.
- Savi, L. A., Caon, T., de Oliveira, A. P., Sobottka, A. M., Werner, W., Reginatto, F. H., Schenkel, E. P., Barardi, C. R. M., Simões, C. M. O., 2010. Evaluation of antirotavirus activity of flavonoids. Fitoterapia xxx, xxx–xxx.
- Saxena, G., Towers, G. H. N., Farmer, S., Hancock, R. E. W., 1995. Use of specific dyes in the detection of antimicrobial compounds from crude plant extracts using thin layer chromatography agar overlay technique. Phytochemical Analysis 6: 125-129.
- Schenk, M., Mueller, C., 2008. The mucosal immune system at the gastrointestinal barrier. Best Practice & Research Clinical Gastroenterology 22, 391–409.
- Schiller, L. R., 1999. Secretory diarrhea. Current Gastroenterology Reports 1, 389-397.
- Schmitz, H., Rokos, K., Florian, P., Gitter, A. H., Fromm, M., Scholz, P., et al., 2002. Supernatants of HIV-infected immune cells affect the barrier function of human HT-29/B6 intestinal epithelial cells. AIDS 16(7), 983–991.
- Schuijer, M., Sies, H., Illek, B., Fischer, H., 2005. Cocoa-Related Flavonoids Inhibit CFTR-Mediated Chloride Transport across T84 Human Colon Epithelia. Journal of Nutrition, 2320-2325.

- Schulzke, J. D., Troger, H., Amasheh, M., 2009. Disorders of intestinal secretion and absorption. *Best Practice & Research Clinical Gastroenterology* 23, 395–406.
- Schwikkard, S., Zhou, B.-N., Glass, T. E., Sharp, J. L., Mattern, M. R., Johnson, R. K., Kingston, D. G. I., 2000. Bioactive compounds from *Combretum erythrophyllum*. *Journal of Natural Products* 63, 457–460.
- Sebothoma, C., 2010. Isolation and characterization of antibacterial compounds from *Searsia (Rhus) leptodictya* Diels (Anacardiaceae). M. Sc. Thesis submitted to University of Limpopo
- Selvarani, V., Hudson, J. B., 2009. Multiple inflammatory and antiviral activities in *Adansonia digitata* (Baobab) leaves, fruit, and seeds. *Journal of Medicinal plant Research* 3(8), 576-582.
- Shai, L. J., McGaw, L. J., Aderogba, L. J., Mdee, L. K., Eloff, J. N., 2008b. Four pentacyclic triterpenoids with antifungal and antibacterial activity from *Curtisia dentata* (Burm.f) C.A. Sm. Leaves. *Journal of Ethnopharmacology* 118, 238–244.
- Shai, L. J., McGaw, L. J., Eloff, J. N., 2009. Extracts of the leaves and twigs of the threatened tree *Curtisia dentata* (Cornaceae) are more active against *Candida albicans* and other microorganisms than the stem bark extract. *South African Journal of Botany* 75, 363–366.
- Shai, L. J., McGaw, L. J., Masoko, P., Eloff J. N., 2008a. Antifungal and antibacterial activity of seven traditionally used South African plant species against *C. albicans*. *South African Journal of Botany* 74, 677-684.
- Shale, T. L., Stirk, W. A., van Staden, J., 2005. Variation in antibacterial and anti-inflammatory activity of different growth forms of *Malva parviflora* and evidence for synergism of the anti-inflammatory compounds. *Journal of Ethnopharmacology* 96, 325-330.
- Sharkey, K. A., Mawe, G. M., 2002. Neuroimmune and epithelial interactions in intestinal inflammation. *Current Opinion in Pharmacology* 2, 669-677.
- Sibandze, G. F., Van Zyl, R. L., Van Vuuren, S. F., 2010. The anti-diarrhoeal properties of *Bretonadia salicina*, *Syzygium cordatum* and *Ozoroa sphaerocarpa* when used in combination in Swazi traditional medicine. *Journal of Ethnopharmacology* 132(2), 506-511.
- Sikander, A., Rana, S. V., Prasad, K. K., 2009. Role of serotonin in gastrointestinal motility and irritable bowel syndrome. *Clinica Chimica Acta* 403, 47–55.
- Simmons, D. L., Botting, R. M., Hla, T., 2004. Cyclooxygenase isozymes: the biology of prostaglandin synthesis and inhibition. *Pharmacological Review* 56, 387-437.
- Simpson, A. J., McNally, D. J., Simpson, M. J., 2011. NMR spectroscopy in environmental research: From molecular interactions to global processes. *Progress in Nuclear Magnetic Resonance Spectroscopy* 58, 97–175.
- Slater, T. F., Cheeseman, K. H., Davies, M. J., Proudfoot, K., Xin, W., 1987. Free radical mechanisms in relation to tissue injury. *Proceedings of the Nutrition Society* 46, 1-12.
- Smid, S. D., Svensson, K. M., 2009. Inhibition of cyclooxygenase-2 and EP1 receptor antagonism reduces human colonic longitudinal muscle contractility *in vitro*. *Prostaglandins and other lipid mediators* 88, 117-121.
- Smirnoff, N., Cumbes, O. J., 1996. Hydroxyl radical scavenging activity of compatible solutes. *Phytochemistry* 28, 1057–1060.
- Soderholm, J. D., Perdue M. H., 2006. Effect of stress on intestinal mucosal function. *Physiology of the gastrointestinal tract, Fourth Edition*, edited by Leonard R. Johnson. Academic Press.
- Sousa, A., Ferreira, I. C. F. R., Calhelha R., 2006. Phenolic and antimicrobial activity of traditional stoned table olives “alcaparra”. *Bioorganic and Medicinal Chemistry* 14, 8533-8538.

- Spickett, A. M. Van Der Merwe, D., Matthe, O., 2007. The effect of orally administered *Aloe marlothii* leaves on Boophilus decoloratus tick burdens on cattle. *Experimental and Applied Acarol* 41, 139–146.
- Spiller, R. C., 2004. Inflammation as a basis for functional GI disorder. *Best Practice and Research Clinical Gastroenterology* 18, 641–661.
- Spiller, R., Garsed, K., 2009. Infection, inflammation and irritable bowel syndrome. *Digestive and Liver disease*, 41, 844-849.
- Sprague, A. H., Khalil, R. A., 2009. Inflammatory cytokines in vascular dysfunction and vascular disease. *Biochemical Pharmacology* 78, 539–552.
- Springfield, E. P., Amabeoku, G., Weitz, F., Mabusela, W., Johnson, Q., 2003. An assessment of two *Carpobrotus* species extracts as potential antimicrobial agents. *Phytomedicine* 10, 434–439.
- Stables, M. J., Gilroy, D. W., 2010. Old and new generation lipid mediators in acute inflammation and resolution. *Progress in Lipid Research*, doi:10.1016/j.plipres.2010.07.005.
- Stanzel, R. D. P., Lourenssen, S., Blennerhassett, M. G., 2008. Inflammation causes expression of NGF in epithelial cells of the rat colon. *Experimental Neurology* 211, 203–213.
- Steenkamp, V., Gouws, M. C., 2006. Cytotoxicity of Six South African medicinal plant extracts used in the traditional of Cancer. *South African Journal of Botany* 72(4), 630-633.
- Steenkamp, V., Gouws, M. C., Gulumian, M., Elgorashi, E. E., Van Staden, J., 2006. Studies on antibacterial, antiinflammatory and antioxidant activity of herbal remedies used in the treatment of benign prostatic and prostatitis. *Journal of Ethnopharmacology* 103(1), 71-75.
- Steenkamp, V., Mathivha, E., Gouws, M. C., Van Rensburg, C. E. J., 2004. Studies on antibacterial, antioxidant and fibroblast growth stimulation of wound healing remedies from South Africa. *Journal of Ethnopharmacology* 95(2-3), 353-357.
- Stehbens, W. E., 2004. Oxidative stress in viral hepatitis and AIDS. *Experimental and Molecular Pathology* 121-132.
- Sticher, O., 2008: Natural product isolation. *Natural product report* 25, 517-554.
- Stoddart, B., Wilcox, M. H., 2002. *Clostridium difficile*. *Current Opinion in Infectious Diseases* 15, 513–518.
- Stojiljkovic, V., Todorovic, A., Pejic, S., Kasapovic, J., Saicic, Z. S., Radlovic, N., Pajovic, S., 2009. Antioxidant status and lipid peroxidation in small intestinal mucosa of children with celiac disease. *Clinical Biochemistry* 42, 1431–1437
- Suleiman, M. M., McGaw, L. J., Naidoo, V., Eloff, J. N., 2010. Evaluation of several tree species for activity against the animal fungal pathogen *Aspergillus fumigatus*. *South African Journal of Botany* 76, 64-71.
- Sun, J., Hu, X.-L., Le, G.-W., Shi, Y.-L., 2010. Lactobacilli prevent hydroxy radical production and inhibit *Escherichia coli* and *Enterococcus* growth in system mimicking colon fermentation. *Letters of Applied Microbiology* 50, 264-269.
- Suzuki, T., Hara, H., 2011. Role of flavonoids in intestinal tight junction regulation. *Journal of Nutritional Biochemistry* 22, 401–408.
- Svenningsen, A. B., Madsen, K. D., Liljefors, T., Stafford, G. I., Staden, J., Jäger, A. K., 2006. Biflavones from *Rhus* species with affinity for the GABA(A)/benzodiazepine receptor. *Journal of Ethnopharmacology* 103, 276–280.
- Taganna, J. C., Quanico, J. P., Perono, R. M. G., Amor, E. C., Rivera, W. L., 2011. Tannin-rich fraction from *Terminalia catappa* inhibits quorum sensing (QS) in *Chromobacterium violaceum* and the QS-controlled biofilm maturation and LasA staphylolytic activity in *Pseudomonas aeruginosa*. *Journal of Ethnopharmacology* 134, 865–871
- Taguri, T., Tanaka, T., Kouno, I., 2004. Antimicrobial activity of 10 different plant polyphenols against bacteria causing food-borne disease. *Biol. Pharm. Bull.* 27(12) 1965-1969.

- Taiwo, O. B., Olajide, O. A., Soyannwo, O. O., Makinde, J. M., 2000. Antiinflammatory, antipyretic and antispasmodic properties of *Chromomaela odorata*. *Pharmaceutical Biology* 38(5), 367-370.
- Takeuchi K, Miyazawa T, Tanaka A, Kato S, Kunikata T., 2002. Pathogenic importance of intestinal hypermotility in NSAID-induced small intestinal damage in rats. *Digestion* 66, 30–41.
- Takeuchi, K., Tanaka, A., Kato, S., Amagase, K., Satoh, H., 2010. Roles of COX inhibition in pathogenesis of NSAID-induced small intestinal damage. *Clinica Chimica Acta* 411, 459–466.
- Takeuchi, T., Fujinami, K., Goto, H., Fujita, A., Taketo, M.M., Manabe, T., Matsui, M., Hata, F., 2005. Roles of M2 and M4 muscarinic receptors in regulating acetylcholine release from myenteric neurons of mouse ileum. *Journal of Neurophysiology* 93, 2841–2848.
- Taraporewala, I. B., Kauffman, J. M., 1990. Synthesis and structure-activity relation-ship of anti-inflammatory 9,10-dihydro 9-oxo-2-acridine alkanolic acids and 4-(2-carboxyphenyl) aminobenzenealkanoic acids. *Journal of Pharmacology Science* 79, 173–178.
- Taylor, J. L. S., van Staden, J., 2001. COX-1 inhibitory activity in extracts from *Eucomis* L'Herit. species. *Journal of Ethnopharmacology* 75, 257–265.
- Teke, G. N., Kuate, J. R., Ngouateu, O. B., Gatsing, D., 2007. Antidiarrhoeal and antimicrobial activities of *Emilia coccinea* (Sims) G. Don extracts. *Journal of Ethnopharmacology* 112, 278-283.
- Thamburan, S., Klaasen, J., Mabusela, W. T., Cannon, J. F., Folk, W., Johnson, Q., 2006. *Tulbaghia alliacea* Phytotherapy: A Potential Anti-infective Remedy for Candidiasis. *Phytotherapy Research* 20, 844–850
- Thapar, N., Sanderson, I. R., 2004. Diarrhoea in children: an interface between developing and developed countries. *The Lancet* 363, 641–53.
- Thiagarajah, J. R., Verkman, A. S., 2005. New drug targets for cholera therapy. *TRENDS in Pharmacological Sciences* 26(4), 172-174.
- Thring, T. S. A., Springfield, E. P., Weitz, F. M., 2007. Antimicrobial activities of four plant species from the Southern Overberg region of South Africa. *African Journal of Biotechnology* 6(15), 1779-1784.
- Thring, T. S. A., Weitz, F. M., 2006. Medicinal plant use in the Bredasdorp/Elim region of the Southern Overberg in the Western Cape Province of South Africa. *Journal of Ethnopharmacology* 103, 261–275.
- Tobin, G., Giglio, D., Lundgren, O., 2009. Muscarinic receptor subtypes in the alimentary tract. *Journal of Physiology and Pharmacology* 60 (1), 3–21.
- Todd, E. C. D., Notermans, S., 2011. Surveillance of listeriosis and its causative pathogen, *Listeria monocytogenes*. *Food Control* 22, 1484-1490.
- Tona, L., Cimanga, R. K., Mesia, K., Musuamba, C.T., De Bruyne, T., Apers, S., Hernans, N., Van Miert, S., Pieters, L., Totte, J., Vlietinck, A.J., 2000. Antiamoebic and spasmolytic activities of extracts from some antidiarrhoeal traditional preparations used in Kinshasa, Congo. *Phytomedicine* 2000 7(1), 31-8.
- Tona, L., Kambu, K., Mesia, K., Cimanga, K., De Bruyne, T., Pieters, L., Totté, J., Vlietinck, A.J., 1999. Biological screening of traditional preparations from some medicinal plants used as antidiarrhoeal in Kinshasa, Congo. *Phytomedicine* 6, 59–66.
- Tona, L., Kambu, K., Ngimbi, N., Cimanga, R. K., Vlietinck, A. J., 1998. Antiamoebic and phytochemical screening of some Congolese medicinal plants. *Journal of Ethnopharmacology* 61(1), 57-65.
- Troeger, H., Epple, H. J., Schneider, T., Wahnschaffe, U., Ullrich, R., Burchard, G. D., *et al.*, 2007. Effect of chronic *Giardia lamblia* infection on epithelial transport and barrier function in human duodenum. *Gut* 56(3), 328-335.

- Troeger, H., Loddenkemper, C., Schneider, T., Schreier, E., Epple, H. J., Zeitz, M., et al., 2009. Structural and functional changes of the duodenum in human norovirus infection. *Gut* 58, 1070-1077.
- Tshibangu, J. N., Wright, A. D., Konig, G. M., 2003. HPLC isolation of the anti-plasmodially active bisbenzylisoquinone alkaloids present in roots of *Cissampelos mucronata*. *Phytochemical Analysis* 14(1),13-22.
- Tshikalange, T. E., Hussein, A. A., 2010. Cytotoxicity activity of compounds from *Elaeodendron transvaalense* ethanol extract. *Journal of Medicinal Plant Research* 4(16), 1695-1697.
- Tshikalange, T. E., Meyer, J. J. M., Hussein, A. A., 2005. Antimicrobial activity, toxicity and the isolation of a bioactive compound from plants used to treat sexually transmitted diseases. *Journal of Ethnopharmacology* 96(3):), 515-519.
- Turel, I., Ozbek, H., Erten, R., Oner, A. C., Cengiz, N., Yilmaz, O., 2009. Hepatoprotective and anti-inflammatory activities of *Plantago major* L. *Indian Journal of Pharmacology* 41(3), 120-124.
- Uchida, K., 1999. Current status of acrolein as a lipid peroxidation product. *Trends Cardiovascular Medicine* 9, 109–113.
- UNAIDS, 2006. Report on the Global AIDS Epidemic, May 2006.
- Unno, T., Matsuyama, H., Sakamoto, T., Uchiyama, M., Izumi, Y., Okamoto, H., Yamada, M., Wess, J., Komori, S., 2005. M2 and M3 muscarinic receptor-mediated contractions in longitudinal smooth muscle of the ileum studied with receptor knockout mice. *British Journal of Pharmacology* 146, 98–108.
- Valeur, J., Lappalainen J., Rita, H., Lin, A. H., Kovanen, P. T., Berstad, A., Eklund, K., Vaali K., 2009. Food allergy alters jejuna circular muscle contractility and induces local inflammatory cytokines expression in a mouse model. *BMC gastroenterology* 9:33.
- Valko, M., Leibfritz, D., Moncol, J., Cronin, M. T. D., Mazur, M., Telser, J., 2007. Free radical and antioxidants in normal physiological functions and human diseases. *The International Journal of Biochemistry and Cell Biology* 39, 44-84.
- Valko, M., Morris, H., & Cronin, M. T. D., (2005). Metals, toxicity and oxidative stress. *Current Medicinal Chemistry* 12, 1161–1208.
- Van Damme, I. Habib, I., De Zutter, L., 2010. “*Yersinia enterocolitica* in slaughter pig tonsils: enumeration and detection by enrichment versus direct plating culture,” *Food Microbiology*, 27(1), 158–161.
- Van der Merwe, D., Swan, G. E., Botha, C. J., 2001. Use of ethnoveterinary medicinal plants in cattle by Setswana-speaking people in the Madikwe area of the North West Province of South Africa. *Journal of the South African Veterinary Association* 72, 189-196.
- Van der Watt, E., Pretorius, J. C., 2001. Purification and identification of active antibacterial components in *Carpobrotus edulis*. L. *Journal of Ethnopharmacology* 76, 87-91.
- Van Dyk, S., Griffitto, S., Van Zyl, R. L., Malan, S. F., 2009. The importance of including toxicity assays when screening plant extracts for antimalarial activity. *African Journal of Biotechnology* 8(20), 5595-5601.
- Van Wyk, B., Van Wyk, P., 1997. *Field Guide to Trees of Southern Africa*. Struik Publishers (A Division of New Holland Publishing, South Africa), Cape Town, South Africa, p. 496.
- Van Wyk, B., Wink, M., 2004. Health disorders and Medicinal plants. In: *Medicinal plants of the world*. Published by Briza Publications CK90/11690/23, P O Box 56569, Arcadia 0007, Pretoria, South Africa Pp 351-370.
- Van Wyk, B.-E., 2008a. A review of Khoi-San and Cape Dutch medical ethnobotany. *Journal of Ethnopharmacology* 119, 331–341.
- Van Wyk, B.-E., 2008b. A broad review of commercially important Southern African medicinal plants. *Journal of Ethnopharmacology* 119, 342–355.

- Van Wyk, B.-E., de Wet, H., Van Heerden, F. R., 2008. An ethnobotanical survey of medicinal plants in the Southeastern Karoo, South Africa. *South African Journal of Botany* 74, 696–704.
- Van Wyk, B.-E., Gericke, N., 2000. *People's Plants: A Guide to Useful Plants of Southern Africa*. Briza Publications, Pretoria, ISBN: 978 1 875093 19 9
- Velazquez, C., Calzada, F., Esquivel, B., Barbosa, E., Calzada, S., 2009. Antisecretory activity from the flower of *Chiranthodendron pentadactylon* and its flavonoids in intestinal fluid accumulation induced by *Vibrio cholerae* toxin in rats. *Journal of Ethnopharmacology*, 126, 455-458.
- Velázquez, C., Calzada, F., Torres, J., González, F., Ceballos, G., 2006. Antisecretory activity of plants used to treat gastrointestinal disorders in Mexico. *Journal of Ethnopharmacology* 103, 66–70.
- Venter, F., Venter, J.-A., 2002. *Making the Most of Indigenous Trees*, rev. edition. Briza Publications, Pretoria, South Africa, p. 28
- Vermerris, W., and Nicholson, R., 2006. *Phenolic Compounds Biochemistry*, Publisher Springer, New York. Pp 151-196.
- Verschaeve, L., Van Staden, J. 2008. Mutagenic and antimutagenic properties of extracts from African South traditional medicinal plants. *Journal of Ethnopharmacology* 119(3), 575-587.
- Vital, P. G., Rivera, W. L., 2009. Antimicrobial activity and cytotoxicity of *Chromolaena odorata* (L. F.) King and *Uncaria perrottetii* (A. Rich) Merr. Extracts. *Journal of Medicinal plant Research* 3(7), 511-518.
- Vitali, F., Fonte, G., Saija, A., Tita, B., 2006. Inhibition of intestinal motility and secretion by extracts of *Epilobium* spp. in mice. *Journal of Ethnopharmacology* 107, 342-348.
- Viviers, P., Kolodziej, H., Young, D., Ferreira, D., Roux, D., 1983. *Journal of Chemical Society, Perkin Trans 1*, 2555.
- Voirin, B.; Jay, M.; Hauteville, M., 1975. Isoetine, nouvelle flavone isolee de *Isoetes delilei* et de *Isoetes durieui*. *Phytochemistry* 14, 257-259.
- Von Koenen, E., 2001. *Medicinal Poisonous and Edible Plants in Namibia*. Klaus Hess Publishers, Windhoek and Göttingen, ISBN: Namibia 99916-747-4-8; Germany 3-9804518-7-9.
- Voravuthikunchai, S. P., and Limsuwan, S., 2006. Medicinal plant Extracts as anti-*Escherichia coli* 0157:H7 agent and their effects on Bacterial Cell aggregation. *Journal of Food Protection* 69(10), 2336-2341.
- Vundac, V. B., Brantner, A. H., Plazibat, M., 2007. Content of Polyphenolic constituent and antioxidant activity of some stachys taxa. *Food Chemistry* 104, 1277-1281.
- Wafo, P., Kamdeen, R. S. T., Ali, Z., Anjuin, S., Begum, A., Oluyemisi, O. O., Khan, S. N., Ngadjui, B. T., Etoa, X. F., Choudhany, M. I., Kaurene-type diterpenoids from *Chromolaena odorata*, their X-ray diffraction studies and potent α -glycosidase inhibition of 16-kauren-19-oic acid.
- Wang, G. J., Chen, Y. M., Wang, T. M., Lee, C. K., Chen, K. J., Lee, T. H., 2008. Flavonoids with iNOS inhibitory activity from *Pogonatherum crinitum*. *Journal of Ethnopharmacology* 118, 71–78.
- Wang, M., Takeda, K., Shiraishi, Y., Okamoto, M., Dakhama, A., Joetham, A., Gelfrand, E. W., 2010. Peanut-induced intestinal allergy is mediated through a mast cell-IgE-Fc ϵ RI-IL-13 pathway. *Journal of Allergy and Clinical Immunology* 126(2), 301-316.
- Wanjala, C.C.W., Juma, B.F., Bojase, G., Gashe, B.A., Majinda, R.R.T., 2002. Erythraline alkaloids and antimicrobial flavonoids from *Erythrina latissima*. *Planta Medica* 68, 640–642
- Watt, J. M., Breyer-Brandwijk, M. G., 1962. *The Medicinal and Poisonous Plants of Southern and Eastern Africa*, second edition, Livingstone, London.

- Watt, J. M., Breyer-Brandwijk, M. G., 1996. *The Medicinal and Poisonous Plants of Southern and Eastern Africa*, 2nd ed. E. & S. Livingstone Ltd., London.
- Weigenand, O., Hussein, A. A., Lall, N., Meyer, J. J. M., 2004. Antibacterial activities of Naphthoquinones and Triterpenoids from *Euclea natalensis*. *Journal of Natural Product* 67, 1936-1938.
- Wernes, S. W., and Lucchesi, R., 1990. Free radicals and ischemic tissue injury. *Trends in Pharmacological Sciences* 11, 161-166.
- Wettasinghe, M., Shahidi, F., Amarowicz, R., Abou-Zaid, M. M., 2001. Phenolic acids in defatted seeds of borage (*Borago officinalis* L.). *Food Chemistry* 75, 49-56.
- Wistrom, S. R., Norrby, E. B., Myhre, E. B., *et al.*, 2001. Frequency of antibiotic-associated diarrhoea in 2462 antibiotic treated hospitalized patients: a prospective study. *Journal of Antimicrobial Chemotherapy* 47, 43-50.
- Wood, J. D., 2004. Enteric Neuroimmunophysiology and pathophysiology. *Gastroenterology* 127, 635-657.
- World Health Organization (WHO) and United Nation Children Fund (UNICEF), 2004. *Clinical World Health Organization (WHO) and United Nation Children Fund (UNICEF), 2004: Clinical Management of Acute Diarrhea. WHO/FCH/CAH/04.7 World Health Organization, Geneva. Management of Acute Diarrhea. WHO/FCH/CAH/04.7 World Health Organization, Geneva.*
- Wouters, M. M., Farrugia, G., Schemann, M., 2007. 5-HT receptors on interstitial cells of Cajal, smooth muscle and enteric nerves. *Neurogastroenterology Motility* 19 (Suppl 2), 5-12.
- Wu, S. W., Dornbusch, K., Kronval, G., 1999. Genetic characterization of resistance to extended-spectrum betalactams in *Klebsiella oxytoca* isolates recovered from patients with septicemia at hospitals in the Stockholm area. *Antimicrobial Agents and Chemotherapy* 43, 1294-1297.
- Wynn, S. G., Fougere, B. J., 2007. A systems-based approach: Diarrhea-Therapeutic rationale. In: *Veterinary Herbal medicine*. (Edited by Wynn GS and Fougere BJ) Mosby Elsevier 11830 Westline Industrial Drive, St Louis, Missouri 63146: 33336-337.
- Xu, Y., Oliverson, B. G., Simmons, D. L., 2007. Trifunctional inhibition of COX-2 by extracts of *Lonicera japonica*: Direct inhibition, transcriptional and post-transcriptional down regulation. *Journal of Ethnopharmacology* 111:667-670.
- Xue, C., Tada, Y., Dong, X., Heitman, J., 2007. The human fungal pathogen *Cryptococcus* can complete its sexual cycle during a pathogenic association with plants. *Cell Host and Microbes* 1, 2633-273.
- Yadav, A. K., Tangpu, V., 2009. Therapeutic efficacy of *Biden pilosa* L. Var. *radiata* and *Galinsoga parviflora* Cav. in experimentally induced diarrhoea in mice. *Phytopharmacology and Therapeutic values* V, 35-45.
- Yakubu, M.T., Afolayan, A.J., 2009. Reproductive toxicologic evaluations of *Bulbine natalensis* baker stem extract in albino rats. *Theriogenology* 72, 322-332.
- Yamagiwa, Y., Ohashi, K., Sakamoto, Y., Hirakawa, S., Kamikawa, T., 1987. Syntheses of anacardic acids and ginkgoic acid. *Tetrahedron* 43(15), 3387.
- Yarnell, E., 2007. Plant chemistry in Veterinary medicine: Medicinal constituents and their mechanism of action. In: *Veterinary Herbal medicine*. (Edited by Wynn GS and Fougere BJ) Mosby Elsevier 11830 Westline Industrial Drive, St Louis, Missouri 63146: 167-168.
- Yff, B. T. S., Lindsey, K. L., Taylor, M. B., Erasmus, D. G., Jager, A. K., 2002. The pharmacological screening of *Pentanisia pruelloides* and the isolation of the antibacterial compound palmitic acid. *Journal of Ethnopharmacology* 79(1), 101-107.

- Yokozawa, T., Dong, E., Chung, H. Y., Oura, H., Nakagawa, H., 1997. Inhibitory effect of green tea on injury to a cultured renal epithelial cell line, LLC-PK₁. *Bioscience Biotechnology and Biochemistry* 61, 204-206.
- Zschocke, S., Van Staden, J., Paulus, K., Bauer, R., Horn, M. M., Munro, O. Q., Brown, N. J., Drewes, S. E., 2000. Stereostructure and anti-inflammatory activity of three diastereomers of Ocobullenone from *Ocotea bullata*. *Phytochemistry* 54, 591-595.
- Zwenger, S., Basu, C., 2008. Plant terpenoids: applications and future potentials. *Biotechnology and Molecular Biology Reviews* 3, 001-007.

Appendix 2.1: ETHNOBOTANICAL AND LITERATURE INFORMATION OF MEDICINAL PLANT SPECIES USED TRADITIONALLY FOR TREATING DIARRHOEA IN SOUTH AFRICA

Family/Plant species	Local names	Part used	Ethnopharmacological information	Biological activities investigated	Bioactive compound(s) isolated
Aizoaceae					
<i>Carpobrotus acinaciformis</i> (L.) L. Bolus	perdevy	Leaf juice	Sore throat, dysentery, mouth infection (van Wyk, 2008)	Antibacterial (Oskey et al., 2009)	2-descarboxy-betanidin (Dembitsky, 2005)
<i>Carpobrotus edulis</i> (L.) L. Bolus	Ikhambi-lamabulawo. Umgongozi	leaves	Diarrhoea, digestive problems, allergy (Thring and Weitz, 2006); dysentery (van Wyk, 2008)	Antibacterial (Van der Watt et al., 2001)	Rutin, hyperoside, neohesperidin, catechin, ferulic acid (van der Watt et al., 2001)
<i>Carpobrotus muiirii</i> (L.) L. Bolus		Leaves	Dysentery, digestive problem, mouth ulcers, thrush (Thring and Weitz, 2006)	Antimicrobial (Springfield et al., 2003)	-
Alliaceae					
<i>Agapanthus praecox</i> Willd.	uMkhondo (X)	Roots	Diarrhoea in sheep and goat (Dold and Cocks, 2001, McGaw and Eloff., 2008)	-	-
<i>Tulbaghia alliacea</i> L.f.	Umwelala X, ivimba-mpunzi X, Sikwa Z	bulb	Stomach ache, fever, tuberculosis, influenza (Bisi-Johnson et al., 2010)	Antimycobacterial (Bamuamba et al., 2008), Mutagenicity and antimutagenicity (Reid et al., 2006), anticandidiasis (Thamburan et al., 2006)	-
Amaranthaceae					
<i>Guilleminea densa</i> Moq	Sepatho (S)	Root	Decoction for diarrhoea (Mathabe et al., 2006)	-	-
<i>Hermbstaedia odorata</i> Wild	Ubuphuphu (X, Z)	leaves	Food and infusion for diarrhoea (Bisi-Johnson et al., 2010); Root cleansing stomach wash alone or with <i>Acaccia xanthophloea</i> and (Hutchings et al., 1996).	-	-
Amaryllidaceae					
<i>Scadoxus puniceus</i> (L.) Friis and Nordal	Umpompho-wezinja, Isiphompho umgola Z	Bulb and root	Stomach ache, diarrhoea, nausea (Bisi-Johnson et al., 2010)	Antimicrobial, anti-inflammatory, acetylcholinesterase inhibition and mutagenic activities (Ndhala et al., 2010)	-
Anacardiaceae					
<i>Mangifera indica</i> L.	Umango	Leaves, bark	Diarrhoea (de Wet et al., 2010)	Antidiarrhoeal (Sairam et al., 2003), antidiabetic (Aderibigbe et al., 2001)	Gallotannins (Engels et al., 2010), mangiferrin (Singh et al., 2009).
<i>Ozoroa insignis</i> Delile	Monoko	Stem bark	Decoction for diarrhoea (Mathabe et al., 2006), vinearal diseases, parasites, kidney trouble (Liu and Abreu, 2006)	Antibacterial (Mathabe et al., 2006); antigardial (Johns et al., 1995), antimalarial (Asase et al., 2005), Cytotoxicity (Rea et al., 2003), antischistosomiasis (Molgaard et al., 2001; Ndamba et al., 1994).	6-pentadecylsalicylic acid (antifouling), tirucallane triterpenes (Liu and Abreu, 2006)

<i>Ozoroa mucronata</i> (Bernh.ex C.Krauss) R.fern & A. Fern		root	Diarrhoea, intestinal parasites and stomach trouble (Yamagiwa <i>et al.</i> , 1987)	LOX inhibition, PG synthase inhibition (Kubo <i>et al.</i> , 1987)	Anarcadic acid (LOX inhibition) (Ha and Kubo, 2005), Moronic acid (Hotestmann Kaldas and Nakanishi, 1979)
<i>Ozoroa paniculosa</i> (Sond.) R. & A. Fernandes	Mubandulakhali, Mudumbula (V)	Bark, root bark	Abdominal problems in animal (Hutchings <i>et al.</i> , 1996), Diarrhoea, sweating sickness (Van der Merwe <i>et al.</i> , 2001; McGaw <i>et al.</i> , 2008)	Antioxidant (Motlhanka, 2008), antimicrobial and antimycobacterial	-
<i>Ozoroa schaeerocarpa</i> R. Fern & A. Fern	Mudumbula (V)	Bark	Infusion for diarrhoea (Sibandze <i>et al.</i> , 2010)	Antiescherichial (Sibandze <i>et al.</i> , 2010)	-
<i>Protorhus longifolia</i> (Bernh.ex C. Krauss) Engl.	i(u)Zntlwa, ikubalo, umkupati X	Bark	Heartwater and diarrhoea in cows (Dold and Cocks, 2001, McGaw and Eloff, 2008); Heart burn and stomach bleeding (Hutchings <i>et al.</i> , 1996)	Antimicrobial (Suleiman <i>et al.</i> , 2010)	-
<i>Sclerocarya birrea</i> (A. Rich.) Hochst. subsp. <i>caffra</i> (Sond.)	Mufula (V)	Leaves, bark, roots	Diarrhoea and fractures (Van der Merwe <i>et al.</i> , 2001; McGaw <i>et al.</i> , 2008)	Mutagenicity, antimutagenicity (Elgorashi <i>et al.</i> , 2003), Antibacterial, antihelmintic and cytotoxicity (McGaw <i>et al.</i> , 2007), anti-diarrhoea (Galvez <i>et al.</i> , 1991), antibacterial (Eloff, 2001), anti-inflammation (Ojewole, 2010), antioxidant (Braca <i>et al.</i> , 2003), anti-diabetic (Ojewole, 2004)	Gallotannin, tannic, mallic, gallic and citric acid, triterpene, flavonoid, coumarins (Ojewole <i>et al.</i> , 2010)
<i>Searsia gueinzii</i> Sond (Syn <i>Rhus gueinii</i> Sond)	Mushakaladza (V)	root	Gastrointestinal infections (Elgorashi <i>et al.</i> , 2003)	Mutagenicity, antimutagenicity (Elgorashi <i>et al.</i> , 2003)	-
<i>Searsia incisa</i> L.f.	uNongquthu	Root and bark decoction	Shock and diarrhoea (Dold and Cocks, 2001, McGaw <i>et al.</i> , 2008)		-
<i>Searsia lancea</i> L.f.	Mushakaladza (V)		Diarrhoea and gallsickness (Van der Merwe <i>et al.</i> , 2001; McGaw <i>et al.</i> , 2008)	Antibacterial, antihelmintic and cytotoxicity (McGaw <i>et al.</i> , 2007)	-
<i>Searsia leptodictya</i> Diels	Mushakaladza (V)	leaves	Brower, gall sickness in cattle, infectious disease, chest and abdominal pain (Sebothoma, 2010)	Antimicrobial (Sebothoma, 2010)	(-)-leucofisetinidin, (-)-epicatechin and [4,8]-(+)-fisetinidol(-)-epicatechin (Viviers <i>et al.</i> , 1983)
<i>Searsia pendulina</i> Jacq.	-	Leaves	Stomach ailment, enema in children (Coates-Palgrave, 2002)	-	-
<i>Searsia pentheri</i> Zahlbr.	Muthasiri (V)	leaves	Epilepsy (Svenningsen <i>et al.</i> , 2006)	GABA _A /benzodiazepine receptor affinity (Svenningsen <i>et al.</i> , 2006)	Apeginin, agathisflavone (Svenningsen <i>et al.</i> , 2006)
<i>Searsia rogersii</i> Schonland	Muthasiri (V)	Bark	Pain, watery diarrhoea (Samie <i>et al.</i> , 2010)	Antifungal (Samie <i>et al.</i> , 2010), Antimycobacterium (Green <i>et al.</i> , 2010)	-
Annonaceae					
<i>Annona</i>	Muembe (V)	Bark	Toothache, venereal, diarrhoea (Mabogo,	Antidiarrhoeal (Suleiman <i>et al.</i> ,	Annosenegalina (cytotoxic and antiparasitic), Annonacin

<i>senegalensis</i> Pers.			1990; More <i>et al.</i> , 2008)	2008), antivenom (Adzu <i>et al.</i> , 2005), antimalaria (Okokon <i>et al.</i> , 2006)	(cytotoxic agent, insecticidal, mutagenic activity) immunosuppressant), senegalene (cytotoxic agent), 17, 19-kauranediol (ent-16β)-form . Dicarboxylic acid, 19-Methyl ester (toxic to brime shrimp)
<i>Uvaria chamae</i> P. Beauv		Root	Catarrh, dysentery, fever, hematemesis, inflammation, jaundice, wounds, yellow fever (Reid <i>et al.</i> , 2006)	Antimalaria (Okokon <i>et al.</i> , 2006); mutagenic and antimutagenic (Reid <i>et al.</i> , 2006)	-
Apiaceae					
<i>Alepidea amatymbica</i> Eckl. & Zeyh.	Iqwili, Ikhathazo (Z)	Root	Decoction for diarrhoea (Appidi <i>et al.</i> , 2008)	Antimicrobial, anti-inflammatory and genotoxicity (Mulauzi <i>et al.</i> , 2009)	Rosmarinic acid, Dehydrokaurenoic acid, Kaurenoic acid, kaurene lactone, acetoxyl kaurene lactone (Holzapfel <i>et al.</i> , 1995)
<i>Centella asiatica</i> (L.) Urb.		Root	Chronic diarrhoea and dysentery; diaphoretic (van Wyk, 2008)	Modulator of nitric oxide production and TNF- α (Nhlem <i>et al.</i> , 2011), , lipid peroxidation (Kumar and Muller, 1999)	Asiaticoside G, asiaticoside, asiaticoside F, asiatic acid, quadranoside IV, 2a,3b,6b-trihydroxyolean-12-en-28-oic acid 28-O-[α -L-rhamnopyranosyl-(1 \rightarrow 4)- β -D-glucopyranosyl-(1 \rightarrow 6)- β -D-glucopyranosyl] ester, kaempferol, quercetin, astragalol, and isoquercetin (Nhlem <i>et al.</i> , 2011)
<i>Centella glabrata</i> L.		Root and stalk	Chronic diarrhoea and dysentery, diaphoretic (van Wyk, 2008b)	-	-
<i>Foeniculum vulgare</i> Mill.		Leaf	Flatulence, cough, diuretic, digestive problem, diarrhoea, stomach ache and cramps (Watt and Breeyer-Brandwijk, 1962; van Wyk, 1997)	Antimicrobial (Bacillus cereus, Clostridium botulinum, Salmonella enteritidis, Staphylococcus aureus, Yersinia enterocolitica) (Ceylan and Fung, 2004)	Falcarindiol (antifungal, antibiotic and analgesic, antinociceptive, DNA topoisomerase inhibitor, phytotoxic, allelochemical, antimutagenic and antiproliferative agents), 1-(4-hydroxyphenyl)-1,2-propanediol form, 4' methyl ether (phytotoxin, antiparasitic, nematocidal agent)
Apocynaceae					
<i>Acokanthera oblongifolia</i> (Hochst.) Codd	inHlungunyembe Intlungunyembe (X, Z)	leaves	Severe gastrointestinal irritation (Verschaeve and Van Staden, 2008), Decoction for stomach ache, diarrhoea (Bisi-Johnson <i>et al.</i> , 2010)	Genotoxicity (Elgorashi <i>et al.</i> , 2003); Epilepsy and convulsion (Risa <i>et al.</i> , 2004)	Acolongifloroside K and H (antineoplastic agent)
<i>Catharanthus roseus</i> (L.) G.Don	Imbali, Ikhwinini, Isishushlungu (Z)	Leaves, stem and root	Diarrhoea (de Wet <i>et al.</i> , 2010)	Antimicrobial (van Vuuren and Naidoo, 2010)	Serpentine (antitumour activity); apparicine (cytotoxin, weak antibacterial, antiviral agent active against Polio virus, analeptic properties); β-carboline (induced mutagenicity, antiparasitic, antitrypanosomal agent); Catharanthamine (antitumour); Trichosetin (antibacteria); 16-Epi-2-isositsirikine antineoplastic); Leurosine (antihyperglycaemic); Lochnerinine (antitumour); Pericyclivine (weak cytotoxic activity); 15',20'-anhydroviriblastine (antineoplastic agent); Vindoline (antineoplastic); Vindolinine (antiglycaemic agent, antifungi); Vingamine (cytotoxic); yohimbine (selective α_2 – adrenoceptor antagonist, antidepressant, antihypotensive, , antidiuretic activity, aphrodisiac, anxiogenic activity in rodent)
<i>Sarcostemma viminale</i>	Umbelebele, Ingotshwa	Stem	Infusion for diarrhoea (de Wet <i>et al.</i> , 2010), Increase livestock productivity	Antibacterial, anti-inflammatory and mutagenic effects (Luseba <i>et al.</i> ,	-

(L) R. Br subsp. <i>viminale</i>			(Kunene and Fossey, 2006)	2007)	
Aquifoliaceae					
<i>Ilex mitis</i> (L.) Radlk.	Monamane (S), Mutanzwa-khamelo (V)	Root bark	Decoction for diarrhoea (Mathabe <i>et al.</i> , 2006)	Antimalaria and cytotoxicity (Rasoanaivo <i>et al.</i> , 2004)	-
Araliaceae					
<i>Cussonia arborea</i> Hochst ex A. Rich		Root, leaves	Decoction for diarrhoea (De Villiers <i>et al.</i> , 2010)	antimicrobial and antimalarial (De Villiers <i>et al.</i> , 2010)	Arboreaside A, Arboreaside B, Arboreaside C, Arboreaside D, Arboreaside E, ciwujianoside C3 and 23-hydroxyursolic acid 28-O- α -L-rhamnopyranosyl-(1 \rightarrow 4)- β -D-glucopyranosyl-(1 β)- β -D-glucopyranosyl ester (Kougan <i>et al.</i> , 2009)
Asclepiadaceae					
<i>Asclepias fruticosa</i> L.	iGwada (X), Mutshulwa (V), Lebegana (S)	Root, leaves	Diarrhoea and stomach pain in children (Lewu and Afolayan, 2009)	Antimycobacterium (Green <i>et al.</i> , 2010), antifungal (Samie <i>et al.</i> , 2010), antimicrobial, anti-inflammatory, anticholinesterase and mutagenic activities (Ndhala <i>et al.</i> , 2010)	-
<i>Secamone filiformis</i> (L.f) J. H. Ross	iMbijela	Stem	Diarrhoea in cattle (Dold and Cocks, 2001, McGaw <i>et al.</i> , 2008)	Anthelmintic, antibacterial and cytotoxicity (McGaw <i>et al.</i> , 2007)	-
<i>Xysmalobium undulatum</i> (L.) W.T. Aiton	Ishongwe (X, Z)	Roots	Diarrhoea, dysentery, stomach cramps, headache, oedema, dysmenorrhoea (Bisi-Johnson <i>et al.</i> , 2010)	Antibacterial (Rabe and Van Staden, 1997), PG inhibition (Jager <i>et al.</i> , 1996), Serotonin re-uptake modulatory activity (Nielsen <i>et al.</i> , 2004), antidepressant (Pedersen <i>et al.</i> , 2008)	-
Asparagaceae					
<i>Asparagus cooperi</i> Bak.	Lefatshana (S)	Whole plant	Decoction for diarrhoea (Mathabe <i>et al.</i> , 2006)	Antibacterial (Mathabe <i>et al.</i> , 2006)	-
Asphodelaceae					
<i>Aloe arborescens</i> Miller	Inhlaba, Tshikhopho (V)	leaves	Diarrhoea and sore (Mlambo, 2008)	Anti-inflammatory (Lindsey <i>et al.</i> , 2002); immunomodulator, anti-inflammatory (Imanishi, 1993),	Aloctin A (Imanishi, 1993); aloenin, 2'-O- <i>p</i> -coumaroylaloenin, 2'-O-feruloylaloenin, isobarbaloin, and barbaloin (Beppu <i>et al.</i> , 2003)
<i>Aloe candelabrum</i> Berger	Ikhala Inkalane (X) Uphondonde (Z)	leaves	Decoction for diarrhoea (Bisi-Johnson <i>et al.</i> , 2010)	-	-
<i>Aloe greatheadii</i> Schonl.	Sekgopho (S)	Leaves	Decoction for diarrhoea (Mathabe <i>et al.</i> , 2006)	antioxidant (Botes <i>et al.</i> , 2008), antiplasmodial and cytotoxicity (Van Dyk <i>et al.</i> , 2009)	-
<i>Aloe marlothii</i> Berger	Bindamutsho, Tshikhopho (V)	Leaves	Gallsickness, parasites, diarrhoea, constipation, retain placenta, dystocia maggots (Van der Merwe <i>et al.</i> , 2001);	antimalaria (Pillay <i>et al.</i> , 2008), antibacterial, anthelmintic, anti-amoebic (McGaw <i>et al.</i> , 2000),	-

			McGaw <i>et al.</i> , 2008)	antitick (Spicket <i>et al.</i> , 2007)	
<i>Bulbine abyssinica</i> A. Rich	Utswelana Intelezi (X) Ibhucu (Z), Incelwane (X)	Leaves, tubers	Vomiting, diarrhoea, tuberculosis (Bisi- Johnson <i>et al.</i> , 2010)	Antileukemia, antiplasmodial, cytotoxicity (Bringmann <i>et al.</i> , 2002)	Chrysophanol, aloe-emodin, knipholone, isoknipholone, Bulbine-knipholone (Bringmann <i>et al.</i> , 2002)
<i>Bulbine asphodeloides</i> (L.) Willd		tuber and leaves	rashes, sores wounds, dysentery and diarrhoea (lwalewa <i>et al.</i> , 2007)	-	-
<i>Bulbine fructescens</i> Wild	Intelezi (X)	leaf, root and rhizome	Diarrhoea, burns, rashes, blisters, insect bites, cracked lips and mouth ulcers (Coopoosamy, 2011)	Antibacterial (Coopoosamy, 2011), antiplasmodial (Mutanyatta <i>et al.</i> , 2005)	knipholone, 4-O-demethylknipholone-4-β-D-glucopyranoside (Mutanyatta <i>et al.</i> , 2005)
<i>Bulbine latifolia</i> (L.f) Roem et Schult	Irooiwater	Root	Decoction for diarrhoea (Appidi <i>et al.</i> , 2008)	Antibacterial (Coopoosamy, 2011)	Knipholone (antiplasmodial activity, cytotoxic agent)
<i>Bulbine natalensis</i> (Bak. Cf. roowortel	Ibhucu (Z)	leaves	Decoction for diarrhoea (Mathabe <i>et al.</i> , 2006); vomiting, diarrhoea, convulsion, venereal diseases, diabetes and rheumatism (Pujol, 1990)	Sexuality behaviour (Yakubu and Afolayan, 2008), Toxicity (Afolayan and Yakubu 2008)	-
Asteraceae					
<i>Acanthospermum glabratum</i> (DC) Wild	Inamathela	Whole plant	Diarrhoea (de Wet <i>et al.</i> , 2010)	-	-
<i>Acanthospermum australe</i> (Loefl.) O. Kuntze	Umgwaqeni (Z)	Whole plant	Diarrhoea (Mlambo, 2008)	Antiherpesvirus and antipoliavirus (Rocha Martin <i>et al.</i> , 2010)	Acanthoaustralide, quercetin and chryso-splenol (Rocha Martin <i>et al.</i> , 2010)
<i>Artemisia absinthium</i> L.		leaves	Diarrhoea (Van Wyk <i>et al.</i> , 2008)	Antimycobacterium (Gautam <i>et al.</i> , 2007)	-
<i>Bidens bipinnata</i> L.	Uvelemampo ndweni uvelegoli	leaves	Infusion for diarrhoea (Bisi-Johnson <i>et al.</i> , 2010), haemorrhage, reduce cancer, flu, cold, fever (Pooley, 1998)	Antidiarrhoea (Atta and Mounair, 2005)	-
<i>Bidens pilosa</i> L.	iSanama, Mushidzhi (V)	Root or leaves, flowers	Stomach pain (Lewu and Afolayan, 2009); diarrhoea, inflammation, female infertility, excessive menstruation (Dold and Cocks, 2000)	Antidiarrhoeal (Yadav and Tangpu, 2009), amoebicidal (Moundipa <i>et al.</i> , 2005), immunomodulator (Chang <i>et al.</i> , 2007, Chiang <i>et al.</i> , 2007)	Centaurein, centaureidin, cytopiloyne (Chang <i>et al.</i> , 2007, Chiang <i>et al.</i> , 2007)
<i>Brachylaena ilicifolia</i> (Lam.) Phill. & Schweick	uMgqh	Leaves	Diarrhoea in lambs (Dold and Cocks, 2001; McGaw <i>et al.</i> , 2008)	-	-
<i>Brachylaena transvaalensis</i> E. Philips and Schweick	Iphahlalehlathi	Leaves and bark	Diarrhoea (de Wet <i>et al.</i> , 2010)	--	-
<i>Callilepis laureola</i> Hutch	Impila (Z)	Roots	Diarrhoea (Mlambo, 2008)	-	-
<i>Chromolaena</i>	Usandanezwe (Z)	Leaves	Diarrhoea (Mlambo, 2008)	Anti-inflammatory, antipyretic	15-angeloyloxy-16,17-epoxy-19-kauronic acid, 16-kauren-19-

<i>odorata</i> L.				antispasmodic (Taiwo <i>et al.</i> , 2000), antidiabetic (Wafo <i>et al.</i> , 2011), antimicrobial and cytotoxicity (Vital and Rivera, 2009)	oic acid, 6'-hydroxy-2',3',4,4'-tetramethoxychalcone, isosakuranetin, acacetin, and kaempferide (Wafo <i>et al.</i> 2011)
<i>Conyza scabrida</i> DC.		Herb	Cold, influenza, inflammation, diarrhoea, fever, diabetes, stomach affliction (Thring <i>et al.</i> , 2007)	Antimicrobial (Thring <i>et al.</i> , 2007)	-
<i>Dicoma anomala</i> Sond.	Umuna (Z), Inyongana (X)	Roots	Decoction for diarrhoea, stomach cramp and skin lesion (Shale <i>et al.</i> , 1999)	Antibacterial, antioxidant, fibroblast growth stimulant (Steenkamp <i>et al.</i> , 2004)	-
<i>Dicoma capensis</i> Less.		Herb	Bitter tonic and diuretic; kidney; bladder; back pain; nausea; influenza; colds; cancer; diarrhoea (van Wyk, 2008)	Cytotoxicity (Steenkamp and Gouws, 2006)	-
<i>Helichrysum adenocarpum</i> DC		Root decoction	Diarrhoea and vomiting in children (Lourens <i>et al.</i> , 2008)		-
<i>Helichrysum calophyllum</i> Klatt		Root	Hyperfunction of lower gastrointestinal tract (Lourens <i>et al.</i> , 2008)	-	-
<i>Helichrysum ecklonis</i> Sond		Root decoction	Diarrhoea in children (Lourens <i>et al.</i> , 2008)	-	-
<i>Helichrysum odoratissimum</i> (L.)	Imphepho (Z)	Whole plant	Diarrhoea (Mlambo, 2008)	Antimicrobial (Puyvelde <i>et al.</i> , 1989)	3,5-dihydroxy-6,7,8-trimethoxyflavone and 3-O-methylquercetin, helichrysetin (Puyvelde <i>et al.</i> , 1989)
<i>Pentzia incana</i> (Thunb.) Kuntze			Diarrhoea (Van Wyk <i>et al.</i> , 2008)	-	-
<i>Schkuhria pinnata</i> (Lam.) Thell.	Unsakansaka (Z)	Aerial parts	Pneumonia, diarrhoea, eye infections, heartwater (Van der Merwe <i>et al.</i> , 2001; McGaw <i>et al.</i> , 2008)	Antibacterial, anti-inflammatory mutagenicity (Luseba <i>et al.</i> , 2007)	-
<i>Senecio quinquelobus</i> DC.	Usinini (Z)	Leaves	Diarrhoea (Mlambo, 2008)	-	-
<i>Vernonia glaberrima</i> Welw		Leaves	Decoction for diarrhoea (De Villiers <i>et al.</i> , 2010)	Antibacterial and antimalaria (De Villiers <i>et al.</i> , 2010)	-
<i>Vernonia kotschyana</i> Sch. Bip. ex Walp. (<i>Baccharoides adoensis</i> var. <i>kotschyana</i> (Sch. Bip. ex Walp.) M.A. Isawumi, G.El-Ghazaly & B. Nordenstam)	Inyathelo (Z)	leaves	Diarrhoea (Mlambo, 2008)	Immunomodulating activity (Nergard <i>et al.</i> , 2004); antibacterial activity (Deeni and Hussain, 1994)	pectic arabinogalactan (Nergard <i>et al.</i> , 2004)
<i>Vernonia natalensis</i> Sch. Bip. ex Walp.	Uhlambihloshane, Isibhaha	Leaves, stem	Decoction for stomach cramps, nervous spasms of the stomach (Fawole <i>et al.</i> ,	Anti-inflammatory (Fawole <i>et al.</i> , 2009a), antimicrobial, mutagenicity	-

			2009b; (Hutching <i>et al.</i> , 1996)	(Fawole <i>et al.</i> , 2009b)	
<i>Vernonia oligocephala</i> Sch. Bip	lihlunguhlungu	Roots	Infusion for diarrhoea (Amusan <i>et al.</i> , 2007)	-	-
<i>Vernonia myriantha</i> Hook. F (syn <i>Vernonia stipulacea</i> Klatt)	Mululudza (V)	Roots	Diarrhoea, fever, flu, contraceptive (Bessong <i>et al.</i> , 2005; Obi <i>et al.</i> , 2003)	-	-
<i>Vernonia tigna</i> Klatt syn <i>V. corymbosa</i>	Uhlunguhlungu (Z), Phathaphathane (V)	Leaves	Diarrhoea (Mlambo, 2008)	-	-
Balanitaceae					
<i>Balanites maughamii</i> Sprague		leaves	Diarrhoea in cattle (Luseba and Van der Merwe, 2006; McGaw <i>et al.</i> , 2008)	Antiplasmodial and cytotoxicity (Prozesky <i>et al.</i> , 2001)	-
Balanophoraceae					
<i>Sarcophyte sanguine</i> Sparrm		whole plant	Amenorrhoea, dysentery, diarrhoea and swellings growth (Iwalewa <i>et al.</i> , 2007)	Antibacterial and antifungal (Naidoo <i>et al.</i> , 1992)	Eriodictyol, naringenin, triandrin, n-pinitol (ID-4-O-methyl chiroinositol), trans-p-coumaraldehyde, Exocarpic acid (13E-octadecene-9,11-diyonic acid)
Bignoniaceae					
<i>Markhamia sessilis</i> Sprague		Leaves	Decoction for diarrhoea (De Villiers <i>et al.</i> , 2010)	Antiplasmodial and cytotoxicity (Mbatchi <i>et al.</i> , 2006), antimicrobial and antimalaria (De Villiers <i>et al.</i> , 2010)	-
<i>Kigelia africana</i> (Lam.) Benth.		Bark	Dysentery and stomach ailments (van Wyk, 2008b)	Antidiarrhoea (Akah, 1996), analgesic and anti-inflammatory (Owolabi and Omogbai, 2007), antifungal and antibacterial (Owolabi <i>et al.</i> , 2007)	Verminoside and Verbascoside (Picerno <i>et al.</i> , 2005)
<i>Tecomaria capensis</i> Spach		Bark	fever, diarrhea and dysentery, pains, sleeplessness, stomach and chest pains (Iwalewa <i>et al.</i> , 2007)	Antimicrobial (Saini <i>et al.</i> , 2011)	-
Bombacaceae					
<i>Adansonia digitata</i> L.	Muvhuyu (V)	Leaves, bark, root fruit	Fever, diarrhoea, haemoptysis, hiccup remedy (van Wyk, 2008b)	Anti-inflammatory, antiviral (Selvarani and Hudson, 2009), antihyperglycemic and hypolipidemic (Bhargav <i>et al.</i> , 2009), Antimicrobial (Mulaudzi <i>et al.</i> , 2011)	Epicatchin, procyanidin B2, procyanidin B5 (Kinghorn <i>et al.</i> , 2011)
Bursareceae					
<i>Commiphora harveyi</i> (Engl.) Engl.		Leaves	Disinfectant for wound, anthelmintic and snakebite (Watt and Breyer-Brandwijk, 1962)	Antimicrobial (Suleiman <i>et al.</i> , 2010)	-

Capparaceae					
<i>Capparis tomentosa</i>	Umqoqolo (Z), Muoba-dali (V)	Root infusions and decoctions	Diarrhoea in cattle, stomach ailments in animals (Watt and Breyer-Brandwijk, 1962, Pujol, 1990, McGaw <i>et al.</i> , 2008)	Antimicrobial (Ramalivhana <i>et al.</i> , 2010), antifungal (Samie <i>et al.</i> , 2010)	Stachydrine L-form (Systolic depressant, rheumatism)
Caricaceae					
<i>Carica papaya</i> L.	Papawe (V)	Leaves, seed	Amoebic dysentery, fever, gastric problems, asthma, immune-stimulant (Green <i>et al.</i> , 2010; Aruoma <i>et al.</i> , 2006)	Antiamoebic (Tona <i>et al.</i> , 1998), anthelmintic (Kermanshai <i>et al.</i> , 2001)	Alternariol Carpamine (cardiotonic agent, CNS depressant), Chymopapain ; Glycerol triacetate (antifungal and adjuvant); Papain ; 2,4'-Dihydroxy-3',5'-dimethoxyacetophenone (antifungal), Benyl isothiocyanate (Kermanshai <i>et al.</i> , 2001)
Caryophyllaceae					
<i>Krauseola mossambicina</i> (Moss.) Pax & K. Hoffm.	Isihlaza, Isihlazi		Diarrhoea (de Wet <i>et al.</i> , 2010)	-	-
Celastraceae					
<i>Elaeodendron transvaalense</i> (Burt Davy) R.H. Archer syn <i>Cassine transvaalensis</i>	Mulumanamana Mukuhazwhi, Umgugudo (Z)	Bark	Cough, piles, venereal diseases, diarrhoea, stomach ache, laxative (Samie <i>et al.</i> , 2010)	Antimicrobial (Tshikalange <i>et al.</i> , 2005), hypoglycaemic (Deutschlander <i>et al.</i> , 2009), Cytotoxicity (Tshikalange and Hussein, 2010)	lup-20(30)-ene-3,29-diol , lup-20(29)-ene-30-hydroxy-3-one-(2), ψ- taraxastanonol, β-sitosterol and 4' -O-methylepigallocatechin (Tshikalange and Hussein, 2010)
<i>Gomphocarpus fruticosus</i> Dryand.		Leaf infusion	Diarrhoea and stomach ache in children (Hutchings <i>et al.</i> , 1996; Fouche <i>et al.</i> , 2008)	-	Gomphoside (cardiotonic agent)
<i>Gymnosporia senegalensis</i> (Lam.) Loes	Ubuhlangwe		Diarrhoea (de Wet <i>et al.</i> , 2010)		-
<i>Maytenus heterophylla</i> Eckl. & Zeyh.) Robson	Isibhubu (Z), Tshiphandwa (V)	Bark and leaf infusions	Diarrhoea in stock animals (Watt and Breyer-Brandwijk, 1962; McGaw <i>et al.</i> , 2008)	Antimicrobial (Orabi <i>et al.</i> , 2001), anti-inflammatory and cytotoxicity (Da Silva <i>et al.</i> , 2010), anticytomegalovirus (Murayama <i>et al.</i> , 2007)	1β-acetoxy-9α-benzoyloxy-2β,6α-dinicotinoyloxy-β-dihydroagarofuran, β-amyrin, maytenfolic acid, 3α-hydroxy-2-oxofriedelane-20α-carboxylic acid, lup-20(29)-ene-1β,3β-diol, (-)-4'-methylepigallocatechin, and (-)-epicatechin (Da Silva <i>et al.</i> , 2010) , pristimerin, lupeol and 2-acetylphenol-1-β-D-glucopyranosyl (1→6)-β-D-xylpyranoside (acetophenol glycoside) (Murayama <i>et al.</i> , 2007)
<i>Maytenus peduncularis</i> (Sond.) Loes.	Mukwatule (V)	root	Backache, pain (Gonzalez <i>et al.</i> , 2000)	-	-
<i>Maytenus procumbens</i> (L.f.) Loes.	-	-	-	-	-

<i>Maytenus senegalensis</i> (Lam.) Exell	Tshiphandwa (V)		Root used for chest pain, rheumatism, snakebites, diarrhoea and fever. Leaves for eye infection (Matu and van Staden, 2003)	Antimicrobial and anti-inflammatory (Matu and van Staden, 2003); anti-inflammatory and cytotoxicity (Da Silva et al., 2010)	Wilforine (insecticidal), β -amyrin, lupenone, maytenoic acid, β -sitosterol, pristimerin (Da Silva et al., 2010)
<i>Maytenus undata</i> (Thunb.) Blakelock	Tshinembane (V)	Leaves		Antimicrobial, anti-inflammatory and antioxidant (Muhammed et al., 2000), antimalaria (Muthaura et al., 2007)	3-oxo-11 α -methoxyolean-12-ene-30-oic acid, 3-oxo-11 α -hydroxyolean-12-ene-30-oic acid, 3-oxo-olean-9(11),12-diene-30-oic acid, 3,4-seco-olean-4(23),12-diene-3,29-dioic acid (20-epikoetjapic acid), 3,11-dioxoolean-12-ene-30-oic acid (3-oxo-18 β -glycyrrhetic acid), koetjapic acid, 12-oleanene artifact 3-oxo-11 α -ethoxyolean-12-ene-30-oic acid (Muhammed et al., 2000)
Chenopodiaceae					
<i>Atriplex nummularia</i> Lindl.		Leaves, flower	Diarrhoea (Van Wyk et al., 2008)	Antitumorigenic activity (Amara et al., 2008)	-
<i>Chenopodium ambrosioides</i> L.	Unakani, Ikhambi	Aerial part	Diarrhoea (de Wet et al., 2010)	Antisecretory against cholera toxin (Velazquez et al., 2006), antiamoeba and antigardia (Calzada et al., 2006)	Ascaridole, quercetin, kaempferol, isorhamnetin, ambroside, malic acid, succinic acid
Clusiaceae					
<i>Garcinia livingstonei</i> T. Anderson	Umphimbi, Muphiphi (V)	leaves	Diarrhoea (de Wet et al., 2010)	Antibacterial (Kaikabo, 2008)	Amentoflavone (Bradykinin antagonist, anti-HIV activity, inhibitor of human cathepsin B, anti-inflammatory properties), amentoflavone and 4"-methoxy amentoflavone (Kaikabo, 2008)
Combretaceae					
<i>Combretum bracteosum</i> (Hochst.) Brandis ex Engl.		leaves	-	anti-inflammatory, anthelmintic and antischistosomal (McGaw et al., 2001)	-
<i>Combretum imberbe</i> Wawra	Mudzwiri (V)	Root	Decoction for diarrhoea (Mathabe et al., 2006)	anti-inflammatory, anthelmintic and antischistosomal (McGaw et al., 2001), antimicrobial (Angeh et al., 2007)	1 α , 23 β -Dihydroxyl-12-Oleanen-29-oic acid-23 β -O- α -4-acetylramnopyranoside; 1, 22-Dihydroxyl-12-Oleanen-30-oic acid; Ethyl cholesta-7, 22,25-trien-O- β -D-glucopyranoside (Angeh et al., 2007), imberbic acid (Katerere et al., 2003)
<i>Combretum molle</i> R. Br.ex G. Don	Mugwiti (V)	Roots	Abdominal pain, fever, snake bite, leprosy and convulsions (Bessong et al., 2005; Mabogo, 1990)	anti-inflammatory, anthelmintic and antischistosomal (McGaw et al., 2001)	Punicalgin, 4- <i>epi</i> -sericoside, sericoside (Asres et al., 2001), β -D-glucopyranosyl 2 α ,3 β ,6 β -trihydroxy-23-galloylolean-12-en-28-oate, combregenin, arjungenin, arjunglucoside I, combreglucoside (Ponou et al., 2008), mollic acid glucoside (Oyewole, 2008)
<i>Combretum padoides</i> Engl. & Diels				anti-inflammatory, anthelmintic and antischistosomal (McGaw et al., 2001); antifungal (Masoko et al., 2007); Antibacterial (Angeh et al., 2007)	1 α ,23 β -dihydroxy-12-oleanen-29-oic-acid-23 β -O- α -4-acetylramnopyranoside, 1,22-dihydroxy-12-oleanen-30-oic acid, 24-ethylcholesta-7,22,25-trien-O- β -D-glucopyranoside (Angeh et al., 2007)
<i>Combretum vendae</i> A.E. van Wyk		Leaves	Leprosy, ophthalmic remedy, and blood purification (Watt and Breyer-Brandwijk,	Antimicrobial (Ahmed et al., 2008; Suleiman et al., 2010)	apigenin (Eloff et al., 2008)

			1962)		
<i>Combretum woodii</i> Dummer				anti-inflammatory, anthelmintic and antischistosomal (McGaw <i>et al.</i> , 2001); antifungal (Masoko <i>et al.</i> , 2007)	Combretastatin B5 (Eloff <i>et al.</i> , 2005)
<i>Combretum zeyheri</i> Sond	Mufhatela-thundu, Mufhatela (V)	Root infusion	Bloody diarrhoea (Hutchings <i>et al.</i> , 1996; Fouche <i>et al.</i> , 2008)	Antibacterial (Breytenbach and Malan, 1989)	
<i>Terminalia laxiflora</i> Engl.		Leaves	Decoction for diarrhoea (De Villiers <i>et al.</i> , 2010)	Antifungal (Batawila <i>et al.</i> , 2005)	
<i>Terminalia phanerophlebia</i> Engl.		Root bark	Diarrhoea and colic (Iwalewa <i>et al.</i> , 2007)	Antimicrobial (Shai <i>et al.</i> , 2008a)	
<i>Terminalia sericea</i> Burch. ex DC.	Mususu (V), Ikonono	Leaves roots	Wound (Luseba and Van der Merwe, 2006); diarrhoea (Van der Merwe <i>et al.</i> , 2001; McGaw <i>et al.</i> , 2008)	Antimicrobial, antidiabetic, cytotoxicity (Moshi and Mbwambo, 2005), COX-1 and COX-2 assays (Eldeen <i>et al.</i> , 2006).	Anolignan B (Eldeen <i>et al.</i> , 2006), Termilignan B, Arjunic acid (Eldeen <i>et al.</i> , 2008), 3'5'-dihydroxy-4-(2-hydroxy-ethoxy) resveratrol-3-O- β -rutinoside, resveratrol-3- β -rutinoside glycoside, 3',4,5'-Trihydroxystilbene (resveratrol), arjungenin (Joseph <i>et al.</i> , 2007)
Convolvulaceae					
<i>Ipomoea batatas</i> (L.) Lam.	Sweet potato	Leaves	Decoction for diarrhoea (De Villiers <i>et al.</i> , 2010)		4,5-Di-transcaffeoyldenoic acid (antioxidant), 6-O-caffeoylsophorose (α -glucosidase inhibitor, antioxidant); 3,5-Di-O-caffeoylquinic acid (active against HIV-1 integrase, antiviral, antihepatotoxic activity); Petrovin B (antibacterial and antitumour)
Cornaceae					
<i>Curtisia dentata</i> (Burm.f.) C.A.Sm.= <i>C. faginea</i> Assegaai	Umlahlani (X, Z) Unsirayi (X), Umgxina	Bark, root	Diarrhoea, stomach ailments (Bisi-Johnson <i>et al.</i> , 2010)	Antimicrobial (Shai <i>et al.</i> , 2008a, Shai <i>et al.</i> , 2009)	Lupeol, betulinic acid, ursolic acid, and 2 α -hydroxyursolic acid (Shai <i>et al.</i> , 2008b)
Crassulaceae					
<i>Crassula ovata</i> (Mill.) Druce	Karkay, karkey (K)	Fresh leave	Diarrhoea (van Wyk, 2008)	-	-
<i>Crassula tetragona</i> L.	Karkay, karkey (K)	Fresh leave	Diarrhoea (van Wyk, 2008)	-	-
Cucurbitaceae					
<i>Cucumis hirsutus</i> Sond.		Leaves, root	Decoction for abdominal pains, diarrhoea (Fawole <i>et al.</i> , 2009); Hutching <i>et al.</i> , 1996)	-	-
<i>Mormodica balsamina</i> L.	Lubavhe (V)	Whole plant	Diabetes, childhood diarrhoea (Samie <i>et al.</i> , 2009)	Shigellocidal (Iwalokun <i>et al.</i> 2001); Cytotoxicity and antiameobic (Samie <i>et al.</i> , 2009)	Balsaminapentaol A, Balsaminol A, Balsaminol B, Cucurbalsaminol A, Cucurbalsaminol B, cucurbita-5,23(E)-diene-3 β ,7 β ,25-triol, karavilagenin E (Ramalhete <i>et al.</i> , 2009)
Ebenaceae					
<i>Diospyros lycioides</i> Desf.	Umbulwa (Z)	Bark, root	Decoction for bloody faeces and dysentery (Fawole <i>et al.</i> , 2009); (Hutching <i>et al.</i> ,	-	Hydroxyisodispyrin (cytotoxic agent)

			1996)		
<i>Diospyros mespiliformis</i>	Musuma	Bark and leaves	Dysentery, fever, ringworm, skin infection, wound healing (Samie <i>et al.</i> , 2010)		Diosquinone, plumbagin (Lajubutu <i>et al.</i> , 1995)
<i>Diospyros pallens</i> (Thunb.) F. White		Root and stem	Stomach arch; diarrhoea (van Wyk, 2008)	-	-
<i>Euclea crispa</i> Thunb Gurke	Ungwali (Z),	leaves	Dysmenorrhoea (Steenkamp, 2003)		
<i>Euclea natalensis</i> A. DC	Mutangule-thavha (V), Umzimane (Z)	root	oral health care, for chest complains, bronchitis, pleurisy, chronic asthma, urinary tract infections, venereal diseases (Lall and Meyer, 2001), Infertility and abortifacient (Arnold and Gulumian, 1984)	Antibacterial (Weigenand <i>et al.</i> , 2004), antimycobacterium (Lall and Meyer, 2001)	Octahydroeuclein, 20(29)-lupene-3 β -isoferulate, shinanolone, lupeol, betulin (Weigenand <i>et al.</i> , 2004); diospyrin (Lall and Meyer, 2001)
Euphorbiaceae					
<i>Antidesma venosum</i> E. Mey. ex Tul.	Mupalakhwali (V)	Leaf	Decoction for abdominal cramps and dysentery (Fawole <i>et al.</i> , 2009; Hutching <i>et al.</i> , 1996)	Antimicrobial (Fawole <i>et al.</i> , 2009)	-
<i>Bridelia micrantha</i> (Hochst.) Baill	Munzere (V)	Bark, leaves, roots	Stomach ache, diarrhoea, abortifacient (Bessong <i>et al.</i> , 2005; Lin <i>et al.</i> , 2002), Gastro-intestinal ailments, painful joints, retained placenta, diabetes mellitus, syphilis prehepatic jaundice, tape worm abdominal pain, conjunctivitis, headache, scabies, bloody diarrhoea, dysentery, emetic, wound infection, coughs, threadworms, tonic for children, sore eyes, epigastric pain, relief of headache, purgative (Nguemem <i>et al.</i> , 2009), diabetes mellitus (Abo <i>et al.</i> , 2008)	Antidiarrhoea (Lin <i>et al.</i> , 2002), beta-lactamase inhibition (Gangoue-Pieboji <i>et al.</i> , 2007); antimalarial (Abo and Ashidi (1999). <i>n</i> -butanol fraction of methanolextract has IC50 of 7.3 μ g/ml against the RNA-dependent DNA polymerization (RDDP) function of HIV-1 RT (Bessong <i>et al.</i> , 2006)	Taraxerol, gallic and ellagic acid, friedelin, delphinidin, methyl salicylate (Nguemem <i>et al.</i> , 2009)
<i>Bridelia mollis</i>	Mukumbakumba	Leaves	Dysentery, burning and itching (Samie <i>et al.</i> , 2010)	Antifungal (Samie <i>et al.</i> , 2010)	
<i>Euphorbia cooperi</i> N. B. Br. Ex. Berger	Umhlonhlo (X)	Root bark	Diarrhoea, stomach disorder (Bisi-Johnson <i>et al.</i> , 2010)	-	-
<i>Euphorbia hirta</i> L.		Leaves	Decoction for diarrhoea (De Villiers <i>et al.</i> , 2010); dysentery, gonorrhoea, jaundice, pimples, digestive problems and tumours, antibacterial, anti-inflammatory, antimalarial, galactogenic, antiasthmatic, antidiarrheal, anticancer, antioxidant, antifertility, antiamoebic, and antifungal activities (Kumar <i>et al.</i> , 2010)	Antiamoebic, spasmolytic (Tona <i>et al.</i> , 2000), Antidiarrhoeal (Galvez <i>et al.</i> , 1993)	β -amyrin, 24-methylenecycloartenol, β -sitosterol, Quercitrin (Galvez <i>et al.</i> , 1993). Quercitol, gallic acid, afzelin, quercitrin, myricitrin, rutin, gallic acid, quercitin, euphorbin-A and euphorbin-B, euphorbin-C, euphorbin-D, β -amyrin, 24-methylenecycloartenol, β -sitosterol, heptacosane, <i>n</i> -nonacosane, shikmic acid, tinyatoxin, choline, camphol, and quercitol (Kumar <i>et al.</i> , 2010)
<i>Jatropha zeyheri</i> Sond.	Xidomeja	Roots	General ailments, diarrhoea (Luseba and Van der Merwe, 2006; McGaw <i>et al.</i> , 2008)	Antimicrobial and Antifungal (Dekker <i>et al.</i> , 1987)	Jaherin (Dekker <i>et al.</i> , 1987)



<i>Ricinus communis</i> L.	Mupfure (T)	leaves	Wound and sores, asthma arthritis, flu, fever, tuberculosis, toothache, diarrhoea, antihelmentic (Bessong <i>et al.</i> , 2005; Grierson and Afolayan, 1999)	-	-
<i>Spirostachys africana</i> Sond	Morekhure (S)	Wood	Stomach ulcers, acute gastritis, headache, rashes, boil, emetic, purgative, diarrhoea, dysentery (Verschaeve and Van Staden, 2008)	Antibacterial and cytotoxicity (Mathabe <i>et al.</i> , 2008)	-
Fabaceae					
<i>Acacia burkei</i> Benth	umkhaya		Diarrhoea (de Wet <i>et al.</i> , 2010)	-	-
<i>Acacia karoo</i> Hayne	uMnga (X), Umunga (Z)	Bark, leaves	Diarrhoea, intestinal parasites in goats, sheep, poultry and pig (Dold and Cocks, 2001; McGaw <i>et al.</i> , 2008) fractures and diarrhoea (Van der Merwe <i>et al.</i> , 2001)	Anti-inflammatory (Adedapo <i>et al.</i> , 2008); Acute toxicity (Adedapo <i>et al.</i> , 2008)	-
<i>Acacia mearnsii</i> De Wild Blackwood	Ublekwana (X) Udywabasi (X, Z) Indywabasi	Bark	Infusion for diarrhoea and dysentery (Bisi-Johnson <i>et al.</i> , 2010)	Protective against acrolein-induced oxidative damage (Huang <i>et al.</i> , 2010)	Robinetinidol-(4 β →8)-epigallocatechin 3-O-gallate (Huang <i>et al.</i> , 2010)
<i>Acacia robusta</i> E. Meyer	Umngamanzi (Z)	leaves	Diarrhoea (Mlambo, 2008)	Antifungal (Hamza <i>et al.</i> , 2006)	
<i>Acacia sieberiana</i> DC.var <i>woodii</i> (Burt Davy) Keay & Brenan	Musaunga, Muunga-luselo (V)	Bark	Enemas, antiseptic, fever, stomach ache, tapeworm, astringent, haemostatic, diarrhoea (Verschaeve and Van Staden, 2008)	Mutagenicity, antimutagenicity; antibacterial, antiinflammatory, anticholinesterase and mutagenic effects (Eldeen <i>et al.</i> , 2005)	-
<i>Acacia tortilis</i> (Forssk.) Hayne	Muunga-khanga, Muswu (V)		Diarrhoea (Van der Merwe <i>et al.</i> , 2001; McGaw <i>et al.</i> , 2008)		-
<i>Bauhinia bowkeri</i> Harv	uMdandlovu	Leaves, bark	Induce vomiting (Ndawonde <i>et al.</i> , 2007)	-	-
<i>Bauhinia galpinii</i> N. E. Br	Mutswiriri (V), Umhuwa (Z)	Bark, leaves	Diarrhoea, infertility (Samie <i>et al.</i> , 2010), infertility (Arnold and Gulumian, 1984), amenorrhoea (Van Wyk and Gericke, 2000)	Antimutagenic (Reid <i>et al.</i> , 2006); antioxidant and cytotoxicity of leaf extracts (Aderogba <i>et al.</i> , 2007); Anticampylobacterial, antiamoebic and cytotoxicity of root extract (Samie <i>et al.</i> , 2009)	Quercetin-3-O- β -glactopyranoside, Myricetin-3-O- β -glactopyranoside, 2"-O-rhamnosylvitexin (Aderogba <i>et al.</i> , 2007)
<i>Bauhinia petersiana</i> Bolle	Mushakule (V)	root	Cold (Coates-Palgrave, 2002); infertility and dysmenorrhoea (Van Wyk and Gericke, 2000)	-	-
<i>Bauhinia variegata</i> Linn		Leaves, bark	Diabetes, goiter, dysentery, diarrhoea (Parekh and Chanda, 2007)	Anti-inflammatory (Rao <i>et al.</i> , 2008); Immunomodulator (Ghaisas <i>et al.</i> , 2009)	kaempferol, ombuin, kaempferol 7,4'-dimethyl ether 3-O- β -D-glucopyranoside, kaempferol 3-O- β -D-glucopyranoside (4), isorhamnetin 3-O- β -D-glucopyranoside, hesperidin, 3 β -trans-(3,4-dihydroxycinnamoyloxy)olean-12-en-28-oic acid (Rao <i>et al.</i> , 2008)

<i>Dichrostachys cinerea</i> (L.) Wight and Am.	Murenzhe (V)	Bark	Diarrhoea and steaming to get ride of acne (Mlambo, 2008)	spasmodic in guinea-pig isolated trachea (Aworet-Samseny <i>et al.</i> , 2011)	dichrostachines A-R (Long <i>et al.</i> , 2009)
<i>Elephantorrhiza burkei</i> Benth.	Umdabu (Z), Tshisese-thavha, Tshisesevhufa (V)	root	abdominal pains, diarrhoea, coughs, bacterial infections (lwalewa <i>et al.</i> , 2007)	Antimicrobial (Mathabe <i>et al.</i> , 2006)	Triterpenoids, α -amyrim, β -sitosterol, alkaloids and saponin
<i>Elephantorrhiza evoluta</i> (Burch.) Skeels	iNtololwane (X, Z)	Roots, aerial part and bulb	Diarrhoea and dysentery in cattle, horse and humans (Watt and Breyer-Brandwijk, 1962; McGaw <i>et al.</i> , 2008)	-	-
<i>Elephantorrhiza elephantina</i> (Burch.) Skeels	Leshitsana	Stem rhizome	Decoction for diarrhoea (Mathabe <i>et al.</i> , 2006)	Antimicrobial (Mathabe <i>et al.</i> , 2006), antiparasitic (Naidoo <i>et al.</i> , 2006), antibabesia (Naidoo <i>et al.</i> , 2005)	
<i>Eriosema psoraleoides</i> (Lam.) G. Don		Leaves	Decoction for diarrhoea (De Villiers <i>et al.</i> , 2010)	Antimicrobial (Khan <i>et al.</i> , 2000)	
<i>Erythrina latissima</i> E. Mey	Muvhale (V)		Sores (Coates-Palgrave, 2002)		erysotrine, erysodine, syringaresinol, vanillic acid, (+)-10,11-dioxoerysotrine, 2-(5'-hydroxy-3'-methoxy phenyl)-6-hydroxy-5-methoxybenzofuran, 7,3'-dihydroxy-4'-methoxy-5'-(γ,γ -dimethylallyl)isoflavone (erylatissin A) (Wanjala <i>et al.</i> , 2002), 7,3'-dihydroxy-6",6"-dimethyl-4",5"-dehydropyrano [2",3": 4',5']isoflavone (erylatissin B), (-)-7,3'-dihydroxy-4'-methoxy-5'-(γ,γ -dimethylallyl)flavanone (erylatissin C) (Chacha <i>et al.</i> , 2004)
<i>Indigofera daleoides</i> Benth. ex Harv & Sond		Whole plant	Decoction for diarrhoea (Mathabe <i>et al.</i> , 2006)	Antimicrobial (Mathabe <i>et al.</i> , 2006)	(6,2-O-[3-nitropropanoyl- β -D-glucopyranose]), (6,3',4'-trihydroxyflavan 5'-O-glucopyranoside) (Mathabe <i>et al.</i> , 2009)
<i>Indigofera jucunda</i> Schrire syn <i>Indigofera cylindrical sensu</i> E. Mey		Root	Intestinal worm (Coates-Palgrave, 2002)	-	-
<i>Indigofera sessilifolia</i> DC.	iKhubalo	Roots	Diarrhoea in calves (Dold and Cocks, 2001; McGaw <i>et al.</i> , 2008)	-	-
<i>Mucuna coriacea</i> Baker	Vhaulada	Roots	Fever, diarrhoea (Bessong <i>et al.</i> , 2005)	Antimicrobial (Samie <i>et al.</i> , 2009)	N.A
<i>Peltophorum africanum</i> Sond.	Musese (T)	Bark, root bark	Tonic, diarrhoea (Van der Merwe <i>et al.</i> , 2001; McGaw <i>et al.</i> , 2008)	Anti parasitic (Bizimenyera <i>et al.</i> , 2006), anti HIV (Bessong <i>et al.</i> , 2005)	Catechin, gallotannin, bergenin (Bessong <i>et al.</i> , 2005)
<i>Rhynchosia adenoids</i> E. & Z.		Roots	Decoction for rheumatic pains, menstrual pains and dysentery (Shale <i>et al.</i> , 1999)	Cyclooxygenase inhibitory (Jager and Van Staden, 2005)	
<i>Senna italic</i> Mill.	Ximbangam bangana	Bark, roots	Diarrhoea and gallsickness diarrhoea, (Luseba and Van der Merwe, 2006; McGaw <i>et al.</i> , 2008)	-	--

<i>Senna occidentalis</i> (L.) Link	Ikhoshokhosh	Leaves, root	Diarrhoea (de Wet <i>et al.</i> , 2010)		
<i>Schotia brachypetala</i> Sond.	Mulubi (V)	Bark	Diarrhoea (Mathabe <i>et al.</i> , 2006), root for dysentery and diarrhoea (Hutching <i>et al.</i> , 1996)	Antibacterial (McGaw <i>et al.</i> , 2002)	Linolenic acid and methyl-5,11,14,17-eicosatetraenoate (McGaw <i>et al.</i> , 2002)
<i>Schotia latifolia</i> Jacq.	Umgxam	bark	Decoction for diarrhoea (Appidi <i>et al.</i> , 2008)	Antibacterial (Masika <i>et al.</i> , 2004)	Epicatechin and catechin (Masika <i>et al.</i> , 2004)
<i>Zornia milneana</i>	Lukandulula (V)	Whole plant	Dysentery and diarrhoea (Samie <i>et al.</i> , 2005)	Anticampylobacterial and antiamoebic (Samie <i>et al.</i> , 2009)	-
Flacourtiaceae					
<i>Oncoba spinosa</i> Lam		root	Dysentery bladder problem (Verschaeve and Van Staden, 2008)	Mutagenicity, antimutagenicity, Epilepsy and convulsion (Risa <i>et al.</i> , 2004)	-
Gentianaceae					
<i>Chironia baccifera</i> L.			treat acne, sores and diarrhoea (Watt and Breyer-Brandwijk, 1962; van Wyk <i>et al.</i> , 1997)	Antibacterial (Thring <i>et al.</i> , 2007)	-
Geraniaceae					
<i>Geranium incanum</i> Burm. f.	Isikhwali (Z)	leaves	Diarrhoea (Amabeoku, 2009; Van Wyk <i>et al.</i> 1997)	Antidiarrhoeic (Amabeoku, 2009); antimicrobial and cytotoxicity (Babajide <i>et al.</i> , 2010)	-
<i>Monsonia emarginata</i> (L.f.) L'Hér.		Herb and root	Diarrhoea, dysentery, cold and inflammation (van Wyk, 2008)	-	-
<i>Monsonia burkeana</i> Planch. Ex Harv.		Herb and root	Diarrhoea, dysentery, cold and inflammation (van Wyk, 2008)	Antioxidant (Mamphiswana <i>et al.</i> , 2010)	-
<i>Pelargonium antidysentericum</i> (Eckl. & Zeyh.) Kostel		Tubers	Used as astringent, diarrhoea, dysenteric fever (Brendler and van Wyk, 2008)	-	-
<i>Pelargonium luridum</i> (Andr.) Sweet	Umsongelo (X) ishwaqa	Leaf, root	Diarrhoea, dysentery, fever and colic (Brendler and van Wyk, 2008)	-	-
<i>Pelargonium reniforme</i> Curtis	iNtololwana, uVendle	Tuberous root	Diarrhoea and dysentery (van Wyk, 2008)	Antibacterial, antifungal and antioxidant (Adewusi and Afolayan, 2009a), Acute toxicity (Adewusi and Afolayan, 2009b)	scopoletin, umckalin, 5,6,7-trimethoxycoumarin, 6,8-dihydroxy-5,7-dimethoxycoumarin, (+)-catechin, gallic acid (Kayser and Kolodziej, 1997)
<i>Pelargonium sidoides</i> DC.	Umsongelo (X)		Tuberculosis, diarrhoea (Brendler and van Wyk, 2008)	Immunomodulatory (Kayser <i>et al.</i> , 2001), antibacterial, antifungal and antitubercular (Mativandlela <i>et al.</i> , 2006)	scopoletin, umckalin, 5,6,7-trimethoxycoumarin, 6,8-dihydroxy-5,7-dimethoxycoumarin, (+)-catechin, gallic acid (Kayser and Kolodziej, 1997)
<i>Pelargonium triste</i>		Tuberous	Diarrhoea and dysentery (van Wyk, 2008)	-	-

(L.) L'Hér.		root			
Hyacinthaceae					
<i>Eucomis autumnalis</i> (Mill.) Chitt.	Ubuhlungu becanti Isithithibala (X) Umathinga (Z)	Bulb	Stomach ache, diarrhoea, back pain, healing of fractures (Bisi-Johnson <i>et al.</i> , 2010)	Anti-inflammatory (Zschocke <i>et al.</i> , 2000)	-
<i>E. regia</i> (L.) L'Herit			Veneral diseases, lumbago, diarrhoea, respiratory conditions especially coughs, biliousness and to prevent premature childbirth (Watt and Breyer-Brandwijk, 1962)	COX-1 assay (Taylor and van Staden, 2001)	-
<i>Ledebouria revoluta</i> (L.f.) Jessop	iKreketsana (X)	Bulb	Bulb infusion for diarrhoea in goat, leaf decoction for gallsickness (Dold and Cocks, 2001; McGaw <i>et al.</i> , 2008)		(3R)-5,7-dihydroxy-3-(4'-methoxybenzyl)-4-chromanone, (3R)-5,7-dihydroxy-3-(4'-hydroxybenzyl)-4-chromanone, 3R)-5-hydroxy-7,8-dimethoxy-3-(4'-hydroxybenzyl)-4-chromanone, (3R)-5,7-dihydroxy-8-methoxy-3-(4'-hydroxybenzyl)-4-chromanone (Moodley <i>et al.</i> , 2006)
<i>Schizocarphus rigidifolius</i>	Ingcino (S)	leaves	Infusion for diarrhoea (Amusan <i>et al.</i> , 2007)	-	-
<i>Scilla nervosa</i> (Burch.) Jessop	Umagaqana, magaqana (X) Imbizankulu ingema (Z)	Root, bulb	All purpose herb. Diarrhoea, tuberculosis (Bisi-Johnson <i>et al.</i> , 2010)	-	-
Hydnoraceae					
<i>Hydnora africana</i>	Ubuklunga (X) Umavumbuka (Z) Umafumbuka (X)	Fruits, tuber leaves	Diarrhoea, dysentery (Bisi-Johnson <i>et al.</i> , 2010)	-	-
Hypoxidaceae					
<i>Hypoxis latifolia</i> Hook.	Inongwe llabateka (X)	Tuber	Decoction for diarrhoea (Bisi-Johnson <i>et al.</i> , 2011)	Antibacterial, antifungal (Buwa and Van Staden, 2006)	
<i>Hypoxis hemerocallidea</i> Fisch. & C. A. Mey	Inongwe llabateka (X)	Tuber	Decoction for diarrhoea (Ojewole <i>et al.</i> , 2009)	Antinociceptive, anti-inflammatory and antidiabetic (Ojewole, 2006), Antidiarrhoeal, acute toxicity test (Ojewole <i>et al.</i> , 2009)	
Iridaceae					
<i>Crocasmia paniculata</i> (Klatt.) Goldbl.	Undwendweni (Z)	corms	Diarrhoea in bovine (Watt and Breyer-Brandwijk, 1962; McGaw <i>et al.</i> , 2008)	-	-
<i>Gladiolus dalenii</i> van Geel		corm	Dysentery, diarrhoea, stomach cramps (Fawole <i>et al.</i> , 2009; Hutching <i>et al.</i> , 1996)	Anti-inflammatory (Fawole <i>et al.</i> , 2009; Ndhala), amoebicidal (Moundipa <i>et al.</i> , 2005), Antimicrobial and mutagenicity (Fawole <i>et al.</i> , 2009; Finnie)	-

<i>Gladiolus sericeo-villosus</i> Hook. F	Umnunge (X), Umlunge (Z)	Corm	Decoction for dysentery, cold, tuberculosis diarrhoea (Bisi-Johnson <i>et al.</i> , 2010)	-	-
<i>Watsonia densiflora</i> Bak.		Corm	Diarrhoea in calves (Watt and Breyer-Brandwijk, 1962; McGaw <i>et al.</i> , 2008)	Antibacterial, antifungal, acetylcholinesterase inhibition, mutagenicity, COX 1 and 2 (Ndhala <i>et al.</i> , 2010)	
<i>Watsonia tabularis</i> Bak		corm	Diarrhoea in human and calves (Fawole <i>et al.</i> , 2009; Hutching <i>et al.</i> , 1996)	Antimicrobial and mutagenicity (Fawole <i>et al.</i> , 2009 Finnie)	
Lamiaceae					
<i>Ballota africana</i> (L.) Berth.					
<i>Cissus quadrangularis</i> (Linn)	Isinwasi (Z), Nyangala (T)	Root, stem	Burns, wounds, gastrointestinal complaints, backache, body- and febrile pain, malaria (Lin <i>et al.</i> , 1999; Hutchings <i>et al.</i> , 1996)	Antibacterial, anti-inflammatory and mutagenicity (Luseba, <i>et al.</i> , 2007)	
<i>Leonotis leonurus</i> (L.) R.Br	Imunyamunya (Z)	leaves and stem bark	feverish headache, dysentery, coughs and colds, and haemorrhoids (Iwalewa <i>et al.</i> , 2007)	Anticonvulsant (Bienvenu <i>et al.</i> , 2002), antinociceptive, anti-inflammatory and hypoglycaemic activities (Ojewole, 2005)	1,2,3-trihydroxy-3,7,11,15-tetramethylhexadecan-1-yl-palmitate, succinic acid, uracil, luteolin 7-O-glucoside, acteoside, geniposidic acid (Agnihotri <i>et al.</i> , 2009)
<i>Leucas capensis</i> (Benth.) Engl.	uPhiphiyo	leaves	Decoction with <i>Aloe forex</i> and <i>Brachylaena ilicifolia</i> for diarrhoea in lambs (Dold and Cocks, 2001)	-	-
<i>Mentha longifera</i> (L.) L.		Leaf	Anti-diarrhoea (Naseri <i>et al.</i> , 2008)	Spasmolyte (Naseri <i>et al.</i> , 2008)	-
<i>Rotheca myricoides</i> (Hochst.) Steane & Mabb.		Root bark	Fever and diarrhoea in cattle (Verschaeve and Van Staden, 2008)	Mutagenicity and antimutagenicity (Verschaeve and Van Staden, 2008)	-
<i>Salvia africana-caerulea</i> L.		Leaf	Coughs, colds, women ailments; diarrhoea (van Wyk, 2008)	Antimicrobial, antioxidant, anti-inflammatory, antiparasitic, cytotoxicity and antituberculosis (Kamatou <i>et al.</i> , 2006)	-
<i>Salvia repens</i> Burch. Ex. Benth		Roots, leaves, whole plant	Sores on the body, stomach problems, diarrhoea (Kamatou <i>et al.</i> , 2008)	Antimicrobial, antioxidant, anti-inflammatory, antiparasitic, cytotoxicity and antituberculosis (Kamatou <i>et al.</i> , 2006)	-
<i>Tetradenia riparia</i> (Hochst.) Codd	Iboza (Z)	leaves	Cough, sore throats, malaria, dengue, dropsy, fever, diarrhoea, haemoptysis, boils, mumps, induce drowsiness (Verschaeve and Van Staden, 2008)	Mutagenicity, antimutagenicity (Verschaeve and Van Staden, 2008); Antibacterial, antifungal, acetylcholinesterase inhibition, mutagenicity, COX 1 and 2 (Ndhala <i>et al.</i> , 2010)	-
<i>Teucrium riparium</i> Hochst	umnunu	Root	Infusion for diarrhoea (Amusan <i>et al.</i> , 2007)	-	-

Lauraceae					
<i>Ocotea bullata</i> (Burch.) Baill.		Bark	Headache, infantile diarrhoea, stomach problems, emetic for emotional and nervous disorder (Verschaeve and Van Staden, 2008)	Mutagenicity , antimutagenicity (Verschaeve and Van Staden, 2008), anti-inflammatory (Zschocke et al., 2000)	Ocobullenone, iso-ocobullenone, sibyllenone (Zschocke et al., 2000)
Loganiaceae					
<i>Strychnos henningsii</i> Gilg.	uMnonono, Umqalothi (Z)	Bark infusion	Heartwater and diarrhoea in cattle (Dold and Cocks, 2001; McGaw et al., 2008)	-	-
Loganiaceae					
<i>Sida alba</i> Forrsk		Leaves	Diarrhoea and dysentery (Samie et al., 2005)	Antibacterial (Samie et al., 2005)	-
<i>Malva parviflora</i> L.	Ujongilanga	leaves	Decoction for diarrhoea (Appidi et al., 2008)	Antibacterial and anti-inflammatory (Shale et al., 2005)	-
Melastomataceae					
<i>Dissotis princeps</i> (Kunth) Triana		Leaves	Infusion for diarrhoea and dysentery (Fawole et al., 2009; Hutching et al., 1996)	Anti-inflammatory (Fawole et al., 2009), Antimicrobial and mutagenicity (Fawole et al., 2009)	-
Meliaceae					
<i>Ekebergia capensis</i> Sparrm		Root, bark	Stomach and intestinal complaints, dysentery, heart burn, purgative, kidney problem, indigestion (Verschaeve and Van Staden, 2008)	Mutagenicity , antimutagenicity (Verschaeve and Van Staden, 2008), antiplasmodial (Murata et al., 2008)	Ekersenin, 4,6-dimethoxy-5-methylcoumarin, oleanolic acid, 3-epioleanolic acid, oleanolic acid (15), 3,11-dioxolean-12-en-28-oic acid, melliferone, 3-oxo-11,13(18)-oleandien-28-oic acid, ekeberin A, (Z)-volkendousin, ekeberin B, 7-deacetoxy-7-oxogedunin, 7-acetylneotrichilenone, proceranolide, mexicanolide, swietenolide, methylangolensate, ekeberins C1, C2, and C3, 2,3,22,23-tetrahydroxy-2,6,10,15,19,23-hexamethyl-6,10,14,18-tetracosatetraene (3R,22R), 2-hydroxymethyl-2,3,22,23-tetrahydroxy-2,6,10,15,19,23-hexamethyl-6,10,14,18-tetracosatetraene (2R,3R,22R), ekeberins D1, D2, D3, D4, and D5 (Murata et al., 2008)
<i>Melia azedarach</i> L.	Umsilinga (Z)	Leaves	Diarrhoea (de Wet et al., 2010)		
<i>Trichilia dregeana</i> Sond.	Umkhuhlu (Z)	Leaves		antibacterial, antiinflammatory, anti-cholinesterase and mutagenic effects (Eldeen et al., 2005)	cycloart-23-ene-3,25-diol (Eldeen et al., 2007)
<i>Trichilia emetica</i> Vahl.	Umkhuhlu (Z)	Leaves	Diarrhoea (de Wet et al., 2010)	Antimicrobial, antioxidant, anti-inflammatory, antimalarial, cytotoxicity (Komane et al., 2011)	sendanin, trichilin, trichilin A, trichilin B, trichilin C, trichilin D, trichilin E, dregeana, nymania 1, rohituka, rohituka, rohituka, Trichilia substance Tr-A, Trichilia substance Tr-B, Trichilia substance Tr-C and seco-A-protolimonoind (Komane et al., 2011)
Menispermaceae					
<i>Albertisia delagoensis</i> (N.E.)	Umgandanganda ,ungandingandi	Root	Diarrhoea , dysentery, cough, colic, bloody stool (De Wet and van Wyk, 2008)	Antiplasmodial and cytotoxicity (De wet et al., 2007)	

Br.) Forman					
<i>Antizoma angustifolia</i> (Burch.) Miers ex Harv		Root	Diarrhoea , dysentery, cough, colic, bloody stool (De Wet and van Wyk, 2008)	-	-
<i>Cissampelos capensis</i> (L.f.) Diels	Umbombo (Z)	Root, rhizome	Purgative, tincture for dysentery (van Wyk, 2008)	-	-
<i>Cissampelos hirta</i> Klotzch	Umanyokane, khalimelo (Z)		Diarrhoea (de Wet <i>et al.</i> , 2010)	-	-
<i>Cissampelos mucronata</i> A. Rich.		Root	Diarrhoea (Giess and Snyman, 1986; Von Koenen, 2001)	Anti-ulcer (Akah and Nwafor, 1999), sedative (Akah <i>et al.</i> , 2002), Antiplasmodial (Tshinbagu <i>et al.</i> , 2003)	Bisbenzylisoquinone alkaloid (Tshinbagu <i>et al.</i> , 2003)
<i>Cissampelos torulosa</i> E.Mey	Lukandulula (V)	Leaves	Diarrhoea and dysentery, sore throat (Mabogo, 1990; Samie <i>et al.</i> , 2005)	Antiamoebic (Samie <i>et al.</i> , 2009), antibacterial (Samie <i>et al.</i> , 2005)	-
Moraceae					
<i>Ficus capensis</i> Thunb.	Infusion	Fruit	Diarrhoea (Pallant and Steenkamp, 2008)	intestinal motility modulation (Ayinde and Owolabi, 2009)	-
<i>Ficus craterostoma</i> Mildbr. & Burret	Muumo (V), inTendekwane, umThombe(X)		Stomach-ache (Bhats and Jacobs , 1995)	-	-
<i>Ficus glumosa</i> Delile		Bark	Decoction for diarrhoea (Venter and Venter , 2002)	-	-
<i>Ficus sur</i> Forssk	Umkhiwane (Z)	Leaves	Diarrhoea (Mlambo, 2008)	Spasmolytic and gastrointestinal protection (Kunle <i>et al.</i> , 1999)	-
Myrtaceae					
<i>Psidium guajava</i> L.	Ugwava (X, Z)	Leaves	Infusion for bloody diarrhoea (Bisi-Johnson <i>et al.</i> , 2010)	Antidiarrhoeal Tona <i>et al.</i> , 1999; Lutterodt, 1992); antispasmodic (Conde <i>et al.</i> , 2003), antirotavirus (Goncalves <i>et al.</i> , 2005), antimicrobial intestinal adhesion (Coutino <i>et al.</i> , 2001)	Limonene, copaene, Asiatic acid, β -carotene, morin-3-O- α -L-arabinopyranoside, avicularin, gajiverin, quercitin ellargic acid (Gutierrez <i>et al.</i> , 2008)
<i>Syzygium cordatum</i> Hochst. Ex. C. Krauss	Umdoni, Mutu (V)	Leaves, bark	Respiratory disorders, tuberculosis, stomach complaints, emetics, diarrhoea, cold, fever (Verschaeve and Van Staden, 2008)	Mutagenicity , antimutagenicity (Verschaeve and Van Staden, 2008); antimycobacterium (Mativandela <i>et al.</i> , 2008); Antiescherichia (Sibandze <i>et al.</i> , 2010)	-
<i>Syzygium paniculatum</i> Gaertner	-	-	-	--	-
Olacaceae					
<i>Ximenia caffra</i> Sond	Mutswili (V)	leaves	Diarrhoea and dysentery (Green <i>et al.</i> , 2010; Fabry <i>et al.</i> , 1996)	Antigardial (John <i>et al.</i> , 1995), Antiamoebic (Samie <i>et al.</i> , 2009),	

<i>Punica granatum</i> L.	iRhamathi (X)	Fruit rind, roots	Diarrhoea and dysentery (van Wyk, 2008; Dold and Cocks, 2000)	antifungal (Samie <i>et al.</i> , 2010) Antidiarrhoeal (Pillai, 1992; Qnais <i>et al.</i> , 2007), Anti-inflammatory (Lansky and Newman, 2007)	Ellagitannins, anthocyanins, flavone glucosides, flavones, flavonol, flavonol glucosides, hydroxycinnamic acid, hydroxybenzoic acid, flavan-3-ols, alkaloids, sterol, triterpenoids (Lansky and Newman, 2007)
Oleaceae					
<i>Olea europaea</i> L. Subsp <i>africana</i> (Mill.) P.S.Green	Uzintwa (X), uMnquma (X)		Anti-hypertensive, diuretic, tonic, diarrhoea, sore throat (Amabeoku and Bamuamba, 2010)	Antidiarrhoeal (Amabeoku and Bamuamba, 2010)	-
Orchidaceae					
<i>Polystachya ottoniana</i> Rchb.f.			Diarrhoea (Chinsamy <i>et al.</i> , 2010)	-	-
Pedaliaceae					
<i>Ceratotheca triloba</i> (Bernh.) Hook	Udonqabathwa (Z)	Leaf	Infusion for diarrhoea and gastrointestinal cramps (Watt and Breyer-Brandwijk, 1962; Roberts, 1990)	5-lipoxygenase inhibitory and antioxidant (Akula and Odhav, 2008), α -amylase inhibitory (Odhav <i>et al.</i> , 2010).	-
Plantaginaceae					
<i>Plantago major</i> L.		Seed, root	Diarrhoea (van Wyk, 2008)	Antidiarrhoeal (Atta and Mounair, 2005), Hepatoprotective and anti-inflammatory (Turel <i>et al.</i> , 2007), wound healing activity, anti-inflammatory, analgesic, antioxidant, weak antibiotic, immuno modulating and antiulcerogenic activity (Samuelsen, 2000)	Indicain, plantagonin, baicalein, hispidulin, plantagin, aucubin, fumaric acid, syringic acid, vanillic acid, <i>p</i> -hydroxy benzoic acid, ferulic acid, <i>p</i> -coumaric acid, gentisic acid, salicylic acid, benzoic acid, cinnamic acid oleanolic acid, ursolic acid, 18 β -glycyrrhetic acid and sitosterol (Samuelsen, 2000)
<i>Plantago lanceolata</i> L.		Seed	Diarrhoea (van Wyk, 2008)	-	-
Plumbaginaceae					
<i>Plumbago auriculata</i>	uTshintshini	Roots	Diarrhoea in cow (Dold and Cocks, 2001; McGaw <i>et al.</i> , 2008)	Antibacterial and anti cancer (Bisi-Johnson <i>et al.</i> , 2011)	-
Polygonaceae					
<i>Rumex lanceolatus</i> Thunb	Idololenkonyane (Z), Idolonyana (X)	Roots	Infantile diarrhoea, tapeworm, wound and sores (Dold and Cocks, 2000)	-	-
<i>Rumex obtusifolius</i>	Idololenkonyane (X, Z)	leaves	Diarrhoea (Bisi-Johnson <i>et al.</i> , 2010)	-	-
Portulacaceae					
<i>Portulacaria afra</i> Jacq.	Umdibili (Z)	Leaves	Diarrhoea (Mlambo, 2008)	-	-
Proteaceae					
<i>Protea caffra</i> Meisn.	Tshidzungu (V)	Root bark decoction	Calves with bloody diarrhoea (Hutching <i>et al.</i> , 1996)	-	-



<i>Protea nitida</i> Mill.		Bark	Astringent for diarrhoea (van Wyk, 2008)	-	-
<i>Protea simplex</i>		Root, bark	Decoction and infusion for diarrhoea, dysentery, stomach pain in human (Fawole et al., 2009 ; Hutching et al., 1996)	Anti-inflammatory (Fawole et al., 2009 Ndhalala), Antimicrobial and mutagenicity (Fawole et al., 2009 Finnie)	-
<i>Protea welwitschii</i> Engl.			Dysentery, diarrhoea in calves and humans (Watt and Breyer-Brandwijk, 1962 ; McGaw et al., 2008)	-	-
Punicaceae					
<i>Punica granatum</i> L.	Mokgranata	Root	Decoction for diarrhoea (Mathabe et al., 2006)	-	-
Rhamnaceae					
<i>Ziziphus mucronata</i> Willd.	Mukhalu, Mutshetshete (V)	Leaves, bark, roots	Boils, sores, grandular swelling, diarrhoea, dysentery, cough (Verschaeve and Van Staden, 2008 ; Green et al., 2010)	Anti-inflammatory (Fawole et al., 2009 Ndhalala), Antimicrobial and mutagenicity (Fawole et al., 2009 Finnie)	-
<i>Ziziphus zeyheriana</i> Sond.		Root-stock	Diarrhoea, internal parasites, general ailments (Van der Merwe et al., 2001 ; McGaw et al., 2008)	-	-
Rosaceae					
<i>Prunus africana</i> (Hook.f) Kalkman Red Stinkwood	Umkhakhazi (X), Umkakase (X)	Root	Diarrhoea, abdominal ailments (Bisi-Johnson et al., 2010)		
<i>Prunus persica</i> (L.) Batsch.	Ipesika	Leaf decoctions	Diarrhoea in lamb and kid goats (Dold and Cocks, 2001 ; McGaw et al., 2008)		
Rubiaceae					
<i>Breonadia salicina</i> (Vahl) Hepper & J. R. I. Wood		Bark decoctions	Diarrhoea, bloody stool, colic (Neuwinger, 1996 ; Venter and Venter, 2002)	Antiescherichial (Sibandze et al., 2010)	
<i>Nauclea latifolia</i> Smith		Root	Dysentery, dyspepsia, fever, gastritis, gonorrhoea, malaria, leprosy, measles, piles, toothache (Reid et al., 2006)	Antiamoebic (Tona et al., 1998 ; Moundipa et al., 2005); antidiarrhoeal (Owolabi et al., 2010)	
<i>Pentanisia prunelloides</i> (Klotzsch exEckl. & Zeyh) Walp	Icishamlilo, Icimamilo (X, Z)	Root, leaves, bulb	Vomiting, diarrhoea in children(Bisi-Johnson et al., 2010)	Antibacterial (Yff et al., 2002)	Palmitic acid (Yff et al., 2002)
<i>Psychotria capensis</i> (Eckl.) Vatke	Ishithitibala (Z), UmGono-gono (X)	Fruits	Diarrhoea and vomiting (Bisi-Johnson et al., 2010)	-	-
<i>Rubia petiolaris</i> DC.		Root	Diarrhoea and dysentery (van Wyk, 2008)	-	-
<i>Rubus pinnatus</i> Willd.	iQunube	Roots	Diarrhoea, haemorrhoids, epilepsy (van Wyk, 2008)	-	-
<i>Rubus rigidus</i>		Root	Diarrhoea and dysentery, toothache,	-	-

			coughs and colds (Iwalewa et al., 2007).		
<i>Vangueria infausta</i> Burch. subsp. <i>infausta</i>	Umviyo		Diarrhoea (de Wet <i>et al.</i> , 2010)	Antibacterial and antifungal (de Boer et al., 2005)	-
Rutaceae					
<i>Agathosma betulina</i> (Bergius) Pillans			Antispasmodic, antipyretic, cough, Kidney and urinary tract infection, cholera and stomach ailment (Molla and Viljoen, 2008)	Antidiarrhoea and antibacterial (Lis-Balchin et al., 2001); anti-inflammatory and antioxidant (Steenkamp et al., 2006)	-
<i>Agathosma crenulata</i> (L.) Pillans			Antispasmodic, antipyretic, cough, Kidney and urinary tract infection, cholera and stomach ailment (Molla and Viljoen, 2008)	Antidiarrhoea and antibacterial (Lis-Balchin et al., 2001)	-
Rutaceae					
<i>Clausena anisata</i> (Willd.) Hook.F. Ex Benth.		Bark infusion	Dysentery in cattle (Hutching et al., 1996)	-	-
<i>Ruta graveolens</i> L.	iVendrit (X)	Leaves	Fever, convulsion, epilepsy, diarrhoea, cardiac asthma, jaundice (Dold and Cocks, 2000)	-	-
Sapindaceae					
<i>Hippobromus pauciflorus</i> (L.f.) Radlk.	Ulwathile (X)	Bark, root, leaves	Heartwater and diarrhoea in cattle (Dold and Cocks, 2001; McGaw <i>et al.</i> , 2008), Diarrhoea and dysentery (Bisi-Johnson et al., 2010)	Acute toxicity (Pendota et al., 2010), anti inflammatory, analgesic antipyretic (Pendota et al., 2009)	-
Scrophulariaceae					
<i>Physalis peruviana</i> L.	Igquzu (X)	Leaves	Stomach disorder (Bisi-Johnson et al., 2010)	-	-
<i>Physalis viscosa</i> L.	Umqumqumu (Z)	Leaves	Diarrhoea (Mlambo, 2008)	Antibacterial (Ovenden et al., 2004)	Physaloside A (Ovenden et al., 2004)
<i>Jamesbrittenia atropurpurea</i> (Benth.) Hilliard		Herb	Antispasmodic, stimulant; convulsions; cough; bronchitis (van Wyk, 2008)	-	-
<i>Xanthium strumarium</i> L.		Root	Cancer, dysentery, catarrh, leprosy (Watt and Breyer-Brandwijk, 1996 ; Fouche et al., 2008)	Anti-inflammatory and analgesic (Han et al., 2007)	1-O-caffeoylquinic acid, 3-O-caffeoylquinic acid, chlorogenic acid, 4-O-caffeoylquinic acid, cynarin, 1,4-O-dicaffeoylquinic acid, 1,5-O-dicaffeoylquinic acid, 1,5-O-dicaffeoylquinic acid, 1,3,5-O-tricaffeoylquinic acid, 3,4,5-O-tricaffeoylquinic acid (Han et al., 2007)
Solanaceae					
<i>Solanum aculeastrum</i> Dun	umthuma (X, Z)	Fruit, root, leaves	Fruit decoction for haemorrhoids, dysentery, fruit as enema for diarrhoea (Bisi-Johnson et al., 2010)	Antimicrobial (Koduru et al., 2006); Anticancer (Koduru et al., 2007)	tomatidine and solasodine (Koduru et al., 2007)
<i>Solanum incanum</i> L.	uMthuma, <i>intfuma</i> (S)	Root	Root infusion for back ache (Amusan, 2007)	-	-

<i>Solanum mauritianum</i>	Umtotovane (Z)	Leaf	Infusion for dysentery and diarrhoea (Watt and Breyer-Brandwijk, 1962)	-	-
<i>Solanum panduriforme</i> E. Mey	Thuthula	Fruit sap	Diarrhoea (Van der Merwe <i>et al.</i> , 2001; McGaw <i>et al.</i> , 2008)	-	-
<i>Solanum supinum</i> Dun.	Thola (S)	Root	Decoction for diarrhoea (Mathabe <i>et al.</i> , 2006)	-	-
Sterculiaceae					
<i>Withania somnifera</i> (L.) Dun	uBuvimba		Fever, cold and flu, abdominal discomfort, diarrhoea, worms sedative and hypnotic (van Wyk and Gericke, 2000; Fouche <i>et al.</i> , 2008)	Anti-inflammatory, antitumor, immunomodulatory (Mishra <i>et al.</i> , 2000); antichorelae (Acharya <i>et al.</i> , 2009)	Isopelletierine, anferine, withanolides, withaferins, sitoindosides (Mishra <i>et al.</i> , 2000)
<i>Dombeya rotundifolia</i> (Hochst.) Planch.	Tshiluvhari (V)	Root, bark, wood	Internal ulcers, haemorrhoids, diarrhoea, stomach problems, nausea, chest pain (Verschaeve and Van Staden, 2008)	Mutagenicity , antimutagenicity	-
<i>Hermannia incana</i> Cav	Mavulakuvaliwe	leaves	Crushed with cold water and taken orally for diarrhoea (Appidi <i>et al.</i> , 2008)	Toxicological assay (Appidi <i>et al.</i> , 2009); antimicrobial, anti-inflammatory, antioxidant and cytotoxicity (Essop <i>et al.</i> , 2008)	-
<i>Waltheria indica</i> L.		Whole plant	Decoction for diarrhoea (Mathabe <i>et al.</i> , 2006)	Antibacterial, antifungal and antiviral (Maregesi <i>et al.</i> , 2008)	-
Strychnaceae					
<i>Strychnos madagascariensis</i> Pior.	Umkwakwa, Mukwakwa (V)		Diarrhoea (de Wet <i>et al.</i> , 2010)	-	-
Urticaceae					
<i>Pouzolzia mixta</i> solms	Muthanzwa	Root, leaves	Dysentery (Verschaeve and Van Staden, 2008); diarrhoea (Samie <i>et al.</i> , 2010)	-	-
Verbenaceae					
<i>Clerodendrum glabrum</i> E. Mey	Umqangazani Uqangazana (X), iNunkisiqaga (X) Umqangazane	leaves	Bloody stool, chest infections (Bisi-Johnson <i>et al.</i> , 2010)	-	--
Viscaceae					
<i>Viscum capense</i> L. F.	Iphakama (Z)		Diarrhoea (Forbes, 1986; Van Wyk <i>et al.</i> , 2008)	-	-
Vitaceae					
<i>Lippia javanica</i> (Burm.f.) Spreng	Musudzungwane (V)	Leaf infusion	Prophylactics against dysentery, diarrhoea and malaria (Mabogo, 1990; Fouche <i>et al.</i> , 2008)	-	-



<i>Rhoicissus tridentata</i> (L.F.) Wild & Drums.	Umthwazi (Z), Murumbulamudzana (V)	Tuber decoction	Diarrhoea in goat and sheep (Dold and Cocks, 2001; McGaw <i>et al.</i> , 2008)	Antispasmodic (Katsoulis et al., 2000)	-
<i>Cyphostemma cirrhosum</i> (Thunb.)	Udekane (Z)	Leaves	Diarrhoea (Mlambo, 2008)	-	-
Zingiberaceae					
<i>Aframomum latifolium</i> (Afzel.) K. Schum		Leaves	Decoction for diarrhoea (De Villiers et al., 2010)	-	-
<i>Elytropappus rhinocerotis</i> (L.f) Less.		Twigs	Bitter for dyspepsia, indigestion, diarrhoea (van Wyk, 2008)	-	-
					<i>ent-15β-seneciolyoxy-16,17-epoxy-kauran-18-oic acid</i>

V=Vhavenda, Z=Zulu, X=Xhosa, S=Swazi

Appendix 9.1: 1D and 2D NMR spectra data of Ursolic acid

Peak number	Hydrogen	¹³ C/DEPT	HSQC	HMBC	LITERATURE
1	1.46-1.6	CH ₂	37	15(C25), 27(C2), 56(C5), 78(C3)	39.2
2	1.38-1.5	CH ₂	27	37(C1), 56(C5), 78(C3)	28.1
3	3.0	CH	78	16(C23), 27(C2)	78.2
4	-	C	39	-	39.6
5	0.64	CH	56	16(C23), 18 (C6), 37(C1)	55.9
6	1.26, 1.44	CH ₂	18	56(C5)	18.8
7	1.24, 1.4	CH ₂	33	56(C5)	33.7
8	-	C	39	-	40.1
9	1.42	CH	48	15(C25), 23(C11), 37 (C1), 38.8(C10), 39(C8)	48.1
10	-	C	37	-	37.5
11	1.76-1.88	CH ₂	23	39 (C8), 125(C12), 140(C13)	23.7
12	5.2	CH	125	42(C14), 48(C9), 53(C18)	125.7
13	-	C	140	-	139.3
14	-	C	42	-	42.6
15	0.95, 1.4	CH ₂	28	24(C16), 48(C17)	28.8
16	1.5, 1.9	CH ₂	24	28 (C15), 42(C14), 48(C17), 53(C18), 178(C28)	25
17	-	C	48	-	48.1
18	2.08	CH	53	17.5(29), 24(16), 38.8(19), 42(14), 37(C20), 125(12), 140(13), 178(28)	53.6
19	0.9	CH	38.6	39(C19), 37(C20)	39.5
20	1.29	CH	37		39.4
21	1.48-1.58	CH ₂	30		31
22	0.87, 1.48	CH ₂	38.9	24(C16), 53(C18)	37.4
23	0.64	CH ₃	16	29(C24), 39(C4), 56(C5), 78(C3)	16.5
24	0.87	CH ₃	29	16(C23), 78(C3)	28.8
25	0.85	CH ₃	15	37(C1), 56(C5)	15.7
26	0.72	CH ₃	17.3	33(C7), 39(C8), 42(C14), 48(C9)	17.5
27	1.02	CH ₃	23.4	28(C15), 39(C8), 42(C14), 140(C13)	24
28	-	C	178	-	179.7
29	0.78	CH ₃	17.5	38.8(C19), 53(C18)	17.5
30	0.90	CH ₃	22	31(C21), 48(C20)	21.4

Appendix 9.2: 1D and 2D NMR spectra data of mixture of corosolic acid and maslinic acid

Peak number	¹ H	¹³ C/DEPT	HSQC (Corosolic acid)	HSQC (Maslinic acid)	HMBC (H→C)	LITERATURE
1	1.7-1.8, 0.73-0.79	CH ₂	47.75	47.75	16.92(C25), 38.27(C10), 55.35(C5), 68(C2), 83.49(C3)	46.8
2	3.4	CH	68.41	68.41	83.49(C3)	68.9
3	2.7	CH	83.49	83.49	68.41(C2), 29.87(C24)	83.8
4	-	C	39.61	39.61	-	39.1
5	0.68-0.75	CH	55.35	55.35	39.61(C4), 38.25(C10), 29.43(C24)	55.4
6	1.41-1.47, 1.26-1.33	CH ₂	18.71	18.71	38.12(C10), 39.61(C4)	18.4
7	1.36-1.45, 1.56-1.61	CH ₂	33	33	17.49(C26), 55.35(C5), 18.71(C6)	32.9
8	-	C	39.75	39.75	-	39.6
9	1.5	CH	47.53	47.53	23.74 (11) (24.01)	47.5
10	-	C	38.25	38.25	-	38.3
11	1.8-1.9, 1.43-1.47	CH ₂	23.74	24.01	47.53(C9), 125.55(C12) (122.72), 139.43(C13) (145.09)	23.4
12	5.11	CH	125.55	122.72	23.74(C11) (24.01), 42.19(C14) (41.92), 47.53(C9), 53.05(C18) (41)	125.3
13	-	C	139.42	145.09	-	138.1
14	-	C	42.19	41.92	-	42.1
15	0.92-0.99, 1.75-1.79	CH ₂	28.21	28.21	39.57(C8), 23.60(C27) (26.8), 42.19(C14) (41.92)	28.0
16	1.26-1.33, 1.47-1.52	CH ₂	24.51	27.94	178 (C28) (179.24), 28.21(C15), 53.05(C18), (41), 48.13(C17)	24.3
17	-	C	48.33	48.33	-	48.1
18	2.09 (2.7)	CH	53.05	41	17.75(C29), 24.51(C16) (27.94), 39.10(C19) (27.94), 42.19(C14) (41.92), 48.13(C17), 37(C22), 125.55(C12) (122.72), 139.42(C13) (145.09), 178 (C28) (179.24)	52.8
19	0.88-0.92	CH	39.07	27.94 (CH ₂)		39.1
20	1.25-1.31	CH	39.02 (CH)	46.13 (C)		38.9
21	1.48-1.58	CH ₂	30.6	37	24, 31	30.7
22	1.58, 1.00-1.09	CH	37	47.53	178(C28) (179.24), 24.51(C16) (27.94)	36.7
23	0.90	CH ₃	29.2	29.2	17.76(C24), 55.35(C5), 39.61(C4)	28.7
24	0.68	CH ₃	17.76	17.76	29.43(C23), 39.61(C4), 55.35(C5)	17.0
25	0.88	CH ₃	16.94	16.94	38.25(C10), 47.53(C9)	17.0
26	0.72	CH ₃	17.4	17.4	47.53(C9), 42.19(C14) (41.92), 33.01(C7)	17.0
27	1.00	CH ₃	23.60	26.8	28.21(C15), 39.75(C8), 42.19(C14) (41.92), 139.42(C13) (145.09)	23.7
28	-	C	178	179.70	-	177.9
29	0.79	CH ₃	17.5	17.5	39.07(C19) (27.94), 53.05(C18) (41)	17.0
30	0.88	CH ₃	21.6	21.6	39.02(C20) (46.13)	21.2

Appendix 9.3: 1D and 2D NMR spectra data of mixture of asiatic acid and arjunolic acid

Peak number	¹ H	¹³ C/DEPT	HSQC (Asiatic acid)	HSQC (Arjunolic acid)	HMBC (H→C)	LITERATURE
1	0.69, 1.73	CH ₂	48.31	48.31	18.04(C25), 68.67(C2), 76.50(C3)	46.8
2	3.45	CH	68.67	68.67	76.50(C3)	68.9
3	3.13	CH	76.50	76.50	68.67(C2), 14.30(C24), 65.01(C23), 43.62(5)	83.8
4	-	C	43.62	43.62	-	39.1
5	1.14	CH	47.27	47.27	76.50(C3), 47.27(C4), 65.01(C23), 14.30(C24), 33.40 (C7), 18.04 (C25)	55.4
6	1.33, 1.18	CH ₂	18.03	18.03	38.12(C10), 39.61(C4)	18.4
7	1.44, 1.20	CH ₂	33.40	33.40	40.49 (C8)	32.9
8	-	C	40.49	40.49	-	39.6
9	1.5	CH	47.51	47.54	23.74 (11) (24.01)	47.5
10	-	C	38.40	38.40	-	38.3
11	1.78, 1.42	CH ₂	23.00	24.01	125.19(C12) (122.72), 138.92(C13) (145.09)	23.4
12	5.09 (5.12)	CH	125.19	122.28	42.21(C14) (41.92), 47.51(C9) (47.54), 52.62(C18) (41.73)	125.3
13	-	C	138.71	144.68	-	138.1
14	-	C	42.21	41.92	-	42.1
15	1.73, 1.58	CH ₂	28.07	28.21	39.57(C8), 23.60(C27) (26.8), 42.19(C14) (41.92)	28.0
16	1.88, 1.48	CH ₂	24.31	27.94	179.13 (C28) (179.24),	24.3
17	-	C	47.99	48.33	-	48.1
18	2.05 (2.72)	CH	52.62	41.73	18.16(C29), 24.31(C16) 39.44(C19) (27.94), 42.21 (C14) (42.52), 47.99(C17) (47.01), 125.19(C12) (122.28), 138.92(C13) (144.68), 179.13 (C28) (179.24)	52.8
19	0.88-0.92	CH	39.44	24.31(CH ₂)		39.1
20	1.25-1.31	CH	38.90(CH)	47.27 (C)		38.9
21	1.38, 1.20	CH ₂	31.42	31	24, 31	30.7
22	1.55, 1.39	CH ₂	33	33	24.31(C16), 31.61 (C21)	36.7
23	3.25, 3.00	CH ₂	65.01	65.01	14.91(C24), 43.62(C5), 47.27(C4), 76.50 (C3)	28.7
24	0.50	CH ₃	14.91	14.91	65.01(C23), 47.27(C4), 43.62(C5)	17.0
25	0.65	CH ₃	18.04	18.04	40.49(C10), 43.62(C5), 48.31(C1)	17.0
26	0.68	CH ₃	17.04	17.04	47.53(C9), 42.19(C14) (41.92), 33.18 (C7)	17.0
27	0.99 (1.04)	CH ₃	23.40	26.8	28.48(C15), 40.49(C8) (39.96), 42.21(C14) (42.57), 138.92(C13) (144.68)	23.7
28	-	C	179.13	179.24	-	177.9
29	0.77 (0.82)	CH ₃	18.16	26.0	39.44(C19) (27.94), 53.05(C18) (41)	17.0
30	0.87 (0.82)	CH ₃	23.0	33	39.02(C20) (46.13)	21.2

Appendix 9.4: 1D and 2D NMR spectra data of combretastatin B5-2'-O- glucopyranoside

Peak number	¹ H	¹³ C/DEPT	HSQC ^a	HMBC (H→C)	LITERATURE DATA HSQC ^a
1		C	132.72		132.43
2	6.5	CH	106.26	C1, C3, C6, C1a	105.96
3		C	144.35		144.77
4		C	133.79		133.44
5		C	148.25		147.77
6	6.5	CH	106.26	C1, C5, C2, C1a	105.96
1a	2.7	CH ₂	36.96	C1, C2, C1a', C1'	36.58
1a'	2.9, 3.0	CH ₂	31.73	C1, C1', C2', C6', C1a	31.35
1''		C	128.55		128.22
2''		C	144.35		143.95
3''		C	139.71		139.29
4''		c	147.26		146.90
5''	6.7	CH	109.40	C1', C3', C4', C6'	109.10
6''	6.6	CH	118.95	C2', C4', C1a'	118.68
1'''	4.5	CH ₂	106.16	C2'', C2'', C5''	105.79
2'''	3.3	CH	74.45	C1'', C3''	74.04
3'''	3.2	CH	76.62	C4'', C5''	76.14
4'''	3.2	C	70.15	C3'', C5''	69.71
5'''	3.2	CH	77.85	C3''	77.44
6'''	3.5, 3.7	CH ₂	61.34	C4'', C5''	60.89
3-OCH ₃	3.5	CH ₃	56.34	C3	56.0
4-OCH ₃	-	CH ₃	-		-
5-OCH ₃	3.5	CH ₃	56.34	C5	56.0
4'-OCH ₃	3.5	CH ₃	56.30	C4'	56.0

^a Pelizzoni Francesca, 1994: Combretastatin derivatives with antitumoral activity and process for the preparation thereof. Patent Cooperation Treaty (PCT), WO 94/05682, CO7H 15/203, CO7C 43/23, A61K 31/70, 31/085

Appendix 9.5: 1D and 2D NMR spectra data of combretastatin B1-2'-O- glucopyranoside

Peak number	¹ H	¹³ C/DEPT	HSQC ^b	HMBC (H→C)	LITERATURE DATA HSQC ^b
1		C	138.54		138.10
2	6.5	CH	105.60	C1, C1a, C3, C4	105.63
3		C	152.69		152.65
4		C	135.53		135.45
5		C	152.72		152.65
6	6.5	CH	105.60	C1, C1a, C4, C5	105.63
1a	2.69, 2.79	CH ₂	36.92	C1, C1', C2, C6, C1a'	36.91
1a'	2.92, 3.20	CH ₂	31.34	C1a, C1', C2', C6'	31.18
1''		C	128.27		128.03
2''		C	143.78		143.93
3''		C	139.13		139.35
4''		c	147.02		146.95
5''	6.7	CH	108.54	C1', C3', C4',	108.99
6''	6.6	CH	119.24	C1a', C2', C4'	118.56
1'''	4.5	CH ₂	105.69		105.77
2'''	3.3	CH	74.21		74.09
3'''		CH	76.63		76.26
4'''		C	69.72		69.76
5'''	3.2	CH	77.03		77.51
6'''	3.5, 3.7	CH ₂	60.97		60.96
3-OCH ₃	3.5	CH ₃	55.14	C3	55.78
4-OCH ₃	3.5	CH ₃	59.69	C4	60.04
5-OCH ₃	3.5	CH ₃	55.34	C5	55.78
4'-OCH ₃	3.5	CH ₃	55.42	C4'	55.94

^b Pelizzoni Francesca, 1994: Combretastatin derivatives with antitumoral activity and process for the preparation thereof. Patent Cooperation Treaty (PCT), WO 94/05682, CO7H 15/203, CO7C 43/23, A61K 31/70, 31/085

Appendix 9.6: 1D and 2D NMR spectra data of 3 β -ethoxy sitosterol

Peak number	¹ H	¹³ C/DEPT	HSQC	HMBC (H→C)	LITERATURE
1	1.036, 1.817	CH ₂	37.47		37.3
2	2.233	CH ₂	31.8	36.73 (C10), 72.04 (C3), 121.93 (C6), 140.97 (C5)	31.9
3	3.502	CH	72.04		71.8
4		CH ₂	40.00		40.5
5	-	C	140.97		140.7
6	5.33	CH	121.93	32.12 (C7), 36.73 (C10)	121.7
7	1.427	CH ₂	32.12		31.9
8		CH	29.91		31.6
9		CH	50.35		50.2
10	-	C	36.73		36.5
11		CH ₂	21.30		21.1
12	2.194	CH ₂	40		39.8
13	-	C	42.54		42.3
14	1.058	CH	56.99		56.8
15	0.988	CH ₂	24.52		24.3
16	1.234	CH ₂	28.47		28.3
17	1.058	CH	56.27		56.1
18	0.988	CH ₃	12.08	140.97 (C5), 37.47 (C1), 50.35 (C9)	11.9
19	0.659	CH ₃	20.03	56.99 (C14), 56.23 (C17), 42.52 (C12), 42.54 (C13),	19.5
20	1.326	CH	36.36		36.2
21	0.904	CH ₃	19.24	56.27 (C17), 36.36 (C20), 34.16 (C22)	18.9
22		CH ₂	34.16		33.9
23		CH ₂	26.29		26.1
24		CH	46.05		45.8
25		CH	29.37		29.1
26	0.988	CH ₃	19.61		19.4
27	0.796	CH ₃	19.00	29.37 (C25), 46.05 (C24), 19.61 (C26)	19.1
28		CH ₂	23.28		23.1
29	0.822	CH ₃	12.20	23.25 (C28)	12.0
CH ₃ CH ₂		CH ₃			
CH ₃ CH ₂		CH ₂	76.86		

Appendix 9.7: 1D and 2D NMR spectra data of Quercetin

Peak number	¹ H	¹³ C/DEPT	HSQC	HMBC (H→C)	LITERATURE
2		C	147.96	-	146.8
3		C	137.23	-	135.8
4		C	177.31	-	175.9
5		C	162.50	-	160.8
6	6.17	CH	99.28	162.50 (C5), 104.51 (C10), 94.39 (C8)	98.2
7		C	165.55	-	163.9
8	6.37	CH	94.39	165.55 (C7), 158.21 (C9), 104.51 (10), 99.28 (C6)	93.4
9		C	158.21	-	156.2
10		C	104.51	-	1103.0
1'		C	124.13	-	122.0
2'	7.87	CH	116.30	147.94 (C2), 144 (C3'), 121 (C6')	115.1
3'		C	147.96	-	145.1
4'		C	148.75	-	147.7
5'	6.88	CH	116.78	144 (C3'), 121 (C6')	115.6
6'	7.72	CH	121.65		120.6

Appendix 9.8: 1D and 2D NMR spectra data of Myricetin

Peak number	¹ H	¹³ C/DEPT	HSQC	HMBC (H→C)	LITERATURE

2		C	146.57	-	148.2
3		C	135.92	-	137.5
4		C	175.86	-	177.5
5		C	161.05	-	162.6
6	6.17	CH	97.80	164.17 (C7), 161.05 (C5), 103.06 (C10), 99 (C6)	99.5
7		C	164.17	-	165.8
8	6.37	CH	92.95	164.17 (C7), 158 (C9), 103.06 (C10), 99 (C6)	94.6
9		C	156.76	-	158.4
10		C	103.06	-	104.7
1'		C	121.65	-	123.3
2'	7.38	CH	107.10	145.29 (C3'), 135.51 (C4'), 121.65 (C1'), 107.10 (C6')	108.8
3'		C	145.29	-	146.9
4'		C	135.51	-	137.1
5'		C	145.29	-	146.9
6'	7.38	CH	107.10	145.29 (C5'), 135.51 (C4'), 121.65 (C1'), 107.10 (C2')	108.8

Appendix 9.9: 1D and 2D NMR spectra data of Isoetin 2' methyl ether/ Isoetin 4' methyl ether

Peak number	¹ H	¹³ C/DEPT		HMBC (H→C)	LITERATURE	
2	-	C	163.09	C3, C6'	163.2 ^a	161.70 ^b
3	7.2(s)	CH	108.85	C6'	108.7	106.77
4	-	C	184.24		183.5	181.74
5	-	C	163.67		162.7	161.41
6	6.2(d, J=)	CH	99.97		99.8	98.48
7	-	C	166.22		165.3	163.88
8	6.4 (d, J=)	CH	94.86	C6	95.1	93.48
9	-	C	159.43		159	157.19
10	-	C	105.09	C3, C8	105.1	103.46
1'	-	C	110.17	C3'	108.9	106.97
2'	-	C	153.27	OCH ₃ , C3', C6'	154.2	150.50
3'	6.65 (s)	CH	101.36		105.4	104.20
4'	-	C	153.27	OCH ₃ , C3', C6'	152.8	151.60
5'	-	C	140.86	C3', C6'	142.8	138.73
6'	7.38 (s)	CH	114.44		112.6	113.44
OCH ₃	3.8 (s)		56.34 at C2' or C4'		57.6 at C5'	-

^aIsoetin 5'methyl ester (AbdurRahman and Moon, 2007), ^bisoetin (Gluchoff-Fiasson *et al.*, 1991)

Appendix 9.10: 1D and 2D NMR spectra data of Quercetin-3-O- β -galactopyranoside

Peak number	^1H	$^{13}\text{C}/\text{DEPT}$	HSQC	HMBC (H \rightarrow C)	LITERATURE
2	-	C	156.76	-	158.3
3	-	C	133.85	-	135.8
4	-	C	177.83	-	179.4
5	-	C	161.42	-	163.0
6		CH	99.17	162.50 (C5), 104.51 (C10), 94.39 (C8)	99.8
7	-	C	164.58	-	166.0
8		CH	94.09	165.55 (C7), 158.21 (C9), 104.51 (10), 99.28 (C6)	94.7
9	-	C	156.80	-	158.8
10	-	C	102.36	-	104.2
1'	-	C	122.14	-	123.2
2'		CH	115.66	147.94 (C2), 144 (C3'), 121 (C6')	117.8
3'	-	C	144.97	-	145.8
4'	-	C	148.75	-	149.9
5'		CH	116.47		116.1
6'		CH	121.42		122.9
1''	5.18 (d)	CH	104.24	133.85(C2)	105.4
2''	3.83 (t)	CH	71.55	104.12 (C1''), 73.70 (C3'')	73.2
3''	3.56 (m)	CH	73.40	104.12 (C1''), 71.87 (C2''),	75.1
4''	3.87 (s)	CH	68.30	144 (C3'), 121 (C6')	70.0
5''	3.94 (t)	CH	75.96		77.2
6''	3.66 (dd), 3.5 (m)	CH ₂	60.45	75.80 (5''), 68.61 (C4'')	61.9

Appendix 9.11: 1D and 2D NMR spectra data of Myricetin-3-O- β -galactopyranoside

Peak number	^1H	$^{13}\text{C}/\text{DEPT}$	HSQC	HMBC (H \rightarrow C)	LITERATURE
2		C	156.95	-	156.4
3		C	134.56	-	135.4
4		C	177.97	-	177.6
5		C	161.54	-	161.4
6	6.19	CH	98.47	164.70 (C7), 161.54 (C5), 104.12 (C10), 93.25 (C8)	98.7
7		C	164.70	-	164.4
8	6.38	CH	93.25	164.70 (C7), 157.24(C9), 104.12 (C10), 98.47 (C6)	93.4
9		C	157.24	-	156.4
10		C	104.14	-	103.9
1'		C	120.26	-	120.0
2'	7.37	CH	108.52	156.95 (C2), 144.95 (C3'), 136.71 (C4'), 120.26 (C1'), 108.52 (C6')	108.6
3'		C	144.95	-	145.5
4'		C	136.71	-	136.8
5'		C	144.95	-	145.5
6'	7.37	CH	108.52	156.95 (C2), 144.95 (C5'), 136.71 (C4'), 120.26 (C1'), 108.52 (2')	108.6
1''	5.18 (d)	CH	104.12	134.56 (C2)	105.4
2''	3.83 (t)	CH	71.87	104.12 (C1''), 73.70 (C3'')	73.2
3''	3.56 (m)	CH	73.70	104.12 (C1''), 71.87 (C2''),	75.1
4''	3.87 (s)	CH	68.61		70.0
5''	3.94 (t)	CH	75.80		77.2
6''	3.66 (dd), 3.5 (m)	CH ₂	60.52	75.80 (5''), 68.61 (C4'')	61.9