



Chapter 3

Cylindrocladium pauciramosum,
the dominant fungal pathogen
in *Eucalyptus* clonal propagation
nurseries in South Africa



Abstract

The implementation of clonal forestry in South Africa has resulted in substantial changes to the *Eucalyptus* nursery system. Thus, large clonal nurseries needed to be established. In these systems, fungal pathogens can be responsible for losses during the rooting of cuttings. The aim of this study was to identify the dominant fungal pathogens responsible for these losses in South African *Eucalyptus* clonal nurseries. A survey was conducted at four of the major nurseries that produce cuttings and which are located in KwaZulu Natal. At each nursery, five of the most commonly produced hybrid clones were selected for sampling. Cuttings were collected at each nursery and immediately examined for symptoms and then placed in moist chambers to enhance sporulation of fungi. Several well-known fungal pathogens were collected and amongst these, *Cylindrocladium* isolates were dominant. All *Cylindrocladium* isolates were identified as *C. pauciramosum* based on morphological characteristics and this was confirmed using DNA sequence data comparisons based on part of the β -tubulin gene. The pathogenicity of *C. pauciramosum* was tested on two – month – old plants representing four commercial hybrid clones. The pathogen gave rise to infection on all clones inoculated with little difference noted between these clones.

Introduction

During the course of the past two decades, the *Eucalyptus* nursery system in South Africa has undergone major changes to accommodate the implementation of clonal forestry. This forestry system involves the utilization of a limited number of selected clones and hybrids propagated by means of vegetative cuttings or tissue culture (Denison & Kietzka 1993b, White 1995). The result is large-scale uniform plantations of selected tropical and sub – tropical *Eucalyptus* clones and interspecific clonal hybrids (Kulkarni & Lal 1995, Wilson 1998). Generally, clones and hybrids are intensively selected for growth and form, disease resistance, wood properties and adaptation to environmental and climatic conditions (Denison & Kietzka 1993a, Aimers – Halliday *et al.* 1999).

Eucalyptus hybrid clones, along with *Pinus* spp., have emerged as amongst the most widely planted exotic forestry trees in South Africa (Denison & Kietzka 1993b). This is due to their rapid growth, adaptability and valuable end products. It is thus imperative to ensure an adequate supply of raw material, not influenced by factors such as diseases that could affect both quality and quantity of the end products. To support these clonal plantations, it is necessary to raise and maintain healthy *Eucalyptus* plants in nurseries.

Fungi cause most of the nursery diseases known on *Eucalyptus* species and include foliar, stem and root diseases (Viljoen *et al.* 1992, Brown & Ferreira

2000). Among the most important diseases are damping – off, root rot, shoot or web blight, stem cankers and leaf spots (Sharma *et al.* 1985, Viljoen *et al.* 1992, Brown & Ferreira 2000). Adequate control of these fungi relies on sound management strategies in nurseries that require knowledge of their development and spread.

Development of diseases in cuttings can be attributed to several factors during production. The optimal environmental requirements for rooting of *Eucalyptus* cuttings are similar to those required for fungal growth and infection (Alfenas *et al.* 1997). Close spacing of plants combined with regular watering and heavy use of fertilizers provides a microclimate between the plants that is ideal for the growth and infection of plants by pathogens. Wounding caused during cutting production or handling also facilitates infection.

Several reviews and surveys have been published on diseases of *Eucalyptus* spp. in nurseries (Sharma *et al.* 1985, Crous *et al.* 1991, Viljoen *et al.* 1992, Alfenas *et al.* 1997, Brown & Ferreira 2000). These have included diseases in both seedling and clonal cutting systems. There has, however, been no overall evaluation of diseases in clonal cutting nurseries in South Africa. The aim of this study was, therefore, to conduct a survey of *Eucalyptus* cutting nurseries and to determine which pathogens are most important in this environment.

Materials & Methods

Survey of Eucalyptus hybrid clones

Hybrid clonal *Eucalyptus* cuttings were collected from the four largest cutting nurseries in South Africa. All four forestry nurseries are located in KwaZulu Natal, South Africa. One nursery was situated in central KwaZulu Natal (Pietermaritzburg area) and the other three on the KwaZulu Natal north coast (Richards Bay area). At each nursery, five of the most commonly produced hybrid clones were selected for sampling. The hybrid clones included a pure *Eucalyptus grandis* Hill (TAG) clone, *E. grandis* x *camaldulensis* Dehnh. (GxC), *E. grandis* x *urophylla* Blake (GxU), *E. grandis* x *nitens* (Deane & Maiden) Maiden (GxN) and hybrid *E. nitens* x *grandis* (NxH). For each of the clones, 50 cuttings showing symptoms of disease were collected. Samples were placed in brown paper bags and transferred to the laboratory for further study. These collections were made on three different occasions at three-month intervals. Thus, 250 samples were collected from each nursery on each of three occasions, resulting in a total sample of 3000 cuttings.

Isolations were made directly from the cuttings on 2% Malt Extract Agar (MEA; Biolab, Midrand, South Africa) from half of the cuttings sampled. The remaining 25 cuttings were placed in moist chambers and incubated at room temperature for 48h in the dark to promote sporulation of fungi. Conidia and fungal structures

produced on the cuttings were transferred to 2% MEA and incubated for seven days at 25°C under continuous near – ultraviolet light. Preliminary identifications were made with light microscopy and the most commonly isolated fungal pathogen was selected for further study. The cultures derived from this study have been maintained in the culture collection (CMW) of the Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, South Africa.

Morphological characteristics

A large number of isolates resembling *Cylindrocladium* spp. were obtained from the cuttings and these were the most common probable pathogens isolated (Table 1). These *Cylindrocladium* isolates were grouped into five groups based on colony morphology and growth rate (results not shown). For each group, ten isolates were randomly selected for more comprehensive identification based on morphology.

Single conidial isolates were transferred onto Carnation Leaf Agar (CLA) plates (Crous 2002) to induce production of both anamorph and teleomorph structures. The plates were incubated at 25°C under continuous near – ultraviolet light and examined after 14 days. Morphological characteristics described by Crous (2002) were used to identify cultures. Macroconidiophores were mounted in lactophenol and twenty measurements of vesicles, stipes and conidia were made for each of the selected isolates, using a light microscope. Measurements for these

structures are presented as (min)-(average – standard deviation) – (average + standard deviation)(-max).

DNA sequence comparisons

One isolate (Table 2) was randomly selected from each of the five morphological groups of *Cylindrocladium* isolates for DNA sequence comparisons. The single conidial isolates were grown on 2% MEA plates from which mycelium was collected and freeze dried. The freeze – dried mycelium was ground to a fine powder in liquid nitrogen with a mortar and pestle. DNA was extracted using the technique described by Möller *et al.* (1992).

A 495 base pairs fragment of the β - tubulin gene was amplified using primers T1 (O'Donnell & Cigelnik 1997) and Bt2b (Glass & Donaldson 1995). The PCR reaction of 25 μ l comprised of 2.5 units Taq enzyme (Roche Molecular Biochemicals, Alameda, California, USA), 10x buffer, 1mM MgCl₂ (as supplied by manufacturer), 0.25 mM deoxynucleoside triphosphate (dNTPs), 0.5 μ l primers and approximately 30 ng of fungal DNA as target. The amplified fragments were purified using a High Pure PCR Product Purification Kit (Roche Molecular Biochemicals, Alameda, California, USA).

Both DNA strands of the PCR products was sequenced using the primers used for the PCR amplification. Sequence reactions were done using an ABI

PRISM™3100 DNA Autosequencer (Applied BioSystems). Sequence data were processed using Sequence Navigator version 1.0.1 (Applied BioSystems). The nucleotide sequences were aligned manually by inserting gaps where necessary and phylogenetic relationships were determined using PAUP version 4.0b10 (Swofford 2002). Gaps were treated as missing data and confidence intervals were determined using 1000 bootstrap replications. To establish the phylogenetic relationships and identities of the *Cylindrocladium* isolates from the cuttings, 15 sequences of known *Cylindrocladium* species (Table 2) obtained by Schoch *et al.* (2001) and Crous (2002) were taken from GenBank and included in the alignment. *Cylindrocladiella infestans* Boesew. (AF320190) was used as the outgroup taxon in the analyses.

Evaluation of pathogenicity

Plants selected to assess pathogenicity of *Cylindrocladium* isolates represented four of the five hybrid clones included in the disease survey. These were thus TAG, GxC, GxU and GxN. The two-month-old plants were maintained in a greenhouse with a constant environment at 25°C prior to and during the trial. Thirty plants of each hybrid clone were used for the trial, which included ten controls for each clone.

One isolate of *C. pauciramosum* (CMW 9108), which was identified as the dominant potential pathogen from the surveys, was selected for inoculation of

Eucalyptus hybrid clones. This isolate was selected because of its rapid growth on MEA (results not shown) and its ability to produce abundant fruiting structures. The isolate was transferred to 100 2% MEA plates and incubated for seven days at 25°C under continuous near – ultraviolet light.

A spore suspension was prepared by adding 1 – 2 ml of sterile water to the plates and dislodging conidia with a sterile glass rod. The spore suspension was drained through a layer of cheesecloth and the concentration adjusted to 3.3×10^5 conidia/ml. Tween 80 (Merck – Schuchardt, Hohenbunn, Germany) was added to the suspension to ensure that conidia adhered to the plant tissue surface.

The spore suspension was sprayed onto the surface of the leaves to just before run – off. The plants were watered and covered with plastic bags for 48h to ensure sufficient humidity for infection. Controls were sprayed with sterile water and otherwise treated in the same manner as the treated plants. Lesions were scored after removing the plastic bags.

Scoring of symptoms was achieved using a scale of 0 – 4 with 0 representing 0% infection, 1 representing less than 25% infection, 2 representing 26 – 50% infection, 3 representing 51 – 75% infection and 4 representing above 76% infection. Statistical analyses of the infection data were carried out using SAS analytical programmes (1990). The analysis produced means for each scale with

a 95% confidence limit for each mean using the General Linier Method. Re – isolations were made from the lesions to confirm the presence of the test fungus.

Results

Survey of Eucalyptus hybrid clones

Several well-known *Eucalyptus* nursery pathogens, represented by 2659 fungal isolates, were collected from the *Eucalyptus* cuttings at all four forestry nurseries surveyed (Table1). *Cylindrocladium* isolates were the most common potential pathogens at all four forestry nurseries and the next most common pathogen was *Botrytis cinerea* Pers. (Table 1). The pathogens were isolated from the roots, stems and leaves of the cuttings and all have been reported previously in South Africa.

Morphological characteristics of Cylindrocladium isolates

White macroconidiophores characteristic of those of *Cylindrocladium* spp. (Fig. 1A – B) were common on the surface of the *Eucalyptus* cuttings collected at the five forestry nurseries. Cultures on 2% MEA resulting from isolations from these structures differed in colony morphology. Thus, grouping according to colony morphology and growth rate was necessary.

All *Cylindrocladium* isolates were morphologically similar. The stipes and extensions to the stipes (Fig. 2 A – B) were septate, straight, hyaline, (85)103 – 130(147) μm in length and terminated in obpyriform to ellipsoidal vesicles, (11)14 – 19(23) \times (4)5 – 7 μm (Fig. 2 A – D). Each terminal branch of the fertile branches produced approximately 4 – 6 phialides (Fig. 2B). Phialides were doliiform to reniform, hyaline and aseptate. Conidia were cylindrical, straight, (40)46 – 53(59) \times 4 – 5(6) μm , 1 – septate and held in parallel cylindrical clusters by colourless slime (Fig. 2E). These characteristics are typical of *C. pauciramosum* C. L. Schoch & Crous as described by Schoch *et al.* 1999.

DNA sequence comparisons for Cylindrocladium isolates

A data set of 15 in-group taxa and one out-group taxon, *Cylindrocladiella infestans*, was analyzed. The alignment of the β -tubulin gene fragments gave rise to a data set of 495 characters of which 316 were constant, 62 were parsimony – uninformative and 17 parsimony – informative (Appendix II). One tree from two most parsimonious trees was chosen for presentation (Fig. 4). The trees had a consistency index (CI) = 0.817, retention index (RI) = 0.813, and rescaled consistency index (RC) = 0.678 while gaps were treated as missing data. The phylogenetic tree (Fig.4) clearly showed that all five randomly selected *Cylindrocladium* isolates grouped in the clade representing *C. pauciramosum* (74% bootstrap support).

Pathogenicity trial

First symptoms of infection were observed immediately after removal of the plastic bags at 48 hours (Fig. 3A – D). Initial symptoms on the leaves appeared as water – soaked lesions that caused a pink to purple discolouration of the leaf laminas. Lesions later became light brown and surrounded by a dark red to purple halo and varied in shape from round to irregular depending on the hybrid clone inoculated (Fig. 3A – D). Regression coefficients over the 22 days of evaluation were then determined for the mean of each scale (Table 3). All four clones were susceptible to *C. pauciramosum* infection (Fig. 3A – D) with the sub – tropical hybrid clone GxC (Fig. 3A) being the most susceptible. No symptoms were observed on the control plants.

Discussion

This study represents the first survey of pathogens in *Eucalyptus* clonal production nurseries in South Africa. A relatively large number of well – known pathogens were found and all have previously been recorded in the country (Lundquist & Baxter 1985, Crous *et al.* 1991, Viljoen *et al.* 1992). An interesting result was that a single species of *Cylindrocladium* was present and this was also the dominant pathogen in all nurseries surveyed.

Cylindrocladium spp. are well-known and important pathogens of *Eucalyptus* in nurseries around the world. In South Africa only *C. scoparium* Morgan, *C. candelabrum* Viégas, *C. clavatum* Hodges & May and *C. colhounii* Peerally have been reported from *Eucalyptus* spp. in forestry nurseries and plantations (Crous *et al.* 1991, Crous *et al.* 1993). This study, however, showed that only *C. pauciramosum* occurs in cutting production nurseries at the present time. It is possible that other *Cylindrocladium* spp. could be important on *Eucalyptus* cuttings in nurseries in the future, but the widespread occurrence of a single species suggests that *C. pauciramosum* is the dominant pathogen.

In this study, we used β – tubulin sequence data to confirm the identification of the *Cylindrocladium* isolates. Schoch *et al.* (1999) found that sequencing the Inter Transcribed spacer (ITS) including the 5.8S ribosomal DNA (rDNA) of *Cylindrocladium* spp. yielded very few variable characters in the region, to such an extent that the mating populations in the *C. candelabrum* species complex including *C. pauciramosum*, could not be distinguished from each other. DNA sequences obtained from part of the β – tubulin gene yielded more variable characters and distinguished between these fungi (Schoch *et al.* 2001). This was confirmed in the present study and supported our identifications made based on morphology.

Pathogenicity tests conducted in this study represent the first that have been done with *C. pauciramosum* on *Eucalyptus* hybrid plants. These tests showed

that most of the commonly used hybrids in the forestry industry are susceptible to infection by this pathogen. The extent of leaf lesions varied between the hybrids as has been shown previously with other *Cylindrocladium* spp. (Brown & Ferreira 2000). This implies that it might be possible to select disease tolerant plants based on nursery screening, but many more clones have to be used in tests to verify this observation.

Several pathogens are responsible for diseases of *Eucalyptus* cuttings in forestry nurseries of South Africa. The most important pathogen of *Eucalyptus* cuttings in South Africa is clearly *C. pauciramosum*. In this study, *C. pauciramosum* was the only *Cylindrocladium* sp. present in the forestry nurseries where the surveys were conducted. This pathogen is capable of rapid infection and can thus cause severe epidemics in forestry nurseries, if it is not controlled. Future studies are thus planned to gain a greater understanding of the genetic structure of the pathogen in South Africa and to consider relative susceptibility of important planting stock.

References

- Aimers – Halliday, J., Dibley, M. J., Hong, S. O. & Menzies, M. I. (1999) Cutting propagation of hybrid *Eucalyptus grandis* x *E. nitens*. *New Zealand Journal of Forestry Science* **29**: 238 – 250.

- Alfenas, A. C., Silveira, S. F. da & Stowasser, E. A. S. V. (1997) Current status and control strategies of diseases associated to clonal propagation of *Eucalyptus* in Brazil. *IUFRO Conference on Silviculture and Improvement of Eucalyptus, Salvador, Brazil, 24 – 29 August.*
- Brown, B. N. & Ferreira, F. A. (2000) Disease during propagation of eucalypts. In *Diseases and pathogens of eucalypts*, Chapter 8: 119-151.(Ed. P. J. Keane, G. A. Kile, F. D. Podger, B. N. Brown). CSIRO Publishing, Collingwood, Australia.
- Crous, P. W. (2002) Taxonomy and pathology of *Cylindrocladium* (*Calonectria*) and allied genera. APS Press, St. Paul, Minnesota, USA.
- Crous, P.W., Phillips, A. L. J. & Wingfield, M. J. (1991) The genera *Cylindrocladium* and *Cylindrocladiella* in South Africa, with special reference to forest nurseries. *South African Journal of Forestry* **157**: 69 – 85.
- Crous, P.W., Phillips, A. L. J. & Wingfield, M. J. (1993) New records of *Cylindrocladium* and *Cylindrocladiella* spp. in South Africa. *Plant Pathology* **42**: 302 – 305.

- Denison, N. P. & Kietzka, J. E. (1993a) The use and importance of hybrid intensive forestry in South Africa. *South African Forestry Journal* **162**: 55 – 60.
- Denison, N. P. & Kietzka, J. E. (1993b) The development and utilization of vegetative propagation in Mondi for commercial afforestation programmes. *South African Forestry Journal* **162**: 47 – 54.
- Glass, N. L. & Donaldson, G. (1995) Development of primer sets designed for use with PCR to amplify conserved genes from filamentous ascomycetes. *Applied and Environmental Microbiology* **61**: 1323 – 1330.
- Kulkarni, H. D. & Lal, P. (1995) Performance of *Eucalyptus* clones at ITC Bhadrachlm, India. CRC for Temperate Hardwood Forestry, IUFRO, Hobart.
- Lundquist, J. E. & Baxter, A. P. (1985) Fungi associated with *Eucalyptus* in South Africa. *South African Forestry Journal* **138**: 9 – 19.
- Möller, E. M., Bahnweg, G., Sandermann, H. & Geiger, H. H. (1992) A simple and efficient protocol for isolation of high molecular weight DNA from filamentous fungi, fruit bodies, and infected plant tissues. *Nucleic Acid Research* **20**: 6115 – 6116.

O'Donnell, K. & Cigelnik, E. (1997) Two divergent intragenomic rDNA ITS2 types within a monophyletic lineage of the fungus *Fusarium* are nonorthologous. *Molecular Phylogenetics Evolution* **7**: 103 – 116.

Schoch, C. L., Crous, P. W. Wingfield, B. D. & Wingfield, M. J. (1999) The *Cylindrocladium candelabrum* species complex includes four distinct mating populations. *Mycologia* **91**: 286 – 298.

Schoch, C. L., Crous, P. W. Wingfield, B. D. & Wingfield, M. J. (2001) Phylogeny of *Calonectria* based on comparisons of β - tubulin DNA sequences. *Mycological Research* **105**: 1045 – 1052.

Sharma, J. K. Mohanan, C. Florence, E. J. M. (1985) Disease survey in nurseries and plantations of forest tree species grown in Kerala. *Research report 36*. (Kerala Forest Research Institute: Peechi, Kerala).

Swofford, D. L. (2002) PAUP*. Phylogenetic analysis using parsimony (* and other methods), 4.0b10. Computer programme. Sunderland, Massachusetts, U.S.A.: Sinauer Associates.

- Viljoen, A. Wingfield, M. J. & Crous, P. W. (1992) Fungal pathogens in *Pinus* and *Eucalyptus* seedling nurseries in South Africa: A review. *South African Forestry Journal* **161**: 45-51.
- White, T. L. (1995) Prototype breeding strategies for clonal forestry: A case study of *Eucalyptus grandis*. CRC for Temperate Hardwood Forestry, IUFRO, Hobart.
- Wilson, P. J. (1998) Developing clones from *Eucalyptus globulus* and hybrid seedlings by stem cuttings propagation. *New Zealand Journal of Forestry Science* **28**: 293 – 303.

Table 1: List of 2659 fungal isolates obtained during the survey of *Eucalyptus* hybrid clone cuttings from all four forestry nurseries expressed as percentages.

FUNGUS	Percentage of isolates
<i>Alternaria</i> sp.	3
<i>Botryosphaeria</i> sp.	0.6
<i>Botrytis cinerea</i> Pers.	15
<i>Cladosporium</i> sp.	0.1
<i>Colletotrichum gloeosporioides</i>	8
<i>Cylindrocladiella</i> sp1.	2
<i>Cylindrocladiella</i> sp2.	1
<i>Cylindrocladium</i> sp.	21
<i>Epicoccum</i> sp.	5
<i>Fusarium</i> sp1.	5
<i>Fusarium</i> sp2.	2
<i>Fusarium</i> sp3.	2
<i>Fusarium</i> sp4.	4
<i>Gliocladium</i> sp.	5
<i>Hainesia lythri</i>	4
<i>Mycosphaerella</i> spp.	a
<i>Pestalotiopsis</i> sp.	7
<i>Phomopsis</i> sp.	3
<i>Pseudocercospora</i> sp.	a
<i>Rhizoctonia</i> sp.	5
<i>Trichoderma</i> sp.	7
<i>Verticillium</i> sp.	0.3

(a) Observed but not isolated from *Eucalyptus* hybrid clone cuttings.

Table 2: List of *Cylindrocladium* spp. used in DNA sequence comparisons.

Species	Isolate number	GenBank Assencion number
<i>Cylindrocladium candelabrum</i>	STE-U 1677	AF210858
	STE-U 1674	AF210857
<i>C. clavatum</i>	ATCC 46300	AF232850
<i>C. colhounii</i>	STE-U 307	AF232855
	STE-U 705	AF232854
<i>C. gracile</i>	STE-U 623	AF333405
	IMI 167580	AF333404
<i>C. ilicola</i>	STE-U 723	AF333413
	CBS 190.50	AF333412
<i>C. insulare</i>	STE-U 3211	AF44951
	STE-U 3219	AF44950
<i>C. pauciramosum</i>	^a CMW 8741	
	^a CMW 9108	
	^a CMW 9109	
	^a CMW 9189	
	^a CMW 9190	
	STE-U 3207	AF449448
	CSL 2021133	AY162320
<i>C. scoparium</i>	STE-U 1722	AF210875
	STE-U 1720	AF210874

^a *Cylindrocladium* isolates from *Eucalyptus* cuttings that were sequenced.

Table 3: Regression coefficients for the means of each scale to indicate rate of infection (leaves/day) for each *Eucalyptus* hybrid clone used in the pathogenicity test with *C. pauciramosum*.

Hybrid clone	0*	1*	2*	3*	4*
G x C	1.90 ^b	-0.41 ^a	0.37 ^b	0.73 ^a	1.20 ^a
G x N	1.49 ^a	-0.69 ^b	-0.56 ^a	0.52 ^b	0.69 ^a
G x U	1.80 ^a	-0.15 ^b	-0.21 ^b	0.39 ^b	0.68 ^b
TAG	1.16 ^a	-0.43 ^a	0.68 ^a	0.72 ^b	-5.42 ^b

^a Infection rate significant ($P < 0.05$).

^b Infection rate not significant ($P > 0.05$).

* Scales used to score disease symptoms.

Figure 1: Macroconidiophores of *Cylindrocladium pauciramosum*. A. Macroconidiophores on the stem of a *Eucalyptus* cutting collected during the survey. B. Macroconidiophores on a *Eucalyptus* leaf.





Figure 2: Macroconidiophores, vesicles and conidia of *Cylindrocladium pauciramosum*. A – B. Macroconidiophores. C – D. Vesicles. E. Conidia (Scale bars = 10 µm).







Figure 3: Leaf lesions on *Eucalyptus* hybrid clones used in pathogenicity trial.

A. *Eucalyptus grandis* x *camaldulensis*; B. *E. grandis* x *nitens*; C. *E. grandis* x *urophylla*; D. *E. grandis*






Figure 4: The most parsimonious tree obtained from a subset of *Cylindrocladium* isolates (495 steps, CI = 0.817, RC = 0.678, RI = 0.813) generated with a heuristic search in PAUP version 4.0b1 from aligned sequences of the 5' end of the β - tubulin gene. Gaps were treated as missing. Clade stability was assessed with 1000 bootstrap replications and values above 50% are shown above branches and decay indices below. A *Cylindrocladiella infestans* (GenBank accession number AF320190) was used as outgroup.





Chapter 4

Cylindrocladium pauciramosum
infecting *Acacia mearnsii*
seedlings in South Africa



Abstract

The versatility of *Acacia mearnsii* makes this a popular tree to cultivate in commercial plantations in South Africa. The timber has a wide variety of uses and the bark contains vegetable tannins that are important in the leather and adhesives industry. The tree is native to Australia and in South Africa has been affected by a number of serious diseases in plantations and nurseries. This study arose from an outbreak of *Cylindrocladium* blight in an *A. mearnsii* nursery. The aim was thus to identify the causal agent of the disease. *Cylindrocladium* isolates were collected from diseased seedlings and identified as *C. pauciramosum*, based on morphological characteristics as well as DNA sequence comparisons. A pathogenicity test was conducted, where *A. mearnsii* seedlings were inoculated using three different isolates. Comparison of lesion lengths and re-isolation of the inoculated fungus from lesions showed that the fungus was pathogenic and there was little difference between isolates.

Introduction

Acacia mearnsii de Wild (black wattle) was first introduced into South Africa from Australia in 1864 (Sherry 1971) to provide firewood, shelterbelts and shade (Dunlop 2002). The first plantations of *A. mearnsii* were established in South Africa in 1888 after it was discovered that the bark was rich in vegetable tannins that could be used in the leather industry (Dunlop 2002). The demand for leather products during World War II led to increased wattle plantings in the mid 1900's and these now make up about 7% of the South African commercial forestry areas (Dunlop 2002, Anonymous 2003).

In South Africa, *A. mearnsii* has been damaged by several pathogens and pests (Wingfield & Roux 2000). Several reviews on this topic have been provided (Roux *et al.* 1995, Old *et al.* 2003, Roux 2002). There is, however, a general lack of knowledge regarding disease problems associated with *A. mearnsii* in forestry nurseries (Roux 2002).

Roux (2002) reported that the most common disease problems of *A. mearnsii* in nurseries are caused by a *Cylindrocladium* sp. Symptoms included root rot (Crous *et al.* 1991), stem cankers and shoot blight (Roux 2002). These symptoms were initially believed to be caused by *C. scoparium* Morgan (Crous *et al.* 1991, Roux & Wingfield 1997) but recent studies indicate that this pathogen is restricted to the Northern hemisphere (Overmeyer *et al.* 1996, Crous 2002). Other *Cylindrocladium* spp. that have been reported on *A. mearnsii* and other *Acacia* spp. elsewhere in the world include *C. crotolariae* (Loos) Bell & Sobers, *C. candelabrum* Viegas, *C. illicicola* (Hawley) Boedjin &

Reitsma, *C. quinqueseptatum* Boedjin & Reitsma, *C. reteaudii* (Bugn.) Boesew. and *C. theae* (Petch) Subramanian (Old *et al.* 2003).

In South Africa, it has been suggested that the *Cylindrocladium* sp. responsible for diseases of *A. mearnsii* in nurseries resides in the *C. candulabrum* species complex (Schoch *et al.* 1999) and could be *C. pauciramosum* Schoch & Crous (Roux 2002). The aim of this study was to identify the *Cylindrocladium* sp. responsible for losses of *A. mearnsii* in nurseries in South Africa.

Material & Methods

Plant material and isolations

Acacia mearnsii seedlings were collected at a forestry nursery in KwaZulu Natal, South Africa. The 25 plants selected for study were girdled and had stem cankers. Isolations were made from symptomatic tissue directly on 2% Malt Extract Agar (MEA; Biolab, Midrand, South Africa) and incubated for seven days at 25°C under continuous near – ultraviolet light. All *Cylindrocladium* isolates obtained are maintained in the culture collection (CMW) of the Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, South Africa.

Morphological characteristics

Single conidial isolates were transferred onto Carnation Leaf Agar (CLA) plates (Crous 2002) to induce production of both anamorph and teleomorph structures. The plates were incubated at 25°C under continuous near – ultraviolet light and examined after 14 days. Morphological characteristics described by Crous (2002) were used to identify cultures. Macroconidiophores were mounted in lactophenol and 20 measurements of vesicles, stipes and conidia were made for each isolate using a light microscope. Measurements are presented as (min-)(average – standard deviation) – (average + standard deviation)(-max).

DNA sequence comparisons

Five isolates (Table 1) were randomly selected and utilized in the DNA sequence comparisons. Single conidial isolates were grown on 2% MEA plates from which mycelium was collected and freeze dried. The freeze – dried mycelium was ground to a fine powder in liquid nitrogen using a mortar and pestle with a mortar. DNA was extracted using the technique described by Möller *et al.* (1992).

A 507 bp. fragment of the β - tubulin gene was amplified using primers T1 (O'Donnell & Cigelnik 1997) and Bt2b (Glass & Donaldson 1995). The PCR reaction of 25 μ l comprised of 2.5 units Taq enzyme (Roche Molecular Biochemicals, Alameda, California, USA), 10x buffer, 1mM MgCl₂ (as supplied

by manufacturer), 0.25 mM deoxynucleoside triphosphate, 0.5 μ m primers and approximately 30 ng of fungal DNA as target. The amplified fragments were purified using a High Pure PCR Product Purification Kit (Taq (Roche Molecular Biochemicals, Alameda, California, USA).

The PCR product was sequenced in both directions using the same primers used for the PCR amplification. Sequence reactions were done using the ABI PRISM™3100 DNA Autosequencer (Applied BioSystems). Sequence data were processed using Sequence Navigator version 1.0.1 (Applied BioSystems, Foster City, California, USA). The nucleotide sequences were aligned manually by inserting gaps where necessary and phylogenetic relationships were determined using PAUP version 4.0b10 (Swofford 2002). Gaps were treated as missing data and confidence intervals were determined using 1000 bootstrap replications. To establish the phylogenetic relationships and identities of the *Cylindrocladium* isolates from the *A. mearnsii* seedlings, 12 sequences of known *Cylindrocladium* spp. (Table 1) were obtained from GenBank and included in the alignment. *Cylindrocladiella infestans* Boesew. (AF320190) was used as the out – group taxon in the analysis.

Evaluation of Pathogenicity

Cylindrocladium isolates CMW 9156, CMW 9164 and CMW 9171 were randomly selected for use in glasshouse inoculations on *A. mearnsii*. Isolates were grown on 2% MEA for seven days prior to inoculation. Glasshouse

conditions were set with day/night lighting and with an average temperature of approximately 25°C.

Twenty *A. mearnsii* seedlings were inoculated for each isolate and an equal number were used for controls. A 5 mm diameter wound was made on the stem of each seedling by removing the bark with a cork borer and exposing the cambium. Plugs (5mm diam) were cut from the surface of Petri dishes, overgrown with the test isolates and placed in the wounds with the mycelial surface facing the cambium. All inoculations were covered with parafilm to prevent desiccation of the inoculum and wounds. For control inoculations, sterile 2% MEA plugs were used. Lesion lengths were assessed after six weeks and statistical differences in lesion length for each isolate were determined using SAS analytical programmes (2002). Re – isolations were made from the lesions to confirm the presence of the test fungus.

Results

Morphological characteristics

All *Cylindrocladium* isolates were morphologically similar. The stipes and extensions to the stipes (Fig. 1A) were septate, straight, hyaline, (84)102 – 140(179) μm in length and terminated in obpyriform to ellipsoidal vesicles, (10)14 – 22(32) \times (3)5 – 7 μm (Fig. 2D - F). Each terminal branch of the conidiophores produced approximately 4 – 6 phialides (Fig. 1A). Phialides were doliiform to reniform, hyaline and aseptate. Conidia were cylindrical,

straight, $(37)46 - 55(64) \times (3)4 - 5(7) \mu\text{m}$, 1 – septate and held in parallel cylindrical clusters by colourless slime (Fig. 1B). These characteristics are typical of *C. pauciramosum* C. L. Schoch & Crous as described by Schoch *et al.* (1999).

DNA sequence comparisons

A data set of 17 in-group taxa and one out-group taxon, *Cylindrocladiella infestans*, was analyzed. The alignment of the β -tubulin gene fragments (Fig. 4) gave rise to a data set of 507 characters of which 344 were constant, 70 were parsimony – uninformative and 93 parsimony – informative (Fig.2). One tree from 4 most parsimonious trees was chosen for presentation (Fig.2). The trees had a consistency index (CI) = 0.917, retention index (RI) = 0.929, and rescaled consistency index (RC) = 0.852 while gaps were treated as missing data. The phylogenetic tree (Fig.2) clearly showed that all five randomly selected *Cylindrocladium* isolates from *A. mearnsii* grouped in the clade representing *C. pauciramosum* (94% bootstrap support).

Pathogenicity tests

After six weeks, inoculated seedlings displayed symptoms of die-back and wilting. Exposure of the cambium showed that the vascular tissue had a streaked appearance (Fig. 3). Isolations from the lesions on the inoculated plants consistently yielded isolates of *Cylindrocladium* and no pathogens were re-isolated from the control plants.

Cylindrocladium isolate CMW 9164 produced the longest lesions on *A. mearnsii* seedlings ($X = 105.7$ mm). Isolates CMW 9156 and CMW.9171 produced lesions with mean lengths of 80.4 mm and 102.7mm respectively. (Fig.3). However, there was no significant difference between the three isolates used in the trial (Table 2). All isolates produced lesions significantly ($P=0.0001$) larger than those of the controls that had an average lesion length of 25.5 mm (Fig. 3).

Discussion

In this study, we have shown that *Cylindrocladium pauciramosum* is the species responsible for disease of *A. mearnsii* in South African nurseries. Identification of the fungus was based on morphological characteristics and this was further confirmed using DNA sequence comparisons. Although *Cylindrocladium* has previously been recognized as a pathogen of *A. mearnsii* seedlings in nurseries, this is the first robust identification of the pathogen. This study also provides the first pathogenicity tests using *C. pauciramosum* on *A. mearnsii*.

DNA sequence comparisons were used in this study to confirm the identity of the *Cylindrocladium* sp. associated with *A. mearnsii* infections. This was necessary because species in the *Cylindrocladium candelabrum* complex including *C. pauciramosum* are morphologically similar and difficult to identify (Crous 2002). It is possible that other species in this group might appear on *A. mearnsii* in the future. However, recent surveys in *Eucalyptus* nurseries have

shown that *C. pauciramosum* is the dominant fungus in those situations (Chapter 3, this thesis).

To the best of our knowledge, pathogenicity tests have not previously been conducted on *A. meamsii*, using *C. pauciramosum*. Our tests clearly showed that this fungus is pathogenic to this tree species and that the three isolates chosen for the test have similar levels of pathogenicity. *Cylindrocladium pauciramosum* appears to be one of the most common species of *Cylindrocladium* causing plant diseases in South Africa (Crous 2002) and it is particularly important in forest nurseries.

Cylindrocladium spp. are soil-borne and produce microsclerotia that enable them to survive in soil and potting medium for long periods of time. To reduce the impact of *C. pauciramosum* in *A. meamsii* nurseries, it will be necessary to eliminate sources of inoculum. In this regard, it will be especially important to ensure that planting substrate is free of the pathogen and that inoculum does not build up in nurseries.

References

Anonymous. (2003) Abstract of South African Forestry Facts for the year 2001/2002. Forestry South Africa, Department Water Affairs & Forestry, South Africa.

- Crous, P. W. (2002) Taxonomy and pathology of *Cylindrocladium* (*Calonectria*) and allied genera. APS Press, St. Paul, Minnesota, USA.
- Crous, P.W., Phillips, A. L. J. & Wingfield, M. J. (1991) The genera *Cylindrocladium* and *Cylindrocladiella* in South Africa, with special reference to forest nurseries. *South African Journal of Forestry* **157**: 69 – 85.
- Dunlop, R. W. (2002) Introduction. In *Black Wattle: The South African research experience*, Chapter 1: 1 – 2 (Ed. R. W. Dunlop & L. A. MacLennan) Institute for Commercial Forestry Research, Pietermaritzburg, South Africa.
- Glass, N. L. & Donaldson, G. (1995) Development of primer sets designed for use with PCR to amplify conserved genes from filamentous ascomycetes. *Applied and Environmental Microbiology* **61**: 1323 – 1330.
- Möller, E. M., Bahnweg, G., Sandermann, H. & Geiger, H. H. (1992) A simple and efficient protocol for isolation of high molecular weight DNA from filamentous fungi, fruit bodies, and infected plant tissues. *Nucleic Acid Research* **20**: 6115 – 6116.

- O'Donnell, K. & Cigelnik, E. (1997) Two divergent intragenomic rDNA ITS2 types within a monophyletic lineage of the fungus *Fusarium* are nonorthologous. *Molecular Phylogenetics Evolution* **7**: 103 – 116.
- Old, K. M., Vercoe, T. K., Floyd, R. B., Wingfield, M. J., Roux, J. & Naser, S. (2003) *Acacia* spp. *FAO/IPGRI Technical Guidelines for the Safe Movement of Germplasm* No. **20**.
- Overmeyer, C. Lünemann, S., Von Wallbrünn, C. & Meinhardt F. (1996) Genetic variability among isolates and sexual offspring of the plant pathogenic fungus *Calonectria morganii* on the basis of random amplification of polymorphic DNA (RAPD) and restriction fragment length polymorphisms (RFLP). *Current Microbiology* **33**: 249 – 255.
- Roux, J. (2002) Threats to wattle plantations III: Diseases. In *Black Wattle: The South African research experience*, Chapter 12: 121 – 126. (Ed. R. W. Dunlop & L. A. MacLennan) Institute for Commercial Forestry Research, Pietermaritzburg, South Africa.
- Roux, J., Kemp, G. H. J. & Wingfield, M. J. (1995) Diseases of black wattle in South Africa – A review. *South African Journal of Forestry* **174**: 35 – 40.

- Roux, J. & Wingfield, M. J. (1997) Survey and virulence of fungi occurring on diseased *Acacia mearnsii* in South Africa. *Forest Ecology and Management* **99**: 327 – 336.
- Sherry, S. P. (1971) *The black wattle (Acacia mearnsii de Wild)*. University of Natal Press, Pietermaritzburg, South Africa. 402p.
- Schoch, C. L., Crous, P. W. Wingfield, B. D. & Wingfield, M. J. (1999) The *Cylindrocladium candelabrum* species complex includes four distinct mating populations. *Mycologia* **91**: 286 – 298.
- Swofford, D. L. (2002) PAUP*. Phylogenetic analysis using parsimony(* and other methods), 4.0b10. Computer programme. Sunderland, Massachusetts, U.S.A.: Sinauer Associates.
- Wingfield, M. J. & Roux, J. (2000) Diseases of Forest Plantations. In *South African Forestry Handbook 2000*. South African Institute of Forestry, Pretoria, South Africa, pp. 241 – 252.

Table 1. List of *Cylindrocladium* isolates used in DNA sequence comparisons

Species	Isolation number	GenBank Assencion number
<i>Cylindrocladium candelabrum</i>	STE-U 1677	AF210858
<i>C. pauciramosum</i>	STE-U 1674	AF210857
	STE-U 3207	AF449448
	CSL 2021133	AY162320
	^{a,b} CMW9156	
	^a CMW9159	
	^{a,b} CMW9164	
<i>C. quinqueseptatum</i>	^a CMW9169	
	^{a,b} CMW9171	
	STE-U 759	AF232869
<i>C. reteaudii</i>	STE-U 516	AF232870
	STE-U 758	AF389847
<i>C. scoparium</i>	STE-U 602	AF389846
	STE-U 1722	AF210875
<i>C. theae</i>	STE-U 1720	AF210874
	UFV 16	AF232862
	ATCC 48895	AF232861

^a Isolates sequenced in this study. All other sequences are those from GenBank.

^b Isolates used in the pathogenicity test.

Table 2. Mean lesion lengths on *A. mearnsii* seedlings 6 weeks after inoculation with three isolates of *C. pauciramosum* and a control in the greenhouse.

Isolates	^aMean lesion length
CMW 9156	80.4
CMW 9164	105.7
CMW 9171	102.7
Control	25.5

^aEach value is the average of 20 measurements for each isolate.

P < 0.0001

CV = 35.94

R – square = 0.517

F = 22.48

Figure 1. Macroconidiophores, vesicles and conidia of *Cylindrocladium pauciramosum* isolated from *Acacia mearnsii* seedlings. A. Macroconidiophore with fertile branches (phialides); B. Conidia; C. Chlamydospores; D – F. Vesicles (Scale bars = 10 μm).






Figure 2. Most parsimonious tree obtained from a subset of *Cylindrocladium* isolates. The most parsimonious tree (507 steps, CI = 0.917, RC = 0.852, RI = 0.929) generated with a heuristic search in PAUP* version 4.0b1 from aligned sequences of the 5' end of the β - tubulin gene. Gaps were treated as missing. Clade stability was assessed with 1000 bootstrap replications and values above 50% are shown above branches and decay indices below. A *Cylindrocladiella infestans* (GenBank accession number AF320190) was used as outgroup.




Figure 3. Streaked appearance of the vascular tissue of *A. mearnsii* stems after inoculation with *C. pauciramosum* including control. A. Control; B. CMW 9164; C. CMW 9171; D. CMW 9156.



Summary

Studies presented in this dissertation highlight the importance of fungal pathogens in forestry nurseries in South Africa. Both *Acacia mearnsii* seedlings and *Eucalyptus* hybrid cuttings are shown to be affected by important nursery pathogens.

Chapter one presents an evaluation of the potential importance of pathogens to *Eucalyptus* hedge plants maintained in hydroponics. Hydroponics is a new technology being used in South African forest nurseries, which allows for the rapid establishment of *Eucalyptus* hedge plants. However, no information is available on pathogens affecting *Eucalyptus* in hydroponics. By applying information on pathogens of other hydroponic crops, several potentially important pathogens were identified and these reside in the genera *Phytophthora*, *Pythium* and *Fusarium*. Possible disease symptoms in *Eucalyptus* caused by these pathogens include wilting, stem cankers and root rots. Implementation of appropriate control measures that include cultural, biological and chemical practices could prevent and/or reduce disease impact in hydroponics.

Chapter two presents the results of a survey of the roots of *Eucalyptus* hedge plants grown in an ebb and flow hydroponic system. An interesting result of the survey was the discovery of *Cylindrocladium pauciramosum* in the hydroponic system. This is the first report of the pathogen in a hydroponic system. Other important pathogens in the genera *Phytophthora* and *Pythium*

were also isolated. Two *Pythium* species, namely *P. dissotocum* and *P. helicoids*, found in the roots and nutrient solution are new to *Eucalyptus*. Several *Fusarium* species were also isolated of which two, namely *F. nygamai* and *F. lateritium*, are also new to *Eucalyptus*.

Chapter three of this dissertation presents the results of a survey of *Eucalyptus* cuttings conducted at four forestry nurseries in KwaZulu – Natal, South Africa. Several well – known *Eucalyptus* nursery pathogens were isolated. *Cylindrocladium pauciramosum* was identified as the dominant pathogen on *Eucalyptus* cuttings. This was confirmed based on morphological characteristics and DNA sequence comparisons. Pathogenicity tests conducted using a spore suspension of *C. pauciramosum* indicated that this pathogen is capable of infecting most commercial *Eucalyptus* clones used in South Africa.


Chapter four considers a serious disease of *Acacia mearnsii* seedlings caused by an unidentified species of *Cylindrocladium*. *Cylindrocladium pauciramosum* was isolated from *A. mearnsii* seedlings showing girdling and stem canker symptoms. The pathogen was identified based on morphological characteristics and DNA sequence comparisons. Pathogenicity tests with *Acacia* seedlings confirmed the susceptibility of this tree to *C. pauciramosum* infection.

This dissertation clearly indicates that *Cylindrocladium pauciramosum* is an important nursery pathogen in South African forestry nurseries. This pathogen

has already been shown to be limiting during production of planting stock. I hope to have highlighted the importance of *C. pauciramosum* and other nursery pathogens in forestry nurseries in South Africa. This study will also hopefully provide information to forestry nursery managers and help them improve production.

**Appendix I – A combined phylogenetic tree of *Cylindrocladium*
pauciramosum isolates used.**

The most parsimonious tree obtained from a subset of *Cylindrocladium* isolates (560 steps, CI = 0.774, RC = 0.602, RI = 0.777) generated with a heuristic search in PAUP version 4.0b1 from aligned sequences of the 5' end of the β - tubulin gene. Gaps were treated as missing. Clade stability was assessed with 1000 bootstrap replications and values above 50% are shown below branches and decay indices above. A *Cylindrocladiella infestans* (GenBank accession number AF320190) was used as out – group.



The most parsimonious tree obtained from the 5' end of the β – tubulin gene DNA sequence of the *C. pauciramosum* isolates used in this dissertation indicates that isolates are similar irrespective of host or environment. A slight difference is evident between the *Eucalyptus* and the *Acacia mearnsii* isolates but this cannot be supported by these DNA sequence comparisons (bootstrap < 50%). The application of further molecular population studies and multigene phylogeny could support and differentiate between *C. pauciramosum* strains.



Appendix II – DNA alignments used.

Chapter 2

Alignment 1. 5' end of the β - tubulin gene of the *Cylindrocladium pauciramosum* isolates obtained from the *Eucalyptus* hedge plants.

CMW 12711		GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	30
CMW 12709		GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
CMW 12697		GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
CMW 12689		GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
CMW 13039		GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
CMW 12702		GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
CMW 12707		GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>C. pauciramosum</i>	AY162320	GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>C. pauciramosum</i>	AF449448	GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>C. candelabrum</i>	AF210857	GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>C. candelabrum</i>	AF210858	GCTT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>C. scoparium</i>	AF210875	GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>C. scoparium</i>	AF210874	GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>C. colhounii</i>	AF232855	GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>C. colhounii</i>	AF232854	GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>C. insulare</i>	AF44951	GCTT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>C. insulare</i>	AF44950	GCTT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>C. gracile</i>	AF333405	GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>C. gracile</i>	AF333404	GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>Cylindrocladiella infestans</i>	AF320190	GCT-GCCCCT	- ATTCTATCC	CGCCGCCCCG	
CMW 12711		GTTTCCACC-	---GCTTCGA	CGACAACAAA	60
CMW 12709		GTTTCCACC-	---GCTTCGA	CGACAACAAA	
CMW 12697		GTTTCCACC-	---GCTTCGA	CGACAACAAA	
CMW 12689		GTTTCCACC-	---GCTTCGA	CGACAACAAA	
CMW 13039		GTTTCCACC-	---GCTTCGA	CGACAACAAA	
CMW 12702		GTTTCCACC-	---GCTTCGA	CGACAACAAA	
CMW 12707		GTTTCCACC-	---GCTTCGA	CGACAACAAA	
<i>C. pauciramosum</i>	AY162320	GTTTCCACC-	---GCTTCGA	CGACAACAAA	
<i>C. pauciramosum</i>	AF449448	GTTTCCACC-	---GCTTCGA	CGACAACAAA	
<i>C. candelabrum</i>	AF210857	GTTTCCACC-	---GCTTCGA	CGACAACAAA	
<i>C. candelabrum</i>	AF210858	GTTTCCACC-	---GCTTCGA	CGACAACAAA	
<i>C. scoparium</i>	AF210875	GTTTCCACC-	---ACATCGA	CGAAAAACAAA	
<i>C. scoparium</i>	AF210874	GTTTCCACC-	---ACATCGA	CGAAAAACAAA	
<i>C. colhounii</i>	AF232855	GTTTCCACC-	---GCTTCGA	CGACAACAAA	
<i>C. colhounii</i>	AF232854	GTTTCCACC-	---GCTTCGA	CGACAACAAA	
<i>C. insulare</i>	AF44951	GTTTCCACC-	---ACCTCGA	CGACAACAAA	
<i>C. insulare</i>	AF44950	GTTTCCACC-	---ACCTCGA	CGACAACAAA	
<i>C. gracile</i>	AF333405	GTTTCCACC-	---GCTCCGA	CGACAACAAA	
<i>C. gracile</i>	AF333404	GTTTCCACC-	---GCTCCGA	CGACAACAAA	
<i>Cylindrocladiella infestans</i>	AF320190	TTTCCACC	ACCGCCTCGA	CGACAACAAA	90
CMW 12711		GCCGCAGCCT	CACGATCATA	ACGAGATATC	
CMW 12709		GCCGCAGCCT	CACGATCATA	ACGAGATATC	
CMW 12697		GCCGCAGCCT	CACGATCATA	ACGAGATATC	
CMW 12689		GCCGCAGCCT	CACGATCATA	ACGAGATATC	
CMW 13039		GCCGCAGCCT	CACGATCATA	ACGAGATATC	
CMW 12702		GCCGCAGCCT	CACGATCATA	ACGAGATATC	
CMW 12707		GCCGCAGCCT	CACGATCATA	ACGAGATATC	
<i>C. pauciramosum</i>	AY162320	GCCGCAGCCT	CACGATCATA	ACGAGATATC	
<i>C. pauciramosum</i>	AF449448	GCCGCAGCCT	CACGATCATA	ACGAGATATC	
<i>C. candelabrum</i>	AF210857	GCCGCAGCCT	CACGATCATG	ACGAGATATC	
<i>C. candelabrum</i>	AF210858	GCCGCAGCCT	CACGATCATG	ACGAGATATC	
<i>C. scoparium</i>	AF210875	GCCGCAGCCT	CACGAACATG	ATGTGATATC	
<i>C. scoparium</i>	AF210874	GCCGCAGCCT	CACGAACATG	ATGTGATATC	
<i>C. colhounii</i>	AF232855	GCCGCAGCCT	CACGAGCATG	ACGAGATATC	
<i>C. colhounii</i>	AF232854	GCCGCAGCCT	CACGAGCATG	ACGAGATATC	
<i>C. insulare</i>	AF44951	GCCGCAGCCT	CACGAACATG	ATGTGATATC	
<i>C. insulare</i>	AF44950	GCCGCAGCCT	CAACAACAAA	ATGTGATATC	
<i>C. gracile</i>	AF333405	GCCGCAGCCT	CACGAGCATG	GCGAGATATC	
<i>C. gracile</i>	AF333404	GCCGCAGCCT	CACGAGCATG	GCGAGATATC	
<i>Cylindrocladiella infestans</i>	AF320190	GCTCGCGATG	CCCACCCACA	TCGTGATATC	



CMW 12711		AGAACAAGAT	T- GCTAACCG	TGTGCTTCTT	120
CMW 12709		AGAACAAGAT	T- GCTAACCG	TGTGCTTCTT	
CMW 12697		AGAACAAGAT	T- GCTAACCG	TGTGCTTCTT	
CMW 12689		AGAACAAGAT	T- GCTAACCG	TGTGCTTCTT	
CMW 13039		AGAACAAGAT	T- GCTAACCG	TGTGCTTCTT	
CMW 12702		AGAACAAGAT	T- GCTAACCG	TGTGCTTCTT	
CMW 12707		AGAACAAGAT	T- GCTAACCG	TGTGCTTCTT	
<i>C. pauciramosum</i>	AY162320	AGAACAAGAT	T- GCTAACCG	TGTGCTTCTT	
<i>C. pauciramosum</i>	AF449448	AGAACAAGAT	T- GCTAACCG	TGTGCTTCTT	
<i>C. candelabrum</i>	AF210857	AGAACAAGAT	T- GCTAACCG	TGTGCTTCTT	
<i>C. candelabrum</i>	AF210858	AGAACAAGAT	T- GCTAACCG	TGTGCTTCTT	
<i>C. scoparium</i>	AF210875	AGAACAAGAT	T- GCTAACCG	TGTGCTTCTT	
<i>C. scoparium</i>	AF210874	AGAACAAGAT	T- GCTAACCG	TGTGCTTCTT	
<i>C. colhounii</i>	AF232855	GAAACAAGAT	TTGCTGACCA	TGTGCTTCTT	
<i>C. colhounii</i>	AF232854	GAAACAAGAT	TTGCTGACCA	TGTGCTTCTT	
<i>C. insulare</i>	AF44951	AGAACAAGAT	T- GCTAACCG	TGTGCTTCTT	
<i>C. insulare</i>	AF44950	AGAACAAGAT	T- GCTAACCG	TGTGCTTCTT	
<i>C. gracile</i>	AF333405	AGAACAAGAT	T- GCTAACCG	TGTGCTTCTT	
<i>C. gracile</i>	AF333404	AGAACAAGAT	T- GCTAACCG	TGTGCTTCTT	
<i>Cylindrocladiella infestans</i>	AF320190	TGAAGACAAT	G- GCTAATTT	TGTG-TGTTT	
CMW 12711		TCTCGATTAT	AGGTCCACCT	CCAGACCGGT	150
CMW 12709		TCTCGATTAT	AGGTCCACCT	CCAGACCGGT	
CMW 12697		TCTCGATTAT	AGGTCCACCT	CCAGACCGGT	
CMW 12689		TCTCGATTAT	AGGTCCACCT	CCAGACCGGT	
CMW 13039		TCTCGATTAT	AGGTCCACCT	CCAGACCGGT	
CMW 12702		TCTCGATTAT	AGGTCCACCT	CCAGACCGGT	
CMW 12707		TCTCGATTAT	AGGTCCACCT	CCAGACCGGT	
<i>C. pauciramosum</i>	AY162320	TCTCGATTAT	AGGTCCACCT	CCAGACCGGT	
<i>C. pauciramosum</i>	AF449448	TCTCGATTAT	AGGTCCACCT	CCAGACCGGT	
<i>C. candelabrum</i>	AF210857	TTTTCGATTAT	AGGTCCACCT	CCAGACCGGT	
<i>C. candelabrum</i>	AF210858	TTTTCGATTAT	AGGTCCACCT	CCAGACCGGT	
<i>C. scoparium</i>	AF210875	TCTCGATTAT	AGGTCCACCT	CCAGACCGGT	
<i>C. scoparium</i>	AF210874	TCTCGATTAT	AGGTCCACCT	CCAGACCGGT	
<i>C. colhounii</i>	AF232855	TTTTCAATTAT	AGGTCCACCT	CCAGACCGGT	
<i>C. colhounii</i>	AF232854	TTTTCAATTAT	AGGTCCACCT	CCAGACCGGT	
<i>C. insulare</i>	AF44951	TCTCGATTAT	AGGTCCACCT	CCAGACCGGT	
<i>C. insulare</i>	AF44950	TCTCGATTAT	AGGTCCACCT	CCAGACCGGT	
<i>C. gracile</i>	AF333405	TCTCGATTAT	AGGTCCACCT	CCAGACCGGT	
<i>C. gracile</i>	AF333404	TCTCGATTAT	AGGTCCACCT	CCAGACCGGT	
<i>Cylindrocladiella infestans</i>	AF320190	CTGCGAATAT	AGGTCCACCT	CCAGACCGGT	
CMW 12711		AGGTCCACCT	CCAGACCGGT	CAGTGCGTAA	180
CMW 12709		AGGTCCACCT	CCAGACCGGT	CAGTGCGTAA	
CMW 12697		AGGTCCACCT	CCAGACCGGT	CAGTGCGTAA	
CMW 12689		AGGTCCACCT	CCAGACCGGT	CAGTGCGTAA	
CMW 13039		AGGTCCACCT	CCAGACCGGT	CAGTGCGTAA	
CMW 12702		AGGTCCACCT	CCAGACCGGT	CAGTGCGTAA	
CMW 12707		AGGTCCACCT	CCAGACCGGT	CAGTGCGTAA	
<i>C. pauciramosum</i>	AY162320	AGGTCCACCT	CCAGACCGGT	CAGTGCGTAA	
<i>C. pauciramosum</i>	AF449448	AGGTCCACCT	CCAGACCGGT	CAGTGCGTAA	
<i>C. candelabrum</i>	AF210857	AGGTCCACCT	CCAGACCGGT	CAGTGCGTAA	
<i>C. candelabrum</i>	AF210858	AGGTCCACCT	CCAGACCGGT	CAGTGCGTAA	
<i>C. scoparium</i>	AF210875	AGGTCCACCT	CCAGACCGGT	CAGTGCGTAA	
<i>C. scoparium</i>	AF210874	AGGTCCACCT	CCAGACCGGT	CAGTGCGTAA	
<i>C. colhounii</i>	AF232855	AGGTCCACCT	CCAGACCGGT	CAGTGCGTAA	
<i>C. colhounii</i>	AF232854	AGGTCCACCT	CCAGACCGGT	CAGTGCGTAA	
<i>C. insulare</i>	AF44951	AGGTCCACCT	CCAGACCGGT	CAGTGCGTAA	
<i>C. insulare</i>	AF44950	AGGTCCACCT	CCAGACCGGT	CAGTGCGTAA	
<i>C. gracile</i>	AF333405	AGGTCCACCT	CCAGACCGGT	CAGTGCGTAA	
<i>C. gracile</i>	AF333404	AGGTCCACCT	CCAGACCGGT	CAGTGCGTAA	
<i>Cylindrocladiella infestans</i>	AF320190	AGGTCCACCT	CCAGACCGGT	CAGTGCGTAA	



				300
CMW 12711		ACCATTTCTG	GCGAGCACGG	TCTCGACAGC
CMW 12709		ACCATTTCTG	GCGAGCACGG	TCTCGACAGC
CMW 12697		ACCATTTCTG	GCGAGCACGG	TCTCGACAGC
CMW 12689		ACCATTTCTG	GCGAGCACGG	TCTCGACAGC
CMW 13039		ACCATTTCTG	GCGAGCACGG	TCTCGACAGC
CMW 12702		ACCATTTCTG	GCGAGCACGG	TCTCGACAGC
CMW 12707		ACCATTTCTG	GCGAGCACGG	TCTCGACAGC
<i>C. pauciramosum</i>	AY162320	ACCATTTCTG	GCGAGCACGG	TCTCGACAGC
<i>C. pauciramosum</i>	AF449448	ACCATTTCTG	GCGAGCACGG	TCTCGACAGC
<i>C. candelabrum</i>	AF210857	ACCATCTCTG	GCGAGCACGG	TCTCGACAGC
<i>C. candelabrum</i>	AF210858	ACCATCTCTG	GCGAGCACGG	TCTCGACAGC
<i>C. scoparium</i>	AF210875	ACCATTTCTG	GCGAGCACGG	TCTCGACAGC
<i>C. scoparium</i>	AF210874	ACCATTTCTG	GCGAGCACGG	TCTCGACAGC
<i>C. colhounii</i>	AF232855	ACCATCTCTG	GCGAGCACGG	TCTCGACAGC
<i>C. colhounii</i>	AF232854	ACCATCTCTG	GCGAGCACGG	TCTCGACAGC
<i>C. insulare</i>	AF44951	ACCATTTCTG	GCGAGCACGG	TCTCGACAGC
<i>C. insulare</i>	AF44950	ACCATTTCTG	GCGAGCACGG	TCTCGACAGC
<i>C. gracile</i>	AF333405	ACCATCTCTG	GCGAGCACGG	TCTCGACAGC
<i>C. gracile</i>	AF333404	ACCATCTCTG	GCGAGCACGG	TCTCGACAGC
<i>Cylindrocladiella infestans</i>	AF320190	ACCATCTCTG	GCGAGCACGG	TCTCGACAGC
				330
CMW 12711		AATGGTGTCT	ACGCCGGTAC	CTCCGAGCTC
CMW 12709		AATGGTGTCT	ACGCCGGTAC	CTCCGAGCTC
CMW 12697		AATGGTGTCT	ACGCCGGTAC	CTCCGAGCTC
CMW 12689		AATGGTGTCT	ACGCCGGTAC	CTCCGAGCTC
CMW 13039		AATGGTGTCT	ACGCCGGTAC	CTCCGAGCTC
CMW 12702		AATGGTGTCT	ACGCCGGTAC	CTCCGAGCTC
CMW 12707		AATGGTGTCT	ACGCCGGTAC	CTCCGAGCTC
<i>C. pauciramosum</i>	AY162320	AATGGTGTCT	ACGCCGGTAC	CTCCGAGCTC
<i>C. pauciramosum</i>	AF449448	AATGGTGTCT	ACGCCGGTAC	CTCCGAGCTC
<i>C. candelabrum</i>	AF210857	AATGGTGTCT	ACGCCGGTAC	CTCCGAGCTC
<i>C. candelabrum</i>	AF210858	AATGGTGTCT	ACGCCGGTAC	CTCCGAGCTC
<i>C. scoparium</i>	AF210875	AATGGTGTCT	ACGCCGGTAC	CTCCGAGCTC
<i>C. scoparium</i>	AF210874	AATGGTGTCT	ACGCCGGTAC	CTCCGAGCTC
<i>C. colhounii</i>	AF232855	AATGGTGTCT	ACGCCGGTAC	CTCCGAGCTC
<i>C. colhounii</i>	AF232854	AATGGTGTCT	ACGCCGGTAC	CTCCGAGCTC
<i>C. insulare</i>	AF44951	AATGGTGTCT	ACGCCGGTAC	CTCCGAGCTC
<i>C. insulare</i>	AF44950	AATGGTGTCT	ACGCCGGTAC	CTCCGAGCTC
<i>C. gracile</i>	AF333405	AATGGTGTCT	ACGCCGGTAC	CTCCGAGCTC
<i>C. gracile</i>	AF333404	AATGGTGTCT	ACGCCGGTAC	CTCCGAGCTC
<i>Cylindrocladiella infestans</i>	AF320190	AATGGTGTCT	ACAACGGCAG	CTCCGAGCTC
				360
CMW 12711		CAGCTCGAGC	GTATGAACGT	CTACTTCAAC
CMW 12709		CAGCTCGAGC	GTATGAACGT	CTACTTCAAC
CMW 12697		CAGCTCGAGC	GTATGAACGT	CTACTTCAAC
CMW 12689		CAGCTCGAGC	GTATGAACGT	CTACTTCAAC
CMW 13039		CAGCTCGAGC	GTATGAACGT	CTACTTCAAC
CMW 12702		CAGCTCGAGC	GTATGAACGT	CTACTTCAAC
CMW 12707		CAGCTCGAGC	GTATGAACGT	CTACTTCAAC
<i>C. pauciramosum</i>	AY162320	CAGCTCGAGC	GTATGAACGT	CTACTTCAAC
<i>C. pauciramosum</i>	AF449448	CAGCTCGAGC	GTATGAACGT	CTACTTCAAC
<i>C. candelabrum</i>	AF210857	CAGCTCGAGC	GTATGAACGT	CTACTTCAAC
<i>C. candelabrum</i>	AF210858	CAGCTCGAGC	GTATGAACGT	CTACTTCAAC
<i>C. scoparium</i>	AF210875	CAGCTCGAGC	GTATGAACGT	CTACTTCAAC
<i>C. scoparium</i>	AF210874	CAGCTCGAGC	GTATGAACGT	CTACTTCAAC
<i>C. colhounii</i>	AF232855	CAGCTCGAGC	GTATGAACGT	CTACTTCAAC
<i>C. colhounii</i>	AF232854	CAGCTCGAGC	GTATGAACGT	CTACTTCAAC
<i>C. insulare</i>	AF44951	CAGCTCGAGC	GTATGAACGT	CTACTTCAAC
<i>C. insulare</i>	AF44950	CAGCTCGAGC	GTATGAACGT	CTACTTCAAC
<i>C. gracile</i>	AF333405	CAGCTCGAGC	GTATGAACGT	CTACTTCAAC
<i>C. gracile</i>	AF333404	CAGCTCGAGC	GTATGAACGT	CTACTTCAAC
<i>Cylindrocladiella infestans</i>	AF320190	CAGCTCGAGC	GCATGAGCGT	CTACTTCAAC



				390
CMW 12711		GAGGTATGTG	AAAACCACTC	GAAGCACTCC
CMW 12709		GAGGTATGTG	AAAACCACTC	GAAGCACTCC
CMW 12697		GAGGTATGTG	AAAACCACTC	GAAGCACTCC
CMW 12689		GAGGTATGTG	AAAACCACTC	GAAGCACTCC
CMW 13039		GAGGTATGTG	AAAACCACTC	GAAGCACTCC
CMW 12702		GAGGTATGTG	AAAACCACTC	GAAGCACTCC
CMW 12707		GAGGTATGTG	AAAACCACTC	GAAGCACTCC
<i>C. pauciramosum</i>	AY162320	GAGGTATGTG	AAAACCACTC	GAAGCACTCC
<i>C. pauciramosum</i>	AF449448	GAGGTATGTG	AAAACCACTC	GAAGCACTCC
<i>C. candelabrum</i>	AF210857	GAGGTATGTG	AAAACCACTC	GAAGCACTCC
<i>C. candelabrum</i>	AF210858	GAGGTATGTG	AAAACCACTC	GAAGCACTCC
<i>C. scoparium</i>	AF210875	GAGGTATGTG	AAAACCACTC	GAAGCACTCC
<i>C. scoparium</i>	AF210874	GAGGTATGTG	AAAACCACTC	GAAGCACTCC
<i>C. colhounii</i>	AF232855	GAGGTATGTG	AAAACCACTC	GAAGCACTCC
<i>C. colhounii</i>	AF232854	GAGGTATGTG	AAAACCACTC	GAAGCACTCC
<i>C. insulare</i>	AF44951	GAGGTATGTG	AAAACCACTC	GAAGCACTCC
<i>C. insulare</i>	AF44950	GAGGTATGTG	AAAACCACTC	GAAGCACTCC
<i>C. gracile</i>	AF333405	GAGGTATGTG	AAAACCACTC	GAAGCACTCC
<i>C. gracile</i>	AF333404	GAGGTATGTG	AAAACCACTC	GAAGCACTCC
<i>Cylindrocladiella infestans</i>	AF320190	GAGGTATGTG	AAAACCACTC	GAAGCACTCC
			ACTATGGCAC	TCA – CATTG
				420
CMW 12711		CTTGACCGAG	AAGCACAAGC	CAACTCACAC
CMW 12709		CTTGACCGAG	AAGCACAAGC	CAACTCACAC
CMW 12697		CTTGACCGAG	AAGCACAAGC	CAACTCACAC
CMW 12689		CTTGACCGAG	AAGCACAAGC	CAACTCACAC
CMW 13039		CTTGACCGAG	AAGCACAAGC	CAACTCACAC
CMW 12702		CTTGACCGAG	AAGCACAAGC	CAACTCACAC
CMW 12707		CTTGACCGAG	AAGCACAAGC	CAACTCACAC
<i>C. pauciramosum</i>	AY162320	CTTGACCGAG	AAGCACAAGC	CAACTCACAC
<i>C. pauciramosum</i>	AF449448	CTTGACCGAG	AAGCACAAGC	CAACTCACAC
<i>C. candelabrum</i>	AF210857	CTTGACCGAG	AAGCACAATC	CGACTCACAC
<i>C. candelabrum</i>	AF210858	CTTGACCGAG	AAGCACAATC	CGACTCACAC
<i>C. scoparium</i>	AF210875	CACG–CCGAG	AGGCACAAGC	AAACTGACAC
<i>C. scoparium</i>	AF210874	CACG–CCGAG	AGGCACAAGC	AAACTGACAC
<i>C. colhounii</i>	AF232855	– TATGTCGAG	AGACGCAAGC	AAACTGACAC
<i>C. colhounii</i>	AF232854	CTATGTCGAG	AGACGCAACT	AAACTGACAC
<i>C. insulare</i>	AF44951	CACG–CCGAG	AGGCACAAGC	AAACTGACAC
<i>C. insulare</i>	AF44950	CACG–CCGAG	AGGCACAAGC	AAACTGACAC
<i>C. gracile</i>	AF333405	CTTGATCGAG	AGGCACAAGC	AAACTGACAC
<i>C. gracile</i>	AF333404	CTTGATCGAG	AGGCACAAGC	AAACTGACAC
<i>Cylindrocladiella infestans</i>	AF320190	CTACTACTGTG	AAATCAGAAT	GTACTCACGC
				450
CMW 12711		CATCATGTAG	GCTTCCGGCA	ACAAGTTCGT
CMW 12709		CATCATGTAG	GCTTCCGGCA	ACAAGTTCGT
CMW 12697		CATCATGTAG	GCTTCCGGCA	ACAAGTTCGT
CMW 12689		CATCATGTAG	GCTTCCGGCA	ACAAGTTCGT
CMW 13039		CATCATGTAG	GCTTCCGGCA	ACAAGTTCGT
CMW 12702		CATCATGTAG	GCTTCCGGCA	ACAAGTTCGT
CMW 12707		CATCATGTAG	GCTTCCGGCA	ACAAGTTCGT
<i>C. pauciramosum</i>	AY162320	CATCATGTAG	GCTTCCGGCA	ACAAGTTCGT
<i>C. pauciramosum</i>	AF449448	CATCATGTAG	GCTTCCGGCA	ACAAGTTCGT
<i>C. candelabrum</i>	AF210857	CATCATGTAG	GCTTCCGGCA	ACAAGTTCGT
<i>C. candelabrum</i>	AF210858	CATCATGTAG	GCTTCCGGCA	ACAAGTTCGT
<i>C. scoparium</i>	AF210875	CAT---GTAG	GCTTCTGGCA	ACAAGTTCGT
<i>C. scoparium</i>	AF210874	CAT---GTAG	GCTTCTGGCA	ACAAGTTCGT
<i>C. colhounii</i>	AF232855	CAT---GTAG	GCTTCCGGCA	ACAAGTTCGT
<i>C. colhounii</i>	AF232854	CAT---GTAG	GCCTTCCGGCA	ACAAGTTCGT
<i>C. insulare</i>	AF44951	CAT---GTAG	GCTTCTGGCA	ACAAGTTCGT
<i>C. insulare</i>	AF44950	CAT---GTAG	GCTTCTGGCA	ACAAGTTCGT
<i>C. gracile</i>	AF333405	CAT---GTAG	GCTTCTGGCA	ACAAGTTCGT
<i>C. gracile</i>	AF333404	CAT---GTAG	GCTTCTGGCA	ACAAGTTCGT
<i>Cylindrocladiella infestans</i>	AF320190	---TCCGTAG	GCTTCTGGCA	ACAAGTATGT



CMW 12711		TCCTCGCGCT	GTCCTCGTCC	ATCTTGAGCC	480
CMW 12709		TCCTCGCGCT	GTCCTCGTCC	ATCTTGAGCC	
CMW 12697		TCCTCGCGCT	GTCCTCGTCC	ATCTTGAGCC	
CMW 12689		TCCTCGCGCT	GTCCTCGTCC	ATCTTGAGCC	
CMW 13039		TCCTCGCGCT	GTCCTCGTCC	ATCTTGAGCC	
CMW 12702		TCCTCGCGCT	GTCCTCGTCC	ATCTTGAGCC	
CMW 12707		TCCTCGCGCT	GTCCTCGTCC	ATCTTGAGCC	
<i>C. pauciramosum</i>	AY162320	TCCTCGCGCT	GTCCTCGTCC	ATCTTGAGCC	
<i>C. pauciramosum</i>	AF449448	TCCTCGCGCT	GTCCTCGTCC	ATCTTGAGCC	
<i>C. candelabrum</i>	AF210857	TCCTCGCGCT	GTCCTCGTCC	ATCTTGAGCC	
<i>C. candelabrum</i>	AF210858	TCCTCGCGCT	GTCCTCGTCC	ATCTTGAGCC	
<i>C. scoparium</i>	AF210875	TCCTCGCGCT	GTCCTCGTCC	ATCTTGAGCC	
<i>C. scoparium</i>	AF210874	TCCTCGCGCT	GTCCTCGTCC	ATCTTGAGCC	
<i>C. colhounii</i>	AF232855	TCCTCGCGCT	GTCCTCGTCC	ATCTTGAGCC	
<i>C. colhounii</i>	AF232854	TCCTCGCGCT	GTCCTCGTCC	ATCTTGAGCC	
<i>C. insulare</i>	AF44951	TCCTCGCGCT	GTCCTCGTCC	ATCTTGAGCC	
<i>C. insulare</i>	AF44950	TCCTCGCGCT	GTCCTCGTCC	ATCTTGAGCC	
<i>C. gracile</i>	AF333405	TCCTCGCGCT	GTCCTCGTCC	ATCTTGAGCC	
<i>C. gracile</i>	AF333404	TCCTCGCGCT	GTCCTCGTCC	ATCTTGAGCC	
<i>Cylindrocladiella infestans</i>	AF320190	CCCTCGCGCC	GTCCTCGTCC	ATCTTGAGCC	
CMW 12711		CGGTACCATG	GACGCCGTCC	GTGCCGGTCC	510
CMW 12709		CGGTACCATG	GACGCCGTCC	GTGCCGGTCC	
CMW 12697		CGGTACCATG	GACGCCGTCC	GTGCCGGTCC	
CMW 12689		CGGTACCATG	GACGCCGTCC	GTGCCGGTCC	
CMW 13039		CGGTACCATG	GACGCCGTCC	GTGCCGGTCC	
CMW 12702		CGGTACCATG	GACGCCGTCC	GTGCCGGTCC	
CMW 12707		CGGTACCATG	GACGCCGTCC	GTGCCGGTCC	
<i>C. pauciramosum</i>	AY162320	CGGTACCATG	GACGCCGTCC	GTGCCGGTCC	
<i>C. pauciramosum</i>	AF449448	CGGTACCATG	GACGCCGTCC	GTGCCGGTCC	
<i>C. candelabrum</i>	AF210857	CGGTACCATG	GACGCCGTCC	GTGCCGGTCC	
<i>C. candelabrum</i>	AF210858	CGGTACCATG	GACGCCGTCC	GTGCCGGTCC	
<i>C. scoparium</i>	AF210875	CGGTACCATG	GACGCCGTCC	GTGCCGGTCC	
<i>C. scoparium</i>	AF210874	CGGTACCATG	GACGCCGTCC	GTGCCGGTCC	
<i>C. colhounii</i>	AF232855	CGGTACCATG	GACGCCGTCC	GTGCCGGTCC	
<i>C. colhounii</i>	AF232854	CGGTACCATG	GACGCCGTCC	GTGCCGGTCC	
<i>C. insulare</i>	AF44951	CGGTACCATG	GACGCCGTCC	GTGCCGGTCC	
<i>C. insulare</i>	AF44950	CGGTACCATG	GACGCCGTCC	GTGCCGGTCC	
<i>C. gracile</i>	AF333405	CGGTACCATG	GACGCCGTCC	GTGCCGGTCC	
<i>C. gracile</i>	AF333404	CGGTACCATG	GACGCCGTCC	GTGCCGGTCC	
<i>Cylindrocladiella infestans</i>	AF320190	CGGTACCATG	GACGCCGTCC	GTGCCGGTCC	
CMW 12711		TTTCGGTCAG	CTCTTCGGCC		520
CMW 12709		TTTCGGTCAG	CTCTTCGGCC		
CMW 12697		TTTCGGTCAG	CTCTTCGGCC		
CMW 12689		TTTCGGTCAG	CTCTTCGGCC		
CMW 13039		TTTCGGTCAG	CTCTTCGGCC		
CMW 12702		TTTCGGTCAG	CTCTTCGGCC		
CMW 12707		TTTCGGTCAG	CTCTTCGGCC		
<i>C. pauciramosum</i>	AY162320	TTTCGGTCAG	CTCTTCGGCC		
<i>C. pauciramosum</i>	AF449448	TTTCGGTCAG	CTCTTCGGCC		
<i>C. candelabrum</i>	AF210857	TTTCGGTCAG	CTCTTCGGCC		
<i>C. candelabrum</i>	AF210858	TTTCGGTCAG	CTCTTCGGCC		
<i>C. scoparium</i>	AF210875	TTTCGGTCAG	CTCTTCGGCC		
<i>C. scoparium</i>	AF210874	TTTCGGTCAG	CTCTTCGGCC		
<i>C. colhounii</i>	AF232855	TTTCGGTCAG	CTCTTCGGCC		
<i>C. colhounii</i>	AF232854	TTTCGGTCAG	CTCTTCGGCC		
<i>C. insulare</i>	AF44951	TTTCGGTCAG	CTCTTCGGCC		
<i>C. insulare</i>	AF44950	TTTCGGTCAG	CTCTTCGGCC		
<i>C. gracile</i>	AF333405	TTTCGGTCAG	CTCTTCGGCC		
<i>C. gracile</i>	AF333404	TTTCGGTCAG	CTCTTCGGCC		
<i>Cylindrocladiella infestans</i>	AF320190	TTTCGGTCAG	CTCTTCGGCC		



Alignment 2. The EF1 – α gene DNA sequence alignment of the *Fusarium* isolates obtained from the *Eucalyptus* hedge plants.

FCC 3110		GAAGTTCGAG	AAGGTTGGT-	CACATCTCCC	30
FCC 3114		GAAGTTCGAG	AAGGTTGGT-	CACATCTCCC	
FCC 3119		GAAGTTCGAG	AAGGTTAGT-	CAATATCCCT	
FCC 3109		GAAGTTCGAG	AAGGTTAGT-	CACTTTCCCT	
FCC 3117		GAAGTTCGAG	AAGGTTAGT-	CAATATCCCT	
FCC 2957		GAAGTTCGAG	AAGGTTAGT-	CACTTTCCCT	
FCC 3115		GAAGTTCGAG	AAGGTTAGT-	CACTTTCCCT	
FCC 2964		GAAGTTCGAG	AAGGTTAGT-	CACTTTCCCT	
FCC 3113		GAAGTTCGAG	AAGGTTAGT-	CACTTTCCCT	
<i>F. avenaceum</i>	AY337423	GAAGTTCGAG	AAGGTTAGT-	CAATATCCCT	
<i>F. equiseti</i>	AY337424	GAAGTTCGAG	AAGGTTAGT-	CAATATCCCT	
<i>F. lateritium</i>	AY337435	GAAGTTCGAG	AAGGTTAGT-	CAATATCCCT	
<i>F. nygamai</i>	AY337445	GAAGTTCGAG	AAGGTTAGT-	CACTTTCCCT	
<i>F. oxysporum</i>	AY337428	GAAGTTCGAG	AAGGTTAGT-	CACTTTCCCT	
<i>F. proliferatum</i>	AY337436	GAAGTTCGAG	AAGGTTAGT-	CACTTTCCCT	
<i>F. sambucinum</i>	AY337422	GAAGTTCGAG	AAGGTTAGT-	CAAAATCCCT	
<i>F. solani</i>	AY337438	GAAGTTCGAG	AAGGTTGGT-	CACATCTCCC	
<i>F. verticillioides</i>	AY337450	GAAGTTCGAG	AAGGTTAGT-	CACTTTCCCT	
<i>Botryosphaeria_dothidea</i>	AY236899	GAAGTTCGAG	AAGGTAAGCA	CACATTTTCT	

FCC 3110		CCGATCGCGC	CT ---- TGCT	ATCCACATC	60
FCC 3114		CCGATCGCGC	CT ---- TGCT	ATCCACATC	
FCC 3119		TCGATTACGC	-----GCG	CTCCCA--TC	
FCC 3109		TCAATCGCGC	GT----CCTT	TGCCCA--TC	
FCC 3117		TCGATTACGC	-----GCG	CTCCCA--TC	
FCC 2957		TCTATCGCGC	GT ---- TCTT	TGCCCA--TC	
FCC 3115		TCAATCGCGC	GT ---- CCTT	TGCCCA--TC	
FCC 2964		TCTATCGCGC	GT ---- TCTT	TGCCCA--TC	
FCC 3113		TCGATCGCGC	GT ---- CCTT	TGTCCA--TC	
<i>F. avenaceum</i>	AY337423	TCGATTACGC	-----GCG	CTCCCA--TC	
<i>F. equiseti</i>	AY337424	TCGATTACGC	-----GCG	CTCCCA--TC	
<i>F. lateritium</i>	AY337435	TCGATTACGC	-----GCG	CTCCCA--TC	
<i>F. nygamai</i>	AY337445	TCGATCGCGC	GT ---- CCTT	TGTCCA--TC	
<i>F. oxysporum</i>	AY337428	TCAATCGCGC	GT ---- CCTT	TGCCCA--TC	
<i>F. proliferatum</i>	AY337436	TCGATCGCGC	GT ---- CCTC	TGCCCA--CC	
<i>F. sambucinum</i>	AY337422	TCGATTGCGC	-----GCG	CTCCCA--TC	
<i>F. solani</i>	AY337438	CCGATCGCGC	CT ---- TGCT	ATCCACATC	
<i>F. verticillioides</i>	AY337450	TCTATCGCGC	GT ---- TCTT	TGCCCA--TC	
<i>Botryosphaeria_dothidea</i>	AY236899	GTGCCTGCAC	GTGTGCTGGG	TTCTGCGCC	

FCC 3110		GAATTCCTCCG	TCGAATTCCC	TCCTCCGCGA	90
FCC 3114		GAATTCCTCCG	TCGAATTCCC	TCCTCCGCGA	
FCC 3119		GATTCCTCCG	ATTCGCTCCC	TCACTCGAAA	
FCC 3109		GATTCCTCC-	-----TA	CGACTCGAAA	
FCC 3117		GATTCCTCCG	ATTCGCTCCC	TCACTCGAAA	
FCC 2957		GATTCCTCCC	-----TA	CGACTCGAAA	
FCC 3115		GATTCCTCC-	-----TA	CGACTCGAAA	
FCC 2964		GATTCCTCCC	-----TA	CGACTCGAAA	
FCC 3113		GATTCCTCCC	-----TC	CGACTCGAAA	
<i>F. avenaceum</i>	AY337423	GATTCCTCCG	ACTCGCTCCC	TCATTGAAA	
<i>F. equiseti</i>	AY337424	GATTCCTCCG	ACTCGCTCCC	TCATTGAAA	
<i>F. lateritium</i>	AY337435	GATTCCTCCG	ATTCGCTCCC	TCACTCGAAA	
<i>F. nygamai</i>	AY337445	GATTCCTCCC	-----TC	CGACTCGAAA	
<i>F. oxysporum</i>	AY337428	GATTCCTCC-	-----TA	CGACTCGAAA	
<i>F. proliferatum</i>	AY337436	GATTCCTCC-	-----TG	CGATTTGAAA	
<i>F. sambucinum</i>	AY337422	GATTCATACG	ACTCGCTCCC	TCACTCGAAA	
<i>F. solani</i>	AY337438	GAATTCCTCCG	TCGAATTCCC	TCCTCCGCGA	
<i>F. verticillioides</i>	AY337450	GATTCCTCCC	-----TA	CGACTCGAAA	
<i>Botryosphaeria_dothidea</i>	AY236899	GAATTCCTCC	TATCA-- CTC	TGGTGAGGGG	



FCC 3110		CACGCTCTGC	GCCCGCTTCT	CCCGAGTCCC	120
FCC 3114		CACGCTCTGC	GCCCGCTTCT	CCCGAGTCCC	
FCC 3119		CACATCCATT	ACCC— —CG	CTCGAGTCC—	
FCC 3109		CGTGCCCGCT	ACCC— —CG	CTCGAGACC—	
FCC 3117		CACATCCATT	ACCC— —CG	CTCGAGTCC—	
FCC 2957		CGTACCCGCT	ACCC— —CG	CTCGAGCCC—	
FCC 3115		CGTGCCCGCT	ACCC— —CG	CTCGAGACC—	
FCC 2964		CGTACCCGCT	ACCC— —CG	CTCGAGCCC—	
FCC 3113		CGTGCCCGCT	ACCC— —CG	CTCGAATTC—	
<i>F. avenaceum</i>	AY337423	CGCATTCATT	ACCC— —CG	CTCAAGTCC—	
<i>F. equiseti</i>	AY337424	CGCATTCATT	ACCC— —CG	CTCAAGTCC—	
<i>F. lateritium</i>	AY337435	CACATCCATT	ACCC— —CG	CTCGAGTCC—	
<i>F. nygamai</i>	AY337445	CGTGCCCGCT	ACCC— —CG	CTCGAATTC—	
<i>F. oxysporum</i>	AY337428	CGTGCCCGCT	ACCC— —CG	CTCGAGACC—	
<i>F. proliferatum</i>	AY337436	CGTGCCCGCT	ACCC— —CG	CTCGAGACC—	
<i>F. sambucinum</i>	AY337422	CGCATTCATT	ACCC— —CG	CTCGAGACC—	
<i>F. solani</i>	AY337438	CACGCTCTGC	GCCCGCTTCT	CTCGAGCCC—	
<i>F. verticillioides</i>	AY337450	CGTACCCGCT	ACCC— —CG	CCCGAGTCCC	
<i>Botryosphaeria_ dothidea</i>	AY236899	CAATTTCTTG	GTGGGGCTGG	CTCGAGCCC—	
				CCCAGTCCC	

FCC 3110		AAAAATTTTG	CGGTCGACC	GTAATTTTTT	150
FCC 3114		AAAAATTTTG	CGGTCGACC	GTAATTTTTT	
FCC 3119		GAAAATTTTG	CGGTGCGACC	GTGATTTTTT	
FCC 3109		AAAAATTTTG	CAATATGACC	GTAATTTTTT	
FCC 3117		GAAAATTTTG	CGGTGCGACC	GTGATTTTTT	
FCC 2957		AAAAATTTTG	CGATACGACC	GTAATTTTTT	
FCC 3115		AAAAATTTTG	CAATATGACC	GTAATTTTTT	
FCC 2964		AAAAATTTTG	CGATACGACC	GTAATTTTTT	
FCC 3113		AAAAATTTTG	CGATATGACC	GTAATTTTTT	
<i>F. avenaceum</i>	AY337423	GAAAATTTTG	CGGTGCGACC	GTGATTTTTT	
<i>F. equiseti</i>	AY337424	GAAAATTTTG	CGGTGCGACC	GTGATTTTTT	
<i>F. lateritium</i>	AY337435	GAAAATTTTG	CGGTGCGACC	GTGATTTTTT	
<i>F. nygamai</i>	AY337445	AAAAATTTTG	CGATATGACC	GTAATTTTTT	
<i>F. oxysporum</i>	AY337428	AAAAATTTTG	CAATATGACC	GTAATTTTTT	
<i>F. proliferatum</i>	AY337436	AAAAATTTTG	CGATATGACC	GTAATTTTTT	
<i>F. sambucinum</i>	AY337422	AAAAATTTTG	CGGTGCGACC	GTGATTTTTT	
<i>F. solani</i>	AY337438	AAAAATTTTG	CGGTCGACC	GTAATTTTTT	
<i>F. verticillioides</i>	AY337450	AAAAATTTTG	CGATACGACC	GTAATTTTTT	
<i>Botryosphaeria_ dothidea</i>	AY236899	GCCTCGTTTG	GTCTTCGGCA	GTAATTTTTT	
				A—AATCTCCG	

FCC 3110		TTGGTGGGGC	AT—TTACCCC	GCCACTCGGG	180
FCC 3114		TTGGTGGGGC	AT—TTACCCC	GCCACTCGGG	
FCC 3119		CTGGTGGGGT	ATCTTACCCC	GCCACTCGAG	
FCC 3109		—TGGTGGGGC	AC—TTACCCC	GCCACTTGAG	
FCC 3117		CTGGTGGGGT	ATCTTACCCC	GCCACTCGAG	
FCC 2957		—TGGTGGGGC	AT—TTATCCC	GCCACTCGAG	
FCC 3115		—TGGTGGGGC	AC—TTACCCC	GCCACTTGAG	
FCC 2964		—TGGTGGGGC	AT—TTATCCC	GCCACTCGAG	
FCC 3113		—TGGTGGGGC	AT—TTACCCC	GCCACTCGAG	
<i>F. avenaceum</i>	AY337423	TTGGTGGGGT	ATCTTACCCC	GCCACTCGAG	
<i>F. equiseti</i>	AY337424	TTGGTGGGGT	ATCTTACCCC	GCCACTCGAG	
<i>F. lateritium</i>	AY337435	CTGGTGGGGT	ATCTTACCCC	GCCACTCGAG	
<i>F. nygamai</i>	AY337445	—TGGTGGGGC	AT—TTACCCC	GCCACTCGAG	
<i>F. oxysporum</i>	AY337428	—TGGTGGGGC	AC—TTACCCC	GCCACTTGAG	
<i>F. proliferatum</i>	AY337436	—TGGTGGGGC	AT—TTACCCC	GCCACTCGAG	
<i>F. sambucinum</i>	AY337422	TTGGTGGGGT	ATCTTACCCC	GCCACTCGAG	
<i>F. solani</i>	AY337438	TTGGTGGGGC	AT—TTACCCC	GCCACTCGGG	
<i>F. verticillioides</i>	AY337450	—TGGTGGGGC	AT—TTATCCC	GCCACTCGAG	
<i>Botryosphaeria_ dothidea</i>	AY236899	CATCTGGATT	TTTTGTGACC	GGCGTGGCAC	



FCC 3110		CGACGTTGGA	CAAAGCCCTG	ATCCCTGCAC	210
FCC 3114		CGACGTTGGA	CAAAGCCCTG	ATCCCTGCAC	
FCC 3119		TCACGGATGC	GCTTGCCCTG	TTCCC— —	
FCC 3109		CGACGGGAGC	GTTTGCCCTC	TTACC-ATTC	
FCC 3117		TCACGGATGC	GCTTGCCCTG	TTCCC— —	
FCC 2957		CGGCGC—GT	TTCTGCCCTC	TC-CC-ATTC	
FCC 3115		CGACGGGAGC	GTTTGCCCTC	TTACC-ATTC	
FCC 2964		CGGCGC—GT	TTCTGCCCTC	TC-CC-ATTC	
FCC 3113		CGGCGC—GT	TTTTGCCCTC	TTCCC-ATTC	
<i>F. avenaceum</i>	AY337423	TGACGGATGC	GCTTGCCCTG	TTCCC— —	
<i>F. equiseti</i>	AY337424	TGACGGATGC	GCTTGCCCTG	TTCCC— —	
<i>F. lateritium</i>	AY337435	TCACGGATGC	GCTTGCCCTG	TTCCC— —	
<i>F. nygamai</i>	AY337445	CGGCGC—T	TTTTGCCCTC	TTCCC-ATTC	
<i>F. oxysporum</i>	AY337428	CGACGGGAGC	GTTTGCCCTC	TTACC-ATTC	
<i>F. proliferatum</i>	AY337436	CGATGGGCGC	GTTTTGCCCTC	TTCC-TGTC	
<i>F. sambucinum</i>	AY337422	TGACGGATGC	GCTTGCCCTG	TTCCC— —	
<i>F. solani</i>	AY337438	CGACGTTGGA	CAAAGCCCTG	ATCCCTGCAC	
<i>F. verticillioides</i>	AY337450	CGGCGC—T	TTCTGCCCTC	TC-CC-ATTC	
<i>Botryosphaeria_dothidea</i>	AY236899	CGACGCGAAC	A-----CCCC	TCACC— —	

FCC 3110		ACAAAAACAC	CA - AACCCCTC	TTGGCGCGCA	240
FCC 3114		ACAAAAACAC	CA - AACCCCTC	TTGGCGCGCA	
FCC 3119		ACAAAACCTT	----ACCACC	CTGTGCGGCA	
FCC 3109		TCAGAACCTC	AATGAGTGCG	TCGTACCGTG	
FCC 3117		ACAAAACCTT	----ACCACC	CTGTGCGGCA	
FCC 2957		-CACAACTC	ACTGAGCTCA	TCGTACCGTG	
FCC 3115		TCAGAACCTC	AATGAGTGCG	TCGTACCGTG	
FCC 2964		-CACAACTC	ACTGAGCTCA	TCGTACCGTG	
FCC 3113		-CACAACTC	ACTGAGCGCA	TCGTACCGTG	
<i>F. avenaceum</i>	AY337423	ACAAAACCTC	----ACCACA	CTGTGCGGCA	
<i>F. equiseti</i>	AY337424	ACAAAACCTC	----ACCACA	CTGTGCGGCA	
<i>F. lateritium</i>	AY337435	ACAAAACCTT	----ACCACC	CTGTGCGGCA	
<i>F. nygamai</i>	AY337445	-CACAACTC	ACTGAGCGCA	TCGTACCGTG	
<i>F. oxysporum</i>	AY337428	TCAGAACCTC	AATGAGTGCG	TCGTACCGTG	
<i>F. proliferatum</i>	AY337436	-CACAACTC	AATGAGCGCA	TTGTACCGTG	
<i>F. sambucinum</i>	AY337422	ACAAAACCTT	----ACTACC	CTGTGCGGCA	
<i>F. solani</i>	AY337438	ACAAAAACAC	CA - AACCCCTC	TTGGCGCGCA	
<i>F. verticillioides</i>	AY337450	-CACAACTC	ACTGAGCTCA	TCGTACCGTG	
<i>Botryosphaeria_dothidea</i>	AY236899	--AAGCTTC	CAGCCACTCA	CGTTCGTCTA	

FCC 3110		TCACGTGGTT	CACAACAGAC	ACTGACTGGT	270
FCC 3114		TCACGTGGTT	CACAACAGAC	ACTGACTGGT	
FCC 3119		CTACATGTCT	T--GCAGTC	ACTAACCA-C	
FCC 3109		TCAAGC--	----AGTC	ACTAACCA-T	
FCC 3117		CTACATGTCT	T--GCAGTC	ACTAACCA-C	
FCC 2957		TCAAGC--	----AGTC	ACTAACCA-T	
FCC 3115		TCAAGC--	----AGTC	ACTAACCA-T	
FCC 2964		TCAAGC--	----AGTC	ACTAACCA-T	
FCC 3113		TCAAGC--	----AGTC	ACTAACCA-T	
<i>F. avenaceum</i>	AY337423	CTATGTCTT -	----GCAGTC	ACTAACCA-C	
<i>F. equiseti</i>	AY337424	CTATGTCTT -	----GCAGTC	ACTAACCA-C	
<i>F. lateritium</i>	AY337435	CTACATGTCT	T--GCAGTC	ACTAACCA-C	
<i>F. nygamai</i>	AY337445	TCAAGC--	----AGTC	ACTAACCA-T	
<i>F. oxysporum</i>	AY337428	TCAAGC--	----AGTC	ACTAACCA-T	
<i>F. proliferatum</i>	AY337436	TCAAGC--	----AGTC	ACTAACCA-T	
<i>F. sambucinum</i>	AY337422	CTATCATATG	TCTTCCAGTC	ACTAACCA-C	
<i>F. solani</i>	AY337438	TCACGTGGTT	CACAACAGAC	ACTGACTGGT	
<i>F. verticillioides</i>	AY337450	TCAAGC--	----AGTC	ACTAACCA-T	
<i>Botryosphaeria_dothidea</i>	AY236899	TGCGAC--	----CATAT	GCTAACCACC	



		280
FCC 3110		TCAACAATAG
FCC 3114		TCAACAATAG
FCC 3119		TGGACAATAG
FCC 3109		TCAACAATAG
FCC 3117		TGGACAATAG
FCC 2957		CCGACAATAG
FCC 3115		TCAACAATAG
FCC 2964		CCGACAATAG
FCC 3113		TCGACAATAG
<i>F. avenaceum</i>	AY337423	TGGACAATAG
<i>F. equiseti</i>	AY337424	TGGACAATAG
<i>F. lateritium</i>	AY337435	TGGACAATAG
<i>F. nygamai</i>	AY337445	TCGACAATAG
<i>F. oxysporum</i>	AY337428	TCAACAATAG
<i>F. proliferatum</i>	AY337436	TCGACAATAG
<i>F. sambucinum</i>	AY337422	TGGACAATAG
<i>F. solani</i>	AY337438	TCAACAATAG
<i>F. verticillioides</i>	AY337450	CCGACAATAG
<i>Botryosphaeria_dothidea</i>	AY236899	GCCACAACAG



Alignment 3. The combined ITS1, 5.8s subunit and ITS2 regions of the genomic RNA gene DNA sequence of the *Phytophthora* isolates obtained from the *Eucalyptus* hedge plants .

CMW 13800		GTTGGGGGTC	TTATTT-GGC	GGCGGCTGCT	30
CMW 13816		GTTGGGGGTC	TTATTT-GGC	GGCGGCTGCT	
CMW 13804		GTTGGGGGCC	TGCTCTGGGC	GGCGTTGTC	
<i>P. cinnamomi</i>	AF266764	GTTGGGGGCC	TGCTCTGGGC	GGCGTTGTC	
<i>P. cryptogea</i>	AF266796	ATTTGGGGGC	TTCCGTCTG-	GCCGGCCG-	
<i>P. infestans</i>	AF266779	GTTGGGGGTC	T-TACTTGGC	GGCGGCTGCT	
<i>P. nicotianae</i>	AF266776	GTTGGGGGTC	TTATTT-GGC	GGCGGCTGCT	
<i>P. palmivora</i>	AF266780	TTTGGGGGTC	T-CTTTCGGC	GGCGGCTGCT	
<i>Pyhtium aphanidermatum</i>	AJ233438	GTTCTGTGCT	CTCTTTCGG-	GAGGGCTG-	
CMW 13800		GGCTTAATTG	TTGGCGGCTG	CTGCTGAGTG	60
CMW 13816		GGCTTAATTG	TTGGCGGCTG	CTGCTGAGTG	
CMW 13804		GATGTCAAAG	TCGACGGTTG	CTGTTGCGTG	
<i>P. cinnamomi</i>	AF266764	GATGTCAAAG	TCGACGGTTG	CTGTTGCGTG	
<i>P. cryptogea</i>	AF266796	-----G	TTTTCGGCTG	-GCTGGGTG	
<i>P. infestans</i>	AF266779	GGCTTTATTG	CTGGCGGCTA	CTGCTG- G	
<i>P. nicotianae</i>	AF266776	GGCTTAATTG	TTGGCGGCTG	CTGCTGAGTG	
<i>P. palmivora</i>	AF266780	GGCTTCATTG	CTGGCGGCTG	CTGTTG- G	
<i>Pyhtium aphanidermatum</i>	AJ233438	-----A	ACGAAGGTGG	- GCTG -	
CMW 13800		AG -- CCCTA	TCAAAAAAAAA	GGCGAACGTT	90
CMW 13816		AG -- CCCTA	TCAAAAAAAAA	GGCGAACGTT	
CMW 13804		GGGGGCCCTA	TCAC --- T	GGCGAGCGTT	
<i>P. cinnamomi</i>	AF266764	GGGGGCCCTA	TCAC --- T	GGCGAGCGTT	
<i>P. cryptogea</i>	AF266796	GCGG- CTCTA	TCA --- T	GGCGACCGCT	
<i>P. infestans</i>	AF266779	GCGAGCCCTA	TCAAA -A	GGCGAGCGTT	
<i>P. nicotianae</i>	AF266776	AG -- CCCTA	TCAAAAAAAAA	GGCGAACGTT	
<i>P. palmivora</i>	AF266780	GAGAGCTCTA	TCA --- T	GGCGAGCGTT	
<i>Pyhtium aphanidermatum</i>	AJ233438	---- CTTAA	TTG --- T	AGT - - - -	
CMW 13800		TGGGCTTC -	--- GGCC	TGATTTAGTA	120
CMW 13816		TGGGCTTC -	--- GGCC	TGATTTAGTA	
CMW 13804		TGGGTCCCCC	TCGGGGGAAC	TGAGCTAGTA	
<i>P. cinnamomi</i>	AF266764	TGGGTCCCCC	TCGGGGGAAC	TGAGCTAGTA	
<i>P. cryptogea</i>	AF266796	TGGGCTTC -	--- GGCC	TGGGCTAGTA	
<i>P. infestans</i>	AF266779	TGGACTTC -	--- GGTC	TGAGCTAGTA	
<i>P. nicotianae</i>	AF266776	TGGGCTTC -	--- GGCC	TGATTTAGTA	
<i>P. palmivora</i>	AF266780	TGGGCTTC -	--- GGTC	TGAAC TAGTA	
<i>Pyhtium aphanidermatum</i>	AJ233438	-----	--- ---	--- CTGCCG	
CMW 13800		GTCTTTTTTT	CTTTTAAACC	CATTC-CTTA	150
CMW 13816		GTCTTTTTTT	CTTTTAAACC	CATTC-CTTA	
CMW 13804		GCCTCTCTTT	T--- AAACC	CATCC-TGTA	
<i>P. cinnamomi</i>	AF266764	GCCTCTCT-	- TTTAAACC	CATCC-TGTA	
<i>P. cryptogea</i>	AF266796	GCGTATTTTT	--- AAACC	CATTC-CTAA	
<i>P. infestans</i>	AF266779	GCTTTTTTAT	- TTTAAACC	CTTTA-CTTA	
<i>P. nicotianae</i>	AF266776	GTCTTTTTTT	CTTTTAAACC	CATTC-CTTA	
<i>P. palmivora</i>	AF266780	GC---TTT-	- TTTAAACC	CATTC-TTTA	
<i>Pyhtium aphanidermatum</i>	AJ233438	ATGTATTTTT	C---AAACC	CATTTACCTA	
CMW 13800		ATACTGAA-T	ATACTGTGGG	GACGAAAGTC	180
CMW 13816		ATACTGAA-T	ATACTGTGGG	GACGAAAGTC	
CMW 13804		ATACTGAA-C	ATACTGTGGG	GACGAAAGTC	
<i>P. cinnamomi</i>	AF266764	ATACTGAA-C	ATACTGTGGG	GACGAAAGTC	
<i>P. cryptogea</i>	AF266796	T TACTGAA-T	ATACTGTGGG	GACGAAAGTC	
<i>P. infestans</i>	AF266779	ATACTGAT-T	ATACTGTGGG	GACGAAAGTC	
<i>P. nicotianae</i>	AF266776	ATACTGAA-T	ATACTGTGGG	GACGAAAGTC	
<i>P. palmivora</i>	AF266780	TAACTGAT-T	ATACTGTAGG	GACGAAAGTC	
<i>Pyhtium aphanidermatum</i>	AJ233438	ATACTGATCT	ATACTCCAAA	AACGAAAGTT	



CMW 13800		TCTGCTTTTA	ACTAGATAGC	AACCTTCAGC	210
CMW 13816		TCTGCTTTTA	ACTAGATAGC	AACCTTCAGC	
CMW 13804		TCTGCTTTTA	ACTAGATAGC	AACCTTCAGC	
<i>P. cinnamomi</i>	AF266764	TCTGCTTTTA	ACTAGATAGC	AACCTTCAGC	
<i>P. cryptogea</i>	AF266796	TCTGCTTTTA	ACTAGATAGC	AACCTTCAGC	
<i>P. infestans</i>	AF266779	TCTGCTTTTA	ACTAGATAGC	AACCTTCAGC	
<i>P. nicotianae</i>	AF266776	TCTGCTTTTA	ACTAGATAGC	AACCTTCAGC	
<i>P. palmivora</i>	AF266780	TCTGCTTTTA	ACTAGATAGC	AACCTTCAGC	
<i>Pyhtium aphanidermatum</i>	AJ233438	TATGGTTTTA	ATCT – ATAAC	AACCTTCAGC	
CMW 13800		AGTGGATGTC	TAGGCTCGCA	CATCGATGAA	240
CMW 13816		AGTGGATGTC	TAGGCTCGCA	CATCGATGAA	
CMW 13804		AGTGGATGTC	TAGGCTCGCA	CATCGATGAA	
<i>P. cinnamomi</i>	AF266764	AGTGGATGTC	TAGGCTCGCA	CATCGATGAA	
<i>P. cryptogea</i>	AF266796	AGTGGATGTC	TAGGCTCGCA	CATCGATGAA	
<i>P. infestans</i>	AF266779	AGTGGATGTC	TAGGCTCGCA	CATCGATGAA	
<i>P. nicotianae</i>	AF266776	AGTGGATGTC	TAGGCTCGCA	CATCGATGAA	
<i>P. palmivora</i>	AF266780	AGTGGATGTC	TAGGCTCGCA	CATCGATGAA	
<i>Pyhtium aphanidermatum</i>	AJ233438	AGTGGATGTC	TAGGCTCGCA	CATCGATGAA	
CMW 13800		GAACGCTGCG	AACTGCGATA	CGTAATGCGA	270
CMW 13816		GAACGCTGCG	AACTGCGATA	CGTAATGCGA	
CMW 13804		GAACGCTGCG	AACTGCGATA	CGTAATGCGA	
<i>P. cinnamomi</i>	AF266764	GAACGCTGCG	AACTGCGATA	CGTAATGCGA	
<i>P. cryptogea</i>	AF266796	GAACGCTGCG	AACTGCGATA	CGTAATGCGA	
<i>P. infestans</i>	AF266779	GAACGCTGCG	AACTGCGATA	CGTAATGCGA	
<i>P. nicotianae</i>	AF266776	GAACGCTGCG	AACTGCGATA	CGTAATGCGA	
<i>P. palmivora</i>	AF266780	GAACGCTGCG	AACTGCGATA	CGTAATGCGA	
<i>Pyhtium aphanidermatum</i>	AJ233438	GAACGCTGCG	AACTGCGATA	CGTAATGCGA	
CMW 13800		ATTGCAGGAT	TCAGTGAGTC	ATCGAAATTT	300
CMW 13816		ATTGCAGGAT	TCAGTGAGTC	ATCGAAATTT	
CMW 13804		ATTGCAGGAT	TCAGTGAGTC	ATCGAAATTT	
<i>P. cinnamomi</i>	AF266764	ATTGCAGGAT	TCAGTGAGTC	ATCGAAATTT	
<i>P. cryptogea</i>	AF266796	ATTGCAGGAT	TCAGTGAGTC	ATCGAAATTT	
<i>P. infestans</i>	AF266779	ATTGCAGGAT	TCAGTGAGTC	ATCGAAATTT	
<i>P. nicotianae</i>	AF266776	ATTGCAGGAT	TCAGTGAGTC	ATCGAAATTT	
<i>P. palmivora</i>	AF266780	ATTGCAGGAT	TCAGTGAGTC	ATCGAAATTT	
<i>Pyhtium aphanidermatum</i>	AJ233438	ATTGCAGAAT	TCAGTGAGTC	ATCGAAATTT	
CMW 13800		TGAACGCATA	TTGCACTTCC	GGGTTAGTCC	330
CMW 13816		TGAACGCATA	TTGCACTTCC	GGGTTAGTCC	
CMW 13804		GGAACGCATA	TTGCACTTCC	GGGTTAGTCC	
<i>P. cinnamomi</i>	AF266764	GGAACGCATA	TTGCACTTCC	GGGTTAGTCC	
<i>P. cryptogea</i>	AF266796	TGAACGCATA	TTGCACTTCC	GGGTTAGTCC	
<i>P. infestans</i>	AF266779	TGAACGCATA	TTGCACTTCC	GGGTTAGTCC	
<i>P. nicotianae</i>	AF266776	TGAACGCATA	TTGCACTTCC	GGGTTAGTCC	
<i>P. palmivora</i>	AF266780	TGAACGCATA	TTGCACTTCC	GGGTTAGTCC	
<i>Pyhtium aphanidermatum</i>	AJ233438	TGAACGCACA	TTGCACTTCC	GGGTTATGCC	
CMW 13800		TGGAAGTATG	CCTGTATCAG	TGTCCGTACA	360
CMW 13816		TGGAAGTATG	CCTGTATCAG	TGTCCGTACA	
CMW 13804		TGGGAGTATG	CCTGTATCAG	TGTCCGTACA	
<i>P. cinnamomi</i>	AF266764	TGGGAGTATG	CCTGTATCAG	TGTCCGTACA	
<i>P. cryptogea</i>	AF266796	TGGGAGTATG	CCTGTATCAG	TGTCCGTACA	
<i>P. infestans</i>	AF266779	TGGAAGTATG	CCTGTATCAG	TGTCCGTACA	
<i>P. nicotianae</i>	AF266776	TGGAAGTATG	CCTGTATCAG	TGTCCGTACA	
<i>P. palmivora</i>	AF266780	TGGGAGTATG	CCTGTATCAG	TGTCCGTACA	
<i>Pyhtium aphanidermatum</i>	AJ233438	TGGAAGTATG	CCTGTATCAG	TGTCCGTACA	



				390
CMW 13800		TAAACTTGA	CTTCTTCT	TCCGTGTAGT
CMW 13816		TAAACTTGA	CTTCTTCT	TCCGTGTAGT
CMW 13804		TAAACTTGG	CTCTCTTCT	TCCGTGTAGT
<i>P. cinnamomi</i>	AF266764	TAAACTTGG	CTCTCTTCT	TCCGTGTAGT
<i>P. cryptogea</i>	AF266796	TAAACTTGG	CTCCCTTCT	TCCGTGTAGT
<i>P. infestans</i>	AF266779	CAAACTTGG	CTTCTCCCT	TCCGTGTAGT
<i>P. nicotianae</i>	AF266776	TAAACTTGA	CTTCTTCT	TCCGTGTAGT
<i>P. palmivora</i>	AF266780	TAAACTTGG	TTTCTTCT	TCCGTGTAGT
<i>Pyhtium aphanidermatum</i>	AJ233438	TAAACTTGC	CTTCTTT-T	TCTGTGTAGT
				420
CMW 13800		CGGTGGA-GG	AGATGT-CAG	ATGTGAAGTG
CMW 13816		CGGTGGA-GG	AGATGT-CAG	ATGTGAAGTG
CMW 13804		CGGTGGATGG	AGGTGC-CAG	ACGTGAGGTG
<i>P. cinnamomi</i>	AF266764	CGGTGGATGG	AGGTGC-CAG	ACGTGAGGTG
<i>P. cryptogea</i>	AF266796	CGGTGGATGG	GGACGCGCAG	ATGTGAAGTG
<i>P. infestans</i>	AF266779	CGGTGG-AGG	AGATGC-CAG	ATGTGAAGTG
<i>P. nicotianae</i>	AF266776	CGGTGGA-GG	AGATGT-CAG	ATGTGAAGTG
<i>P. palmivora</i>	AF266780	CGGTGG-TGG	ATGTGC-CAG	ATGTGAAGTG
<i>Pyhtium aphanidermatum</i>	AJ233438	CAG-GGAGAG	AGATGGCAGA	ATGTGAGGTG
				450
CMW 13800		TCTTGC-GAT	TGGTCTTC-	-GGACCGGC
CMW 13816		TCTTGC-GAT	TGGTCTTC-	-GGACCGGC
CMW 13804		TCTTGC-GGG	CGGTCTTC-	-GGACTGGC
<i>P. cinnamomi</i>	AF266764	TCTTGC-GGG	CGGTCTTC-	-GGACTGGC
<i>P. cryptogea</i>	AF266796	TCTTGC-GGC	TGGTCTTC-	-GGTCCGGC
<i>P. infestans</i>	AF266779	TCTTGC-GGT	TGGTTTCC-	-GGACCGAC
<i>P. nicotianae</i>	AF266776	TCTTGC-GAT	TGGTCTTC-	-GGACCGGC
<i>P. palmivora</i>	AF266780	TCTTGC-GGC	TGGTCTTC-	-GGATCGGC
<i>Pyhtium aphanidermatum</i>	AJ233438	TCTCGCTGGC	TCCCTTTTCG	GAGGAGAAGA
				480
CMW 13800		TGCGAGTCCT	TTTAAATGTA	C-TAAACTGA
CMW 13816		TGCGAGTCCT	TTTAAATGTA	C-TAAACTGA
CMW 13804		TGTGAGTCCC	TTGAAATGTA	C-TGAACTGT
<i>P. cinnamomi</i>	AF266764	TGTGAGTCCC	TTGAAATGTA	C-TGAACTGT
<i>P. cryptogea</i>	AF266796	TGCGAGTCCT	TTGAAATGTA	C-TACACTGT
<i>P. infestans</i>	AF266779	TGCGAGTCCT	TTGAAATGTA	C-TAAACTGT
<i>P. nicotianae</i>	AF266776	TGCGAGTCCT	TTTAAATGTA	C-TAAACTGA
<i>P. palmivora</i>	AF266780	TGTGAGTCCC	TTGAAATGTA	C-TGAACTGT
<i>Pyhtium aphanidermatum</i>	AJ233438	CGCGAGTCCC	TTTAAATGTA	CGTTCGCTCT
				510
CMW 13800		ACTTCTCTTT	GCTCGAAAAG	TGGTGGCGTT
CMW 13816		ACTTCTCTTT	GCTCGAAAAG	TGGTGGCGTT
CMW 13804		ACTTCTCTTT	GCTCGAAAAG	CG-TGACGTT
<i>P. cinnamomi</i>	AF266764	ACTTCTCTTT	GCTCGAAAAG	CG-TGACGTT
<i>P. cryptogea</i>	AF266796	ACTTCTCTTT	GCTCGAAAAG	CG-TGACGTT
<i>P. infestans</i>	AF266779	ACTTCTCTTT	GCTCCAAAAG	TGGTGGCATT
<i>P. nicotianae</i>	AF266776	ACTTCTCTTT	GCTCGAAAAG	TGGTGGCGTT
<i>P. palmivora</i>	AF266780	ACTTCTCTTT	GCTCCAAAAG	CG-TGGCGTT
<i>Pyhtium aphanidermatum</i>	AJ233438	TTCTTGTGTC	TAAGATGAAG	TG-TGAT-TC
				540
CMW 13800		GCTGGTTGTG	AAGG-CTGCT	ATTGTGGCAA
CMW 13816		GCTGGTTGTG	AAGG-CTGCT	ATTGTGGCAA
CMW 13804		GCTGGTTGTG	GAGG-CTGCC	TGTATGGCCA
<i>P. cinnamomi</i>	AF266764	GCTGGTTGTG	GAGG-CTGCC	TGTATGGCCA
<i>P. cryptogea</i>	AF266796	GCTGGTTGTG	GAGG-CTGCC	TGTGTGGCAT
<i>P. infestans</i>	AF266779	GCTGGTTGTG	GACG-CTGCT	ATTGTAGCCA
<i>P. nicotianae</i>	AF266776	GCTGGTTGTG	AAGG-CTGCT	ATTGTGGCAA
<i>P. palmivora</i>	AF266780	GCTGATTGTG	GAGG-CTGCT	TGCGTAGCCA
<i>Pyhtium aphanidermatum</i>	AJ233438	TCGAATCGCG	GTGATCTGTT	TGGATCGCTT



				570
CMW 13800		ATT-GGCGAC	TGGTTTGTCT	GCTGCGGCA-
CMW 13816		ATT-GGCGAC	TGGTTTGTCT	GCTGCGGCA-
CMW 13804		GTC-GGCGAC	CGGTTTGTCT	GCTGCGGCGT
<i>P. cinnamomi</i>	AF266764	GTC-GGCGAC	CGGTTTGTCT	GCTGCGGCGT
<i>P. cryptogea</i>	AF266796	GTC-GGCGAC	CGGTTTGTCT	GCTGCGGCGT
<i>P. infestans</i>	AF266779	GTT-GGCGAC	CGGTTTGTCT	GCTGCGGCGT
<i>P. nicotianae</i>	AF266776	ATT-GGCGAC	TGGTTTGTCT	GCTGCGGCA-
<i>P. palmivora</i>	AF266780	GTCTGGCGAC	CAGTTTGTCT	GCTGTGGCGT
<i>Pyhtium aphanidermatum</i>	AJ233438	TGC-GCATTT	GGGCGACTTC	GGTTAGGACA
				600
CMW 13800		TTAATGGAAG	AGTGTTTCGAT	TCGTGGTATG
CMW 13816		TTAATGGAAG	AGTGTTTCGAT	TCGTGGTATG
CMW 13804		TTAATGGAGG	AGTGTTTCGAT	TCGCGGTATG
<i>P. cinnamomi</i>	AF266764	TTAATGGAGG	AGTGTTTCGAT	TCGCGGTATG
<i>P. cryptogea</i>	AF266796	TTAATGGAGG	AGTGTTTCGAT	TCGCGGTATG
<i>P. infestans</i>	AF266779	T-AATGGAGA	AATGCTTCGAT	TCGTGGTATG
<i>P. nicotianae</i>	AF266776	TTAATGGAAG	AGTGTTTCGAT	TCGTGGTATG
<i>P. palmivora</i>	AF266780	T-AATGGAGG	AGTGTTTCGAT	TCGCGGTATG
<i>Pyhtium aphanidermatum</i>	AJ233438	TTAAAGGAAG	CAACCTCTAT	TGGCGGTATG
				630
CMW 13800		GTTGGCTTCG	GCTGAACAA-	TGCACTTATT
CMW 13816		GTTGGCTTCG	GCTGAACAA-	TGCACTTATT
CMW 13804		GTTGGCTCCG	GCTGAACAA-	AGCGCTTATT
<i>P. cinnamomi</i>	AF266764	GTTGGCTCCG	GCTGAACAA-	AGCGCTTATT
<i>P. cryptogea</i>	AF266796	GTTGGCTTCG	GCTGAACA-	GACGCTTATT
<i>P. infestans</i>	AF266779	GTTGCCTTCG	GCTGAACAA-	TGCGCTTATT
<i>P. nicotianae</i>	AF266776	GTTGGCTTCG	GCTGAACAA-	TGCACTTATT
<i>P. palmivora</i>	AF266780	GTTGGCTTCG	GCTGAACAG-	A-CGCTTATT
<i>Pyhtium aphanidermatum</i>	AJ233438	TTAGGCTTCG	GCCCGACGTT	GCAGCTGACA
				660
CMW 13800		GGACGTTTTT	CC-TGCTGTG	GCGTGATGGA
CMW 13816		GGACGTTTTT	CC-TGCTGTG	GCGTGATGGA
CMW 13804		GGATGTTTCT	CCCTGCTGTG	GCGGTACGGA
<i>P. cinnamomi</i>	AF266764	GGATGTTTCT	CCCTGCTGTG	GCGGTACGGA
<i>P. cryptogea</i>	AF266796	GGGTGCTTTT	CC-TGCTGTG	GCTGGATGGA
<i>P. infestans</i>	AF266779	GGGTGATTTT	CCT-GCTGTG	GCGTGATGGA
<i>P. nicotianae</i>	AF266776	GGACGTTTTT	CC-TGCTGTG	GCGTGATGGA
<i>P. palmivora</i>	AF266780	GAATATTTCT	TCA-GCTGTG	GTGGTATG-A
<i>Pyhtium aphanidermatum</i>	AJ233438	GAGTGTGGTT	—TTCTGTT	CTTTCTTGA
				690
CMW 13800		CTGGTGAACC	ATAGCTCGGT	—GGCTTGGC
CMW 13816		CTGGTGAACC	ATAGCTCGGT	—GGCTTGGC
CMW 13804		TCGGTGAACC	GTAGCTGTGC	TAGGCTTGGC
<i>P. cinnamomi</i>	AF266764	TCGGTGAACC	GTAGCTGTGC	TAGGCTTGGC
<i>P. cryptogea</i>	AF266796	CTGGTGAACC	GTAGCTGTGC	TAGGCTTGGC
<i>P. infestans</i>	AF266779	CTGGTGAACC	ATGGCTCT—	TTAGCTTGGC
<i>P. nicotianae</i>	AF266776	CTGGTGAACC	ATAGCTCGGT	—GGCTTGGC
<i>P. palmivora</i>	AF266780	TTGGTGAACC	GTAGCTATG-	TGAGCTTGGC
<i>Pyhtium aphanidermatum</i>	AJ233438	—GGTGTACC	TGAATTGTGT	GAGGCAATG-
				720
CMW 13800		TTTTGAATTG	GCTTTGCTGT	TGCCAAGTAG
CMW 13816		TTTTGAATTG	GCTTTGCTGT	TGCCAAGTAG
CMW 13804		GTTTGAACCG	GCGGTGTTGT	TGCCAAGTAG
<i>P. cinnamomi</i>	AF266764	GTTTGAACCG	GCGGTGTTGT	TGCCAAGTAG
<i>P. cryptogea</i>	AF266796	GTTTGAACCG	GCGGTG-TGG	TGCCAAGTAG
<i>P. infestans</i>	AF266779	ATTTGAATCG	GCTTTGCTGT	TGCCAAGTAG
<i>P. nicotianae</i>	AF266776	TTTTGAATTG	GCTTTGCTGT	TGCCAAGTAG
<i>P. palmivora</i>	AF266780	TTTTGAATTG	GCTTTGCTGT	TGCCAAGTAG
<i>Pyhtium aphanidermatum</i>	AJ233438	GTCTGGGCAA	ATGGT—TGC	TGTGTAGTAG



				750
CMW 13800		GGTGG—	—	— — — — C —
CMW 13816		GGTGG—	—	— — — — C —
CMW 13804		GGTGG—CGG	CTTC—GGCT	GTCGAGGGTC
<i>P. cinnamomi</i>	AF266764	GGTGG—	—	— — — — C
<i>P. cryptogea</i>	AF266796	GGTGT—CTG	TTCT—GGCG	— — — — TA
<i>P. infestans</i>	AF266779	AGTGG—	—	— — — — C
<i>P. nicotianae</i>	AF266776	GGTGG—	—	— — — — C —
<i>P. palmivora</i>	AF266780	AGTGG—	—	— — — — C
<i>Pyhtium aphanidermatum</i>	AJ233438	GGTTTTGCTG	CTCTTGGACG	CCCTGTTTTC
				780
CMW 13800		AGCTTCGGTT	GTCGAGGGTC	GATCCATTTG
CMW 13816		AGCTTCGGTT	GTCGAGGGTC	GATCCATTTG
CMW 13804		— GATCCATTT	GGGAACCTCTG	TGTCTCTCCG
<i>P. cinnamomi</i>	AF266764	GGCTTCGGCT	GTCGAGGGTC	GATCCATTTG
<i>P. cryptogea</i>	AF266796	AGCTGGGGTG	GACGAGGGTC	GATCCATTTG
<i>P. infestans</i>	AF266779	GGCTTCGGCT	GCCGAGGGTC	GATCCATTTG
<i>P. nicotianae</i>	AF266776	AGCTTCGGTT	GTCGAGGGTC	GATCCATTTG
<i>P. palmivora</i>	AF266780	GGCTTCGGCT	GTCGAGGGTC	GATCCATTTG
<i>Pyhtium aphanidermatum</i>	AJ233438	GGATAGGGTA	AAGGAGGCAA	CACCAATTTG
				810
CMW 13800		GGA—ACTTAA	TGTGTACTTC	GGTATGCATC
CMW 13816		GGA—ACTTAA	TGTGTACTTC	GGTATGCATC
CMW 13804		GCGCACTT—	—GTGTGCTTG	TGGTGGCATC
<i>P. cinnamomi</i>	AF266764	GGA—ACTCTG	TGTCT—CTCC	GGCGCACTTG
<i>P. cryptogea</i>	AF266796	GGAAACGTTG	TGTGCGCTTC	GGCGCGCATC
<i>P. infestans</i>	AF266779	GGA—AATGT—	TGTGTACTTC	GGTATGCATC
<i>P. nicotianae</i>	AF266776	GGA—ACTTAA	TGTGTACTTC	GGTATGCATC
<i>P. palmivora</i>	AF266780	GGA—ACT—TG	TGTATGCTTC	GGCATGCATC
<i>Pyhtium aphanidermatum</i>	AJ233438	GGACTGTTTG	CAATTTATTG	TGAACAACCT
				814
CMW 13800		TCAA		
CMW 13816		TCAA		
CMW 13804		TCAA		
<i>P. cinnamomi</i>	AF266764	TGTG		
<i>P. cryptogea</i>	AF266796	TCAA		
<i>P. infestans</i>	AF266779	TCAA		
<i>P. nicotianae</i>	AF266776	TCAA		
<i>P. palmivora</i>	AF266780	TCAA		
<i>Pyhtium aphanidermatum</i>	AJ233438	TCTA		



Alignment 4. The combined ITS1, 5.8s subunit and ITS2 regions of the genomic RNA gene DNA sequence of the *Pythium* isolates obtained from the *Eucalyptus* hedge plants.

				30
CMW 13819		CCACACC- AA	AAAAA - CTTT	CCACGTGAA-
CMW 13822		CCACACC- TA	AAAAA - CTTT	CCACGTGAA-
CMW 13805		CCACACC- AT	AAAA - CTTT	CCACGTGAA-
CMW 13796		CCACACC- TA	AAAAA - CTTT	CCACGTGAA-
CMW 13820		CCACACC- AA	AAAAA - CTTT	CCACGTGAA-
CMW 13807		CCACACCAAA	AAAA - CTTT	CCACGTGAA-
CMW 13808		CCACACCTAA	AAACATCTTT	CCACGTGAA-
CMW 13815		CCACACC - AT	AAAAA - CTTT	CCACGTGAA-
CMW 13817		CCACACCTAA	AAACATCTTT	CCACGTGAA-
CMW 13810		CCACACC- A	TAAAAACTTT	CCACGTGAA-
<i>P. aphanidermatum</i>	AJ233438	CCACACC- AT	AAAAA - CTTT	CCACGTGAA-
<i>P. deliense</i>	AJ233442	CCACACC- AT	AAAA - CTTT	CCACGTGAA-
<i>P. dissotocum</i>	AJ233443	CCACACC- AA	AAAAA - CTTT	CCACGTGAA-
<i>P. helicoides</i>	AB108061	CCACACCTAA	AAACATCTTT	CCACGTGAA-
<i>P. intermedium</i>	AJ233447	CCACACC- TA	AAAAA - CTTT	CCACGTGAA-
<i>P. irregulare</i>	AJ233448	CCACACCTAA	AAAAA - CTTT	CCACGTGAA-
<i>P. ultimum</i>	AB108064	CCACACT- TT	AAAAAACTGT	CCACGTGAA-
<i>P. vexans</i>	AJ233462	CCACACC- AA	AAAAA - CTTT	CCACGTGAA-
<i>Phytophthora cinnamomi</i>	AF266764	CTGCTCTGGG	CGGCGGTTGT	CGATGTCAAA
				60
CMW 13819		—CCGTT	-TTGTG-CGT	TTTGTGCTTG
CMW 13822		—CTGTC	ATTATT-TGT	TGTGCGCTCT
CMW 13805		—CCGTT	-GAAAT-CAT	GTTCTG—
CMW 13796		—CTGTC	ATTATT-TGT	TGTGCGCTCT
CMW 13820		—CCGTT	-GTAAC-TAT	GTTCTG—
CMW 13807		—CCGTT	GTAACATATGT	TCTGTGCTCT
CMW 13808		—CCGTT	TGTGACATGG	T-TGGGCTTG
CMW 13815		—CCGTT	G-AAATCAT	GTTCTGTG—
CMW 13817		—CCGTT	T-GTGACAT	GGTTGGGC—
CMW 13810		—CCGTT	G-AAATCAT	GTTCTGTG—
<i>P. aphanidermatum</i>	AJ233438	—CCGTT	G-AAATCAT	GTTCTGTG—
<i>P. deliense</i>	AJ233442	—CCGTT	-GAAAT-CAT	GTTCTG—
<i>P. dissotocum</i>	AJ233443	—CCGTT	GTAACATATGT	TCTGTGCTCT
<i>P. helicoides</i>	AB108061	—CCGTT	-TGTGACAT	GGTTGGGC—
<i>P. intermedium</i>	AJ233447	—CTGTC	ATTATT-TGT	TGTGCGCTCT
<i>P. irregulare</i>	AJ233448	—CTGTC	GTTATT-TGT	TGTGTGTGTG
<i>P. ultimum</i>	AB108064	—CTGTA	AGCAAGTCTA	GCGCTGTGAC
<i>P. vexans</i>	AJ233462	—CCGTT	-TTGTG-CGT	TTTGTGCTTG
<i>Phytophthora cinnamomi</i>	AF266764	GTCGACGGTT	GCTGTTGCGT	GGGGGGCC—
				90
CMW 13819		TTGTTTTTCG	ATCTGCTCTC	TGTGCCT—
CMW 13822		CTGCGGTGTC	GGTGGCGTCT	GTTGGCTGTA
CMW 13805		—	—TGCTCT—	—CT—
CMW 13796		CTGCGGTGTC	GGTGGCGTCT	GTTGGCTGTA
CMW 13820		—	—TGCTCT—	—CTT—
CMW 13807		CTTC—	—	—
CMW 13808		TGCG—	—	—
CMW 13815		—	CTCTCTTT—	—
CMW 13817		—	TTGTGCGT—	—
CMW 13810		—	CTCTCTTT—	—
<i>P. aphanidermatum</i>	AJ233438	—	CTCTCTTT—	—
<i>P. deliense</i>	AJ233442	—	—TGCTCT—	—CT—
<i>P. dissotocum</i>	AJ233443	CTTC—	—	—
<i>P. helicoides</i>	AB108061	—	TTGT—	—
<i>P. intermedium</i>	AJ233447	CTGCGGTGTC	GGTGGCGTCT	GTTGGCTGTA
<i>P. irregulare</i>	AJ233448	CGTGTGGTA	GCATGCGTGT	TTGCTTACGC
<i>P. ultimum</i>	AB108064	TGAGCTGGTG	TTTTCATTTT	TGGACACTGG
<i>P. vexans</i>	AJ233462	TTGTTTTTCG	ATCTGCTCTC	TGTGCCT—
<i>Phytophthora cinnamomi</i>	AF266764	—	CTATCACT—	—



CMW 13819		TTCGGGTGT-	G—GAGTGT	GGGGAAGAAA	120
CMW 13822		TTTGATACT-	GCTGGCGGGT	GCGAGCCGGA	
CMW 13805		CTCGGGAG—	_____	_____	
CMW 13796		TTTGATACT-	GCTGGCGGGT	GCGAGCCGGA	
CMW 13820		CTCGGAGAG—	_____	_____	
CMW 13807		-TCGGAGAG—	_____	_____	
CMW 13808		-TTTTCTCT-	_____	_____	
CMW 13815		-CGGGAG—	_____	_____	
CMW 13817		-TTTCTC—	_____	_____	
CMW 13810		-CGGGAG—	_____	_____	
<i>P. aphanidermatum</i>	AJ233438	-CGGGAG—	_____	_____	
<i>P. deliense</i>	AJ233442	CTCGGGAG—	_____	_____	
<i>P. dissotocum</i>	AJ233443	_____	_____	—TCGGA	
<i>P. helicoides</i>	AB108061	-GCGTTTTTC-	_____	_____	
<i>P. intermedium</i>	AJ233447	TTTGATACT-	GCTGGCGGGT	GCGAGCCGGA	
<i>P. irregulare</i>	AJ233448	TTTGGGGGTT	GCGAGTGTGT	GTGTTGTCGG	
<i>P. ultimum</i>	AB108064	AACGGGAGT-	_____	_____	
<i>P. vexans</i>	AJ233462	TTCGGGTGT-	G—GAGTGT	GGGGAAGAAA	
<i>Phytophthora cinnamomi</i>	AF266764	GGCGAGCGT-	_____	_____	

CMW 13819		GCTCCAGGCT	AAACGAAGGC	-TGCGAGTTT	150
CMW 13822		TGCAGAGGCT	GAACGAAGGT	-CG—AGTTG	
CMW 13805		—GGCT	GAACGAAGGT	-GGGCTGCTT	
CMW 13796		TGCAGAGGCT	GAACGAAGGT	-CG—AGTTG	
CMW 13820		—AGCT	GAACGAAGGT	-GGGCTGCTT	
CMW 13807		—AGCT	GAACGAAGGT	-GGGCTGCTT	
CMW 13808		—CTCT	TT-TGTAGGG	-GGGATGCGT	
CMW 13815		—GGCT	GAACGAAGGT	GGG-CTGCTT	
CMW 13817		—TCTC	TTTTGTAGGG	GGG-ATGCGT	
CMW 13810		—GGCT	GAACGAAGGT	GGG-CTGCTT	
<i>P. aphanidermatum</i>	AJ233438	—GGCT	GAACGAAGGT	GGG-CTGCTT	
<i>P. deliense</i>	AJ233442	—GGCT	GAACGAAGGT	-GGGCTGCTT	
<i>P. dissotocum</i>	AJ233443	—GAGAGCT	GAACGAAGGT	-GGGCTGCTT	
<i>P. helicoides</i>	AB108061	—TCTCT	CTTTTGTAGG	GGGATGCGT	
<i>P. intermedium</i>	AJ233447	TGCAGAGGCT	GAACGAAGGT	CGAGTTGCTT	
<i>P. irregulare</i>	AJ233448	TGCGCAGACT	GAACGAAGGT	-CGTGTGTTG	
<i>P. ultimum</i>	AB108064	—CAGCA	GGACGAAGGT	TGGTCTGTTG	
<i>P. vexans</i>	AJ233462	GCTCCAGGCT	AAACGAAGGC	-TGCGAGTTT	
<i>Phytophthora cinnamomi</i>	AF266764	—TTGGG	TCCCCCTCGG	GGGAAC—	

CMW 13819		CGTGCTTG—	_____	_____	180
CMW 13822		CTTTGCTCT-	_____	_____	
CMW 13805		AATTGTGGT-	_____	_____	
CMW 13796		CTTTGCTCT-	_____	_____	
CMW 13820		AATTGTAGT-	_____	_____	
CMW 13807		AATTGTAGT-	_____	_____	
CMW 13808		GCGAGCTAT-	_____	_____	
CMW 13815		AATTGTAGTC	T—	_____	
CMW 13817		—GCGAGC	T—	_____	
CMW 13810		AATTGTAGTC	T—	_____	
<i>P. aphanidermatum</i>	AJ233438	AATTGTAGTC	T—	_____	
<i>P. deliense</i>	AJ233442	AATTGTGGT-	_____	_____	
<i>P. dissotocum</i>	AJ233443	AATTGTAGT-	_____	_____	
<i>P. helicoides</i>	AB108061	—GCGAGC	T—	_____	
<i>P. intermedium</i>	AJ233447	—TGCTCT-	_____	_____	
<i>P. irregulare</i>	AJ233448	CTGTGTGCCT	GCTGCACTGC	TGACTTTGCA	
<i>P. ultimum</i>	AB108064	TAATGCAAGT	TATGATGGAC	_____	
<i>P. vexans</i>	AJ233462	CGTGCTTG—	_____	_____	
<i>Phytophthora cinnamomi</i>	AF266764	—TGAGC	T—	_____	



210

CMW 13819		_____	_____	_____
CMW 13822		_____	_____	_____
CMW 13805		_____	_____	_____
CMW 13796		_____	_____	_____
CMW 13820		_____	_____	_____
CMW 13807		_____	_____	_____
CMW 13808		_____	_____	_____
CMW 13815		_____	_____	_____
CMW 13817		_____	_____	_____
CMW 13810		_____	_____	_____
<i>P. aphanidermatum</i>	AJ233438	_____	_____	_____
<i>P. deliense</i>	AJ233442	_____	_____	_____
<i>P. dissotocum</i>	AJ233443	_____	_____	_____
<i>P. helicoides</i>	AB108061	_____	_____	_____
<i>P. intermedium</i>	AJ233447	_____	_____	_____
<i>P. irregulare</i>	AJ233448	TTGATTTGCA	TGGTGTTGGC	GGAGCGGCGG
<i>P. ultimum</i>	AB108064	_____	_____	_____
<i>P. vexans</i>	AJ233462	_____	_____	_____
<i>Phytophthora cinnamomi</i>	AF266764	_____	_____	_____

240

CMW 13819		_____	_____CGGCC	GATTTATTCT
CMW 13822		_____	_____CGGCT	GACTTATT-T
CMW 13805		_____	_____CTGCC	GATGTATT-T
CMW 13796		_____	_____CGGCT	GACTTATT-T
CMW 13820		_____	_____CTGCC	GATGFACT-
CMW 13807		_____	_____CTGCC	GATGFACT-
CMW 13808		_____	_____CTGTA	AAC-TT-
CMW 13815		_____	_____GCC	GATGTATTTT
CMW 13817		_____	_____ATC	TGTAACTTG
CMW 13810		_____	_____GCC	GATGTATTTT
<i>P. aphanidermatum</i>	AJ233438	_____	_____GCC	GATGTATTTT
<i>P. deliense</i>	AJ233442	_____	_____CTGCC	GATGTATT-T
<i>P. dissotocum</i>	AJ233443	_____	_____CTGCC	GATGFACT-T
<i>P. helicoides</i>	AB108061	_____	_____ATC	TGTAACTTG
<i>P. intermedium</i>	AJ233447	_____	_____CGGCT	GACTTATT-T
<i>P. irregulare</i>	AJ233448	GTGCTGTTGC	ATGCGCGGCT	GACCTATTTT
<i>P. ultimum</i>	AB108064	_____	_____TAGCT	GATGAATTTT
<i>P. vexans</i>	AJ233462	_____	_____CGGCC	GATTTATTCT
<i>Phytophthora cinnamomi</i>	AF266764	_____	_____AGT	AGCCTCTCTT

270

CMW 13819		TT-CAAAC	CCATACATTA	AA-CACTGA
CMW 13822		TT-CAAAC	CCA-ATACCC	AACTTACTGA
CMW 13805		TT-CAAAC	CCATTTACCT	AA-TACTGA
CMW 13796		TT-CAAAC	CCA-ATACCC	AACTTACTGA
CMW 13820		TT-TAAAC	CCATTAAACT	AA-TACTGA
CMW 13807		TT-TAAAC	CCATTAAACT	-AATACTGA
CMW 13808		GT-CAAAC	CCATTCTTTT	TGATAACTGA
CMW 13815		T-CAAAC	CCATT-TACC	TAATA-CTGA
CMW 13817		T-CAAAC	CCATTCTTTT	TGATAACTGA
CMW 13810		T-CAAAC	CCATT-TACC	TAATA-CTGA
<i>P. aphanidermatum</i>	AJ233438	T-CAAAC	CCATT-TACC	TAATA-CTGA
<i>P. deliense</i>	AJ233442	TT-CAAAC	CCATTTACCT	AA-TACTGA
<i>P. dissotocum</i>	AJ233443	TT-AAAC	CCATTAAACT	AA-TACTGA
<i>P. helicoides</i>	AB108061	T-CAAAC	CCATTCTTTT	TGATAACTGA
<i>P. intermedium</i>	AJ233447	TT-CAAAC	CCAATACCCA	AC-TTACTG
<i>P. irregulare</i>	AJ233448	TT-TCAAA	CCCCATACCT	AAATGACTGA
<i>P. ultimum</i>	AB108064	TGTTTTTAAA	CCCTT-ACCT	AAATA-CTGA
<i>P. vexans</i>	AJ233462	TT-CAAAC	CCATACATTA	AA-CACTGA
<i>Phytophthora cinnamomi</i>	AF266764	T-TAAAC	CCATCCTGT-	-AATACTGA



				300
CMW 13819		AGTATACTGT	GAGGACGAAA	GTCCCTTGCTT
CMW 13822		T-TATACTGT	GAGAACGAAA	GTTCTTGCTT
CMW 13805		TCTATACTCC	AAAAACGAAA	GTTTCTGGTT
CMW 13796		T-TATACTGT	GAGAACGAAA	GTTCTTGCTT
CMW 13820		ACTATACTCC	GAAAACGAAA	GTCTTTGGTT
CMW 13807		ACTATACTCC	GAAAACGAAA	GTCTTTGGTT
CMW 13808		AACATACTGT	GGGGACGAAA	GTCTCTGCTT
CMW 13815		TCTATACTCC	AAAAACGAAA	GTTTATGGTT
CMW 13817		AACATACTGT	GGGGACGAAA	GTCTCTGCTT
CMW 13810		TCTATACTCC	AAAAACGAAA	GTTTATGGTT
<i>P. aphanidermatum</i>	AJ233438	TCTATACTCC	AAAAACGAAA	GTTTCTGGTT
<i>P. deliense</i>	AJ233442	TCTATACTCC	AAAAACGAAA	GTTTCTGGTT
<i>P. dissotocum</i>	AJ233443	ACTATACTCC	GAAAACGAAA	GTCTTTGGTT
<i>P. helicoides</i>	AB108061	AACATACTGT	GGGGACGAAA	GTCTCTGCTT
<i>P. intermedium</i>	AJ233447	ATTATACTGT	GAGAACGAAA	GTTCTTGCTT
<i>P. irregulare</i>	AJ233448	T-TATACTGT	GAGAACGAAA	GTTCTTGCTT
<i>P. ultimum</i>	AB108064	TTTATACTGT	GGGGACGAAA	GTCCCTTGCTT
<i>P. vexans</i>	AJ233462	AGTATACTGT	GAGGACGAAA	GTCCCTTGCTT
<i>Phytophthora cinnamomi</i>	AF266764	A-CATACTGT	GGGGACGAAA	GTCTCTGCTT

				330
CMW 13819		TGAACTAGAT	AGCAACTTTC	AGCAGTGGAT
CMW 13822		TTAACTAGAT	AACAAC TTTT	AGCAGTGGAT
CMW 13805		TTAATCC-AT	AACAAC TTTT	AGCAGTGGAT
CMW 13796		TTAACTAGAT	AACAAC TTTT	AGCAGTGGAT
CMW 13820		TTAATCA-AT	AACAAC TTTT	AGCAGTGGAT
CMW 13807		TTAATCA-AT	AACAAC TTTT	AGCAGTGGAT
CMW 13808		TGAACTAGAT	AGCAACTTTC	AGCAGTGGAT
CMW 13815		TTAA-TCTAT	AACAAC TTTT	AGCAGTGGAT
CMW 13817		TGAACTAGAT	AGCAACTTTC	AGCAGTGGAT
CMW 13810		TTAA-TCTAT	AACAAC TTTT	AGCAGTGGAT
<i>P. aphanidermatum</i>	AJ233438	TTAA-TCTAT	AACAAC TTTT	AGCAGTGGAT
<i>P. deliense</i>	AJ233442	TTAATCC-AT	AACAAC TTTT	AGCAGTGGAT
<i>P. dissotocum</i>	AJ233443	TTAATCA-AT	AACAAC TTTT	AGCAGTGGAT
<i>P. helicoides</i>	AB108061	TGAACTAGAT	AGCAACTTTC	AGCAGTGGAT
<i>P. intermedium</i>	AJ233447	TTAACTAGAT	AACAAC TTTT	AGCAGTGGAT
<i>P. irregulare</i>	AJ233448	TTAACTAGAT	AACAAC TTTT	AGCAGTGGAT
<i>P. ultimum</i>	AB108064	TTAC-TAGAT	AACAAC TTTT	AGCAGTGGAT
<i>P. vexans</i>	AJ233462	TGAACTAGAT	AGCAACTTTC	AGCAGTGGAT
<i>Phytophthora cinnamomi</i>	AF266764	TTAACTAGAT	AGCAACTTTC	AGCAGTGGAT

				360
CMW 13819		GTCTAGGCTC	GCACATCGAT	GAAGAACGCT
CMW 13822		GTCTAGGCTC	GCACATCGAT	GAAGAACGCT
CMW 13805		GTCTAGGCTC	GCACATCGAT	GAAGAACGCT
CMW 13796		GTCTAGGCTC	GCACATCGAT	GAAGAACGCT
CMW 13820		GTCTAGGCTC	GCACATCGAT	GAAGAACGCT
CMW 13807		GTCTAGGCTC	GCACATCGAT	GAAGAACGCT
CMW 13808		GTCTAGGCTC	GCACATCGAT	GAAGAACGCT
CMW 13815		GTCTAGGCTC	GCACATCGAT	GAAGAACGCT
CMW 13817		GTCTAGGCTC	GCACATCGAT	GAAGAACGCT
CMW 13810		GTCTAGGCTC	GCACATCGAT	GAAGAACGCT
<i>P. aphanidermatum</i>	AJ233438	GTCTAGGCTC	GCACATCGAT	GAAGAACGCT
<i>P. deliense</i>	AJ233442	GTCTAGGCTC	GCACATCGAT	GAAGAACGCT
<i>P. dissotocum</i>	AJ233443	GTCTAGGCTC	GCACATCGAT	GAAGAACGCT
<i>P. helicoides</i>	AB108061	GTCTAGGCTC	GCACATCGAT	GAAGAACGCT
<i>P. intermedium</i>	AJ233447	GTCTAGGCTC	GCACATCGAT	GAAGAACGCT
<i>P. irregulare</i>	AJ233448	GTCTAGGCTC	GCACATCGAT	GAAGAACGCT
<i>P. ultimum</i>	AB108064	GTCTAGGCTC	GCACATCGAT	GAAGAACGCT
<i>P. vexans</i>	AJ233462	GTCTAGGCTC	GCACATCGAT	GAAGAACGCT
<i>Phytophthora cinnamomi</i>	AF266764	GTCTAGGCTC	GCACATCGAT	GAAGAACGCT



				390
CMW 13819		GCGAACTGCG	ATACGTAATG	CGAATTGCAG
CMW 13822		GCGAACTGCG	ATACGTAATG	CGAATTGCAG
CMW 13805		GCGAACTGCG	ATACGTAATG	CGAATTGCAG
CMW 13796		GCGAACTGCG	ATACGTAATG	CGAATTGCAG
CMW 13820		GCGAACTGCG	ATACGTAATG	CGAATTGCAG
CMW 13807		GCGAACTGCG	ATACGTAATG	CGAATTGCAG
CMW 13808		GCGAACTGCG	ATACGTAATG	CGAATTGCAG
CMW 13815		GCGAACTGCG	ATACGTAATG	CGAATTGCAG
CMW 13817		GCGAACTGCG	ATACGTAATG	CGAATTGCAG
CMW 13810		GCGAACTGCG	ATACGTAATG	CGAATTGCAG
<i>P. aphanidermatum</i>	AJ233438	GCGAACTGCG	ATACGTAATG	CGAATTGCAG
<i>P. deliense</i>	AJ233442	GCGAACTGCG	ATACGTAATG	CGAATTGCAG
<i>P. dissotocum</i>	AJ233443	GCGAACTGCG	ATACGTAATG	CGAATTGCAG
<i>P. helicoides</i>	AB108061	GCGAACTGCG	ATACGTAATG	CGAATTGCAG
<i>P. intermedium</i>	AJ233447	GCGAACTGCG	ATACGTAATG	CGAATTGCAG
<i>P. irregulare</i>	AJ233448	GCGAACTGCG	ATACGTAATG	CGAATTGCAG
<i>P. ultimum</i>	AB108064	GCGAACTGCG	ATACGTAATG	CGAATTGCAG
<i>P. vexans</i>	AJ233462	GCGAACTGCG	ATACGTAATG	CGAATTGCAG
<i>Phytophthora cinnamomi</i>	AF266764	GCGAACTGCG	ATACGTAATG	CGAATTGCAG

				420
CMW 13819		AATTCAGTGA	GTCATCGAAA	TTTTGAACGC
CMW 13822		AATTCAGTGA	GTCATCGAAA	TTTTGAACGC
CMW 13805		AATTCAGTGA	GTCATCGAAA	TTTTGAACGC
CMW 13796		AATTCAGTGA	GTCATCGAAA	TTTTGAACGC
CMW 13820		AATTCAGTGA	GTCATCGAAA	TTTTGAACGC
CMW 13807		AATTCAGTGA	GTCATCGAAA	TTTTGAACGC
CMW 13808		GATTCAGTGA	GTCATCGAAA	TTTTGAACGC
CMW 13815		AATTCAGTGA	GTCATCGAAA	TTTTGAACGC
CMW 13817		GATTCAGTGA	GTCATCGAAA	TTTTGAACGC
CMW 13810		AATTCAGTGA	GTCATCGAAA	TTTTGAACGC
<i>P. aphanidermatum</i>	AJ233438	AATTCAGTGA	GTCATCGAAA	TTTTGAACGC
<i>P. deliense</i>	AJ233442	AATTCAGTGA	GTCATCGAAA	TTTTGAACGC
<i>P. dissotocum</i>	AJ233443	AATTCAGTGA	GTCATCGAAA	TTTTGAACGC
<i>P. helicoides</i>	AB108061	GATTCAGTGA	GTCATCGAAA	TTTTGAACGC
<i>P. intermedium</i>	AJ233447	AATTCAGTGA	GTCATCGAAA	TTTTGAACGC
<i>P. irregulare</i>	AJ233448	AATTCAGTGA	GTCATCGAAA	TTTTGAACGC
<i>P. ultimum</i>	AB108064	AATTCAGTGA	GTCATCGAAA	TTTTGAACGC
<i>P. vexans</i>	AJ233462	AATTCAGTGA	GTCATCGAAA	TTTTGAACGC
<i>Phytophthora cinnamomi</i>	AF266764	GATTCAGTGA	GTCATCGAAA	TTTTGAACGC

				450
CMW 13819		ATATTGCACT	TTCGGGTTAT	GCCTGGAAGT
CMW 13822		ATATTGCACT	TCCGGGTTAT	GCCTGGAAGT
CMW 13805		ACATTGCACT	TTCGGGTTAT	GCCTGGAAGT
CMW 13796		ATATTGCACT	TCCGGGTTAT	GCCTGGAAGT
CMW 13820		ACATTGCACT	TTCGGGTTAT	GCCTGGAAGT
CMW 13807		ACATTGCACT	TTCGGGTTAT	GCCTGGAAGT
CMW 13808		ATATTGCACT	TTCGGGTTAT	GCCTGGAAGT
CMW 13815		ACATTGCACT	TTCGGGTTAT	GCCTGGAAGT
CMW 13817		ATATTGCACT	TTCGGGTTAT	GCCTGGAAGT
CMW 13810		ACATTGCACT	TTCGGGTTAT	GCCTGGAAGT
<i>P. aphanidermatum</i>	AJ233438	ACATTGCACT	TTCGGGTTAT	GCCTGGAAGT
<i>P. deliense</i>	AJ233442	ACATTGCACT	TTCGGGTTAT	GCCTGGAAGT
<i>P. dissotocum</i>	AJ233443	ACATTGCACT	TTCGGGTTAT	GCCTGGAAGT
<i>P. helicoides</i>	AB108061	ATATTGCACT	TTCGGGTTAT	GCCTGGAAGT
<i>P. intermedium</i>	AJ233447	ATATTGCACT	TCCGGGTTAT	GCCTGGAAGT
<i>P. irregulare</i>	AJ233448	ATATTGCACT	TCCGGGTTAT	GCCTGGAAGT
<i>P. ultimum</i>	AB108064	ATATTGCACT	TTCGGGTTAT	GCCTGGAAGT
<i>P. vexans</i>	AJ233462	ATATTGCACT	TTCGGGTTAT	GCCTGGAAGT
<i>Phytophthora cinnamomi</i>	AF266764	ATATTGCACT	TCCGGGTTAT	TCCTGGGAGT



CMW 13819		ATGTCTGTAT	CAGTGTCCGT	ACCTCAACCT	480
CMW 13822		ATGTCTGTAT	CAGTGTCCGT	AAATCAACCT	
CMW 13805		ATGCCTGTAT	CAGTGTCCGT	ACATCAAACCT	
CMW 13796		ATGTCTGTAT	CAGTGTCCGT	AAATCAACCT	
CMW 13820		ATGCCTGTAT	CAGTGTCCGT	ACATCAAACCT	
CMW 13807		ATGCCTGTAT	CAGTGTCCGT	ACATCAAACCT	
CMW 13808		ATGTCTGTAT	CAGTGTCCGT	ACACTAAACT	
CMW 13815		ATGCCTGTAT	CAGTGTCCGT	ACATCAAACCT	
CMW 13817		ATGTCTGTAT	CAGTGTCCGT	ACACTAAACT	
CMW 13810		ATGCCTGTAT	CAGTGTCCGT	ACATCAAACCT	
<i>P. aphanidermatum</i>	AJ233438	ATGCCTGTAT	CAGTGTCCGT	ACATCAAACCT	
<i>P. deliense</i>	AJ233442	ATGCCTGTAT	CAGTGTCCGT	ACATCAAACCT	
<i>P. dissotocum</i>	AJ233443	ATGCCTGTAT	CAGTGTCCGT	ACATCAAACCT	
<i>P. helicoides</i>	AB108061	ATGTCTGTAT	CAGTGTCCGT	ACACTAAACT	
<i>P. intermedium</i>	AJ233447	ATGTCTGTAT	CAGTGTCCGT	AAATCAACCT	
<i>P. irregulare</i>	AJ233448	ATGTCTGTAT	CAGTGTCCGT	AAATCAACCT	
<i>P. ultimum</i>	AB108064	ATGTCTGTAT	CAGAGTCCGT	AAATCAACCT	
<i>P. vexans</i>	AJ233462	ATGTCTGTAT	CAGTGTCCGT	ACCTCAACCT	
<i>Phytophthora cinnamomi</i>	AF266764	ATGCCTGTAT	CAGTGTCCGT	ACATCAAACCT	

CMW 13819		TGCCTTTCTT	T-TCCTGTGT	AGTCAG—	510
CMW 13822		TGCCTTTCTT	CCTTCTGTGT	AGTCAG—	
CMW 13805		TGCCTTTCTT	T-TTCTGTGT	AGTCAG—	
CMW 13796		TGCCTTTCTT	CCTTCTGTGT	AGTCAG—	
CMW 13820		TGCCTTTCTT	T-TTTTGTGT	AGTCAA—	
CMW 13807		TGCCTTTCTT	T-TTTTGTGT	AGTCAA—	
CMW 13808		TGCCCTCTTT	G-CGTCGTGT	AGTCGTCGCG	
CMW 13815		TGCCTTTCTT	TTTCT-GTGT	AGTCAG—	
CMW 13817		TGCCCTCTTT	GCGTC-GTGT	AGTCGTCGCG	
CMW 13810		TGCCTTTCTT	TTTCT-GTGT	AGTCAG—	
<i>P. aphanidermatum</i>	AJ233438	TGCCTTTCTT	TTTCT-GTGT	AGTCAG—	
<i>P. deliense</i>	AJ233442	TGCCTTTCTT	T-TTCTGTGT	AGTCAG—	
<i>P. dissotocum</i>	AJ233443	TGCCTTTCTT	T-TTTTGTGT	AGTCAA—	
<i>P. helicoides</i>	AB108061	TGCCCTCTTT	GCGTC-GTGT	AGTCGTCGCG	
<i>P. intermedium</i>	AJ233447	TGCCTTTCTT	CCTTCTGTGT	AGTCAG—	
<i>P. irregulare</i>	AJ233448	TGCGTTTCTT	CCTTCCGTGT	AGTCGG—	
<i>P. ultimum</i>	AB108064	TGCCTTTCTT	TTTCT-GTGT	AGTCAG—	
<i>P. vexans</i>	AJ233462	TGCCTTTCTT	T-TCCTGTGT	AGTCAG—	
<i>Phytophthora cinnamomi</i>	AF266764	TGGCTCTCTT	CCTTCCGTGT	AGTCGGTG-G	

CMW 13819		—GAGAGGAA	ACGAGCAGAC	TTGAAGTGTC	540
CMW 13822		—TGGAGGAT	GTGGC-AGAC	GTGAAGTGTC	
CMW 13805		—GGAGAGAG	ATGGCAGAAT	GTGAGGTGTC	
CMW 13796		—TGGAGGAT	GTGGC-AGAC	GTGAAGTGTC	
CMW 13820		—GAAGAGAG	ATGGCAGACT	GTGAGGTGTC	
CMW 13807		—GAAGAGAG	ATGGCAGACT	GTGAGGTGTC	
CMW 13808		TTGGAAATTT	GTGGCAGA-T	GTGAGGTGTC	
CMW 13815		—GGAGAGAG	ATGGCAGAAT	GTGAGGTGTC	
CMW 13817		TTGGAAATTT	GTGGCAGA-T	GTGAGGTGTC	
CMW 13810		—GGAGAGAG	ATGGCAGAAT	GTGAGGTGTC	
<i>P. aphanidermatum</i>	AJ233438	—GGAGAGAG	ATGGCAGAAT	GTGAGGTGTC	
<i>P. deliense</i>	AJ233442	—GGAGAGAG	ATGGCAGAAT	GTGAGGTGTC	
<i>P. dissotocum</i>	AJ233443	—GAAGAGAG	ATGGCAGACT	GTGAGGTGTC	
<i>P. helicoides</i>	AB108061	TTGGAAATTT	GTGGCAGA-T	GTGAGGTGTC	
<i>P. intermedium</i>	AJ233447	—TGGAGGAT	GTGGCAGAC-	GTGAAGTGTC	
<i>P. irregulare</i>	AJ233448	—TGGAGGAG	AGTTGCAGAT	GTGAAGTGTC	
<i>P. ultimum</i>	AB108064	—GGATGGAA	TGTGCAGA-T	GTGAAGTGTC	
<i>P. vexans</i>	AJ233462	—GAGAGGAA	ACGAGCAGAC	TTGAAGTGTC	
<i>Phytophthora cinnamomi</i>	AF266764	ATGGAG—	GTGCCAGA-C	GTGAGGTGTC	



CMW 13819		TTTCTACGGA	G-TGTGGCGC	TCGAAAGGCG	660
CMW 13822		TT-CTGCGAG	G-TGCTGTGC	TCGAA-CGCG	
CMW 13805		CTAAGATGAA	G-TGTGATTC	TCGAATCGCA	
CMW 13796		TT-CTGCGAG	G-TGCTGTGC	TCGAA-CGCG	
CMW 13820		TTAAGATGAA	G-TGTGACTT	TCGAA-CGCA	
CMW 13807		TTAAGATGAA	G-TGTGACTT	TCGAA-CGCA	
CMW 13808		GTTGGGTGCC	G-GTGGGCTG	TGGGA-CGCG	
CMW 13815		CTAAGATGAA	G-TGTGATTC	TCGAATCGCG	
CMW 13817		GTTGGGTGCC	G-GTGGGCT	GTGGGACGCG	
CMW 13810		CTAAGATGAA	G-TGTGATTC	TCGAATCGCG	
<i>P. aphanidermatum</i>	AJ233438	CTAAGATGAA	G-TGTGATTC	TCGAATCGCG	
<i>P. deliense</i>	AJ233442	CTAAGATGAA	G-TGTGATTC	TCGAATCGCA	
<i>P. dissotocum</i>	AJ233443	TTAAGATGAA	G-TGTGACTT	TCGAA-CGCA	
<i>P. helicoides</i>	AB108061	GTTGGGTGCC	GGTG-GGCT	GTGGGACGCG	
<i>P. intermedium</i>	AJ233447	TTCTG-CGAG	G-TGCTGTGC	TCGAA-CGCG	
<i>P. irregulare</i>	AJ233448	AT-GTGCGCG	G-TGCTGTGC	GTGAA-CGCG	
<i>P. ultimum</i>	AB108064	TTTCTATGAA	G-TGTAATGG	TTGAAAGGCA	
<i>P. vexans</i>	AJ233462	TTTCTACGGA	G-TGTGGCGC	TCGAAAGGCG	
<i>Phytophthora cinnamomi</i>	AF266764	AAAGCGTGAC	GTTGCTGGTT	GTGGAGGCTG	

CMW 13819		GCGGTTTCCT	TCGGGATCGC	TCGCACTCGG	690
CMW 13822		GTGGTTTTCC	—GGATCGC	TCGCGGC-TG	
CMW 13805		GTGATCTGTT	T—GGATCGC	TTTGCGCATT	
CMW 13796		GTGGTTTTCC	—GGATCGC	TCGCGGC-TG	
CMW 13820		GTGATCTGTT	T—GGATCGC	TTTGCTCGAG	
CMW 13807		GTGATCTGTT	T—GGATCGC	TTTGCTCGAG	
CMW 13808		—TCTGTT	—GACGAG	TCT—	
CMW 13815		GTGATCTGTT	T—GGATCGC	TTTGCGCATT	
CMW 13817		—TCTGTT	G—A—	—CGAGTC	
CMW 13810		GTGATCTGTT	T—GGATCGC	TTTGCGCATT	
<i>P. aphanidermatum</i>	AJ233438	GTGATCTGTT	T—GGATCGC	TTTGCGCATT	
<i>P. deliense</i>	AJ233442	GTGATCTGTT	T—GGATCGC	TTTGCTCGAG	
<i>P. dissotocum</i>	AJ233443	GTGATCTGTT	T—GGATCGC	TTTGCTCGAG	
<i>P. helicoides</i>	AB108061	—TCTGT—	T—GACGAGT	CTGGCGACC—	
<i>P. intermedium</i>	AJ233447	GTGGTTT—T	C—GGATCGC	TC—GCCGCTG	
<i>P. irregulare</i>	AJ233448	GTGGTTTTCC	—GGATCGC	TCGCGGC-TG	
<i>P. ultimum</i>	AB108064	GTGATTT—T	C—GGATTGC	TG—GCGGCTT	
<i>P. vexans</i>	AJ233462	GCGGTTTCCT	TCGGGATCGC	TCGCACTCGG	
<i>Phytophthora cinnamomi</i>	AF266764	—CCTGTA	T—GGCCAGT	C—GGCGACC—	

CMW 13819		ACGGCGAC-T	TTGGCGAATA	CAT-ATGGGA	720
CMW 13822		TTGGCGAC-T	TCGGTGAATG	CATTATGGAG	
CMW 13805		TGGGCGAC-T	TCGGTTAGGA	CATTAAGGA	
CMW 13796		TTGGCGAC-T	TCGGTGAATG	CATTATGGAG	
CMW 13820		TGGGCGAC-T	TCGGTTAGGA	CATTAAGGA	
CMW 13807		TGGGCGAC-T	TCGGTTAGGA	CATTAAGGA	
CMW 13808		—GGCGACCT	TTGGTGCGTG	CATGCTTGG—	
CMW 13815		TGGGCGAC-T	TCGGTTAGGA	CATTAAGGA	
CMW 13817		TGGGCGAC-T	TTGGTGCGTG	CATGCTTGG—	
CMW 13810		TGGGCGAC-T	TCGGTTAGGA	CATTAAGGA	
<i>P. aphanidermatum</i>	AJ233438	TGGGCGAC-T	TCGGTTAGGA	CATTAAGGA	
<i>P. deliense</i>	AJ233442	TGGGCGAC-T	TCGGTTAGGA	CATTAAGGA	
<i>P. dissotocum</i>	AJ233443	TGGGCGAC-T	TCGGTTAGGA	CATTAAGGA	
<i>P. helicoides</i>	AB108061	———T	TTGGTGCG—	TGCATGCTTG	
<i>P. intermedium</i>	AJ233447	TTGGCGAC-T	TCGGTGAATG	CATTATGGAG	
<i>P. irregulare</i>	AJ233448	TCGGCGAC-T	TCGGTGAATG	CATAATGGAG	
<i>P. ultimum</i>	AB108064	TTGGCGAC-T	TCGGTATGAA	CGT—ATGGA	
<i>P. vexans</i>	AJ233462	ACGGCGAC-T	TTGGCGAATA	CAT-ATGGGA	
<i>Phytophthora cinnamomi</i>	AF266764	—GGTTG-T	CTGCTGCGG—	CGTTAATGG	



				750
CMW 13819		AGCAGACTCG	-ACTCGCGGT	ACGTTAGGTG
CMW 13822		TG-GACCTCG	-ATTGCGGGT	ATGTTGGGC-
CMW 13805		AGCAACCTCT	-ATTGCGGGT	ATGTTAGGC-
CMW 13796		TG-GACCTCG	-ATTGCGGGT	ATGTTGGGC-
CMW 13820		AGCAACCTCT	-ATTGCGGGT	ATGTTAGGC-
CMW 13807		AGCAACCTCT	-ATTGCGGGT	ATGTTAGGC-
CMW 13808		-GCACTGTGT	-ATTGCGGGT	ATGTTAGGC-
CMW 13815		AGCAACCTCT	-ATTGCGGGT	ATGTTAGGC-
CMW 13817		-GCACTGTGT	-ATTGCGGGT	ATGTTAGGC-
CMW 13810		AGCAACCTCT	-ATTGCGGGT	ATGTTAGGC-
<i>P. aphanidermatum</i>	AJ233438	AGCAACCTCT	-ATTGCGGGT	ATGTTAGGC-
<i>P. deliense</i>	AJ233442	AGCAACCTCT	-ATTGCGGGT	ATGTTAGGC-
<i>P. dissotocum</i>	AJ233443	AGCAACCTCT	-ATTGCGGGT	ATGTTAGGC-
<i>P. helicoides</i>	AB108061	GGCACTGTGT	-ATTGCGGGT	ATGTTAGGC-
<i>P. intermedium</i>	AJ233447	TG-GACCTCG	-ATTGCGGGT	ATGTTGGGC-
<i>P. irregulare</i>	AJ233448	TG-GACCTCG	-ATTGCGGGT	ATGTTGGGC-
<i>P. ultimum</i>	AB108064	GACTAGCTCA	-ATTGCGGGT	ATGTTAGGC-
<i>P. vexans</i>	AJ233462	AGCAGACTCG	-ACTCGCGGT	ACGTTAGGTG
<i>Phytophthora cinnamomi</i>	AF266764	AGGAGTGTCC	GATTGCGGGT	ATGGTTGGC-
				780
CMW 13819		TGTTGCTTTT	GCGGCGTGCT	GAACAATGTT
CMW 13822		——TTC	G——GCT	GGACAATGTT
CMW 13805		——TTC	G——GCC	CGACGTTGCA
CMW 13796		——TTC	G——GCT	GGACAATGTT
CMW 13820		——TTC	G——GCC	CGACTTTGCA
CMW 13807		——TTC	G——GCC	CGACTTTGCA
CMW 13808		——TGC	G——TTC	GCGCGGCTTT
CMW 13815		——TTC	G——GCC	CGACGTTGCA
CMW 13817		——TGC	G——TTC	G——CGCG
CMW 13810		——TTC	G——GCC	CGACGTTGCA
<i>P. aphanidermatum</i>	AJ233438	——TTC	G——GCC	CGACGTTGCA
<i>P. deliense</i>	AJ233442	——TTC	G——GCC	CGACGTTGCA
<i>P. dissotocum</i>	AJ233443	——TTC	G——GCC	CGACTTTGCA
<i>P. helicoides</i>	AB108061	——TGC	G——TTC	G——CGCG
<i>P. intermedium</i>	AJ233447	——TTC	G——GCT	GGACAATGTT
<i>P. irregulare</i>	AJ233448	——TTC	G——GCT	GGACAATGTT
<i>P. ultimum</i>	AB108064	——TTC	G——GCT	CGACAATGTT
<i>P. vexans</i>	AJ233462	TGTTGCTTTT	GCGGCGTGCT	GAACAATGTT
<i>Phytophthora cinnamomi</i>	AF266764	——TCC	G——GCT	G——
				810
CMW 13819		GCGTGTGTG	GTCTTGTTT	CTGTGTTGCG
CMW 13822		GCTTATTGTG	TGTTTGTT—	CCGCGTTGCG
CMW 13805		GC-TGACGGA	GTGTGGTTTT	CTGTTCTTTC
CMW 13796		GCTTATTGTG	TGTTTGTT—	CCGCGTTGCG
CMW 13820		GCTGACTGGA	G—TTGTTT	CTGTTCTTTC
CMW 13807		GCTGACTGGA	G—TTGTTT	CTGTTCTTTC
CMW 13808		GACAA-TGCA	G—CTGATGC	GTGTGTTTGG
CMW 13815		GC-TGACAGA	GTGTGGTTTT	CTGTTCTTTC
CMW 13817		GC-TTTGACA	ATGCAGCTGA	TGCGTGTGT-
CMW 13810		GC-TGACAGA	GTGTGGTTTT	CTGTTCTTTC
<i>P. aphanidermatum</i>	AJ233438	GC-TGACAGA	GTGTGGTTTT	CTGTTCTTTC
<i>P. deliense</i>	AJ233442	GC-TGACGGA	GTGTGGTTTT	CTGTTCTTTC
<i>P. dissotocum</i>	AJ233443	GCTGACTGGA	GTTGTTTT—	CTGTTCTTTC
<i>P. helicoides</i>	AB108061	GC-TTTGACA	ATGCAGCTGA	TGCGTGTGT
<i>P. intermedium</i>	AJ233447	GCTTATTGTG	TGTTTGTT—	CCGCGTTGCG
<i>P. irregulare</i>	AJ233448	GCTTATTGTG	TGTTTGTT—	CCGCGTTGCG
<i>P. ultimum</i>	AB108064	GCGTAATTGT	GTGTGGTCTT	-TGTTTGTCG
<i>P. vexans</i>	AJ233462	GCGTGTGTG	GTCTTGTTTC	CTGTGTTGCG
<i>Phytophthora cinnamomi</i>	AF266764	——AACA	AAGCGCTTAT	TGGATGTTCC



840

CMW 13819		TTCGAGGTGT	ACTGTCTAAT	GGCTG-TGGG
CMW 13822		CTTGAGGTGT	ACTTTCTGCT	GTGTGCTTGA
CMW 13805		CTTGAGGTGT	AC—CTGAT	TTGTG-TGAG
CMW 13796		CTTGAGGTGT	ACTTTCTGCT	GTGTGCTTGA
CMW 13820		CTTGAGGTGT	AC—CTGTC	TTGTG-TGAG
CMW 13807		CTTGAGGTGT	AC—CTGTC	TTGTG-TGAG
CMW 13808		GCTGTGGTGC	—————	-TGTA-TGGG
CMW 13815		CTTGAGGTGT	AC—CTGAA	TTGTG-TGAG
CMW 13817		-TTGGGCTGT	G——GTG	CTGTA-TGGG
CMW 13810		CTTGAGGTGT	AC—CTGAA	TTGTG-TGAG
<i>P. aphanidermatum</i>	AJ233438	CTTGAGGTGT	AC—CTGAA	TTGTG-TGAG
<i>P. deliense</i>	AJ233442	CTTGAGGTGT	AC—CTGAT	TTGTG-TGAG
<i>P. dissotocum</i>	AJ233443	CTTGAGGTGT	AC—CTGTC	TTGTG-TGAG
<i>P. helicoides</i>	AB108061	TGG—GCTGT	GG—TGCTG	T—A-TGGG
<i>P. intermedium</i>	AJ233447	CTTGAGGTGT	ACTTTCTGCT	GTGTGCTTGA
<i>P. irregulare</i>	AJ233448	CTTGAGGTGT	ACTGATGGCT	GTGGGATTGA
<i>P. ultimum</i>	AB108064	CTTGAGGTGT	AC—TAGAG	GT-TG-TCGG
<i>P. vexans</i>	AJ233462	TTCGAGGTGT	ACTGTCTAAT	GGCTG-TGGG
<i>Phytophthora cinnamomi</i>	AF266764	TCCCTGCTGT	GG—CGGTA	CGG-A-TCGG

851

CMW 13819		TTTCGAACCT	G
CMW 13822		ACTGGGATCT	G
CMW 13805		GCAATGGTCT	G
CMW 13796		ACTGGGATCT	G
CMW 13820		GCAATGGTCT	G
CMW 13807		GCAATGGTCT	G
CMW 13808		TGAACCGGAT	G
CMW 13815		GCAATGGTCT	G
CMW 13817		TGAACCGGAT	G
CMW 13810		GCAATGGTCT	G
<i>P. aphanidermatum</i>	AJ233438	GCAATGGTCT	G
<i>P. deliense</i>	AJ233442	GCAATGGTCT	G
<i>P. dissotocum</i>	AJ233443	GCAATGGTCT	G
<i>P. helicoides</i>	AB108061	TGAACCGGAT	G
<i>P. intermedium</i>	AJ233447	ACTGGGATCT	G
<i>P. irregulare</i>	AJ233448	ACTGGTACT	G
<i>P. ultimum</i>	AB108064	TT—TGAACC	G
<i>P. vexans</i>	AJ233462	TTTCGAACCT	G
<i>Phytophthora cinnamomi</i>	AF266764	TGAACCGTAG	G

Chapter 3.

Alignment 5. 5' end of the β - tubulin gene DNA sequence alignment of selected *Cylindrocladium* isolates from the survey *Eucalyptus* hybrid cuttings.

CMW8734		GCTGCCCTG	ATTCTACCCC	GCCGCCCCGG	30
CMW9108		GCTGCCCTG	ATTCTACCCC	GCCGCCCCGG	
CMW9190		GCTGCCCTG	ATTCTACCCC	GCCGCCCCGG	
CMW9109		GCTGCCCTG	ATTCTACCCC	GCCGCCCCGG	
CMW9189		GCTGCCCTG	ATTCTACCCC	GCCGCCCCGG	
<i>Cylindrocladium pauciramosum</i>	AY162320	GCTGCCCTG	ATTCTACCCC	GCCGCCCCGG	
<i>Cylindrocladium pauciramosum</i>	AF449448	GCTGCCCTG	ATTCTACCCC	GCCGCCCCGG	
<i>Cylindrocladium insulare</i>	AF44951	GCTGCCCTG	ATTCTACCCC	GCCGCCCCGG	
<i>Cylindrocladium insulare</i>	AF44950	GCTGCCCTG	ATTCTACCCC	GCCGCCCCGG	
<i>Cylindrocladium colhounii</i>	AF232855	GCTGCCCTG	ATTCTACCCC	GCCGCCCCGG	
<i>Cylindrocladium colhounii</i>	AF232854	GCTGCCCTG	ATTCTACCCC	GCCGCCCCGG	
<i>Cylindrocladium gracile</i>	AF333405	GCTGCCCTG	ATTCTACCCC	GCCGCCCCGG	
<i>Cylindrocladium gracile</i>	AF333404	GCTGCCCTG	ATTCTACCCC	GCCGCCCCGG	
<i>Cylindrocladium clavatum</i>	AF232850	GCTGCCCTG	ATTCTACCCC	GCCGCCCCGG	
<i>Cylindrocladium candelabrum</i>	AF210858	GCTGCCCTG	ATTCTACCCC	GCCGCCCCGG	
<i>Cylindrocladium candelabrum</i>	AF210857	GCTGCCCTG	ATTCTACCCC	GCCGCCCCGG	
<i>Cylindrocladium scoparium</i>	AF210875	GCTGCCCTG	ATTCTACCCC	GCCGCCCCGG	
<i>Cylindrocladium scoparium</i>	AF210874	GCTGCCCTG	ATTCTACCCC	GCCGCCCCGG	
<i>Cylindrocladium ilicicola</i>	AF333413	GCTGCCCTG	AGCGTACCCC	GCCGCCCCGG	
<i>Cylindrocladium ilicicola</i>	AF333412	GCTGCCCTG	AGCGTACCCC	GCCGCCCCGG	
<i>Cylindrocladiella infestans</i>	AF320190	GCTGCCCTG	ATTCTACCCC	GCCGAATCGT	
CMW8734		TTTCCACC --	-- GCTTCGAC	GACAACAAAG	60
CMW9108		TTTCCACC --	-- GCTTCGAC	GACAACAAAG	
CMW9190		TTTCCACC --	-- GCTTCGAC	GACAACAAAG	
CMW9109		TTTCCACC --	-- GCTTCGAC	GACAACAAAG	
CMW9189		TTTCCACC --	-- GCTTCGAC	GACAACAAAG	
<i>Cylindrocladium pauciramosum</i>	AY162320	TTTCCACC --	-- GCTTCGAC	GACAACAAAG	
<i>Cylindrocladium pauciramosum</i>	AF449448	TTTCCACC --	-- GCTTCGAC	GACAACAAAG	
<i>Cylindrocladium insulare</i>	AF44951	TTTCCACC --	-- ACCTCG --	- ACAACAAAG	
<i>Cylindrocladium insulare</i>	AF44950	TTTCCACC --	-- ACCTCGAC	GACAACAAAG	
<i>Cylindrocladium colhounii</i>	AF232855	TTTCCACC --	-- GCTTCGAC	GACAACAAAG	
<i>Cylindrocladium colhounii</i>	AF232854	TTTCCACC --	-- GCTTCGAC	GACAACAAAG	
<i>Cylindrocladium gracile</i>	AF333405	TTTCCACC --	-- GCTTCGAC	GACAA - AAAG	
<i>Cylindrocladium gracile</i>	AF333404	TTTCCACC --	-- GCTTCGAC	GACAACAAAG	
<i>Cylindrocladium clavatum</i>	AF232850	TTTCCACC --	-- GCTTCGAC	GACAACAAAG	
<i>Cylindrocladium candelabrum</i>	AF210858	TTTCCACC --	-- GCTTCGAC	GACAACAAAG	
<i>Cylindrocladium candelabrum</i>	AF210857	TTTCCACC --	-- GCTTCGAC	GACAACAAAG	
<i>Cylindrocladium scoparium</i>	AF210875	TTTCCACC --	-- ACATCGAC	GACAACAAAG	
<i>Cylindrocladium scoparium</i>	AF210874	TTTCCACC --	-- ACATCGAC	GACAACAAAG	
<i>Cylindrocladium ilicicola</i>	AF333413	TTTCCACC --	-- GCTTCGAC	AACAACAAAG	
<i>Cylindrocladium ilicicola</i>	AF333412	TTTCCACC --	-- GCTTCGAC	AACAACAAAG	
<i>Cylindrocladiella infestans</i>	AF320190	TTTCCACCCA	CCGCCTCGAC	AACAACAAAG	



CMW8734		TCAGTGCCTA	AGTACTCTTC	TCAACTCCAA	180
CMW9108		TCAGTGCCTA	AGTACTCTTC	TCAACTCCAA	
CMW9190		TCAGTGCCTA	AGTACTCTTC	TCAACTCCAA	
CMW9109		TCAGTGCCTA	AGTACTCTTC	TCAACTCCAA	
CMW9189		TCAGTGCCTA	AGTACTCTTC	TCAACTCCAA	
<i>Cylindrocladium pauciramosum</i>	AY162320	TCAGTGCCTA	AGTACTCTTC	TCAACTCCAA	
<i>Cylindrocladium pauciramosum</i>	AF449448	TCAGTGCCTA	AGTACTCTTC	TCAACTCCAA	
<i>Cylindrocladium insulare</i>	AF44951	TCAGTGCCTA	AGTACTCTTC	TCAACTCCAA	
<i>Cylindrocladium insulare</i>	AF44950	TCAGTGCCTA	AGTACTCTTC	TCAACTCCAA	
<i>Cylindrocladium colhounii</i>	AF232855	TCAGTGCCTA	AGTACTCTTC	TCAACTCCAA	
<i>Cylindrocladium colhounii</i>	AF232854	TCAGTGCCTA	AGTACTCTTC	TCAACTCCAA	
<i>Cylindrocladium gracile</i>	AF333405	TCAGTGCCTA	AGTACTCTTC	TCAACTCCAA	
<i>Cylindrocladium gracile</i>	AF333404	TCAGTGCCTA	AGTACTCTTC	TCAACTCCAA	
<i>Cylindrocladium clavatum</i>	AF232850	TCAGTGCCTA	AGTACTCTTC	TCAACTCCAA	
<i>Cylindrocladium candelabrum</i>	AF210858	TCAGTGCCTA	AGTACTCTTC	TCAACTCCAA	
<i>Cylindrocladium candelabrum</i>	AF210857	TCAGTGCCTA	AGTACTCTTC	TCAACTCCAA	
<i>Cylindrocladium scoparium</i>	AF210875	TCAGTGCCTA	AGTACTCTTC	TCAACTCCAA	
<i>Cylindrocladium scoparium</i>	AF210874	TCAGTGCCTA	AGTACTCTTC	TCAACTCCAA	
<i>Cylindrocladium ilicicola</i>	AF333413	TCAGTGCCTA	AGTACTCTTC	TCAACTCCAA	
<i>Cylindrocladium ilicicola</i>	AF333412	TCAGTGCCTA	AGTACTCTTC	TCAACTCCAA	
<i>Cylindrocladiella infestans</i>	AF320190	TCAGTGCCTA	AGTACTCTTC	TCAACTCCAA	
CMW8734		CAAAATTCTC	ACGACGAGAT	TCACTGACAG	210
CMW9108		CAAAATTCTC	ACGACGAGAT	TCACTGACAG	
CMW9190		CAAAATTCTC	ACGACGAGAT	TCACTGACAG	
CMW9109		CAAAATTCTC	ACGACGAGAT	TCACTGACAG	
CMW9189		CAAAATTCTC	ACGACGAGAT	TCACTGACAG	
<i>Cylindrocladium pauciramosum</i>	AY162320	CAAAATTCTC	ACGACGAGAT	TCACTGACAG	
<i>Cylindrocladium pauciramosum</i>	AF449448	CAAAATTCTC	ACGACGAGAT	TCACTGACAG	
<i>Cylindrocladium insulare</i>	AF44951	CAAAATTCTC	ACGACGAGAT	TCACTGACAG	
<i>Cylindrocladium insulare</i>	AF44950	CAAAATTCTC	ACGACGAGAT	TCACTGACAG	
<i>Cylindrocladium colhounii</i>	AF232855	CAAAATTCTC	ACGACGAGAT	TCACTGACAG	
<i>Cylindrocladium colhounii</i>	AF232854	CAAAATTCTC	ACGACGAGAT	TCACTGACAG	
<i>Cylindrocladium gracile</i>	AF333405	CAAAATTCTC	ACGACGAGAT	TCACTGACAG	
<i>Cylindrocladium gracile</i>	AF333404	CAAAATTCTC	ACGACGAGAT	TCACTGACAG	
<i>Cylindrocladium clavatum</i>	AF232850	CAAAATTCTC	ACGACGAGAT	TCACTGACAG	
<i>Cylindrocladium candelabrum</i>	AF210858	CAAAATTCTC	ACGACGAGAT	TCACTGACAG	
<i>Cylindrocladium candelabrum</i>	AF210857	CAAAATTCTC	ACGACGAGAT	TCACTGACAG	
<i>Cylindrocladium scoparium</i>	AF210875	CAAAATTCTC	ACGACGAGAT	TCACTGACAG	
<i>Cylindrocladium scoparium</i>	AF210874	CAAAATTCTC	ACGACGAGAT	TCACTGACAG	
<i>Cylindrocladium ilicicola</i>	AF333413	CAAAATTCTC	ACGACGAGAT	TCACTGACAG	
<i>Cylindrocladium ilicicola</i>	AF333412	CAAAATTCTC	ACGACGAGAT	TCACTGACAG	
<i>Cylindrocladiella infestans</i>	AF320190	CAAAATTCTC	ACGACGAGAT	TCACTGACAG	
CMW8734		T - TGTCGATA	GGGTAACCAA	ATTGGTGCTG	240
CMW9108		T - TGTCGATA	GGGTAACCAA	ATTGGTGCTG	
CMW9190		T - TGTCGATA	GGGTAACCAA	ATTGGTGCTG	
CMW9109		T - TGTCGATA	GGGTAACCAA	ATTGGTGCTG	
CMW9189		T - TGTCGATA	GGGTAACCAA	ATTGGTGCTG	
<i>Cylindrocladium pauciramosum</i>	AY162320	T - TGTCGATA	GGGTAACCAA	ATTGGTGCTG	
<i>Cylindrocladium pauciramosum</i>	AF449448	T - TGTCGATA	GGGTAACCAA	ATTGGTGCTG	
<i>Cylindrocladium insulare</i>	AF44951	T - TATCGACA	GGGTAACCAA	ATTGGTGCTG	
<i>Cylindrocladium insulare</i>	AF44950	T - TATCGACA	GGGTAACCAA	ATTGGTGCTG	
<i>Cylindrocladium colhounii</i>	AF232855	T - TGTCGATA	GGGTAACCAA	ATTGGTGCTG	
<i>Cylindrocladium colhounii</i>	AF232854	A - TGTCGATA	GGGTAACCAA	ATTGGTGCTG	
<i>Cylindrocladium gracile</i>	AF333405	T - TGTCGATA	GGGTAACCAA	ATTGGTGCTG	
<i>Cylindrocladium gracile</i>	AF333404	T - TGTCGATA	GGGTAACCAA	ATTGGTGCTG	
<i>Cylindrocladium clavatum</i>	AF232850	T - TGTCGATA	GGGTAACCAA	ATTGGTGCTG	
<i>Cylindrocladium candelabrum</i>	AF210858	T - TGTCGATA	GGGTAACCAA	ATTGGTGCTG	
<i>Cylindrocladium candelabrum</i>	AF210857	T - TGTCGATA	GGGTAACCAA	ATTGGTGCTG	
<i>Cylindrocladium scoparium</i>	AF210875	T - TGTCGATA	GGGTAACCAA	ATTGGTGCTG	
<i>Cylindrocladium scoparium</i>	AF210874	T - TGTCGATA	GGGTAACCAA	ATTGGTGCTG	
<i>Cylindrocladium ilicicola</i>	AF333413	T - GGCGATCA	GGGTAACCAA	ATTGGTGCTG	
<i>Cylindrocladium ilicicola</i>	AF333412	T - GGCGATCA	GGGTAACCAA	ATTGGTGCTG	
<i>Cylindrocladiella infestans</i>	AF320190	TGTCGATA	GGGTAACCAA	ATTGGTGCTG	



CMW8734		CTTTCTGGCA	GACCATTTCT	GGCGAGCACG	270
CMW9108		CTTTCTGGCA	GACCATTTCT	GGCGAGCACG	
CMW9190		CTTTCTGGCA	GACCATTTCT	GGCGAGCACG	
CMW9109		CTTTCTGGCA	GACCATTTCT	GGCGAGCACG	
CMW9189		CTTTCTGGCA	GACCATTTCT	GGCGAGCACG	
<i>Cylindrocladium pauciramosum</i>	AY162320	CTTTCTGGCA	GACCATTTCT	GGCGAGCACG	
<i>Cylindrocladium pauciramosum</i>	AF449448	CTTTCTGGCA	GACCATTTCT	GGCGAGCACG	
<i>Cylindrocladium insulare</i>	AF44951	CTTTCTGGCA	GACCATTTCT	GGCGAGCACG	
<i>Cylindrocladium insulare</i>	AF44950	CTTTCTGGCA	GACCATTTCT	GGCGAGCACG	
<i>Cylindrocladium colhounii</i>	AF232855	CTTTCTGGCA	GACCATCTCT	GGCGAGCACG	
<i>Cylindrocladium colhounii</i>	AF232854	CTTTCTGGCA	GACCATCTCT	GGCGAGCACG	
<i>Cylindrocladium gracile</i>	AF333405	CTTTCTGGCA	GACCATCTCT	GGCGAGCACG	
<i>Cylindrocladium gracile</i>	AF333404	CTTTCTGGCA	GACCATCTCT	GGCGAGCACG	
<i>Cylindrocladium clavatum</i>	AF232850	CTTTCTGGCA	GACCATCTCT	GGCGAGCACG	
<i>Cylindrocladium candelabrum</i>	AF210858	CTTTCTGGCA	GACCATCTCT	GGCGAGCACG	
<i>Cylindrocladium candelabrum</i>	AF210857	CTTTCTGGCA	GACCATCTCT	GGCGAGCACG	
<i>Cylindrocladium scoparium</i>	AF210875	CTTTCTGGCA	GACCATTTCT	GGCGAGCACG	
<i>Cylindrocladium scoparium</i>	AF210874	CTTTCTGGCA	GACCATTTCT	GGCGAGCACG	
<i>Cylindrocladium ilicicola</i>	AF333413	CTTTCTGGCA	GACCATCTCT	GGCGAGCACG	
<i>Cylindrocladium ilicicola</i>	AF333412	CTTTCTGGCA	GACCATCTCT	GGCGAGCACG	
<i>Cylindrocladiella infestans</i>	AF320190	CTTTCTGGCA	GACCATCTCT	GGCGAGCACG	
CMW8734		GTCTCGACAG	CAATGGTGTC	TACGCCGGTA	300
CMW9108		GTCTCGACAG	CAATGGTGTC	TACGCCGGTA	
CMW9190		GTCTCGACAG	CAATGGTGTC	TACGCCGGTA	
CMW9109		GTCTCGACAG	CAATGGTGTC	TACGCCGGTA	
CMW9189		GTCTCGACAG	CAATGGTGTC	TACGCCGGTA	
<i>Cylindrocladium pauciramosum</i>	AY162320	GTCTCGACAG	CAATGGTGTC	TACGCCGGTA	
<i>Cylindrocladium pauciramosum</i>	AF449448	GTCTCGACAG	CAATGGTGTC	TACGCCGGTA	
<i>Cylindrocladium insulare</i>	AF44951	GTCTCGACAG	CAATGGTGTC	TACACTGGTA	
<i>Cylindrocladium insulare</i>	AF44950	GTCTCGACAG	CAATGGTGTC	TACGCTGGTA	
<i>Cylindrocladium colhounii</i>	AF232855	GTCTCGACAG	CAATGGTGTC	TACGCTGGTA	
<i>Cylindrocladium colhounii</i>	AF232854	GTCTCGACAG	CAATGGTGTC	TACGCCGGTA	
<i>Cylindrocladium gracile</i>	AF333405	GTCTCGACAG	CAATGGTGTC	TACGCCGGTA	
<i>Cylindrocladium gracile</i>	AF333404	GTCTCGACAG	CAATGGTGTC	TACGCCGGTA	
<i>Cylindrocladium clavatum</i>	AF232850	GTCTCGACAG	CAATGGTGTC	TACGCCGGTA	
<i>Cylindrocladium candelabrum</i>	AF210858	GTCTCGACAG	CAATGGTGTC	TACGCCGGTA	
<i>Cylindrocladium candelabrum</i>	AF210857	GTCTCGACAG	CAATGGTGTC	TACGCCGGTA	
<i>Cylindrocladium scoparium</i>	AF210875	GTCTCGACAG	CAATGGTGTC	TACGCCGGTA	
<i>Cylindrocladium scoparium</i>	AF210874	GTCTCGACAG	CAATGGTGTC	TACGCCGGTA	
<i>Cylindrocladium ilicicola</i>	AF333413	GTCTCGACAG	CAATGGTGTC	TACAACGGTA	
<i>Cylindrocladium ilicicola</i>	AF333412	GTCTCGACAG	CAATGGTGTC	TACAACGGTA	
<i>Cylindrocladiella infestans</i>	AF320190	GTCTCGACAG	CAATGGTGTC	TACAACGGTA	
CMW8734		CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG	330
CMW9108		CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG	
CMW9190		CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG	
CMW9109		CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG	
CMW9189		CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG	
<i>Cylindrocladium pauciramosum</i>	AY162320	CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG	
<i>Cylindrocladium pauciramosum</i>	AF449448	CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG	
<i>Cylindrocladium insulare</i>	AF44951	CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG	
<i>Cylindrocladium insulare</i>	AF44950	CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG	
<i>Cylindrocladium colhounii</i>	AF232855	CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG	
<i>Cylindrocladium colhounii</i>	AF232854	CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG	
<i>Cylindrocladium gracile</i>	AF333405	CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG	
<i>Cylindrocladium gracile</i>	AF333404	CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG	
<i>Cylindrocladium clavatum</i>	AF232850	CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG	
<i>Cylindrocladium candelabrum</i>	AF210858	CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG	
<i>Cylindrocladium candelabrum</i>	AF210857	CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG	
<i>Cylindrocladium scoparium</i>	AF210875	CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG	
<i>Cylindrocladium scoparium</i>	AF210874	CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG	
<i>Cylindrocladium ilicicola</i>	AF333413	CCTCCGAGCT	CCAGTTGGAG	CGCATGAACG	
<i>Cylindrocladium ilicicola</i>	AF333412	CCTCCGAGCT	CCAGTTGGAG	CGCATGAACG	
<i>Cylindrocladiella infestans</i>	AF320190	GCTCTGAGCT	CCAGCTCGAG	CGCATGAGCG	



CMW8734		GCAACAAGTT	CGTTCCTCGC	GCTGTCCTCG	450
CMW9108		GCAACAAGTT	CGTTCCTCGC	GCTGTCCTCG	
CMW9190		GCAACAAGTT	CGTTCCTCGC	GCTGTCCTCG	
CMW9109		GCAACAAGTT	CGTTCCTCGC	GCTGTCCTCG	
CMW9189		GCAACAAGTT	CGTTCCTCGC	GCTGTCCTCG	
<i>Cylindrocladium pauciramosum</i>	AY162320	GCAACAAGTT	CGTTCCTCGC	GCTGTCCTCG	
<i>Cylindrocladium pauciramosum</i>	AF449448	GCAACAAGTT	CGTTCCTCGC	GCTGTCCTCG	
<i>Cylindrocladium insulare</i>	AF44951	GCAACAAGTT	CGTTCCTCGC	GCTGTCCTCG	
<i>Cylindrocladium insulare</i>	AF44950	GCAACAAGTT	CGTTCCTCGC	GCTGTCCTCG	
<i>Cylindrocladium colhounii</i>	AF232855	GCAACAAGTT	CGTTCCTCGC	GCTGTCCTCG	
<i>Cylindrocladium colhounii</i>	AF232854	GCAACAAGTT	CGTTCCTCGC	GCTGTCCTCG	
<i>Cylindrocladium gracile</i>	AF333405	GCAACAAGTT	CGTTCCTCGC	GCTGTCCTCG	
<i>Cylindrocladium gracile</i>	AF333404	GCAACAAGTT	CGTTCCTCGC	GCTGTCCTCG	
<i>Cylindrocladium clavatum</i>	AF232850	GCAACAAGTT	CGTTCCTCGC	GCTGTCCTCG	
<i>Cylindrocladium candelabrum</i>	AF210858	GCAACAAGTT	CGTTCCTCGC	GCTGTCCTCG	
<i>Cylindrocladium candelabrum</i>	AF210857	GCAACAAGTT	CGTTCCTCGC	GCTGTCCTCG	
<i>Cylindrocladium scoparium</i>	AF210875	GCAACAAGTT	CGTTCCTCGC	GCTGTCCTCG	
<i>Cylindrocladium scoparium</i>	AF210874	GCAACAAGTT	CGTTCCTCGC	GCTGTCCTCG	
<i>Cylindrocladium ilicicola</i>	AF333413	GCAACAAGTT	CGTTCCTCGC	GCTGTCCTCG	
<i>Cylindrocladium ilicicola</i>	AF333412	GCAACAAGTT	CGTTCCTCGC	GCTGTCCTCG	
<i>Cylindrocladiella infestans</i>	AF320190	GCAACAAGTT	CGTTCCTCGC	GCTGTCCTCG	
CMW8734		TCGATCTTGA	GCCCCGTACC	ATGGACGCCG	480
CMW9108		TCGATCTTGA	GCCCCGTACC	ATGGACGCCG	
CMW9190		TCGATCTTGA	GCCCCGTACC	ATGGACGCCG	
CMW9109		TCGATCTTGA	GCCCCGTACC	ATGGACGCCG	
CMW9189		TCGATCTTGA	GCCCCGTACC	ATGGACGCCG	
<i>Cylindrocladium pauciramosum</i>	AY162320	TCGATCTTGA	GCCCCGTACC	ATGGACGCCG	
<i>Cylindrocladium pauciramosum</i>	AF449448	TCGATCTTGA	GCCCCGTACC	ATGGACGCCG	
<i>Cylindrocladium insulare</i>	AF44951	TCGATCTTGA	GCCCCGTACC	ATGGATGCCG	
<i>Cylindrocladium insulare</i>	AF44950	TCGATCTTGA	GCCCCGTACC	ATGGATGCCG	
<i>Cylindrocladium colhounii</i>	AF232855	TCGATCTTGA	GCCCCGTACC	ATGGATGCCG	
<i>Cylindrocladium colhounii</i>	AF232854	TCGATCTTGA	GCCCCGTACC	ATGGATGCCG	
<i>Cylindrocladium gracile</i>	AF333405	TCGATCTTGA	GCCCCGTACC	ATGGATGCCG	
<i>Cylindrocladium gracile</i>	AF333404	TCGATCTTGA	GCCCCGTACC	ATGGATGCCG	
<i>Cylindrocladium clavatum</i>	AF232850	TCGATCTTGA	GCCCCGTACC	ATGGATGCCG	
<i>Cylindrocladium candelabrum</i>	AF210858	TCGATCTTGA	GCCCCGTACC	ATGGATGCCG	
<i>Cylindrocladium candelabrum</i>	AF210857	TCGATCTTGA	GCCCCGTACC	ATGGACGCCG	
<i>Cylindrocladium scoparium</i>	AF210875	TCGATCTTGA	GCCCCGTACC	ATGGATGCCG	
<i>Cylindrocladium scoparium</i>	AF210874	TCGATCTTGA	GCCCCGTACC	ATGGATGCCG	
<i>Cylindrocladium ilicicola</i>	AF333413	TCGATCTTGA	GCCCCGTACC	ATGGATGCCG	
<i>Cylindrocladium ilicicola</i>	AF333412	TCGATCTTGA	GCCCCGTACC	ATGGATGCCG	
<i>Cylindrocladiella infestans</i>	AF320190	TCGATCTTGA	GCCCCGTACC	ATGGATGCCG	
CMW8734		TCCGTGCCGG	TCCTT		495
CMW9108		TCCGTGCCGG	TCCTT		
CMW9190		TCCGTGCCGG	TCCTT		
CMW9109		TCCGTGCCGG	TCCTT		
CMW9189		TCCGTGCCGG	TCCTT		
<i>Cylindrocladium pauciramosum</i>	AY162320	TCCGTGCCGG	TCCTT		
<i>Cylindrocladium pauciramosum</i>	AF449448	TCCGTGCCGG	TCCTT		
<i>Cylindrocladium insulare</i>	AF44951	TCCGTGCCGG	TCCTT		
<i>Cylindrocladium insulare</i>	AF44950	TCCGTGCCGG	TCCTT		
<i>Cylindrocladium colhounii</i>	AF232855	TCCGTGCCGG	TCCTT		
<i>Cylindrocladium colhounii</i>	AF232854	TCCGTGCCGG	TCCTT		
<i>Cylindrocladium gracile</i>	AF333405	TCCGTGCCGG	TCCTT		
<i>Cylindrocladium gracile</i>	AF333404	TCCGTGCCGG	TCCTT		
<i>Cylindrocladium clavatum</i>	AF232850	TCCGTGCCGG	TCCTT		
<i>Cylindrocladium candelabrum</i>	AF210858	TCCGTGCCGG	TCCTT		
<i>Cylindrocladium candelabrum</i>	AF210857	TCCGTGCCGG	TCCTT		
<i>Cylindrocladium scoparium</i>	AF210875	TCCGTGCCGG	TCCTT		
<i>Cylindrocladium scoparium</i>	AF210874	TCCGTGCCGG	TCCTT		
<i>Cylindrocladium ilicicola</i>	AF333413	TCCGTGCCGG	TCCTT		
<i>Cylindrocladium ilicicola</i>	AF333412	TCCGTGCCGG	TCCTT		
<i>Cylindrocladiella infestans</i>	AF320190	TCCGTGCCGG	TCCTT		

Chapter 4.

Alignment 6. 5' end of the β - tubulin gene DNA sequence alignment of selected *Cylindrocladium* isolates from *Acacia mearnsii* seedlings and other *Cylindrocladium* species.

CMW 9156		GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	30
CMW 9159		GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
CMW 9164		GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
CMW 9169		GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
CMW 9171		GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>C. pauciramosum</i>	AY162320	GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>C. pauciramosum</i>	AY449448	GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>C. candelabrum</i>	AF210858	GCTT GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>C. candelabrum</i>	AF210857	GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>C. quinqueseptatum</i>	AF232869	GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>C. quinqueseptatum</i>	AF232870	GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>C. theae</i>	AF232861	GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>C. theae</i>	AF232862	GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>C. reteaudii</i>	AF 389846	GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>C. reteaudii</i>	AF389847	GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>C. scoparium</i>	AF210874	GCTT GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>C. scoparium</i>	AF210875	GCT-GCCCCT	GATTCTACCC	CGCCGCCCCG	
<i>Cylindrocladiella infestans</i>	AF320190	GCT-GCCCCT	- ATTCTATCC	- GCCGAATCG	

CMW 9156		GTTTCCACC-	— GCTTCGA	CGACAACAAA	60
CMW 9159		GTTTCCACC-	— GCTTCGA	CGACAACAAA	
CMW 9164		GTTTCCACC-	— GCTTCGA	CGACAACAAA	
CMW 9169		GTTTCCACC-	— GCTTCGA	CGACAACAAA	
CMW 9171		GTTTCCACC-	— GCTTCGA	CGACAACAAA	
<i>C. pauciramosum</i>	AY162320	GTTTCCACC-	— GCTTCGA	CGACAACAAA	
<i>C. pauciramosum</i>	AY449448	GTTTCCACC-	— GCTTCGA	CGACAACAAA	
<i>C. candelabrum</i>	AF210858	GTTTCCACC-	— GCTTCGA	CGACAACAAA	
<i>C. candelabrum</i>	AF210857	GTTTCCACC-	— GCTTCGA	CGACAACAAA	
<i>C. quinqueseptatum</i>	AF232869	GTTTCCACC-	— GCTTCGA	CGACAACAAA	
<i>C. quinqueseptatum</i>	AF232870	GTTTCCACC-	— GCTTCGA	CGACAACAAA	
<i>C. theae</i>	AF232861	GTTTCCACC-	— GCCTCGA	TGGCAGCGAA	
<i>C. theae</i>	AF232862	GTTTCCACC-	— GCCTCGA	TGGCAGCGAA	
<i>C. reteaudii</i>	AF 389846	GTTTCCACC-	— GCTTCGA	CGACAACAAA	
<i>C. reteaudii</i>	AF389847	GTTTCCACC-	— GCTTCGA	CGACAACAAA	
<i>C. scoparium</i>	AF210874	GTTTCCACC-	— ACATCGA	CGAAAACAAA	
<i>C. scoparium</i>	AF210875	GTTTCCACC-	— ACATCGA	CGAAAACAAA	
<i>Cylindrocladiella infestans</i>	AF320190	TTTCCACCC	ACCGCCTCGA	CAACAACAAA	

CMW 9156		GCCGCAGCCT	CACGATCATA	A--CGAGATA	90
CMW 9159		GCCGCAGCCT	CACGATCATA	A--CGAGATA	
CMW 9164		GCCGCAGCCT	CACGATCATA	A--CGAGATA	
CMW 9169		GCCGCAGCCT	CACGATCATA	A--CGAGATA	
CMW 9171		GCCGCAGCCT	CACGATCATA	A--CGAGATA	
<i>C. pauciramosum</i>	AY162320	GCCGCAGCCT	CACGATCATA	A--CGAGATA	
<i>C. pauciramosum</i>	AY449448	GCCGCAGCCT	CACGATCATA	A--CGAGATA	
<i>C. candelabrum</i>	AF210858	GCCGCAGCCT	CACGATCATG	A--CGAGATA	
<i>C. candelabrum</i>	AF210857	GCCGCAGCCT	CACGATCATG	A--CGAGATA	
<i>C. quinqueseptatum</i>	AF232869	GCCGCAACAT	CATGAACAAG	A--CGAGATA	
<i>C. quinqueseptatum</i>	AF232870	GCCGCAACAT	CATGAACAAG	A--CGAGATA	
<i>C. theae</i>	AF232861	GCCGCATCCT	CATGAACAAA	AGACGAGGCA	
<i>C. theae</i>	AF232862	GCCGCATCCT	CATGAACAAA	AGACGAGGCA	
<i>C. reteaudii</i>	AF 389846	GCCGCAACAT	CATGAACAA-	- GACGAGATA	
<i>C. reteaudii</i>	AF389847	GCCGCAACAT	CATGAACAA-	- GACGAGATA	
<i>C. scoparium</i>	AF210874	GCCGCAGCCT	CACGAACAT-	- GATGTGATA	
<i>C. scoparium</i>	AF210875	GCCGCAGCCT	CACGAACAT-	- GATGTGATA	
<i>Cylindrocladiella infestans</i>	AF320190	GCTCGCGATG	CCCACCCACA	T--CGTGATA	



				120
CMW 9156		TCAGAACAAG	ATTGCTAACC	GTGTGCTTCT
CMW 9159		TCAGAACAAG	ATTGCTAACC	GTGTGCTTCT
CMW 9164		TCAGAACAAG	ATTGCTAACC	GTGTGCTTCT
CMW 9169		TCAGAACAAG	ATTGCTAACC	GTGTGCTTCT
CMW 9171		TCAGAACAAG	ATTGCTAACC	GTGTGCTTCT
<i>C. pauciramosum</i>	AY162320	TCAGAACAAG	ATTGCTAACC	GTGTGCTTCT
<i>C. pauciramosum</i>	AY449448	TCAGAACAAG	ATTGCTAACC	GTGTGCTTCT
<i>C. candelabrum</i>	AF210858	TCAGAACAAG	ATTGCTAACC	GTGTGCTTCT
<i>C. candelabrum</i>	AF210857	TCAGAACAAG	ATTGCTAACC	GTGTGCTTCT
<i>C. quinqueseptatum</i>	AF232869	TCAGAACAAG	ATTGCTAACC	GTGTGCTTCT
<i>C. quinqueseptatum</i>	AF232870	TCAGAACAAG	ATTGCTAACC	GTGTGCTTCT
<i>C. theae</i>	AF232861	TCAGAACAAG	ATTGCTAACC	GTGTGCTTCT
<i>C. theae</i>	AF232862	TCAGAACAAG	ATTGCTAACC	GTGTGCTTCT
<i>C. reteaudii</i>	AF 389846	TCAGAACAAG	ATTGCTAACC	GTGTGCTTCT
<i>C. reteaudii</i>	AF389847	TCAGAACAAG	ATTGCTAACC	GTGTGCTTCT
<i>C. scoparium</i>	AF210874	TCAGAACAAG	ATTGCTAACC	GTGTGCTTCT
<i>C. scoparium</i>	AF210875	TCAGAACAAG	ATTGCTAACC	GTGTGCTTCT
<i>Cylindrocladiella infestans</i>	AF320190	TCTGAAGACA	ATTGCTAATT	TTGTGTGTTT

				150
CMW 9156		TTCTCGATTA	TAGGTCCACC	TCCAGACCGG
CMW 9159		TTCTCGATTA	TAGGTCCACC	TCCAGACCGG
CMW 9164		TTCTCGATTA	TAGGTCCACC	TCCAGACCGG
CMW 9169		TTCTCGATTA	TAGGTCCACC	TCCAGACCGG
CMW 9171		TTCTCGATTA	TAGGTCCACC	TCCAGACCGG
<i>C. pauciramosum</i>	AY162320	TTCTCGATTA	TAGGTCCACC	TCCAGACCGG
<i>C. pauciramosum</i>	AY449448	TTCTCGATTA	TAGGTCCACC	TCCAGACCGG
<i>C. candelabrum</i>	AF210858	TTTTCGATTA	TAGGTCCACC	TCCAGACCGG
<i>C. candelabrum</i>	AF210857	TTTTCGATTA	TAGGTCCACC	TCCAGACCGG
<i>C. quinqueseptatum</i>	AF232869	TTCTCGATTA	TAGGTCCACC	TCCAGACCGG
<i>C. quinqueseptatum</i>	AF232870	TTCTCGATTA	TAGGTCCACC	TCCAGACCGG
<i>C. theae</i>	AF232861	CTCTGAATTA	TAGGTCCACC	TCCAGACCGG
<i>C. theae</i>	AF232862	CTCTGAATTA	TAGGTCCACC	TCCAGACCGG
<i>C. reteaudii</i>	AF 389846	TTCTCGATTA	TAGGTCCACC	TCCAGACCGG
<i>C. reteaudii</i>	AF389847	TTCTCGATTA	TAGGTCCACC	TCCAGACCGG
<i>C. scoparium</i>	AF210874	TTCTCGATTA	TAGGTCCACC	TCCAGACCGG
<i>C. scoparium</i>	AF210875	TTCTCGATTA	TAGGTCCACC	TCCAGACCGG
<i>Cylindrocladiella infestans</i>	AF320190	CTG-CGAATA	TAGGTCCACC	TCCAGACCGG

				180
CMW 9156		TCAGTGCGTA	AGTACTCTTC	TCAACTCCAA
CMW 9159		TCAGTGCGTA	AGTACTCTTC	TCAACTCCAA
CMW 9164		TCAGTGCGTA	AGTACTCTTC	TCAACTCCAA
CMW 9169		TCAGTGCGTA	AGTACTCTTC	TCAACTCCAA
CMW 9171		TCAGTGCGTA	AGTACTCTTC	TCAACTCCAA
<i>C. pauciramosum</i>	AY162320	TCAGTGCGTA	AGTACTCTTC	TCAACTCCAA
<i>C. pauciramosum</i>	AY449448	TCAGTGCGTA	AGTACTCTTC	TCAACTCCAA
<i>C. candelabrum</i>	AF210858	TCAGTGCGTA	AGTACTCTTC	TCAACTCCAA
<i>C. candelabrum</i>	AF210857	TCAGTGCGTA	AGTACTCTTC	TCAACTCCAA
<i>C. quinqueseptatum</i>	AF232869	TCAGTGCGTA	AGTACTCTTC	TCAACTCCAA
<i>C. quinqueseptatum</i>	AF232870	TCAGTGCGTA	AGTACTCTTC	TCAACTCCAA
<i>C. theae</i>	AF232861	TCAGTGCGTA	AGTACTCTTC	TCAACTCCAA
<i>C. theae</i>	AF232862	TCAGTGCGTA	AGTACTCTTC	TCAACTCCAA
<i>C. reteaudii</i>	AF 389846	TCAGTGCGTA	AGTACTCTTC	TCAACTCCAA
<i>C. reteaudii</i>	AF389847	TCAGTGCGTA	AGTACTCTTC	TCAACTCCAA
<i>C. scoparium</i>	AF210874	TCAGTGCGTA	AGTACTCTTC	TCAACTCCAA
<i>C. scoparium</i>	AF210875	TCAGTGCGTA	AGTACTCTTC	TCAACTCCAA
<i>Cylindrocladiella infestans</i>	AF320190	TCAGTGCGTA	AGTACTCTTC	TCAACTCCAA



				210
CMW 9156		CAAAATTCTC	ACGACGAGAT	TCACTGACAG
CMW 9159		CAAAATTCTC	ACGACGAGAT	TCACTGACAG
CMW 9164		CAAAATTCTC	ACGACGAGAT	TCACTGACAG
CMW 9169		CAAAATTCTC	ACGACGAGAT	TCACTGACAG
CMW 9171		CAAAATTCTC	ACGACGAGAT	TCACTGACAG
<i>C. pauciramosum</i>	AY162320	CAAAATTCTC	ACGACGAGAT	TCACTGACAG
<i>C. pauciramosum</i>	AY449448	CAAAATTCTC	ACGACGAGAT	TCACTGACAG
<i>C. candelabrum</i>	AF210858	CCAAATTCTC	ACGACGAGAT	TCACTGACAG
<i>C. candelabrum</i>	AF210857	CCAAATTCTC	ACGACGAGAT	TCACTGACAG
<i>C. quinquesseptatum</i>	AF232869	ATATGTTCTC	ATGACAAGAT	TCACTGACAG
<i>C. quinquesseptatum</i>	AF232870	ATATGTTCTC	ATGACAAGAT	TCACTGACAG
<i>C. theae</i>	AF232861	CAAAATTCTC	ACGACGGGAT	TCGCTGACAC
<i>C. theae</i>	AF232862	CAAAATTCTC	ACGACGGGAT	TCGCTGACAC
<i>C. reteaudii</i>	AF 389846	ATATGTTCTC	ATGACAAGAT	TCACTGACAG
<i>C. reteaudii</i>	AF389847	ATATGTTCTC	ATGACAAGAT	TCACTGACAG
<i>C. scoparium</i>	AF210874	CAAAATTCTC	ACGACGGGAT	TCACTGACAG
<i>C. scoparium</i>	AF210875	CAAAATTCTC	ACGACGGGAT	TCACTGACAG
<i>Cylindrocladiella infestans</i>	AF320190	CAAGCTTC--	-- GTCAACGG	CTGCTAACGG

				240
CMW 9156		T- TGTCGATA	GGGTAACCAA	ATTGGTGCTG
CMW 9159		T- TGTCGATA	GGGTAACCAA	ATTGGTGCTG
CMW 9164		T- TGTCGATA	GGGTAACCAA	ATTGGTGCTG
CMW 9169		T- TGTCGATA	GGGTAACCAA	ATTGGTGCTG
CMW 9171		T- TGTCGATA	GGGTAACCAA	ATTGGTGCTG
<i>C. pauciramosum</i>	AY162320	T- TGTCGATA	GGGTAACCAA	ATTGGTGCTG
<i>C. pauciramosum</i>	AY449448	T- TGTCGATA	GGGTAACCAA	ATTGGTGCTG
<i>C. candelabrum</i>	AF210858	T- TGTCGATA	GGGTAACCAA	ATTGGTGCTG
<i>C. candelabrum</i>	AF210857	T- TGTCGATA	GGGTAACCAA	ATTGGTGCTG
<i>C. quinquesseptatum</i>	AF232869	T- TGTCGATA	GGGTAACCAA	ATTGGTGCTG
<i>C. quinquesseptatum</i>	AF232870	T- TGTCGATA	GGGTAACCAA	ATTGGTGCTG
<i>C. theae</i>	AF232861	T- CGCGGATA	GGGTAACCAA	ATCGGTGCTG
<i>C. theae</i>	AF232862	T- CGCGGATA	GGGTAACCAA	ATCGGTGCTG
<i>C. reteaudii</i>	AF 389846	T- TGTCGATA	GGGTAACCAA	ATTGGTGCTG
<i>C. reteaudii</i>	AF389847	T- TGTCGATA	GGGTAACCAA	ATTGGTGCTG
<i>C. scoparium</i>	AF210874	T- TATCGACA	GGGTAACCAA	ATTGGTGCTG
<i>C. scoparium</i>	AF210875	T- TATCGACA	GGGTAACCAA	ATTGGTGCTG
<i>Cylindrocladiella infestans</i>	AF320190	TGTCTCGATA	GGGTAACCAA	ATTGGTGCTG

				270
CMW 9156		CTTTCTGGCA	GACCATTCT	GGCGAGCACG
CMW 9159		CTTTCTGGCA	GACCATTCT	GGCGAGCACG
CMW 9164		CTTTCTGGCA	GACCATTCT	GGCGAGCACG
CMW 9169		CTTTCTGGCA	GACCATTCT	GGCGAGCACG
CMW 9171		CTTTCTGGCA	GACCATTCT	GGCGAGCACG
<i>C. pauciramosum</i>	AY162320	CTTTCTGGCA	GACCATTCT	GGCGAGCACG
<i>C. pauciramosum</i>	AY449448	CTTTCTGGCA	GACCATTCT	GGCGAGCACG
<i>C. candelabrum</i>	AF210858	CTTTCTGGCA	GACCATCTCT	GGCGAGCACG
<i>C. candelabrum</i>	AF210857	CTTTCTGGCA	GACCATCTCT	GGCGAGCACG
<i>C. quinquesseptatum</i>	AF232869	CTTTCTGGCA	GACCATCTCT	GGCGAGCACG
<i>C. quinquesseptatum</i>	AF232870	CTTTTTGGCA	GACCATCTCT	GGCGAGCACG
<i>C. theae</i>	AF232861	CTTTCTGGCA	GACCATTCT	GGCGAGCACG
<i>C. theae</i>	AF232862	CTTTCTGGCA	GACCATTCT	GGCGAGCACG
<i>C. reteaudii</i>	AF 389846	CTTTCTGGCA	GACCATCTCT	GGCGAGCACG
<i>C. reteaudii</i>	AF389847	CTTTTTGGCA	GACCATCTCT	GGCGAGCACG
<i>C. scoparium</i>	AF210874	CTTTCTGGCA	GACCATTCT	GGCGAGCACG
<i>C. scoparium</i>	AF210875	CTTTCTGGCA	GACCATTCT	GGCGAGCACG
<i>Cylindrocladiella infestans</i>	AF320190	CTTTCTGGCA	GACCATCTCT	GGCGAGCACG



				300
CMW 9156		GTCTCGACAG	CAATGGTGTC	TACGCCGGTA
CMW 9159		GTCTCGACAG	CAATGGTGTC	TACGCCGGTA
CMW 9164		GTCTCGACAG	CAATGGTGTC	TACGCCGGTA
CMW 9169		GTCTCGACAG	CAATGGTGTC	TACGCCGGTA
CMW 9171		GTCTCGACAG	CAATGGTGTC	TACGCCGGTA
<i>C. pauciramosum</i>	AY162320	GTCTCGACAG	CAATGGTGTC	TACGCCGGTA
<i>C. pauciramosum</i>	AY449448	GTCTCGACAG	CAATGGTGTC	TACGCCGGTA
<i>C. candelabrum</i>	AF210858	GTCTCGACAG	CAATGGTGTC	TACGCCGGTA
<i>C. candelabrum</i>	AF210857	GTCTCGACAG	CAATGGTGTC	TACGCCGGTA
<i>C. quinquesepatum</i>	AF232869	GTCTCGACAG	CAATGGTGTC	TACGCCGGTA
<i>C. quinquesepatum</i>	AF232870	GTCTCGACAG	CAATGGTGTC	TACGCCGGTA
<i>C. theae</i>	AF232861	GTCTCGACAG	CAATGGTGTC	TACGCCGGTA
<i>C. theae</i>	AF232862	GTCTCGACAG	CAATGGTGTC	TACGCCGGTA
<i>C. reteaudii</i>	AF 389846	GTCTCGACAG	CAATGGTGTC	TACGCCGGTA
<i>C. reteaudii</i>	AF389847	GTCTCGACAG	CAATGGTGTC	TACGCCGGTA
<i>C. scoparium</i>	AF210874	GTCTCGACAG	CAATGGTGTC	TACGCCGGTA
<i>C. scoparium</i>	AF210875	GTCTCGACAG	CAATGGTGTC	TACGCCGGTA
<i>Cylindrocladiella infestans</i>	AF320190	GTCTCGACAG	CAATGGTGTC	TACGCCGGTA

				330
CMW 9156		CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG
CMW 9159		CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG
CMW 9164		CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG
CMW 9169		CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG
CMW 9171		CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG
<i>C. pauciramosum</i>	AY162320	CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG
<i>C. pauciramosum</i>	AY449448	CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG
<i>C. candelabrum</i>	AF210858	CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG
<i>C. candelabrum</i>	AF210857	CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG
<i>C. quinquesepatum</i>	AF232869	CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG
<i>C. quinquesepatum</i>	AF232870	CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG
<i>C. theae</i>	AF232861	CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG
<i>C. theae</i>	AF232862	CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG
<i>C. reteaudii</i>	AF 389846	CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG
<i>C. reteaudii</i>	AF389847	CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG
<i>C. scoparium</i>	AF210874	CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG
<i>C. scoparium</i>	AF210875	CCTCCGAGCT	CCAGCTCGAG	CGTATGAACG
<i>Cylindrocladiella infestans</i>	AF320190	GCTCTGAGCT	CCAGCTCGAG	CGTATGAACG

				360
CMW 9156		TCTACTTCAA	CGAGGTATGT	GAAAACCACT
CMW 9159		TCTACTTCAA	CGAGGTATGT	GAAAACCACT
CMW 9164		TCTACTTCAA	CGAGGTATGT	GAAAACCACT
CMW 9169		TCTACTTCAA	CGAGGTATGT	GAAAACCACT
CMW 9171		TCTACTTCAA	CGAGGTATGT	GAAAACCACT
<i>C. pauciramosum</i>	AY162320	TCTACTTCAA	CGAGGTATGT	GAAAACCACT
<i>C. pauciramosum</i>	AY449448	TCTACTTCAA	CGAGGTATGT	GAAAACCACT
<i>C. candelabrum</i>	AF210858	TCTACTTCAA	CGAGGTATGT	GAAAACCACT
<i>C. candelabrum</i>	AF210857	TCTACTTCAA	CGAGGTATGT	GAAAACCACT
<i>C. quinquesepatum</i>	AF232869	TCTACTTCAA	CGAGGTATGC	GAAAAACCAT
<i>C. quinquesepatum</i>	AF232870	TCTACTTCAA	CGAGGTATGC	GAAAAACCAT
<i>C. theae</i>	AF232861	TCTACTTCAA	CGAGGTATGT	GAAAAAGCA
<i>C. theae</i>	AF232862	TCTACTTCAA	CGAGGTATGT	GAAAAAGCA
<i>C. reteaudii</i>	AF 389846	TCTACTTCAA	CGAGGTATGC	GAAAAACCAT
<i>C. reteaudii</i>	AF389847	TCTACTTCAA	CGAGGTATGC	GAAAAACCAT
<i>C. scoparium</i>	AF210874	TCTACTTCAA	CGAGGTATGT	GAAAA-CCAC
<i>C. scoparium</i>	AF210875	TCTACTTCAA	CGAGGTATGT	GAAAA-CCAC
<i>Cylindrocladiella infestans</i>	AF320190	TCTACTTCAA	CGAGGTACGT	GACTATGGCA



				390
CMW 9156		CGAAGCACTC	CCTT -GACCG	AGAAGCACAA
CMW 9159		CGAAGCACTC	CCTT -GACCG	AGAAGCACAA
CMW 9164		CGAAGCACTC	CCTT -GACCG	AGAAGCACAA
CMW 9169		CGAAGCACTC	CCTT -GACCG	AGAAGCACAA
CMW 9171		CGAAGCACTC	CCTT -GACCG	AGAAGCACAA
<i>C. pauciramosum</i>	AY162320	CGAAGCACTC	CCTT -GACCG	AGAAGCACAA
<i>C. pauciramosum</i>	AY449448	CGAAGCACTC	CCTT -GACCG	AGAAGCACAA
<i>C. candelabrum</i>	AF210858	CGAAGCACTC	CCTT -GACCG	AGAAGCACAA
<i>C. candelabrum</i>	AF210857	CGAAGCACTC	CCTT -GACCG	AGAAGCACAA
<i>C. quinqueseptatum</i>	AF232869	GCCTGCGCTC	GCTTTGTCGA	AAAAGCACAA
<i>C. quinqueseptatum</i>	AF232870	GCCTGCGCTC	GCTTTGTCGA	AAAAGCACAA
<i>C. theae</i>	AF232861	CGCACA -GTT	GTG-AAC- -G	CGAAGGACA-
<i>C. theae</i>	AF232862	CGCACA -GTT	GTGTAAC- -G	CGAAGGACA-
<i>C. reteaudii</i>	AF 389846	GCCTGCGCTC	GCTTTGTCGA	AAAAGCACAA
<i>C. reteaudii</i>	AF389847	GCCTGCGCTC	GCTTTGTCGA	AAAAGCACAA
<i>C. scoparium</i>	AF210874	GCGGTGTACT	CACACG-CCG	AGAGGCACAA
<i>C. scoparium</i>	AF210875	GCGGTGTACT	CACACG-CCG	AGAGGCACAA
<i>Cylindrocladiella infestans</i>	AF320190	CTCA - CATT	GCTA- CACTG	TGAAATCAGA

				420
CMW 9156		GCCAACTCAC	ACCATCATGT	AGGCTTCCGG
CMW 9159		GCCAACTCAC	ACCATCATGT	AGGCTTCCGG
CMW 9164		GCCAACTCAC	ACCATCATGT	AGGCTTCCGG
CMW 9169		GCCAACTCAC	ACCATCATGT	AGGCTTCCGG
CMW 9171		GCCAACTCAC	ACCATCATGT	AGGCTTCCGG
<i>C. pauciramosum</i>	AY162320	GCCAACTCAC	ACCATCATGT	AGGCTTCCGG
<i>C. pauciramosum</i>	AY449448	GCCAACTCAC	ACCATCATGT	AGGCTTCCGG
<i>C. candelabrum</i>	AF210858	TCCGACTCAC	ACCATCATGT	AGGCTTCCGG
<i>C. candelabrum</i>	AF210857	TCCGACTCAC	ACCATCATGT	AGGCTTCCGG
<i>C. quinqueseptatum</i>	AF232869	GCAAAGTAC	AC-ACCATGT	AGGCTTCCGG
<i>C. quinqueseptatum</i>	AF232870	GCAAAGTAC	AC-ACCATGT	AGGCTTCCGG
<i>C. theae</i>	AF232861	GCCAACTCAC	ACCA- --TGT	AGGCTTCCGG
<i>C. theae</i>	AF232862	GCCAACTCAC	ACCA- --TGT	AGGCTTCCGG
<i>C. reteaudii</i>	AF 389846	GCAAAGTAC	AC-ACCATGT	AGGCTTCCGG
<i>C. reteaudii</i>	AF389847	GCAAAGTAC	AC-ACCATGT	AGGCTTCCGG
<i>C. scoparium</i>	AF210874	GCAAAGTAC	ACCAT- --GT	AGGCTTCCGG
<i>C. scoparium</i>	AF210875	GCAAAGTAC	ACCAT- --GT	AGGCTTCCGG
<i>Cylindrocladiella infestans</i>	AF320190	ATGTACTCAC	GC- --TCCGT	AGGCTTCCGG

				450
CMW 9156		CAACAAGTTC	GTTCTCGCG	CTGTCCTCGT
CMW 9159		CAACAAGTTC	GTTCTCGCG	CTGTCCTCGT
CMW 9164		CAACAAGTTC	GTTCTCGCG	CTGTCCTCGT
CMW 9169		CAACAAGTTC	GTTCTCGCG	CTGTCCTCGT
CMW 9171		CAACAAGTTC	GTTCTCGCG	CTGTCCTCGT
<i>C. pauciramosum</i>	AY162320	CAACAAGTTC	GTTCTCGCG	CTGTCCTCGT
<i>C. pauciramosum</i>	AY449448	CAACAAGTTC	GTTCTCGCG	CTGTCCTCGT
<i>C. candelabrum</i>	AF210858	CAACAAGTTC	GTTCTCGCG	CTGTCCTCGT
<i>C. candelabrum</i>	AF210857	CAACAAGTTC	GTTCTCGCG	CTGTCCTCGT
<i>C. quinqueseptatum</i>	AF232869	CAACAAGTTC	GTTCTCGCG	CTGTCCTCGT
<i>C. quinqueseptatum</i>	AF232870	CAACAAGTTC	GTTCTCGCG	CTGTCCTCGT
<i>C. theae</i>	AF232861	CAACAAGTTC	GTTCTCGCG	CTGTCCTCGT
<i>C. theae</i>	AF232862	CAACAAGTTC	GTTCTCGCG	CTGTCCTCGT
<i>C. reteaudii</i>	AF 389846	CAACAAGTTC	GTTCTCGCG	CTGTCCTCGT
<i>C. reteaudii</i>	AF389847	CAACAAGTTC	GTTCTCGCG	CTGTCCTCGT
<i>C. scoparium</i>	AF210874	CAACAAGTTC	GTTCTCGCG	CTGTCCTCGT
<i>C. scoparium</i>	AF210875	CAACAAGTTC	GTTCTCGCG	CTGTCCTCGT
<i>Cylindrocladiella infestans</i>	AF320190	CAACAAGTAT	GTTCTCGCG	CCGTCCTCGT

