

CHAPTER 4

EFFECT OF TEMPERATURE AND SOIL MOISTURE CONTENT ON CUTTING ESTABLISHMENT

4.1 ABSTRACT

Effective rooting is essential for successful crop establishment from cuttings. The objective of this study was to determine the effect of temperature and soil water content on root and shoot growth during the early establishment of sweet potato cuttings. Root growth and development were examined in a phytotron at four temperatures (20, 24, 28 and 32 °C constant) and with four types of cuttings (3 node cutting vertically planted, 3 node cutting planted upside down, 3 node cutting horizontally planted and 1 node cutting planted horizontally). The cuttings were allowed to establish and grow for three weeks. The highest total root length (3.59m per plant) was recorded in the 24 °C growth chamber, significantly longer than the roots in the other temperature treatments. The highest root dry mass (0.22 g per plant), shoot dry mass (1.7 g per plant), leaf area (578 cm²) and total dry mass (2.0 g per plant) were also obtained from the 24 °C growth chamber. The three node cuttings planted vertically produced the longest total root length of 3.66 m per plant, significantly longer than those of the other cutting types. The highest root dry mass, shoot dry mass, leaf area, vine length, leaf number, and total dry mass were also obtained from the vertically planted three node cuttings.

A pot experiment was conducted with cuttings sealed in plastic bags containing sandy soil at 100, 80, 60 and 40% of field water capacity. The cuttings were allowed to establish and grow in a plant growth chamber at 28 °C and harvested after 12 and 20 days. The 80% of field capacity moisture regime was found to be the optimum soil moisture content for sweet potato root growth.

With the soil water content at field capacity and at 40% of field capacity root development was somewhat suppressed. The results illustrated the capacity of sweet potato cuttings to establish successfully under a range of ambient temperatures and soil moisture contents.

4.2 INTRODUCTION

Establishment of sweet potato cuttings can be quite variable depending on environmental conditions. Although sweet potato is grown in the tropical, sub tropical and warm temperate regions of the world, it is essentially a warm weather crop (Onwueme, 1978). The thermal optimum is reported to be above 24 °C, compared to 25-30 °C for cassava and yam (Kay, 1973). Differences in thermal responsiveness would be expected among the wide range of sweet potato genotypes. Being a tropical crop, sweet potato is sensitive to low temperatures. Harter & Whitney (1962) reported that sweet potato could not survive temperatures of less than 12 °C, at 15 °C the plants were able to survive but did not grow, above 15 °C growth increased with increasing temperature up to 35 °C, and at 38 °C growth was somewhat depressed. Sark (1978) reported that sweet potato grown in a 10 to 15 °C greenhouse had much reduced vine growth compared to those grown in temperatures of 21 to 25 °C.

Gomes & Carr (2003) studied the effect of water availability and vine harvesting frequency on the productivity of sweet potato in Mozambique, and suggested that the water requirement over the growing season is between 360 and 800 mm. Sweet potato is normally propagated from vine cuttings, and the development of adventitious roots is expected to be sensitive to soil moisture deficits immediately after planting. An adequate moisture supply is probably essential for promoting rapid and uniform root development and good stand establishment. During vegetative development of plants even minor stresses can reduce the rate of leaf expansion and the leaf area

at later stages of development. Although no published information on the effect of soil moisture content on cutting establishment could be found, various publications refer to the negative effect of water stress on growth and yield of sweet potato. During vegetative development the leaf area index increases with increase in soil moisture (Enyi, 1977; Indira & Ramanujam, 1985; Chowdhury & Ravi, 1988). The storage root initiation period is the most sensitive to moisture deficit due to its effect on storage root number (Indira & Kabeerathumma, 1988; Nair & Nair, 1995; Ravi & Indira, 1996). Moisture deficits during the storage root initiation period induce lignification of adventitious root and hampers growth (Ravi & Indira, 1996).

Considering the scarcity of information on factors affecting early root growth and cutting establishment, two experiments were conducted to establish to what extent root and shoot growth of newly planted cuttings is affected by ambient temperature and soil moisture. The objective of this study was to determine the effect of temperature and soil water content on the root and shoot growth during the establishment of cuttings.

4.3 MATERIALS AND METHODS

Temperature Experiment

A pot experiment with four temperature treatments and four types of cuttings was conducted during 2000 in the phytotron on the Experimental Farm of the University of Pretoria. Leaves were removed from the cuttings before planting. The experiment was carried out in four plant growth chambers. The growth chambers were regulated to constant temperatures of 20 °C, 24 °C,

28 °C, and 32 °C. The photoperiod was 12hr with an abrupt light/dark change. The four types of cuttings were:

- N1. Three-node cuttings vertically planted with one node under the soil surface.
- N2. Three-node cuttings planted upside down with one node under the soil surface.
- N3. Three-node cuttings horizontally planted with all the nodes 5-7 cm under the soil surface.
- N4. One-node cuttings planted horizontally.

Pots were filled with fine sifted, heat-sterilized sand and cuttings were planted on 24 October 2000. There were five replications (pots) of each treatment combination. Within the growth chambers the pots were arranged randomly and watered once a day. All pots were harvested three weeks after planting. The stem length, leaf number and leaf area were determined. The roots were carefully removed from the sand by submerging the pot in water to loosen the sand and minimize root breakage. After uprooting the roots were washed to remove the remaining sand. Images of the roots were obtained by scanning the roots with an image analyzing computer programme. The root length was measured using the “GS Root” programme. Roots and shoots were dried in a forced-ventilation oven at 60 °C for 48 hours. Dry matter partitioning was determined from the total dry mass of the shoots and the roots.

Treatments were arranged in a randomized complete block design. The experimental data were subjected to standard analyses of variance using the General Linear Model (GLM) procedure of the Statistical Analysis System (SAS, 1989) to determine the effect of main factors and the

interaction between them. Differences at the $P \leq 0.05$ level were used as a test of significance and means were separated using Tukey's t-test.

Moisture Experiment

A pot experiment with four-soil water content levels was conducted. The four moisture levels were field capacity (FC), 80% of field capacity, 60% of field capacity, and 40% of field capacity. The four moisture regimes were obtained by adding 70 ml, 56 ml, 42 ml, and 28 ml water respectively to 2 kg dry sand in plastic bags. After adding the water the sand in each bag was thoroughly mixed to achieve an even distribution of moisture, and sealed in polyethylene bags to come to equilibrium. After two days three cuttings of identical fresh mass and length were planted vertically through small slits in each bag on 6 September 2000. After planting the containers were placed in a growth chamber at a constant temperature of 28 °C and a photoperiod of 12 hours.

Two harvests were made, the first harvest was 12 days after planting (12DAP) and the second one 20 days after planting (20DAP). At the first harvest three replicates of each treatment were sampled, while at the second harvest there were four replications. The harvesting procedure, data recorded and statistical procedures were similar to those described for the temperature experiment.

4.4 RESULTS

Temperature experiment

Effect of temperature

Root lengths and root dry mass results are presented in Table 4.1. An increase in temperature from 20 °C to 24 °C increased the root length and root dry mass per plant. The highest total root length of 3.95 m per plant was obtained from plants exposed to a 24 °C growing temperature. As the temperature increased from 24 to 32 °C the root length decreased to 1.62 m per plant. The highest root dry mass was obtained from the plants exposed to 24 °C growing temperature and the lowest from the plants exposed to 20 °C. The largest proportion (13.6%) of the total biomass was partitioned to roots in the 32 °C treatment and the lowest (9.5%) at 28 °C.

The effect of temperature on shoot growth is presented in Table 4.2. In general shoot characteristics did not differ significantly between the 24°C and 28°C temperatures, but exposure to 20° C or 32° C affected shoot growth negatively. The highest vine length of 26.7 cm per plant was obtained from the plants exposed to 28 °C growing temperature, and the shortest vine length of 14.1 cm per plant was obtained from plants exposed to 20 °C. The highest leaf number (14) per plant was obtained from the plants grown at 28 °C, and the lowest temperature of 20 °C produced the smallest number of leaves (7) per plant. The highest shoot dry mass of 1.8 g per plant was obtained from the plants exposed to 24 °C and the lowest shoot dry mass (0.6 g per plant) was from the plants grown at 20 °C. The largest leaf area of 578 cm² was attained by plants grown in the 24 °C growth chamber and the smallest leaf area (188 cm²) was obtained from plants grown at 20 °C.

Effect of type of cutting

Results of the effect of type of cutting on root length, root dry mass and fraction of the mass partitioned to roots are presented in Table 4.1. The highest root length of 3.66 m per plant was obtained from the three node cuttings oriented vertically, and the shortest (2.16 m) from the one node cutting oriented horizontally. The three node cuttings planted vertically produced the highest root dry mass of 0.22 g per plant, and the one node cutting planted horizontally produced the lowest (0.08 g per plant). The three node cuttings planted horizontally partitioned the highest fraction of dry mass (12.3%) to the roots while the three-node cuttings planted upside down partitioned the lowest.

Results of the effect of type of cutting on vine length, leaf number, shoot dry mass production and partitioning and leaf area are presented in Table 4.2. The vine length, leaf number and shoot dry mass from the three node cuttings were similar, whether planted vertically, horizontally or even up side down. The leaf area from the three node cuttings was similar, whether planted horizontally or vertically. It is interesting that even when planted upside down sweet potato cuttings retain the ability to establish vigorous root and shoot growth.

Temperature x type of cutting interactions

The temperature x type of cutting interactions were only significant for root length and leaf area. Vertically oriented three node cuttings exposed to 24 °C growing temperature resulted in a much longer root length (6.4 m) than any of the other treatment combinations (Figure 4.1). The one node cutting produced less root growth (1.2 m) than the other cuttings at 28 °C, but had similar root growth at 20, 24 and 32 °C. In Figure 4.2 the temperature x type of cutting interaction for

leaf area shows that vertically planted three node cuttings had a larger leaf area (877 cm²) at 24 °C than the other cuttings, while at 20°, 28° and 32 °C the leaf areas were similar except for the small leaf area of the one node cuttings at 20 °C. Similarity between Figures 4.1 and 4.2 is interesting and may be an indication of the reliability of the data, but an explanation for the interactions is not obvious.

Moisture experiment

The effect of soil water content 12 days after planting on root development is summarized in Table 4.3. The longest total root length (3.46 m) was obtained from cuttings grown in soils at 80% of field capacity. As the soil water content decreased from 80 to 40% of field capacity the root length decreased to 1.5 m per plant. Differences in shoot and total dry mass per plant and percentage dry matter partitioned to the shoot were significant. The largest proportion (86.5%) of the total biomass was partitioned to stems, the highest shoot dry mass of 0.09 gram per plant, and the highest total dry mass of 0.11 gram per plant were recorded from plants grown at 80% of field capacity, and the lowest at 40% of field capacity.

Results of the effect of soil water content 20 days after planting on root development is summarized in Table 4.4. Differences in root length and root dry mass were not statistically significant, although roots in the 80% of field capacity treatment tended to be the longest. Differences in shoot dry mass per plant were not significant, but the soil moisture regimes showed significant differences in dry matter partitioned to the shoots and total dry mass produced per plant. The largest proportion (90.9%) of the total biomass was partitioned to the

stems, and a highest total dry mass of 0.146 g per plant, was recorded from plants grown at 80% of field capacity.

In this study the 80% of field capacity moisture regime was found to be the optimum soil moisture content for root and shoot development. Lower or higher moisture regimes resulted in less root growth, but it is interesting to note that even at an initial moisture content of 40% of field capacity, substantial root development occurred. This is an indication that soil water content is not critical during the establishment of cuttings, provided that the soil is neither too wet nor too dry.

4.5 DISCUSSION AND CONCLUSION

The increase in root length and root dry mass per plant with increasing temperature from 20 to 24 °C, and a decrease in root length and root dry mass with increase in temperature from 28 to 32 °C suggests that 24 °C was the optimum temperature for root growth and development. The increase in leaf number and vine length per plant with increasing temperature from 20 to 28 °C, and decrease in vine length and leaf number with increase in temperature from 28 to 32 °C, indicate that the optimum temperature for shoot growth was approximately 28 °C. The results suggest that temperature ranging from 24 to 28 °C is the most suitable for early root and shoot growth. According to Spence & Humphries (1971) leaf cuttings planted in sand and exposed to root temperatures of 35, 30, 25, 15, and 10 °C indicated that at 15 °C no roots developed into storage roots but there were more fibrous roots than at 25 and 30 °C.

Despite the relative large coefficient of variation a reasonably clear picture emerged from the experiments regarding the effect of temperature and soil moisture content on early root development. Better root growth was achieved 12 and 20 days after planting from cuttings planted at 80% of field capacity. Lower or higher moisture regimes resulted in less root growth, but it is interesting to note that even at an initial moisture content of 40% of field capacity, substantial root development occurred. This is an indication that soil water content is not critical during the establishment of cuttings, provided that the soil is not too wet (field capacity or wetter) nor too dry (less than 40% of field capacity). Using two sweet potato cultivars Bok (1998) conducted a pot experiment where cuttings were planted in soil at 100%, 70%, 50%, and 30% of field capacity. He reported that the cultivars performed similarly under different soil water regimes, with respect to root length and root dry mass production. The 70% of field capacity moisture regime was found to be the optimum for root development of the cultivars. This result is similar to my data, which indicated that the 80% of field capacity moisture regime was the most favorable for root development. An explanation for the somewhat depressed root growth at field capacity reported by Bok (1998), and observed in this experiment, is not clear. The possibility of partly anaerobic conditions in the polyethylene bags containing the sandy substrate cannot be excluded, and should be investigated.

The results confirm the capacity of sweet potato cuttings to successfully establish under a range of soil moisture conditions and ambient temperatures. This is one of the features of sweet potato cuttings contributing to its suitability as propagating material, even under relatively unfavorable environmental conditions.

Table 4.1 Effect of temperature and orientation of cuttings on sweet potato root development

Treatment		Per plant		
		Root length (m)	Root dry mass (g)	% dry mass partitioned to root
Temperature	20 °C	2.51	0.08	10.9
	24 °C	3.95	0.22	11.0
	28 °C	2.92	0.17	9.5
	32 °C	1.62	0.15	13.6
	LSD _T	0.75	0.05	1.6
Cutting type	N1	3.66	0.22	12.0
	N2	2.32	0.13	9.7
	N3	2.86	0.18	12.3
	N4	2.16	0.08	11.0
	LSD _T	0.75	0.05	1.6
Mean		2.75	0.16	11.3
CV%		43.2	55.0	22.2

Table 4.2 Effect of temperature and orientation of cuttings on sweet potato shoot development

Treatment		Vine length cm	Leaf number	Leaf area cm ²	Shoot dry mass g/plt	Total dry mass g/plt
Temperature	20 °C	14.1	7.2	187.5	0.61	0.7
	24 °C	25.6	13.7	578.2	1.75	2.0
	28 °C	26.7	14.4	519.0	1.50	1.7
	32 °C	20.9	11.7	246.5	1.00	1.1
	LSD _T	3.2	2.3	107.3	0.4	0.4
Cutting type	N1	24.1	13.8	510.4	1.59	1.8
	N2	22.7	12.7	375.8	1.21	1.3
	N3	22.9	12.8	425.3	1.34	1.5
	N4	17.6	7.7	219.7	0.70	0.8
	LSD _T	3.2	2.3	107.3	0.4	0.4
Mean		21.8	11.8	382.8	1.21	1.35
CV%		23.1	30.6	44.3	47.6	47.8




Figure 4.1 Interaction between temperature and orientation of cutting on root length




Figure 4.2 Interaction between temperature and orientation of cutting on leaf area per plant

Table 4.3 Effect of soil water content on the shoot and root growth of sweet potato 12 days after planting

Soil water content (%FC)	Root length in (m)	Root dry mass g/plant	Shoot dry mass g/plant	Total dry mass g/plant	% DM partitioned to shoot
100	2.10	0.011	0.067	0.078	85.47
80	3.46	0.015	0.093	0.108	86.53
60	2.46	0.013	0.050	0.063	78.26
40	1.53	0.010	0.034	0.044	75.35
LSD_T	1.86	0.010	0.044	0.053	10.51
Mean	2.39	0.0123	0.061	0.073	81.40
C.V. (%)	41.37	41.53	38.31	38.10	6.86

Table 4.4 Effect of soil water content on the shoot and root growth of sweet potato 20 days after planting

Soil water content (%FC)	Root length in (m)	Root dry mass g/plant.	Shoot dry mass g/plant	Total dry mass g/plant	%DM partitioned to shoot
100	2.40	0.009	0.086	0.095	90.26
80	3.80	0.012	0.118	0.146	90.92
60	3.21	0.010	0.094	0.103	90.83
40	3.02	0.015	0.083	0.098	85.27
LSD_T	1.66	0.006	0.040	0.039	3.72
Mean	3.11	0.012	0.095	0.111	89.32
C.V. (%)	37.72	32.61	27.24	23.10	2.70

4.6 REFERENCES

- BOK, I., 1998. Response of sweet potato to different soil moisture regimes. Msc. thesis, University of Pretoria, Pretoria, South Africa.
- CHOWDHURY, S.R. & RAVI, V., 1988. Physiology of tuberization in sweet potato with references to moisture stress and seasonal influence. In: *Ann. Rept. Central Tuber Crops Res. Inst., Trivandrum, India*, pp. 89-90.
- ENYI, B.A.C., 1977. Analysis of growth and tuber yield in sweet potato (*Ipomoea batatas*) cultivars. *J. Agr. Sci.* 88, 421-428.
- GOMES, F. & CARR, M. K. V., 2003. Effect of water availability and vine harvesting frequency on the productivity of sweet potato in Southern Mozambique. II. Crop water use. *J. Expl. Agric.* 39, 39-54.
- HARTER, L.L. & WHITNEY, W.A., 1962. Influence of soil temperature and soil moisture on the infection of sweet potato by the black rot fungus. *J. Agric. Res.* 32, 1153.
- INDIRA, P. & KABEERATHUMMA, S., 1988. Physiological response of sweet potato under water stress: 1. Effect of water stress during different phases of tuberization. *J. Root Crops* 14, 37-40.
- INDIRA, P. & RAMANUJAM, T., 1985. Leaf area index, net assimilation rate and crop growth rate of five sweet potato genotypes. (Cited by Ravi, V. & Indira, P., 1999).
- KAY, D.E., 1973. *Root Crops product digest 2: Root crops*. Tropical Product Institute, London.

- NAIR, G.M. & NAIR, V.M., 1995. Influence of irrigation and fertilizers on the growth attributes of sweet potato. *J. Root Crops* 21, 17-23.
- ONWUEME, I. C., 1978. The tropical tuber crops: Yams, Cassava, Sweet potato, and Cocoyams. John Wiley & Sons. New York.
- RAVI, V. & INDIRA, P., 1996. Anatomical studies on tuberization in sweet potato under water deficit stress and stress free conditions. *J. Root Crops* 22, 105-111.
- SARK, E.S.M., 1978. Effect of temperature on yield of sweet potato. *Proc. Am. Soc. Hort. Sci.* 42. 517
- SAS INSTITUTE INC., 1989. SAS User's guide: Statistics. 1989. Cary, North Carolina: SAS Institute.
- SPENCE, J.A. & HUMPHRIES, E.C., 1971. Effect of moisture supply, root temperature and growth regulators on photosynthesis of isolated rooted leaves of sweet potato (*Ipomoea batatas*) *Ann. Bot.* 36, 115-121.