

Phylogeny and Taxonomy of *Calonectria* and its *Cylindrocladium* anamorphs

By

Lorenzo Lombard

Submitted in partial fulfilment of the requirements for the degree

PHILOSOPHIAE DOCTOR

In the Faculty of Natural and Agricultural Science

Department of Microbiology and Plant Pathology

University of Pretoria

Pretoria

April 2010

Supervisor: Prof. M.J. Wingfield

Co-supervisors: Prof. P.W. Crous

Prof. B.D. Wingfield



Declaration

I, the undersigned, hereby declare that the thesis submitted herewith for the degree Philosophiae Doctor to the University of Pretoria contains my own independent work. This work has hitherto not been submitted for any degree at any other University.

Lorenzo Lombard

April 2010

SUMMARY

Species in the genus *Calonectria* (anamorph: *Cylindrocladium*) are euascomycetes in the order *Hypocreales* and are important pathogens of a wide range of plant hosts globally. At the outset, this thesis considers the literature pertaining to species of *Calonectria* and especially the importance of the biological, morphological and phylogenetic species concepts on the taxonomy of this group. It is clear that DNA sequence comparisons have revolutionised the taxonomy of *Calonectria* and literature also highlights the importance of a polyphasic approach to species identification. Studies in this thesis treat a number of forest nursery disease problems caused by *Calonectria* spp. and new species are consequently described based on DNA sequence comparisons, morphological characteristics and sexual compatibility tests. As a consequence several cryptic species were also identified in the genus. Therefore, a multigene genealogy was constructed for all *Calonectria* spp. for which cultures were available and shown to group together in 13 subclades also supported by morphological similarities. As a consequence all *Cylindrocladium* spp. were circumscribed to the genus *Calonectria*, regardless whether the teleomorph state was present or not, based on new nomenclature regulations stated in Article 59.



TABLE OF CONTENTS

Acknowledgements	1
Preface	3
Chapter One: Species concepts in <i>Calonectria</i> (<i>Cylindrocladium</i>)	5
Chapter Two: <i>Calonectria</i> (<i>Cylindrocladium</i>) species associated with dying <i>Pinus</i> cuttings	42
Chapter Three: <i>Calonectria</i> species associated with cutting rot of <i>Eucalyptus</i>	70
Chapter Four: Multigene phylogeny and mating tests reveal three cryptic species related to <i>Calonectria pauciramosa</i>	107
Chapter Five: Phylogeny and systematics of the genus <i>Calonectria</i>	147
Summary	244

ACKNOWLEDGEMENTS

I would like to express my sincere gratitude and appreciation to the following people and institutions that have assisted me in preparing this thesis:

My supervisors: Prof. M. J. Wingfield, Prof. P. W. Crous and Prof. B. D. Wingfield, for their advice, guidance and patience and without whom this thesis would not have been completed.

Dr. H. F. Glen for providing the Latin descriptions and helping with assigning names to the new *Calonectria* spp.

Prof. P. W. Crous and the Evolutionary Phytopathology group for giving me the opportunity to conduct part of my study at the Centraalbureau voor Schimmelcultures (CBS), Utrecht, The Netherlands.

Staff and students at CBS, for their help and advice during my nine month stay there.

My colleagues and friends in FABI, especially those in the Denison lab.

The administrative staff in FABI, especially Rose Visser, Helen Doman, Eva Muller and Jenny Hale, for their assistance with administrative issues.

The culture collection staff at FABI and CBS, who were always willing to help with culture issues.

My closest friends, Lotte van Belzen, Gavin Hunter, Lawrie Wright, Almuth Hammerbacher, Cecile Gueidan, Tinieke van Doorn, Ewald and Marizeth Groenewald, for their support, advice and encouragement during the last few years of study.

My parents, Lorenzo Z. C. Lombard and Lee E. Lombard, for all their love, support and never-ending encouragement as well as financial support during my studies.

My brothers and sister, Vincent Lombard, Riaan Lombard, Claudie Brand and extended families for always being there for me.

The University of Pretoria (UP), National Research Foundation (NRF), members of the Tree Protection Co-operative Programme (TPCP), Centre of Excellence in Tree



Health Biotechnology and CBS for financial support that allowed me to conduct and complete this study.

PREFACE

Fungal species in the genus *Calonectria* (anamorph *Cylindrocladium*) are important plant pathogens of a wide variety of hosts particularly in tropical and sub-tropical regions of the world. For most of the taxonomic history of *Calonectria* spp., their identification has depended deeply on the morphology of the anamorph state, most frequently encountered in nature, and more recently the sexual compatibility of isolates. Advances in molecular technology and DNA sequencing have, however, resulted in the recognition of several species complexes that include numerous cryptic species.

Several studies focussing on the taxonomy of selected groups of *Calonectria* spp., have recently incorporated DNA sequence comparisons, morphology and sexual compatibility to identify cryptic species in these groups. Application of the Phylogenetic, Morphological and Biological Species Concepts has revolutionised the taxonomy of *Calonectria*. The first chapter of this thesis critically reviews and analyses published research on the application of these three species concepts as they pertain to the taxonomic history of *Calonectria* and places this research undertaken in this thesis in context.

The experimental part of this thesis is based on isolates collected over a period of five years from various geographical regions on plant hosts and in soil. In Chapter two, the mortality of rooted *Pinus* cuttings in a commercial forest nursery in Colombia was investigated. The *Calonectria* spp. responsible for plant deaths were identified using DNA sequence and morphological comparisons. Furthermore, pathogenicity of these *Calonectria* spp. was tested on *P. maximinoi* and *P. tecunumanii* in greenhouse studies.

Eucalyptus hybrid clones have emerged as the most widely planted forestry trees in China. The result has been large-scale uniform plantations to accommodate a high-demand for wood and wood-products. To support these *Eucalyptus* plantations, it is necessary to raise and maintain healthy plants in nurseries. Decline in *Eucalyptus* cutting production in a forest nursery in GuangDong Province, China, has been linked to cutting rot associated with several *Calonectria* spp. In Chapter three of this thesis, these *Calonectria* spp. were identified using DNA sequence and morphological

comparisons. Species in the *Ca. reteaudii* complex were re-considered and shown to accommodate two cryptic species.

Calonectria pauciramosa belongs to the *Ca. scoparia* complex and is a well-known plant pathogen of various hosts, worldwide. In some countries, it is regarded as the dominant pathogen in commercial forest nurseries. Several studies, in the past, have suggested that it accommodates cryptic species. These studies, however, included small numbers of isolates. The aim of chapter four was to identify these cryptic taxa in *Ca. pauciramosa*. This was accomplished using a large number of isolates subjected to DNA sequence comparisons, morphological observations and sexual compatibility tests.

Over a period of five years a large number of *Calonectria* isolates were collected from infected plant material or baited from soils. These isolates were obtained from many different geographical regions of the world. In Chapter 5, I identify these *Calonectria* isolates based on DNA sequence comparisons, morphological characteristics and sexual compatibility. This resulted in the identification of numerous new taxa in *Calonectria*. Subsequently, a multigene phylogeny was constructed for all *Calonectria* spp. for which cultures are available and particularly to determine the phylogenetic relationships within *Calonectria*.



CHAPTER 1

Species concepts in *Calonectria* (*Cylindrocladium*)

A Literature Review

1.0 INTRODUCTION

The genus *Calonectria* (*Ca.*) was erected in 1867 by De Notaris, based on *Ca. daldiniana* collected on leaves of *Magnolia grandiflora* (*Magnoliaceae*), in Daldini, Italy (Rossman 1979a). Rossman (1979a) later reduced *Ca. daldiniana* to synonymy under *Ca. pyrochroa*, and defined this nectrioid fungus as having an ascocarp wall structure that is brightly coloured, changing to blood-red in 3% KOH solution, warty to scaly and with a *Cylindrocladium* (*Cy.*) anamorph (Rossman 1993, Rossman *et al.* 1999). However, due to the restricted morphological characteristics of the teleomorph (Rossman 1979b, 1983), specimens can in many cases only be identified to species level if the anamorph is present (Schoch *et al.* 2000b, Crous 2002).

The anamorph genus *Cylindrocladium*, which is based on *Cy. scoparium*, was first described by Morgan (1892) in the U.S.A., where it was found growing as saprobe on a pod of *Gleditsia triacanthos*. Although Morgan (1892) failed to mention the stipe extension terminating in a vesicle of characteristic shape, he defined the genus as having branched conidiophores producing cylindrical conidia. This fungus has a wide distribution in sub-tropical and tropical regions of the world, and species are pathogenic to numerous plants (Crous 2002).

The aim of this review is to present an overview of published research on the genus *Calonectria*. More specifically, the application of three types of species concepts is considered as they pertain to the taxonomic history of this genus up to 2008. Although several species concepts (Mayden 1997) have been proposed, only the Morphological Species Concept (MSC), the Biological Species Concept (BSC) and the Phylogenetic Species Concept (PSC) are treated, as these have been most widely applied to *Calonectria*. Several reviews (Rossman 1996, Brasier 1997, Harrington & Rizzo 1999, Taylor *et al.* 1999, 2000, Seifert *et al.* 2000; Kohn 2005) have treated the various species concepts applied to the taxonomy of fungi and this topic is not treated other than in the manner in which it applies to *Calonectria*.

2.0 TAXONOMIC HISTORY

Calonectria resides in the Nectriaceae, one of three families in Hypocreales, an order that has been reviewed extensively (Rogerson 1970, Rossman 1983, Rossman *et al.* 1996, 1999). The Nectriaceae is circumscribed as having uniloculate ascomata that are orange to purple and not immersed in well-developed stromata (Rossman *et al.* 1999). The family includes approximately 20 genera of socio-economic importance and of these, *Calonectria* are more clearly distinguished from the others by their *Cylindrocladium* anamorphs and relevance as plant pathogens.

The first monograph of *Cylindrocladium*, by Boedjin & Reitsma (1950), introduced seven *Cylindrocladium* species with a *Calonectria* connection to one of these species. Later, in her treatment of *Calonectria*, Rossman (1983) recognized five species including the novel *Ca. ophiospora*. However, this species description did not include the anamorph state. The circumscribed type, *Ca. pyrochoa*, was also incorrectly reduced to synonymy with several other species based only on the teleomorph morphology. Peerally (1991a) highlighted this in a monograph of *Cylindrocladium*, where he regarded the anamorph morphology as important in distinguishing species of *Calonectria*. He subsequently recognized 10 *Calonectria* species with their *Cylindrocladium* anamorphs, including an additional 16 *Cylindrocladium* species not associated with a teleomorph. However, he mistakenly reduced *Cylindrocladiella*, a genus that accommodates *Cylindrocladium*-like species with small conidia (Boesewinkel 1982), to synonymy with *Cylindrocladium*.

The monograph of *Cylindrocladium* by Crous & Wingfield (1994) entrenched the importance of anamorph characteristics in the taxonomy of *Calonectria* spp. In this monograph, 22 *Cylindrocladium* species and one variety were recognised, associated with 16 *Calonectria* species. Five species were assigned to the genus *Cylindrocladiella* based on morphological characters of the holomorph. The focus on anamorph characteristics is perpetuated in the most recent monograph (Crous 2002), which recognized 28 *Calonectria* species, all associated with *Cylindrocladium* anamorphs and an additional 18 *Cylindrocladium* species for which teleomorph states were not known. Of the latter group, seven taxa were of doubtful authenticity.

Presently, 32 *Calonectria* and 52 *Cylindrocladium* species are recognized (Table 1; Crous 2002, Crous *et al.* 2004b, 2006, Gadgil & Dick 2004).

A general search on MycoBank (www.mycobank.org; Crous *et al.* 2004a, Roberts *et al.* 2005) and Index Fungorum (www.indexfungorum.org) resulted in a total of 256 and 258 name records respectively for *Calonectria*. A similar search for *Cylindrocladium* species on both electronic databases indicated a total of 103 and 94 names respectively.

3.0 NOMENCLATURE OF CALONECTRIA

The nomenclature of pleomorphic fungi has been a topic of substantial debate during the course of the past two decades (Gams 1991, Cannon & Kirk 2000, Hawksworth 2004, 2005). The separate naming of anamorphs (mitotic morphs) and teleomorphs (meiotic morphs) has resulted in confusion, especially for non-taxonomists (Cannon & Kirk 2000). This is especially evident where teleomorph species epithets are different to those of their anamorphs and also where more than one anamorph (synanamorph) is found. The naming of fungal morphs based on the International Code of Botanical Nomenclature (ICBN; McNeill *et al.* 2005) and in particular following strict interpretation of Article 59 of the Code has now been unsatisfactory for many fungal groups due to our ability to connect morphs using molecular evidence, and there are increasing calls for further changes to be made.

According to Article 59.4, the teleomorph name takes precedence over the anamorph name when both types belong to the same holomorph taxon. Further, the earliest available legitimate name typified (Article 59.1) should be regarded as the correct name after 1 January 2008 (Hawksworth 2004). Following these rules, the name *Calonectria* typified in 1867, takes precedence over *Cylindrocladium* typified in 1892 (Morgan 1892). Although there are several *Cylindrocladium* species without *Calonectria* connections (Crous 2002, Crous *et al.* 2004b, 2006), we believe that new species should be described in *Calonectria* irrespective of whether a teleomorph is known or not. This follows a clear view based on phylogenetic inference that *Cylindrocladium* spp. all are derived from the same common ancestor as the *Calonectria* spp. (Schoch *et al.* 1999, 2000a, 2000b, Crous 2002, Crous *et al.* 2004b,

2006). Thus, for taxonomic purposes, *Cylindrocladium* species with known teleomorph states are referred to as *Calonectria* in this review.

4.0 IMPORTANCE OF CALONECTRIA

The genus *Calonectria* was initially regarded as a saprobe as no disease symptoms could be induced by inoculating a suspected host (Graves 1915). The first proof of pathogenicity of these fungi was provided by Massey (1917), and subsequently by Anderson (1919), who proved pathogenicity of *Ca. morganii*. Subsequently, *Calonectria* species have been associated with a wide range of disease symptoms on a large number of hosts worldwide (Crous 2002; Table 2). In the past, several authors have indicated that *Calonectria* species cause disease on plants residing in approximately 30 plant families (Booth & Gibson 1973, French & Menge 1978, Peerally 1991a, Wiapara *et al.* 1996, Schoch *et al.* 1999). Upon closer inspection, the number of plant host families is actually closer to 100 (Table 2) that include approximately 335 plant species (Crous 2002). These hosts include important forestry, agricultural and horticultural crops. This suggests that the impact of these plant pathogens has been underestimated in the past.

The majority of disease reports associated with *Calonectria* species in forestry include hosts in 5 plant families, of which the most important are associated with Fabaceae (*Acacia* spp.), Myrtaceae (*Eucalyptus* spp.) and Pinaceae (*Pinus* spp.). Disease symptoms include cutting rot (Crous *et al.* 1991, Crous 2002), damping-off (Batista 1951, Cox 1953, Terashita & Itô 1956, Sharma & Mohanan 1982, Sharma *et al.* 1984, Crous *et al.* 1991, Brown & Ferreira 2000, Crous 2002, Taniguchi *et al.* 2008) leaf diseases (Cox 1953, Hodges & May 1972, Barnard 1984, Sharma *et al.* 1984, El-Gholl *et al.* 1986, Peerally *et al.* 1991a, Crous *et al.* 1993b, Crous & Wingfield 1994, Crous *et al.* 1998b, Schoch & Crous 1999, Schoch *et al.* 1999, Booth *et al.* 2000, Park *et al.* 2000, Crous & Kang 2001, Gadgil & Dick 2004), shoot blight (Sharma *et al.* 1984, Crous *et al.* 1991, Crous *et al.* 1998b, Crous & Kang 2001), stem cankers (Cox 1953, Sharma *et al.* 1984, 1985, Crous *et al.* 1991) and root rot (Cox 1953, Hodges & May 1972, Cordell & Skilling 1975, Mohanan & Sharma 1985, Crous *et al.* 1991). The majority of these diseases are associated with seedling and cutting production in forestry nurseries, but in a few cases *Cylindrocladium* species have also been reported

from commercial plantations. In these cases the pathogens have been reported to cause leaf diseases and shoot blight resulting in defoliation of trees leading to loss of growth vigour (Hodges & May 1972, Sharma *et al.* 1985, Booth *et al.* 2000, Park *et al.* 2000, Crous & Kang 2001, Crous 2002, Old *et al.* 2003, Rodas *et al.* 2005).

In agriculture, *Calonectria* species have been reported to cause diseases on several economically important crops. Several plant families of agricultural importance are susceptible to *Calonectria* infections, of which the most significant fall in Fabaceae, and Solanaceae. Important diseases in these families include *Cylindrocladium* black rot of *Arachis hypogea* (peanut) and red crown rot of *Glycine max* (soybean) caused by *Ca. ilicicola* and *Ca. pyrochroa* in the USA (Bell & Sobers 1966, Beute & Rowe 1973, Rowe *et al.* 1973, Sobers & Littrell 1974, Rowe & Beute 1975, Phipps *et al.* 1976, Johnson 1985, Dianese *et al.* 1986, Berner *et al.* 1988, Berner *et al.* 1991, Culbreath *et al.* 1991, Porter *et al.* 1991, de Varon 1991, Hollowell *et al.* 1998, Kim *et al.* 1998) and *Cylindrocladium* tuber rot of *Solanum tuberosum* (potato) (Boedjin & Reitsma 1950, Bolkan *et al.* 1980, 1981) by *Cy. gracile* in Brazil. Other diseases associated with *Calonectria* species on agricultural crops include root rot and leaf diseases of fruit bearing and spice plants (Jauch 1943, Wormald 1944, Sobers & Seymour 1967, Nishijima & Aragaki 1973, Milholland 1974, Krausz & Caldwell 1987, Hutton & Sanewski 1989, Anandaraj & Sarma 1992, Risède 1994, Jayasinghe & Wijesundera 1996, Risède & Simoneau 2001, Vitale & Polizzi 2008), post-harvest diseases of fruits (Fawcett & Klotz 1937, Boedjin & Reitsma 1950, Sepiah 1990, Fitzell & Peak 1992, Vaidya & Roa 1992, Sivapalan *et al.* 1998), root and crown rot of *Medicago sativa* (alfalfa) (Ooka & Uchida 1982, Hwang & Flores 1987), and sheath net blotch of *Oryza sativa* (rice) (Crous 2002).

On horticultural crops, *Calonectria* species have been reported mostly from the Northern Hemisphere, especially in gardens and ornamental commercial nurseries in Europe and Asia (Polizzi & Crous 1999, Polizzi 2000, Crous 2002, Henricot & Culham 2002, Pérez-Sierra *et al.* 2007, Polizzi *et al.* 2007a, 2007b, Hirooka *et al.* 2008). Hosts in this sector include ornamental trees, shrubs and cut-flowers in several plant families, most commonly in Arecaceae, Asteraceae, Ericaceae and Rosaceae. A wide range of disease symptoms are recorded including crown-, collar- and root rot, leaf spots, and cutting rot (Massey 1917, Anderson 1919, Aragaki *et al.* 1972, 1988,

Peerally 1991b, Uchida & Kadooka 1997, Polizzi & Crous 1999, Polizzi 2000, Crous 2002, Henricot & Culham 2002, Henricot & Beales 2003, Poltronieri *et al.* 2004, Lane *et al.* 2006, Polizzi *et al.* 2006a, 2006b, 2007a, 2007b, Pérez-Sierra *et al.* 2007, Vitale & Polizzi 2007, Hirooka *et al.* 2008, Vitale *et al.* 2008).

5.0 MORPHOLOGY

Morphological or phenotypic characters have played a major role in the description of fungal species (Brasier 1997, Taylor *et al.* 2000) and form the basis of new fungal descriptions as required by the ICBN (McNeill *et al.* 2005). In recent years, the use of morphological characters alone to delimit new species has been set aside, to a large extent, with more focus being placed on biological and phylogenetic characters (Rossman 1996, Brasier 1997, Taylor *et al.* 2000). This trend is also evident in recent studies on *Calonectria* species (Crous *et al.* 2004b, 2006).

The morphology of *Calonectria* and to a greater extent its anamorph, *Cylindrocladium*, has been important in the taxonomic history of these fungi. Prior to the 1990's, identification of species was based on morphological characteristics and to a lesser extent on sexual compatibility using standardised media (Boedjin & Reitsma 1950, Peerally 1991a, Crous *et al.* 1992, Crous & Wingfield 1994, Crous 2002). This resulted in the establishment of several species complexes, as many *Cylindrocladium* species are morphologically very similar. These include the *Ca. scoparia* complex (Schoch *et al.* 1999), *Cy. gracile* complex (Crous *et al.* 2004b) and *Ca. kyotensis* complex (Crous *et al.* 2006). Characteristics (Fig. 1) of the anamorphs that are extensively employed in identifications include vesicle shape (Fig. 1C–F), stipe extension length (Fig. 1A–B) and macroconidial septation and dimensions (Fig. 1G–J; Boesewinkel 1982, Peerally 1991a, Crous & Wingfield 1994, Crous 2002). The morphological characteristics of the teleomorph that are important for identifications are ascospore (Fig. 1M–N) septation and dimensions. The perithecia of *Calonectria* species are morphologically very similar and these are typically not very useful in identifications (Crous & Wingfield 1994, Crous 2002).

Biochemical techniques can also be used in phenotypic characterization. These include substrate utilization and cell wall polysaccharide analysis. The use of aminopeptidase specificity (Stevens *et al.* 1990) and utilization of specific nitrogen

and carbon (Hunter & Barnett 1978, Sharma *et al.* 1992) have been used successfully to separate several *Cylindrocladium* species. The use of polysaccharides obtained from cell walls of *Cylindrocladium* positively identified linkages between asexual species and their respective *Calonectria* teleomorphs (Ahrazem *et al.* 1997). However, this method has been found to have a limited value as it does not distinguish between some species in complexes (Crous 2002).

6.0 MATING COMPATIBILITY

Mating strategies have been employed in the taxonomy of *Calonectria* and have played an important role in identifying new species of the genus (Schoch *et al.* 1999, Crous 2002). Based on these studies, there are approximately 18 homothallic and 34 heterothallic species of *Calonectria* (Crous 2002, Crous *et al.* 2004b, Gadgil & Dick 2004, Crous *et al.* 2006), with the heterothallic species showing a biallelic mating system (Schoch *et al.* 1999). Studies in the female fertility of *Cylindrocladium* by Schoch *et al.* (1999, 2000a, 2001a) have also shown that several species are self-sterile hermaphrodites requiring fertilization from an opposite mating type. This is typical of heterothallic ascomycetes (Leslie & Klein 1996).

Several difficulties associated with applying the BSC have been highlighted (Brasier 1997, Taylor *et al.* 1999, Taylor *et al.* 2000, Kohn 2005). The most relevant underlying problem occurs where genetically isolated fungal strains retain the ancestral ability to recombine to produce viable progeny (Brasier 1997). This phenomenon has also been found with several phylogenetic species that are closely related in *Calonectria*. Crous (2002), for example, showed that *Cy. hawksworthii*, *Ca. insularis* and *Ca. morganii* were capable of recombining, but that the progeny had low levels of fertility. Other mating studies done by Overmeyer *et al.* (1996) and Neubauer & Zinkernagel (1995) have found that induction of fertile perithecia requires the presence of an additional isolate that, however, does not contribute to the genetic make-up of the progeny. This clearly highlights the need for further studies regarding the mechanism of perithecial formation and recombination in *Calonectria*.

7.0 PHYLOGENY

Phylogenetic studies on *Calonectria*, and its *Cylindrocladium* anamorphs have substantially influenced the taxonomy of these genera. Application of molecular techniques and particularly DNA sequence comparisons to distinguish between species has resulted in the recognition of numerous cryptic species. Several molecular approaches have been employed that include total protein electrophoresis (Crous *et al.* 1993a, El-Gholl *et al.* 1993a), isozyme electrophoresis (El-Gholl *et al.* 1992, 1997, Crous *et al.* 1998a), random amplification of polymorphic DNA (RAPD) (Overmeyer *et al.* 1996, Victor *et al.* 1997, Schoch *et al.* 2000a, Riséde & Simoneau 2004) restriction fragment length polymorphisms (RFLP) (Crous *et al.* 1993b, Crous *et al.* 1995, Crous *et al.* 1997b, Jeng *et al.* 1997, Victor *et al.* 1997; Riséde & Simoneau 2001) and DNA hybridization (Crous *et al.* 1993b, 1995, 1997a, Victor *et al.* 1997). Although the above-mentioned techniques have been useful, DNA sequence comparisons and associated phylogenetic inference have had the most dramatic impact on the taxonomy of *Calonectria* and are most widely applied today.

In the first study using 5.8S ribosomal RNA gene and flanking internally transcribed spacers (ITS) sequences Jeng *et al.* (1997) were able to distinguish between *C. scoparium* and *C. floridanum* isolates. Subsequently, it was found that this gene region contains few informative characters (Crous *et al.* 1999, Schoch *et al.* 1999, Riséde & Simoneau 2001, Schoch *et al.* 2001b). Therefore, the β -tubulin (Schoch *et al.* 2001b) and histone H3 (Kang *et al.* 2001a) gene regions have been applied in order to allow for improved resolution in separating species.

The first complete DNA sequence-based phylogenetic study using partial β -tubulin gene sequences (Schoch *et al.* 2001b) compared phenotypic, biological and phylogenetic concepts used in the taxonomy of *Cylindrocladium*. This also highlighted the fact that *Calonectria* represents a monophyletic lineage (Schoch *et al.* 2000b, 2001b). Subsequently, combined DNA sequence data for the ITS, β -tubulin and histone H3 gene regions have been widely used in studies relating to taxonomic issues surrounding *Cylindrocladium* and *Calonectria* (Crous *et al.* 1999, Schoch *et al.* 2000a, 2000b, Crous & Kang 2001, Kang *et al.* 2001a, 2001b, Henricot & Culham 2002, Crous *et al.* 2004b, 2006). Other partial gene sequences recently used include

translation elongation 1-alpha (TEF-1 α) and calmodulin (Crous *et al.* 2004b). However, insufficient data are currently available for these gene regions on GenBank (www.ncbi.nlm.nih.gov) to make them particularly valuable for comparative analysis.

In a search of GenBank, a total of 734 partial gene sequences was obtained for *Calonectria* and *Cylindrocladium*. These include 311 for β -tubulin, 177 for histone H3, 159 for ITS, 39 for calmodulin, 36 for TEF-1 α , five for large subunit RNA gene (LSU), three each for the high mobility group (HMG) box and peptidase synthetase and one for the small subunit RNA (SSU) gene. For *Cylindrocladium* and *Calonectria*, there are only four studies (Kang *et al.* 2001a, 2001b; Crous *et al.* 2004b, 2006) that provide files on TreeBase (www.treebase.org).

8.0 FUTURE RESEARCH

8.1 Population Biology

Most studies on *Calonectria* have focused on the taxonomy, phylogeny and pathology of species. There have in contrast been relatively few studies treating the population biology of these fungi. This is unfortunate as population dynamics contributes considerable knowledge to a better understanding of population structure, distribution of genetic diversity, gene flow, centres of origin and mating strategies (McDonald 1997, Linde *et al.* 2002, Grünwald *et al.* 2003). An understanding of the population dynamics of *Calonectria* would contribute in determining the natural spread of these fungi as well as assist in phytosanitary and quarantine regulations. Another important aspect surrounding knowledge of *Calonectria* population dynamics is that this would contribute to plant breeding programmes and thus control of the many diseases that are caused by these fungi (McDonald 1997, Wright *et al.* 2006, 2007).

Limited research has been conducted on the population dynamics of *Calonectria*. To date only two studies (Wright *et al.* 2006, 2007) have reported on the development of polymorphic markers to characterise simple sequence repeats (SSRs) in loci of *Ca. ilicicola* (Wright *et al.* 2006) and *Ca. pauciramosa* (Wright *et al.* 2007). However, no study has yet been published on the population biology of either of these important pathogens using these markers. There is clearly a gap in this area of research concerning *Calonectria* spp. and future research in this area should be encouraged.

8.2 Whole genome sequences

A relatively new and innovative technology employed in fungal genetics is the use of whole genome sequences of filamentous fungi. Whole genome sequencing has become relatively inexpensive and thus common in recent years. This revolutionary technology will promote our understanding of the mechanisms of gene function, conidiation, pathogenesis and sexual reproduction at the genotype level (Kupfer *et al.* 1997, Prade 1998, Yoder & Turgeon 2001, Foster *et al.* 2006, Cuomo *et al.* 2007). It is estimated that most filamentous fungi have a genome size of 30 to 40 Mb, containing approximately 8000 to 9000 genes (Kupfer *et al.* 1997, Prade 1998, Foster *et al.* 2006). There are currently several completed fungal genome sequences (<http://www.broad.mit.edu/annotation/fungi/fgi/>, Foster *et al.* 2006, Baker *et al.* 2008) that include the model yeast *Saccharomyces cerevisiae* (Goffeau *et al.* 1996), plant pathogens and spoilage fungi such as *Aspergillus flavus* (Payne *et al.* 2006), *Fusarium graminearum* (<http://www.broad.mit.edu>, Cuomo *et al.* 2007), *Magnaporthe grisea* (Dean *et al.* 2005) and the model filamentous fungus *Neurospora crassa* (Galagan *et al.* 2003). Although there are currently over 300 ongoing filamentous fungal genome sequencing projects (<http://www.genomesonline.org>, Baker *et al.* 2008, Liolios *et al.* 2008), none include species of *Calonectria*.

The most closely related plant pathogen to *Calonectria* species currently being sequenced is *Haematonectria haematococca* (<http://www.ncbi.nlm.gov>). When the first *Calonectria* species is selected for whole genome sequencing, comparisons with *H. haematococca* could help to identify some important genes in pathogenesis and sexual reproduction. Some *Calonectria* species that could be considered for genome sequencing include *Ca. pauciramosa*, based on its pathogenicity and importance on several plant hosts worldwide (Crous 2002), and *Ca. reteaudii*, one of the most important forest pathogens of South East Asia (Booth *et al.* 2000, Old *et al.* 2003).

9.0 CONCLUSIONS

Early studies on the taxonomy of *Calonectria* and *Cylindrocladium* focused on the use of MSC in combination with BSC. More recently, the wide availability of molecular techniques and particularly DNA sequence data have revolutionised the taxonomy of *Calonectria* and *Cylindrocladium*. Today, it is well accepted that the

morphology of the *Cylindrocladium* state contributes most information to naming species and that these fungi all reside in *Calonectria*.

The first study to combine MSC, BSC and PSC concepts by Schoch *et al.* (1999) resulted in the identification of four species within a single species complex. Subsequently, several studies including the MSC, BSC and PSC have elucidated cryptic species in the genus (Kang *et al.* 2001a, 2001b; Henricot & Culham 2002; Crous *et al.* 2004b, 2006). Application of the BSC in the taxonomy of *Calonectria* has been found to be unreliable in some instances (Crous 2002). However, the implementation of MSC and PSC in combination provides powerful tool for taxonomic studies of these genera and it is likely that this will continue in future studies. Although several species complexes have been identified in *Calonectria*, more research is needed on the population level in order to study the gene flow between populations. Additional to this, more gene regions need to be identified and widely used in PSC. With the identification of several new species since 2002, an updated monograph is required to facilitate ease of identification.

LITERATURE SITED

- Ahrazem O, Prieto A, Leal JA, Gomez-Miranda B, Domenech J, Jimenez-Barbero J, Bernabe M. (1997). Structural elucidation of acidic fungal polysaccharides isolated from the cell-wall of genera *Cylindrocladium* and *Calonectria*. *Carbohydrate Research* **303**: 67–72.
- Alfieri SA, Linderman RG, Morrison RH, Sobers EK. (1972). Comparative pathogenicity of *Calonectria theae* and *Cylindrocladium scoparium* to leaves and roots of azalea. *Phytopathology* **62**: 647–650.
- Anandaraj M, Sarma YR. (1992). A new leaf rot in *Pimenta dioica*. *Indian Phytopathology* **45**: 276–277.
- Anderson PJ. (1919). Rose canker and its control. *Massachusetts Agricultural Experiment Station Bulletin* **183**: 11–46.
- Aragaki M, Laemmlen FF, Nishijima WT. (1972). Collar rot of Koa caused by *Calonectria crotalariae*. *Plant Disease Reporter* **56**: 73–74.
- Aragaki M, Yahata PS, Uchida JY. (1988). Heliconia root rot caused by *Cylindrocladium spathiphylli* f. sp. *heliconiae*. *Phytopathology* **78**: 1614.
- Baker SE, Thykaer J, Adney WS, Brettin TS, Brockman FJ, D’Haeseleer P, Martinez AD, Miller RM, Rokhsar DS, Schadt CW, Torok T, Tuskan G, Bennett J, Berka RM, Briggs SP, Heitman J, Taylor J, Turgeon BG, Werner-Washburne M, Himmel ME. (2008). Fungal genome sequencing and bioenergy. *Fungal Biology Reviews* **22**: 1–5.
- Barnard EL. (1984). Occurrence, impact and fungicide control of girdling stem cankers caused by *Cylindrocladium scoparium* on *Eucalyptus* seedlings in a south Florida nursery. *Plant Disease* **68**: 471–473.
- Batista AC. (1951). *Cylindrocladium scoparium* Morgan var. *brasiliensis* Batista & Ciferri, a new fungus on *Eucalyptus*. *Boletim da Secretaria de Agricultura, Industria e Comercio do Estado de Pernambuco* **18**: 188–191.
- Bell DK, Sobers EK. (1966). A peg, pod and root necrosis of peanuts caused by a species of *Calonectria*. *Phytopathology* **56**: 1361–1364.
- Berner DK, Berggren GT, Snow JP, White EP. (1988). Distribution and management of red crown rot of soybean in Louisiana, U.S.A. *Applied Agricultural Research* **3**: 160–166.

- Berner DK, Berggren GT, Snow JP. (1991). Effects of glyphosate on *Calonectria crotalariae* and red crown rot of soybean. *Plant Disease* **75**: 809–813.
- Beute MK, Rowe RC. (1973). Studies on the biology and control of *Cylindrocladium* black rot (CBR) of peanut. *Journal of the American Peanut Research Educational Associations* **5**: 197.
- Boedjin KB, Reitsma J. (1950). Notes on the genus *Cylindrocladium*. *Reinwardtia* **1**: 51–60.
- Boesewinkel HJ. (1982). Heterogeneity within *Cylindrocladium* and its teleomorphs. *Transactions of the British Mycological Society* **78**: 553–556.
- Bolkan HA, Dianese JC, Ribeiro WRC, Almeida OC de. (1980). Disease caused by *Cylindrocladium* on potato tubers in Brazil. *Plant Disease* **64**: 225.
- Bolkan HA, Ribeiro WRC, Almeida OC de. (1981). Pathogenicity of *Cylindrocladium clavatum* causing potato tuber rot. *Plant Disease* **65**: 47–49.
- Booth C. (1966). The genus *Cylindrocarpon*. *Mycological Papers* **104**: 1–56.
- Booth C, Gibson IAS. (1973). *Cylindrocladium scoparium*. *CMI Descriptions of Pathogenic Fungi and Bacteria No. 362*.
- Booth TH, Jovanovic T, Old KM, Dubzinski MJ. (2000). Climatic mapping to identify high-risk areas for *Cylindrocladium quinquesseptatum* leaf blight on eucalypts in mainland South East Asia and around the world. *Environmental Pollution* **108**: 365–372.
- Booth C, Murray JS. (1960). *Calonectria hederæ* Arnaud and its *Cylindrocladium* conidial state. *Transactions of the British Mycological Society* **43**: 69–72.
- Brasier CM. (1997). Fungal species in practice: identifying species units in fungi. In: *Species: The units of Biodiversity* (Claridge MF, Dawah HA, Wilson MR, eds). Chapman & Hall, U.K.: 135–170.
- Brown BB, Ferreira FA. (2000). Disease during propagation of eucalypts. In: *Diseases and pathogens of eucalypts* (Keane PJ, Kile GA, Podger FD, Brown BN, eds.). CSIRO publishing, Australia: 119–151.
- Cannon PF, Kirk PM. (2000). The philosophy and practicalities of amalgamating anamorph and teleomorph concepts. *Studies in Mycology* **45**: 19–25.
- Cordell CE, Skilling DD. (1975). Forest nursery diseases in the U.S.A. 7. *Cylindrocladium* root rot. *U.S.D.A. Forest Service Agricultural Handbook No. 470* : 23–26.

- Cox RS. (1953). Etiology and control of a serious complex of diseases of conifer seedlings. *Phytopathology* **43**: 469.
- Crous PW. (2002) Taxonomy and pathology of *Cylindrocladium* (*Calonectria*) and allied genera. APS Press, St. Paul, Minnesota, U.S.A.
- Crous PW, Alfenas AC, Junghans TG. (1998a). Variability within *Calonectria ovate* and its anamorph *Cylindrocladium ovatum* from Brazil. *Sydowia* **50**: 1–13.
- Crous PW, Alfenas AC, Wingfield MJ. (1993a). *Calonectria scoparia* and *Calonectria morganii* sp. nov., and variation among isolates of their *Cylindrocladium* anamorphs. *Mycological Research* **97**: 701–708.
- Crous, PW, Gams W, Staplers JA, Roberts V, Stegehuis G. (2004a). MycoBank: an online initiative to launch mycology into the 21st century. *Studies in Mycology* **50**: 19–22.
- Crous PW, Groenewald JZ, Hill CF. (2002) *Cylindrocladium pseudonaviculatum* sp. nov. from New Zealand, and new *Cylindrocladium* records from Vietnam. *Sydowia* **54**: 23–33.
- Crous PW, Groenewald JZ, Risède J-M, Simoneau P, Hyde KD. (2006). *Calonectria* species and their *Cylindrocladium* anamorphs: species with clavate vesicles. *Studies in Mycology* **55**: 213–226.
- Crous PW, Groenewald JZ, Risède J-M, Simoneau P, Hywel-Jones NL. (2004b). *Calonectria* species and their *Cylindrocladium* anamorphs: species with sphaeropedunculate vesicles. *Studies in Mycology* **50**: 415–430.
- Crous PW, Janse BJH, Victor D, Marais GF, Alfenas AC. (1993b). Molecular characterization of *Cylindrocladium* spp. with three-septate conidia and ovoid-like vesicles. *Systematic and Applied Microbiology* **16**: 266–273.
- Crous PW, Kang JC. (2001). Phylogenetic confirmation of *Calonectria spathulata* and *Cylindrocladium leucothoes* based on morphology, and sequence data of the β -tubulin and ITS rRNA genes. *Mycoscience* **42**: 51–57.
- Crous PW, Kang JC, Schoch CL, Mchau GRA. (1999). Phylogenetic relationships of *Cylindrocladium pseudogracile* and *Cylindrocladium rumohrae* with morphologically similar taxa, based on morphology and DNA sequences of internal transcribed spacers and β -tubulin. *Canadian Journal of Botany* **77**: 1813–1820.

- Crous PW, Krof A, Van Zyl WH. (1995). Nuclear DNA polymorphisms of *Cylindrocladium* species with 1-septate conidia and clavate vesicles. *Systematic and Applied Microbiology* **18**: 224–250.
- Crous PW, Mchau GRA, Van Zyl WH, Wingfield MJ. (1997a). New species of *Calonectria* and *Cylindrocladium* isolated from soil in the tropics. *Mycologia* **89**: 653–660.
- Crous PW, Phillips AJL, Wingfield MJ. (1991). The genera *Cylindrocladium* and *Cylindrocladiella* in South Africa, with special reference to forestry nurseries. *South African Forestry Journal* **157**: 69–85.
- Crous PW, Phillips AJL, Wingfield MJ. (1992). Effects of cultural conditions on vesicle and conidium morphology in species of *Cylindrocladium* and *Cylindrocladiella*. *Mycologia* **84**: 497–504.
- Crous PW, Schoch CL, El-Gholl NE, Schubert TS, Leahy RM. (2000). *Cylindrocladium angustatum* sp. nov., a new leaf spot pathogen of *Tillandsia capitata* from Florida U.S.A. *Mycosience* **41**: 521–526.
- Crous PW, Theron L, Van Zyl WH. (1997b). Delineation of *Cylindrocladium* species with 1–3-septate conidia and clavate vesicles based on morphology and rDNA RFLPs. *Mycological Research* **101**: 210 – 214.
- Crous PW, Wingfield MJ. (1994). A monograph of *Cylindrocladium*, including anamorphs of *Calonectria*. *Mycotaxon* **51**: 341–435.
- Crous PW, Wingfield MJ, Alfenas AC. (1993e). Additions to *Calonectria*. *Mycotaxon* **46**: 217–234.
- Crous PW, Wingfield MJ, Alfenas AC. (1993d). *Cylindrocladium parasiticum* sp. nov., a new name for *C. crotalariae*. *Mycological Research* **97**: 889–896.
- Crous PW, Wingfield MJ, Alfenas AC, Silveira SF. (1994). *Cylindrocladium naviculatum* sp. nov., and two new vesiculate hyphomycete genera, *Falcocladium* and *Vesiculomyces*. *Mycotaxon* **50**: 441–458.
- Crous PW, Wingfield MJ, Mohammed C, Yuan ZQ. (1998b). New foliar pathogens of *Eucalyptus* from Australia and Indonesia. *Mycological Research* **102**: 527–532.
- Culbreath AK, Beute MK, Campbell CL. (1991). Spatial and temporal aspects of epidemics of *Cylindrocladium* black rot in resistant and susceptible peanut genotypes. *Phytopathology* **81**: 144–150.

- Cuomo CA, Güldener U, Xu J-R, Trial F, Turgeon BG, Di Pietro A, Walton JD, Ma J-L, Baker SE, Rep M, Adam G, Antoniw J, Baldwin T, Calvo S, Chang Y-L, DeCaprio D, Gale LR, Gnerre S, Goswami RS, Hammond-Kosack K, Harris LJ, Hilburn K, Kennell JC, Kroken S, Magnuson JK, Mannhaupt G, Mauceli E, Mewes H-W, Mitterbauer R, Muelbauer G, Münsterkötter M, Nelson D, O'Donnell K, Ouellet T, Qi W, Quesneville H, Roncero MIG, Seong K-Y, Tetko IV, Urban M, Waalwijk C, Ward TJ, Yao J, Birren BW, Kistler HC. (2007). The *Fusarium graminearum* genome reveals a link between localized polymorphism and pathogen specialization. *Science* **317**: 1400–1402.
- Dean RA, Talbot NJ, Ebbole DJ, Farman ML, Mitchell TK, Orbach MJ, Thon M, Kulkarni R, Xu J-R, Pan H, Read ND, Lee Y-H, Carbone I, Brown D, Oh YY, Donofrio N, Jeong JS, Soanes DM, Djonovic S, Kolomiets E, Rehmeyer C, Li W, Harding M, Kim S, Leburn M-H, Bohnert H, Coughlan S, Butler J, Calvo S, Ma L-J, Nicol R, Purcell S, Nusbaum C, Galagan JE, Birren BW. (2005). The genome sequence of the rice blast fungus *Magnaporthe grisea*. *Nature* **434**: 980–986.
- Dianese JC, Ribeiro WRC, Urban AF. (1986). Root rot of soybean caused by *Cylindrocladium clavatum* in central Brazil. *Plant Disease* **70**: 977–980.
- El-Gholl NE, Alfenas AC, Crous PW, Schubert TS. (1993a). Description and pathogenicity of *Cylindrocladium ovatum* sp. nov. *Canadian Journal of Botany* **71**: 466–470.
- El-Gholl NE, Alfenas AC, Junghans, DT, Schubert TS, Miller JW, Leahy EM. (1997). Description of *Calonectria rumohrae* sp. nov. (anamorph = *Cylindrocladium rumohrae* sp. nov.) *Mycotaxon* **64**: 467–484.
- El-Gholl NE, Alfieri SA, Barnard EL. (1993b). Description and pathogenicity of *Calonectria clavata* sp. nov. *Mycotaxon* **48**: 201–216.
- El-Gholl NE, Kimbrough JW, Barnard EL, Alfieri SA, Schouties CL. (1986). *Calonectria spathulata* sp. nov. *Mycotaxon* **26**: 151–164.
- El-Gholl NE, Leahy RM, Schubert TS. (1989). *Cylindrocladium leucothoeae* sp. nov. *Canadian Journal of Botany* **67**: 2529–2532.
- El-Gholl NE, Uchida JY, Alfenas AC, Schubert TS, Alfieri SA, Chase AR. (1992). Induction and description of perithecia of *Calonectria spathiphylli* sp. nov. *Mycotaxon* **45**: 285–300.

- Fawcett HS, Klotz LJ. (1937). A new species of *Candelospora* causing decay of citrus fruit. *Mycologia* **29**: 207–215.
- Fitzell RD, Peak CM. (1992). Field evaluation of benomyl to control *Cylindrocladium* fruit spot of custard apple. *Australasian Plant Pathology* **21**: 16–17.
- Foster SJ, Monahan BJ, Bradshaw RE. (2006). Genomics of the filamentous fungi – moving from the shadow of the bakers yeast. *Mycologist* **20**: 10–14.
- French DW, Menge JA. (1978). Survival of *Cylindrocladium floridanum* in naturally and artificially infested forest tree nurseries. *Plant Disease Reporter* **62**: 806–810.
- Gadgil PD, Dick MA. (2004). Fungi silvicolae novazelandiae: 5. *New Zealand Journal of Forestry Science* **34**: 316–323.
- Galagan JE, Calvo SE, Borkovich KA, Selker EU, Read, ND, Jaffe D, Fitzhugh W, Ma L-J, Smirnov S, Purcell S, Rehman B, Elkins T, Engels R, Wang A, Nielsen CB, Butler J, Endrizzi M, Qui D, Ianakiev P, Bell-Pedersen D, Nelson MA, Werner-Washburne M, Selitrennikoff CP, Kinsey JA, Braun JA, Zelter A, Schulte U, Kothe GO, Jedd G, Mewes W, Staben C, Marcotte E, Greenberg D, Roy A, Foley K, Naylor J, Stange-Thomann N, Barrett R, Gnerre S, Kamal M, Kamvysselis M, Mauceli E, Bielke C, Rudd S, Frishman D, Krystofova S, Rasmussen C, Metzenberg RL, Perkins DD, Kroken S, Cogoni C, Macino G, Catcheside D, Li W, Pratt RJ, Osmani SA, DeSouza CPC, Glass L, Orbach MJ, Berglund JA, Voelker R, Yarden O, Plamann M, Seiler S, Dunlap J, Radford A, Aramayo R, Natvig DO, Alex LA, Mannhaupt G, Ebbole DJ, Freitag M, Paulsen I, Sachs MS, Lander ES, Nusbaum C, Birren B. (2003). The genome sequence of the filamentous fungus *Neurospora crassa*. *Nature* **422**: 859–868.
- Gams W. (1991). What are names in current use? *Mycotaxon* **40**: 319–322.
- Gill DL, Alfieri SA, Sobers EK. (1971). A new leaf disease of *Ilex* spp. caused by *Cylindrocladium avesciculatum* sp. nov. *Phytopathology* **61**: 58–60.
- Goffeau A, Barrell BG, Bussey H, Davis RW, Dujon B, Feldmann H, Galibert F, Hoheisel JD, Jacq C, Johnston M, Louis EJ, Mewes HW, Murakami Y, Philippsen P, Tettelin H, Oliver SG. (1996). Life with 6000 genes. *Science* **274**: 546–567.
- Graves AH. (1915). Root rot of coniferous seedlings. *Phytopathology* **5**: 213–217.
- Grünwald NJ, Goodwinj SB, Milgroom MG, Fry WE. (2003). Analysis of genotypic diversity data for populations of microorganisms. *Phytopathology* **93**: 738–746.

- Harrington TC, Rizzo DM. (1999). Defining species in the fungi. In *Structure and Dynamics of Fungal Populations* (Worrall JJ, ed.). Kluwer Academic, Dordrecht: 43–70
- Hawksworth DL. (2004). Limitation of dual nomenclature for pleomorphic fungi. *Taxon* **53**: 596–598.
- Hawksworth DL. (2005). Two major changes in fungal nomenclature enacted in Vienna. *Mycological Research* **109**: 1061–1062.
- Henricot B, Beales P. (2003). First record of *Cylindrocladium pauciramosum* on myrtle (*Myrtus communis*) in Portugal. *Plant Pathology* **52**: 420.
- Henricot B, Culham A. (2002). *Cylindrocladium buxicola*, a new species affecting *Buxus* spp., and its phylogenetic status. *Mycologia* **94**: 980–997.
- Hirooka Y, Takeuchi J, Horie H, Natsuaki KT. (2008). *Cylindrocladium* brown leaf spot on *Howea belmoreana* caused by *Calonectria ilicicola* (anamorph: *Cylindrocladium parasiticum*) in Japan. *Journal of General Plant Pathology* **74**: 66–70.
- Hodges CS, May LC. (1972). A root disease of pine, *Araucaria*, and *Eucalyptus* in Brazil caused by a new species of *Cylindrocladium*. *Phytopathology* **62**: 898–901.
- Hollowell JE, Shew BB, Beute MK, Abad ZG. (1998). Occurrence of pod rot pathogens in peanuts grown in North Carolina. *Plant Disease* **82**: 1345–1349.
- Hunter BB, Barnett HL. (1978). Growth and sporulation of species and isolates of *Cylindrocladium* in culture. *Mycologia* **70**: 614–635.
- Hutton DG, Sanewski GM. (1989). *Cylindrocladium* leaf and fruit spot of custard apple in Queensland. *Australasian Plant Pathology* **18**: 15–16.
- Hwang SF, Flores G. (1987). Effects of *Cylindrocladium gracile*, *Fusarium roseum* and *Plenodomus meliloti* on crown and root rot, foliage yield and winterkill of alfalfa in north-eastern Alberta. *Canadian Plant Disease Survey* **67**: 31–33.
- Jauch C. (1943). The presence of *Cylindrocladium scoparium* in Argentina. *Revista Argentina de Agronomia* **10**: 355–360.
- Jayasinghe CK, Wijesundera RLC. (1996). Morphological, cultural and pathogenic variation among the clove isolates of *Cylindrocladium quinquesepatum*. *Journal of Plantation Crops* **24**: 34–42.

- Jeng RS, Dumas M, Liu FH, Wang CL, Hubbes M. (1997). DNA analysis of *Cylindrocladium floridanum* isolates from selected forest nurseries. *Mycological Research* **101**: 285–291.
- Johnson GI. (1985). Occurrence of *Cylindrocladium crotalariae* on peanut (*Arachis hypogaea*) seed. *Plant Disease* **69**: 434–436.
- Kang JC, Crous PW, Old KM, Dubzinski MJ. (2001a). Non-conspecificity of *Cylindrocladium quinqueseptatum* and *Calonectria quinqueseptata* based on a β -tubulin gene phylogeny and morphology. *Canadian Journal of Botany* **79**: 1241–1247.
- Kang JC, Crous PW, Schoch CL. (2001b). Species concepts in the *Cylindrocladium floridanum* and *Cy. spathiphylli* complexes (*Hypocreaceae*) based on multi-allelic sequence data, sexual compatibility and morphology. *Systematic Applied Microbiology* **24**: 206–217.
- Kim KD, Russin JS, Snow JP. (1998). Susceptibility to *Calonectria ilicicola* in soybean grown in greenhouse and field. *Korean Journal of Crop Science* **43**: 239–244.
- Kohn LM. (2005). Mechanisms of fungal speciation. *Annual Review of Phytopathology* **43**: 279–308.
- Krausz JP, Caldwell JD. (1987). *Cylindrocladium* root rot of kiwifruit. *Plant Disease* **71**: 374–375.
- Kupfer DM, Reece CA, Clifton SW, Roe BA, Prade RA. (1997). Multicellular ascomycetous fungal genomes contain more than 8000 genes. *Fungal Genetics and Biology* **21**: 364–372.
- Lane CR, Beales PA, Henricot B, Holden A. (2006). First record of *Cylindrocladium pauciramosum* on *Ceanothus* in the UK. *Plant Pathology* **55**: 582.
- Leahy RM, Schubert TS, El-Gholl NE. (2000). *Cylindrocladium gordoniae* sp. nov. *Mycotaxon* **76**: 77–83.
- Leslie JF, Klein KK. (1996). Female fertility and mating type effects on effective population size and evolution in filamentous fungi. *Genetics* **144**: 557–567.
- Liolios K, Mavromatis K, Tavernarakis N, Kyrpides NC. (2008). The Genome On Line Database (GOLD) in 2007: status of genomic and metagenomic projects and their associated metadata. *Nucleic Acids Research* **36**: D475–D479.

- Linde CC, Zhan J, McDonald BA. (2002). Population structure of *Mycosphaerella graminicola*: from lesions to continents. *Phytopathology* **92**: 946–955.
- Massey LM. (1917). The crown canker disease of rose. *Phytopathology* **7**: 408–417.
- Mayden RL. (1997). A hierarchy of species concepts: the denouement in the saga of the species problem. In: *Species: The units of Biodiversity* (Claridge MF, Dawah HA, Wilson MR, eds). Chapman & Hall, UK.: 381–424.
- McDonald BA. (1997). The population genetics of fungi: tools and techniques. *Phytopathology* **87**: 448–453.
- McNeill J, Stuessy TF, Turland NJ, Hörandl E. (2005). XVII International Botanical Congress: preliminary mail vote and report of Congress action on nomenclature proposals. *Taxon* **54**: 1057–1064.
- Milholland RD. (1974). Stem and root rot of blueberry caused by *Calonectria crotalariae*. *Phytopathology* **64**: 831–834.
- Mohan C, Sharma JK. (1985). *Cylindrocladium* causing seedling diseases of *Eucalyptus* in Kerala, India. *Transactions of the British Mycological Society* **84**: 538–539.
- Morgan AP. (1892). Two new genera of hyphomycetes. *Botanical Gazette* **17**: 190–192.
- Neubauer C, Zinkernagel V. (1995). *Calonectria morganii* (Crous, Alfenas and Wingfield), the sexual stage of *Cylindrocladium scoparium* Morgan. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* **102**: 323–325.
- Nishijima WT, Aragaki M. (1973). Pathogenicity and further characterization of *Calonectria crotalariae* causing collar rot of papaya. *Phytopathology* **63**: 553–558.
- Old KM, Wingfield MJ, Yuan ZQ. (2003). A manual of diseases of eucalypts in South-East Asia. Center for International Forestry Research, Jakarta, Indonesia. Pp. 98.
- Ooka JJ, Uchida JY. (1982). *Cylindrocladium* root and crown rot of alfalfa in Hawaii. *Plant Disease* **66**: 947–948.
- Overmeyer C, Lünneberg S, Von Wallburgn C, Meinhardt F. (1996). Genetic variability among isolates and sexual offspring of the plant pathogenic fungus *Calonectria morganii* on the basis of random amplification of polymorphic DNA (RAPD) and restriction fragment length polymorphism (RFLP). *Current Microbiology* **33**: 249–255.

- Park RF, Keane PJ, Wingfield MJ, Crous PW. (2000). Fungal diseases of eucalypt foliage. In: *Diseases and pathogens of eucalypts*. (Keane PJ, Kile GA, Podger FD, Brown BN, eds.). CSIRO publishing, Australia: 153 – 239.
- Payne GA, Nierman WC, Wortman JR, Pritchard BL, Brown D, Dean RA, Bhatnagar D, Cleveland TE, Machida M, Yu J. (2006). Whole genome comparison of *Aspergillus flavus* and *A. oryzae*. *Medical Mycology* **44**: S9–S11.
- Peerally A. (1973). *Calonectria colhounii* sp. nov., a common parasite of tea in Mauritius. *Transactions of the British Mycological Society* **61**: 89–93.
- Peerally A. (1991a). The classification and phytopathology of *Cylindrocladium* species. *Mycotaxon* **40**: 367–366.
- Peerally A. (1991b). *Cylindrocladium hawksworthii* sp. nov. pathogenic to water-lilies in Mauritius. *Mycotaxon* **40**: 367–376.
- Pérez-Sierra A, Alvarez LA, Henricot B, Garcia-Jimenez J, Armengol J. (2006). *Cylindrocladium pauciramosum* causes root and collar rot of *Polygala myrtifolia* in Spain. *Plant Pathology* **55**: 298.
- Pérez-Sierra A, Alvarez LA, Leon M, Abad-Campos P, Armengol J, Garcia-Jimenez J. (2007). First report of leaf spot, blight and stem lesions caused by *Cylindrocladium pauciramosum* on *Callistemon* in Spain. *Plant Disease* **91**: 1057.
- Phipps PM, Beute MK, Barker KR. (1976). An elutriation method for quantitative isolation of *Cylindrocladium crotalariae* microsclerotia from peanut field soil. *Phytopathology* **66**: 1255–1259.
- Polizzi G. (2000). Prime esperienze di lotta chimica nei confronti del marciume del colletto e delle radici di *Polygala myrtifolia* causato da *Cylindrocladium pauciramosum*. *Informatore Fitopatologico* **11**: 39–47.
- Polizzi G, Crous PW. (1999). Root and collar of milkwort caused by *Cylindrocladium pauciramosum*, a new record for Europe. *European Journal of Plant Pathology* **105**: 407–411.
- Polizzi G, Grasso FM, Vitale A, Aiello D. (2007b). First occurrence of *Calonectria* leaf spot on mexican blue palm in Italy. *Plant Disease* **91**: 1057.
- Polizzi G, Vitale A, Aiello D, Dimartino MA, Parlavecchio G. (2007a). First report of damping-off and leaf spot caused by *Cylindrocladium scoparium* on different accessions of bottlebrush cuttings in Italy. *Plant Disease* **91**: 769.

- Polizzi G, Vitale A, Aiello D, Parlavecchio G. (2006a). First record of crown and root rot caused by *Cylindrocladium pauciramosum* on California lilac in Italy. *Plant Disease* **90**: 1459.
- Polizzi G, Vitale A, Castello I, Groenewald JZ, Crous PW. (2006b). *Cylindrocladium* leaf spot, blight and crown rot, new diseases of mastic tree seedlings caused by *Cylindrocladium scoparium*. *Plant Disease* **90**: 1110.
- Poltronieri LS, Silva JF da, Alfenas AC, Zauza EAV, Trindade DR. (2004). *Eugenia brachypoda*, new host of *Cylindrocladium pteridis* in the State of Pará, Brazil. *Fitopatologia Brasileira* **29**: 102–103.
- Porter DM, Wright FS, Taber RA, Smith DH. (1991). Colonization of peanut seed by *Cylindrocladium crotalariae*. *Phytopathology* **81**: 896–900.
- Prade RA. (1998). Fungal genomics – one per week. *Fungal Genetics and Biology* **25**: 76–78.
- Riséde JM. (1994). Partial characterization of *Cylindrocladium* sp., a root pathogen of banana in Martinique. *Fruits (Paris)* **49**: 167–178.
- Riséde JM, Simoneau P. (2001). Typing *Cylindrocladium* species by analysis of ribosomal DNA spacers polymorphism: application to field isolates from the banana rhizosphere. *Mycologia* **93**: 494–504.
- Riséde JM & Simoneau P. (2004). Pathogenic and genetic diversity of soilborne isolates of *Cylindrocladium* from banana cropping systems. *European Journal of Plant Pathology* **110**: 139–154.
- Roberts V, Stegehuis G, Stalpers J. (2005). The MycoBank engine and related databases. <http://www.mycobank.org>
- Rodas CA, Lombard L, Gryzenhout M, Slippers B, Wingfield MJ. (2005). *Cylindrocladium* blight of *Eucalyptus grandis* in Colombia. *Australasian Plant Pathology* **34**: 134–149.
- Rogerson CT. (1970). The Hypocrealean fungi (Ascomycetes, Hypocreales). *Mycologia* **62**: 865–910.
- Rossmann AY. (1979a). *Calonectria* and its type species, *C. daldiniana*, a later synonym of *C. pyrochroa*. *Mycotaxon* **8**: 321–328.
- Rossmann AY. (1979b). A preliminary account of the taxa described in *Calonectria*. *Mycotaxon* **8**: 485–558.

- Rossmann AY. (1983). The phragmosporous species of *Nectria* and related genera. *Mycological Papers* **150**: 1–164.
- Rossmann AY. (1993). Holomorphic hypocrealean fungi: *Nectria sensu stricto* and telemorphs of *Fusarium*. In: *The fungal holomorph: mitotic, meiotic and pleomorphic speciation in fungal systematics*. (Reynolds DR, Taylor JW, eds.). CAB International, Wallingford, U.K.: 149 – 160.
- Rossmann AY. (1996). Morphological and molecular perspectives on systematics of the Hypocreales. *Mycologia* **88**: 1–19.
- Rossmann AY, Samuels GJ, Rogerson CT, Lowen R. (1999). Genera of Bionectriaceae, Hypocreaceae and Nectriaceae (Hypocreales, Ascomycetes). *Studies in Mycology* **42**: 1–248.
- Rowe RC, Beute MK. (1975). Variability in virulence of *Cylindrocladium crotalariae* isolates on peanut. *Phytopathology* **65**: 422–425.
- Rowe RC, Beute MK, Wells JC. (1973). *Cylindrocladium* black rot of peanuts in North Carolina – 1972. *Plant Disease Reporter* **57**: 387–389.
- Schoch CL, Crous PW. (1999). First report of *Cylindrocladium* root and petiol rot on *Spathiphyllum* in South Africa. *South African Journal of Botany* **65**: 67–72.
- Schoch CL, Crous PW, Polizzi G, Koike ST. (2001a). Female fertility and single nucleotide polymorphism comparisons in *Cylindrocladium pauciramosum*. *Plant Disease* **85**: 941–946.
- Schoch CL, Crous PW, Wingfield BD, Wingfield MJ. (1999). The *Cylindrocladium candelabrum* species complex includes four distinct mating populations. *Mycologia* **91**: 286–298.
- Schoch CL, Crous PW, Wingfield BD, Wingfield MJ. (2001b). Phylogeny of *Calonectria* based on comparisons of β -tubulin DNA sequences. *Mycological Research* **105**: 1045–1052.
- Schoch CL, Crous PW, Cronwright G, Witthuhn RC, El-Gholl NE, Wingfield BD. (2000a). Recombination in *Calonectria morganii* and phylogeny with other heterothallic small-spored *Calonectria* species. *Mycologia* **92**: 665–673.
- Schoch CL, Crous PW, Wingfield MJ, Wingfield BD. (2000b). Phylogeny of *Calonectria* and selected hypocrealean genera with cylindrical macroconidia. *Studies in Mycology* **45**: 45–62.

- Schoulties CL, El-Gholl NE, Alfieri SA. (1982). *Cylindrocladium spathiphylli* sp. nov. *Mycotaxon* **16**: 265–272.
- Schubert TS, El-Gholl NE, Alfieri SA, Schoulties CL. (1989). *Calonectria avesiculata* sp. nov. *Canadian Journal of Botany* **67**: 2414–2419.
- Seifert KA, Gams W, Crous PW, Samuels GJ. (2000). Molecules, morphology and classification: Towards monophyletic genera in the Ascomycetes. *Studies in Mycology* **45**: 1–4.
- Sepiah M. (1990). New storage disease of guava fruit caused by *Cylindrocladium scoparium*. *Plant Disease* **74**: 253.
- Sharma JK, Mohanan C. (1982). *Cylindrocladium* spp. associated with various diseases of *Eucalyptus* in Kerala. *European Journal of Forest Pathology* **12**: 129–136.
- Sharma JK, Mohanan C, Florence EJM. (1984). Nursery diseases of *Eucalyptus* in Kerala. *European Journal of Forest Pathology* **14**: 77–89.
- Sharma JK, Mohanan C, Florence EJM. (1985). Disease survey in nurseries and plantations of forest tree species grown in Kerala. *Kerala Forest Research Institute, Research Report* **36**: 1–268.
- Sharma JK, Mohanan C, Rugimini P. (1992). Cultural characters and growth of *Cylindrocladium quinqueseptatum* isolates. *European Journal of Forest Pathology* **22**: 217–226.
- Sivapalan A, Metussin R, Hamdan F, Zain RM. (1998). Fungi associated with postharvest fruit rots of *Durio graveolens* and *D. kutejensis* in Brunei Darussalam. *Australasian Plant Pathology* **27**: 274–277.
- Sobers EK, Littrell RH. (1974). Pathogenicity of three species of *Cylindrocladium* to select hosts. *Plant Disease Reporter* **58**: 1017–1019.
- Sobers EK, Seymour CP. (1967). *Cylindrocladium floridanum* sp. nov. associated with decline of peach trees in Florida. *Phytopathology* **57**: 389–393.
- Stevens C, Palmer MA, McRoberts RE. (1990). Use of aminopeptidase substrate specificities to identify species of *Cylindrocladium* in Wisconsin nurseries. *Mycologia* **82**: 436–443.
- Taniguchi T, Tanaka C, Tamai S, Yamanaka N, Futai K. (2008). Identification of *Cylindrocladium* sp. causing damping-off disease of Japanese black pine (*Pinus*

- thunbergii*) and factor affecting the disease severity in a black locust (*Robinia pseudoacacia*)-dominated area. *Journal of Forest Research* **13**: 233–240.
- Taylor JW, Jacobson DJ, Fisher MC. (1999). The evolution of asexual fungi: Reproduction, speciation and classification. *Annual Review of Phytopathology* **37**: 197–246.
- Taylor JW, Jacobson DJ, Kroken SM, Kasuga T, Geiser DM, Hibbett DS, Fisher MC. (2000). Phylogenetic species recognition and species concepts in fungi. *Fungal Genetics and Biology* **31**: 21–32.
- Terashita T. (1968). A new species of *Calonectria* and its conidial state. *Transactions of the Mycological Society of Japan* **8**: 124–129.
- Terashita T, Itô K. (1956). Some notes on *Cylindrocladium scoparium* in Japan. *Bulletin of the Forestry Experiment Station Tokyo* **87**: 33–47.
- Tubaki K. (1958). Studies on Japanese Hyphomycetes. 5. Leaf & stem group with a discussion of the classification of Hyphomycetes and their perfect stages. *Journal of Hattori Botanical Laboratory* **20**: 142–144.
- Uchida JY, Kadooka CY. (1997). Blight of leatherfern caused by *Calonectria theae*, and *Cylindrocladium* spp. *Phytopathology* **87**: S98–S99.
- Vaidya P, Rao VG. (1992). Three undescribed post-harvest diseases of fruits from Maharashtra. *Journal of Economic and Taxonomic Botany* **16**: 241–244.
- Varon AF de. (1991). *Cylindrocladium scoparium* associated with drying up and early death of soybean plants. *Fitopatologia Colombiana* **15**: 2–7.
- Victor D, Crous PW, Janse BJH, Wingfield MJ. (1997). Genetic variation in *Cylindrocladium floridanum* and other morphologically similar *Cylindrocladium* species. *Systematic and Applied Microbiology* **20**: 268–285.
- Vitale A, Aiello D, Castello I, Parlavecchio G, Polizzi G. (2008). First report of crown rot and root rot caused by *Cylindrocladium pauciramosum* on Feijoa (*Feijoa settoiana*) in Italy. *Plant Disease* **92**: 1590.
- Vitale A, Polizzi G. (2007). First record of the perfect stage *Calonectria pauciramosa* on mastic tree in Italy. *Plant Disease* **91**: 328.
- Vitale A, Polizzi G. (2008). First record of leafspots and stem lesions on *Pistacia lentiscus* caused by *Cylindrocladium pauciramosum* and *C. scoparium* in Italy. *Plant Pathology* **57**: 384.

- Wiapara NW, Di Menna ME, Cole ALJ, Skipp RA. (1996). Pathogenicity of *Cylindrocladium scoparium* to pasture clover and grass species. *Australasian Plant Pathology* **25**: 205–211.
- Wolf FA. (1926). Brown leaf spot of leather fern. *Journal of the Elisha Mitchell Scientific Society* **42**: 55–62.
- Wormald H. (1944). A *Cylindrocladium* as the cause of a shoot wilt of varieties of plum and cherry used for rootstocks. *Transactions of the British Mycological Society* **27**: 71–80.
- Wright LP, Wingfield BD, Crous PW, Brenneman T, Wingfield MJ. (2006). Isolation and characterization of microsatellite loci in *Cylindrocladium parasiticum*. *Molecular Ecology Notes* **6**: 110–112.
- Wright LP, Wingfield BD, Crous PW, Wingfield MJ. (2007). Isolation and characterization of microsatellite loci in *Cylindrocladium pauciramsum*. *Molecular Ecology Notes* **7**: 343–345.
- Yoder OC, Turgeon BG. (2001). Fungal genomics and pathogenicity. *Current Opinion in Plant Biology* **4**: 315–321.



Table 1. List of recognized *Calonectria* species and their respective *Cylindrocladium* anamorph species.

Teleomorph	Reference	Anamorph	Reference
<i>Calonectria acicola</i> Gadgil & Dick	Gadgil & Dick 2004	<i>Cylindrocladium acicola</i> Gadgil & Dick	Gadgil & Dick 2004
<i>Calonectria asiatica</i> Crous & Hywel-Jones	Crous <i>et al.</i> 2004b	<i>Cylindrocladium asiaticum</i> Crous & Hywel-Jones	Crous <i>et al.</i> 2004b
<i>Calonectria avesiculata</i> T.S. Schubert, Ell-Gholl, Alfieri & Schoult.	Schubert <i>et al.</i> 1989	<i>Cylindrocladium avesiculatum</i> D.L. Gill, Alfieri & Sobers	Gill <i>et al.</i> 1971
<i>Calonectria clavata</i> Alfieri, El-Gholl, & E.L. Barnard	El-Gholl <i>et al.</i> 1993b	<i>Cylindrocladium flexuosum</i> Crous	Crous <i>et al.</i> 1995
<i>Calonectria colhounii</i> Peerally	Peerally 1973	<i>Cylindrocladium colhounii</i> Peerally	Peerally 1973
<i>Calonectria colombiensis</i> Crous	Crous <i>et al.</i> 2004b	<i>Cylindrocladium colombiense</i> Crous	Crous <i>et al.</i> 2004b
<i>Calonectria gracilipes</i> Crous & G.R.A. Mchau	Crous <i>et al.</i> 1997a	<i>Cylindrocladium graciloideum</i> Crous & G.R.A. Mchau	Crous <i>et al.</i> 1997a
<i>Calonectria gracilis</i> Crous, M.J. Wingf. & Alfenas	Crous <i>et al.</i> 1997b	<i>Cylindrocladium pseudogratile</i> Crous M.J. Wingf. & Alfenas	Crous <i>et al.</i> 1997b
<i>Calonectria hederæ</i> C. Booth & J.S. Murray	Booth & Murray 1960	<i>Cylindrocladium hederæ</i> Peerally	Peerally 1991a
<i>Calonectria hongkongensis</i> Crous	Crous <i>et al.</i> 2004b	<i>Cylindrocladium hongkongense</i> Crous	Crous <i>et al.</i> 2004b



Table 1. (Continued)

Teleomorph	Reference	Anamorph	Reference
<i>Calonectria ilicicola</i> Boedjin & Reitsma	Boedjin & Reitsma 1950	<i>Cylindrocladium parasiticum</i> Crous, M.J. Wingf. & Alfenas	Crous <i>et al.</i> 1993d
<i>Calonectria indusiata</i> Crous	Crous 2002	<i>Cylindrocladium theae</i> (Petch) Alfieri & Sobers	Alfieri <i>et al.</i> 1972
<i>Calonectria insularis</i> C.L. Schoch & Crous	Schoch <i>et al.</i> 1999	<i>Cylindrocladium insulare</i> C.L. Schoch & Crous	Schoch <i>et al.</i> 1999
<i>Calonectria kyotensis</i> Terashita	Terashita 1968	<i>Cylindrocladium floridanum</i> Sobers & C.P. Seymour	Sobers & Seymour 1967
<i>Calonectria leguminum</i> Crous	Crous 2002	<i>Cylindrocladium leguminum</i> Crous	Crous 2002
<i>Calonectria macroconidialis</i> Crous	Crous <i>et al.</i> 1999	<i>Cylindrocladium macroconidiale</i> Crous	Crous <i>et al.</i> 1999
<i>Calonectria madagascariensis</i> Crous	Crous 2002	<i>Cylindrocladium madagascariense</i> Crous	Crous 2002
<i>Calonectria mexicana</i> C.L. Schoch & Crous	Schoch <i>et al.</i> 1999	<i>Cylindrocladium mexicanum</i> C.L. Schoch & Crous	Schoch <i>et al.</i> 1999
<i>Calonectria morganii</i> Crous, Alfenas & M.J. Wingf.	Crous <i>et al.</i> 1993a	<i>Cylindrocladium scoparium</i> Morgan	Morgan 1892
<i>Calonectria multiseptata</i> Crous & M.J. Wingf.	Crous <i>et al.</i> 1998b	<i>Cylindrocladium multiseptatum</i> Crous & M.J. Wingf.	Crous <i>et al.</i> 1998b



Table 1. (Continued)

Teleomorph	Reference	Anamorph	Reference
<i>Calonectria naviculata</i> Crous & M.J. Wingf.	Crous <i>et al.</i> 1994a	<i>Cylindrocladium naviculatum</i> Crous & M.J. Wingf.	Crous <i>et al.</i> 1994a
<i>Calonectria ovata</i> D. Victor & Crous	Victor <i>et al.</i> 1997	<i>Cylindrocladium ovatum</i> El-Gholl , Alfenas, Crous & T.S. Schubert	El-Gholl <i>et al.</i> 1993a
<i>Calonectria pauciramosa</i> C.L. Schoch & Crous	Schoch <i>et al.</i> 1999	<i>Cylindrocladium pauciramosum</i> C.L. Schoch & Crous	Schoch <i>et al.</i> 1999
<i>Calonectria pseudospathiphylli</i> J.C. Kang, Crous & C.L. Schoch	Kang <i>et al.</i> 2001b	<i>Cylindrocladium pseudospathiphylli</i> J.C. Kang, Crous & C.L. Schoch	Kang <i>et al.</i> 2001b
<i>Calonectria pteridis</i> Crous, M.J. Wingf. & Alfenas	Crous <i>et al.</i> 1993c	<i>Cylindrocladium pteridis</i> F.A. Wolf	Wolf 1926
<i>Calonectria pyrochroa</i> Saccardo	Rossmann 1979a	<i>Cylindrocladium ilicicola</i> Boedjin & Reitsma	Boedjin & Reitsma 1950
<i>Calonectria reteaudii</i> C. Booth	Booth 1966	<i>Cylindrocladium reteaudii</i> Boesewinkel	Boesewinkel 1982
<i>Calonectria rumohrae</i> El-Gholl & Alfenas	El-Gholl <i>et al.</i> 1997	<i>Cylindrocladium rumohrae</i> El-Gholl & Alfenas	El-Gholl <i>et al.</i> 1997
<i>Calonectria scoparia</i> Ribeiro & Matsuoka ex Peerally	Peerally 1991	<i>Cylindrocladium candelabrum</i> Viegas	Crous 2002



Table 1. (Continued)

Teleomorph	Reference	Anamorph	Reference
<i>Calonectria spathiphylli</i>	El-Gholl, J.Y. Uchida, Alfenas, T.S. Schubert, Alfieri & A.R. Chase	<i>Cylindrocladium spathiphylli</i>	Schoutties <i>et al.</i> 1982
<i>Calonectria spathulata</i>	El-Gholl, Kimbr. & E.L. Barnard	<i>Cylindrocladium spathulatum</i>	El-Gholl, Kimbr. & E.L. Barnard
<i>Calonectria variabilis</i>	Crous, B.J.H. Janse, D. Victor, G.F. Marais & Alfenas	<i>Cylindrocladium variabile</i>	Crous, B.J.H. Janse, D. Victor, G.F. Marais & Alfenas
		<i>Cylindrocladium angustatum</i>	Crous & El-Gholl
		<i>Cylindrocladium australiense</i>	Crous & K.D. Hyde
		<i>Cylindrocladium canadense</i>	J.C. Kang, Crous & C.L. Schoch.
		<i>Cylindrocladium chinense</i>	Crous
		<i>Cylindrocladium citri</i>	Boedjin & Reitsma
		<i>Cylindrocladium curvatum</i>	Boedjin & Reitsma



Table 1. (Continued)

Teleomorph	Reference	Anamorph	Reference
		<i>Cylindrocladium curvisporum</i> Crous & D. Victor	Victor <i>et al.</i> 1997
		<i>Cylindrocladium ecuadoriae</i> Crous & M.J. Wingf.	Crous <i>et al.</i> 2006
		<i>Cylindrocladium gordoniae</i> Leahy, T.S. Schubert & El-Gholl	Leahy <i>et al.</i> 2000
		<i>Cylindrocladium gracile</i> Boesewinkel	Boesewinkel 1982
		<i>Cylindrocladium hawksworthii</i> Peerally	Peerally 1991b
		<i>Cylindrocladium hurae</i> (Linder & Whetzel) Crous	Crous 2002
		<i>Cylindrocladium indonesiae</i> Crous	Crous <i>et al.</i> 2004b
		<i>Cylindrocladium leucothoes</i> El-Gholl, Leahy & T.S. Schubert	El-Gholl <i>et al.</i> 1989
		<i>Cylindrocladium malesianum</i> Crous	Crous <i>et al.</i> 2004b
		<i>Cylindrocladium multiphialidicum</i> Crous, P. Simoneau & J-M. Riséde	Crous <i>et al.</i> 2004b



Table 1. (Continued)

Teleomorph	Reference	Anamorph	Reference
		<i>Cylindrocladium pacificum</i> J.C. Kang, Crous & C.L. Schoch	Kang <i>et al.</i> 2001b
		<i>Cylindrocladium penicilloides</i> Tubaki	Tubaki 1958
		<i>Cylindrocladium pseudonaviculatum</i> Crous	Crous <i>et al.</i> 2002
		<i>Cylindrocladium sumatrense</i> Crous	Crous <i>et al.</i> 2004b



Table 2. Plant families that are hosts to *Calonectria* species and number of plant host species in each family.


Host Plant family	Host species	Host Plant family	Host species	Host Plant family	Host species	Host Plant family	Host species
Actinidiaceae	2	Cornaceae	1	Malpighiaceae	2	Polypodiaceae	1
Altingiaceae	1	Crassulaceae	1	Malvaceae	6	Proteaceae	7
Anacardiaceae	3	Cupressaceae	4	Meliaceae	2	Pteridaceae	1
Annonaceae	4	Curcubitaceae	3	Moraceae	2	Rhamnaceae	1
Aparagaceae	1	Cycadaceae	1	Musaceae	2	Rhizophoraceae	1
Apiaceae	1	Davalliaceae	1	Myristicaceae	1	Rosaceae	10
Apocynaceae	2	Dennstaedtiaceae	1	Myrsinaceae	1	Rubiaceae	2
Aquifoliaceae	4	Dilleniaceae	1	Myrtaceae	31	Ruscaceae	1
Araceae	5	Dipterocarpaceae	1	Nelumbonaceae	1	Rutaceae	3
Araliaceae	2	Dryopteridaceae	2	Nepenthaceae	1	Salicaceae	3
Arecaceae	20	Ebenaceae	1	Nothofagaceae	1	Sapindaceae	4
Armacariaceae	2	Ericaceae	14	Nymphaeaceae	1	Sapotaceae	3
Aspleniaceae	1	Euphorbiaceae	6	Oleaceae	1	Sarraceniaceae	1



Table 2. (Continued).

Host Plant family	Host species	Host Plant family	Host species	Host Plant family	Host species	Host Plant family	Host species
Asteraceae	5	Fabaceae	57	Onagraceae	2	Saxifragaceae	1
Berberidaceae	2	Fagaceae	4	Orchidaceae	1	Solanaceae	4
Betulaceae	1	Ginkgoaceae	1	Oryzaceae	1	Sterculiaceae	2
Bixaceae	1	Juglandaceae	2	Phoeniceae	1	Strelitziaceae	2
Bromeliaceae	3	Lauraceae	6	Phytolaccaceae	1	Theaceae	1
Buxaceae	1	Laxmanniaceae	1	Pinaceae	17	Ulmaceae	1
Caricaceae	2	Lecythidaceae	1	Piperaceae	1	Verbenaceae	1
Caryophyllaceae	1	Leeaceae	1	Platanaceae	1	Vitaceae	2
Celastraceae	1	Linaceae	1	Plumbaginaceae	1	Vochysiaceae	1
Chenopodiaceae	1	Lomariopsidaceae	1	Poaceae	5	Xanthorrhoeaceae	1
Combretaceae	3	Lythraceae	1	Polygalaceae	1	Zingiberaceae	1
Convolvulaceae	1	Magnoliaceae	2	Polygonaceae	3		

Fig. 1. Morphological characteristic used for identification of *Calonectria* (= *Cylindrocladium*) species. A, B. Macroconidiophores. A. Macroconidiophore of *Ca. pauciramosa*. B. Macroconidiophore of *Ca. hongkongensis*. C–F. Vesicles. C. Obpyriform vesicle of *Ca. pauciramosa*. D. Sphaeropenduculate vesicle of *Ca. hongkongensis*. E, F. Clavate vesicle of *Cy. gracile*. G–I. Macroconidia. G. Macroconidia of *Ca. pauciramosa*. H. Macroconidia of *Ca. hongkongensis*. I. Macroconidia of *Ca. reteaudii*. J. Microconidia of *Ca. reteaudii*. K, L. Fertile branches. K. Fertile branches with doliiform to reniform phialides of *Ca. pauciramosa*. L. Fertile branches of *Ca. reteaudii* with cylindrical to allantoid phialides. M. Asci of *Ca. hongkongensis* with ascospore. N. Ascospores of *Ca. hongkongensis*. Scale bars A, B, M = 20 μm , C–L, N = 10 μm





CHAPTER 2

Calonectria (Cylindrocladium) species associated with dying Pinus cuttings

Published as: Lombard L, Rodas CA, Crous PW, Wingfield BD, Wingfield MJ. (2009). *Calonectria (Cylindrocladium) species associated with dying Pinus cuttings. Persoonia* **23**: 41–47.

ABSTRACT

Calonectria (*Ca.*) species and their *Cylindrocladium* (*Cy.*) anamorphs are well-known pathogens of forest nursery plants in subtropical and tropical areas of the world. An investigation of the mortality of rooted *Pinus* cuttings in a commercial forest nursery in Colombia led to the isolation of two *Cylindrocladium* anamorphs of *Calonectria* species. The aim of this study was to identify these species by using DNA sequence data and morphological comparisons. Two species were identified, namely one undescribed species, and *Cy. gracile*, which is allocated to *Calonectria* as *Ca. brassicae*. The new species, *Calonectria brachiatica* sp. nov., resides in the *Ca. brassicae* species complex. Pathogenicity tests with *Ca. brachiatica* and *Ca. brassicae* showed that both are able to cause disease on *Pinus maximinoi* and *P. tecunumanii*. An emended key is provided to distinguish between *Calonectria* species with clavate vesicles and 1-septate macroconidia.

Taxonomic novelties: *Calonectria brassicae* (Panwar & Bohra) L. Lombard, M.J. Wingf. & Crous comb. nov., *Calonectria brachiatica* L. Lombard, M.J. Wingf. & Crous. sp. nov.

INTRODUCTION

Species of *Calonectria* (anamorph *Cylindrocladium*) are plant pathogens associated with a large number of agronomic and forestry crops in temperate, sub-tropical and tropical climates, worldwide (Crous & Wingfield 1994, Crous 2002). Infection by these fungi gives rise to symptoms including cutting rot (Crous *et al.* 1991), damping-off (Sharma *et al.* 1984, Ferreira 1995), leaf spot (Sharma *et al.* 1984, Ferreira *et al.* 1995, Crous *et al.* 1998), shoot blight (Crous *et al.* 1991, Crous *et al.* 1998), stem cankers (Sharma *et al.* 1984, Crous *et al.* 1991) and root disease (Mohanani & Sharma 1985, Crous *et al.* 1991) on various forest tree species.

The first report of *Ca. morganii* (as *Cy. scoparium*) infecting *Pinus* species was by Graves (1915), but he failed to re-induce the symptoms and assumed that it was a saprobe. There have subsequently been several reports of *Cylindrocladium* spp. infecting *Pinus* and other conifers, leading to root rot, stem cankers and needle blight (Jackson 1938, Cox 1953, Thies & Patton 1970, Sober & Alfieri 1972, Cordell & Skilling 1975, Darvas *et al.* 1978, Crous *et al.* 1991, Crous 2002). Most of these reports implicated *Ca. morganii* and *Ca. pteridis* (as *Cy. macrosporum* or *Cy. pteridis*) as the primary pathogens (Thies & Patton 1970, Ahmad & Ahmad 1982). However, as knowledge of these fungi has grown, together with refinement of their taxonomy applying DNA sequence comparisons (Crous *et al.* 2004, 2006), several additional *Cylindrocladium* species have been identified as causal agents of disease on different conifer species. These include *Ca. acicola*, *Ca. colhounii*, *Ca. kyotensis* (= *C. floridanum*), *Ca. pteridis*, *Cy. canadense*, *Cy. curvisporum*, *Cy. gracile* and *Cy. pacificum* (Hodges & May 1970, Crous 2002, Gadgil & Dick 2004, Taniguchi *et al.* 2008).

In a recent survey, wilting, collar and root rot symptoms were observed in Colombian nurseries generating *Pinus* spp. from cuttings. Isolations from these diseased plants consistently yielded *Cylindrocladium* anamorphs of *Calonectria* species, and hence the aim of this study was to identify them, and to determine if they were the causal agents of the disease in Colombian nurseries.

MATERIAL AND METHODS

Isolates

Pinus maximinoi and *P. tecunumanii* rooted cutting plants showing symptoms of collar and root rot (Fig. 1) were collected from a nursery close to Buga in Colombia. Isolations were made directly from lesions on the lower stems and roots on fusarium selective medium (FSM; Nelson *et al.* 1983) and malt extract agar (MEA, 2 % w/v; Biolab, Midrand, South Africa). After 5 d of incubation at 25 °C, fungal colonies of *Calonectria* spp. were transferred on to MEA and incubated further for 7 d. For each isolate, single conidial cultures were prepared on MEA, and representative strains are maintained in the culture collection (CMW) of the Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria, South Africa and the Centraalbureau voor Schimmelcultures (CBS), Utrecht, The Netherlands.

Taxonomy

For morphological identification of *Calonectria* isolates, single conidial cultures were prepared on MEA and synthetic nutrient-poor agar (SNA; Nirenburg 1981). Inoculated plates were incubated at room temperature and examined after 7 d. Gross morphological characteristics were assessed by mounting fungal structures in lactic acid. Thirty measurements at $\times 1\,000$ magnification were made for each isolate. The 95 % confidence levels were determined for the pooled measurements of the respective species studied and extremes for structure sizes are given in parentheses. Optimal growth temperatures were determined between 6–36 °C at 6 °C intervals in the dark on MEA for each isolate. Colony reverse colours were determined after 7 d on MEA at 24 °C in the dark, using the colour charts of Rayner (1970) for comparison.

DNA phylogeny

Calonectria isolates were grown on MEA for 7 d. Mycelium was then scraped from the surfaces of the cultures, freeze-dried, and ground to a powder in liquid nitrogen, using a mortar and pestle. DNA was extracted from the powdered mycelium as described by Lombard *et al.* (2008). A fragment of the β -tubulin gene region was amplified and sequenced using primers T1 (O'Donnell & Cigelnik 1997) and CYLTUB1R (Crous *et al.* 2004) and a fragment for the histone H3 (HIS3) gene region was sequenced using primers CYLH3F and CYLH3R (Crous *et al.* 2004).

The PCR reaction mixture used to amplify the different loci consisted of 2.5 units FastStart *Taq* polymerase (Roche Applied Science, USA), 10× PCR buffer, 1–1.5 mM MgCl₂, 0.25 mM of each dNTP, 0.5 μM of each primer and approximately 30 ng of fungal genomic DNA, made up to a total reaction volume of 25 μL with sterile distilled water.

Amplified fragments were purified using High Pure PCR Product Purification Kit (Roche, USA) and sequenced in both directions. For this purpose, the BigDye terminator sequencing kit v. 3.1 (Applied Biosystems, USA) and an ABI PRISM™ 3100 DNA sequencer (Applied Biosystems) were used. All PCRs and sequencing reactions were performed on an Eppendorf Mastercycler Personal PCR (Eppendorf AG, Germany) with cycling conditions as described in Crous *et al.* (2006) for each locus.

Sequences generated were added to other sequences obtained from GenBank (<http://www.ncbi.nlm.nih.gov>) and were assembled and aligned using Sequence Navigator v. 1.0.1 (Applied Biosystems) and MAFFT v. 5.11 (Kato *et al.* 2005), respectively. The aligned sequences were then manually corrected where needed.

PAUP (Phylogenetic Analysis Using Parsimony, v. 4.0b10; Swofford 2002) was used to analyse the DNA sequence datasets. A partition homogeneity test (Farris *et al.* 1994) and a 70 % reciprocal bootstrap method (Mason-Gamer & Kellog 1996) were applied to evaluate the feasibility of combining the data sets. Phylogenetic relationships were estimated by heuristic searches based on 1 000 random addition sequences and tree bisection-reconnection, with the branch swapping option set on ‘best trees’ only.

All characters were weighted equally and alignment gaps were treated as missing data. Measures calculated for parsimony included tree length (TL), consistency index (CI), retention index (RI) and rescaled consistence index (RC). Bootstrap analysis (Hillis & Bull 1993) was based on 1 000 replications. All sequences for the isolates studied were analysed using the Basic Local Alignment Search Tool for Nucleotide sequences (BLASTN, Altschul *et al.* 1990). The phylogenetic analysis included 19 partial gene sequences per gene, representing eight *Calonectria* species (Table 1) closely related to the isolates studied. *Calonectria colombiensis* was used as the outgroup taxon. All sequences were deposited in GenBank and the alignments in TreeBASE (<http://treebase.org>).

A Markov Chain Monte Carlo (MCMC) algorithm was used to generate phylogenetic trees with Bayesian probabilities using MrBayes v. 3.1.1 (Ronquist & Huelsenbeck 2003). Models of nucleotide substitution for each gene were determined using Mrmodeltest (Nylander 2004) and included for each gene partition. Four MCMC chains were run simultaneously from random trees for one million generations and sampled every 100 generations. The first 800 trees were discarded as the burn-in phase of each analysis and posterior probabilities determined from the remaining trees.

Pathogenicity tests

In order to test the pathogenicity of the *Calonectria* spp. collected in this study, profusely sporulating isolates CMW 25293, representing *Ca. brachiatica*, CMW 25296 and CMW 25297, both representing *Ca. brassicae*, were used for inoculations onto rooted cuttings of *P. maximinoi*. Isolate CMW 25299, representing *Ca. brassicae* and isolates CMW 25302 and CMW 25307 representing *Ca. brachiatica* were used for inoculations onto rooted cuttings of *P. tecunumanii*. Trees used for inoculation were between 0.5–1 m in height and 10–50 mm diam at the root collar. Trees were maintained in a greenhouse under controlled conditions prior to inoculation, so that they could become acclimatised and to ensure that they were healthy. Sixty trees for each *Pinus* spp. were used and an additional 60 trees were used as controls. This resulted in a total of 180 trees in the pathogenicity tests.

Inoculations were performed in the greenhouse by making a 5 mm diam wound on the main stems of plants with a cork borer to expose the cambium. The cambial discs were replaced with an MEA disc overgrown with the test fungi taken from 7 d old cultures. The inoculum discs were placed mycelium side facing the cambium and the inoculation points were sealed with Parafilm to reduce contamination and desiccation. Control trees were treated in a similar fashion but inoculated with a sterile MEA plug.

Six weeks after inoculation, lesion lengths on the stems of the plants were measured. The results were subsequently analysed using SAS Analytical Programmes v. 2002. Re-isolations were made from the edges of lesions on the test trees to ensure the presence of the inoculated fungi.

RESULTS

DNA phylogeny

For the β -tubulin gene region, approx. 580 bases were generated for each of the isolates used in the study (Table 1). The adjusted alignment included 19 taxa with the outgroup, and 523 characters including gaps after uneven ends were removed from the beginning of each sequence. Of these characters, 459 were constant and uninformative. For the analysis, only the 64 parsimony informative characters were included. Parsimony analysis of the aligned sequences yielded five most parsimonious trees (TL = 231 steps; CI = 0.870; RI = 0.799; RC = 0.695; results not shown).

Sequences for the histone gene region consisted of approx. 460 bases for the isolates used in the study and the adjusted alignment of 19 taxa including the outgroup, consisted of 466 characters including gaps. Of these characters, 391 were excluded as constant and parsimony uninformative and 79 parsimony informative characters included. Analysis of the aligned data yielded one most parsimonious tree (TL = 290 steps; CI = 0.845; RI = 0.807; RC = 0.682; results not shown).

The partition homogeneity test showed that the β -tubulin and histone data set could be combined ($P = 0.245$). The 70 % reciprocal bootstrap method indicated no conflict in tree topology among the two partitions, resulting in a combined sequence data set consisting of 993 characters including gaps for the 19 taxa (including outgroup). Of these, 850 characters were constant and parsimony uninformative and excluded from the analysis. There were 143 characters in the analysis that were parsimony informative. Parsimony analysis of the combined alignments yielded one most parsimonious tree (TL = 526 steps; CI = 0.848; RI = 0.791; RC = 0.670), which is presented in Fig. 2 (TreeBase S2568).

All the isolates obtained from the *Pinus* spp. used in this study grouped in the *Ca. brassicae* species complex with a bootstrap (BP) value of 96 and a low Bayesian posterior probability (PP) of 0.70. This clade was further subdivided into two clades. The first clade (BP = 64, PP below 0.70) representing *Ca. brassicae*, included the type of *Cy. gracile* and *Cy. clavatum*. It also included three isolates (CMW 25297, CMW 25296 and CMW 25299) from *P. maximinoi* and *P. tecunumanii*. The second clade (BP = 98, PP = 0.82) accommodated *Calonectria* isolates (CMW 25293, CMW 25298, CMW 25302 and CMW 25307), representing what we recognize as a distinct species. The consensus tree obtained with Bayesian analysis showed topographical similarities with the most parsimonious tree as indicated in Fig. 2.

Pathogenicity tests

All plants inoculated with *Calonectria* spp. in this study developed lesions. Lesions included discolouration of the vascular tissue with abundant resin formation, 6 wk after inoculation. Lesions on the control trees were either non-existent or small, representing wound reactions. There were significant ($p < 0.0001$) differences in lesion lengths associated with individual isolates used on *P. maximinoi* (Fig. 3). Comparisons of the lesion lengths clearly showed that *Ca. brassicae* (CMW 25297) produced the longest average lesions (av. = 30.04 mm) compared to the undescribed *Calonectria* sp. (CMW 25293) (av. = 14.41mm). The other *Ca. brassicae* isolate (CMW 25296) produced an average lesion length of 15.30 mm. Lesions on the control trees were an average of 8.84 mm and significantly ($p < 0.0001$) smaller than those on any of the trees inoculated with the test fungi (Fig. 5).

Results of inoculations on *P. tecunumanii* were similar to those on *P. maximinoi*. Thus, *Ca. brassicae* (CMW 25299) (av. = 20.64 mm) produced the longest lesions compared with the undescribed *Calonectria* sp. (CMW 25302; av. = 18.63 mm and CMW 25307; av. = 15.20 mm). The lesions on the *P. tecunumanii* control trees were also significantly ($p < 0.0001$) smaller (av. = 8.82 mm) than those on any of the trees inoculated with the test fungi. Re-isolations from the test trees consistently yielded the inoculated fungi and no *Calonectria* spp. were isolated from the control trees.

Taxonomy

Isolates CMW 25296, CMW 25297 and CMW 25299 clearly represent *Ca. brassicae* based on morphological observations (Crous 2002) and comparisons of DNA sequence data. Isolates CMW 25293, CMW 25298, CMW 25302 and CMW 25307 represent an undescribed species closely related to *Ca. brassicae* but morphologically distinct. Species of *Cylindrocladium* (1892) represent anamorph states of *Calonectria* (1867) (Rossman *et al.* 1999), and therefore this fungus is described as a new species of *Calonectria*, which represents the older generic name for these holomorphs:

Calonectria brachiatica L. Lombard, M.J. Wingf. & Crous, **sp. nov.** MycoBank MB512998.
Fig. 4

Etymology: Name refers to the lateral branches on the conidiophore stipes .

Stipa extensiones septatum, hyalinum, 134–318 μm , in vesiculam clavatum, 5–7 μm diam terminans. Conidia cylindrica, hyalina, 1–2-septata, utrinque obtusa, (37–)40–48(–50) \times 4–6 μm .

Teleomorph unknown. *Conidiophores* with a stipe bearing penicillate suites of fertile branches, stipe extensions, and terminal vesicles; stipe septate, hyaline, smooth, $32\text{--}67 \times 6\text{--}8 \mu\text{m}$; stipe extensions septate, straight to flexuous, $134\text{--}318 \mu\text{m}$ long, $4\text{--}5 \mu\text{m}$ wide at the apical septum, terminating in a clavate vesicle, $5\text{--}7 \mu\text{m}$ diam; lateral stipe extensions (90° to the axis) also present. *Conidiogenous apparatus* $40\text{--}81 \mu\text{m}$ long, and $35\text{--}84 \mu\text{m}$ wide; primary branches aseptate or 1-septate, $15\text{--}30 \times 4\text{--}6 \mu\text{m}$; secondary branches aseptate, $10\text{--}23 \times 3\text{--}5 \mu\text{m}$; tertiary branches and additional branches (-5), aseptate, $10\text{--}15 \times 3\text{--}4 \mu\text{m}$, each terminal branch producing 2–6 phialides; phialides doliform to reniform, hyaline, aseptate, $10\text{--}15 \times 3\text{--}4 \mu\text{m}$; apex with minute periclinal thickening and inconspicuous collarete. *Macroconidia* cylindrical, rounded at both ends, straight, $(37\text{--})40\text{--}48(-50) \times 4\text{--}6 \mu\text{m}$ (av. = $44 \times 5 \mu\text{m}$), 1(-2)-septate, lacking a visible abscission scar, held in parallel cylindrical clusters by colourless slime. *Mega-* and *microconidia* not seen.

Specimens examined : **Colombia**, Valle del Cauca, Buga, from *Pinus maximinoi*, July 2007, M.J. Wingfield, Herb. PREM 60197, **holotype** of *Ca. brachiatica*, culture ex-type CMW 25298 = CBS 123700; Valle del Cauca, Buga, from *P. tecunumanii*, July 2007, M.J. Wingfield, culture CMW 25303 = CBS 123699; Valle del Cauca, Buga, from *P. tecunumanii*, July 2007, M.J. Wingfield, (Herb. PREM 60198) culture CMW 25341 = CBS 123703.

Cultural characteristics: Colonies fast growing with optimal growth temperature at 24°C (growth at $12\text{--}30^\circ\text{C}$) on MEA, reverse amber (13k) to sepia (13i) brown after 7 d; abundant white aerial mycelium with moderate to extensive sporulation; chlamydospores extensive throughout the medium.

Substrate: *Pinus maximinoi*, *P. tecunumanii*.

Distribution: Colombia.

Notes: The anamorph state of *Ca. brachiatica* can be distinguished from *C. gracile*, *C. pseudogracile* and *C. graciloideum* by its shorter macroconidia. Another characteristic distinguishing *Ca. brachiatica* is the formation of lateral branches not reported for *C. gracile* or other closely related species.

Calonectria brassicae (Panwar & Bohra) L. Lombard, M.J. Wingf. & Crous, **comb. nov.**
Mycobank MB513423. Fig. 5

Basionym. Cyandrocladium brassicae Panwar & Bohra, Indian Phytopathology 27: 425. 1974.

= *Cylindrocarpon gracile* Bugnic., Encycl. Mycologique 11: 162. 1939.

≡ *Cyandrocladium gracile* (Bugnic.) Boesew., Trans. Brit. Mycol. Soc. 78: 554. 1982.

= *Cyandrocladium clavatum* Hodges & L.C. May, Phytopathology 62: 900. 1972.

Notes: Both the names *Ca. clavata* and *Ca. gracilis* and are already occupied, hence the oldest available epithet is that of *Cy. brassicae* (Crous 2002).

DISCUSSION

Results of this study show that *Calonectria* species are important pathogens in pine cutting nurseries in Colombia. In this case, two species were discovered, the one newly described here as *Ca. brachiatica* and the other representing *Ca. brassicae*. Both of the species were pathogenic on *P. maximinoi* and *P. tecunumanii*.

The description of *Ca. brachiatica* from *P. maximinoi* and *P. tecunumanii* adds a new species to the *Ca. brassicae* species complex, which already includes six other *Calonectria* species (Crous 2002, Crous *et al.* 2006). This species can be distinguished from the other species in the complex by the formation of lateral branches on the macroconidiophores and the presence of a small number of 2-septate macroconidia. Macroconidial dimensions (av. = $44 \times 5 \mu\text{m}$) are also smaller than those of *Ca. brassicae* (av. $53 \times 4.5 \mu\text{m}$).

A recent study of *Calonectria* spp. with clavate vesicles by Crous *et al.* (2006) attempted to resolve the taxonomic status of these species, and added two new species to the group. Crosses among isolates of *Ca. brachiatica* and isolates of *Ca. brassicae*, did not result in sexual structures in the present study, and teleomorphs are rarely observed in this species complex.

Hodges & May (1972) reported *Ca. brassicae* (as *Cy. clavatum*) from several *Pinus* species in nurseries and plantations in Brazil. Subsequent studies based on comparisons of DNA sequence data revealed *Cy. clavatum* to be a synonym of *Cy. gracile* (Crous *et al.* 1995, 1999, Schoch *et al.* 2001). *Calonectria brassicae* (as *Cy. gracile*) is a well-known pathogen of numerous plant hosts in subtropical and tropical areas of the world. However, in Colombia,

this plant pathogen has been isolated only from soil (Crous 2002, Crous *et al.* 2006). This study thus represents the first report of *Ca. brassicae* infecting *Pinus* spp. in Colombia.

Pathogenicity tests with isolates of *Ca. brachiatica* and *Ca. brassicae* clearly showed that they are able to cause symptoms similar to those observed in naturally infected plants. Both *P. maximinoi* and *P. tecunumanii* were highly susceptible to infection by *Ca. brassicae*. This supports earlier work of Hodges & May (1972) in Brazil, where they reported a similar situation. In their study, seven *Pinus* spp. were wound-inoculated with *Ca. brassicae* and this resulted in mortality of all test plants within 2 wk. Although they did not include *P. maximinoi* and *P. tecunumanii* in the study, they concluded that the pathogen is highly virulent and regarded it as unique in causing disease symptoms in established plantations of *Pinus* spp. No disease symptoms associated with *C. brachiatica* or *Ca. brassicae* were seen in established plantations in the present study and we primarily regard these fungi as nursery pathogens, of which the former species is more virulent than the latter.

The use of SNA (Nirenburg 1981) rather than carnation leaf agar (CLA; Fisher *et al.* 1982) for morphological descriptions of *Calonectria* species represents a new approach employed in this study. Previously, species descriptions for *Calonectria* have typically been conducted on carnation leaf pieces on tap water agar (Crous *et al.* 1992). However, carnation leaves are not always readily available for such studies and SNA, a low nutrient medium, also used for the related genera *Fusarium* and *Cylindrocarpon* species identification (Halleen *et al.* 2006, Leslie & Summerell 2006), provides a useful medium for which the chemical components are readily available. Another advantage of using SNA is its transparent nature, allowing direct viewing through a compound microscope as well as on mounted agar blocks for higher magnification (Leslie & Summerell 2006). In this study, it was found that the *Calonectria* isolates sporulate profusely on the surface of SNA and comparisons of measurements for structures on SNA and those on CLA showed no significant difference. However, CLA remains important to induce the formation of teleomorph structures in homothallic isolates or heterothallic isolates for which both mating types are present.

Key to *Calonectria* species with clavate vesicles and predominantly 1-septate macroconidia (To be inserted in Crous 2002, p. 56, couplet no. 2)

- 2. Stipe extension thick-walled; vesicle acicular to clavate *Ca. avesculata*
- 2. Stipe extension not thick-walled; vesicle clavate 3



3. Teleomorph unknown	4
3. Teleomorph readily formed	7
4. Macroconidia always 1-(2)-septate	5
4. Macroconidia 1-(3)-septate	6
5. Macroconidia 1-septate, (38-)40-55(-65) × (3.5-)4-5(-6) μm, av. = 53 × 4.5 μm; lateral stipe extensions absent	<i>Ca. brassicae</i>
5. Macroconidia 1-(2)-septate, (37-)40-48(-50) × 4-6 μm, av. = 44 × 5 μm; lateral stipe extensions present	<i>Ca. brachiatica</i>
6. Macroconidia (48-)57-68(-75) × (6-)6.5(-7) μm, av. = 63 × 6.5 μm	<i>Cy. australiense</i>
6. Macroconidia (45-)48-55(-65) × (4-)4.5(-5) μm, av. = 51 × 4.5 μm ..	<i>Cy. ecuadoriae</i>
7. Macroconidial state absent; megaconidia and microconidia present	<i>Ca. multiseptata</i>
7. Macroconidial state present	8
8. Teleomorph homothallic	9
8. Teleomorph heterothallic	10
9. Perithecia orange; macroconidia av. size = 45 × 4.5 μm	<i>Ca. gracilipes</i>
9. Perithecia red; macroconidia av. size = 56 × 4.5 μm	<i>Ca. gracilis</i>
10. Perithecia orange; macroconidia av. size = 32 × 3 μm	<i>Ca. clavata</i>
10. Perithecia red-brown; macroconidia av. size = 30 × 3 μm	<i>Ca. pteridis</i>

REFERENCES

- Ahmad N, Ahmad S. (1982). Needle disease of pine caused by *Cylindrocladium macrosporum*. *The Malaysian Forester* **45**: 84–86.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. (1990). Basic Local Alignment Search Tool. *Journal of Molecular Biology* **215**: 403–410.
- Cordell CE, Skilling DD. (1975). Forest nursery diseases in the U.S.A. 7. *Cylindrocladium* root rot. U.S.D.A. *Forest Service Handbook* No. **470**: 23–26.
- Cox RS. (1953). Etiology and control of a serious complex of diseases of conifer seedlings. *Phytopathology* **43**: 469.
- Crous PW. (2002). Taxonomy and pathology of *Cylindrocladium* (*Calonectria*) and allied genera. APS Press, St. Paul, Minnesota, USA.
- Crous PW, Groenewald JZ, Risède J-M, Simoneau P, Hyde KD. (2006). *Calonectria* species and their *Cylindrocladium* anamorphs: species with clavate vesicles. *Studies in Mycology* **55**: 213–226.
- Crous PW, Groenewald JZ, Risède J-M, Simoneau P, Hywel-Jones N. (2004). *Calonectria* species and their *Cylindrocladium* anamorphs: species with sphaeropedunculate vesicles. *Studies in Mycology* **50**: 415–430.
- Crous PW, Kang JC, Schoch CL, Mchua GRA. (1999). Phylogenetic relationships of *Cylindrocladium pseudogracile* and *Cylindrocladium rumohrae* with morphologically similar taxa, based on morphology and DNA sequences of internal transcribed spacers and β -tubulin. *Canadian Journal of Botany* **77**: 1813–1820.
- Crous PW, Korf A, Zyl WH van. (1995). Nuclear DNA polymorphisms of *Cylindrocladium* species with 1-septate conidia and clavate vesicles. *Systematic and Applied Microbiology* **18**: 224–250.
- Crous PW, Phillips AJL, Wingfield MJ. (1991). The genera *Cylindrocladium* and *Cylindrocladiella* in South Africa, with special reference to forest nurseries. *South African Journal of Forestry* **157**: 69–85.
- Crous PW, Phillips AJL, Wingfield MJ. (1992). Effects of cultural conditions on vesicle and conidium morphology in species of *Cylindrocladium* and *Cylindrocladiella*. *Mycologia* **84**: 497–504.
- Crous PW, Wingfield MJ. (1994). A monograph of *Cylindrocladium*, including anamorphs of *Calonectria*. *Mycotaxon* **51**: 341–435.

- Crous PW, Wingfield MJ, Mohammed C, Yuan ZQ. (1998). New foliar pathogens of *Eucalyptus* from Australia and Indonesia. *Mycological Research* **102**: 527–532.
- Darvas JM, Scott DB, Kotze JM. (1978). Fungi associated with damping-off in coniferous seedlings in South African nurseries. *South African Forestry Journal* **104**: 15–19.
- Farris JS, Källersjö M, Kluge AG, Bult C. (1994). Testing significance of incongruence. *Cladistics* **10**: 315–320.
- Ferreira FA, Alfenas AC, Moreira AM, Demuner NLJ. (1995). Foliar eucalypt disease in tropical regions of Brasil caused by *Cylindrocladium pteridis*. *Fitopatologia Brasileira* **20**: 107–110.
- Fisher NL, Burgess LW, Toussoun TA, Nelson PE. (1982). Carnation leaves as a substrate and for preserving cultures of *Fusarium* species. *Phytopathology* **72**: 151–153.
- Gadgil PD, Dick MA. (2004). Fungi silvicolae novazelandiae: 5. *New Zealand Journal of Forestry Science* **34**: 316–323.
- Graves AH. (1915). Root rot of coniferous seedlings. *Phytopathology* **5**: 213–217.
- Halleen F, Schroers H-J, Groenewald JZ, Rego C, Oliveira H, Crous PW. (2006). *Neonectria liriodendri* sp. nov., the main causal agent of black foot disease of grapevines. *Studies in Mycology* **55**: 227–234.
- Hillis DM, Bull JJ. (1993). An empirical test of bootstrapping as a method for assessing confidence in phylogenetic analysis. *Systematic Biology* **42**: 182–192.
- Hodges CS, May LC. (1972). A root disease of pine, *Araucaria*, and *Eucalyptus* in Brazil caused by a new species of *Cylindrocladium*. *Phytopathology* **62**: 898–901.
- Jackson LWR. (1938). *Cylindrocladium* associated with diseases of tree seedlings. *Plant Disease Reporter* **22**: 84–85.
- Katoh K, Kuma K, Toh H, Miyata T. (2005). MAFFT version 5: improvement in accuracy of multiple sequence alignment. *Nucleic Acid Research* **33**: 511–518.
- Leslie JF, Summerell BA. (2006). *The Fusarium laboratory manual*. Blackwell Publishing, Iowa, USA.
- Lombard L, Bogale M, Montenegro F, Wingfield BD, Wingfield MJ. (2008). A new bark canker disease of the tropical hardwood tree *Cedrelinga cateniformis* in Ecuador. *Fungal Diversity* **31**: 73–81.
- Mason-Gamer RJ & Kellogg EA. (1996). Testing for phylogenetic conflict among molecular data sets in the tribe *Triticeae* (*Gramineae*). *Systematic Biology* **45**: 524–545.

- Mohan C, Sharma JK. (1985). *Cylindrocladium* causing seedling diseases of *Eucalyptus* in Kerala, India. *Transactions of the British Mycological Society* **84**: 538–539.
- Nelson PE, Toussoun TA, Marasas WF0. (1983). *Fusarium* species – an illustrated manual for identification, p. 5–18. The Pennsylvania State University Press, University Park, PA.
- Nirenburg HI. (1981). A simplified method for identifying *Fusarium* spp. occurring on wheat. *Canadian Journal of Botany* **59**: 1599–1609.
- Nylander JAA. (2004). MrModeltest v. 2. Programme distributed by the author. Evolutionary Biology Centre, Uppsala University.
- O'Donnell K, Cigelnik E. (1997). Two divergent intragenomic rDNA ITS2 types within a monophyletic lineage of the fungus *Fusarium* are nonorthologous. *Molecular Phylogenetics and Evolution* **7**: 103–116.
- Panwar KS & Bohra A. (1974). A new species of *Cylindrocladium* from India. *Indian Phytopathology* **27**: 425.
- Rayner RW. (1970). A mycological colour chart. Commonwealth Mycological Institute, Kew, Surrey. British Mycological Society.
- Ronquist F, Heuleisenbeck JP. (2003). MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* **19**: 1572–1574.
- Rossman AY, Samuels GJ, Rogerson CT, Lowen R. (1999). Genera of *Bionectriaceae*, *Hypocreaceae* and *Nectriaceae* (*Hypocreales*, Ascomycetes). *Studies in Mycology* **42**: 1–248.
- Schoch CL, Crous PW, Wingfield BD, Wingfield MJ. (2001). Phylogeny of *Calonectria* based on comparisons of β -tubulin DNA sequences. *Mycological Research* **105**: 1045–1052.
- Sharma JK, Mohan C, Florence EJM. (1984). Nursery diseases of *Eucalyptus* in Kerala. *European Journal of Forest Pathology* **14**: 77–89.
- Sobers EK, Alfieri SA. (1972). Species of *Cylindrocladium* and their hosts in Florida and Georgia. *Proceedings of the Florida State Horticultural Society* **85**: 366–369.
- Swofford DL. (2002). PAUP*. Phylogenetic analysis using parsimony (* and other methods), 4.0b10. Computer programme. Sunderland, Massachusetts, USA: Sinauer Associates.
- Taniguchi T, Tanaka C, Tamai S, Yamanaka N, Futai K. (2008). Identification of *Cylindrocladium* sp. causing damping-off disease of Japanese black pine (*Pinus thunbergii*) and factors affecting the disease severity in a black locust (*Robinia pseudoacacia*)-dominated area. *Journal of Forest Research* **13**: 233–240.



Thies WF, Patton RF. (1970). The biology of *Cylindrocladium scoparium* in Wisconsin forest tree nurseries. *Phytopathology* **60**: 1662–1668.



Table 1. Strains of *Calonectria* (*Cylindrocladium*) species included in the phylogenetic analyses (TreeBase S2568)

Species	Isolate number ¹	β -tubulin ²	Histone H3 ²	Host	Origin	Collector
<i>Ca. avesiculata</i>	CBS 313.92 ^T	AF333392	DQ190620	<i>Ilex vomitoria</i>	USA	S.A. Alfieri
<i>Ca. brachiatica</i> sp. nov.	CMW 25293	FJ716710	FJ716714	<i>P. maximinoi</i>	Colombia	M.J. Wingfield
	CMW 25298 (=	FJ696388	FJ696396	<i>P. maximinoi</i>	Colombia	M.J. Wingfield
	CBS 123700) ^T					
	CMW 25302	FJ716708	FJ716712	<i>P. tecunumanii</i>	Colombia	M.J. Wingfield
	CMW 25307	FJ716709	FJ716713	<i>P. tecunumanii</i>	Colombia	M.J. Wingfield
<i>Ca. brassicae</i> com. nov.	CBS 111869 ^T	AF232857	DQ190720	<i>Argyreia</i> sp.	South East Asia	
	CBS 111478	DQ190611	DQ190719	Soil	Brazil	A.C. Alfenas
	CMW 25296	FJ716707	FJ716711	<i>P. maximinoi</i>	Colombia	M.J. Wingfield
	CMW 25297;	FJ696387	FJ696395	<i>P. maximinoi</i>	Colombia	M.J. Wingfield
	CBS123702					
	CMW 25299;	FJ696390	FJ696398	<i>P. tecunumanii</i>	Colombia	M.J. Wingfield
CBS123701						
<i>Ca. clavata</i>	CBS 114557 ^T	AF333396	DQ190623	<i>Callistemon viminalis</i>	USA	N.E. El-Gholl
	CBS 114666	DQ190549	DQ190624	<i>Pinus caribaea</i>	Brazil	C.S. Hodges



Table 1. (Continued)

Species	Isolate number ¹	β -tubulin ²	Histone H3 ²	Host	Origin	Collector
<i>Cy. clavatum</i> (= <i>Cy. gracile</i>)	CBS111776 ^T	AF232850	DQ190700	<i>Pinus caribaea</i>	Brazil	C.S. Hodges
<i>Ca. colombiensis</i>	CBS 12221	AY725620	AY725663	Soil	Colombia	M.J. Wingfield
<i>Cy. ecuadoriae</i>	CBS 111406 ^T	DQ190600	DQ190705	Soil	Ecuador	M.J. Wingfield
<i>Ca. gracilipes</i>	CBS 111141 ^T	DQ190566	DQ190644	<i>Eucalyptus</i> sp.	Colombia	M.J. Wingfield
	CBS 115674	AF333406	DQ190645	Soil	Colombia	M.J. Wingfield
<i>Ca. gracilis</i>	CBS 111284	DQ190567	DQ190647	<i>Manilkara</i> sp.	Brazil	P.W. Crous
	CBS 111807 ^T	AF232858	DQ190646		Brazil	

¹CBS: Centraalbureau voor Schimmelcultures, Utrecht, The Netherlands; CMW: culture collection of the Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria Pretoria, South Africa; ²GenBank accession numbers. ^TEx-type culture.

Fig. 1. Collar and root rot on *Pinus maximinoi* and *P. tecunumanii*. A. Girdled stem of *P. maximinoi*. B. Exposed *P. maximinoi* root collar showing discolouration and resin exudation. C, D. Exposed *P. tecunumanii* root collars showing girdling and discolouration of the cambium.



Fig. 2. The most parsimonious tree obtained from a heuristic search with 1000 random addition of the combined β -tubulin and Histone H3 sequence alignments. Scale bar shows 10 changes and bootstrap support values from 1 000 replicates are shown above the branches. Bayesian posterior probabilities are indicated below the branches. Red lines indicates bootstrap support value of 100 and posterior probability value of 1.00. Bold lines indicate branches present in the Bayesian consensus tree. The tree was rooted with *Calonectria colombiensis* (CBS 112221).






Fig. 3. Histogram showing mean lesion lengths induced by each isolate on *P. maximinoi* and *P. tecunumanii*. *Calonectria brassicae* is represented by CMW 25296, CMW 25297 and CMW 25299. *Ca. brachiatica* is represented by CMW 25293, CMW 25302 and CMW 25307.






Fig. 4. *Calonectria brachiatica*. A. Macroconidiophore with lateral branching stipe extensions. B, C. Clavate vesicles. D. Fertile branches. E. Macroconidia. Scale bars A = 20 μm , B–E = 10 μm .







Fig. 5. *Calonectria brassicae*. A. Macroconidiophore on SNA. B. Macroconidia. C. Fertile branches. D, E. Clavate vesicles. Scale bars A = 20 μm , B–E = 10 μm .





CHAPTER 3

Calonectria species associated with cutting rot of *Eucalyptus*

ABSTRACT

Decline in the productivity of *Eucalyptus* hybrid cutting production in the GuangDong Province of China is linked to cutting rot associated with several *Calonectria* spp. The aim of this study was to identify these fungi using morphological and DNA sequence comparisons. Two previously undescribed *Calonectria* spp., *Ca. pseudoreteaudii* sp. nov. and *Ca. cerciana* sp. nov. were identified as well as *Ca. pauciramosa*. *Calonectria pseudoreteaudii* resides in the *Ca. reteaudii* complex and *Ca. cerciana* is closely related to *Ca. morgani*. Connected to the discovery of *Ca. pseudoreteaudii*, species in the *Ca. reteaudii* complex were re-considered and the group is shown to accommodate two cryptic species. These were collected in Australia and are described as *Ca. queenslandica* sp. nov. and *Ca. terrae-reginae* sp. nov.

Taxonomic novelties: *Calonectria cerciana* L. Lombard, M.J. Wingf. & Crous sp. nov., *Calonectria pseudoreteaudii* L. Lombard, M.J. Wingf. & Crous sp. nov., *Calonectria queenslandica* L. Lombard, M.J. Wingf. & Crous sp. nov., *Calonectria terrae-reginae* L. Lombard, M.J. Wingf. & Crous sp. nov.

INTRODUCTION

Species of *Calonectria* (*Ca.*), and their *Cylindrocladium* (*Cy.*) anamorphs, are important plant pathogens worldwide (Crous 2002). Past taxonomic studies on these pathogens have focused on morphology and sexual compatibility to delimit new species (Peerally 1991, Crous & Wingfield 1994). More recently, DNA sequence comparisons have resulted in the recognition of several species complexes in *Calonectria* (Schoch *et al.* 1999, Crous *et al.* 2004, 2006a).

One of the newly recognised groups in *Calonectria* is the *Ca. reteaudii* complex (Crous & Kang 2001, Kang *et al.* 2001). The complex encompasses several *Calonectria* spp. with *Cylindrocladium* anamorphs morphologically similar to the *Ca. reteaudii* anamorph state, having clavate vesicles with multiseptate macroconidia. These include *Cy. angustatum*, *Cy. hurae*, *Ca. leguminum* and *Ca. rumohrae* (Crous 2002).

The *Ca. morganii* species complex (Crous *et al.* 1993, Schoch *et al.* 2001) includes *Ca. morganii*, *Ca. insularis* and *Cy. hawksworthii* (Schoch *et al.* 1999, 2001, Crous 2002). Species in this complex are characterised by ellipsoidal to obpyriform to clavate vesicles, 1-septate conidia and orange to red perithecia producing 1-septate ascospores (Peerally 1991, Schoch *et al.* 1999, Crous 2002).

Species in both the *Ca. reteaudii* and the *Ca. morganii* complexes are responsible for a wide variety of disease symptoms on several plant hosts in sub-tropical and tropical regions of the world (Bolland *et al.* 1985, Peerally 1991, Sharma & Mohanan 1991, Booth *et al.* 2000, Crous 2002, Rodas *et al.* 2005). Disease symptoms include leaf blight (Sharma & Mohanan 1991, Booth *et al.* 2000, Rodas *et al.* 2005) and cutting rot (Sharma & Mohanan 1982, Sharma *et al.* 1984, Schoch *et al.* 1999, Crous 2002). Of these, leaf blight is most devastating in the tropical regions of South-East Asia and South America and is particularly serious on *Eucalyptus* spp. (Booth *et al.* 2000, Crous & Kang 2001, Crous 2002, Rodas *et al.* 2005).

Decline in *Eucalyptus* hybrid cutting production due to cutting rot has recently been observed in a commercial forest nursery in the GuangDong Province of China. Initial investigations indicated that the causal agents were unknown *Calonectria* spp. that represented species in the *Ca. reteaudii* and *Ca. morganii* complexes (Zhou *et al.* 2008). The aim of this study was to identify these *Calonectria* spp. using morphological characteristics and phylogenetic

inference. In addition, the taxonomic status and circumscription of species in the *Ca. reteaudii* species complex were re-considered.

MATERIALS AND METHODS

Isolates

Hybrid clonal *Eucalyptus* cuttings showing symptoms of cutting rot were collected from the commercial forestry nursery of the China Eucalypt Research Centre (CERC) in GuangDong Province, China. Diseased cuttings were placed in moist chambers and incubated for 48 h at room temperature to induce sporulation. Direct isolations were made onto malt extract agar (2 % w/v; MEA; Biolab, Midrand, South Africa) and cultures were incubated for 7 d at 25 °C under continuous near-ultraviolet light. For each isolate, single conidial cultures were prepared on MEA and representative strains are maintained in the culture collection (CMW) of the Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria, South Africa, the Centraalbureau voor Schimmelcultures (CBS), Utrecht, The Netherlands, and CERC, ZhanJiang Province, China. Isolates of *Ca. reteaudii* were obtained from the culture collection of CBS, including representative isolates used in the study of Kang *et al.* (2001).

DNA sequence comparisons

Calonectria isolates were grown on MEA for 7 d. Mycelium was then scraped from the surfaces of Petri dishes, freeze-dried, and ground to a powder in liquid nitrogen, using a mortar and pestle. DNA was extracted from the powdered mycelium as described by Lombard *et al.* (2008). Three loci were amplified and sequenced. These included a fragment of the β -tubulin (BT) gene region using primers T1 (O'Donnell & Cigelnik 1997) and CYLTUB1R (Crous *et al.* 2004), a fragment of the histone H3 (HIS3) gene region using primers CYLH3F and CYLH3R (Crous *et al.* 2004) and a fragment of the translation elongation factor-1 α (TEF-1 α) gene region using primers EF1-728F (Carbone & Kohn 1999) and EF2 (O'Donnell *et al.* 1998). The PCR reaction mixture used to amplify the different loci consisted of 2.5 units FastStart *Taq* polymerase (Roche Applied Science, USA), 10 \times PCR buffer, 1–1.5 mM MgCl₂, 0.25 mM of each dNTP, 0.5 μ M of each primer and approximately 30 ng of fungal genomic DNA, made up to a total reaction volume of 25 μ L with sterile distilled water.

Amplified fragments were purified using High Pure PCR Product Purification Kit (Roche, USA) and sequenced in both directions. For this purpose, the BigDye terminator sequencing kit v. 3.1 (Applied Biosystems, USA) and an ABI PRISMTM 3100 DNA sequencer (Applied Biosystems) were used. All PCRs and sequencing reactions were performed on an Eppendorf Mastercycler Personal PCR (Eppendorf AG, Germany) with cycling conditions as described in Crous *et al.* (2004, 2006a) for each locus.

Generated sequences were added to other sequences obtained from GenBank (<http://www.ncbi.nlm.nih.gov>) and these were assembled and aligned using Sequence Navigator v. 1.0.1 (Applied Biosystems) and MAFFT v. 5.11 (Katoh *et al.* 2005), respectively. The aligned sequences were then manually corrected where needed.

To determine whether the sequence data sets for the separate loci are congruent, tree topologies of 70 % reciprocal Neighbour-Joining bootstrap with Maximum Likelihood distances were compared visually to identify conflicts between partitions (Mason-Gamer & Kellogg 1996, Gueidan *et al.* 2007). Molecular evolution models for the separate partitions were determined in Modeltest v. 3.7 (Posada & Crandall 1998) and bootstrap analyses were run for 10k replicates.

PAUP (Phylogenetic Analysis Using Parsimony, v. 4.0b 10, Swofford 2002) was used to analyse the DNA sequence datasets. Phylogenetic relationships were estimated by heuristic searches with 1 000 random addition sequences and tree bisection-reconnection was used, with the branch swapping option set on 'best trees' only.

All characters were weighted equally and alignment gaps were treated as missing data. Measures calculated for parsimony included tree length (TL), consistency index (CI), retention index (RI) and rescaled consistence index (RC). Bootstrap analysis (Hillis & Bull 1993) was based on 1 000 replications. All sequences for the isolates studied were analysed using the Basic Local Alignment Search Tool for Nucleotide sequences (BLASTN, Altschul *et al.* 1990). The phylogenetic analysis was done on two separate sequence data sets.

Datasets were separated based on morphological characteristics to allow combinations of gene regions. The first dataset included 40 partial gene sequences per gene, representing 13 *Calonectria* spp. that form part of the *Ca. reteaudii* species complex and other morphologically similar species (Table 1). These included isolates used by Kang *et al.*

(2001), and are *Calonectria* species with anamorph states producing large, multiseptate macroconidia and stipe extensions terminating in clavate vesicles. The second data set consisted of 20 partial gene sequences per gene, representing 12 *Calonectria* spp. of the *Ca. morganii* species complex, and other morphologically similar species. This group of *Calonectria* species is characterised by smaller, 1–3-septate macroconidia, and stipe extensions terminating in ellipsoidal to obpyriform vesicles. *Ca. colombiensis* (CBS 112221) and *Cy. chinense* (CBS 112744) were used as outgroup taxa in both analyses (Lombard *et al.* 2009). All sequences were deposited in GenBank and the alignments in TreeBASE (<http://www.treebase.org>).

A Markov Chain Monte Carlo (MCMC) algorithm was used to generate phylogenetic trees with Bayesian probabilities using MrBayes v. 3.1.1 (Ronquist & Huelsenbeck 2003) for the combined sequence datasets. Models of nucleotide substitution for each gene were determined using MrModeltest v. 2.3 (Nylander 2004) and included for each gene partition. Three analyses of four MCMC chains were run from random trees for one million generations and sampled every 100 generations. All runs converged on the same likelihood score and tree topology and therefore the first 7600 trees for the *Ca. reteaudii* complex and 2000 trees for the *Ca. morganii* complex were discarded as the burn-in phase of each analysis and posterior probabilities determined from the remaining trees.

Sexual compatibility

A total of 29 single conidial *Ca. reteaudii*-like isolates (Table 1), originating from various geographical regions were crossed in all possible combinations including mating tester isolates CBS 112144 (+) and CBS 112147 (-) (Kang *et al.* 2001). Crosses were made as described in Schoch *et al.* (1999) on carnation leaf agar (CLA; Fisher *et al.* 1982, Crous *et al.* 1993). Control inoculations consisted of isolates crossed with themselves to determine whether they had a heterothallic or homothallic mating system. The plates were stacked in plastic containers and incubated at 22 °C for 6 wk. Crosses were regarded as successful when isolate combinations produced perithecia extruding viable ascospores.

Taxonomy

For morphological identification of *Calonectria* isolates, single conidial cultures were prepared on MEA and synthetic nutrient-poor agar (SNA; Nirenburg 1981, Lombard *et al.* 2009). Inoculated plates were incubated at room temperature and examined after 7 d. Gross

morphological characteristics of the anamorph state were determined by mounting fungal structures in lactic acid and 30 measurements at $\times 1\ 000$ magnification were made for each isolate. The 95 % confidence levels were determined and extremes of conidial measurements are given in parentheses. For other structures, only extremes are presented. Optimal growth conditions for cultures were determined in the dark on MEA for each isolate, at temperatures ranging from 5–35 °C at 5 °C intervals. Colony colours were determined after 7 d on MEA at 25 °C in the dark, using the colour charts of Rayner (1970).

RESULTS

DNA phylogeny

Amplicons of approximately 500 bases (BT and TEF-1 α) and 450 bases (HIS3) were obtained for all isolates. The pooled sequence datasets for all three loci showed conflict in tree topology for the 70 % reciprocal bootstrap trees, with *Ca. spathulata* (CBS 112689) and *Ca. multiseptata* (CBS 112682) grouping within different clusters for all three gene regions considered (results not shown). This conflict was resolved by separating the sequence datasets into two datasets representing morphological and phylogenetically closely related species to *Ca. reteaudii* and *Ca. morgani*, and sequence datasets for BT, HIS3 and TEF-1 α were combined for the two separate datasets. Sequence alignments have been deposited in TreeBASE as SN4542.

The combined sequence dataset for isolates representing *Ca. reteaudii* and other closely related species consisted of 1540 characters, of which 926 were constant, 178 were parsimony uninformative and 436 were parsimony informative. Parsimony analysis of the alignment yielded six most parsimonious trees (TL = 1316 steps; CI = 0.703; RI = 0.782; RC = 0.549), one of which is presented in Fig. 1. For Bayesian analyses, a HKY+G model was selected for BT, GTR+I+G for HIS3 and GTR+G for TEF-1 α , and incorporated in the analyses. The consensus tree obtained from Bayesian analyses confirmed the tree topology obtained with parsimony as well as bootstrap support (Fig. 1).

The isolates obtained from the *Eucalyptus* cuttings grouped in the *Ca. reteaudii* cluster, which formed a monophyletic group with a bootstrap value (BP) of 100 and a Bayesian posterior probability (PP) value of 1.00 (Fig. 1). This cluster segregated into two separate clades. The first of these, with a BP of 100 and PP value of 1.00 included the isolates from the *Eucalyptus* cuttings in a clade (BP = 100, PP = 1.00) separate from *Ca. reteaudii*,

possibly representing a distinct species. The second clade (BP = 100, PP = 0.99) was comprised of two sub-clades (BP = 79, PP = 1.00 and BP = 62, PP = 0.86, respectively) representing isolates from the study of Kang *et al.* (2001), and also suggested the existence of distinct species.

For the isolates that are closely related to *Ca. morganii*, the combined dataset consisted of 1518 characters. Of these characters, 969 were constant, 161 were parsimony uninformative and 388 were parsimony informative. Parsimony analysis of the alignment yielded three most parsimonious trees (TL = 1065 steps; CI = 0.736; RI = 0.783; RC = 0.577), one of which is represented in Fig. 2. For Bayesian analyses, a HKY+G model was selected for BT, GTR+I+G for HIS3, and GTR+G for TEF-1 α , and incorporated in the analyses. The consensus tree obtained from Bayesian analyses confirmed the tree topology obtained with parsimony as well as bootstrap support (Fig. 1).

In the tree (Fig. 2), isolates from the *Eucalyptus* cuttings grouped in the cluster representing species in the *Ca. morganii* complex (BP = 89, PP = 0.95), and some in the cluster representing *Ca. pauciramosa* (BP = 100, PP = 1.00). However, those isolates grouping in the *Ca. morganii* cluster formed a separate clade (BP = 100, PP = 1.00), suggesting that they might represent a distinct species.

Sexual compatibility

Protoperithecia formed within 3 wk and mating tests produced viable perithecia within 6 wk on CLA. Mating isolates CBS 112146, CBS 112151, CBS 112155 and CBS 112634 with isolates of *Ca. reteaudii* (represented in the current phylogenetic study), failed to produce perithecia in any combination tested. Crossing isolates CMW 25284, CMW 25285, CMW 25292 and CMW 25310 with isolates of *Ca. reteaudii* also failed to produce perithecia. However, crossed isolates of *Ca. reteaudii* produced perithecia with viable ascospores as reported by Kang *et al.* (2001). Isolate CBS 582.50 did not cross with any of the other *Ca. reteaudii* isolates, possibly due to loss of fertility.

Taxonomy

Morphological observations and DNA sequence comparisons (Fig. 2) showed that isolates CMW 25311 and CMW 25283 clearly represent the anamorph state of *Ca. pauciramosa*. Isolates CMW 25309, CMW 25290 and CMW 25288 represent an undescribed species closely related to other species in the *Ca. morganii* species complex, and it is consequently

described as new. Isolates CMW 25310, CMW 25292, CMW25285 and CMW 25284 obtained from the *Eucalyptus* cuttings are morphologically very similar to the anamorph state of *Ca. reteaudii*. However, some morphological differences were found and this fungus is also treated as a new species. Based on DNA sequence analysis, mating strategies and morphological observations, isolates CBS 112146 (= CPC 3213) and CBS 112155 (= CPC 3210), which were previously regarded as *Ca. reteaudii* (Kang *et al.* 2001), are shown to represent a distinct species. A similar situation was found for isolates CBS 112151 (= CPC 3202) and CBS 112634 (= CPC 4233), and they are also treated as a new species.

Calonectria cerciana L. Lombard, M.J. Wingf. & Crous, **sp. nov.** MycoBank 513263. Fig. 3

Etymology: Name refers to the China Eucalypt Research Centre (CERC), a research institution that is pioneering the study of *Eucalyptus* diseases in China.

Stipa extensiones septatae, rectae vel flexuosae, 148–222 μm longae, ad septum apicale 5–6 μm latae, vesiculo fusiforme vel obpyriforme 8–13 μm diametro terminantes. *Macroconidia* cylindrica, in extremitatibus ambabus rotundata, recta (37–)41–46(–49) \times 5–6 μm , 1-septata.

Teleomorph unknown. *Conidiophores* with a stipe bearing a suit of penicillate, fertile branches, a stipe extension, and a terminal vesicle; stipe septate, hyaline, smooth, 48–90 \times 6–10 μm ; stipe extensions septate, straight to flexuous, 148–222 μm long, 5–6 μm wide at the apical septum, terminating in fusiform to obpyriform vesicles, 8–13 μm diam. *Conidiogenous apparatus* 70–98 μm long, and 62–113 μm wide; primary branches aseptate or 1-septate, 21–31 \times 5–7 μm ; secondary branches aseptate, 15–22 \times 4–5 μm ; tertiary and additional branches (–4), aseptate, 10–20 \times 4–5 μm , each terminal branch producing 2–6 phialides; phialides doliiform to reniform, hyaline, aseptate, 9–12 \times 3 μm ; apex with minute periclinal thickening and inconspicuous collarete. *Macroconidia* cylindrical, rounded at both ends, straight, (37–)41–46(–49) \times 5–6 μm (av. = 44 \times 5 μm), 1-septate, lacking a visible abscission scar, held in parallel cylindrical clusters by colourless slime. *Mega-* and *microconidia* not seen.

Specimens examined: **China**, GuangDong Province, CERC nursery, on stems of *E. urophylla* \times *grandis* hybrid cutting, Nov. 2007, M.J. Wingfield & X.D. Zhou, Herb. PREM 60241, **holotype** of *Calonectria cerciana*, culture ex-type CMW 25309 = CBS 123693; GuangDong Province, CERC nursery, on stems of *E. urophylla* \times *grandis* hybrid cutting, Nov. 2007, M.J. Wingfield & X.D. Zhou (Herb. PREM 60242), culture CMW 25290 = CBS 123695; CERC

nursery, on stems of *E. urophylla* × *grandis* hybrid cutting, Nov. 2007, M.J. Wingfield & X.D. Zhou, culture CMW 25288.

Culture characteristics: Colonies fast growing with optimal growth temperature at 25 °C (growth at 15–30 °C) on MEA, reverse sepia-brown (13i) after 7 d; abundant white aerial mycelium and sporulation; chlamydospores abundant throughout the medium, forming microsclerotia.

Substrate: *Eucalyptus urophylla* × *grandis* hybrid cuttings

Distribution: China

Notes: *Calonectria cerciana* can be distinguished from *Cy. hawksworthii* (conidia av. 56 × 4 μm), *Ca. morgani* (conidia av. 45 × 4 μm) and *Ca. insularis* (conidia av. 45 × 4 μm) based on its fusoid to ellipsoidal vesicles. Macroconidia of *Ca. cerciana* are also slightly smaller (av. 44 × 5 μm) than those of the above-mentioned species.

Calonectria pseudoreteauidii L. Lombard, M.J. Wingf. & Crous, **sp. nov.** MycoBank 513264. Fig. 4

Etymology: Name reflects the fact that the species resembles the anamorph state of *Ca. reteauidii*.

Stipa extensiones septatae, rectae vel flexuosae, 193–313 μm longae, ad septum apicale 5–6 μm latae, vesiculo anguste clavato, 3–5 μm diametro terminantes. *Macroconidia* cylindrica, apice rotundata, basi complanata, recta (88–96–112(–119) × 7–9(–10) μm, 5–8-septata. *Microconidia* cylindrica, apice rotundata, basi complanata, (30–)34–54(–68) × 3–5(–6) μm, 1–3-septata, cum muco in glomerulis.

Teleomorph unknown. *Macroconidiophores* with a stipe bearing a suite of penicillate fertile branches, a stipe extension, and a terminal vesicle; stipe septate, hyaline, smooth, 43–111 × 5–9 μm; stipe extensions septate, straight to flexuous, 193–313 μm long, 5–6 μm wide at the apical septum, terminating in a narrowly clavate vesicle, 3–5 μm diam. *Conidiogenous apparatus* 45–103 μm long and 26–82 μm wide; primary branches aseptate or 1-septate, 29–42 × 5–6 μm; secondary branches aseptate, 20–36 × 3–6 μm; tertiary branches aseptate, 15–24 × 4–5 μm, each terminal branch producing 1–3 phialides; phialides cylindrical to allantoid, obpyriform when carried singly, hyaline, aseptate, 16–25 × 3–5 μm; apex with minute periclinal thickening and inconspicuous collarette. *Macroconidia* cylindrical, rounded at the apex, flattened at the base, straight, (88–)96–112(–119) × 7–9(–10) μm (av. = 104 × 8

µm), 5–8-septate, lacking a visible abscission scar, held in parallel cylindrical clusters by colourless slime. *Microconidiophores* simple with some lateral branching, comprising a stipe and a penicillate or subverticillate arrangement of fertile branches. *Stipe* septate, hyaline, thin-walled, $27\text{--}70 \times 2\text{--}3$ µm; primary branches aseptate, subcylindrical, straight to curved, $14\text{--}19 \times 3\text{--}4$ µm, terminating in 1–3 phialides that are straight to slightly curved, $11\text{--}18 \times 3\text{--}4$ µm; apex with minute periclinal thickening and collarete. *Microconidia* cylindrical, straight, rounded at the apex, flattened at the base, $(30\text{--})34\text{--}54(\text{--}68) \times 3\text{--}5(\text{--}6)$ µm (av. 44×4 µm), 1–3-septate, held in fascicles by colourless slime. *Megaconidia* not seen.

Specimens examined: **China**, GuangDong Province, CERC nursery, on stems of *E. urophylla* × *grandis* hybrid cutting, Nov. 2007, M.J. Wingfield & X.D. Zhou, Herb. PREM 60290, **holotype** of *Calonectria pseudoreteaudii*, culture ex-type CMW 25310 = CBS 123694; GuangDong Province, CERC nursery, on stems of *E. urophylla* × *grandis* hybrid cutting, Nov. 2007, M.J. Wingfield & X.D. Zhou (Herb. PREM 60291), culture CMW 25292 = CBS 123696; CERC nursery, on stems of *E. urophylla* × *grandis* hybrid cutting, Nov. 2007, M.J. Wingfield & X.D. Zhou, culture CMW 25284; CERC nursery, on stems of *E. urophylla* × *grandis* hybrid cutting, Nov. 2007, M.J. Wingfield & X.D. Zhou, culture CMW 25285.

Culture characteristics: Colonies fast growing with optimal growth temperature at 25 °C (growth at 15–30 °C) on MEA, reverse amber (13k) to sepia-brown (13i) after 7 d; abundant white aerial mycelium with moderate sporulation; chlamydospores abundant throughout the medium, forming microsclerotia.

Substrate: *Eucalyptus urophylla* × *grandis* hybrid cuttings

Distribution: China

Notes: The anamorph state of *Ca. pseudoreteaudii* can be distinguished from that of *Ca. reteaudii* based on its larger macroconidia (av. = 104×8 µm vs. 84×6.5 µm), as well as larger microconidia (av. = 44×4 µm vs. 30×3 µm). The microconidiophores of *Ca. pseudoreteaudii* do not produce stipe extensions, a feature which is common in *Ca. reteaudii*.

Calonectria queenslandica L. Lombard, M.J. Wingf. & Crous, **sp. nov.** MycoBank 513265.
Fig. 5

Etymology: Name refers to Queensland, Australia where the fungus was collected.

Stipa extensiones septatae, rectae vel flexuosae, 105–156 μm longae, ad septum apicale 4–5 μm latae, vesiculo anguste clavato 3–4 μm diametro terminantes. *Macroconidia* cylindrica, in extremitatibus ambabus rotundata, recta (61–)65–73(–78) \times (4–)5–6(–7) μm , 4–6-septata.

Teleomorph unknown. *Conidiophores* consisting of a stipe bearing a suit of penicillate fertile branches, a stipe extension, and a terminal vesicle; stipe septate, hyaline, smooth, 38–58 \times 6–7 μm ; stipe extensions septate, straight to flexuous, 105–156 μm long, 4–5 μm wide at the apical septum, terminating in narrowly clavate vesicle, 3–4 μm diam. *Conidiogenous apparatus* 39–64 μm long, and 27–68 μm wide; primary branches aseptate or 1-septate, 14–26 \times 4–6 μm ; secondary branches aseptate, 11–22 \times 3–5 μm ; tertiary branches aseptate, 13–17 \times 3–5 μm , each terminal branch producing 1–3 phialides; cylindrical to allantoid, obpyriform when carried singly, hyaline, aseptate, 10–16 \times 3–5 μm ; apex with minute periclinal thickening and inconspicuous collarette. *Macroconidia* cylindrical, rounded at both ends, straight, (61–)65–73(–78) \times (4–)5–6(–7) μm (av. = 69 \times 6 μm), 4–6-septate, lacking a visible abscission scar, held in parallel cylindrical clusters by colourless slime. *Mega-* and *microconidia* not seen.

Specimens examined: **Australia**, Queensland, Lannercost, on leaves of *E. urophylla*, 15 Apr. 1991, B. Brown, Herb. PREM 60243, **holotype** of *Calonectria queenslandica*, culture ex-type CMW 30604 = CBS 112146 = CPC 3213 = DFRI00147; Queensland, Lannercost, on leaves of *E. pellita*, 10 Mar. 1999, P.Q. Thu & K.M. Old (Herb. PREM 60244), culture CMW 30603 = CBS 112155 = CPC 3210 = DFRI00172.

Culture characteristics: Colonies fast growing, with optimal growth temperature at 25 °C (growth at 15–30 °C) on MEA; reverse sepia-brown (13i) after 7 d; abundant white aerial mycelium and sporulation; chlamydospores abundant throughout the medium, forming microsclerotia.

Substrate: *Eucalyptus urophylla*, *E. pellita*.

Distribution: Australia

Notes: *Calonectria queenslandica* can be distinguished from *Ca. reteaudii* and *Ca. pseudoreteaudii* based on its smaller macroconidia (av. = 69 \times 6 μm) and shorter stipe extensions of the anamorph state. No microconidiophores were observed in *Ca.*

queenslandica, although *Ca. reteaudii* and *Ca. pseudoreteaudii* readily produce these structures on SNA.

Calonectria terrae-reginae L. Lombard, M.J. Wingf. & Crous, *sp. nov.* MycoBank 513266.

Fig. 6

Etymology: Name refers to Queensland, Australia, from where this fungus was isolated.

Stipa extensiones septatae, rectae vel flexuosae, 127–235 μm longae, ad septum apicale 4–6 μm latae, vesiculo anguste clavato 3–5 μm diametro terminantes. *Macroconidia* cylindrica, in extremitatibus ambabus rotundata, recta, 60–83(–87) \times (4–)5–7(–8) μm , 4–6-septata.

Teleomorph unknown. *Conidiophores* consisting of a stipe bearing a suite of penicillate fertile branches, a stipe extension, and a terminal vesicle; stipe septate, hyaline, smooth, 58–86 \times 4–8 μm ; stipe extensions septate, straight to flexuous, 127–235 μm long, 4–6 μm wide at the apical septum, terminating in narrowly clavate vesicle, 3–5 μm diam. *Conidiogenous apparatus* 35–54 μm long, and 33–48 μm wide; primary branches aseptate, 16–25 \times 4–6 μm ; secondary branches aseptate, 13–18 \times 3–6 μm ; tertiary branches aseptate, 10–14 \times 3–5 μm , each terminal branch producing 1–3 phialides; cylindrical to allantoid, obpyriform when carried singly, hyaline, aseptate, 10–17 \times 2–4 μm ; apex with minute periclinal thickening and inconspicuous collarette. *Macroconidia* cylindrical, rounded at both ends, straight, 60–83(–87) \times (4–)5–7(–8) μm (av. = 76 \times 6) μm , 4–6-septate, lacking a visible abscission scar, held in parallel cylindrical clusters by colourless slime. *Megaconidia* and *microconidia* not seen.

Specimens examined: **Australia**, Queensland, Cardwell, Meunga, on leaves of *E. urophylla*, 11 Apr. 1997, C. Hanwood, Herb. PREM 60239, **holotype** of *Calonectria terrae-reginae*, culture ex-type CMW 30601 = CBS 112151 = CPC 3202 = DFRI00150; Victoria, on *Xanthorrhoea australis*, T. Baigent (Herb. PREM 60240), culture CMW 30602 = CBS 112634 = CPC 4233 = Lynfield 417.

Culture characteristics: Colonies fast growing with optimal growth temperature at 25 °C (growth at 15–30 °C) on MEA; reverse sepia-brown (13i) after 7 d; abundant white aerial mycelium and sporulation; chlamydospores abundant throughout the media, forming microsclerotia.

Substrate: *Eucalyptus urophylla*, *Xanthorrhoea australis*

Distribution: Australia

Notes: *Calonectria terrae-reginae* is distinct from *Ca. queenslandica* in having larger macroconidia (av.= $76 \times 6 \mu\text{m}$), although they are smaller than those of *Ca. reteaudii* and *Ca. pseudoreteaudii*.

DISCUSSION

This study emerged from the collection of isolates of *Calonectria* spp. from infected *Eucalyptus* cuttings in the GuangDong Province of China. The isolates were shown to represent three species including *Ca. pauciramosa*, and two new species that have been described as *Ca. cerciana* and *Ca. pseudoreteaudii*. The former species is related to taxa in the *Ca. morganii* complex, while the latter species resides in the *Ca. reteaudii* complex.

Taxonomic placement of *Ca. pseudoreteaudii* required a re-evaluation of the *Ca. reteaudii* complex. It was consequently found that the group has been poorly defined, and that it encompasses a number of cryptic species. This resulted in the description of *Ca. pseudoreteaudii*, *Ca. queenslandica* and *Ca. terrae-reginae* as three new sibling species in the *Ca. reteaudii* complex based on phylogenetic inference and morphological comparisons with the ex-type culture of *Ca. reteaudii* (CBS 112144). Species in the *Ca. reteaudii* complex are important pathogens of *Eucalyptus* spp. causing *Cylindrocladium* leaf blight and cutting rot in Australia, South-East Asia and South America (Pikethley 1976, Bolland 1985, Sharma & Mohanan 1991, Sharma & Mohanan 1992, Booth *et al.* 2000, Crous & Kang 2001, Crous 2002, Rodas *et al.* 2005) and their refined taxonomy presented here will contribute to efforts to manage this disease.

Discovery of *Ca. queenslandica* and *Ca. terrae-reginae* was serendipitous as isolates used in the phylogenetic component of the study were largely the same as those used in a previous study by Kang *et al.* (2001). Although the latter study focused on the taxonomic position of *Cy. quinqueseptatum* (= *Cy. reteaudii*) and *Ca. quinqueseptata* (= *Ca. leguminum*), it employed a single gene region, and could thus not adequately define the variation in *Ca. reteaudii*, which was later recognised in multi-gene analyses (Crous *et al.* 2006a). Mating tests undertaken by Kang *et al.* (2001) indicated that 15 of the 20 *Ca. reteaudii* isolates used were capable of producing viable progeny. Mating tests conducted in the present study were unsuccessful in crossing strains of *Ca. queenslandica*, *Ca. terrae-reginae*, *Ca. reteaudii* and

Ca. pseudoreteauidi, and only isolates of *Ca. reteaudii* could be induced to produce perithecia with viable ascospores.

Calonectria pauciramosa (anamorph: *Cy. pauciramosum*) is a well-known pathogen in *Eucalyptus* cutting nurseries (Schoch *et al.* 1999, Crous 2002). This pathogen resides in the *Ca. scoparia* species complex and is regarded as the dominant nursery pathogen of various host plants in countries such as Australia, Italy, South Africa and the U.S.A. (Koike *et al.* 1999, Polizzi & Crous 1999, Schoch *et al.* 1999, 2001, Koike & Crous 2001). The present study represents the first report of this pathogen in China, but pathogenicity tests and disease surveys will be required to determine its relevance in that country.

The description of *Ca. cerciana* from *Eucalyptus* cuttings adds a new species to the *Ca. morganii* species complex. *Calonectria cerciana* can be distinguished from the other species in the complex based on its smaller macroconidia and the formation of fusiform to ellipsoidal vesicles. Crous (2002) and Schoch *et al.* (1999) found that there was a low level of fertility among species in this complex, and Schoch *et al.* (2001) used BT to show that they were closely related.

It is unknown whether *Ca. cerciana* and *Ca. pseudoreteauidi* are pathogens of *Eucalyptus*. Other species in the *Ca. reteaudii* (Sharma & Mohanan 1982, Sharma & Mohanan 1991, Sharma & Mohanan 1992) and *Ca. morganii* (Mohanan & Sharma 1985, Crous 2002) species complexes are known to be *Eucalyptus* pathogens, and this is probably also true for *Ca. cerciana* and *Ca. pseudoreteauidi*. However, the pathogenicity of these two *Calonectria* spp. must be tested and these studies would logically also consider the susceptibility of different *Eucalyptus* clones and hybrids being deployed in plantations.

Although no teleomorph states for the four newly described *Calonectria* spp. could be induced in this study, they have all been placed in *Calonectria*, and not in the anamorph genus *Cylindrocladium* (Lombard *et al.* 2009). The decision to use the oldest generic name for a well-defined clade of fungi (*Calonectria*) is consistent with the approach taken previously by Lombard *et al.* (2009), and has also been followed in other groups of fungi such as *Botryosphaeriaceae* (Crous *et al.* 2006b, 2008, Phillips *et al.* 2008), *Mycosphaerellaceae* and *Teratosphaeriaceae* (Crous *et al.* 2009a, 2009b) to name a few. Although it might be considered a broad interpretation, it is allowed by the International Code of Botanical Nomenclature (Hawksworth 2005, McNeill *et al.* 2005). Based on these



regulations and the fact that the anamorph genus *Cylindrocladium* is linked to the single teleomorph genus *Calonectria* (Rossman *et al.* 1999), all new species have been accommodated in *Calonectria*.

REFERENCES

- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. (1990). Basic Local Alignment Search Tool. *Journal of Molecular Biology* **215**: 403–410.
- Bolland L, Tierney JW, Tierney BJ. (1985). Studies on leaf spot and shoot blight of *Eucalyptus* caused by *Cylindrocladium quinqueseptatum*. *European Journal of Forest Pathology* **15**: 385–397.
- Booth TH, Jovanovic T, Old KM, Dudzinski MJ. (2000). Climatic mapping to identify high-risk areas for *Cylindrocladium quinqueseptatum* leaf blight on eucalypts in main land South East Asia and around the world. *Environmental Pollution* **108**: 365–372.
- Carbone I, Kohn LM. (1999). A method for designing primer sets for speciation studies in filamentous ascomycetes. *Mycologia* **91**: 553–556.
- Crous PW. (2002). Taxonomy and pathology of *Cylindrocladium* (*Calonectria*) and allied genera. APS Press, St. Paul, Minnesota, USA.
- Crous PW, Alfenas AC, Wingfield MJ. (1993). *Calonectria scoparia* and *Calonectria morganii* sp. nov., and variations among isolates of their *Cylindrocladium* anamorphs. *Mycological Research* **97**: 701–708.
- Crous PW, Groenewald JZ, Risède J-M, Simoneau P, Hyde KD. (2006a). *Calonectria* species and their *Cylindrocladium* anamorphs: species with clavate vesicles. *Studies in Mycology* **55**: 213–226.
- Crous PW, Groenewald JZ, Risède J-M, Simoneau P, Hywel-Jones N. (2004). *Calonectria* species and their *Cylindrocladium* anamorphs: species with sphaeropedunculate vesicles. *Studies in Mycology* **50**: 415–430.
- Crous PW, Kang J-C. (2001). Phylogenetic confirmation of *Calonectria spathulata* and *Cylindrocladium leucothoes* based on morphology, and sequence data of the β -tubulin and ITS rRNA genes. *Mycoscience* **42**: 51–57.
- Crous PW, Slippers B, Wingfield MJ, Rheeder J, Marasas WFO, Phillips AJL, Alves A, Burgess T, Barber P, Groenewald JZ. (2006b). Phylogenetic lineages in the *Botryosphaeriaceae*. *Studies in Mycology* **55**: 235–253.
- Crous PW, Summerell BA, Carnegie AJ, Wingfield MJ, Groenewald JZ. (2009a). Novel species of *Mycosphaerellaceae* and *Teratosphaeriaceae*. *Persoonia* **23**: 119–146.

- Crous PW, Summerell BA, Carnegie AJ, Wingfield MJ, Hunter GC, Burgess TI, Andjic V, Barber PA, Groenewald JZ. (2009b). Unravelling *Mycosphaerella*: do you believe in genera? *Persoonia* **23**: 99–118.
- Crous PW, Wingfield MJ. (1994). A monograph of *Cylindrocladium*, including anamorphs of *Calonectria*. *Mycotaxon* **51**: 341–435.
- Crous PW, Wood AR, Okada G, Groenewald JZ. (2008). Foliicolous microfungi occurring on *Encephalartos*. *Persoonia* **21**: 135–146.
- Fisher NL, Burgess LW, Toussoun TA, Nelson PE. (1982). Carnation leaves as a substrate and for preserving cultures of *Fusarium* species. *Phytopathology* **72**: 151–153.
- Gueidan C, Roux C, Lutzoni F. (2007). Using multigene phylogeny analysis to assess generic delineation and character evolution in Verrucariaceae (Verrucariales, Ascomycota). *Mycological Research* **111**: 1145–1168.
- Hawksworth DL. (2005). Two major changes in fungal nomenclature enacted in Vienna. *Mycological Research* **109**: 1061–1062.
- Hillis DM, Bull JJ. (1993). An empirical test of bootstrapping as a method for assessing confidence in phylogenetic analysis. *Systematic Biology* **42**: 182–192.
- Kang J-C, Crous PW, Old KM, Dudzinski MJ. (2001). Non-conspecificity of *Cylindrocladium quinqueseptatum* and *Calonectria quinqueseptata* based on a β -tubulin gene phylogeny and morphology. *Canadian Journal of Botany* **79**: 1241–1247.
- Katoh K, Kuma K, Toh H, Miyata T. (2005). MAFFT version 5: improvement in accuracy of multiple sequence alignment. *Nucleic Acid Research* **33**: 511–518.
- Koike ST, Crous PW. (2001). First report of a root and crown rot disease of myrtle in California caused by *Cylindrocladium pauciramosum*. *Plant Disease* **85**: 448.
- Koike ST, Henderson DM, Crous PW, Schoch CL, Tjosvold SA. (1999). A new root and crown rot disease of heath in California caused by *Cylindrocladium pauciramosum*. *Plant Disease* **83**: 589.
- Lombard L, Bogale M, Montenegro F, Wingfield BD, Wingfield MJ. (2008). A new bark canker disease of the tropical hardwood tree *Cedrelinga cateniformis* in Ecuador. *Fungal Diversity* **31**: 73–81.
- Lombard L, Rodas CA, Crous PW, Wingfield BD, Wingfield MJ. (2009). *Calonectria* (*Cylindrocladium*) species associated with dying *Pinus* cuttings. *Persoonia* **23**: 41–47.
- Mason-Gamer R, Kellogg E. (1996). Testing for phylogenetic conflict among molecular datasets in the tribe *Tiriceae* (*Graminae*). *Systematic Biology* **45**: 524–545.

- McNiell J, Stuessy TF, Turland NJ, Hörandl E. (2005). XVII International Botanical Congress: preliminary mail vote and report of Congress action on nomenclature proposals. *Taxon* **54**: 1057–1064.
- Mohanan C, Sharma JK. (1985). *Cylindrocladium* causing seedling diseases of *Eucalyptus* in Kerala, India. *Transactions of the British Mycological Society* **84**: 538–539.
- Nirenburg HI. (1981). A simplified method for identifying *Fusarium* spp. occurring on wheat. *Canadian Journal of Botany* **59**: 1599–1609.
- Nylander JAA. (2004). MrModeltest v.2. Programme distributed by the author. Evolutionary Biology Centre, Uppsala University.
- O'Donnell K, Cigelnik E. (1997). Two divergent intragenomic rDNA ITS2 types within a monophyletic lineage of the fungus *Fusarium* are nonorthologous. *Molecular Phylogenetics and Evolution* **7**: 103–116.
- O'Donnell K, Kistler HC, Cigelnik E, Plötz RC. (1998). Multiple evolutionary origins of the fungus causing Panama disease of banana: Concordant evidence from nuclear and mitochondrial gene genealogies. *Proceedings of the National Academy of Sciences of the United States of America* **95**: 2044–2049.
- Peerally A. (1991). The classification and phytopathology of *Cylindrocladium* species. *Mycotaxon* **40**: 323–366.
- Phillips AJL, Alves A, Pennycook SR, Johnston PR, Ramaley A, Akulov A, Crous PW. (2008). Resolving the phylogenetic and taxonomic status of dark-spored teleomorph genera in the Botryosphaeriaceae. *Persoonia* **21**: 29–55.
- Pikethley RN. (1976). *Cylindrocladium quinquesepatum* on myrtaceous tree seedlings. *Australian Plant Pathology Newsletter* **5**: 57.
- Polizzi G, Crous PW. (1999). Root and Collar Rot of Milkwort Caused by *Cylindrocladium pauciramosum*, a new record for Europe. *European Journal of Plant Pathology* **105**: 407–411.
- Posada D, Crandall KA. (1998). Modeltest: testing the model of DNA substitution. *Bioinformatics* **14**: 817–818.
- Rayner RW. (1970). A mycological colour chart. Commonwealth Mycological Institute, Kew, Surrey. British Mycological Society.
- Rodas CA, Lombard L, Gryzenhout M, Slippers B, Wingfield MJ. (2005). *Cylindrocladium* blight of *Eucalyptus* in Colombia. *Australasian Plant Pathology* **34**: 143–149.

- Ronquist F, Heulsenbeck JP. (2003). MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* **19**: 1572–1574.
- Rossmann AY, Samuels GJ, Rogerson CT, Lowen R. (1999). Genera of Bionectriaceae, Hypocreaceae and Nectriaceae (Hypocreales, Ascomycetes). *Studies in Mycology* **42**: 1–248.
- Schoch CL, Crous PW, Wingfield BD, Wingfield MJ. (1999). The *Cylindrocladium candelabrum* species complex includes four distinct mating populations. *Mycologia* **91**: 286–298.
- Schoch CL, Crous PW, Wingfield BD, Wingfield MJ. (2001). Phylogeny of *Calonectria* based on comparisons of β -tubulin DNA sequences. *Mycological Research* **105**: 1045–1052.
- Sharma JK, Mohanan C. (1982). *Cylindrocladium* spp. associated with various diseases of *Eucalyptus* in Kerala. *European Journal of Forest Pathology* **12**: 129–136.
- Sharma JK, Mohanan C. (1991). Pathogenic variation in *Cylindrocladium quinqueseptatum* causing leaf blight of *Eucalyptus*. *European Journal of Forest Pathology* **21**: 210–217.
- Sharma JK, Mohanan C. (1992). Relative susceptibility of *Eucalyptus* provenances to *Cylindrocladium* leaf blight in Kerala, India. *European Journal of Forest Pathology* **22**: 257–265.
- Sharma JK, Mohanan C, Florence EJM. (1984). Nursery diseases of *Eucalyptus* in Kerala. *European Journal of Forest Pathology* **14**: 77–89.
- Swofford DL. (2002). PAUP*. Phylogenetic analysis using parsimony (* and other methods), 4.0b10. Computer programme. Sunderland, Massachusetts, USA: Sinauer Associates.
- Zhou XD, Xie YJ, Chen SF, Wingfield MJ. (2008). Diseases of eucalypt plantations in China: challenges and opportunities. *Fungal Diversity* **32**: 1–7.



Table 1 Strains of *Calonectria* and *Cylindrocladium* used in the phylogenetic study.

Species	Isolate number ¹	GenBank accession no.			Host	Origin	Collector
		BT	HIS3	TEF-1 α			
<i>Cy. angustatum</i>	CBS 109169	DQ190593	DQ190695	FJ918552	<i>Tillandsia capitata</i>	U.S.A.	R.M. Leahy
	CBS 109065 ^T	AF207543	DQ190656	FJ918551	<i>T. capitata</i>	U.S.A.	R.M. Leahy
<i>Ca. cerciana</i> sp. nov. (= CBS 123693) ^T	CMW 25309	FJ918510	FJ918528	FJ918559	<i>Eucalyptus urophylla</i>	× China	M.J. Wingfield & X.D. Zhou
	(= CBS 123693) ^T				<i>grandis</i> cutting		
	CMW 25290 (= CBS 123695)	FJ918511	FJ918529	FJ918560	<i>E. urophylla</i>	× China	M.J. Wingfield & X.D. Zhou
	CMW 25288	GQ25288	GQ267243	GQ267288	<i>E. urophylla</i>	× China	M.J. Wingfield & X.D. Zhou
<i>Ca. colombiensis</i>	CBS112221	AY725620	AY725663	AY725712	Soil	Colombia	M.J. Wingfield
<i>Cy. chinense</i>	CBS112744	AY725618	AY725660	AY725709	Soil	China	E.C.Y. Liew
<i>Ca. brassicae</i>	CBS 111869 ^T	AF232857	DQ190720	FJ918567	<i>Argyrea</i> sp.	South East Asia	
	CBS 111478	DQ190611	DQ190719	FJ918567	Soil	Brazil	A.C. Alfenas
<i>Cy. hawksworthii</i>	CBS 111870 ^T	AF333407	DQ190649	FJ918558	<i>Nelumbo nucifera</i>	Mauritius	A. Peerally
<i>Cy. hurae</i>	CBS 114551	AF333408	DQ190728	FJ918548	<i>Rumhra</i>	Brazil	A.C. Alfenas
					<i>adiantiformis</i>		



Table 1. (Continued)

Species	Isolate number ¹	GenBank accession no.			Host	Origin	Collector
		BT	HIS3	TEF-1 α			
<i>Ca. insularis</i>	CBS 114558	AF210861	FJ918526	FJ918556	Soil	Madagascar	P.W. Crous
	CBS 114559	AF210862	FJ918525	FJ918555	Soil	Madagascar	C.L. Schoch
<i>Ca. leguminum</i>	CBS 728.68 ^T	AF389837	DQ190654	FJ918547	<i>Annona squamosa</i>	Brazil	M.B. Figueiredo
<i>Cy. leucothoës</i>	CBS 109166 ^T	FJ918508	FJ918523	FJ918553	<i>Leucothoë axillaris</i>	USA	N.E. El-Gholl
<i>Cy. multiseptatum</i>	CBS 112682 ^T	DQ190573	DQ190659	FJ918535	<i>Eucalyptus</i> sp.	Indonesia	M.J. Wingfield
<i>Ca. pauciramosa</i>	CMW 5683	FJ918514	FJ918531	FJ918565	<i>Eucalyptus</i> sp.	Brazil	A.C. Alfenas
	CMW 30823	FJ918515	FJ918532	FJ918566	<i>E. grandis</i>	South Africa	P.W. Crous
	CMW 25311	FJ918516	FJ918533	FJ918569	<i>E. urophylla</i> × <i>grandis</i> cutting	China	M.J. Wingfield & X.D. Zhou
	CMW 25283	FJ918517	FJ918534	FJ918570	<i>E. urophylla</i> × <i>grandis</i> cutting	China	M.J. Wingfield & X.D. Zhou
<i>Ca. pseudoreteaudii</i> sp. nov.	CMW 25310 (= CBS 123694) ^T	FJ918504	FJ918519	FJ918541	<i>E. urophylla</i> × <i>grandis</i> cutting	China	M.J. Wingfield & X.D. Zhou



Table 1. (Continued)

Species	Isolate number ¹	GenBank accession no.			Host	Origin	Collector
		BT	HIS3	TEF-1 α			
<i>Ca. pseudoreteaudii</i> sp. nov.	CMW 25292 (= CBS 123696)	FJ918505	FJ918520	FJ918542	<i>E. urophylla</i> × <i>grandis</i> cutting	China	M.J. Wingfield & X.D. Zhou
	CMW 25284	GQ267205	GQ267244	GQ267289	<i>E. urophylla</i> × <i>grandis</i> cutting	China	M.J. Wingfield & X.D. Zhou
	CMW 25285	GQ267206	GQ267245	GQ267290	<i>E. urophylla</i> × <i>grandis</i> cutting	China	M.J. Wingfield & X.D. Zhou
<i>Ca. pseudospathiphylli</i>	CBS 109165	FJ918513	AF348241	FJ918562	Soil	Ecuador	M.J. Wingfield
<i>Ca. pteridis</i>	CBS 111793	DQ190578	DQ190679	FJ918563	<i>Arachnoides</i> <i>adiantiformis</i>	USA	
	CBS 111871	DQ190579	DQ190680	FJ918564	<i>Pinus</i> sp.	Spain	T.L. Krugner
<i>Ca. queenslandica</i> sp. nov.	CMW 30604 (= CBS 112146 = CPC 3213 = DRFI000147) ^T	AF389835	FJ918521	FJ918543	<i>E. urophylla</i>	Australia	B. Brown
	CMW 30603 (= CBS 112155 = CPC 3210 = DFRI00172)	AF389834	DQ190667	FJ918544	<i>E. pellita</i>	Australia	K.M. Old



Table 1. (Continued)

Species	Isolate number ¹	GenBank accession no.			Host	Origin	Collector
		BT	HIS3	TEF-1 α			
<i>Ca. reteaudii</i>	CBS 582.50	AF389836	DQ190673	FJ918540	<i>Hibiscus sabdariffa</i>	Indonesia	—
	CBS 112143	GQ240642	DQ190660	FJ918536	<i>E. camaldulensis</i>	Vietnam	M.J. Dudzinski
	CBS 112144 ^T	AF389833	DQ190661	FJ918537	<i>E. camaldulensis</i>	Vietnam	M.J. Dudzinski
	CBS 112147	AF389830	DQ190663	FJ918539	<i>E. camaldulensis</i>	Vietnam	M.J. Dudzinski
	CBS 112153	AF389831	FJ918518	FJ918538	<i>E. camaldulensis</i>	Vietnam	M.J. Dudzinski
	CBS 113582	GQ240643	GQ240659	GQ240675	<i>Eucalyptus</i> sp.	Thailand	—
	CBS 113583	GQ240644	GQ240660	GQ240676	<i>Eucalyptus</i> sp.	Madagascar	P.W. Crous
	CMW 18446	GQ240649	GQ240665	GQ240681	<i>E. urophylla</i>	Indonesia	M.J. Wingfield
	CMW 18448	GQ240650	GQ240666	GQ240682	<i>E. urophylla</i>	Indonesia	M.J. Wingfield
	CMW 18450	GQ240651	GQ240667	GQ240683	<i>E. grandis</i>	Thailand	M.J. Wingfield
	CMW 18458	GQ240654	GQ240670	GQ240686	<i>E. urophylla</i>	Indonesia	M.J. Wingfield
	CMW 18462	GQ240653	GQ240669	GQ240685	<i>E. urophylla</i>	Indonesia	M.J. Wingfield
	CMW 18463	GQ240646	GQ240662	GQ240678	<i>E. urophylla</i>	Indonesia	M.J. Wingfield
	CMW 20597	GQ240645	GQ240661	GQ240677	<i>E. grandis</i>	Thailand	M.J. Wingfield
	CMW 31177	GQ240657	GQ240673	GQ240689	<i>Eucalyptus</i> sp.	Indonesia	M.J. Wingfield
	CMW 31178	GQ240656	GQ240672	GQ240688	<i>Eucalyptus</i> sp.	Indonesia	M.J. Wingfield
	CMW 31179	GQ240655	GQ240671	GQ240687	<i>Eucalyptus</i> sp.	Indonesia	M.J. Wingfield



Table 1. (Continued)

Species	Isolate number ¹	GenBank accession no.			Host	Origin	Collector
		BT	HIS3	TEF-1 α			
<i>Ca. reteaudii</i>	CMW 31188	GQ240658	GQ240674	GQ240690	<i>Eucalyptus</i> sp.	Indonesia	M.J. Wingfield
	CMW 31189	GQ240647	GQ240663	GQ240679	<i>Eucalyptus</i> sp.	Indonesia	M.J. Wingfield
	CMW 31190	GQ240648	GQ240664	GQ240680	<i>Eucalyptus</i> sp.	Indonesia	M.J. Wingfield
	CMW 31191	GQ240652	GQ240668	GQ240684	<i>Eucalyptus</i> sp.	Thailand	M.J. Wingfield
<i>Ca. rumohrae</i>	CBS 111431 ^T	AF232871	DQ190675	FJ918549	<i>Rumohra adiantiformis</i>	Panama	J.W. Miller
	CBS 109062	AF232873	DQ190676	FJ918550	<i>Adiantum</i> sp.	The Netherlands	R. Pieters
<i>Ca. morganii</i>	CBS 110666	FJ918509	FJ918527	FJ918557	<i>Rosa</i> sp.	USA	N.E. Ell-Gholl
<i>Ca. spathiphylli</i>	CBS 116168 ^T	FJ918512	FJ918530	FJ918561	<i>Spathiphyllum</i> sp.	USA	C.L. Schoulties
<i>Ca. spathulata</i>	CBS 112689	AF308463	FJ918524	FJ918554	<i>E. viminalis</i>	Brazil	N.E. Ell-Gholl
	CMW 30601						
<i>Ca. terrae-reginae</i> sp. nov.	(= CBS 112151 = CPC 3202 = DFRI00150 = Lynfield 417) ^T	FJ918506	FJ918522	FJ918545	<i>E. urophylla</i>	Australia	C. Hanwood
	CMW 30602 (= CBS 112634)	FJ918507	DQ190668	FJ918546	<i>Xanthorrhoea australis</i>	Australia	T. Baigent

CBS: Centraalbureau voor Schimmelcultures, Utrecht, The Netherlands; CMW: culture collection of the Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria Pretoria, South Africa; ^T All ex-type cultures.

Fig. 1 One of six most parsimonious trees obtained from a heuristic search with 1 000 random addition of the combined sequences of β -tubulin, histone H3 and translation elongation factor-1 alpha sequence alignments of the *Ca. reteaudii* complex and other closely related species. Scale bar shows 10 changes and bootstrap support values from 1 000 replicates are shown above the nodes in bold. Bayesian posterior probability values are indicated below the nodes. Red lines indicate bootstrap support values of 100 and posterior probability values of 1.00. Thickened lines indicate branches in the strict consensus tree and the consensus tree of the Bayesian analyses. The tree was rooted to *Ca. colombiensis* (CBS 112221) and *Cy. chinense* (CBS 112744).




Fig. 2 One of three most parsimonious trees obtained from a heuristic search with 1 000 random addition sequences of the combined sequences of β -tubulin, histone H3 and translation elongation factor-1 alpha sequence alignments of the *Ca. morgani* complex and other closely related species. Scale bar shows 10 changes and bootstrap support values from 1 000 replicates are above the nodes in bold. Bayesian posterior probability values are indicated below the nodes. Red lines indicate bootstrap support values of 100 and posterior probability values of 1.00. Thickened lines indicate branches in the strict consensus tree and the consensus tree of the Bayesian analyses. The tree was rooted to *Ca. colombiensis* (CBS 112221) and *Cy. chinense* (CBS 112744).






Fig. 3 *Calonectria cerciana*. A. Macroconidiophore. B, C. fusiform to ellipsoidal vesicles. D, E. Fertile branches with reniform to doliiform phialides. F, G. Macroconidia. Scale bars A = 20 μm , B–G = 10 μm .






Fig. 4 *Calonectria pseudoreteauidii*. A. Macroconidiophore. B, C. Clavate vesicle, D, E. Macroconidia, F, G. Fertile branches with cylindrical to allantoid phialides. H, I. Microconidiophores. J. Microconidia. K. Comparison of macroconidia and microconidia. Scale bars A, H, I, K = 20 μm , B–G, J = 10 μm .






Fig. 5 *Calonectria queenslandica*. A. Macroconidiophore. B, C. Clavate vesicles. D, E. Fertile branches with cylindrical to allantoid phialides. F, G. Macroconidia. Scale bars A = 20 μm , B–G = 10 μm .







Fig. 6 *Calonectria terrae-reginae*. A. Macroconidiophore. B, C. Clavate vesicles. D, E. Fertile branches with cylindrical to allantoid phialides. F, G. Macroconidia. Scale bars A = 20 μm , B–G = 10 μm .





Chapter 4

Multigene phylogeny and mating tests reveal three cryptic species related to *Calonectria pauciramosa*

Submitted as: Lombard L, Crous PW, Wingfield BD, Wingfield MJ. (2009). Multigene phylogeny and mating tests reveal three cryptic species related to *Calonectria pauciramosa*. *Studies in Mycology*.

ABSTRACT

Calonectria pauciramosa is a pathogen of numerous plant hosts worldwide. Recent studies have suggested that it accommodates cryptic species and the aim of this study was to identify these taxa. Isolates from various geographical origins were collected and compared based on morphology, DNA sequence data of the β -tubulin, histone H3 and translation elongation factor-1 α regions and mating compatibility. Comparisons of the DNA sequence data and mating compatibility revealed three new species. These included *Ca. colombiana* sp. nov. from Colombia, *Ca. zuluensis* sp. nov. from South Africa and *Ca. polizzii* sp. nov. from Italy, all of which had distinguishing morphological features. Based on DNA sequence data, *Ca. brasiliensis* is also elevated to species level.

Taxonomic novelties: *Calonectria brasiliensis* (Bat. & Cif.) L. Lombard, M.J. Wingf. & Crous comb. nov., *Calonectria colombiana* L. Lombard, M.J. Wingf. & Crous sp. nov., *Calonectria polizzii* L. Lombard, M.J. Wingf. & Crous sp. nov., *Calonectria zuluensis* L. Lombard, M.J. Wingf. & Crous sp. nov.

INTRODUCTION

Several past studies have focused on the taxonomy of *Calonectria* spp. with small, 1-septate macroconidia and ellipsoidal to obpyriform vesicles (Crous *et al.* 1993, Overmeyer *et al.* 1996, Schoch *et al.* 1999, 2000). These *Calonectria* spp. were initially regarded as either *Ca. morganii* (= *Cy. scoparium*) or *Ca. scoparia* (= *Cy. candelabrum*) based on their morphological similarities. However, the anamorph state of *Ca. morganii* was circumscribed as having ellipsoidal to pyriform vesicles and *Ca. scoparia* having ellipsoidal to obpyriform vesicles by Crous *et al.* (1993). Later studies, incorporating DNA sequence data, have shown that *Ca. morganii* is restricted to the Northern Hemisphere and Brazil (Crous *et al.* 1993, Overmeyer *et al.* 1996, Schoch *et al.* 2000). In contrast, *Ca. scoparia* is found worldwide and forms part of a species complex consisting of four mating groups, each representing a different *Calonectria* species that includes *Ca. pauciramosa* (anamorph: *Cy. pauciramsum*), *Ca. scoparia*, *Ca. mexicana* (anamorph: *Cy. mexicanum*) and *Ca. insularis* (anamorph: *Cy. insulare*) (Schoch *et al.* 1999).

Calonectria pauciramosa has been reported worldwide on numerous plant hosts (Schoch *et al.* 1999, Koike *et al.* 1999, Koike & Crous 2001, Polizzi & Crous 1999, Polizzi 2000, Polizzi & Catara 2001, Polizzi & Vitale 2001, Crous 2002, Polizzi *et al.* 2006, 2007), where it causes diseases such as cutting rot, damping-off, root rot and leaf blight. In South Africa and Australia, *Ca. pauciramosa* is regarded as the dominant pathogen in commercial forest nurseries (Crous 2002) and it is also found on various horticultural crops in commercial nurseries in Italy and the U.S.A. (Schoch *et al.* 2001, Crous 2002).

Schoch *et al.* (2001) considered female fertility in populations of *Ca. pauciramosa* from various geographical regions to determine the ratio of mating types present, and based on these data suggested that *Ca. pauciramosa* could be endemic to South America. The latter study also indicated that *Ca. pauciramosa* isolates from California were represented by only one mating type, supporting the view that this represented an introduced pathogen. Isolates from Italy showed higher ratios of hermaphrodites, also indicative of a recent introduction, although some variation was observed in the β -tubulin sequences. In contrast, South African isolates had close to a 1:1 mating type ratio and showed variation in β -tubulin sequence data (Schoch *et al.* 1999, 2001), suggesting that this was either a native pathogen or that there had been multiple introductions into the country.

Initial investigations using DNA sequence comparisons and mating studies on *Ca. pauciramosa* isolates from South Africa and Colombia showed some variation amongst isolates. These findings and those of Schoch *et al.* (2001) suggested that *Ca. pauciramosa* might accommodate a number of cryptic species. The aim of this study was, therefore, to consider the phylogenetic relationships, morphological characters and mating compatibility of available isolates of *Ca. pauciramosa* and to determine whether this species represented an assemblage of cryptic taxa.

MATERIALS AND METHODS

Isolates

Isolates of *Ca. pauciramosa* were obtained from culture collections (Table 1) or were isolated from infected plant material and soil samples following the methods of Crous (2002). For each isolate, single conidial cultures were prepared on 2 % (w/v) malt extract agar (MEA, Biolab, Midrand, South Africa). Representative strains are maintained in the culture collection (CMW) of the Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria, South Africa and the Centraalbureau voor Schimmelcultures (CBS), Utrecht, The Netherlands.

Sexual compatibility

A total of 57 single conidial *Ca. pauciramosa*-like isolates (Table 1), originating from various geographic regions and hosts were crossed in all possible combinations. Mating-tester strains CMW 30823 (= STE-U 416) and CMW 5683 (= STE-U 971) for *Ca. pauciramosa* defined by Schoch *et al.* (2001) were also crossed with these isolates. Matings were done as described in Schoch *et al.* (1999) on carnation leaf agar (CLA; Fisher *et al.* 1982, Crous *et al.* 1993) and on minimal salt agar (MSA; Guerber & Correll 2001, Halleen *et al.* 2006) with sterile toothpicks placed on the surface of the agar. Control tests, where isolates were crossed with themselves, were undertaken to determine whether strains had a heterothallic or homothallic mating system. The plates were stacked in plastic containers and incubated at 22 °C for 6 wk. Matings were regarded as successful when isolate combinations produced perithecia extruding viable ascospores.

DNA sequence comparisons

Calonectria pauciramosa-like isolates were grown on MEA for 7 d. Mycelium was then scraped from the surface of the cultures, freeze-dried, and ground to a powder in liquid

nitrogen, using a mortar and pestle. DNA was extracted from the powdered mycelium as described by Lombard *et al.* (2008). Three loci including fragments of the β -tubulin (BT), histone H3 (HIS3) and translation elongation factor-1 alpha (TEF-1 α) gene regions were sequenced. Primers used to sequence these regions were T1 (O'Donnell & Cigelnik 1997) and CYLTUB1R (Crous *et al.* 2004b) for the BT region, CYLH3F and CYLH3R (Crous *et al.* 2004b) for the HIS3 region and EF1-728F (Carbone & Kohn 1999) and EF2 (O'Donnell *et al.* 1998) for the TEF-1 α region. The PCR reaction mixture used to amplify the different loci consisted of 2.5 units FastStart *Taq* polymerase (Roche Applied Science, USA), 10 \times PCR buffer, 1–1.5 mM MgCl₂, 0.25 mM of each dNTP, 0.5 μ M of each primer and approximately 30 ng of fungal genomic DNA, made up to a total reaction volume of 25 μ L with sterile distilled water.

Amplified fragments were purified using High Pure PCR Product Purification Kit (Roche, USA) and sequenced in both directions. For this purpose, the BigDye terminator sequencing kit v. 3.1 (Applied Biosystems, USA) and an ABI PRISMTM 3100 DNA sequencer (Applied Biosystems) were used. All PCRs and sequencing reactions were performed on an Eppendorf Mastercycler Personal PCR (Eppendorf AG, Germany) with cycling conditions as described in Crous *et al.* (2006) for BT and HIS3. The same cycling conditions for HIS3 were used for TEF-1 α amplifications.

The generated sequences were added to other sequences of closely related *Calonectria* spp. obtained from GenBank (<http://www.ncbi.nlm.nih.gov>) and these were assembled and aligned using Sequence Navigator v. 1.0.1 (Applied Biosystems) and MAFFT v. 5.11 (Katoh *et al.* 2005), respectively. The aligned sequences were then manually corrected where needed. Single nucleotide polymorphisms (SNP'S) were determined for each gene region analysed using DnaSP v. 5.00.07 (Librado & Rozas 2009).

To determine whether the DNA sequence datasets for the three gene regions were congruent, a 70 % reciprocal bootstrap method using Neighbour-Joining with Maximum Likelihood distance was employed (Mason-Gamer & Kellogg 1996, Gueidan *et al.* 2007). Models of evolution were estimated in Modeltest v. 3.7 (Posada & Crandall 1998) using the Akaike Information Criterion for each separate gene region. The bootstrap analyses were run in PAUP (Phylogenetic Analysis Using Parsimony v. 4.0b10, Swofford 2002) for 10 000 replicates. Resulting tree topologies were compared visually for conflicts between the

separate gene regions. Phylogenetic relationships were estimated in PAUP, by heuristic searches based on 1 000 random addition sequences and tree bisection-reconnection was used, with the branch swapping option set on “best trees” only.

All characters were weighted equally and alignment gaps were treated as missing data. Measures calculated for parsimony included tree length (TL), consistency index (CI), retention index (RI) and rescaled consistence index (RC). Bootstrap analysis (Hillis & Bull 1993) was based on 1 000 replications. All sequences for the isolates studied were analysed using the Basic Local Alignment Search Tool for Nucleotide sequences (BLASTN, Altschul *et al.* 1990). The phylogenetic analysis included 73 partial gene sequences per gene, representing 11 *Calonectria* and *Cylindrocladium* species (Table 1). *Calonectria colombiensis* (CBS 112221) and *Cy. chinense* (CBS 112744) were used as outgroup taxa (Lombard *et al.* 2009). Novel sequences were deposited in GenBank and all alignments in TreeBASE (<http://treebase.org>) as SN4773.

A Markov Chain Monte Carlo (MCMC) algorithm was used to generate phylogenetic trees with Bayesian probabilities using MrBayes v. 3.1.1 (Ronquist & Huelsenbeck 2003). Models of nucleotide substitution for each gene were determined using Mrmodeltest (Nylander 2004) and included for each gene partition. Two analyses of four MCMC chains were run from random trees for 1 000 000 generations and sampled every 100 generations. Both runs converged on the same likelihood score and tree topology. Therefore, the first 1 000 trees were discarded as the burn-in phase of each analysis and posterior probabilities were determined from the remaining trees.

Taxonomy

For morphological identification of the anamorphs, single conidial cultures were prepared on synthetic nutrient-poor agar (SNA; Nirenburg 1981, Lombard *et al.* 2009). Inoculated plates were incubated at room temperature and examined after 7d. Gross morphological characteristics were determined by mounting fungal structures in lactic acid and 30 measurements at $\times 1\,000$ magnification were made for each isolate. Teleomorph morphology was determined by mounting perithecia obtained from the sexual compatibility tests in Leica mountant (Setpoint Premier, Johannesburg, South Africa) and hand-sectioned with a Leica CM1100 cryostat (Setpoint Technologies) at $-20\text{ }^{\circ}\text{C}$. The $10\text{ }\mu\text{m}$ sections were mounted in lactophenol or 3 % KOH. Gross morphological characteristics were observed as above. The

95 % confidence levels were calculated and extreme measurements of conidia are given in parentheses. For other structures, only the extremes are indicated. Optimal growth temperatures were determined for each isolate on MEA at 5–35 °C in 5 °C intervals in the dark. Colony colours were determined after 7 d on MEA at 25 °C in the dark, using the colour charts of Rayner (1970) for comparison. Descriptions, nomenclature, and illustrations were deposited in MycoBank (Crous *et al.* 2004a).

RESULTS

Sexual compatibility

Protoperithecia formed within 3 wk and successful matings produced perithecia with viable ascospores within 6 wk on both CLA and MSA. A total of 1 649 crosses were made using the 57 putative *Ca. pauciramosa* isolates and mating tester strains for *Ca. pauciramosa s. str.* This resulted in 642 tests where perithecia produced viable ascospores. Self-self crosses indicated that 11 of the 57 isolates were self-fertile (homothallic). These included the Colombian isolates CBS 111041, CBS 111136, CBS 115127, CBS 115638, CBS 115694 and CMW 9058, and South African isolates CMW 9115, CMW 9188, CMW 9208, CMW 9215 and CMW 9896. Sixteen of the 57 putative *Ca. pauciramosa* did not cross with the mating tester strains for that species or with any other isolate included in this study. These included isolates CMW 7578 from Argentina; CBS 114257, CBS 116078, CBS 116076, CBS 116081, CMW 31505, CMW 31507 and CMW 31508, from Brazil; CMW 7804, CMW 10151 and CBS 123402 from Italy, CMW 30814 and CMW 30815 from Kenya; CMW 30817 from New Zealand; CMW 1786 and CMW 30815 from South Africa. The remaining 30 isolates produced perithecia containing viable ascospores when crossed with the *Ca. pauciramosa* mating tester strains and between them. This resulted in 203 successful heterothallic matings (Table 2).

DNA sequence comparisons

Amplicons of approx. 500 bp were generated for the BT and TEF-1 α gene regions and those for the HIS3 region were approx. 450 bp. Comparing the tree topologies of the 70 % reciprocal bootstrap trees indicated no conflicts. Subsequently, the datasets were combined and this resulted in a data set consisting of 1 529 characters including gaps. Of these characters, 1 151 were constant and parsimony uninformative. The 378 parsimony informative characters included in the parsimony analyses yielded eight most parsimonious

trees (TL = 993, CI = 0.732, RI = 0.903, RC = 0.661), one of which is presented (Fig. 1). For Bayesian analyses, a HKY+I model was selected for BT, GTR+I+G model for HIS3 and a GTR+G model for TEF-1 α and incorporated into the analyses. The consensus tree obtained for the Bayesian analyses confirmed the tree topology obtained with parsimony as well as bootstrap support (Fig. 1).

The majority of the *Ca. pauciramosa* isolates grouped together to form a monophyletic cluster with a bootstrap (BP) value of 100 and a Bayesian posterior probability (PP) value of 1.00. Within this cluster, two separate clades could be distinguished. The first (BP = 66, PP = 0.92) represented isolates obtained from South Africa (Table 1) and analyses of the SNP's (Table 3) showed one fixed allele for BT, two for HIS3 and one indel for TEF-1 α . The second clade (BP = 97, PP = 1.00) represented isolates from Italy (Table 1) that were closely related to *Ca. pauciramosa* and have a number of shared fixed polymorphisms; five BT and two HIS3 (Table 3). Isolates from Colombia (Table 1) grouped together (BP = 100, PP = 1.00), separate from the *Ca. pauciramosa* cluster and SNP analyses show that six BT, 13 HIS3 and nine TEF-1 α shared fixed alleles including three indels are characteristic for this group (Table 3). These isolates were closely related to *Ca. spathulata*. Isolates from Brazil grouped together with isolate CBS 230.51 (ex-type of *Cy. brasiliensis*; BP = 100, PP = 1.00), closely related to *Ca. morgani* and *Ca. insularis*, but separate from both of these species. Analyses of the SNP's for the isolates from Brazil compared to *Ca. morgani* and *Ca. insulare* also show several fixed alleles for these isolates, which include the ex-type culture of *Cy. brasiliensis* (CBS 230.51) (Table 4). The DNA sequence data for the three gene regions used in the present study showed 16 fixed alleles between *Ca. brasiliensis*, *Ca. insularis* and *Ca. morgani* (Table 4). An additional 10 fixed alleles were shared between *Cy. brasiliensis* and *Ca. insularis* and distinguished both species from *Ca. morgani*.

Taxonomy

Isolates CMW 9115, CMW 9188, CMW 9208, CMW 9215 and CMW 9896 represent a distinct species closely related to *Ca. pauciramosa*, based on phylogenetic inference. Mating studies also showed that these isolates have a homothallic mating system, distinguishing them from *Ca. pauciramosa s. str.* A similar situation was found for the isolates CBS 111136, CBS 115127, CBS 115638 and CBS 115694 from Colombia and they are also treated as a new species based on their homothallic mating system and phylogenetic inference. Furthermore, isolates CBS 123402, CMW 7804 and CMW 10151 from Italy are closely related to *Ca.*

pauciramosa and failed to cross with the mating tester strains of that species. However, morphological observations and DNA sequence data indicate that these isolates represent an undescribed taxon.

Species of *Cylindrocladium* (1892) represent anamorph states of *Calonectria* (1867) (Rossman *et al.* 1999). In this study, these fungi are described as new species of *Calonectria*, which represents the older generic name for these holomorphs. This is irrespective whether the teleomorph states of these fungi have been found or not. This is consistent with the new regulations on fungal nomenclature as proposed by Hawksworth (2005) stating that for all newly described pleomorphic fungal species, the teleomorph name takes precedence over the anamorph name when both types belong to the same holomorph taxon.

Calonectria brasiliensis (Bat. & Cif.) L. Lombard, M.J. Wingf. & Crous, **comb. nov.** MycoBank MB 515110. Fig. 2.

Basionym: *Cylindrocladium brasiliensis* (Bat. & Cif.) Peeraly, (as *braziliensis*) CMI Descriptions of Pathogenic Fungi and Bacteria 427. 1974.

≡ *Cylindrocladium scoparium* var. *brasiliensis* Bat. & Cif., (as *brasiliense*) Boletim de SA.I.C. Pernambuco 18: 188–191. 1951.

Teleomorph unknown. *Conidiophores* with a stipe bearing a penicillate suite of fertile branches, stipe extensions, and terminal vesicles. *Stipe* septate, hyaline, smooth, 63–103 × 7–14 μm; stipe extensions septate, straight to flexuous, 204–266 μm long, 6–7 μm wide at the apical septum, terminating in a broadly clavate to ellipsoidal to fusiform vesicle, 7–11 μm diam. *Conidiogenous apparatus* 58–90 μm long, and 81–103 μm wide; primary branches aseptate or 1-septate, 25–34 × 5–8 μm; secondary branches aseptate, 14–25 × 4–7 μm; tertiary branches aseptate, 8–20 × 3–5 μm, each terminal branch producing 2–6 phialides; phialides doliiiform to reniform, hyaline, aseptate, 8–12 × 2–4 μm; apex with minute periclinal thickening and inconspicuous collarette. *Conidia* cylindrical, rounded at both ends, straight, (35–)36–40(–41) × 3–5 μm (av. = 38 × 3.5 μm), 1-septate, lacking a visible abscission scar, held in parallel cylindrical clusters by colourless slime. *Megaconidia* and *microconidia* not seen.

Specimens examined: **Brazil**, Ceara State, *Eucalyptus* sp., Sep. 1948, T.R. Ciferri, ex-type culture CBS 230.51 = IMI 299576 = CMW 23671. Aracruz, *Eucalyptus* sp., June 1998, A.C.

Alfenas, CBS 114257 = CMW 32949. Rio de Janeiro, *E. citriodora*, A.O. Carvalho, CBS 116078 = CMW 32950. Champion nursery, *Eucalyptus* sp., P.W. Crous, CPC 602 = CMW 31507. Aracruz, *Eucalyptus* sp., P.W. Crous, CPC 1943 = CMW 31508.

Culture characteristics: Colonies fast growing with optimal growth temperature at 25 °C (growth at 10–30 °C) on MEA, reverse amber (13k) to sepia-brown (13i) after 7 d; sparse white aerial mycelium with sparse sporulation; chlamydospores moderate throughout the medium, forming microsclerotia.

Substrates: *Eucalyptus* spp.

Distribution: Brazil.

Notes: Based on morphological observations, Crous & Wingfield (1994) reduced *Ca. brasiliensis* to synonymy with *Ca. morganii*. However, phylogenetic inference in this study has shown that the ex-type culture of *Ca. brasiliensis* (CBS 230.51) is distinct from *Ca. morganii* (CBS 110666). Morphological observations in this study also indicated that conidia of *Ca. brasiliensis* (av. $38 \times 3.5 \mu\text{m}$) are smaller than those of *Ca. morganii* (av. $45 \times 4 \mu\text{m}$). *Calonectria brasiliensis* only produces up to three branches per conidiophore, whereas *Ca. morganii* can have up to six branches per conidiophore.

Calonectria colombiana L. Lombard, Crous & M.J. Wingf., **sp. nov.** MycoBank MB515065, Fig. 3.

Etymology: Name refers to Colombia, the country this fungus was isolated from.

Telomorpha *Calonectriae pauciramosa* similis, sed ascosporis brevioribus, (28–)31–36(–40) \times 3–5 μm (in medio 34 \times 4 μm). Culturae homothallicae. Anamorpha *Cylindrocladio pauciramoso* simile, sed vesiculis obpyriforme vel fusiforme (8–12 μm diam.) et conidiis maioribus (33–)35–39(–40) \times 3–4 μm , in medio 37 \times 3 μm .

Perithecia solitary or in groups, orange to red, becoming red-brown with age; in section, apex and body yellow to orange, base red-brown, sub-globose to ovoid, 270–410 μm high, 175–285 μm diam, body turning dark red, and base dark red-brown (KOH+). Perithecial walls rough, consisting of 2 thick-walled layers: outside layer of *textura globulosa*, 24–90 μm wide; becoming more compressed towards inner layer of *textura angularis*, 18–22 μm wide; becoming thin-walled and hyaline towards the center, outer cells, 38–55 \times 16–40 μm ; inner cells, 3–12 \times 3–7 μm ; perithecial base up to 114 μm wide; consisting of dark red, angular

cells; merging with an erumpent stroma, cells of the outer wall layer continuing into the pseudoparenchymatous cells of the erumpent stroma. *Asci* 8-spored, clavate, $87\text{--}162 \times 12\text{--}18 \mu\text{m}$, tapering to a long thin stalk. *Ascospores* aggregated in the upper third of the ascus, hyaline, gluttulate, fusoid with rounded ends, straight to slightly curved, 1-septate, not or slightly constricted at the septum, $(28\text{--})31\text{--}36(\text{--}40) \times 3\text{--}5 \mu\text{m}$ (av. = $34 \times 4 \mu\text{m}$). Cultures homothallic. *Conidiophores* with a stipe bearing a penicillate suite of fertile branches, stipe extensions, and terminal vesicles. *Stipe* septate, hyaline, smooth, $45\text{--}126 \times 6\text{--}9 \mu\text{m}$; stipe extensions septate, straight to flexuous, $143\text{--}173 \mu\text{m}$ long, $5\text{--}7 \mu\text{m}$ wide at the apical septum, terminating in an obpyriform to fusiform vesicle, $8\text{--}12 \mu\text{m}$ diam. *Conidiogenous apparatus* $38\text{--}115 \mu\text{m}$ long, and $35\text{--}91 \mu\text{m}$ wide; primary branches aseptate or 1-septate, $19\text{--}37 \times 5\text{--}8 \mu\text{m}$; secondary branches aseptate, $9\text{--}17 \times 4\text{--}5 \mu\text{m}$; tertiary and additional branches (–4), aseptate, $8\text{--}13 \times 3\text{--}4 \mu\text{m}$, each terminal branch producing 2–6 phialides; phialides doliiform to reniform, hyaline, aseptate, $9\text{--}12 \times 3\text{--}4 \mu\text{m}$; apex with minute periclinal thickening and inconspicuous collarete. *Conidia* cylindrical, rounded at both ends, straight, $(33\text{--})35\text{--}39(\text{--}40) \times 3\text{--}4 \mu\text{m}$ (av. = $37 \times 3 \mu\text{m}$), 1-septate, lacking a visible abscission scar, held in parallel cylindrical clusters by colourless slime. *Megaconidia* and *microconidia* not seen.

Specimen examined: **Colombia**, La Selva, from soil, June 1995, M.J. Wingfield, Herb. PREM 60295, **holotype** of *Calonectria colombiana*, cultures ex-type CBS 115127 = CMW 30871 = CPC 1160; La Selva, June 1995, M.J. Wingfield, CBS 111041 = CMW 30767 = CPC 1163; La Selva, June 1995, M.J. Wingfield, CBS 111136 = CMW 30812 = CPC 1151; CBS 115638 = CMW 30766 = CPC 1161 (Herb. PREM 60296); La Selva, June 1995, M.J. Wingfield, CBS 115694 = CMW 30813 = CPC 1162; La Selva, June 1995, M.J. Wingfield, CMW 9058.

Culture characteristics: Colonies fast growing with optimal growth temperature at $25 \text{ }^\circ\text{C}$ (growth at $10\text{--}30 \text{ }^\circ\text{C}$) on MEA, reverse amber (13k) to sepia-brown (13i) after 7 d; abundant white aerial mycelium with sparse sporulation; chlamydospores extensive throughout the medium, forming microsclerotia.

Substrate: Soil.

Distribution: Colombia.

Notes: Isolates of *Ca. colombiana* were previously regarded as either *Ca. pauciramosa* or *Ca. scoparia* (Crous 2002) based on the morphological similarity of the anamorph states of these species. Based on macroconidial dimensions, *Ca. colombiana* (av. $37 \times 3 \mu\text{m}$) can be distinguished from *Ca. pauciramosa* (av. $50 \times 4.5 \mu\text{m}$) and *Ca. scoparia* (av. $60 \times 4.5 \mu\text{m}$) in having smaller, 1-septate macroconidia. Both *Ca. pauciramosa* and *Ca. scoparia* have a biallelic, heterothallic mating system (Schoch *et al.* 1999, 2001), whereas *Ca. colombiana* is homothallic.

Calonectria polizzii L. Lombard, Crous & M.J. Wingf., **sp. nov.** MycoBank MB515066, Fig. 4.

Etymology: The name honours Prof. dr. Giancarlo Polizzi, who isolated the fungus in Italy.

Teleomorpha ignota. *Cylindrocladio pauciramoso* simile, sed vesiculis clavato vel obpyriforme ($6\text{--}9 \mu\text{m}$ diam.) et conidiis maioribus ($31\text{--}32\text{--}42\text{--}49$) \times $3\text{--}5 \mu\text{m}$, in medio $37 \times 4 \mu\text{m}$.

Teleomorph unknown. *Conidiophores* with a stipe bearing a penicillate suite of fertile branches, stipe extensions, and terminal vesicles. *Stipe* septate, hyaline, smooth, $58\text{--}108 \times 5\text{--}7 \mu\text{m}$; stipe extensions septate, straight to flexuous, $111\text{--}167 \mu\text{m}$ long, $5\text{--}6 \mu\text{m}$ wide at the apical septum, terminating in a broadly clavate to pyriform vesicle, $6\text{--}9 \mu\text{m}$ diam. *Conidiogenous apparatus* $27\text{--}57 \mu\text{m}$ long, and $28\text{--}51 \mu\text{m}$ wide; primary branches aseptate or 1-septate, $15\text{--}35 \times 4\text{--}6 \mu\text{m}$; secondary branches aseptate, $12\text{--}26 \times 3\text{--}5 \mu\text{m}$; tertiary branches aseptate, $10\text{--}15 \times 4\text{--}5 \mu\text{m}$, each terminal branch producing 2–6 phialides; phialides doliiform to reniform, hyaline, aseptate, $8\text{--}13 \times 3\text{--}4 \mu\text{m}$; apex with minute periclinal thickening and inconspicuous collarette. *Conidia* cylindrical, rounded at both ends, straight, ($31\text{--}32\text{--}42\text{--}49$) \times $3\text{--}5 \mu\text{m}$ (av. = $37 \times 4 \mu\text{m}$), 1-septate, lacking a visible abscission scar, held in parallel cylindrical clusters by colourless slime. *Megaconidia* and *microconidia* not seen.

Specimen examined: **Italy**, Sicily, Carrubba, on *Arbutus unedo*, 1997, G. Polizzi, Herb. PREM 60297, **holotype** of *Calonectria polizzii*, cultures ex-type CBS 123402 = CMW 30872; *Callistemon citrinus*, 1997, G. Polizzi, CMW 7804 = CPC 2681 = CBS 125270; *Callistemon citrinus*, 1997, G. Polizzi CMW 10151 = CPC 2771 = CBS 125271 (Herb. PREM 60298).

Culture characteristics: Colonies fast growing with optimal growth temperature at $25 \text{ }^\circ\text{C}$ (growth at $10\text{--}30 \text{ }^\circ\text{C}$) on MEA, reverse amber (13k) to sepia-brown (13i) after 7 d; abundant

white aerial mycelium with sparse sporulation; chlamydospores extensive throughout the medium, forming microsclerotia.

Substrates: Arbutus unedo, Callistemon citrinus.

Distribution: Italy.

Notes: Calonectria polizzii is morphologically similar to *Ca. pauciramosa* and *Ca. zuluensis*. The macroconidia of *Ca. polizzii* (av. $37 \times 4 \mu\text{m}$) are smaller to those of *Ca. pauciramosa* (av. $50 \times 4.5 \mu\text{m}$). Mating tests also showed that *Ca. polizzii* does not mate with either of the tester strains of *Ca. pauciramosa* (Schoch *et al.* 2001) used in this study. However, the isolates of *Ca. polizzii* tested might represent a single mating type, or might have lost their ability to mate, and further studies incorporating more isolates will be required to confirm this.

Calonectria zuluensis L. Lombard, Crous & M.J. Wingf., **sp. nov.** MycoBank MB515067, Fig. 5.

Etymology: Name refers to KwaZulu-Natal, South Africa, the province where the fungus was isolated.

Telomorpha *Calonectriae pauciramosa* similis, sed ascosporis brevioribus, (26–)29–34(–38) \times 4–5 μm (in medio 32 \times 4 μm). Culturae homothallicae. Anamorpha *Cylindrocladio pauciramoso* simile, sed vesiculis clavato vel obpyriforme (6–10 μm diam) et conidiis maioribus (31–)34–38(–40) \times 3–5 μm , in medio 36 \times 4 μm .

Perithecia solitary or in groups, orange to red, becoming red-brown with age; in section apex and body yellow to orange, base red-brown, sub-globose to ovoid, 292–394 μm high, 170–285 μm diam, body turning dark red, and base dark red-brown (KOH+). Perithecial walls rough, consisting of 2 thick-walled layers: outside layer of *textura globulosa*, 30–80 μm wide; becoming more compressed towards inner layer of *textura angularis*, 20–22 μm wide; becoming thin-walled and hyaline towards the center, outer cells, 40–50 \times 18–40 μm ; inner cells, 4–12 \times 3–5 μm : perithecial base up to 116 μm wide; consisting of dark red, angular cells; merging with an erumpent stroma, cells of the outer wall layer continuing into the pseudoparenchymatous cells of the erumpent stroma. *Asci* 8-spored, clavate, 92–140 \times 10–16 μm , tapering to a long thin stalk. *Ascospores* aggregate in the upper third of the ascus, hyaline, gluttulate, fusoid with rounded ends, straight to slightly curved, 1-septate, not or slightly constricted at the septum, (26–)29–34(–38) \times 4–5 μm (av. = 32 \times 4 μm). Cultures

homothallic. *Conidiophores* with a stipe bearing penicillate clusters of fertile branches, stipe extensions, and terminal vesicles. *Stipe* septate, hyaline, smooth, $57\text{--}84 \times 6\text{--}9 \mu\text{m}$; stipe extensions septate, straight to flexuous, $110\text{--}171 \mu\text{m}$ long, $5\text{--}8 \mu\text{m}$ wide at the apical septum, terminating in a broadly clavate to obpyriform vesicle, $6\text{--}10 \mu\text{m}$ diam. *Conidiogenous apparatus* $35\text{--}67 \mu\text{m}$ long, and $37\text{--}70 \mu\text{m}$ wide; primary branches aseptate or 1-septate, $16\text{--}28 \times 4\text{--}6 \mu\text{m}$; secondary branches aseptate, $11\text{--}20 \times 3\text{--}5 \mu\text{m}$; tertiary branches aseptate, $8\text{--}13 \times 3\text{--}4 \mu\text{m}$, each terminal branch producing 2–6 phialides; phialides doliiiform to reniform, hyaline, aseptate, $10\text{--}13 \times 3\text{--}4 \mu\text{m}$; apex with minute periclinal thickening and inconspicuous collarette. *Conidia* cylindrical, rounded at both ends, straight, $(31\text{--})34\text{--}38(\text{--}40) \times 3\text{--}5 \mu\text{m}$ (av. = $36 \times 4 \mu\text{m}$), 1-septate, lacking a visible abscission scar, held in parallel cylindrical clusters by colourless slime. *Megaconidia* and *microconidia* not seen.

Specimen examined: **South Africa**, KwaZulu-Natal, Kwambonambi, from *Eucalyptus grandis* clonal cutting, Feb. 2001, L. Lombard, Herb. PREM 60292, **holotype** of *Calonectria zuluensis*, cultures ex-type CBS 125268 = CMW 9188; KwaZulu-Natal, Kwambonambi, *E. grandis* \times *urophylla* hybrid cutting, Feb. 2001, L. Lombard, CMW 9115, CMW 9208 (Herb. PREM 60293), CMW 9215, Pietermaritzburg, *E. grandis* \times *urophylla* hybrid cutting, Mar. 2001, L. Lombard, CMW 9896 = CBS 125272.

Culture characteristics: Colonies fast growing with optimal growth temperature at $25 \text{ }^\circ\text{C}$ (growth at $10\text{--}30 \text{ }^\circ\text{C}$) on MEA, reverse amber (13k) to sepia-brown (13i) after 7 d; abundant white aerial mycelium with sparse sporulation; chlamydospores extensive throughout the medium, forming microsclerotia.

Substrate: *Eucalyptus grandis* rooted cuttings, *E. grandis* \times *urophylla* rooted cuttings

Distribution: South Africa.

Notes: *Calonectria zuluensis* can be distinguished from *Ca. pauciramosa* and *Ca. scoparia* based on its homothallic mating system. Macroconidia of *Ca. zuluensis* (av. $36 \times 4 \mu\text{m}$) are also smaller than those of *Ca. pauciramosa* (av. $50 \times 4.5 \mu\text{m}$) and *Ca. scoparia* (av. $60 \times 4.5 \mu\text{m}$). This species is morphologically very similar to *Ca. colombiana*. However, *Ca. zuluensis* can be distinguished from *Ca. colombiana* based on the fact that it has a broadly clavate to obpyriform vesicle as compared with a obpyriform to fusiform vesicle in *Ca.*

colombiana. Furthermore, *Ca. zuluensis* can easily be distinguished based on phylogenetic inference.

DISCUSSION

Considerable variation observed amongst isolates of “*Ca. pauciramosa*” from different geographical localities was illustrated in this study. Morphological characteristics, phylogenetic inference and mating studies revealed the presence of three cryptic species accommodated in cultures that have collectively been treated as *Ca. pauciramosa*. This is consistent with the results of previous studies (Schoch *et al.* 1999, 2001), which noted variation within *Ca. pauciramosa*, although at that time the sample size was inordinately small to consider the matter further. Schoch *et al.* (2001) also noted a high level of variation among isolates from South America, but concluded that this most likely reflected diversity consistent with an endemic population.

Crous (2002) suggested that mating isolates with recognised mating tester strains represented an important step in identifying isolates of *Ca. pauciramosa*. Various studies (Crous *et al.* 1993, Crous & Wingfield 1994, Crous *et al.* 1998, Schoch *et al.* 1999, 2001, Crous 2002) have used CLA as standardised medium to study sexual compatibility amongst isolates of *Cylindrocladium*. However, CLA has its limitations in that carnation leaf pieces are not always available and the present study used both CLA and MSA amended with sterile tooth picks, which proved to be very successful. Effective application of the latter technique to induce teleomorphs in culture has also been achieved for various other plant pathogenic genera, including *Glomerella* (Geurber & Correll 2001) and *Neonectria* (Halleen *et al.* 2006).

The descriptions of *Ca. colombiana*, *Ca. zuluensis* and *Ca. polizzii* add three new species to the *Ca. scoparia* species complex. This complex is characterised by species having ellipsoidal to obpyriform vesicles and producing 1-septate macroconidia (Schoch *et al.* 1999, Crous 2002). The complex was previously regarded as having a biallelic, heterothallic mating system (Schoch *et al.* 1999, 2001). However, both the newly described *Ca. colombiana* and *Ca. zuluensis* are homothallic. The occurrence of both heterothallic and homothallic *Calonectria* species in a single complex is not unique, having previously been found in the *Ca. kyotensis* species complex (Crous *et al.* 2004b).

Schoch *et al.* (2001) considered female fertility of *Ca. pauciramosa*, and found variation in BT sequence data for isolates from Italy. This variation has most likely been captured in the description of *Ca. polizzii* in the present study. This new species has thus been shown as unique based on morphological, phylogenetic inference and biological characteristics, separating it from *Ca. pauciramosa*. Morphologically, *Ca. polizzii* can be distinguished from *Ca. pauciramosa* by its smaller 1-septate macroconidia. Isolates of *Ca. polizzii* were also not capable of mating with the *Ca. pauciramosa* mating-tester strains or other *Ca. pauciramosa* isolates from different geographic regions.

Schoch *et al.* (2001), noted variation amongst isolates of *Ca. pauciramosa* from South America, and suggested that the fungus could be native to that continent. Results of the present study, including isolates from Colombia, led to the description of *Ca. colombiana*. This fungus is distinct from *Ca. pauciramosa* in having a homothallic mating system, smaller macroconidia and quaternary branches on the conidiophores. Although *Ca. insularis* also forms conidiophores with quaternary branches (Schoch *et al.* 1999), *Ca. colombiana* can easily be distinguished from it based on DNA sequence comparisons and its homothallic mating system.

Various species of *Calonectria* have been recorded from South Africa (Crous *et al.* 1991, Crous *et al.* 1993, Schoch *et al.* 1999, Crous 2002) and the description of *Ca. zuluensis* adds another species to those already reported from the country. *Calonectria zuluensis* has a homothallic mating system, which is different from *Ca. pauciramosa* with a biallelic, heterothallic mating system (Schoch *et al.* 2001). The two species can also easily be distinguished from each other based on DNA sequence comparisons.

In the analyses of the SNP's for the three gene regions used in this study, several fixed and shared SNP alleles were found for *Ca. colombiana*, *Ca. polizzii* and *Ca. zuluensis*. The majority of the fixed SNPs are shared between *Ca. polizzii* and *Ca. zuluensis*, indicating that these are sibling species, and that genetic isolation between them occurred recently (Taylor *et al.* 2000). For *Ca. colombiana*, fewer of the fixed SNPs are shared with *Ca. polizzii* and *Ca. zuluensis*, indicating that speciation occurred less recently than that of *Ca. polizzii* and *Ca. zuluensis*. These three species do not share the same alleles with *Ca. pauciramosa*, clearly distinguishing it from them.

Calonectria brasiliensis has been elevated to species level based on phylogenetic inference. Although Peerally (1974) indicated that the macroconidia of *Ca. brasiliensis* ($24\text{--}38 \times 2\text{--}3 \mu\text{m}$) are smaller than those of *Ca. morgani* (av. $45 \times 4 \mu\text{m}$), Crous & Wingfield (1994) reduced *Ca. brasiliensis* to *Ca. morgani*, based on similar conidial dimensions and vesicles morphology observed in culture. It is possible, however, that the original ex-type strain of *Ca. brasiliensis* was in fact morphologically degenerated, appearing atypical for the species. Several isolates from Brazil, previously identified as *Ca. pauciramosa*, grouped with the ex-type strain of *Ca. brasiliensis* (CBS 230.51). Previous DNA sequence comparisons and mating studies with *Ca. morgani* (Crous *et al.* 1993, Overmeyer *et al.* 1996, Schoch *et al.* 2000, 2001) failed to include the ex-type strain CBS 230.51 of *Ca. brasiliensis*, as this species was seen as a synonym of *Ca. morgani* (Crous 2002).

This study has shown the importance of combining morphological, biological and phylogenetic data to identify cryptic species of *Calonectria*. Although the biological species concept is regarded as insufficient for this purpose and needs to be clearly defined in *Calonectria* (Crous 2002), this study has shown that it has some use in identifying cryptic species within *Ca. pauciramosa*. However, morphology in combination with phylogenetic inference provides the most useful approach to identify cryptic species in *Calonectria* (Lombard *et al.* 2009). The present study has also shown the importance of the multi-gene approach in studying the phylogenetic relationships of phenotypic closely related *Calonectria* spp.

REFERENCES

- Altschul SF, Gish, W, Miller W, Myers EW, Lipman DJ. (1990). Basic Local Alignment Search Tool. *Journal of Molecular Biology* **215**: 403–410.
- Carbone I, Kohn LM. (1999). A method for designing primer sets for speciation studies in filamentous ascomycetes. *Mycologia* **91**: 553–556.
- Crous PW. (2002) Taxonomy and pathology of *Cylindrocladium* (*Calonectria*) and allied genera. APS Press, St. Paul, Minnesota, U.S.A.
- Crous PW, Alfenas AC, Junghans TG. (1998). Variability within *Calonectria ovata* and its anamorph *Cylindrocladium ovatum* from Brazil. *Sydowia* **50**: 1–13.
- Crous PW, Alfenas AC, Wingfield MJ. (1993). *Calonectria scoparia* and *Calonectria morganii* sp. nov., and variation among isolates of their *Cylindrocladium* anamorphs. *Mycological Research* **97**: 701–708.
- Crous PW, Gams W, Stalpers JA, Robert V, Stegehuis G. (2004a). MycoBank: an online initiative to launch mycology into the 21st century. *Studies in Mycology* **50**: 19–22.
- Crous PW, Groenewald JZ, Risède J-M, Simoneau P, Hyde KD. (2006). *Calonectria* species and their *Cylindrocladium* anamorphs: species with clavate vesicles. *Studies in Mycology* **55**: 213–226.
- Crous PW, Groenewald JZ, Risède J-M, Simoneau P, Hywel-Jones N. (2004b). *Calonectria* species and their *Cylindrocladium* anamorphs: species with sphaeropedunculate vesicles. *Studies in Mycology* **50**: 415–430.
- Crous PW, Phillips AJL, Wingfield MJ. (1991). The genera *Cylindrocladium* and *Cylindrocladiella* in South Africa, with special reference to forest nurseries. *South African Forestry Journal* **157**: 69–89.
- Crous PW, Wingfield MJ. (1994). A monograph of *Cylindrocladium*, including anamorphs of *Calonectria*. *Mycotaxon* **51**: 341–435.
- Fisher NL, Burgess LW, Toussoun TA, Nelson PE. 1982. Carnation leaves as a substrate and for preserving cultures of *Fusarium* species. *Phytopathology* **72**: 151–153.
- Geurber JC, Correll JC. (2001). Characterization of *Glomerella acutata*, the teleomorph of *Colletotrichum acutatum*. *Mycologia* **93**: 216–229.
- Gueidan C, Roux C, Lutzoni F. (2007). Using multigene phylogeny analysis to assess generic delineation and character evolution in *Verrucariaceae* (*Verrucariales*, *Ascomycota*). *Mycological Research* **111**: 1145–1168.

- Hawksworth DL. (2005). Two major changes in fungal nomenclature enacted in Vienna. *Mycological Research* **109**: 1061–1062.
- Halleen F, Schroers H-J, Groenewald JZ, Rego C, Oliveira H, Crous PW. (2006). *Neonectria liriodendra* sp. nov., the main causal agent of black foot disease of grapevine. *Studies in Mycology* **55**: 227–234.
- Hillis DM, Bull JJ. (1993). An empirical test of bootstrapping as a method for assessing confidence in phylogenetic analysis. *Systematic Biology* **42**: 182–192.
- Katoh K, Kuma K, Toh H, Miyata T. (2005). MAFFT version 5: improvement in accuracy of multiple sequence alignment. *Nucleic Acid Research* **33**: 511–518.
- Kioke ST, Crous PW. (2001). First report of root and crown rot of myrtle in California caused by *Cylindrocladium pauciramosum*. *Plant Disease* **85**: 448.
- Kioke ST, Henderson DM, Crous PW, Tjosvold SA. (1999). A new root and crown rot disease of heath in California caused by *Cylindrocladium pauciramosum*. *Plant Disease* **83**: 589.
- Librado P, Rozas J. (2009). DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* **25**: 1451–1452.
- Lombard L, Bogale M, Montenegro F, Wingfield BD, Wingfield MJ. (2008). A new bark canker disease of the tropical hardwood tree *Cedrelinga cateniformis* in Ecuador. *Fungal Diversity* **31**: 73–81.
- Lombard L, Rodas CA, Crous PW, Wingfield BD, Wingfield MJ. (2009). *Calonectria* (*Cylindrocladium*) species associated with dying *Pinus* cuttings. *Persoonia* **23**: 41–47.
- Mason-Gamer R, Kellogg E. (1996). Testing for phylogenetic conflict among molecular datasets in the tribe *Tiriceae* (*Graminae*). *Systematic Biology* **45**: 524–545.
- Nirenburg HI. (1981). A simplified method for identifying *Fusarium* spp. occurring on wheat. *Canadian Journal of Botany* **59**: 1599–1609.
- Nylander JAA. (2004). MrModeltest v.2. Programme distributed by the author. Evolutionary Biology Centre, Uppsala University.
- O'Donnell K, Cigelnik E. (1997). Two divergent intragenomic rDNA ITS2 types within a monophyletic lineage of the fungus *Fusarium* are nonorthologous. *Molecular Phylogenetics and Evolution* **7**: 103–116.
- O'Donnell K, Kistler HC, Cigelnik E, Ploetz RC. (1998). Multiple evolutionary origins of the fungus causing Panama disease of banana: Concordant evidence from nuclear and

- mitochondrial gene genealogies. *Proceedings of the National Academy of Sciences of the United States of America* **95**: 2044–2049.
- Overmeyer C, Lünemann S, von Wallburnn C, Meinhardt F. (1996). Genetic variability among isolates and sexual offspring of the plant pathogenic fungus *Calonectria morganii* on the basis of random amplification of polymorphic DNA (RAPD) and restriction fragment length polymorphism (RFLP). *Current Microbiology* **33**: 249–255.
- Peerally A. (1974). *Cylindrocladium brasiliense*. *CMI Descriptions of Pathogenic Fungi and Bacteria* no. **427**.
- Polizzi G. (2000). Prime esperienze di lotta chimica nei confronti del marciume del colletto e delle radici di *Polygala myrtifolia* causato da *Cylindrocladium pauciramosum*. *Informatore Fitopatologico* **11**: 39–47.
- Polizzi G, Catara V. (2001). First report of leaf spot caused by *Cylindrocladium pauciramosum* on *Acacia retinodes*, *Arbutus unedo*, *Feijoa sellowiana* and *Dodonaea viscosa* in Southern Italy. *Plant Disease* **85**: 803.
- Polizzi G, Crous PW. (1999). Root and collar of milkwort caused by *Cylindrocladium pauciramosum*, a new record for Europe. *European Journal of Plant Pathology* **105**: 407–411.
- Polizzi G, Grasso FM, Vitale A, Aiello D. (2007). First occurrence of *Calonectria* leaf spot on Mexican blue palm in Italy. *Plant Disease* **91**: 1057.
- Polizzi G, Vitale A. (2001). First report of the prevalence of benzimidazole-resistant isolates in a population of *Cylindrocladium pauciramosum* in Italy. *Plant Disease* **85**: 1210.
- Polizzi G, Vitale A, Aiello D & Parlavecchio G. (2006). First record of crown and root rot caused by *Cylindrocladium pauciramosum* on California lilac in Italy. *Plant Disease* **90**: 1459.
- Posada D, Crandall KA. (1998). Modeltest: testing the model of DNA substitution. *Bioinformatics* **14**: 817–818.
- Rayner RW. (1970). A mycological colour chart. Commonwealth Mycological Institute, Kew, Surrey. British Mycological Society.
- Ronquist F, Heulsenbeck JP. (2003). MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* **19**: 1572–1574.
- Rossmann AY, Samuels GJ, Rogerson CT, Lowen R. (1999). Genera of Bionectriaceae, Hypocreaceae and Nectriaceae (Hypocreales, Ascomycetes). *Studies in Mycology* **42**: 1–248.

- Schoch CL, Crous PW, Cronwright G, Witthuhn RC, El-Gholl NE, Wingfield BD. (2000). Recombination in *Calonectria morganii* and phylogeny with other heterothallic small-spored *Calonectria* species. *Mycologia* **92**: 665–673.
- Schoch CL, Crous PW, Polizzi G, Koike ST. (2001). Female fertility and single nucleotide polymorphism comparisons in *Cylindrocladium pauciramosum*. *Plant Disease* **85**: 941–946.
- Schoch CL, Crous PW, Wingfield BD, Wingfield MJ. (1999). The *Cylindrocladium candelabrum* species complex includes four distinct mating populations. *Mycologia* **91**: 286–298.
- Swofford DL. (2002). PAUP*. Phylogenetic analysis using parsimony (* and other methods), 4.0b10. Computer programme. Sunderland, Massachusetts, USA: Sinauer Associates.
- Taylor JW, Jacobson DJ, Kroken SM, Kasuga T, Geiser DM, Hibbett DS, Fisher MC. (2000). Phylogenetic species recognition and species concepts in fungi. *Fungal Genetics and Biology* **31**: 21–32.



Table 1. Isolates of *Calonectria pauciramosa* and other *Calonectria* species studied.

Species	Isolate	Mating system	GenBank Assession nr.			Host	Origin	Collector
			β -tubulin	Histone H3	TEF-1 α			
<i>Ca. brasiliensis</i>	CBS 230.51 ^T (= IMI 299576)		GQ267241	GQ267259	GQ267328	<i>Eucalyptus</i> sp.	Brazil	T.R. Ciferri
	CBS 114257		GQ267242	GQ267260	GQ267329	Leaf litter	Brazil	A.C. Alfenas
	CBS 116078 (= UFO 202)		GQ421772	GQ421780	GQ421788	<i>E. citriodora</i>	Brazil	A.O. Carvalho
	CMW 31505 (= CPC 2581)		GQ421775	GQ421783	GQ421791	<i>Prunus</i> sp.	South Africa	C. Linde
	CMW 31507 (= CPC 602)		GQ421773	GQ421781	GQ421789	<i>Eucalyptus</i> sp.	Brazil	P.W. Crous
	CMW 31508 (= CPC 1943)		GQ421774	GQ421782	GQ421790	Leaf litter	Brazil	A.C. Alfenas
<i>Ca. colombiana</i> sp. nov.	CBS 111136	homothallic	FJ972424	FJ972443	FJ972493	Soil	Colombia	M.J. Wingfield
	CBS 115127 ^T	homothallic	FJ972423	FJ972442	FJ972492	Soil	Colombia	M.J. Wingfield
	CBS 115638	homothallic	FJ972422	FJ972441	FJ972491	Soil	Colombia	M.J. Wingfield



Table 1. (Continued)

Species	Isolate	Mating Type	GenBank Assession nr.			Host	Origin	Collector
			β -tubulin	Histone H3	TEF-1 α			
	CBS 115694	homothallic	FJ972425	FJ972444	FJ972494	Soil	Colombia	M.J. Wingfield
	CMW 9058	homothallic	FJ972420	FJ972439	FJ972489	Soil	Colombia	M.J. Wingfield
<i>Ca. colombiensis</i>	CBS 112221		AY725620	AY725663	AY725712	Soil	Colombia	M.J. Wingfield
<i>Ca. insularis</i>	CBS 114558		AF210861	FJ918526	FJ918556	Soil	Madagascar	P.W. Crous
	CBS 114559		AF210862	FJ918525	FJ918555	Soil	Madagascar	C. L. Schoch
<i>Ca. mexicana</i>	CBS 110918 ^T		AF210863	FJ972460	FJ972526	Soil	Mexico	M.J. Wingfield
<i>Ca. morganii</i>	CBS 110666		FJ918509	FJ918527	FJ918557	<i>Ilex vomitoria</i>	USA	N.E. El-Gholl
	CBS 119669		DQ521599	DQ521601	GQ421796	<i>Pistacia lentiscus</i>	Italy	G. Polizzi
	CBS 119670		DQ521600	DQ521602	GQ421797	<i>Pistacia lentiscus</i>	Italy	G. Polizzi
	CMW 31506 (= P94-4359)		AF210875	GQ421787	GQ421795	<i>Dodenaea vicosa</i>	USA	N.E. El-Gholl
<i>Ca. pauciramosa</i>	CMW 1786	Unknown	FJ972378	FJ972445	FJ972495	<i>Eucalyptus smithii</i>	South Africa	M.J. Wingfield
	CMW 2151	Mat1-2	FJ972400	FJ972468	FJ972517	<i>E. nitens</i>	South Africa	M.J. Wingfield
	CMW 5683 ^T	Mat1-2	FJ918514	FJ918531	FJ918565	<i>E. grandis</i>	South Africa	P.W. Crous



Table 1. (Continued)

Species	Isolate	Mating Type	GenBank Assession nr.			Host	Origin	Collector
			β -tubulin	Histone H3	TEF-1 α			
<i>Ca. pauciramosa</i>	CMW 7592	Mat1-1	FJ972380	FJ972447	FJ972497	<i>E. grandis</i>	Uruguay	M.J. Wingfield
	CMW 7597	Mat1-1	FJ972406	FJ972474	FJ972523	<i>E. grandis</i>	Uruguay	M.J. Wingfield
	CMW 7600	Mat1-1	FJ972405	FJ972473	FJ972522	<i>E. grandis</i>	Uruguay	M.J. Wingfield
	CMW 7826	Mat1-2	FJ972392	FJ972459	FJ972509	Soil	Australia	P.W. Crous
	CMW 7827	Mat1-2	FJ972385	FJ972452	FJ972502	Soil	Australia	P.W. Crous
	CMW 7828	Mat1-2	FJ972391	FJ972458	FJ972508	Soil	Australia	P.W. Crous
	CMW 7849	Mat1-2	FJ972383	FJ972450	FJ972500	<i>Erica</i> sp.	USA	S.T. Koike
	CMW 7851	Mat1-2	FJ972382	FJ972449	FJ972499	<i>Myrtus communis</i>	USA	S.T. Koike
	CMW 7852	Mat1-2	FJ972381	FJ972448	FJ972498	<i>M. communis</i>	USA	S.T. Koike
	CMW 8061	Mat1-2	FJ972386	FJ972453	FJ972503	Soil	Australia	P.W. Crous
	CMW 9151	Mat1-2	FJ972384	FJ972451	FJ972501	<i>Acacia mearnsii</i>	South Africa	L. Lombard
	CMW 9172	Mat1-2	FJ972379	FJ972446	FJ972496	<i>A. mearnsii</i>	South Africa	L. Lombard
	CMW 10148	Mat1-2	FJ972387	FJ972454	FJ972504	<i>Erica</i> sp.	USA	S.T. Koike



Table 1. (Continued)

Species	Isolate	Mating Type	GenBank Assession nr.			Host	Origin	Collector
			β -tubulin	Histone H3	TEF-1 α			
<i>Ca. pauciramosa</i>	CBS 102296	Mat1-2	FJ972404	FJ972472	FJ972521	<i>Vriessea</i> sp.	New Zealand	H.M. Dance
	CBS 110945	Mat1-1	FJ972389	FJ972456	FJ972506	<i>Podocarpus</i> sp.	South Africa	P.W. Crous
	CBS 111873	Mat1-1	FJ972399	FJ972467	FJ972516	<i>Prunus</i> sp.	South Africa	C. Linde
	CBS 114861	Mat1-1	FJ972403	FJ972471	FJ972520	<i>Eucalyptus</i> sp.	South Africa	P.W. Crous
	CBS 115670	Mat1-1	FJ972393	FJ972461	FJ972510	<i>Pinus</i> sp.	South Africa	P.W. Crous
	CBS 115893	Unknown	FJ972411	FJ972430	FJ972480			
	CMW 30819	Mat1-2	FJ972402	FJ972470	FJ972519	<i>E. grandis</i>	South Africa	P.W. Crous
	CMW 30875	Mat1-1	FJ972390	FJ972457	FJ972507	<i>Eucalyptus</i> sp.	South Africa	P.W. Crous
	CMW 30823	Mat1-1	FJ918515	FJ918532	FJ918566	<i>E. grandis</i>	South Africa	P.W. Crous
	CMW 30814	Unknown	FJ972408	FJ972427	FJ972477	<i>Eucalyptus</i> sp.	Kenya	J. Roux
	CMW 30822	Unknown	FJ972409	FJ972428	FJ972478	<i>Eucalyptus</i> sp.	Kenya	J. Roux
	CMW30873	Mat1-2	FJ972388	FJ972455	FJ972505	<i>Eucalyptus</i> sp.	South Africa	L. Lombard
	CMW 27203	Mat1-2	FJ972398	FJ972466	FJ972515	<i>Eucalyptus</i> sp.	China	S. Chen



Table 1. (Continued)

Species	Isolate	Mating Type	GenBank Assession nr.			Host	Origin	Collector
			β -tubulin	Histone H3	TEF-1 α			
<i>Ca. pauciramosa</i>	CMW 27206	Mat1-2	FJ972396	FJ972464	FJ972513	<i>Eucalyptus</i> sp.	China	S. Chen
	CMW 27283	Mat1-2	FJ972397	FJ972465	FJ972514	<i>Eucalyptus</i> sp.	China	S. Chen
	CMW 30878	Mat1-1	FJ972401	FJ972469	FJ972518	<i>Prunus</i> sp.	South Africa	C. Linde
	CMW 30818	Mat1-2	FJ972395	FJ972463	FJ972512	<i>Limonium</i> sp.	New Zealand	I. Brice
	CMW 30817	Unknown	FJ972394	FJ972462	FJ972511	<i>Rhododendron</i> sp.	New Zealand	R.A.J. White
	CMW 30879	Mat1-2	FJ972407	FJ972475	FJ972524	<i>Azalea</i> sp.	Germany	G. Hagedorn
	CMW 30815	Unknown	FJ972410	FJ972429	FJ972479	<i>Eucalyptus</i> sp.	South Africa	P.W. Crous
<i>Ca. polizzii</i> sp. nov.	CBS 123402 ^T		FJ972419	FJ972438	FJ972488	<i>Arbustus unedo</i>	Italy	G. Polizzi
	CMW 7804		FJ972417	FJ972436	FJ972486	<i>Callistemon citrinus</i>	Italy	G. Polizzi
	CMW 10151		FJ972418	FJ972437	FJ972487	<i>A. unedo</i>	Italy	G. Polizzi
<i>Ca. scoparia</i>	CMW 31000		FJ972426	FJ972476	FJ97252	<i>Eucalyptus</i> sp.	Brazil	A.C. Alfenas
	CMW 31001		GQ421779	GQ267246	GQ267246	<i>Eucalyptus</i> sp.	Brazil	A.C. Alfenas
	CBS 116076		GQ421776	GQ421784	GQ421792	<i>Eucalyptus</i> sp.	Brazil	P.W. Crous



Table 1. (Continued)

Species	Isolate	Mating Type	GenBank Assession nr.			Host	Origin	Collector
			β -tubulin	Histone H3	TEF-1 α			
<i>Ca. scoparia</i>	CBS 116081		GQ421777	GQ421785	GQ421793	Soil	Brazil	M.J. Wingfield
	CMW 7578		GQ421778	GQ421786	GQ421794	<i>E. grandis</i>	Argentina	L. Lombard
<i>Ca. spathulata</i>	CBS 112689		AF308463	FJ918524	FJ918554	<i>E. viminalis</i>	Brazil	N.E. El-Gholl
	CBS555.92 ^T		GQ267215	GQ267261	GQ267331	<i>Araucaria angustifolia</i>	Brazil	C. Hodges
<i>Ca. zuluensis</i> sp. nov.	CMW 9115	homothallic	FJ972413	FJ972432	FJ972482	<i>Eucalyptus</i> sp.	South Africa	L. Lombard
	CMW 9188 ^T	homothallic	FJ972414	FJ972433	FJ972483	<i>Eucalyptus</i> sp.	South Africa	L. Lombard
	CMW 9208	homothallic	FJ972412	FJ972431	FJ972481	<i>Eucalyptus</i> sp.	South Africa	L. Lombard
	CMW 9215	homothallic	FJ972416	FJ972435	FJ972485	<i>Eucalyptus</i> sp.	South Africa	L. Lombard
	CMW 9896	homothallic	FJ972415	FJ972434	FJ972484	<i>Eucalyptus</i> sp.	South Africa	L. Lombard
<i>Cy. chinense</i>	CBS 112744		AY725618	AY725660	AY725709	Soil	China	M.J. Wingfield
<i>Cy. hawksworthii</i>	CBS 111870 ^T		AF333407	DQ190649	FJ918558	<i>Nelumbo nucifera</i>	Mauritius	A. Peeraly
<i>Cy. leucothoës</i>	CBS 109166 ^T		FJ918508	FJ918523	FJ918553	<i>Leucothoë axillaris</i>	USA	N.E. El-Gholl

CBS: Centraalbureau voor Schimmelcultures, Utrecht, The Netherlands; CMW: culture collection of the Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria Pretoria, South Africa; ^T Ex-type cultures.



Table 2. Results of mating studies between isolates of *Calonectria pauciramosa* from various geographic regions.

	CBS 102296	CBS 110945	CBS 111873	CBS 114861	CBS 115670	CMW 2151	CMW 5683	CMW 7592	CMW 7597	CMW 7600	CMW 7826	CMW 7827	CMW 7828	CMW 7849	CMW 7851	CMW 7852	CMW 8061	CMW 9151	CMW 9172	CMW 10148	CMW 27203	CMW 27206	CMW 27283	CMW 30817	CMW 30818	CMW 30819	CMW 30823	CMW 30873	CMW 30875	CMW 30878	CMW 30879			
CBS 102296	-																																	
CBS 110945	+	-																																
CBS 111873	+	-	-																															
CBS 114861	+	-	-	-																														
CBS 115670	-	+	+	+	-																													
CMW 2151	-	+	+	+	-	-																												
CMW 5683	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
CMW 7592	+	-	-	-	+	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
CMW 7597	+	-	-	-	+	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
CMW 7600	+	-	-	-	+	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
CMW 7826	-	+	+	+	-	+	-	+	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
CMW 7827	-	+	+	+	-	+	-	+	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
CMW 7828	-	+	+	+	-	+	-	+	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
CMW 7849	-	+	+	+	-	+	-	+	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
CMW 7851	-	+	+	+	-	+	-	+	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
CMW 7852	-	+	+	+	-	+	-	+	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
CMW 8061	-	+	+	+	-	+	-	+	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
CMW 9151	-	+	+	+	-	+	-	+	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
CMW 9172	-	+	+	+	-	+	-	+	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
CMW 10148	-	+	+	+	-	+	-	+	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
CMW 27203	-	+	+	+	-	+	-	+	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
CMW 27206	-	+	+	+	-	+	-	+	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
CMW 27283	-	+	+	+	-	+	-	+	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
CMW 30817	-	+	+	+	-	+	-	+	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
CMW 30818	-	+	+	+	-	+	-	+	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
CMW 30819	-	+	+	+	-	+	-	+	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
CMW 30823	+	-	-	-	+	-	+	-	-	-	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	-	-	-	-	-	-
CMW 30873	-	+	+	+	-	+	-	+	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
CMW 30875	+	-	-	-	+	-	+	-	-	-	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	-	-	-	-	-	-
CMW 30878	+	-	-	-	+	-	+	-	-	-	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	-	-	-	-	-	-
CMW 30879	-	+	+	+	-	+	-	+	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

Isolates in bold indicate *Ca. pauciramosa* mating tester strains. + indicates formation of perithecia with viable ascospores; - indicates no perithecial formation.

Table 3. Single nucleotide polymorphisms from the β -tubulin, histone H3 and translation elongation factor-1 α sequence data of *Calonectria* isolates from Colombia, Italy and South Africa.

Species	Isolate no.	β -tubulin													Histone H3													Translation elongation factor-1 α														
		87	187	200	201	208	336	375	382	385	401	406	496	514	36	39	76	209	251	257	266	274	276	281	282	292	329	350	53	86	94	95	103	126	127	128	269	423	426	515		
<i>Ca. pauciramosa</i>	CMW 5683	A	A	C	G	A	C	C	C	G	C	C	T	T	G	C	G	C	A	T	G	C	A	C	T	C	C	T	T	G	C	A	A	T	T	-	A	C	C	T		
	CMW 30823	A	A	C	G	A	C	C	C	G	C	C	T	T	G	C	G	C	A	T	G	C	A	C	T	C	C	T	T	G	C	A	A	T	T	-	A	C	C	T		
<i>Ca. colombiana</i>	CBS 111136	G	C	C	A	G	C	C	C	A	C	A	C	T	C	T	A	T	G	C	A	T	G	T	A	T	T	C	C	T	-	-	G	A	A	G	T	T	T	-	-	-
	CBS 115127	G	C	C	A	G	C	C	C	A	C	A	C	T	C	T	A	T	G	C	A	T	G	T	A	T	T	C	C	T	-	-	G	A	A	G	T	T	T	-	-	-
	CBS 115638	G	C	C	A	G	C	C	C	A	C	A	C	T	C	T	A	T	G	C	A	T	G	T	A	T	T	C	C	T	-	-	G	A	A	G	T	T	T	-	-	-
	CBS 115694	G	C	C	A	G	C	C	C	A	C	A	C	T	C	T	A	T	G	C	A	T	G	T	A	T	T	C	C	T	-	-	G	A	A	G	T	T	T	-	-	-
<i>Ca. polizii</i>	CMW 9058	G	C	C	A	G	C	C	C	A	C	A	C	T	C	T	A	T	G	C	A	T	G	T	A	T	T	C	C	T	-	-	G	A	A	G	T	T	T	-	-	-
	CBS 123402	G	A	C	G	A	T	G	T	G	G	C	T	C	C	C	G	G	A	T	G	C	A	C	T	C	C	T	T	G	C	A	A	T	T	-	A	C	C	-		
<i>Ca. zuluensis</i>	CMW 7804	G	A	C	G	A	T	G	T	G	G	C	T	C	C	C	G	G	A	T	G	C	A	C	T	C	C	T	T	G	C	A	A	T	T	-	A	C	C	-		
	CMW 10151	G	A	C	G	A	T	G	T	G	G	C	T	C	C	C	G	G	A	T	G	C	A	C	T	C	C	T	T	G	C	A	A	T	T	-	A	C	C	-		
	CMW 9115	G	A	T	G	A	C	C	C	G	C	C	T	T	C	C	G	G	A	T	G	C	A	C	T	C	C	T	T	G	C	A	A	T	T	-	A	C	C	-		
	CMW 9188	G	A	T	G	A	C	C	C	G	C	C	T	T	C	C	G	G	A	T	G	C	A	C	T	C	C	T	T	G	C	A	A	T	T	-	A	C	C	-		
	CMW 9208	G	A	T	G	A	C	C	C	G	C	C	T	T	C	C	G	G	A	T	G	C	A	C	T	C	C	T	T	G	C	A	A	T	T	-	A	C	C	-		
CMW 9215	G	A	T	G	A	C	C	C	G	C	C	T	T	C	C	G	G	A	T	G	C	A	C	T	C	C	T	T	G	C	A	A	T	T	-	A	C	C	-			
CMW 9896	G	A	T	G	A	C	C	C	G	C	C	T	T	C	C	G	G	A	T	G	C	A	C	T	C	C	T	T	G	C	A	A	T	T	-	A	C	C	-			

Yellow = unique SNP's; Green = shared SNP's



Table 4. Single nucleotide polymorphisms from the sequence data of β -tubulin, histone H3 and translation elongation factor-1 α of *Ca. brasiliensis*, *Ca. insulare* and *Ca. morgani* used in this study.

Species	Isolate no.	β -tubulin					Histone H3										Translation elongation factor-1 α																							
		53	61	117	360	472	9	10	68	69	70	94	204	252	257	359	389	416	451	460	9	10	20	47	48	102	110	112	113	114	115	142	280	378	406	407	408	414	437	479
<i>Ca. brasiliensis</i>	CBS 230.51	C	C	C	A	T	T	G	-	-	-	G	T	G	T	C	T	T	A	T	A	G	T	A	C	G	C	-	-	-	-	T	T	T	T	C	C	T	G	T
	CBS 114257	C	C	C	A	T	T	G	-	-	-	G	T	G	T	C	T	T	A	T	A	G	T	A	C	G	C	-	-	-	-	T	T	T	T	C	C	T	G	T
	CBS 116078	C	C	C	A	T	T	G	-	-	-	G	T	G	T	C	T	T	A	T	A	G	T	A	C	G	C	-	-	-	-	T	T	T	T	C	C	T	G	T
	CMW 31508	C	C	C	A	T	T	G	-	-	-	G	T	G	T	C	T	T	A	T	A	G	T	A	C	G	C	-	-	-	-	T	T	T	T	C	C	T	G	T
	CMW 31507	C	C	C	A	T	T	G	-	-	-	G	T	G	T	C	T	T	A	T	A	G	T	A	C	G	C	-	-	-	-	T	T	T	T	C	C	T	G	T
CMW 31505	C	C	C	A	T	T	G	T	T	C	A	T	G	T	C	T	T	A	T	A	G	T	A	C	G	C	-	-	-	-	T	T	T	T	C	C	T	G	T	
<i>Ca. insulare</i>	CBS114558	C	C	T	G	C	T	G	T	T	C	A	T	G	T	C	C	C	A	C	G	T	G	C	C	A	A	C	A	C	C	-	C	C	G	A	T	C	A	C
	CBS 114559	C	C	T	G	C	T	G	T	T	C	A	T	G	T	C	C	C	A	C	G	T	G	C	C	A	A	C	A	C	C	-	C	C	G	A	T	C	A	C
<i>Ca. morgani</i>	CBS 110666	A	A	T	A	T	C	T	-	-	-	A	C	T	C	T	C	C	C	C	G	T	T	A	G	G	A	-	-	-	C	-	T	C	G	A	T	C	A	C
	CBS 119669	A	A	T	A	T	C	T	-	-	-	A	C	T	C	T	C	C	C	C	G	T	T	A	G	G	A	-	-	-	C	-	T	C	G	A	T	C	A	C
	CBS 119670	A	A	T	A	T	C	T	-	-	-	A	C	T	C	T	C	C	C	C	G	T	T	A	G	G	A	-	-	-	C	-	T	C	G	A	T	C	A	C
	CMW 31506	A	A	T	A	T	C	T	-	-	-	A	C	T	C	T	C	C	C	C	G	T	T	A	G	G	A	-	-	-	C	-	T	C	G	A	T	C	A	C

Yellow = unique SNP's; Green = shared SNP's

Fig. 1. One of eight most parsimonious trees obtained from a heuristic search with 1 000 random addition of the combined BT, HIS3 and TEF-1 α sequence alignments. Scale bar shows 10 changes and bootstrap support values from 1 000 replicates are shown above the nodes in bold. Bayesian posterior probability values are indicated below the nodes. Red lines indicate bootstrap support values of 100 and posterior probability values of 1.00. Thickened lines indicate branches in the strict consensus and Bayesian consensus tree. The tree was rooted to *Calonectria colombiensis* (CBS 112221) and *Cylindrocladium chinense* (CBS 112744). Mating tester strains of *Ca. pauciramosa* used in this study are indicated in bold.



Fig. 2. *Calonectria brasiliensis*. A–E. Anamorph state of *Ca. brasiliensis*. A. Macroconidiophore. B–C. Vesicles. D. Fertile branches of the conidiophore with doliiform to reniform phialides. E. Macroconidia. Scale bars in A = 20 μm , B–E = 10 μm .




Fig. 3. *Calonectria colombiana*. A–F. Teleomorph state of *Ca. colombiana*. G–L. Anamorph state of *Ca. colombiana*. A. Perithecium on toothpick.. B. A 10 µm thick vertical section through perithecium. C. Section through perithecial wall. D. Ostiolar region of perithecium. E. Asci and ascospores. F. Ascospore. G–H. Macroconidiophores of *Ca. colombiana*. I–J. Vesicles. K. Fertile branches of the conidiophore with doliiform to reniform phialides. L. Macroconidia. Scale bars: A = 50 µm, B, D = 20 µm, C, E–H = 10 µm.






Fig. 4. *Calonectria polizzii*. A–E. Anamorph state of *Ca. polizzii*. A. Macroconidiophore. B–C. Vesicles. D. Fertile branches of the conidiophore with doliiform to reniform phialides. E. Macroconidia. Scale bars in A = 20 μm , B–E = 10 μm .





Fig. 5. *Calonectria zuluensis*. A–F. Teleomorph state of *Ca. zuluensis*. G–L. Anamorph state of *Ca. zuluensis*. A. Perithecium on toothpick.. B. A 10 µm thick vertical section through perithecium. C. Section through perithecial wall. D. Ostiolar region of perithecium. E. Asci and ascospores. F. Ascospore. G–H. Macroconidiophores of *Ca. zuluensis*. I–J. Vesicles. K. Fertile branches of the conidiophore with doliiform to reniform phialides. L. Macroconidia. Scale bars: A = 50 µm, B, D = 20 µm, C, E–H = 10 µm.





Chapter 5

Phylogeny and systematics of the genus *Calonectria*

Submitted as: Lombard L, Crous PW, Wingfield BD, Wingfield MJ. (2009). Phylogeny and systematics of the genus *Calonectria*. *Studies in Mycology*.

ABSTRACT

Species of *Calonectria* are important plant pathogens, several of which have a worldwide distribution. Contemporary taxonomic studies on these fungi have chiefly relied on DNA sequence comparisons of the β -tubulin gene region. Despite many new species being described, there has been no phylogenetic synthesis for the group since the last monographic study almost a decade ago. In this study, the identity of a large collection of *Calonectria* isolates from various geographic regions was determined using morphological and DNA sequence comparisons. This resulted in the discovery of seven new species; *Ca. densa*, *Ca. eucalypti*, *Ca. humicola*, *Ca. orientalis*, *Ca. pini*, *Ca. pseudoscoparia* and *Ca. orientalis*, bringing the total number of currently accepted *Calonectria* species to 68. A multigene phylogeny was subsequently constructed for all available *Calonectria* spp., employing seven gene regions, namely actin, β -tubulin, calmodulin, histone H3, the internal transcribed spacer regions 1 and 2 and the 5.8S gene of the ribosomal RNA, 28S large subunit RNA gene and translation elongation 1-alpha. Based on these data 13 phylogenetic groups could be distinguished within the genus *Calonectria* that correlated morphological features. Dichotomous and synoptic keys to all *Calonectria* spp. currently recognised are also provided.

Taxonomic novelties: New combinations *Calonectria angustata* (Crous & El-Gholl) L. Lombard, M.J. Wingf. & Crous, *Calonectria australiensis* (Crous & H.D. Hyde) L. Lombard, M.J. Wingf. & Crous, *Calonectria canadensis* (J.C. Kang, Crous & C.L. Schoch) L. Lombard, M.J. Wingf. & Crous, *Calonectria chinensis* (Crous) L. Lombard, M.J. Wingf. & Crous, *Calonectria citri* (Boedjin & Reitsma) L. Lombard, M.J. Wingf. & Crous, *Calonectria curvata* (Boedjin & Reitsma) L. Lombard, M.J. Wingf. & Crous, *Calonectria curvispora* (Crous & D. Victor) L. Lombard, M.J. Wingf. & Crous, *Calonectria ecuadoriae* (Crous & M.J. Wingf.) L. Lombard, M.J. Wingf. & Crous, *Calonectria gordoniae* (Leahy, T.S. Schub. & El-Gholl) L. Lombard, M.J. Wingf. & Crous, *Calonectria hawksworthii* (Peerally) L. Lombard, M.J. Wingf. & Crous, *Calonectria hurae* (Linder & Whetzel) L. Lombard, M.J. Wingf. & Crous, *Calonectria indonesiae* (Crous) L. Lombard, M.J. Wingf. & Crous,

Calonectria leucothoës (El-Gholl, Leahy & T.S. Schub.) L. Lombard, M.J. Wingf. & Crous,
Calonectria malesiana (Crous) L. Lombard, M.J. Wingf. & Crous, *Calonectria*
multiplialidica (Crous, P. Simoneau & J.-M. Risède) L. Lombard, M.J. Wingf. & Crous,
Calonectria pacifica (J.C. Kang, Crous & C.L. Schoch) L. Lombard, M.J. Wingf. & Crous,
Calonectria penicilloides (Tubaki) L. Lombard, M.J. Wingf. & Crous, *Calonectria*
pseudonaviculata (Crous, J.Z. Groenew. & C.F. Hill) L. Lombard, M.J. Wingf. & Crous,
Calonectria sumatrensis (Crous) L. Lombard, M.J. Wingf. & Crous. **New species:**
Calonectria densa L. Lombard, M.J. Wingf. & Crous, *Calonectria eucalypti* L. Lombard,
M.J. Wingf. & Crous, *Calonectria humicola* L. Lombard, M.J. Wingf. & Crous, *Calonectria*
orientalis L. Lombard, M.J. Wingf. & Crous, *Calonectria pini* L. Lombard, M.J. Wingf. &
Crous, *Calonectria pseudoscoparia* L. Lombard, M.J. Wingf. & Crous, *Calonectria*
sulawesiensis L. Lombard, M.J. Wingf. & Crous.

INTRODUCTION

The genus *Calonectria* (*Ca.*) was first described in 1867, with *Ca. daldiniana* as the type. This species was later reduced to synonymy with *Ca. pyrochroa* based on morphological comparisons done by Rossman (1979). *Calonectria* spp. are Euascomycetes in the order *Hypocreales* (Hibbett *et al.* 2007, Schoch *et al.* 2009) and are characterised by their yellow to dark red perithecia, with scaly to warty ascocarp walls giving rise to long-stalked, clavate asci with 1-multi-septate ascospores and *Cylindrocladium* (*Cy.*) anamorph states (Rossman 1993, Crous 2002). The genus *Cylindrocladium* was described by Morgan (1892), and is characterised by branched conidiophores with stipe extensions terminating in characteristic vesicles and producing cylindrical, 1-multi-septate conidia (Crous & Wingfield 1994, Crous 2002). Morphologically, the anamorph state provides the greatest number of distinguishing characters for *Calonectria* and it is also the state most frequently encountered in nature (Peerally 1991, Crous & Wingfield 1994, Schoch *et al.* 2001b, Crous 2002).

Species of *Calonectria* are primarily distinguished based anamorph characters, such as vesicle shape, stipe extension length, conidial septation, and dimensions on a standardised medium under defined growth conditions (Boesewinkel 1982, Peerally 1991, Crous & Wingfield 1994, Crous 2002). Despite the use of standardised conditions, taxonomic confusion can result because some intraspecific variation in vesicle shape and conidial dimension is common (Crous & Peerally 1996, Crous *et al.* 1998a). Although the reliability of vesicle shape as a distinguishing morphological character has been questioned (Sober & Alfieri 1972, Hunter & Barnett 1978, Rossman 1983), Crous *et al.* (1992) demonstrated experimentally that the shape of this structure can be influenced by the osmotic potential of the medium and the age of the culture, but that it remains a reliable morphological feature if these aspects are standardised. In the original description of *Ca. morganii* (= *Cy. scoparium*), the type of the anamorph state, Morgan (1892) failed to include details of the stipe extension and terminal vesicle, which is a defining characteristic in distinguishing anamorphs of *Calonectria* (Boesewinkel 1982, Peerally 1991, Crous & Wingfield 1994, Crous 2002).

Calonectria spp. produce three different morphological forms of conidia, of which the macroconidia are present in all but *Ca. multiseptata* (Peerally 1991, Crous & Wingfield 1994, Crous *et al.* 1998b, Crous 2002). Mega- and microconidia are less frequently encountered and

these are not regarded as important characters to distinguish between species (Sober 1971, Crous & Wingfield 1994, Crous & Seifert 1998, Crous 2002).

Both homothallic and heterothallic mating systems are found amongst species of *Calonectria* (Alfieri *et al.* 1982, Schubert *et al.* 1989, Crous & Wingfield 1994, Crous 2002). Heterothallic *Calonectria* spp. have a biallelic heterothallic mating system with the female structures (protoperithecia) spermatized by conidia or hyphae of an opposite mating type strain (Schoch *et al.* 1999, 2000a, 2001a). Some *Calonectria* spp. have retained the ability to recombine with other closely related *Calonectria* spp., although the progeny from these crosses have low levels of fertility (Crous 2002). This has complicated the application of the biological species concept for *Calonectria*, although it has been useful for some species (Schoch *et al.* 1999, Lombard *et al.* 2009a).

Several molecular approaches have been employed to identify *Calonectria* spp. These include total protein electrophoresis (Crous *et al.* 1993a, El-Gholl *et al.* 1993), isozyme electrophoresis (El-Gholl *et al.* 1992, 1997, Crous *et al.* 1998a), random amplification of polymorphic DNA (RAPD) (Overmeyer *et al.* 1996, Victor *et al.* 1997, Schoch *et al.* 2000a, Riséde & Simoneau 2004) restriction fragment length polymorphisms (RFLP) (Crous *et al.* 1993b, Crous *et al.* 1995, Crous *et al.* 1997, Jeng *et al.* 1997, Victor *et al.* 1997; Riséde & Simoneau 2001) and DNA hybridization (Crous *et al.* 1993a, 1995, 1997, Victor *et al.* 1997). However, DNA sequence comparisons and associated phylogenetic inference has had the most significant impact on the taxonomy of the group. It is also most widely applied in contemporary species descriptions. The 5.8S ribosomal RNA gene and flanking internally transcribed spacer (ITS) sequences made it possible for Jeng *et al.* (1997) to distinguish between *Cy. scoparium* and *Cy. floridanum* isolates. Subsequently, it was found that this gene region contains few informative characters for members of the genus (Crous *et al.* 1999, Schoch *et al.* 1999, Riséde & Simoneau 2001, Schoch *et al.* 2001b). As a consequence, this resulted in the β -tubulin (BT) (Schoch *et al.* 2001b) and histone H3 (HIS3) (Kang *et al.* 2001b) gene regions being widely employed to improve the resolution of phylogenetic trees for species of *Calonectria*.

The first complete DNA sequence-based phylogenetic study using partial BT gene sequences (Schoch *et al.* 2001b) compared phenotypic, biological and phylogenetic species concepts used in the taxonomy of *Calonectria*. Results showed that the genus represents a well

resolved monophyletic lineage. Subsequently, combined DNA sequence data for the ITS, BT and HIS3 gene regions have been used to resolve taxonomic questions for *Calonectria* (Schoch *et al.* 2000a, Henricot & Culham 2002, Crous *et al.* 2004b, 2006). Other DNA sequences recently used to distinguish between species include the translation elongation factor 1-alpha (TEF-1 α) and calmodulin (CAL) gene regions (Crous *et al.* 2004b). However, sequence data for these regions on GenBank (www.ncbi.nlm.nih.gov) are incomplete for the group, substantially reducing their value.

The aim of this study was to consider the identity of a large collection of previously unidentified *Calonectria* isolates collected over a five year period from various parts of the world. Morphological characteristics, phylogenetic inference and mating compatibility were employed for this purpose. Subsequently, the phylogenetic relationships between *Calonectria* spp. were re-evaluated by constructing a multigene phylogeny for seven gene regions and considering these results together with morphological features for all species in the genus.

MATERIALS AND METHODS

Isolates

Plant material showing symptoms of *Calonectria* infections as well as soil samples were collected from various geographical regions over a period of five years. Diseased plant material was placed in moist chambers and incubated for 48 h at room temperature to induce sporulation. Direct isolations were made onto malt extract agar (2 % w/v; MEA; Biolab, Midrand, South Africa) and cultures were incubated for 7 d at 25 °C under continuous near-ultraviolet light. Baiting, using *Medicago sativa* seed, was applied for the soil samples following the technique of Crous (2002). For each isolate, single conidial cultures were prepared on MEA. Representative strains are maintained in the culture collection (CMW) of the Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria, South Africa and the Centraalbureau voor Schimmelcultures (CBS), Utrecht, The Netherlands (Table 1).

DNA extraction and amplification

Identification of unknown *Calonectria* isolates: Total genomic DNA was extracted from 7 d old *Calonectria* cultures using the methods presented in Lombard *et al.* (2008). Three loci were amplified and sequenced. These included a fragment of the BT gene region using primers T1 (O'Donnell & Cigelnik 1997) and CYLTUB1R (Crous *et al.* 2004b), a fragment

of the HIS3 gene region using primers CYLH3F and CYLH3R (Crous *et al.* 2004b) and a fragment of the TEF-1 α gene region using primers EF1-728F (Carbone & Kohn 1999) and EF2 (O'Donnell *et al.* 1998).

Phylogenetic relationships amongst *Calonectria* spp.: Total genomic DNA was extracted as above. Seven loci were amplified including the ITS gene region using primers V9G (De Hoog & van den Ende 1998) and ITS4 (White *et al.* 1990); the 28S large subunit RNA gene (LSU) using primers LR0R (Moncalvo *et al.* 1995) and LR5 (Vilgalys & Hester 1990); and parts of the TEF-1 α gene region; the BT gene region, the HIS3 gene region with the same primer sets mentioned previously, the actin (ACT) gene region using primers ACT-512F and ACT-783R (Carbone & Kohn 1999) and CAL gene region using primers CAL-228F and CAL-737R (Carbone & Kohn 1999).

The PCR reaction mixture used to amplify the different loci consisted of 2.5 units FastStart *Taq* polymerase (Roche Applied Science, USA), 10 \times PCR buffer, 1–1.5 mM MgCl₂, 0.25 mM of each dNTP, 0.5 μ M of each primer and approximately 30 ng of fungal genomic DNA, made up to a total reaction volume of 25 μ L with sterile deionised water. Amplified fragments were purified using High Pure PCR Product Purification Kit (Roche, U.S.A.)

DNA sequencing and analysis

Amplified fragments were sequenced in both directions using the same primer pairs used for amplification. For this purpose, the BigDye terminator sequencing kit v. 3.1 (Applied Biosystems, U.S.A.) and an ABI PRISMTM 3100 DNA sequencer (Applied Biosystems) were used. All PCRs and sequencing reactions were performed on an Eppendorf Mastercycler Personal PCR (Eppendorf AG, Germany) with cycling conditions as described in Crous *et al.* (2006) for all loci amplified.

In addition to the sequences generated in this study, *Calonectria* spp. sequences were obtained from GenBank. All sequences were assembled and aligned using Sequence Navigator v. 1.0.1 (Applied Biosystems) and MAFFT v. 5.11 (Katoh *et al.* 2005), respectively. The aligned sequences were then manually corrected where necessary. Single nucleotide polymorphisms (SNP's) were determined for the aligned DNA sequences of each gene region using DnaSP v. 5.00.06 (Librado & Rozas 2009)

To determine whether the DNA sequence data sets were congruent, a partition homogeneity test (PHT; Farris *et al.* 1994) of all possible combinations, with a 1 000 replications on all informative characters was conducted in PAUP (Phylogenetic Analysis Using Parsimony, v. 4.0b10; Swofford 2002). A 70 % reciprocal bootstrap method using Neighbour-Joining with Maximum Likelihood distance (Mason-Gamer & Kellogg 1996; Gueidan *et al.* 2007) was also employed. Models of evolution were estimated in Modeltest v. 3.7 (Posada & Crandall 1998) using the Akaike Information Criterion (AIC) for each gene region. The bootstrap analyses were run in PAUP for 10 000 replicates. Resulting tree topologies were compared visually for conflict between the separate gene regions.

Maximum-parsimony genealogies, for single genes and the combined genes, were estimated in PAUP, by heuristic searches based on 1 000 random addition sequences and tree bisection-reconnection, with the branch swapping option set on “best trees” only. All characters were weighted equally and alignment gaps were treated as missing data. Statistics calculated for parsimony included tree length (TL), consistency index (CI), retention index (RI) and rescaled consistence index (RC). Bootstrap analysis (Hillis & Bull 1993) was based on 1 000 replications. All sequences for the isolates studied were analysed using the Basic Local Alignment Search Tool for Nucleotide sequences (BLASTN, Altschul *et al.* 1990).

A Markov Chain Monte Carlo (MCMC) algorithm was used to generate phylogenetic trees for each gene region and combined sequence data subsets with Bayesian probabilities using MrBayes v. 3.1.1 (Ronquist & Huelsenbeck 2003). Models of nucleotide substitution for each gene were determined using MrModeltest (Nylander 2004) and included for each gene partition. Four MCMC chains were run simultaneously from random trees for one million generations, sampled every 100 generations and repeated twice. Both runs converged on the same likelihood score and tree topology for each gene. The first 1 000 trees were, therefore, discarded as the burn-in phase of each analysis and posterior probabilities were determined from the remaining trees.

Sexual compatibility

Based on the results of the DNA sequence analyses, single conidial isolates of *Calonectria* spp. of unknown identity were crossed with closely related species in all possible combinations. Where available, mating tester strains defined in previous studies were also used. Crosses were made as described in Schoch *et al.* (1999) on carnation leaf agar (CLA;

Fisher *et al.* 1982, Crous *et al.* 1993a) and minimal salt agar (MSA; Guerber & Correll 2001, Halleen *et al.* 2006) with sterile toothpicks placed on the surface of the agar (Lombard *et al.* 2009a). Controls were of isolates crossed with themselves, making it possible to distinguish between those having heterothallic or homothallic mating systems. Isolates CBS 125273-125276 from Indonesia were mated with *Ca. macroconidialis* (CBS 114880). Colombian isolates CBS 123698 and CBS 125523 and Indonesian isolates CBS 125258-125260 were crossed with *Ca. brachiatica* (CBS 123700 and CMW 25302) and *Ca. brassicae* (CBS 111478 and CBS 111869) in all possible combinations. Isolates CBS 125248, CBS 125253, CBS 125277 and CMW 14883 were crossed with *Ca. cerciana* (CBS 123693 and CBS 123695), *Ca. brasiliensis* (CBS 230.51 and CBS 114257) and mating tester strains of *Ca. insularis* (CBS 114558 and CBS 114559; Schoch *et al.* 1999). Similarly, isolates CBS 125249-125252, CBS 125261 and CBS 125269 were crossed with mating tester strains of *Ca. spathiphylli* (CBS 114540 and CBS 116168; Crous 2002). Isolates CBS 125254-125257 were crossed with mating tester strains of *Ca. scoparia* (CMW 31000 and CMW 31001; Lombard *et al.* 2009a) and *Ca. pauciramosa* (CMW 5683 and CMW 30823; Schoch *et al.* 2001a). The plates were stacked in plastic containers and incubated at 22 °C for 6–8 wk. Crosses were regarded as successful when isolate combinations produced numerous perithecia extruding viable ascospores.

Taxonomy

For identification of *Calonectria* isolates based on morphology, single conidial cultures were prepared on MEA and synthetic nutrient-poor agar (SNA; Nirenburg 1981, Lombard *et al.* 2009b). Inoculated plates were incubated at room temperature and examined after 7 d. Gross morphological characteristics of the anamorph structures were determined by mounting fungal structures in lactic acid and 30 measurements at ×1 000 magnification were made for all taxonomically informative characters for each isolate. Teleomorph morphology was determined by mounting perithecia resulting from the sexual compatibility tests in Leica mountant (Setpoint Premier, Johannesburg, South Africa) and making sections using a Leica CM1100 cryostat (Setpoint Technologies) at –20 °C. The 10 µm sections were mounted in lactophenol or 3 % KOH. Gross morphological characteristics were determined in the same manner as for the anamorph states. The 95 % confidence levels were determined and extremes of conidial measurements are given in parentheses. For other structures, only extremes are presented in the descriptions.

Optimal growth conditions for cultures were determined in the dark on MEA for each isolate, at temperatures ranging from 5–35 °C at 5 °C intervals with three replicate plates for each temperature tested. Two measurements of culture diameter perpendicular to each other were made daily for 7 d. Colony colours were determined after 7 d on MEA at 25 °C in the dark, using the colour charts of Rayner (1970). Descriptions, nomenclature and illustrations were deposited in MycoBank (Crous *et al.* 2004a).

RESULTS

DNA sequencing and analysis

Identification of unknown *Calonectria* isolates: Amplicons of approx. 500 bp were generated for the BT and TEF-1 α gene regions and those for the HIS3 region were approx. 450 bp in length. Based on preliminary BT sequence comparisons and morphological characteristics, the sequence data sets for the unknown *Calonectria* spp. were divided into four separate data sets representing the *Ca. colhounii*, *Ca. brassicae*, *Ca. scoparia* and *Ca. morganii* complexes and other closely related species in each data set. These data sets were analysed separately with *Ca. colombiensis* (CBS 112221) and *Ca. chinensis* (CBS 112744) as outgroup taxa. For Bayesian analyses, a HKY+I+G model was selected for BT and TEF-1 α , and GTR+I+G for HIS3 for all four data sets, which was incorporated in the analyses. The consensus trees obtained from the Bayesian analyses confirmed the tree topologies obtained with maximum-parsimony as well as bootstrap support. Therefore, only maximum-parsimony trees are presented with bootstrap values and posterior probabilities shown for well-supported branches.

The partition homogeneity tests for all possible combinations of the three gene regions used, consistently yielded a P-value of 0.001 for the four separate data sets. The 70 % reciprocal bootstrap trees showed no conflict in tree topologies for the three gene regions in each of the four separate data sets. Based on the tree topologies of the 70 % reciprocal bootstrap trees and a P-value of 0.001 in the PHT test (Cunningham 1997, Dettman *et al.* 2003) the DNA sequences for the three gene regions were combined for each of the four separate data sets.

The combined sequence data set representing the *Ca. colhounii* complex, with 10 taxa including outgroups, consisted of 1 497 characters, including gaps. Of these characters, 1 051 were constant, 133 were parsimony-uninformative and 313 characters were parsimony informative. Parsimony analysis of the aligned sequences yielded one most parsimonious tree

(Fig. 1; TL = 649 steps; CI = 0.888; RI = 0.891; RC = 0.791). In the tree, isolates CBS 125273, CBS 125274, CBS 125275 and CBS 125276, from Indonesia, grouped close to but separate from *Ca. colhounii* (CBS 293.79 and CBS 114704) with 100 % bootstrap support (BP) and a posterior probability (PP) of 0.97. The SNP analyses showed 16 unique alleles for the Indonesian isolates with one shared unique allele with *Ca. madagascariensis* (CBS 114571 and CBS 114572) and two shared alleles with *Ca. macroconidialis* (CBS 114880) for the three gene regions analysed (Table 2). These unique alleles, however, distinguish the Indonesian isolates from *Ca. colhounii*, *Ca. macroconidialis* and *Ca. madagascariensis*.

The data set representing the *Ca. brassicae* complex consisted of 15 taxa including the outgroups, while the combined sequence alignment was made up of 1 509 characters, including gaps. These characters represented 1 092 constant, 127 parsimony-uninformative and 290 parsimony-informative characters. Parsimony analysis yielded one most parsimonious tree (Fig. 2; TL = 569 steps; CI = 0.931; RI = 0.918; RC = 0.855). In the tree, Colombian isolates CBS 123698 and CBS 125523 clustered close to *Ca. brassicae* (CBS 111869 and CBS 111478) and *Ca. brachiatica* (CBS 123700 and CMW 25302) but separately from both these species with high support (BP = 100 and PP = 1.00). Similarly, isolates CBS 125258, CBS 125259 and CBS 125260, from Indonesia, clustered together closely related to *Ca. brassicae* and *Ca. brachiatica*. These Indonesian isolates were also closely related to the Colombian isolates but grouped separately from them in a clade with high support (BP = 97 and PP = 1.00). The SNP analyses showed that isolates CBS 123698 and CBS 125523 have 18 unique alleles and isolates CBS 125258, CBS 125259 and CBS 125260 have four unique alleles distinguishing them from each other for the three gene regions analysed. These isolates also share 14 unique alleles, distinguishing them from *Ca. brassicae* and *Ca. brachiatica* (Table 3).

The third data set, represented by 16 ingroup taxa residing in the *Ca. scoparia* complex and closely related species, consisted of 1 530 characters including gaps for the three gene regions analysed. Of these characters, 1 114 were constant, 138 were parsimony-uninformative and 278 characters were parsimony informative. Parsimony analysis of the aligned sequences yielded two most parsimonious trees (TL = 551 steps; CI = 0.902; RI = 0.925; RC = 0.834), one of which is presented in Fig 3. In the tree, isolates CBS 125254, CBS 125255, CBS 125256 and CBS 125257 from Ecuador, clustered closely but separately from *Ca. scoparia* (CMW 31000 and CMW 31001) and other species in the *Ca. pauciramosa*

complex with low support (BP = 63 and PP = 1.00). The Ecuadorian isolates also had three unique alleles separating them from *Ca. scoparia* and *Ca. pauciramosa* (CMW 5683 and CMW 30823) for the BT and TEF-1 α regions, but there were no unique alleles for these isolates in the HIS3 region (Table 4).

The aligned sequence data set for the *Ca. morganii* complex included 25 ingroup taxa consisting of 1 535 characters. Of these characters, 975 were constant, 211 were parsimony-uninformative and 349 characters were parsimony-informative. Parsimony analysis of the aligned sequences yielded three most parsimonious trees (TL = 977 steps; CI = 0.784; RI = 0.825; RC = 0.647), one of which is presented in Fig 4. In the tree, isolates CBS 125249, CBS 125250, CBS 125251, CBS 125252, CBS 125261 and CBS 125269 from Ecuador clustered in a clade (BP = 99 and PP = 1.00) with *Ca. spathiphylli* (CBS 114540 and CBS 116168) and *Ca. pseudospathiphylli* (CBS 109165), whereas isolates CBS 125248, CBS 125253, CBS 125277 and CMW 14883 from Indonesia clustered close to *Ca. brasiliensis* (CBS 230.51 and CBS 114257) but with low support (BP = 52; PP = 0.90) in a separate, well-supported clade (BP = 100; PP = 1.00). Isolates CBS 125249, CBS 125250 and CBS 125261 clustered together in a well-supported clade (BP = 93; PP = 1.00) separate from CBS 125251, CBS 125252 and CBS 125269, that also clustered together in a well-supported clade (BP = 81; PP = 1.00). Both clades were separate from *Ca. spathiphylli* and *Ca. pseudospathiphylli* but closely related to these species. The SNP analyses showed that isolates CBS 125249, CBS 125250 and CBS 125261 shared four unique alleles and CBS 125251, CBS 125252 and CBS 125269 shared seven unique alleles for the three gene regions. These isolates also shared an additional 33 alleles, distinguishing them from *Ca. spathiphylli* (Table 5). Isolates CBS 125248, CBS 125253, CBS 125277 and CMW 14883 shared eight unique alleles, distinguishing them from *Ca. brasiliensis* (CBS 230.51 and CBS 114257), *Ca. cerciana* (CBS 123693 and CBS 123695) and *Ca. insularis* (CBS 114558 and CBS 114559) (Table 6).

Phylogenetic relationships amongst *Calonectria* spp.: Approximately 250 bases were determined for ACT, 450 bases for HIS3, 500 for BT, CAL and TEF-1 α , 700 for ITS and 880 for LSU. The adjusted sequence alignments for each gene region consisted of 122 ingroup taxa with *Cylindrocladiella lageniformis* (CBS 112898) and *C. peruviana* (CPC 5614) as outgroup taxa for each gene region. For Bayesian analyses, a K80+G model was selected for ACT, HKY+I+G for BT, CAL and TEF-1 α , GTR+I+G for HIS3 and LSU, and SYM+I+G

for ITS and incorporated in the analyses. The consensus trees obtained from the Bayesian analyses confirmed the tree topologies obtained with maximum-parsimony as well as bootstrap support.

Individual analyses of the gene regions showed similar tree topologies for the protein coding regions (ACT, BT, CAL, HIS3 and TEF-1 α) with well-supported clades for *Calonectria* spp. with similar morphological characteristics. In contrast, the non-coding gene regions (ITS and LSU) provided little or no support for the clades that emerged from the protein coding regions, with several *Calonectria* spp. clustering together with no significant similarities. The trees for the ITS and LSU regions showed a single monophyletic clade for all *Calonectria* spp. and did not reveal the two clades observed for the coding gene regions. The phylogeny constructed based on CAL sequences showed the best resolution of the species and it had the highest support for the individual clades, followed by TEF-1 α gene region. Statistical data for the individual trees (not shown) are presented in Table 7.

The partition homogeneity tests for all possible combinations of the seven gene regions used, consistently yielded a P-value of 0.001. The 70 % reciprocal bootstrap trees showed no conflict in tree topologies for the five coding gene regions (ACT, BT, CAL, HIS3 and TEF-1 α), however conflicts were observed between the non-coding gene regions (ITS and LSU) and the coding gene regions. Based on the tree topologies and a P-value of 0.001 (Cunningham 1997, Dettman *et al.* 2003) the sequence data for coding gene regions were combined. The data for the ITS and LSU datasets were treated separately, but these are not presented because they add little taxonomic value. However, all ITS and LSU sequences generated in this study have been deposited in GenBank and TreeBase as SN4777 (Table 1).

The combined sequence alignment of the five coding gene regions consisted of 2 472 characters, including gaps. Of these characters, 925 were constant, 267 were parsimony-uninformative and 1 280 characters were parsimony informative. Parsimony analysis of the aligned sequences yielded 24 most parsimonious trees (TL = 7319 steps; CI = 0.397; RI = 0.820; RC = 0.326), one of which is presented in Fig. 5. The tree topology obtained with the combined sequence dataset was similar to that obtained for the individual gene regions analysed and therefore the only tree presented is that of the combined dataset.

In the tree (Fig. 5), the *Calonectria* spp. were found to clearly reside in two main clades which was consistent for the analyses for these gene regions separately. One of these clades

(BP = 82, PP = 0.62) which we refer to as representing the Prolate Group, includes *Calonectria* spp. with clavate to pyriform to ellipsoidal vesicles. This clade (Fig. 5) is made up of two sub-clades, one (BP = 81, PP = 1.00) of which includes 10 minor clades representing *Calonectria* spp. that have vesicles and conidia that have similar morphology. The second sub-clade (BP = 99, PP = 1.00) representing the Prolate Group includes taxa represented by single isolates and for which there were no obvious unifying morphological characters.

The second main clade (BP = 65, PP = 0.64) which is referred to as the Sphaero-Naviculate Group of species included *Calonectria* spp. characterised by sphaeropedunculate and naviculate vesicles and these were also seen in the analyses based on the individual gene regions. This clade is further sub-divided into two clades. The first of these sub-clades (BP = 65, PP = 1.00) includes *Calonectria* spp. characterised by sphaeropedunculate vesicles. The second sub-clade (BP = 93, PP = 0.86) accommodates *Calonectria* spp. with naviculate vesicles.

Sexual compatibility

The only isolates in the mating tests that yielded perithecia were CBS 125273, CBS 125274, CBS 125275 and CBS 125276 (Fig. 6). These isolates all produced perithecia containing viable ascospores within 6 wk when mated with themselves, indicating that they are self-fertile (homothallic). All other control inoculations with the selected isolates failed to yield perithecia, indicating that they were either self-sterile (heterothallic) and non-compatible, or that they had lost the ability to undergo sexual recombination.

Taxonomy

Based on morphological observations, phylogenetic inference and mating, numerous isolates of unknown *Calonectria* spp. included in this study represent undescribed species. Species of *Cylindrocladium* (1892) represent anamorph states of *Calonectria* (1867) (Rossmann *et al.* 1999). In an attempt to move to a single nomenclature for pleomorphic fungi (Hawksworth 2005), the teleomorph name takes precedence over the anamorph name when both types belong to the same holomorph. Subsequently, the species below are described as new species in *Calonectria*, which represents the older generic name for these holomorphs. All *Cylindrocladium* species without a *Calonectria* state, are subsequently also transferred to *Calonectria*.

Calonectria densa L. Lombard, M.J. Wingf. & Crous, **sp. nov.** MycoBank MB515529, Fig. 7.

Etymology: Name refers to the fact that lateral stipe extensions are readily formed in this species and that are absent from the most closely related species, giving it a bushy appearance.

Teleomorpha ignota. Anamorpha *Cy. spathiphylli* similis sed extensiones laterales stiparum facit, macroconidiis cylindricis utrinque rotundatis rectis (47–)50–58(–62) × 5–6 µm mediocriter 54 × 6 µm, semel septatis, sine cicatrice abscissionis manifesta, cum muco hyalino in fasciculis parallelis cylindricis differ.

Teleomorph unknown. *Conidiophores* with a stipe bearing penicillate suites of fertile branches, stipe extensions, and terminal vesicles; stipe septate, hyaline, smooth, 54–90 × 6–10 µm; stipe extensions septate, straight to flexuous, 149–192 µm long, 5–6 µm wide at the apical septum, terminating in a globose to ovoid to sphaeropedunculate vesicle, 10–12 µm diam; lateral stipe extensions (90° to the axis) also present. *Conidiogenous apparatus* 49–78 µm long, and 63–123 µm wide; primary branches aseptate, 20–29 × 5–6 µm; secondary branches aseptate, 16–20 × 4–6 µm; tertiary and additional branches (–4) aseptate, 9–16 × 3–5 µm, each terminal branch producing 2–4 phialides; phialides doliiform to reniform, hyaline, aseptate, 11–16 × 2–4 µm; apex with minute periclinal thickening and inconspicuous collarete. *Macroconidia* cylindrical, rounded at both ends, straight, (47–)50–58(–62) × 5–6 µm (av. = 54 × 6 µm), 1-septate, lacking a visible abscission scar, held in parallel cylindrical clusters by colourless slime. *Mega-* and *microconidia* not seen.

Specimens examined: **Ecuador**, Pichincha Province, Las Golondrinas, from soil, Dec. 2004, M.J. Wingfield, Herb. PREM 60302, **holotype** of *Ca. densa*, culture ex-type CMW 31182 = CBS 125261; Pichincha Province, Las Golondrinas, from soil, Dec. 2004, M.J. Wingfield, cultures CMW 31184 = CBS 125249; Las Golondrinas, from soil, Dec. 2004, M.J. Wingfield, culture CMW 31185 = CBS 125250.

Culture characteristics: Colonies fast growing with optimal growth temperature at 25 °C (growth at 15–35 °C) on MEA, reverse umber (15m) to verona-brown (13''k) after 7 d; moderate white aerial mycelium with moderate sporulation; chlamydospores extensive throughout the medium forming microsclerotia.

Substrate: Soil.

Distribution: Ecuador.

Notes: Morphologically, *Ca. densa* is very similar to *Ca. spathiphylli* and *Ca. pseudospathiphylli*. However, macroconidia of *Ca. densa* (av. $54 \times 6 \mu\text{m}$) are smaller than those of *Ca. spathiphylli* (av. $70 \times 6 \mu\text{m}$), but slightly larger and broader than those of *Ca. pseudospathiphylli* (av. $52 \times 4 \mu\text{m}$). *Calonectria densa* also readily forms lateral stipe extensions, not reported for the other two species.

Calonectria eucalypti L. Lombard, M.J. Wingf. & Crous, **sp. nov.** MycoBank MB515530, Fig. 8.

Etymology: Name refers to *Eucalyptus* from which the fungus was isolated.

Teleomorpha *Ca. colhounii* similis sed ascocarpo flavo vel aurantiaco differt. Anamorpha *Cy. colhounii* similis sed macroconidiis cylindricis utrinque rotundatis rectis (66 –) 69 – 75 (– 80) \times 5 – $6 \mu\text{m}$ mediocriter $72 \times 6 \mu\text{m}$, ter septatis, sine cicatrice abscissionis manifesta, cum muco hyalino in fasciculis parallelis cylindricis, differt.

Perithecia solitary or in groups, yellow to orange, becoming brown with age; in section apex and body yellow to orange, base red-brown, sub-globose to ovoid, 325 – $510 \mu\text{m}$ high, 285 – $360 \mu\text{m}$ diam, body turning dark red, and base dark red-brown (KOH+). Perithecial walls rough consisting of 2 thick-walled layers: outside layer of *textura globulosa*, 45 – $90 \mu\text{m}$ wide; becoming more compressed towards inner layer of *textura angularis*, 12 – $18 \mu\text{m}$ wide; becoming thin-walled and hyaline towards the centre, outer cells 24 – 50×10 – $40 \mu\text{m}$; inner cells 6 – 19×3 – $6 \mu\text{m}$: perithecial base up to $125 \mu\text{m}$ wide; consisting of dark red, angular cells; merging with an erumpent stroma, cells of the outer wall layer continuing into the pseudoparenchymatous cells of the erumpent stroma. *Asci* 4-spored, clavate, 92 – 188×10 – $27 \mu\text{m}$, tapering to a long thin stalk. *Ascospores* aggregated in the upper third of the ascus, hyaline, guttulate, fusoid with rounded ends, straight to slightly curved, (1–)3-septate, not or slightly constricted at the septum, (25 –) 30 – 36 (– 56) \times (3 –) 5 – 6 (– 8) μm (av. = $33 \times 6 \mu\text{m}$). Cultures were homothallic. *Conidiophores* with a stipe bearing penicillate suites of fertile branches, a stipe extension, and a terminal vesicle; Stipe septate, hyaline, smooth, 45 – 91×7 – $10 \mu\text{m}$; stipe extensions septate, straight to flexuous, 110 – $235 \mu\text{m}$ long, 5 – $6 \mu\text{m}$ wide at the apical septum, terminating in a broadly clavate vesicle, 4 – $6 \mu\text{m}$ diam. *Conidiogenous apparatus* 52 – $82 \mu\text{m}$ long, and 40 – $95 \mu\text{m}$ wide; primary branches aseptate or 1-septate, 21 – 29×5 – $6 \mu\text{m}$; secondary branches aseptate, 14 – 21×3 – $5 \mu\text{m}$; tertiary branches and additional branches (–5), aseptate, 11 – 16×3 – $5 \mu\text{m}$, each terminal branch producing 2–6 phialides;

phialides doliiform to reniform, hyaline, aseptate, $10\text{--}14 \times 3\text{--}4 \mu\text{m}$; apex with minute periclinal thickening and inconspicuous collarete. *Macroconidia* cylindrical, rounded at both ends, straight, $(66\text{--})69\text{--}75\text{--}(80) \times 5\text{--}6 \mu\text{m}$ (av. = $72 \times 6 \mu\text{m}$), 3-septate, lacking a visible abscission scar, held in parallel cylindrical clusters by colourless slime. *Mega-* and *microconidia* not seen.

Specimens examined: **Indonesia**, Sumatra Utara, Aek Nauli, on leaf of *Eucalyptus grandis*, May 2005, M.J. Wingfield, Herb. PREM 60298, **holotype** of *Ca. eucalypti*, culture ex-type CMW 18444 = CBS 125275; Aek Nauli, on leaf of *E. grandis*, May 2005, M.J. Wingfield, PREM 60299, culture CMW 14890 = CBS 125273; Aek Nauli, on leaf of *E. grandis*, May 2005, M.J. Wingfield, culture CMW 18443 = CBS 125274, Aek Nauli, on leaf of *E. grandis*, May 2005, M.J. Wingfield, culture CMW 18445 = CBS 125276.

Culture characteristics: Colonies fast growing with optimal growth temperature at $25 \text{ }^\circ\text{C}$ (growth at $15\text{--}30 \text{ }^\circ\text{C}$) on MEA, reverse colour tawny brown (13'i) after 7 d; abundant white aerial mycelium and sporulation; chlamydospores abundant throughout the medium, forming microsclerotia.

Substrate: *Eucalyptus grandis*.

Distribution: Indonesia.

Notes: The perithecia of *Ca. eucalypti* can be distinguished from *Ca. colhounii* and *Ca. macroconidialis* based on their yellow to orange colour in KOH. Macroconidia of *Ca. eucalypti* (av. $72 \times 6 \mu\text{m}$) are also larger than those of *Ca. colhounii* (av. $55 \times 6 \mu\text{m}$) and *Ca. madagascariensis* (av. $55 \times 4.5 \mu\text{m}$), but smaller than those of *Ca. macroconidialis* (av. $90 \times 6.5 \mu\text{m}$). Mating tests (Fig. 5) also showed that *Ca. eucalypti* is homothallic, a characteristic shared by *Ca. colhounii* and *Ca. madagascariensis* but not with *Ca. macroconidialis*, which is heterothallic (Crous 2002).

Calonectria humicola L. Lombard, M.J. Wingf. & Crous, **sp. nov.** MycoBank MB515531, Fig. 9.

Etymology: Name refers to the fact that this fungus was isolated from soil.

Teleomorpha ignota. Anamorpha *Cy. spathiphylli* similis sed macroconidiis cylindricis utrinque rotundatis rectis (45–)48–54(–56) × 4–5 μm mediocriter 51 × 5 μm, semel septatis, sine cicatrice abscissionis manifesta, cum muco hyalino in fasciculis parallelis cylindricis differt

Teleomorph unknown. *Conidiophores* with a stipe bearing penicillate suites of fertile branches, stipe extensions, and terminal vesicles; stipe septate, hyaline, smooth, 44–90 × 6–8 μm; stipe extensions septate, straight to flexuous, 126–157 μm long, 4–5 μm wide at the apical septum, terminating in a globose to ovoid to sphaeropedunculate vesicle, 10–12 μm diam. *Conidiogenous apparatus* 43–71 μm long, and 42–49 μm wide; primary branches aseptate, 20–29 × 4–6 μm; secondary branches aseptate, 12–19 × 3–5 μm; tertiary branches aseptate, 9–16 × 3–5 μm, each terminal branch producing 2–4 phialides; phialides doliiform to reniform, hyaline, aseptate, 10–15 × 3–4 μm; apex with minute periclinal thickening and inconspicuous collarete. *Macroconidia* cylindrical, rounded at both ends, straight, (45–)48–54(–56) × 4–5 μm (av. = 51 × 5 μm), 1-septate, lacking a visible abscission scar, held in parallel cylindrical clusters by colourless slime. *Mega-* and *microconidia* not seen.

Specimens examined: **Ecuador**, Pichincha Province, Las Golondrinas, from soil, Dec. 2004, M.J. Wingfield, Herb. PREM 60369, **holotype** of *Ca. humicola*, culture ex-type CMW 31183 = CBS 125251; Las Golondrinas, from soil, Jan. 2006, L. Lombard, culture CMW 31186 = CBS 125252; Las Golondrinas, from soil, Jan. 2006, L. Lombard, (Herb. PREM 60368) culture CMW 31187 = CBS 125269.

Culture characteristics: Colonies fast growing with optimal growth temperature at 25 °C (growth at 15–35 °C) on MEA, reverse umber (15m) to verona-brown (13''k) after 7 d; moderate white aerial mycelium with moderate sporulation; chlamydospores extensive throughout the medium, forming microsclerotia.

Substrate: Soil.

Distribution: Ecuador.

Notes: *Calonectria humicola* is morphologically very similar to *Ca. densa*, *Ca. pseudospathiphylli* and *Ca. spathiphylli*. However, no lateral stipe extensions occur in this species, whereas these are common in *Ca. densa*. Macroconidia of *Ca. humicola* (av. 51 × 5 μm) are slightly smaller than those of *Ca. densa* (av. 54 × 6 μm) and *Ca. spathiphylli* (av. 70 × 6 μm), but slightly broader than those of *Ca. pseudospathiphylli* (av. 52 × 4 μm).

Calonectria orientalis L. Lombard, M.J. Wingf. & Crous, **sp. nov.** MycoBank MB515532, Fig. 10.

Etymology: Name refers to the East Asian region, where the fungus was isolated.

Teleomorpha ignota. Anamorpha Cy. candelabro similis sed macroconidiis cylindricis utrinque rotundatis rectis (43–)46–50(–53) × 4–5 µm mediocriter 48 × 4 µm, semel septatis, sine cicatrice abscissionis manifesta, cum muco hyalino in fasciculis parallelis cylindricis differt.

Teleomorph unknown. *Conidiophores* with a stipe bearing penicillate suites of fertile branches, stipe extensions, and terminal vesicles; stipe septate, hyaline, smooth, 60–169 × 6–12 µm; stipe extensions septate, straight to flexuous, 90–218 µm long, 5–10 µm wide at the apical septum, terminating in a fusiform to pyriform or broadly clavate vesicle, 8–12 µm diam. *Conidiogenous apparatus* 54–174 µm long, and 67–92 µm wide; primary branches aseptate, 19–30 × 4–7 µm; secondary branches aseptate, 16–29 × 4–6 µm; tertiary and additional branches (–5) aseptate, 10–20 × 5–5 µm, each terminal branch producing 2–6 phialides; phialides doliiform to reniform, hyaline, aseptate, 10–19 × 2–5 µm; apex with minute periclinal thickening and inconspicuous collarette. *Macroconidia* cylindrical, rounded at both ends, straight, (43–)46–50(–53) × 4–5 µm (av. = 48 × 4 µm), 1-septate, lacking a visible abscission scar, held in parallel cylindrical clusters by colourless slime. *Mega-* and *microconidia* not seen.

Specimen examined: **Indonesia**, Langam, from soil, June 2005, M.J. Wingfield, Herb. PREM 60303, **holotype** of *Ca. orientalis*, culture ex-type CMW 20291 = CBS 125260; Teso East, from soil, June 2005, M.J. Wingfield culture CMW 20273 = CBS 125259; Teso East, from soil, June 2005, M.J. Wingfield culture CMW 20272 = CBS 125258.

Culture characteristics: Colonies fast growing with optimal growth temperature at 25 °C (growth at 15–35 °C) on MEA, reverse sepia-brown (13i) after 7 d; abundant white aerial mycelium with moderate to extensive sporulation; chlamydospores extensive throughout the medium, forming microsclerotia.

Substrate: Soil.

Distribution: Indonesia.

Notes: *Calonectria orientalis* is closely related to *Calonectria* spp. in the *Ca. brassicae* complex, based on phylogenetic inference and SNP analyses. However, morphological comparisons showed that it is most similar to species in the *Ca. scoparia* and *Ca. morganii* complexes with stipe extensions terminating in fusiform to pyriform or broadly clavate vesicles. As with *Ca. pini*, perithecia could not be induced when this species was mated with *Ca. brachiatica* and *Ca. brassicae*, highlighting the rarity of teleomorph structures for this group of fungi.

Calonectria pini L. Lombard, M.J. Wingf. & Crous, **sp. nov.** MycoBank MB515533, Fig. 11.

Etymology: Name refers to *Pinus*, the host from which the fungus was isolated.

Teleomorpha ignota. Anamorpha *Ca. brachiaticae* similis sed ramis conidiophorae tres vel minus sine extensionibus lateralibus stipae, macroconidiis cylindricis utrinque rotundatis rectis (37–)40–48(–50) × 4–6 μm mediocriter 44 × 5 μm, semel septatis, sine cicatrice abscissionis manifesta, cum muco hyalino in fasciculis parallelis cylindricis differt

Teleomorph unknown. *Conidiophores* with a stipe bearing penicillate suites of fertile branches, stipe extensions, and terminal vesicles; stipe septate, hyaline, smooth, 40–99 × 6–7 μm; stipe extensions septate, straight to flexuous, 121–266 μm long, 5–7 μm wide at the apical septum, terminating in a clavate vesicle, 4–6 μm diam. *Conidiogenous apparatus* 49–81 μm long, and 35–84 μm wide; primary branches aseptate, 20–30 × 4–6 μm; secondary branches aseptate, 13–22 × 3–5 μm; tertiary branches aseptate, 11–15 × 3–4 μm, each terminal branch producing 2–6 phialides; phialides doliiform to reniform, hyaline, aseptate, 10–15 × 3–4 μm; apex with minute periclinal thickening and inconspicuous collarete. *Macroconidia* cylindrical, rounded at both ends, straight, (37–)40–48(–50) × 4–6 μm (av. = 44 × 5 μm), 1-septate, lacking a visible abscission scar, held in parallel cylindrical clusters by colourless slime. *Mega-* and *microconidia* not seen.

Specimen examined: **Colombia**, Valle del Cauca, Buga, from *Pinus patula*, Sep. 2007, C.A. Rodas, Herb. PREM 60304, **holotype** of *Ca. pini*, culture ex-type CMW 31209 = CBS 123698. Buga, from *P. patula*, Sep. 2007, C.A. Rodas; Buga, from *P. patula*, Sep. 2007, C.A. Rodas, culture CMW 31210 = CBS 125523.

Culture characteristics: Colonies fast growing with optimal growth temperature at 25 °C (growth at 15–30 °C) on MEA, reverse amber (13k) to sepia-brown (13i) after 7 d; abundant white aerial mycelium with moderate to extensive sporulation; chlamydospores extensive throughout the medium forming microsclerotia.

Substrate: *Pinus patula*.

Distribution: Colombia.

Notes: *Calonectria pini* is very similar to *Ca. brachiatica*, but can be distinguished morphologically by the fact that it has three or fewer conidiophore branches and no lateral stipe extensions (Lombard *et al.* 2009b). Macroconidia of *Ca. pini* (av. $44 \times 5 \mu\text{m}$) are shorter than those of *Ca. brassicae* (av. $53 \times 4.5 \mu\text{m}$) and *Ca. gracilis* ($56 \times 4.5 \mu\text{m}$). This species also has fewer conidiophore branches than those mentioned above. *Calonectria pini* failed to produce perithecia when crossed with *Ca. brachiatica* and *Ca. brassicae*. This supports the findings of Crous *et al.* (2004b) and Lombard *et al.* (2009b), that teleomorph structures are rarely observed in members of the *Ca. brassicae* complex.

Calonectria pseudoscoparia L. Lombard, M.J. Wingf. & Crous, **sp. nov.** MycoBank MB515534, Fig. 12.

Etymology: Name reflects the fact that the species resembles the anamorph state of *Ca. scoparia*.

Teleomorpha ignota. Anamorpha *Ca. scopario* similis sed phialidibus elongato-doliiformibus vel reniformibus hyalinis non septatis $7\text{--}11 \times 2\text{--}4 \mu\text{m}$ apice minute periclinale incrassatis colliculo inconspicuo, macroconidiis cylindricis utrinque rotundatis rectis $(41\text{--})45\text{--}51\text{--}(52) \times 3\text{--}5 \mu\text{m}$ mediocriter $48 \times 4 \mu\text{m}$, semel septatis, sine cicatrice abscissionis manifesta, cum muco hyalino in fasciculis parallelis cylindricis differt

Teleomorph unknown. *Conidiophores* with a stipe bearing penicillate suites of fertile branches, stipe extensions, and terminal vesicles; stipe septate, hyaline, smooth, $56\text{--}107 \times 6\text{--}10 \mu\text{m}$; stipe extensions septate, straight to flexuous, $124\text{--}201 \mu\text{m}$ long, $4\text{--}6 \mu\text{m}$ wide at the apical septum, terminating in a obpyriform to ellipsoidal vesicle, $6\text{--}10 \mu\text{m}$ diam. *Conidiogenous apparatus* $34\text{--}87 \mu\text{m}$ long, and $52\text{--}74 \mu\text{m}$ wide; primary branches aseptate, $26\text{--}38 \times 4\text{--}7 \mu\text{m}$; secondary branches aseptate, $17\text{--}28 \times 4\text{--}6 \mu\text{m}$; tertiary branches and additional branches (–4) aseptate, $14\text{--}19 \times 3\text{--}4 \mu\text{m}$, each terminal branch producing 2–6 phialides; phialides elongate-doliiform to reniform, hyaline, aseptate, $7\text{--}11 \times 2\text{--}4 \mu\text{m}$; apex

with minute periclinal thickening and inconspicuous collarete. *Macroconidia* cylindrical, rounded at both ends, straight, $(41\text{--})45\text{--}51(\text{--}52) \times 3\text{--}5 \mu\text{m}$ (av. = $48 \times 4 \mu\text{m}$), 1-septate, lacking a visible abscission scar, held in parallel cylindrical clusters by colourless slime. *Mega-* and *microconidia* not seen.

Specimens examined: **Ecuador**, Pichincha Province, Las Golondrinas, Buenos Aires Nursery, from *Eucalyptus grandis* cutting, Dec. 2004, M.J. Wingfield, Herb. PREM 60305, **holotype** of *Ca. pseudoscoparia*, culture ex-type CMW 15218 = CBS 125257; Buenos Aires Nursery, from *E. grandis* cutting, Dec. 2004, M.J. Wingfield, Herb. PREM 60306, cultures from different cuttings, CMW 15214 = CBS 125254, CMW 15215 = CBS 125255, CMW 15216 = CBS 125256.

Culture characteristics: Colonies fast growing with optimal growth temperature at 25 °C (growth at 10–30 °C) on MEA, reverse amber (13k) to sepia-brown (13i) after 7 d; colony margins irregular with sparse to moderate white aerial mycelium with moderate sporulation; chlamydospores extensive throughout the medium forming microsclerotia.

Substrate: *Eucalyptus grandis*.

Distribution: Ecuador.

Notes: *Calonectria pseudoscoparia* (conidia av. $48 \times 4 \mu\text{m}$) can be distinguished from *Ca. scoparia* (conidia av. $60 \times 4.5 \mu\text{m}$) based on smaller macroconidia and the fact that it has elongated-doliiform to reniform phialides unlike those of *Ca. pauciramosa* and *Ca. scoparia*. Mating tests between this fungus and *Ca. scoparia* and *Ca. pauciramosa* failed to produce perithecia. Control crosses with both *Ca. pauciramosa* (CMW 5683 and CMW 30823) and *Ca. scoparia* tester isolates (CMW 31000 and CMW 31001) produced perithecia with viable ascospores (Fig. 6) showing that culture conditions were appropriate for mating.

Calonectria sulawesiensis L. Lombard, M.J. Wingf. & Crous, **sp. nov.** MycoBank MB515535, Fig. 13.

Etymology: Name refers to the Indonesian island of Sulawesi, where the fungus was collected.

Teleomorpha ignota. Anamorpha *Ca. morganii* similis sed vesiculo terminali late clavato vel ellipsoideo $5\text{--}7 \mu\text{m}$ diametro, macroconidiis cylindricis utrinque rotundatis rectis $(41\text{--})45\text{--}51(\text{--}54) \times (3\text{--})4\text{--}6 \mu\text{m}$ mediocriter 48×4

μm , semel septatis, sine cicatrice abscissionis manifesta, cum muco hyalino in fasciculis parallelis cylindricis differt.

Teleomorph unknown. *Conidiophores* with a stipe bearing penicillate suites of fertile branches, stipe extensions, and terminal vesicles; stipe septate, hyaline, smooth, $37\text{--}139 \times 5\text{--}11 \mu\text{m}$; stipe extensions septate, straight to flexuous, $113\text{--}262 \mu\text{m}$ long, $5\text{--}7 \mu\text{m}$ wide at the apical septum, terminating in a broadly clavate to ellipsoidal vesicle, $5\text{--}7 \mu\text{m}$ diam. *Conidiogenous apparatus* $41\text{--}79 \mu\text{m}$ long, and $43\text{--}81 \mu\text{m}$ wide; primary branches aseptate, $17\text{--}41 \times 3\text{--}6 \mu\text{m}$; secondary branches aseptate, $10\text{--}27 \times 3\text{--}6 \mu\text{m}$; tertiary branches and additional branches (–5), aseptate, $9\text{--}15 \times 3\text{--}5 \mu\text{m}$, each terminal branch producing 2–6 phialides; phialides doliiform to reniform, hyaline, aseptate, $9\text{--}15 \times 2\text{--}5 \mu\text{m}$; apex with minute periclinal thickening and inconspicuous collarete. *Macroconidia* cylindrical, rounded at both ends, straight, $(41\text{--})45\text{--}51(\text{--}54) \times (3\text{--})4\text{--}6 \mu\text{m}$ (av. = $48 \times 4 \mu\text{m}$), 1-septate, lacking a visible abscission scar, held in parallel cylindrical clusters by colourless slime. *Mega-* and *microconidia* not seen.

Specimens examined: **Indonesia**, Sulawesi, from leaf of *Eucalyptus* sp., July 2003, M.J. Wingfield, Herb. PREM 60300, **holotype** of *Ca. sulawesiensis*, culture ex-type CMW 14878 = CBS 125277; Sulawesi, from leaf of *Eucalyptus* sp., July 2003, M.J. Wingfield, PREM 60301 culture CMW 14883; from different leaves, culture CMW 14859 = CBS 125248, CMW 14879 = CBS 125253.

Culture characteristics: Colonies fast growing with optimal growth temperature at $25 \text{ }^\circ\text{C}$ (growth at $15\text{--}30 \text{ }^\circ\text{C}$) on MEA, reverse amber (13k) to sepia-brown (13i) after 7 d; abundant white aerial mycelium with moderate to extensive sporulation; chlamydospores extensive throughout the medium, forming microsclerotia.

Substrate: *Eucalyptus* sp.

Distribution: Indonesia.

Notes: There are a few morphological differences distinguishing *Ca. sulawesiensis* from other species in the *Ca. morganii* complex. Macroconidia of *Ca. sulawesiensis* (av. $48 \times 4 \mu\text{m}$) are slightly larger than those of *Ca. brasiliensis* (av. $30 \times 4 \mu\text{m}$), *Ca. cerciana* (av. $44 \times 5 \mu\text{m}$), *Ca. insularis* (av. $45 \times 4 \mu\text{m}$) and *Ca. morganii* (av. $45 \times 4 \mu\text{m}$), but smaller than those of *Ca. hawksworthii* (av. $56 \times 4 \mu\text{m}$), *Ca. leucothoës* (av. $73 \times 5 \mu\text{m}$) and *Ca. variabilis* (av. 73×5

µm). Mating tests where *Ca. sulawesiensis* was crossed with *Ca. brasiliensis*, *Ca. cerciana* and *Ca. insularis* failed to produce perithecia, or produced perithecia without viable ascospores.

Calonectria angustata (Crous & El-Gholl) L. Lombard, M.J. Wingf. & Crous, **comb. nov.** MycoBank MB515536.

Basionym: *Cylindrocladium angustatum* Crous & El-Gholl, Mycoscience 41: 522. 2000.

Calonectria australiensis (Crous & K.D. Hyde) L. Lombard, M.J. Wingf. & Crous, **comb. nov.** MycoBank MB515537.

Basionym: *Cylindrocladium australiense* Crous & K.D. Hyde, Stud. Mycol. 55: 221. 2006.

Calonectria canadensis (J.C. Kang, Crous & C.L. Schoch) L. Lombard, M.J. Wingf. & Crous, **comb. nov.** MycoBank MB515538.

Basionym: *Cylindrocladium canadense* J.C. Kang, Crous & C.L. Schoch, Syst. Appl. Microbiol. 24: 210. 2001.

Calonectria chinensis (Crous) L. Lombard, M.J. Wingf. & Crous, **comb. nov.** MycoBank MB515539.

Basionym: *Cylindrocladium chinense* Crous, Stud. Mycol. 50: 420. 2004.

Calonectria citri (Boedjin & Reitsma) L. Lombard, M.J. Wingf. & Crous, **comb. nov.** MycoBank MB515540.

Basionym: *Cylindrocladium citri* (H.S. Fawc. & Klotz) Boedjin & Reitsma, Reinwardtia 1: 57. 1950.

≡ *Candelospora citri* H.S. Fawc. & Klotz, Mycologia 29: 213. 1937.

Calonectria curvata (Boedjin & Reitsma) L. Lombard, M.J. Wingf. & Crous, **comb. nov.** MycoBank MB515541

Basionym: *Cylindrocladium curvatum* Boedjin & Reitsma, Reinwardtia 1: 54. 1950.

Calonectria curvispora (Crous & D. Victor) L. Lombard, M.J. Wingf. & Crous, **comb. nov.** MycoBank MB515542

Basionym: *Cylindrocladium curvisporum* Crous & D. Victor, Syst. Appl. Microbiol. 20: 283. 1997.

Calonectria ecuadoriae (Crous & M.J. Wingf.) L. Lombard, M.J. Wingf. & Crous, **comb. nov.** MycoBank MB515543.

Basionym: *Cylindrocladium ecuadoriae* Crous & M.J. Wingf., Stud. Mycol. 55:222. 2006.

Calonectria gordoniae (Leahy, T.S. Schub. & El-Gholl) L. Lombard, M.J. Wingf. & Crous, **comb. nov.** MycoBank MB515544.

Basionym: *Cylindrocladium gordoniae* Leahy, T.S. Schub. & El-Gholl, Mycotaxon 76: 80. 2000.

Calonectria hawksworthii (Peerally) L. Lombard, M.J. Wingf. & Crous, **comb. nov.** MycoBank MB515545.

Basionym: *Cylindrocladium hawksworthii* Peerally, Mycotaxon 40: 375. 1991.

Calonectria hurae (Linder & Whetzel) L. Lombard, M.J. Wingf. & Crous, **comb. nov.** MycoBank MB515546.

Basionym: *Cercospora hurae* Linder & Whetzel, Mycologia 29: 656. 1937

≡ *Cylindrocladiopsis hurae* (Linder & Whetzel) U. Braun, Mycotaxon 51: 40. 1994

≡ *Cylindrocladium hurae* (Linder & Whetzel) Crous, Taxonomy and pathology of *Cylindrocladium* (*Calonectria*) and allied genera: 185. 2002.

= *Cylindrocladium heptaseptatum* Sober, Alfieri & Knauss, Phytopathology 65: 333. 1975.

= *Cylindrocladiopsis lagerstroemiae* J.M. Yen, Mycotaxon 8: 236. 1979.

Calonectria indonesiae (Crous) L. Lombard, M.J. Wingf. & Crous, **comb. nov.** MycoBank MB515547.

Basionym: *Cylindrocladium indonesiae* Crous, Stud. Mycol. 50: 424. 2004.

Calonectria leucothoës (El-Gholl, Leahy & T.S. Schub.) L. Lombard, M.J. Wingf. & Crous, **comb. nov.** MycoBank MB515548.

Basionym: *Cylindrocladium leucothoës* El-Gholl, Leahy & T.S. Schub., Canad. J. Bot. 67: 2530. 1989.

= *Cylindrocladium perseae* T.S. Schub., Leahy & El-Gholl, Mycotaxon 73: 474. 1999.

Calonectria malesiana (Crous) L. Lombard, M.J. Wingf. & Crous, **comb. nov.** MycoBank MB515549.

Basionym: *Cylindrocladium malesianum* Crous, Stud. Mycol. 50: 425. 2004

Calonectria multiphialidica (Crous, P. Simoneau & J.-M. Risède) L. Lombard, M.J. Wingf. & Crous, **comb. nov.** MycoBank MB515550.

Basionym: *Cylindrocladium multiphialidicum* Crous, P. Simoneau & J.-M. Risède, Stud. Mycol. 50: 425. 2004.

Calonectria pacifica (J.C. Kang, Crous & C.L. Schoch) L. Lombard, M.J. Wingf. & Crous, **comb. nov.** MycoBank MB515551.

Basionym: *Cylindrocladium pacificum* J.C. Kang, Crous & C.L. Schoch, Syst. Appl. Microbiol. 24: 213. 2001.

Calonectria penicilloides (Tubaki) L. Lombard, M.J. Wingf. & Crous, **comb. nov.** MycoBank MB515552

Basionym: *Candelospora penicilloides* Tubaki, Nogaoa 2: 58. 1952.

≡ *Cylindrocladium penicilloides* (Tubaki) Tubaki, J. Hattori Bot. Lab. 20: 154. 1958.

Calonectria pseudonaviculata (Crous, J.Z. Groenew. & C.F. Hill) L. Lombard, M.J. Wingf. & Crous, **comb. nov.** MycoBank MB515554

Basionym: *Cylindrocladium pseudonaviculatum* Crous, J.Z. Groenew. & C.F. Hill, Sydowia 54: 26. 2002.

= *Cylindrocladium buxicola* Henricot, Mycologia 94: 993. 2002.

Calonectria sumatrensis (Crous) L. Lombard, M.J. Wingf. & Crous, **comb. nov.** MycoBank MB515555.

Basionym: *Cylindrocladium sumatrense* Crous, Stud. Mycol. 50: 426. 2004.

DISCUSSION

In this study, a collection of isolates of unknown identity were shown to represent seven new species of *Calonectria*. These species, provided with the names *Ca. eucalypti*, *Ca. orientalis* and *Ca. sulawesiensis* from Indonesia, *Ca. densa*, *Ca. humicola* and *Ca. pseudoscoparia* from Ecuador and *Ca. pini* from Colombia were recognised based on morphological characteristics and phylogenetic inference. Recognition of a relatively large number of new species, mainly from soil samples collected in areas not previously intensively sampled, suggests that many more species of *Calonectria* remain to be discovered, particularly from the tropics and Southern Hemisphere.

Calonectria eucalypti, isolated from the leaves of *E. grandis*, adds a new species to *Ca. colhounii* complex (Crous 2002, Crous *et al.* 2006), which includes *Ca. colhounii*, *Ca. macroconidialis* and *Ca. madagascariensis*. Members of this complex are characterised by their unique yellow perithecia (Crous 2002). Although *Ca. eucalypti* was isolated from lesions typical of *Cylindrocladium* leaf blight, its importance as a pathogen is unknown. *Calonectria eucalypti* was shown to be homothallic, which is a characteristic that this species shares with *Ca. colhounii* and *Ca. madagascariensis*.

The descriptions of *Ca. pini* and *Ca. orientalis* add two species to the *Ca. brassicae* complex (Crous *et al.* 2006, Lombard *et al.* 2009b). *Calonectria pini* was isolated from *P. patula* rooted cuttings with symptoms similar to those associated with root and collar infections caused by *Ca. brassicae* and *Ca. brachiatica* on other *Pinus* spp. (Lombard *et al.* 2009b). In contrast, *Ca. orientalis* was isolated from soils collected in Indonesia and nothing is known regarding its pathogenicity. Phylogenetic inference and SNP analyses showed that these are closely related sibling species (Taylor *et al.* 2000) with genetic isolation having apparently occurred recently. Crosses between isolates of *Ca. pini* and *Ca. orientalis* as well as those with themselves and other *Calonectria* spp. in the group failed to produce perithecia. This is consistent with the observations of Crous *et al.* (2006) and Lombard *et al.* (2009b), that *Calonectria* spp. in this complex rarely produce teleomorph structures in culture.

Calonectria sulawesiensis resides in the *Ca. morganii* complex, closely related to *Ca. brasiliensis* and *Ca. insularis*. Morphologically, *Ca. sulawesiensis* can be distinguished from other species in the complex based only on macroconidia dimensions. Therefore phylogenetic inference based on DNA sequence data is necessary to distinguish it from other members of the *Ca. morganii* complex. Members of this complex are well-known pathogens of various hosts world-wide (Crous 2002), but nothing is known regarding the pathogenicity of *Ca. sulawesiensis*.

Calonectria pseudoscoparia is a new species in the *Ca. scoparia* complex (Schoch *et al.* 1999), isolated from *E. grandis* cuttings collected in Ecuador and displaying basal rot symptoms. *Calonectria* spp. in this group are well known causal agents of cutting rot in commercial forestry nurseries worldwide (Crous *et al.* 1991, Crous 2002, Lombard *et al.* 2009c). However, the pathogenicity of *Ca. pseudoscoparia* is only assumed based on the symptoms with which the fungus was associated.

The two newly described species, *Ca. densa* and *Ca. humicola*, isolated from Ecuadorian soils reside in the *Ca. spathiphylli* complex as defined by Kang *et al.* (2001b). *Calonectria pseudospathiphylli* and *Ca. spathiphylli*, that define this complex, are not easily distinguished based on morphology and DNA sequence comparisons are required for their identification. They can, however, be distinguished based on their mating strategies, with *Ca. pseudospathiphylli* being homothallic and *Ca. spathiphylli* being heterothallic (Kang *et al.* 2001b, Crous 2002). The mating strategies of *Ca. densa* and *Ca. humicola* could not be determined in this study. This complex of species appears to originate from Central and South America (Chase & Poole 1987, Kang *et al.* 2001b, Crous 2002).

DNA sequence data for the ITS, BT and HIS3 have been used more extensively to explore phylogenetic relationships amongst *Calonectria* spp. (Schoch *et al.* 1999, Kang *et al.* 2001a, 2001b, Henricot & Culham 2002, Crous *et al.* 2004b, 2006). In this regard, BT is the gene region that provides the most valuable insights into relationships between all species of *Calonectria* (Schoch *et al.* 2001b, Crous 2002, Henricot & Culham 2002). Application of the CAL and TEF-1 α partial gene sequences has only recently been introduced for *Calonectria* spp. (Crous *et al.* 2004b, 2006) but data for these gene regions have been available for only a small sub-set of species. The present study has attempted to address this problem and also introduce the ACT and LSU gene sequences that have not been employed previously for *Calonectria* spp. It has also provided sequence data for all seven gene regions for all accepted species in the genus.

The ITS and LSU sequences provided little valuable information to separate *Calonectria* spp. In contrast, sequence data for the protein-coding gene regions ACT, BT, CAL, HIS3 and TEF-1 α provided good resolution of *Calonectria* spp., confirming the results of previous studies (Schoch *et al.* 1999, 2001a, Crous 2002, Henricot & Culham 2002, Crous *et al.* 2004b, 2006). This study also introduced sequence data for the ACT gene region, although it had few informative sites, consistent with the results of previous studies on other fungi (Helgason *et al.* 2003, Hunter *et al.* 2006). Phylogenetic analyses of the individual coding gene regions and single nucleotide polymorphisms showed that CAL sequence data provide the best resolution distinguishing *Calonectria* spp. from each other followed by sequence data for the TEF-1 α , HIS3, BT and ACT gene regions.

In addition to identifying the most useful gene regions to accurately identify species of *Calonectria*, an important goal of this study was to re-consider the phylogenetic relationships between all the species in this genus. Having determined that the ACT, BT, CAL, HIS3 and TEF-1 α gene regions give the best resolution when identifying species of *Calonectria*, a phylogenetic tree for the genus was generated. This showed that the group includes two major clades and that these define morphologically similar groups of *Calonectria* spp. These two major clades have substantial sub-structure with all of the 66 species of *Calonectria* residing in one of 13 sub-clades. Eleven of these sub-clades, that include 50 species, represent the Prolate Group of isolates and two sub-clades that include 16 species representing the Sphaero-Naviculate Group of isolates.

The Prolate group of isolates incorporates the majority of the plant pathogenic *Calonectria* spp. and includes the type species for *Calonectria* (*Ca. pyrochoa*) and *Cylindrocladium* (*C. scoparium*), respectively. Most of these pathogenic species have been reported from forestry crops (Peerally 1991, Crous & Wingfield 1994, Crous 2002, Crous *et al.* 2006) but a few have also been found to infect horticultural and agronomic crops (Boedjin & Reitsma 1950, Kim *et al.* 1998, Crous 2002, Polizzi *et al.* 2007, Vitale *et al.* 2008). None of the sub-clades in this group could, however, be correlated with any specific host type.

The geographic distribution of the *Calonectria* spp. representing the various sub-clades of the unifying Prolate Group of isolates shows some correlation in their distribution. *Calonectria* spp. in the sub-clade representing the *Ca. reteaudii* complex (Sub-clade I) have been reported only from Australia, China, Indonesia and New Zealand (Crous 2002, Gadgill & Dick 2004, Crous *et al.* 2006, Lombard *et al.* 2009c). Another sub-clade of isolates that appears to have geographical structure resides in the *Ca. brassicae* complex (Sub-clade IV). Species in this sub-clade, with the exception of *Ca. orientalis*, have all been reported from South and Central America (Crous 2002, Crous *et al.* 2004b, Lombard *et al.* 2009b). Isolates in other sub-clades appeared to have broad geographic distribution and not to occur in any defined part of the world.

Species residing in the Sphaero-Naviculate Group had no obvious patterns of pathogenicity, or distribution. This group consisted of two sub-clades in which only vesicle morphology was a consistent character. The majority of the species in the *Ca. kyotensis* complex (Sub-clade XII) have been isolated from debris and soil (Crous *et al.* 2004b) but a few such as *Ca.*

kyotensis, *Ca. ilicicola* and *Ca. pacifica* are important pathogens of agronomic and forestry crops (Crous 2002, Crous *et al.* 2004b). Members of this sub-clade also had a broad distribution with the majority reported from Asia (Crous *et al.* 2004b) and they included both heterothallic and homothallic species (Crous 2002, Crous *et al.* 2004b).

The second sub-clade in the Sphaero-Naviculate Group of isolates (Sub-clade XIII) included only three *Calonectria* spp., only two of which have morphological similarities. *Calonectria multiphialidica* is morphologically similar to the *Calonectria* spp. in Sub-clade XII but there were no obvious patterns of distribution and pathogenicity for this group.

The intention of this phylogenetic study was to include all *Calonectria* spp. recognised to date. *Calonectria curvata* and *Ca. hederiae* were, however, not included because there are no cultures for them as has previously been mentioned by Crous (2002). Furthermore, *Ca. rajasthanensis*, *Cy. avesiculatum* var. *microsporum*, *Cy. bambusae*, *Cy. couratarii*, *Cy. crataegi*, *Cy. intermedium* and *Cy. musae* were not included due either to the fact that they have not been validly described or not recognised as true species of *Calonectria* (Crous 2002). Based on the results of this study, 68 *Calonectria* spp. are recognised as valid and cultures are available for 66 of them.

The teleomorph state has not been seen for several species of *Calonectria*. Nonetheless *Cylindrocladium* spp., irrespective of whether their perithecial states are known or not, have been provided names in *Calonectria*. This is consistent with the view that all newly described pleomorphic fungal species, the teleomorph name or the oldest typified name takes precedence over the anamorph or more recent name when both types belong to the same holomorph taxon (Hawksworth 2005, McNiell *et al.* 2005). It has already been established that *Calonectria* spp. have only *Cylindrocladium* anamorphs (Rossman *et al.* 1999, Schoch *et al.* 2001b), with micro- and megaconidial states that have thus far not been named. The name *Calonectria* was typified in 1867 (Rossman 1979) whereas that of *Cylindrocladium* was typified in 1892 (Morgan 1892). Therefore *Calonectria* takes preference above *Cylindrocladium* and should henceforth be used for all species irrespective of whether the perithecial state has been found.

KEYS

Both synoptic and dichotomous keys to species of *Calonectria* are presented. In the synoptic key, numbers grouped with each character refer to the species that are alphabetically arranged below:

1. *Ca. acicola* P.D. Gadgill & M.A. Dick
2. *Ca. angustata* (Crous & El-Gholl) L. Lombard, M.J. Wingf. & Crous
3. *Ca. asiatica* Crous & N.L.Hywel-Jones
4. *Ca. australiensis* (Crous & K.D. Hyde) L. Lombard, M.J. Wingf. & Crous
5. *Ca. avesiculata* T.S. Schub., El-Gholl, Alfieri & Schoult.
6. *Ca. brachiatica* L. Lombard, M.J. Wingf. & Crous
7. *Ca. brassicae* (Panwar & Borha) L. Lombard, M.J. Wingf. & Crous
8. *Ca. brasiliensis* (Peerally) L. Lombard, M.J. Wingf. & Crous
9. *Ca. canadensis* (J.C. Kang, Crous & C.L. Schoch) L. Lombard, M.J. Wingf. & Crous
10. *Ca. cerciana* L. Lombard, M.J. Wingf. & Crous
11. *Ca. chinensis* (Crous) L. Lombard, M.J. Wingf. & Crous
12. *Ca. citri* (Boedjin & Reitsma) L. Lombard, M.J. Wingf. & Crous
13. *Ca. clavata* Alfieri, El-Gholl & E.L. Barnard
14. *Ca. colhounii* Peerally
15. *Ca. colombiana* L. Lombard, M.J. Wingf., Crous
16. *Ca. colombiensis* Crous
17. *Ca. curvata* (Boedjin & Reitsma) L. Lombard, M.J. Wingf. & Crous
18. *Ca. curvispora* (Crous & D. Victor) L. Lombard, M.J. Wingf. & Crous
19. *Ca. densa* L. Lombard, M.J. Wingf. & Crous
20. *Ca. ecuadoriae* (Crous & M.J. Wingf.) L. Lombard, M.J. Wingf. & Crous
21. *Ca. eucalypti* L. Lombard, M.J. Wingf. & Crous
22. *Ca. gracilipes* Crous & G.R.A. Mchau
23. *Ca. gracilis* Crous, M.J. Wingf. & Alfenas
24. *Ca. gordoniae* (Leahy, T.S. Schub. & El-Gholl) L. Lombard, M.J. Wingf. & Crous
25. *Ca. hawksworthii* (Peerally) L. Lombard, M.J. Wingf. & Crous
26. *Ca. hederæ* C. Booth & J.S. Murray
27. *Ca. hongkongensis* Crous
28. *Ca. humicola* L. Lombard, M.J. Wingf. & Crous

29. *Ca. hurae* (Linder & Whetzel) L. Lombard, M.J. Wingf. & Crous
30. *Ca. ilicicola* Boedjin & Reitsma
31. *Ca. indonesiae* (Crous) L. Lombard, M.J. Wingf. & Crous
32. *Ca. indusiata* (Seaver) Crous
33. *Ca. insularis* C.L. Schoch & Crous
34. *Ca. kyotensis* Tersh.
35. *Ca. leguminum* (Rehm) Crous
36. *Ca. leucothoës* (El-Gholl, Leahy & T.S. Schub.) L. Lombard, M.J. Wingf. & Crous
37. *Ca. macroconidialis* (Crous, M.J. Wingf. & Alfenas) Crous
38. *Ca. madagascariensis* Crous
39. *Ca. malesiana* (Crous) L. Lombard, M.J. Wingf. & Crous
40. *Ca. mexicana* C.L. Schoch & Crous
41. *Ca. morganii* Crous, Alfenas & M.J. Wingf.
42. *Ca. multiphialidica* (Crous, P. Simoneau & J.-M. Risède) L. Lombard, M.J. Wingf. & Crous
43. *Ca. multiseptata* Crous & M.J. Wingf.
44. *Ca. naviculata* Crous & M.J. Wingf.
45. *Ca. orientalis* L. Lombard, M.J. Wingf. & Crous
46. *Ca. ovata* D. Victor & Crous
47. *Ca. pacifica* (J.C. Kang, Crous & C.L. Schoch) L. Lombard, M.J. Wingf. & Crous
48. *Ca. pauciramosa* C.L. Schoch & Crous
49. *Ca. penicilliodes* (Tubaki) L. Lombard, M.J. Wingf. & Crous
50. *Ca. pini* L. Lombard, M.J. Wingf. & Crous
51. *Ca. polizzii* L. Lombard, M.J. Wingf. & Crous
52. *Ca. pseudonaviculata* (Crous, J.Z. Groenew. & C.F. Hill) L. Lombard, M.J. Wingf. & Crous
53. *Ca. pseudoreteaudii* L. Lombard, M.J. Wingf. & Crous
54. *Ca. pseudoscoparia* L. Lombard, M.J. Wingf. & Crous
55. *Ca. pseudospathiphylli* J.C. Kang, Crous & C.L. Schoch
56. *Ca. pteridis* Crous, M.J. Wingf. & Alfenas
57. *Ca. pyrochoa* (Desm.) Sacc.
58. *Ca. queenslandica* L. Lombard, M.J. Wingf. & Crous
59. *Ca. reteaudii* (Bugn.) C. Booth

60. *Ca. rumohrae* El-Gholl & Alfenas
 61. *Ca. scoparia* Peerally
 62. *Ca. spathiphylli* El-Gholl, J.Y. Uchida, Alfenas, T.S. Schub., Alfieri & A.R. Chase
 63. *Ca. spathulata* El-Gholl, Kimbr., E.L. Barnard, Alfieri & Schoult.
 64. *Ca. sulawesiensis* L. Lombard, M.J. Wingf. & Crous
 65. *Ca. sumatrensis* (Crous) L. Lombard, M.J. Wingf. & Crous
 66. *Ca. terrae-reginae* L. Lombard, M.J. Wingf. & Crous
 67. *Ca. variabilis* Crous, B.J.H. Janse, D. Victor, G.F. Marias & Alfenas
 68. *Ca. zuluensis* L. Lombard, M.J. Wingf. & Crous

Synoptic key to *Calonectria* species

1. Teleomorph:
 - a. Teleomorph state known
 1, 3, 5, 13, 14, 15, 16, 21, 22, 23, 26, 27, 29, 30, 31, 32, 33, 34, 35, 37, 38, 40, 41, 43, 44, 46, 48, 55, 56, 57, 59, 60, 61, 62, 63, 67, 68
 - b. Teleomorph state unknown
 2, 4, 6, 7, 8, 9, 10, 11, 12, 17, 18, 19, 20, 24, 25, 28, 36, 39, 42, 45, 47, 49, 50, 51, 52, 53, 54, 58, 64, 65, 66
2. Ascocarps:
 - a. Red-brown to red in colour, changing to dark-red in 3 % KOH
 1, 23, 44, 56, 61, 67
 - b. Orange to red in colour, changing to dark-red in 3 % KOH
 3, 5, 15, 16, 22, 26, 30, 32, 33, 34, 40, 43, 55, 62, 68
 - c. Orange to red-brown in colour, changing to dark-red in 3 % KOH
 13, 27, 35, 46, 48, 57, 59, 60, 63
 - d. Yellow to orange in colour, only base and stroma changing to dark-red in 3 % KOH
 14, 21, 37, 38, 41
3. Asci:
 - a. 8-spored and clavate
 1, 3, 5, 13, 15, 16, 22, 23, 26, 27, 30, 32, 33, 34, 35, 38, 40, 41, 43, 44, 46, 48, 55, 56, 57, 59, 60, 61, 62, 63, 67, 68

- b. 4-spored and clavate
14, 21, 37
- 4. Ascospore septation:
 - a. 1-septate
3, 15, 16, 22, 23, 27, 33, 34, 40, 41, 48, 61, 68
 - b. (1–)3-septate
5, 13, 14, 21, 26, 30, 32, 35, 37, 38, 44, 46, 55, 56, 57, 59, 62, 63, 67
 - c. (3–)4-septate
1
 - d. (1–)3–6(–9) septate
43, 60
- 5. Ascospore width (av. in μm)
 - a. 4–5
15, 16, 22, 34, 44, 62, 67, 68
 - b. 5.5–6
1, 3, 5, 13, 14, 21, 26, 27, 30, 33, 37, 38, 40, 41, 46, 55, 56, 57, 59, 61, 63
 - c. 6.5–7
22, 32, 35, 43, 48, 60
- 6. Ascospore length (av. in μm)
 - a. 30–39
3, 15, 16, 21, 22, 23, 27, 33, 34, 41, 48, 68
 - b. 40–49
5, 13, 30, 44, 55, 57, 61, 62, 67
 - c. 50–59
14, 26, 32, 37, 38, 40, 56, 63
 - d. 60–69
46
 - e. 70 and above
1, 35, 43, 59, 60

7. Stipe length (av. in μm)

a. 40–100

1, 5, 6, 9, 10, 16, 18, 20, 21, 27, 30, 31, 33, 34, 36, 38, 40, 44, 47, 48,
49, 50, 57, 58, 61, 63, 65, 66, 68

b. 101–150

4, 7, 11, 13, 15, 24, 32, 41, 42, 51, 53, 54, 60, 62, 64,

c. 151–200

2, 3, 12, 14, 19, 22, 23, 28, 29, 35, 39, 45, 46, 52, 56, 67

d. above 200

25, 26, 37, 55, 59

8. Stipe extension length (av. in μm)

a. Less than 100

1

b. 100–200

9, 11, 12, 15, 16, 18, 19, 25, 27, 28, 31, 34, 39, 41, 44, 51, 52, 57, 58,
68

c. 201–300

2, 3, 10, 13, 14, 21, 22, 24, 26, 30, 33, 35, 36, 40, 45, 46, 47, 48, 50,
54, 55, 56, 61, 62, 63, 64, 65, 66, 67

d. Above 300

4, 5, 6, 7, 20, 23, 29, 32, 37, 38, 42, 53, 59, 60

9. Vesicle shape

a. Avesiculate to clavate

5

b. Clavate

1, 2, 4, 6, 7, 13, 14, 20, 21, 22, 23, 24, 29, 32, 35, 37, 38, 43, 50, 53,
56, 58, 59, 60, 64, 66

c. Ellipsoidal to pyriform to obovoid

8, 12, 25, 26, 41, 45, 55, 61, 63

d. Ellipsoidal to ovoid

46

- e. Ellipsoidal to obpyriform
10, 15, 33, 36, 40, 48, 51, 54, 57, 68
- f. Sphaeropedunculate
3, 9, 11, 16, 17, 18, 27, 30, 31, 34, 39, 42, 47, (49), 64, 67
- g. Globose
19, 28, 62
- h. Naviculate
44, 52

10. Shape of phialides on macroconidiophore

- a. Reniform to doliiform
3, 6, 7, 8, 9, 10, 12, 15, 17, 19, 20, 21, 22, 23, 24, 25, 26, 28, 33, 34, 36, 40, 41, 44, 45, 46, 48, 49, 50, 51, 52, 54, 57, 61, 63, 64, 68
- b. Elongate reniform to doliiform
5, 11, 13, 14, 16, 18, 27, 30, 31, 39, 42, 47, 55, 56, 62, 65, 67
- c. Cylindrical to allantoids
1, 2, 4, 29, 32, 35, 37, 38, 53, 58, 59, 60, 66

11. Number of fertile branches on macroconidiophore

- a. 1–3
1, 5, 8, 9, 11, 12, 17, 18, 28, 30, 46, 48, 49, 50, 51, 52, 53, 57, 58, 60, 63, 66, 67, 68
- b. 4–6
2, 3, 4, 6, 7, 14, 16, 19, 21, 24, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 44, 45, 46, 47, 54, 55, 56, 59, 61, 62, 64, 65
- c. More than 6
20, 27, 42

12. Microconidia

- a. Microconidia absent
2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 25, 26, 27, 28, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 44, 45, 47, 48, 49, 50, 51, 52, 54, 55, 57, 58, 61, 63, 64, 65, 66, 68

b. Microconidia present

1, 13, 24, 29, 30, 43, 46, 53, 56, 59, 60, 62, 67

13. Microconidia septation

a. 1-septate

13, 29, 30, 46, 56, 62, 67

b. 1(-3)-septate

24, 59, 60

c. 1-3-septate

1, 43, 53

14. Microconidia width (mean in μm)

a. Up to 3

13, 29, 43, 46, 56, 59

b. Up to 4

24, 53, 62, 67

c. Up to 5

1, 30, 60

15. Microconidia length (mean in μm)

a. Below 20

29

b. 20-30

1, 30, 46, 56, 59, 60, 67

c. 31-40

13, 24, 62

d. above 40

43, 53

16. Macroconidial septation

a. 1-septate

3, 6, 7, 8, 9, 10, 11, 12, 15, 17, 19, 22, 25, 27, 28, 31, 33, 34, 39, 40,
41, 42, 44, 45, 47, 48, 50, 51, 52, 54, 61, 64, 65, 68

b. 1(-3)-septate

5, 13, 16, 18, 20, 23, 24, 36, 46, 53, 55, 56, 62

c. (1-)3-septate

4, 14, 21, 30, 32, 38, 49, 57,

d. (1-)3(-6)-septate

26, 37, 58, 66

e. (1-)5(-6)-septate

1, 26, 35, 59, 60

f. (1-)7(-8)-septate

29

g. More than 8-septate

2

17. Macroconidia width (av. in μm)

a. 3-4

8, 9, 11, 12, 15, 17, 25, 27, 31, 33, 34, 39, 40, 41, 44, 45, 51, 54, 55,
63, 64, 68

b. 4.5-5

3, 5, 6, 7, 10, 13, 14, 16, 18, 20, 22, 23, 24, 28, 35, 36, 38, 42, 46, 47,
48, 49, 50, 52, 61, 65, 67

c. 5.5-6

19, 21, 26, 30, 32, 56, 57, 58, 62, 66

d. 6.5-7

1, 4, 37, 59

e. above 7

2, 29, 53, 60

18. Macroconidia length (av. in μm)

a. Less than 40

8, 15, 51, 68

b. 40-46

6, 10, 11, 17, 22, 30, 33, 34, 40, 41, 44, 50

c. 47-55

3, 7, 9, 14, 16, 19, 20, 27, 28, 31, 38, 39, 42, 45, 47, 48, 49, 52, 54, 55, 63, 64

d. 56–66

4, 5, 12, 13, 18, 23, 24, 25, 26, 35, 57, 61, 65

e. 67–75

1, 21, 36, 46, 58, 62, 67

f. 76–95

32, 37, 56, 59, 66

g. above 95

29, 53, 60

Dichotomous key to *Calonectria* species

The following key is an adaptation of the key provided by Crous (2002) to include all *Calonectria* spp. described subsequent to 2002. Measurements and observations are those of Crous (2002) and other authors who have described species subsequent to 2002 (Table 1). Only average conidial dimensions, where available, and a few distinguishing characters are presented in the key. Complete descriptions should be consulted to determine species variations. *Calonectria penicilloides* has been omitted from the keys, due to the fact that there is little morphological information available for this species.

1. Stipe extensions thick-walled; acicular to clavate vesicles 2
1. Stipe extensions and vesicles not as above 27
2. Stipe extensions thick-walled, terminating in a acicular to clavate vesicle; fertile branches –3; phialides elongate-doliiform to reniform; macroconidia 1(–3)-septate, 64 × 5 µm; perithecia orange to red; ascospores 1(–3) septate, 40 × 6 µm *Ca. avesiculata*
2. Stipe extensions not thick-walled and vesicles clavate 3
3. Teleomorph state unknown 4
3. Teleomorph state known 13



4. Macroconidia 1-septate 5
4. Macroconidia more than 1-septate 6

5. Stipe extensions terminating in a clavate vesicle; fertile branches –5; phialides doliiform to reniform; macroconidia 1-septate, $53 \times 4.5 \mu\text{m}$ *Ca. brassicae*
5. Stipe extensions terminating in clavate vesicle; fertile branches –3; phialides doliiform to reniform; macroconidia 1-septate, $44 \times 5 \mu\text{m}$ *Ca. pini*

6. Macroconidia 1(–2)-septate, $44 \times 5 \mu\text{m}$; stipe extensions terminating in a clavate vesicle; lateral stipe extensions present; fertile branches –5; phialides doliiform to reniform *Ca. brachiatica*
6. Macroconidia 1(–3)-septate to multi-septate 7

7. Macroconidia longer than $100 \mu\text{m}$ 8
7. Macroconidia shorter than $100 \mu\text{m}$ 9

8. Macroconidia 1(–3)-septate, $104 \times 8 \mu\text{m}$; stipe extension terminate in clavate vesicle; fertile branches –3; phialides cylindrical to allantoid; microconidiophores lacking stipe extension; microconidia 1–3-septate, $44 \times 4 \mu\text{m}$ *Ca. pseudoreteaudii*
8. Macroconidia more than 1-septate 10

9. Macroconidia (1–)3-septate, $63 \times 6.5 \mu\text{m}$; stipe extensions terminating in clavate vesicle; fertile branches –6; phialides cylindrical to allantoid *Ca. australiensis*
9. Macroconidia 1(–3)-septate, $51 \times 4.5 \mu\text{m}$; stipe extensions terminating in clavate vesicles; fertile branches –7; phialides doliiform to reniform *Ca. ecuadoriae*

10. Macroconidia longer than $100 \mu\text{m}$ with more than 6 septa 11
10. Macroconidia shorter than $100 \mu\text{m}$ with 6 or less septa 12



11. Stipe extensions terminating in narrowly clavate vesicles; fertile branches –4; phialides cylindrical; macroconidia (1–)7–10(–12)-septate with slight swelling in the middle, $110 \times 10 \mu\text{m}$; Mega- and microconidia absent *Ca. angustata*
11. Stipe extensions terminating in narrowly clavate vesicles; fertile branches –3; phialides cylindrical; microconidia present, 1-septate, $18 \times 3 \mu\text{m}$; macroconidia (1–)7(–8)-septate, $120 \times 7.5 \mu\text{m}$; megaconidia present, 9–16-septate, bent or curved, $(150–)200–250(–270) \times 6–7(–8) \mu\text{m}$ *Ca. hurae*
12. Stipe extensions terminating in narrowly clavate vesicles; fertile branches –3; phialides cylindrical to allantoid, obpyriform when carried singly; macroconidia 4–6-septate, $69 \times 6 \mu\text{m}$ *Ca. queenslandica*
12. Stipe extensions terminating in a narrowly clavate vesicles; fertile branches –3; phialides cylindrical to allantoid, obpyriform when carried singly; macroconidia 4–6-septate, $76 \times 6 \mu\text{m}$ *Ca. terrae-reginae*
13. Macroconidial state unknown; megaconidiophores with stipe extensions terminating in clavate vesicles when present; megaconidia 6–10-septate, boomerang-shaped or curved, $(120–)150–170(–220) \times 8–9 \mu\text{m}$; microconidia 1–3-septate, straight or curved, $20–65 \times 2.5–3.5 \mu\text{m}$ *Ca. multiseptata*
13. Macroconidial state known 14
14. Teleomorph state known and macroconidia 1-septate to 1(–3)septate 15
14. Teleomorph state known and macroconidia multi-septate 18
15. Teleomorph homothallic 16
15. Teleomorph heterothallic 17
16. Perithecia orange with a red apex; ascospores 1-septate, $35 \times 6.5 \mu\text{m}$; stipe extensions terminating in clavate vesicles; fertile branches –4; phialides doliiform to reniform; macroconidia 1-septate, $45 \times 4.5 \mu\text{m}$ *Ca. gracilipes*



16. Perithecia red; ascospores 1-septate, $37 \times 5 \mu\text{m}$; stipe extensions terminating in narrowly clavate vesicles; fertile branches –4; phialides doliiform to reniform; macroconidia 1(–3)-septate, $56 \times 4.5 \mu\text{m}$ *Ca. gracilis*
17. Perithecia orange; ascospores 1(–3)-septate, $44 \times 5.5 \mu\text{m}$; stipe extensions terminating in narrowly clavate vesicles; fertile branches –4; phialides elongate-doliiform to reniform; macroconidia 1(–3)-septate, $65 \times 5 \mu\text{m}$; microconidia 1-septate, $32 \times 3 \mu\text{m}$ *Ca. clavata*
17. Perithecia red-brown; ascospores 1(–3)-septate, $52 \times 6 \mu\text{m}$; stipe extensions terminating in clavate to narrowly ellipsoidal vesicles; fertile branches –5; phialides elongate-doliiform to reniform; macroconidia 1(–3)-septate, $82 \times 5.5 \mu\text{m}$; microconidia 1-septate, $30 \times 3.5 \mu\text{m}$ *Ca. pteridis*
18. Macroconidia 3-septate 19
18. Macroconidia 3- to multi-septate 23
19. Perithecia yellow to orange 20
19. Perithecia yellow 21
20. Teleomorph state homothallic; perithecia yellow to orange; ascospores (1–)3-septate, $33 \times 6 \mu\text{m}$; stipe extensions terminating in broadly clavate vesicles; fertile branches –5; phialides doliiform to reniform; macroconidia 3-septate, $72 \times 6 \mu\text{m}$ *Ca. eucalypti*
20. Teleomorph state homothallic; perithecia orange to red; ascospores (1–)3-septate, $53 \times 7 \mu\text{m}$; stipe extensions terminating in narrowly clavate vesicle; fertile branches –5; phialides allantoid to reniform; macroconidia (1–)3-septate, $81 \times 6 \mu\text{m}$; megaconidia 7–9(–14)-septate, boomerang-shaped to curved, $130\text{--}200 \times 5\text{--}6 \mu\text{m}$ *Ca. indusiata*
21. Macroconidia and ascospores shorter than $65 \mu\text{m}$; teleomorph state homothallic; perithecia bright yellow; ascospores (1–)3-septate, $50 \times 5.5 \mu\text{m}$; stipe extensions terminating in clavate vesicles; fertile branches –5; phialides allantoid to cylindrical; macroconidia (1–)3-septate, $55 \times 4.5 \mu\text{m}$ *Ca. madagascariensis*

21. Macroconidia and ascospores longer than 65 μm 22
22. Teleomorph state homothallic; perithecia bright yellow; ascospores (1–)3-septate, 55 \times 6 μm ; stipe extensions terminating in clavate vesicles; fertile branches –5; phialides elongate-doliiform to reniform; macroconidia (1–)3-septate, 65 \times 5 μm *Ca. colhounii*
22. Teleomorph state heterothallic; perithecia dirty yellow, ascospores (1–)3-septate, 55 \times 6 μm ; stipe extensions terminating in clavate vesicles; fertile branches –5; phialides allantoid to cylindrical; macroconidia (1–)3(–4)-septate, 90 \times 6.5 μm *Ca. macroconidialis*
23. Macroconidiophore branches –2 or less 24
23. Macroconidiophore with more than 2 series of branches 25
24. Teleomorph state homothallic; perithecia orange-brown; ascospores 3–6(–9)-septate, 90 \times 6.5 μm ; stipe extensions terminating in narrowly clavate vesicles; fertile branches –2; phialides cylindrical; microconidia 1(–3)-septate, (8–)15–30(–50) \times 3–5 μm ; macroconidia 5(–7)-septate, 110 \times 9 μm ; megaconidia 7–13-septate, bent or curved, (120–)180–230 \times (8–)10–11(–13) μm *Ca. rumohrae*
24. Teleomorph state homothallic; perithecia red to red-brown; ascospores 3–4-septate, 70 \times 6 μm ; stipe extensions, when present, terminating in narrowly clavate vesicles; fertile branches –1; macroconidia 5–7-septate, 75 \times 7 μm ; microconidia 1–3-septate, 10–30 \times 3–5 μm *Ca. acicola*
25. Teleomorph state homothallic; perithecia orange to red-brown; ascospores (1–)3-septate, 70 \times 6.5 μm ; stipe extensions terminating in narrowly clavate vesicles; fertile branches –6; phialides cylindrical to allantoid; macroconidia (1–)3–5(–6)-septate, 60 \times 5 μm *Ca. leguminum*
26. Teleomorph state heterothallic; perithecia orange to red-brown; ascospores (1–)5(–6)-septate, 70 \times 5.5 μm ; stipe extensions terminating in clavate vesicles; fertile branches –6; phialides cylindrical to allantoid; macroconidia (1–)5(–6)-septate, 84 \times 6.5 μm ; microconidia 1(–3)-septate, 30 \times 3 μm *Ca. reteaudii*

27. Vesicles sphaeropedunculate, globose or ovoid	28
27. Vesicles not as above	48
28. Vesicles consistently ovate; teleomorph state heterothallic; perithecia orange; ascospores 1–3(–7)-septate, 60 × 5.5 µm; fertile branches –3; phialides doliiform to reniform; macroconidia straight or curved, 1(–3)-septate, 70 × 5 µm; microconidia 1-septate, 21 × 3 µm	<i>Ca. ovata</i>
28. Vesicles not consistently ovate	29
29. Macroconidia 1(–3)-septate	30
29. Macroconidia only 1-septate	35
30. Teleomorph state unknown; stipe extensions terminating in sphaeropedunculate vesicle; fertile branches –3; phialides elongate-doliiform to reniform; macroconidia 1(–3)-septate, 60 × 5 µm	<i>Ca. curvispora</i>
30. Teleomorph state known	31
31. Perithecia red-brown; teleomorph state homothallic; ascospores 1(–3)-septate, 42 × 5 µm; stipe extensions terminating in sphaeropedunculate to ovoid or ellipsoidal to clavate vesicles; fertile branches –3; phialides elongate-doliiform to reniform; macroconidia (1–)3(–4)-septate, 73 × 5 µm; microconidia 1-septate, 27 × 4 µm	<i>Ca. variabilis</i>
31. Perithecia orange to red	32
32. Teleomorph state heterothallic; perithecia orange to red; ascospores 1(–3)-septate, 45 × 5 µm; stipe extensions terminating in globoid or ellipsoid to obpyriform vesicles; fertile branches –5; phialides elongate-doliiform to reniform; macroconidia 1(–3)-septate, 70 × 6 µm; microconidia 1-septate, 39 × 4 µm	<i>Ca. spathiphylli</i>
32. Teleomorph state homothallic	33



33. Lateral stipe extensions abundant; perithecia orange; ascospores 1-septate, $33 \times 5 \mu\text{m}$; stipe extensions terminating in sphaeropedunculate vesicles; fertile branches –5; phialides elongate-doliiform to reniform; macroconidia 1(–3)-septate, $53 \times 4.5 \mu\text{m}$ *Ca. colombiensis*
33. Lateral stipe extensions absent 34
34. Ascospores 1(–3)-septate, $42 \times 5.5 \mu\text{m}$; stipe extensions terminating in sphaeropedunculate to ellipsoidal vesicles; fertile branches –4; phialides elongate-doliiform to reniform; macroconidia 1(–3)-septate, $52 \times 4 \mu\text{m}$ *Ca. pseudospathiphylli*
34. Ascospores 1(–3)-septate, $45 \times 6 \mu\text{m}$; stipe extensions terminating in sphaeropedunculate vesicle; fertile branches –3; phialides elongate-doliiform to reniform; macroconidia (1–)3-septate, $62 \times 6 \mu\text{m}$; microconidia 1-septate, $30 \times 4.5 \mu\text{m}$ *Ca. ilicicola*
35. Stipe thick-walled; teleomorph state unknown; stipe extensions terminating in clavate to sphaeropedunculate vesicle; fertile branches –8; phialides elongate-doliiform to reniform; macroconidia 1-septate, $53 \times 4.5 \mu\text{m}$ *Ca. multiphialidica*
35. Stipe thin-walled 36
36. Teleomorph state known 37
36. Teleomorph state unknown 39
37. Macroconidiophore branches –8; perithecia orange; teleomorph state homothallic; perithecia orange; ascospores 1-septate, $31 \times 6 \mu\text{m}$; stipe extensions terminating in sphaeropedunculate vesicle; phialides elongate-doliiform to reniform; macroconidia 1-septate, $46.5 \times 4 \mu\text{m}$ *Ca. hongkongensis*
37. Macroconidiophore branches –5 38
38. Teleomorph state homothallic; perithecia orange; ascospores 1-septate, $33 \times 6 \mu\text{m}$; stipe extensions terminating in sphaeropedunculate vesicles, lateral stipe extensions

- abundant; phialides doliiform to reniform; macroconidia 1-septate, $53 \times 5 \mu\text{m}$
..... *Ca. asiatica*
38. Teleomorph state homothallic; perithecia orange to red; ascospores 1-septate, $35 \times 5 \mu\text{m}$; stipe extensions terminating in sphaeropedunculate vesicles, lateral stipe extensions abundant; phialides doliiform to reniform; macroconidia 1-septate, $40 \times 3.5 \mu\text{m}$ *Ca. kyotensis*
39. Lateral stipe extensions absent 40
39. Lateral stipe extensions present 42
40. Macroconidia curved, 1-septate, $40\text{--}46 \times 3\text{--}4 \mu\text{m}$; stipe extensions terminating in sphaeropedunculate vesicle; fertile branches –2 *Ca. curvata*
40. Macroconidia straight 41
41. Stipe extensions terminating in globose to ovoid to sphaeropedunculate vesicle; fertile branches –3; phialides doliiform to reniform; macroconidia 1-septate, $51 \times 5 \mu\text{m}$
..... *Ca. humicola*
41. Stipe extensions terminating in sphaeropedunculate vesicles; fertile branches –5; phialides elongate-doliiform to reniform; macroconidia 1-septate, $50.5 \times 4 \mu\text{m}$
..... *Ca. indonesiae*
42. Lateral stipe extensions rare; stipe extensions terminating in pyriform to sphaeropedunculate vesicles; fertile branches – 3; phialides doliiform to reniform; macroconidia 1-septate, $50 \times 4 \mu\text{m}$ *Ca. canadensis*
42. Lateral stipe extensions abundant 43
43. Macroconidiophore branches 4–6 44
43. Macroconidiophore branches –3 45
44. Macroconidiophore branches –4; stipe extension terminating in globose to ovoid to sphaeropedunculate vesicles; phialides doliiform to reniform; macroconidia 1-septate, $54 \times 6 \mu\text{m}$ *Ca. densa*

44. Macroconidiophore branches –6; stipe extensions terminating in sphaerpedunculate vesicles; phialides elongate-doliiform to reniform; macroconidia 1-septate, $47.5 \times 4 \mu\text{m}$ *Ca. malesiana*
45. Macroconidia $45 \times 4 \mu\text{m}$, 1-septate; stipe extensions terminating in sphaeropedunculate vesicles; phialides elongate-doliiform to reniform *Ca. chinensis*
45. Macroconidia longer than $45 \mu\text{m}$ 46
46. Stipe extensions terminating in sphaeropedunculate vesicle; phialides elongate-doliiform to reniform; macroconidia 1-septate, $55 \times 4.5 \mu\text{m}$ *Ca. pacifica*
46. Stipe extensions terminating in sphaeropedunculate vesicles; phialides elongate-doliiform to reniform; macroconidia 1-septate, $58 \times 5 \mu\text{m}$ *Ca. sumatrensis*
47. Vesicles pyriform to ellipsoidal or clavate, rarely ovoid, never obpyriform 48
47. Vesicles not as above 54
48. Macroconidia more than 1-septate 49
48. Macroconidia 1-septate 50
49. Teleomorph state unknown; stipe extensions terminating in narrowly ellipsoidal to pyriform or ovoid to sphaeropedunculate vesicles; fertile branches –3; phialides doliiform to reniform; macroconidia (1–)3-septate, $58 \times 4 \mu\text{m}$ *Ca. citri*
49. Teleomorph state homothallic; perithecia orange-red; ascospores 1(–3)-septate, $33.5\text{--}69 \times 4.5\text{--}7 \mu\text{m}$; stipe extensions terminating in clavate to ovoid or ellipsoidal vesicles; fertile branches –4; phialides doliiform to reniform; macroconidia (1–)3(–5)-septate, $(44\text{--})50\text{--}70(–102) \times 5\text{--}7(–8) \mu\text{m}$ *Ca. hederæ*
50. Macroconidia curved, 1-septate, $56 \times 4 \mu\text{m}$, stipe extensions terminating in ellipsoidal to clavate vesicles; fertile branches –4; phialides doliiform to reniform; teleomorph state unknown *Ca. hawksworthii*



50. Macroconidia straight	51
51. Stipe extensions up to 200 μm long	52
51. Stipe extensions longer than 200 μm	53
52. Teleomorph state heterothallic; perithecia yellow to orange; ascospores 1-septate, 37 \times 6 μm ; stipe extensions terminating in ellipsoidal to pyriform or clavate vesicle; fertile branches –6; phialides doliiform to reniform; macroconidia 1-septate, 45 \times 4 μm	<i>Ca. morgani</i>
52. Teleomorph state unknown; stipe extensions terminating in oval to ellipsoidal to fusiform vesicle; fertile branches –3; phialides doliiform to reniform; macroconidia 1-septate, 38 \times 3.5 μm	<i>Ca. brasiliensis</i>
53. Teleomorph state unknown; stipe extensions terminating in fusiform to pyriform or clavate vesicle; fertile branches –5; phialides doliiform to reniform; macroconidia 1-septate, 48 \times 4 μm	<i>Ca. orientalis</i>
53. Teleomorph state unknown; stipe extensions terminating in broadly clavate to ellipsoidal vesicles; fertile branches –5; phialides doliiform to reniform; macroconidia 1-septate, 48 \times 4 μm	<i>Ca. sulawesiensis</i>
54. Vesicles obpyriform to ellipsoidal	55
54. Vesicles naviculate	66
55. Macroconidia 1-septate	56
55. Macroconidia more than 1-septate	64
56. Macroconidiophore branches –3	57
56. Macroconidiophore branches 4–6	59

57. Teleomorph state homothallic; perithecia orange to red; ascospores 1-septate, 32×4 μm ; stipe extensions terminating in broadly clavate to obpyriform vesicle; phialides doliiform to reniform; macroconidia 1-septate, 36×4 μm *Ca. zuluensis*
57. Teleomorph state heterothallic 58
58. Perithecia orange to red-brown; ascospores 1-septate, 35×6.5 μm ; stipe extensions terminating in obpyriform to ellipsoidal vesicles; phialides doliiform to reniform; macroconidia 1-septate, 50×4.5 μm *Ca. pauciramosa*
58. Teleomorph state unknown; stipe extensions terminating in broadly clavate to obpyriform vesicles; phialides doliiform to reniform; macroconidia 1-septate, 37×4 μm *Ca. polizzii*
59. Macroconidia up to 45 μm long 60
59. Macroconidia longer than 45 μm 63
60. Macroconidiophore branches –6; teleomorph state heterothallic; perithecia orange to red; ascospores 1-septate, 33×6 μm ; stipe extensions terminating in obpyriform to broadly ellipsoidal vesicle; phialides doliiform to reniform; macroconidia 1-septate, 45×4 μm *Ca. insularis*
60. Macroconidiophore branches –4 61
61. Vesicles broadly ellipsoidal with a papillate apex; phialides doliiform to reniform; macroconidia 1-septate, 45×4 μm ; teleomorph state heterothallic; perithecia orange to red; ascospores 1-septate, 50×5.5 μm *Ca. mexicana*
61. Vesicles fusiform to obpyriform 62
62. Teleomorph state homothallic; perithecia yellow to orange; ascospores 1-septate, 34×4 μm ; phialides doliiform to reniform; macroconidia 1-septate, 37×3 μm *Ca. colombiana*
62. Teleomorph state unknown; phialides doliiform to reniform; macroconidia 1-septate, 44×5 μm *Ca. cerciana*



63. Teleomorph state heterothallic; perithecia red-brown; ascospores 1-septate, 48×5.5 μm ; stipe extensions terminating in ellipsoidal to narrowly obpyriform; fertile branches –5; phialides doliiform to reniform; macroconidia 1-septate, 60×4.5 μm *Ca. scoparia*
63. Teleomorph state unknown; stipe extensions terminating in obpyriform to ellipsoidal vesicles; fertile branches –4; phialides doliiform to reniform; macroconidia 1-septate, 48×4 μm *Ca. pseudoscoparia*
64. Macroconidiophore branches –6; stipe extensions terminating in ellipsoidal to obpyriform vesicles; phialides cylindrical, straight or doliiform to reniform; macroconidia 1(–3)-septate, 73×5 μm *Ca. leucothoës*
64. Macroconidiophore branches –3 65
65. Teleomorph state homothallic; perithecia orange to red-brown; ascospores 1(–3)-septate, 50×5.5 μm ; stipe extensions terminating in obpyriform to broadly ellipsoidal vesicles; phialides doliiform to reniform; macroconidia (1–)3-septate, $50\text{--}70 \times 5\text{--}6$ μm *Ca. pyrochoa*
65. Teleomorph state homothallic; perithecia orange; ascospores (1–)3-septate, 50×5.5 μm ; stipe extensions terminating in ellipsoidal to obpyriform or clavate vesicles; phialides cylindrical, straight or doliiform to reniform; macroconidia (1–)3(–6)-septate, 55×4 μm *Ca. spathulata*
66. Teleomorph state heterothallic; perithecia red-brown; ascospores 1(–3)-septate, 40×5 μm ; fertile branches –5; phialides doliiform to reniform; macroconidia 1-septate, 45×3 μm *Ca. naviculata*
66. Teleomorph state unknown; fertile branches –3; phialides doliiform to reniform; macroconidia 1-septate, $42\text{--}68 \times 4\text{--}6$ μm *Ca. pseudonaviculata*

REFERENCES

- Alfieri SA, El-Gholl NE, Schoulties CL. (1982). Homothallism in *Calonectria ilicicola*. *Mycologia* **74**: 513–514.
- Altschul SF, Gish, W, Miller, W, Myers, EW, Lipman DJ. (1990). Basic Local Alignment Search Tool. *Journal of Molecular Biology* **215**: 403–410.
- Boedjin KB, Reitsma J. (1950). Notes on the genus *Cylindrocladium*. *Reinwardtia* **1**: 51–60.
- Boesewinkel HJ. (1982). Heterogeneity within *Cylindrocladium* and its teleomorphs. *Transactions of the British Mycological Society* **78**: 553–556.
- Carbone I, Kohn L.M. (1999). A method for designing primer sets for speciation studies in filamentous ascomycetes. *Mycologia* **91**: 553–556.
- Chase AR, Poole RT. (1987). Effects of potting medium pH and air temperature on severity of *Cylindrocladium* root and petiole rot of *Spathiphyllum* sp. *Plant Disease* **71**: 509–511.
- Crous PW. (2002) Taxonomy and pathology of *Cylindrocladium* (*Calonectria*) and allied genera. APS Press, St. Paul, Minnesota, U.S.A.
- Crous PW, Alfenas AC, Junghans TG. (1998a). Variability within *Calonectria ovata* and its anamorph *Cylindrocladium ovatum* from Brazil. *Sydowia* **50**: 1–13.
- Crous PW, Alfenas AC, Wingfield MJ. (1993a). *Calonectria scoparia* and *Calonectria morganii* sp. nov., and variation among isolates of their *Cylindrocladium* anamorphs. *Mycological Research* **97**: 701–708.
- Crous, PW, Gams W, Staplers JA, Roberts V, Stegehuis G. (2004a). MycoBank: an online initiative to launch mycology into the 21st century. *Studies in Mycology* **50**: 19–22.
- Crous PW, Groenewald JZ, Hill CF. (2002). *Cylindrocladium pseudonaviculatum* sp. nov. from New Zealand, and new *Cylindrocladium* records from Vietnam. *Sydowia* **54**: 23–33.
- Crous PW, Groenewald JZ, Risède J-M, Simoneau P, Hyde KD. (2006). *Calonectria* species and their *Cylindrocladium* anamorphs: species with clavate vesicles. *Studies in Mycology* **55**: 213–226.
- Crous PW, Groenewald JZ, Risède J-M, Simoneau P, Hywel-Jones NL. (2004b). *Calonectria* species and their *Cylindrocladium* anamorphs: species with sphaeropedunculate vesicles. *Studies in Mycology* **50**: 415–430.
- Crous PW, Janse BJH, Victor D, Marais GF, Alfenas AC. (1993b). Molecular characterization of *Cylindrocladium* spp. with three-septate conidia and ovoid-like vesicles. *Systematic and Applied Microbiology* **16**: 266–273.

- Crous PW, Kang JC, Schoch CL, Mchau GRA. (1999). Phylogenetic relationships of *Cylindrocladium pseudogracile* and *Cylindrocladium rumohrae* with morphologically similar taxa, based on morphology and DNA sequences of internal transcribed spacers and β -tubulin. *Canadian Journal of Botany* **77**: 1813–1820.
- Crous PW, Krof A, Van Zyl WH. (1995). Nuclear DNA polymorphisms of *Cylindrocladium* species with 1-septate conidia and clavate vesicles. *Systematic and Applied Microbiology* **18**: 224–250.
- Crous PW, Peeraly A. (1996). *Gliocladiopsis irregular* sp. nov. and notes on *Cylindrocladium spathiphylli*. *Mycotaxon* **58**: 119–128.
- Crous PW, Phillips AJL & Wingfield MJ. (1991). The genera *Cylindrocladium* and *Cylindrocladiella* in South Africa, with special reference to forest nurseries. *South African Forestry Journal* **157**: 69–89.
- Crous PW, Phillips AJL, Wingfield MJ. (1992). Effects of cultural conditions on vesicle and conidium morphology in species of *Cylindrocladium* and *Cylindrocladiella*. *Mycologia* **84**: 497–504.
- Crous PW, Seifert KA. (1998). Megaconidia as an additional taxonomic character in *Cylindrocladium*, with a note on *Cylindrocladiopsis*. *Fungal Diversity* **1**: 51–62.
- Crous PW, Theron L, Van Zyl WH. (1997). Delineation of *Cylindrocladium* species with 1–3-septate conidia and clavate vesicles based on morphology and rDNA RFLPs. *Mycological Research* **101**: 210–214.
- Crous PW, Wingfield MJ. (1994). A monograph of *Cylindrocladium*, including anamorphs of *Calonectria*. *Mycotaxon* **51**: 341–435.
- Crous PW, Wingfield MJ, Mohammed C, Yuan ZQ. (1998b). New foliar pathogens of *Eucalyptus* from Australia and Indonesia. *Mycological Research* **102**: 527–532.
- Cunningham CW. (1997). Can three incongruency tests predict when data should be combined? *Molecular Biology and Evolution* **14**: 733–740.
- De Hoog GS, Van den Ende AHG. (1998). Molecular diagnostics of clinical strains of filamentous Basidiomycetes. *Mycoses* **41**: 183–189.
- Dettman JR, Jacobson DJ, Taylor JW. (2003). A multilocus genealogical approach to phylogenetic species recognition in the model eukaryote *Neurospora*. *Evolution* **57**: 2703–2720.
- El-Gholl NE, Alfenas AC, Crous PW, Schubert TS. (1993). Description and pathogenicity of *Cylindrocladium ovatum* sp. nov. *Canadian Journal of Botany* **71**: 466–470.

- El-Gholl NE, Alfenas AC, Junghans, DT, Schubert TS, Miller JW, Leahy EM. (1997). Description of *Calonectria rumohrae* sp. nov. (anamorph = *Cylindrocladium rumohrae* sp. nov.) *Mycotaxon* **64**: 467–484.
- El-Gholl NE, Uchida JY, Alfenas AC, Schubert T S, Alfieri SA, Chase AR. (1992). Induction and description of perithecia of *Calonectria spathiphylli* sp. nov. *Mycotaxon* **45**: 285–300.
- Farris JS, Källersjö M, Kluge AG, Bult C. (1994). Testing significance of incongruence. *Cladistics* **10**: 315–320.
- Fisher NL, Burgess LW, Toussoun TA, Nelson PE. 1982. Carnation leaves as a substrate and for preserving cultures of *Fusarium* species. *Phytopathology* **72**: 151–153.
- Gadgill PD, Dick MA. (2004). Fungi silvicolae novazelandiae: 5. *New Zealand Journal of Forestry Science* **34**: 316–323.
- Geurber JC, Correll JC. (2001). Characterization of *Glomerella acutata*, the teleomorph of *Colletotrichum acutatum*. *Mycologia* **93**: 216–229.
- Gueidan C, Roux C, Lutzoni F. (2007). Using multigene phylogeny analysis to assess generic delineation and character evolution in *Verrucariaceae* (*Verrucariales*, *Ascomycota*). *Mycological Research* **111**: 1145–1168.
- Halleen F, Schroers H-J, Groenewald JZ, Rego C, Oliveira H, Crous PW. (2006). *Neonectria liriodendra* sp. nov., the main causal agent of black foot disease of grapevine. *Studies in Mycology* **55**: 227–234.
- Hawksworth DL. (2005). Two major changes in fungal nomenclature enacted in Vienna. *Mycological Research* **109**: 1061–1062.
- Helgason T, Watson IJ, Young PW. (2003). Phylogeny of the *Glomerales* and *Diversisporales* (Fungi: *Glomeromycota*) from actin and elongation factor 1-alpha sequences. *FEMS Microbiology Letters* **229**: 127–132.
- Henricot B & Culham A. (2002). *Cylindrocladium buxicola*, a new species affecting *Buxus* spp., and its phylogenetic status. *Mycologia* **94**: 980–997.
- Hibbett DS, Binder M, Bischoff JF, Blackwell M, Cannon PF, *et al.* (2007). A higher-level phylogenetic classification of the Fungi. *Mycological Research* **111**: 509–547.
- Hillis DM, Bull JJ. 1993. An empirical test of bootstrapping as a method for assessing confidence in phylogenetic analysis. *Systematic Biology* **42**: 182–192.
- Hunter BB, Barnett HL. (1978). Growth and sporulation of species and isolates of *Cylindrocladium* in culture. *Mycologia* **70**: 614–635.

- Hunter GC, Wingfield BD, Crous PW, Wingfield MJ. (2006). A multi-gene phylogeny for species of *Mycosphaerella* occurring on *Eucalyptus* leaves. *Studies in Mycology* **55**: 147–161.
- Jeng RS, Dumas M, Liu FH, Wang CL, Hubbes M. (1997). DNA analysis of *Cylindrocladium floridanum* isolates from selected forest nurseries. *Mycological Research* **101**: 285–291.
- Kang JC, Crous PW, Old KM, Dubzinski MJ. (2001a). Non-conspecificity of *Cylindrocladium quinqueseptatum* and *Calonectria quinqueseptata* based on a β -tubulin gene phylogeny and morphology. *Canadian Journal of Botany* **79**: 1241–1247.
- Kang JC, Crous PW, Schoch CL. (2001b). Species concepts in the *Cylindrocladium floridanum* and *Cy. spathiphylli* complexes (*Hypocreaceae*) based on multi-allelic sequence data, sexual compatibility and morphology. *Systematic and Applied Microbiology* **24**: 206–217.
- Katoh K, Kuma K, Toh H, Miyata T. (2005). MAFFT version 5: improvement in accuracy of multiple sequence alignment. *Nucleic Acid Research* **33**: 511–518.
- Kim KD, Russin JS, Snow JP. (1998). Susceptibility to *Calonectria ilicicola* in soybean grown in greenhouse and field. *Korean Journal of Crop Science* **43**: 239–244.
- Librado P, Rozas J. (2009). DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* **25**: 1451–1452.
- Lombard L, Bogale M, Montenegro F, Wingfield BD, Wingfield MJ. (2008). A new bark canker disease of the tropical hardwood tree *Cedrelinga cateniformis* in Ecuador. *Fungal Diversity* **31**: 73–81.
- Lombard L, Crous PW, Wingfield BD, Wingfield MJ. (2009a). Multigene phylogeny and mating tests reveal three cryptic species related to *Calonectria pauciramosa*. *Studies in Mycology*: submitted.
- Lombard L, Rodas CA, Crous PW, Wingfield BD, Wingfield MJ. (2009b). *Calonectria* (*Cylindrocladium*) species associated with dying *Pinus* cuttings. *Persoonia*: **23**: 41–47.
- Lombard L, Zhou XD, Crous PW, Wingfield BD, Wingfield MJ. (2009c). *Calonectria* species associated with cutting rot of *Eucalyptus*. *Persoonia*: submitted.
- Mason-Gamer R, Kellogg E. (1996). Testing for phylogenetic conflict among molecular datasets in the tribe *Tiriceae* (*Graminae*). *Systematic Biology* **45**: 524–545.

- McNiell J, Stuessy TF, Turland NJ, Hörandl E. (2005). XVII International Botanical Congress: preliminary mail vote and report of Congress action on nomenclature proposals. *Taxon* **54**: 1057–1064.
- Moncalvo JM, Wang HH, Hseu RS. (1995). Phylogenetic relationships in *Ganoderma* inferred from the internal transcribed spacers and 25S ribosomal DNA sequences. *Mycologia* **87**: 223–238.
- Morgan AP. (1892). Two new genera of hyphomycetes. *Botanical Gazette* **17**: 190–192.
- Nirenburg HI. (1981). A simplified method for identifying *Fusarium* spp. occurring on wheat. *Canadian Journal of Botany* **59**: 1599–1609.
- Nylander JAA. (2004). MrModeltest v.2. Programme distributed by the author. Evolutionary Biology Centre, Uppsala University.
- Overmeyer C, Lünneberg S, Von Wallburnn C, Meinhardt F. (1996). Genetic variability among isolates and sexual offspring of the plant pathogenic fungus *Calonectria morganii* on the basis of random amplification of polymorphic DNA (RAPD) and restriction fragment length polymorphism (RFLP). *Current Microbiology* **33**: 249–255.
- O'Donnell K, Cigelnik E. (1997). Two divergent intragenomic rDNA ITS2 types within a monophyletic lineage of the fungus *Fusarium* are nonorthologous. *Molecular Phylogenetics and Evolution* **7**: 103–116.
- O'Donnell K, Kistler HC, Cigelnik E, Ploetz RC. (1998). Multiple evolutionary origins of the fungus causing Panama disease of banana: Concordant evidence from nuclear and mitochondrial gene genealogies. *Proceedings of the National Academy of Science of the USA* **95**: 2044–2049.
- Peerally A. (1991). The classification and phytopathology of *Cylindrocladium* species. *Mycotaxon* **40**: 367–366.
- Polizzi G, Grasso FM, Vitale A, Aiello D. (2007). First occurrence of *Calonectria* leaf spot on mexican blue palm in Italy. *Plant Disease* **91**: 1057.
- Posada D, Crandall KA. (1998). Modeltest: testing the model of DNA substitution. *Bioinformatics* **14**: 817–818.
- Rayner RW. (1970). *A mycological colour chart*. Commonwealth Mycological Institute, Kew, Surrey. British Mycological Society.
- Riséde JM, Simoneau P. (2001). Typing *Cylindrocladium* species by analysis of ribosomal DNA spacers polymorphism: application to field isolates from the banana rhizosphere. *Mycologia* **93**: 494–504.

- Riséde JM, Simoneau P. (2004). Pathogenic and genetic diversity of soilborne isolates of *Cylindrocladium* from banana cropping systems. *European Journal of Plant Pathology* **110**: 139–154.
- Ronquist F, Heulsenbeck JP. (2003). MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* **19**: 1572–1574.
- Rossmann AY. (1979). *Calonectria* and its type species, *C. daldiniana*, a later synonym of *C. pyrochroa*. *Mycotaxon* **8**: 321–328.
- Rossmann AY. (1983). The phragmosporous species of *Nectria* and related genera. *Mycological Papers* **150**: 1–164.
- Rossmann AY. (1993). Holomorphic hypocrealean fungi: *Nectria* sensu stricto and telemorphs of *Fusarium*. In: *The fungal holomorph: mitotic, meiotic and pleomorphic speciation in fungal systematics*. (Reynolds DR, Taylor JW, eds.). CAB International, Wallingford, U.K.: 149–160.
- Rossmann AY, Samuels GJ, Rogerson CT, Lowen R. (1999). Genera of Bionectriaceae, Hypocreaceae and Nectriaceae (Hypocreales, Ascomycetes). *Studies in Mycology* **42**: 1–248.
- Schoch CL, Crous PW, Polizzi G, Koike ST. (2001a). Female fertility and single nucleotide polymorphism comparisons in *Cylindrocladium pauciramosum*. *Plant Disease* **85**: 941–946.
- Schoch CL, Crous PW, Wingfield BD, Wingfield MJ. (1999). The *Cylindrocladium candelabrum* species complex includes four distinct mating populations. *Mycologia* **91**: 286–298.
- Schoch CL, Crous PW, Wingfield BD, Wingfield MJ. (2001b). Phylogeny of *Calonectria* based on comparisons of β -tubulin DNA sequences. *Mycological Research* **105**: 1045–1052.
- Schoch CL, Crous PW, Cronright G, Witthuhn RC, El-Gholl NE, Wingfield BD. (2000a). Recombination in *Calonectria morganii* and phylogeny with other heterothallic small-spored *Calonectria* species. *Mycologia* **92**: 665–673.
- Schoch CL, Crous PW, Wingfield MJ, Wingfield BD. (2000b). Phylogeny of *Calonectria* and selected hypocrealean genera with cylindrical macroconidia. *Studies in Mycology* **45**: 45–62.

- Schoch CL, Sung G-H, López-Giráldez F, Townsend JP, Miadlikowska J *et al.* (2009). The Ascomycota Tree of Life: A phylum-wide phylogeny clarifies the origin and evolution of fundamental reproductive and ecological traits. *Systematic Biology* **58**: 224–239.
- Schubert TS, El-Gholl NE, Alfieri SA, Schouties CL. (1989). *Calonectria avesiculata* sp. nov. *Canadian Journal of Botany* **67**: 2414–2419.
- Sober EK. (1971). A macro-conidial form of *Cylindrocladium theae* occurring on glycerol-water agar. *Georgia Academy of Science Bulletin* **29**: 98
- Sobers EK, Alfieri SA. (1972). Species of *Cylindrocladium* and their hosts in Florida and Georgia. *Proceedings of the Florida State Horticultural Society* **85**: 366–369.
- Swofford DL. (2002). PAUP*. Phylogenetic analysis using parsimony (* and other methods), 4.0b10. Computer programme. Sunderland, Massachusetts, USA: Sinauer Associates.
- Taylor JW, Jacobson DJ, Kroken SM, Kasuga T, Geiser DM, Hibbett DS, Fisher MC. (2000). Phylogenetic species recognition and species concepts in fungi. *Fungal Genetics and Biology* **31**: 21–32.
- Victor D, Crous PW, Janse BJH, Wingfield MJ. (1997). Genetic variation in *Cylindrocladium floridanum* and other morphologically similar *Cylindrocladium* species. *Systematic and Applied Microbiology* **20**: 268–285.
- Vilgalys R, Hester M. (1990). Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* **172**: 4238–4246.
- Vitale A, Polizzi G. (2008). First record of leaf spots and stem lesions on *Pistacia lentiscus* caused by *Cylindrocladium pauciramosum* and *C. scoparium* in Italy. *Plant Pathology* **57**: 384.
- White TJ, Burns T, Lee S, Taylor J. (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: *PCR protocols: a guide to methods and applications*. (Innis MA, Gelfand DH, Snisky JJ, White TJ, eds.). Academic Press, U.S.A.: 282–287.



Table 1. Isolates of *Calonectria* spp. studied.

Species	Isolate number ¹	Other collections ¹	GenBank accession nr. ²							Reference ³
			ACT	BT	CAL	HIS3	ITS	LSU	TEF-1 α	
<i>Ca. acicola</i>	CBS 114812		GQ280424	DQ190590	GQ267359	DQ190692	GQ280546	GQ280668	GQ267291	Gadgill & Dick 2004
	CBS 114813 ^T	CMW 30996	GQ280425	DQ190591	GQ267360	DQ190693	GQ280547	GQ280669	GQ267292	
<i>Ca. angustata</i>	CBS 109065 ^T	CMW 30990 = CPC 2347 = P99-0454	GQ280426	AF207543	GQ267361	DQ190696	GQ280548	GQ280670	FJ918551	Crous 2002
	CBS 109169	CMW 30983 = CPC 3152 = P99-1321	GQ280427	DQ190593	GQ267362	DQ190695	GQ280549	GQ280670	FJ918552	
<i>Ca. asiatica</i>	CBS 112711	CPC 3898 = SFE 744	GQ280429	AY725613	AY725738	AY725655	GQ280551	GQ280673	AY725702	Crous <i>et al.</i> 2004b
	CBS 114073 ^T	CMW 23782 = CPC 3900 = SFE 726	GQ280428	AY725616	AY725741	AY725658	GQ280550	GQ280672	AY725705	
<i>Ca. australiensis</i>	CBS 112954 ^T	CMW 23669 = CPC 4714	GQ280430	DQ190596	GQ267363	DQ190699	GQ280552	GQ280674	GQ267293	Crous <i>et al.</i> 2006
<i>Ca. avesiculata</i>	CBS 313.92 ^T	CMW 23670 = CPC 2373 = ATCC 38226	GQ280431	AF333392	GQ267364	DQ190620	GQ280553	GQ280675	GQ267294	Crous 2002
<i>Ca. brachiatica</i>	CBS 123700 ^T	CMW 25298	GQ280433	FJ696388	GQ267366	FJ696396	GQ280555	GQ280677	GQ267296	Lombard <i>et al.</i> 2009b
	CMW 25302		GQ280432	FJ716708	GQ267365	FJ716712	GQ280554	GQ280676	GQ267295	
<i>Ca. brassicae</i>	CBS 111478	CMW 30981	GQ280455	DQ190611	GQ267383	DQ190719	GQ280577	GQ280699	FJ918567	Crous 2002
	CBS 111869 ^T	CMW 30982 = CPC 2409 = PC 551197	GQ280454	AF232857	GQ267382	DQ190720	GQ280576	GQ280698	FJ918566	
<i>Ca. brasiliensis</i>	CBS 230.51 ^T	CMW 23670 = CPC 2390	GQ280502	GQ267241	GQ267421	GQ267259	GQ280624	GQ280746	GQ267328	Lombard <i>et al.</i> 2009a



Table 1. (Continued)

Species	Isolate number ¹	Other collections ¹	GenBank accession nr. ²							Reference ³
			ACT	BT	CAL	HIS3	ITS	LSU	TEF-1 α	
<i>Ca. brasiliensis</i>	CBS 114257	CMW 32949 = CPC 1944	GQ280503	GQ267242	GQ267422	GQ267260	GQ280625	GQ280747	GQ267329	
<i>Ca. canadensis</i>	CBS 110817 ^T	CMW 23673 = CPC 499	GQ280434	AF348212	AY725743	AF348228	GQ280556	GQ280678	GQ267297	Crous 2002
<i>Ca. cerciana</i>	CBS 123693 ^T	CMW 25309	GQ280437	FJ918510	GQ267369	FJ918528	GQ280559	GQ280681	FJ918559	Lombard <i>et al.</i> 2009c
	CBS 123695	CMW 25290	GQ280438	FJ918511	GQ267370	FJ918529	GQ280560	GQ280682	FJ918560	
<i>Ca. chinensis</i>	CBS 112744	CMW 30986 = CPC 4104	GQ280440	AY725618	AY725746	AY725660	GQ280562	GQ280684	AY725709	Crous <i>et al.</i> 2004b
	CBS 114827 ^T	CMW 23674 = CPC 4101	GQ2804390	AY725619	AY725747	AY725661	GQ280561	GQ280683	AY725710	
<i>Ca. citri</i>	CBS 186.36 ^T	CMW 23675	GQ280441	AF333393	GQ267371	GQ267247	GQ280563	GQ280685	GQ267299	Crous 2002
<i>Ca. clavata</i>	CBS 114557 ^T	CMW 23690 = CPC 2536 = ATCC 66389	GQ280449	AF333396	GQ267377	DQ190623	GQ280571	GQ280693	GQ267305	Crous 2002
	CBS 114666 ^T	CMW 30994 = CPC 2537	GQ280450	DQ190549	GQ267378	DQ190624	GQ280572	GQ280694	GQ267306	
<i>Ca. colhounii</i>	CBS 293.79 ^T	CMW 30999	GQ280443	DQ190564	GQ267373	DQ190639	GQ280565	GQ280687	GQ267301	Crous 2002
	CBS 114704		GQ280442	DQ190563	GQ267372	DQ190638	GQ280564	GQ280686	GQ267300	
<i>Ca. colombiensis</i>	CBS 112220 ^T	CMW 23676 = CPC 723	GQ280444	GQ267207	AY725748	AY725662	GQ280566	GQ280688	AY725711	Crous <i>et al.</i> 2004b
	CBS 112221	CMW 30985 = CPC 724	GQ280445	AY725620	AY725749	AY725663	GQ280567	GQ280689	AY725712	Crous 2002
<i>Ca. curvispora</i>	CBS 116159 ^T	CMW 23693	GQ280446	AF333396	GQ267374	AY725664	GQ280568	GQ280690	GQ267302	



Table 1. (Continued)

Species	Isolate number ¹	Other collections ¹	GenBank accession nr. ²							Reference ³
			ACT	BT	CAL	HIS3	ITS	LSU	TEF-1 α	
<i>Ca. densa</i>	CBS 125249	CMW 31184	GQ280523	GQ267230	GQ267442	GQ267279	GQ280645	GQ280767	GQ267350	This study
	CBS 125250	CMW 31185	GQ280524	GQ267231	GQ267443	GQ267280	GQ280646	GQ280768	GQ267351	
	CBS 125261 ^T	CMW 31182	GQ280525	GQ267232	GQ267444	GQ267281	GQ280647	GQ280769	GQ267352	
<i>Ca. ecuadoriae</i>	CBS 111394	CMW 30980 = CPC 1628	GQ280448	DQ190599	GQ267376	DQ190704	GQ280570	GQ280692	GQ267304	Crous <i>et al.</i> 2006
	CBS 111406 ^T	CMW 23677 = CPC 1635	GQ280447	DQ190600	DQ267375	DQ190705	GQ280569	GQ280691	GQ267303	
<i>Ca. eucalypti</i>	CBS 125273	CMW 14890	GQ280510	GQ267217	GQ267429	GQ267266	GQ280632	GQ280754	GQ267337	This study
	CBS 125274	CMW 18443	GQ280509	GQ267216	GQ267428	GQ267265	GQ280631	GQ280753	GQ267336	
	CBS 125275 ^T	CMW18444	GQ280511	GQ267218	GQ267430	GQ267267	GQ280633	GQ280755	GQ267338	
	CBS 125276	CMW 18445	GQ280512	GQ267219	GQ267431	GQ267268	GQ280634	GQ280756	GQ267339	
<i>Ca. gracilipes</i>	CBS 111141 ^T		GQ280457	DQ190566	GQ267385	DQ190644	GQ280579	GQ280701	GQ267311	Crous 2002
	CBS 115674		GQ280456	AF333406	GQ267384	DQ190645	GQ280578	GQ280700	GQ267310	
<i>Ca. gracilis</i>	CBS 111284		GQ280489	DQ190567	GQ267408	DQ190647	GQ280611	GQ280733	GQ267324	Crous 2002
	CBS 111807		GQ280488	AF232858	GQ267407	DQ190646	GQ280610	GQ280734	GQ267323	
<i>Ca. hawksworthii</i>	CBS 111870 ^T	CPC 2405 = MUCL 30866	GQ280458	AF333407	GQ267386	DQ190649	GQ280580	GQ280702	FJ918558	Crous 2002
<i>Ca. hongkongensis</i>	CBS 114711	CMW 30995	GQ280460	AY725621	AY725754	AY725666	GQ280582	GQ280704	AY725716	Crous <i>et al.</i> 2004b
	CBS 114828 ^T		GQ280459	AY725622	AY725755	AY725667	GQ280581	GQ280703	AY725717	
<i>Ca. humicola</i>	CBS 125251 ^T	CMW 31183	GQ280526	GQ267233	GQ267445	GQ267282	GQ280648	GQ280770	GQ267353	This study
	CBS 125252	CMW 31186	GQ280527	GQ267234	GQ267446	GQ267283	GQ280649	GQ280771	GQ267354	
	CBS 125269	CMW31187	GQ280528	GQ267235	GQ267447	GQ267284	GQ280650	GQ280772	GQ267355	



Table 1. (Continued)

Species	Isolate number ¹	Other collections ¹	GenBank accession nr. ²							Reference ³
			ACT	BT	CAL	HIS3	ITS	LSU	TEF-1 α	
<i>Ca. hurae</i>	CBS 114551	CMW 16720 = CPC 2344	GQ280461	AF333408	GQ267387	DQ190728	GQ280583	GQ280705	FJ918548	Crous 2002
<i>Ca. ilicicola</i>	CBS 190.50 ^T	CMW 30998 = CPC 2482 = IMI 299389	GQ280483	AY725631	AY725764	AY725676	GQ280605	GQ280727	AY725726	Crous 2002
	CBS 115897		GQ280484	AY725647	GQ267403	GQ267256	GQ280606	GQ280728	AY725729	
<i>Ca. indonesiae</i>	CBS 112823 ^T	CMW 23683 = CPC 4508	GQ280463	AY725623	AY725756	AY725668	GQ280585	GQ280707	AY725718	Crous <i>et al.</i> 2004b
	CBS 112840	CPC 4547	GQ280464	AY725625	AY725758	AY725670	GQ280586	GQ280708	AY725720	
<i>Ca. industriata</i>	CBS 144.36	CMW 23699	GQ280536	GQ267239	GQ267453	GQ267262	GQ280658	GQ280780	GQ267332	Crous 2002
	CBS 114684	CPC 2446 = UFV 16A	GQ280537	AF232862	GQ267454	DQ190652	GQ280659	GQ280781	GQ2673333	
<i>Ca. insularis</i>	CBS 114558 ^T	CMW 30991	GQ280465	AF210861	GQ267389	FJ918526	GQ280587	GQ280709	FJ918556	Crous 2002
	CBS 114559	CMW 30992	GQ280466	AF210862	GQ267390	FJ918525	GQ280588	GQ280710	FJ918555	
<i>Ca. kyotensis</i>	CBS 170.77	CMW 23679 = IMI 299388	GQ280452	GQ267209	GQ267380	GQ267249	GQ280574	GQ280696	GQ267308	Crous 2002
	CBS 413.67	CMW 23678 = CPC 2391	GQ280451	GQ267208	GQ267379	GQ267248	GQ280573	GQ280695	GQ267307	
<i>Ca. leguminum</i>	CBS 728.68 ^T	CMW 23684 = IMI 299578	GQ280467	AF389837	GQ267391	DQ190654	GQ280589	GQ280711	FJ918547	Crous 2002
<i>Ca. leucothoë</i>	CBS 109166	CMW 30977 = CPC 3612 = P97-2605	GQ280468	FJ918508	GQ267392	FJ918523	GQ280590	GQ280712	FJ918553	Crous 2002
<i>Ca. macroconidialis</i>	CBS 114880 ^T	CPC 307	GQ280469	AF232855	GQ267393	DQ190655	GQ280591	GQ280713	GQ267313	Crous 2002
<i>Ca. madagascariensis</i>	CBS 114571	CMW 30993 = CPC 2253	GQ280471	DQ190571	GQ267395	DQ190657	GQ280593	GQ280715	GQ267315	Crous 2002



Table 1. (Continued)

Species	Isolate number ¹	Other collections ¹	GenBank accession nr. ²							Reference ³
			ACT	BT	CAL	HIS3	ITS	LSU	TEF-1 α	
<i>Ca. madagascariensis</i>	CBS 114572 ^T	CMW 23686 = CPC 2252	GQ280471	DQ190572	GQ267394	DQ190658	GQ280592	GQ280714	GQ267314	
<i>Ca. malesiana</i>	CBS 112710	CPC 3899	GQ280473	AY725626	AY725759	AY725671	GQ280595	GQ280717	AY725721	Crous <i>et al.</i> 2004b
	CBS 112752 ^T	CMW 23687 = CPC 4223	GQ280472	AY725627	AY725760	AY725672	GQ280594	GQ280716	AY725722	
<i>Ca. mexicana</i>	CBS 110918 ^T	CMW 9055	GQ280474	AF210863	GQ267396	FJ972460	GQ280596	GQ280718	FJ972526	Crous 2002
<i>Ca. morgani</i>	CBS 110666	CMW 30978 = P90.1479 =	GQ280504	FJ918509	GQ267423	FJ918527	GQ280626	GQ280748	FJ9188557	Crous 2002
<i>Ca. multiphialidica</i>	CBS 112678	CMW 23688 =	GQ280475	AY725628	AY725761	AY725673	GQ280597	GQ280719	AY725723	Crous <i>et al.</i> 2004b
<i>Ca. multiseptata</i>	CBS 112682	CMW 23692 = CPC 1589	GQ280476	DQ190573	GQ267397	DQ190659	GQ280598	GQ280720	FJ918535	Crous 2002
<i>Ca. naviculata</i>	CBS 101121 ^T	CMW 30974 = INIFAT C98/19	GQ280478	GQ267211	GQ267399	GQ267252	GQ280600	GQ280722	GQ267317	Crous 2002
	CBS 116080	CMW 16723	GQ280477	AF333409	GQ267398	GQ267251	GQ280599	GQ280721	GQ267316	
<i>Ca. orientalis</i>	CBS 125258	CMW 20272	GQ280531	GQ267238	GQ267450	GQ267287	GQ280653	GQ280775	GQ267358	This study
	CBS 125259	CMW 20273	GQ280530	GQ267237	GQ267449	GQ267286	GQ280652	GQ280774	GQ267357	
	CBS 125260 ^T	CMW 20291	GQ280529	GQ267236	GQ267448	GQ267285	GQ267651	GQ280773	GQ267356	
<i>Ca. ovata</i>	CBS 111299	CMW 16724	GQ280479	GQ267212	GQ267400	GQ267253	GQ280601	GQ280723	GQ267318	Crous 2002
	CBS111307	CMW 30979	GQ280480	AF210868	GQ267401	GQ267254	GQ280602	GQ280724	GQ267319	
<i>Ca. pacifica</i>	CBS 109063	CMW 16726 = IMI 35428	GQ280481	GQ267213	AY725762	GQ267255	GQ280603	GQ280725	AY725724	Crous 2002
	CBS 114038	CMW 30988	GQ280482	AY725630	GQ267402	AY725675	GQ280604	GQ280726	GQ267320	



Table 1. (Continued)

Species	Isolate number ¹	Other collections ¹	GenBank accession nr. ²							Reference ³
			ACT	BT	CAL	HIS3	ITS	LSU	TEF-1 α	
<i>Ca. pauciramosa</i>	CMW 5683 ^T	CPC 971	GQ280486	FJ918514	GQ267405	FJ918531	GQ280608	GQ280730	FJ918565	Crous 2002
	CMW30823	CPC 416	GQ280485	FJ918515	GQ280404	FJ918532	GQ280607	GQ280729	FJ918566	
<i>Ca. penicilloides</i>	CBS 174.55 ^T	CMW 23696	GQ280487	AF333414	GQ267406	GQ267257	GQ280609	GQ280731	GQ267322	Crous 2002
<i>Ca. pini</i>	CBS 123698 ^T	CMW 31209	GQ280517	GQ267224	GQ267436	GQ267273	GQ280639	GQ280761	GQ267344	This study
	CBS 125523	CMW 31210	GQ280518	GQ267225	GQ267437	GQ267274	GQ280640	GQ280672	GQ267345	
<i>Ca. polizzii</i>	CBS 125270	CMW 7804	GQ280544	FJ972417	GQ267461	FJ972436	GQ280666	GQ280788	FJ972486	Lombard <i>et al.</i> 2009a
	CBS 125271	CMW 10151	GQ280545	FJ972418	GQ267462	FJ972437	GQ280667	GQ280789	FJ972487	
<i>Ca. pseudonaviculata</i>	CBS 11441 ^T	CMW 23672	GQ280490	GQ267214	GQ267409	GQ267258	GQ280612	GQ280734	GQ267325	Crous <i>et al.</i> 2002
<i>Ca. pseudoreteaudii</i>	CBS 123694 ^T	CMW 25310	GQ280492	FJ918504	GQ267411	FJ918519	GQ280614	GQ280736	FJ918541	Lombard <i>et al.</i> 2009c
	CBS 123696	CMW 25292	GQ280491	FJ918505	GQ267410	FJ918520	GQ280613	GQ280735	FJ918542	
<i>Ca. pseudoscoparia</i>	CBS 125254	CMW 15214	GQ280519	GQ267226	GQ267438	GQ267275	GQ280641	GQ280763	GQ267346	This study
	CBS 125255	CMW 15215	GQ280520	GQ267227	GQ267439	GQ267276	GQ280642	GQ280764	GQ267347	
	CBS 125256	CMW 15216	GQ280521	GQ267228	GQ267440	GQ267277	GQ280643	GQ280765	GQ267348	
	CBS 125257 ^T	CMW 15218	GQ280522	GQ267229	GQ267441	GQ267278	GQ280644	GQ280766	GQ267349	
<i>Ca. pseudospathiphylli</i>	CBS 109162 ^T	CMW 30976 = CPC 1623	GQ280493	FJ918513	GQ267412	AF348241	GQ280615	GQ280737	FJ918562	Crous 2002
<i>Ca. pteridis</i>	CBS 111793 ^T	CMW 16736 = CPC 2372 = ATCC 34395	GQ280494	DQ190578	GQ267413	DQ190679	GQ280616	GQ280738	FJ918563	Crous 2002
	CBS 111871	CMW 30982 = CPC 2443	GQ280495	DQ190579	GQ267414	DQ190680	GQ280617	GQ280739	FJ918564	
<i>Ca. pyrochoa</i>	CBS 749.70 ^T	CMW 23682	GQ280462	GQ267210	GQ267388	GQ267250	GQ280584	GQ280706	GQ267312	Crous <i>et al.</i> 2006



Table 1. (Continued)

Species	Isolate number ¹	Other collections ¹	GenBank accession nr. ²							Reference ³
			ACT	BT	CAL	HIS3	ITS	LSU	TEF-1 α	
<i>Ca. queenslandica</i>	CBS 112146 ^T	CMW 30604 = CPC 3213	GQ280496	AF389835	GQ267415	FJ918521	GQ280618	GQ280740	FJ918543	Lombard <i>et al.</i> 2009c
	CBS 112155	CMW 30603 = CPC 3210	GQ280497	AF389834	GQ267416	DQ190667	GQ280619	GQ280741	FJ918544	
<i>Ca. reteaudii</i>	CBS 112143	CMW 16738 = CPC 3200	GQ280499	GQ240642	GQ267418	DQ190660	GQ280621	GQ280743	FJ918536	Crous 2002
	CBS 112144 ^T	CMW 30984 = CPC 3201	GQ280498	AF389833	GQ267417	DQ190661	GQ280620	GQ280742	FJ918537	
<i>Ca. rumohrae</i>	CBS 109062	CMW 30989 = CPC 1603	GQ280501	AF232873	GQ267420	DQ190676	GQ280623	GQ280745	FJ918550	Crous 2002
	CBS 111431 ^T	CMW 23697 = CPC 1716	GQ285000	AF232871	GQ267419	DQ190675	GQ280622	GQ280744	FJ918549	
<i>Ca. scoparia</i>	CMW 31000	CPC 1675 = UFV 117	GQ280435	FJ972426	GQ267367	FJ972476	GQ280557	GQ280679	FJ972525	Crous 2002
	CMW 31001	UFV 126	GQ280436	FJ972427	GQ267368	GQ267246	GQ280558	GQ280680	GQ267246	
<i>Ca. spathiphylli</i>	CBS 114540	CMW 16742	GQ280505	AF348214	GQ267424	AF348230	GQ280627	GQ280749	GQ267330	Crous 2002
	CBS 116168	CMW 30997	GQ280506	FJ918512	GQ267425	FJ918530	GQ280628	GQ280750	FJ918561	
<i>Ca. spathulata</i>	CBS 555.92	CMW 16744	GQ280508	GQ267215	GQ267427	GQ267261	GQ280630	GQ280752	GQ267331	Crous 2002
	CBS 112689	CMW 16745	GQ280507	AF308463	GQ267426	FJ918524	GQ280629	GQ280751	FJ918554	
<i>Ca. sulawesiensis</i>	CBS 125248	CMW 14857	GQ280516	GQ267223	GQ267435	GQ267272	GQ280638	GQ280760	GQ267343	This study
	CBS 125253	CMW 14879	GQ280513	GQ267220	GQ267432	GQ267269	GQ280635	GQ280757	GQ267340	
	CBS 125277 ^T	CMW 14878	GQ280515	GQ267222	GQ267434	GQ267271	GQ280637	GQ280759	GQ267342	



Table 1. (Continued)

Species	Isolate number ¹	Other collections ¹	GenBank accession nr. ²							Reference ³
			ACT	BT	CAL	HIS3	ITS	LSU	TEF-1 α	
<i>Ca. sulawesiensis</i>	CMW 14883		GQ280514	GQ267221	GQ267433	GQ267270	GQ280636	GQ280758	GQ267341	
<i>Ca. sumatrensis</i>	CBS 112829 ^T	CMW 23698 = CPC4518	GQ280532	AY725649	AY725771	AY725696	GQ280654	GQ280776	AY725733	Crous <i>et al.</i> 2004b
	CBS 112934	CMW 30987 = CPC 4516	GQ280533	AY725651	AY725773	AY725798	GQ280655	GQ280777	AY725735	
<i>Ca. terrae-reginae</i>	CBS 112151 ^T	CMW 30601 = CPC 3202	GQ280534	FJ918506	GQ267451	FJ918522	GQ280656	GQ280778	FJ918545	Lombard <i>et al.</i> 2009b
	CBS 112634	CMW 30602 = CPC 4233	GQ280535	FJ918507	GQ267452	DQ190668	GQ280657	GQ280779	FJ918546	
<i>Ca. colombiana</i>	CBS 115127 ^T	CMW 30871 = CPC 1160	GQ280538	FJ972423	GQ267455	FJ972442	GQ280660	GQ280782	FJ972492	Lombard <i>et al.</i> 2009c
	CBS 115638	CMW 30766 = CPC 1161	GQ280539	FJ972422	GQ267456	FJ972441	GQ280661	GQ280783	FJ972491	
<i>Ca. variabilis</i>	CBS 112691	CMW 2914	GQ280541	GQ267240	GQ267458	GQ267264	GQ280663	GQ280785	GQ267335	Crous 2002
	CBS 114677	CMW 3187	GQ280541	AF333424	GQ267457	GQ267263	GQ280662	GQ267784	GQ267334	
<i>Ca. zuluensis</i>	CBS 125268	CMW 9188 ^T	GQ280542	FJ972414	GQ267459	FJ972433	GQ280664	GQ280786	FJ972483	Lombard <i>et al.</i> 2009c
	CBS 125272	CMW 9896	GQ280543	FJ972415	GQ267460	FJ972434	GQ280665	GQ280787	FJ972484	

¹ CBS: Centraalbureau voor Schimmelcultures, Utrecht, The Netherlands; CPC: Pedro Crous working collection housed at CBS; IMI: International Mycological Institute, CABI-Bioscience, Egham, Bakenham Lane, U.K.; ATCC: American Type Culture Collection, Virginia, U.S.A.; UFV: Universidade Federal de Vicosa, Brazil. ² ACT = Actin, BT = β -tubulin, CAL = Calmodulin, HIS3 = Histone H3, ITS = Internal transcribed spacer regions 1 and 2 and the 5.8S gene of the ribosomal RNA, LSU = 28S large subunit RNA, TEF-1 α = Translation elongation factor 1-alpha. ³ References used for species descriptions. ^T Ex-type cultures.

Table 2. Single nucleotide polymorphisms comparisons between *Ca. eucalypti* and *Ca. colhounii*.

Species	Isolate no.	β -tubulin				Histone H3					TEF-1 α								
		167	207	398	507	58	290	362	454	455	43	105	106	107	108	109	264	457	472
<i>Ca. colhounii</i>	CBS 293.79	A	G	A	C	A	A	C	A	C	C	A	C	A	A	C	G	C	C
	CBS 114704	A	G	A	C	A	A	C	A	C	C	A	C	A	A	C	G	C	C
<i>Ca. eucalypti</i>	CBS 125273	G	T	G	T	-	T	T	C	A	A	-	-	-	-	-	A	T	T
	CBS 125274	G	T	G	T	-	T	T	C	A	A	-	-	-	-	-	A	T	T
	CBS 125275	G	T	G	T	-	T	T	C	A	A	-	-	-	-	-	A	T	T
	CBS 125276	G	T	G	T	-	T	T	C	A	A	-	-	-	-	-	A	T	T
<i>Ca. macroconidialis</i>	CBS 114880	C	G	A	C	A	A	T	A	C	C	C	A	A	C	C	C	T	C
<i>Ca. madagascariensis</i>	CBS 114571	C	G	A	T	T	A	G	A	C	C	C	C	A	C	C	C	C	A
	CBS 114572	C	G	A	T	T	A	G	A	C	C	C	C	A	C	C	C	C	A



Table 3. Single nucleotide polymorphisms from the sequence datasets for *Ca. pini* and *Ca. orientalis* compared to *Ca. brachiatica* and *Ca. brassicae*.

Species	Isolate no.	β-tubulin								Histone H3										TEF-1α																					
		84	91	121	202	380	382	395	518	12	58	59	61	62	65	71	105	255	268	270	4	12	49	61	62	65	79	93	124	141	142	186	194	195	196	197	198	199	200	201	
<i>Ca. brachiatica</i>	CBS 123700	A	G	A	A	T	C	A	-	T	-	T	C	A	T	C	C	T	A	A	C	T	C	G	C	C	C	A	T	T	T	G	T	-	-	-	-	C	A	T	
	CMW 25302	A	G	A	A	T	C	A	-	T	-	T	C	A	T	C	C	T	A	A	C	T	C	G	C	C	C	A	T	T	T	G	T	-	-	-	-	C	A	T	
<i>Ca. brassicae</i>	CBS 111478	A	G	C	G	G	T	G	-	T	A	T	C	C	C	C	C	C	A	T	C	-	G	C	C	C	A	T	-	-	G	T	-	-	-	-	C	A	T		
	CBS 111869	A	G	C	G	G	T	G	-	T	A	T	C	C	C	C	C	C	A	T	C	-	G	C	C	C	A	T	-	-	G	T	-	-	-	-	C	A	T		
<i>Ca. pini</i>	CBS 123698	A	C	C	G	G	T	G	C	G	-	-	T	C	C	-	A	C	C	A	G	T	C	C	A	G	C	G	A	T	T	A	A	A	A	A	A	A	G	C	C
	CMW 31210	A	C	C	G	G	T	G	C	G	-	-	T	C	C	-	A	C	C	A	G	T	C	C	A	G	C	G	A	T	T	A	A	A	A	A	A	A	G	C	C
<i>Ca. orientalis</i>	CBS 125258	G	G	C	G	G	T	G	-	T	A	T	C	C	C	C	C	C	C	G	T	T	C	C	C	C	T	G	T	T	T	T	G	A	-	-	-	-	G	C	C
	CBS 125259	G	G	C	G	G	T	G	-	T	A	T	C	C	C	C	C	C	C	G	T	T	C	C	C	C	T	G	T	T	T	T	G	A	-	-	-	-	G	C	C
	CBS 125260	G	G	C	G	G	T	G	-	T	A	T	C	C	C	C	C	C	C	G	T	T	C	C	C	C	T	G	T	T	T	T	G	A	-	-	-	-	G	C	C

Table 3. (Continue)

Species	Isolate no.	TEF-1α											
		236	240	246	259	273	428	447	448	449	465	473	493
<i>Ca. brachiatica</i>	CBS 123700	C	T	G	T	C	G	C	C	A	G	G	-
	CMW 25302	C	T	G	T	C	G	C	C	A	G	G	-
<i>Ca. brassicae</i>	CBS 111478	C	T	G	T	C	G	C	C	A	C	G	C
	CBS 111869	C	T	G	T	C	G	C	C	A	C	G	C
<i>Ca. pini</i>	CBS 123698	T	A	C	C	C	A	-	-	-	G	A	C
	CMW 31210	T	A	C	C	C	A	-	-	-	G	A	C
<i>Ca. orientalis</i>	CBS 125258	T	A	C	C	T	A	-	-	-	G	G	C
	CBS 125259	T	A	C	C	T	A	-	-	-	G	G	C
	CBS 125260	T	A	C	C	T	A	-	-	-	G	G	C



Table 4. Single nucleotide polymorphisms comparisons between *Ca. scoparia* and *Ca. pseudoscoparia*.

Species	Isolate no.	β -tubulin			TEF-1 α		
		193	288	490			
<i>Ca. scoparia</i>	CMW 31000	T	-	-			
	CMW 31001	T	-	-			
<i>Ca. pauciramosa</i>	CMW 5683	T	-	-			
	CMW 30823	T	-	-			
<i>Ca. pseudoscoparia</i>	CBS 125254	C	C	C			
	CBS 125255	C	C	C			
	CBS 125256	C	C	C			
	CBS 125257	C	C	C			



Table 5. Single nucleotide polymorphisms from the sequence datasets for *Ca. densa* and *Ca. humicola* compared to *Ca. spathiphylli*.

Species	Isolate no.	β-tubulin												Histone H3																	
		8	74	103	151	193	220	225	234	235	241	388	393	515	524	527	71	83	101	103	105	127	209	253	256	261	262	266	279	459	460
<i>Ca. spathiphylli</i>	CBS 114540	A	T	G	C	C	G	C	T	T	C	T	G	C	T	A	C	C	A	C	T	T	T	T	A	C	T	T	T	A	C
	CBS 116168	A	T	G	C	C	G	C	T	T	C	T	G	C	T	A	C	C	A	C	T	T	T	T	A	C	T	T	T	A	C
<i>Ca. densa</i>	CBS 125249	A	-	A	G	A	A	T	C	C	T	T	C	T	C	T	T	G	G	T	C	C	C	C	G	T	C	C	C	A	C
	CBS 125250	A	-	A	G	A	A	T	C	C	T	T	C	T	C	T	T	G	G	T	C	C	C	C	G	T	C	C	C	A	C
	CBS 125261	A	-	A	G	A	A	T	C	C	T	T	C	T	C	T	T	G	G	T	C	C	C	C	G	T	C	C	C	A	C
<i>Ca. humicola</i>	CBS 125251	T	-	A	G	A	A	T	T	T	T	C	C	T	C	T	T	C	G	T	C	C	C	C	G	T	C	C	C	A	A
	CBS 125252	T	-	A	G	A	A	T	T	T	T	C	C	T	C	T	T	C	G	T	C	C	C	C	G	T	C	C	C	A	A
	CBS 125269	T	-	A	G	A	A	T	T	T	T	C	C	T	C	T	T	C	G	T	C	C	C	C	G	T	C	C	C	A	A

Table 5. (Continued)

Species	Isolate no.	TEF-1α													
		49	72	84	100	102	104	113	114	115	116	207	262	454	469
<i>Ca. spathiphylli</i>	CBS 114540	T	T	-	G	A	G	-	-	-	-	T	A	C	T
	CBS 116168	T	T	-	G	A	G	-	-	-	-	T	A	C	T
<i>Ca. densa</i>	CBS 125249	C	C	-	A	G	A	C	A	-	-	C	T	T	G
	CBS 125250	C	C	-	A	G	A	C	A	-	-	C	T	T	G
	CBS 125261	C	C	-	A	G	A	C	A	-	-	C	T	T	G
<i>Ca. humicola</i>	CBS 125251	C	C	C	A	G	A	C	A	C	A	T	A	T	G
	CBS 125252	C	C	C	A	G	A	C	A	C	A	T	A	T	G
	CBS 125269	C	C	C	A	G	A	C	A	C	A	T	A	T	G



Table 6. Single nucleotide polymorphisms comparisons between *Ca. brasiliensis*, *Ca. insularis* and *Ca. sulawesiensis*.

Species	Isolate no.	β -tubulin					Histone H3							TEF-1 α									
		117	360	395	472	509	95	100	253	259	260	390	417	452	98	100	103	104	105	109	143	263	439
<i>Ca. brasiliensis</i>	CBS 230.51	C	A	A	T	C	G	C	G	A	C	T	T	A	G	-	-	-	-	G	T	C	G
	CBS 114257	C	A	A	T	C	G	C	G	A	C	T	T	A	G	-	-	-	-	G	T	C	G
<i>Ca. cerciana</i>	CBS 123693	T	A	A	T	T	A	C	C	A	C	C	T	C	G	-	-	C	G	A	-	C	G
	CBS 123695	T	A	A	T	T	A	C	C	A	C	C	T	C	G	-	-	C	G	A	-	C	G
<i>Ca. insularis</i>	CBS 114558	T	G	A	C	C	A	C	G	A	C	C	C	A	G	C	A	C	A	A	-	C	A
	CBS 114559	T	G	A	C	C	A	C	G	A	C	C	C	A	G	C	A	C	A	A	-	C	A
<i>Ca. sulawesiensis</i>	CBS 125248	T	A	G	T	T	A	T	T	G	T	T	T	C	C	G	A	C	G	A	-	T	A
	CBS 125253	T	A	G	T	T	A	T	T	G	T	T	T	C	C	G	A	C	G	A	-	T	A
	CBS 125277	T	A	G	T	T	A	T	T	G	T	T	T	C	C	G	A	C	G	A	-	T	A
	CMW 14883	T	A	G	T	T	A	T	T	G	T	T	T	C	C	G	A	C	G	A	-	T	A



Table 7. Statistical information on the sequence dataset and maximum parsimony trees for each locus.

	ACT	BT	CAL	HIS3	ITS	LSU	TEF-1α
Aligned characters	290	532	531	499	706	887	596
Variable characters	15	42	39	62	32	10	57
Informative characters	151	268	323	223	112	37	337
Most parsimonious trees	2622	91	1000	372	1000	100	9970
Tree length	573	1454	1282	1843	296	91	1641
CI	0.490	0.431	0.467	0.352	0.618	0.538	0.477
RI	0.867	0.840	0.849	0.793	0.882	0.913	0.871
RC	0.425	0.569	0.397	0.648	0.545	0.492	0.416

Fig. 1. The most parsimonious trees obtained from a heuristic search with 1000 random additions of the combined BT, HIS3 and TEF-1 α sequence alignments of the *Ca. colhounii* complex. Scale bar shows 10 changes and bootstrap support values (bold) from 1000 replicates and Bayesian posterior probability values are indicated at the nodes. Red lines indicate bootstrap values of 100 and posterior probabilities of 1.00. Thickened lines indicate branches in the strict consensus tree and the consensus tree of the Bayesian analyses.




Fig. 2. The most parsimonious trees obtained from a heuristic search with 1000 random additions of the combined BT, HIS3 and TEF-1 α sequence alignments of the *Calonectria brassicae* complex. Scale bar shows 10 changes and bootstrap support values (bold) from 1000 replicates and Bayesian posterior probability values are indicated at the nodes. Red lines indicate bootstrap values of 100 and posterior probabilities of 1.00. Thickened lines indicate branches in the strict consensus tree and the consensus tree of the Bayesian analyses.




Fig. 3. One of two most parsimonious trees obtained from a heuristic search with 1000 random additions of the combined BT, HIS3 and TEF-1 α sequence alignments of the *Calonectria scoparia* complex. Scale bar shows 10 changes and bootstrap support values (bold) from 1000 replicates and Bayesian posterior probability values are indicated at the nodes. Red lines indicate bootstrap values of 100 and posterior probabilities of 1.00. Thickened lines indicate branches in the strict consensus tree and the consensus tree of the Bayesian analyses.




Fig. 4. One of three most parsimonious trees obtained from a heuristic search with 1000 random additions of the combined BT, HIS3 and TEF-1 α sequence alignments of the *Calonectria morganii* complex. Scale bar shows 10 changes and bootstrap support values (bold) from 1000 replicates and Bayesian posterior probability values are indicated at the nodes. Red lines indicate bootstrap values of 100 and posterior probabilities of 1.00. Thickened lines indicate branches in the strict consensus tree and the consensus tree of the Bayesian analyses.




Fig. 5. One of 24 most parsimonious trees obtained from a heuristic search with 1000 random additions of the combined actin, β -tubulin, calmodulin, histone H3 and translation elongation factor 1-alpha sequence alignments of the *Calonectria* spp. Scale bar shows 10 changes and bootstrap support values (bold) from 1000 replicates and Bayesian posterior probability values are indicated at the nodes. Thickened lines indicate branches in the strict consensus tree and the consensus tree of the Bayesian analyses. Red lines indicate bootstrap values of 100 and posterior probabilities of 1.00. The tree was rooted to *C. lageniformis* (CBS 112898) and *C. peruviana* (CPC 5614). Phylogenetic groups are indicated on the right.




Fig 6. Results of sexual compatibility tests. Successful matings are indicated by (+) and unsuccessful matings is indicated with (-). Blue highlighted blocks indicate homothallic matings. Yellow blocks highlight unsuccessful self-self matings. Green blocks indicate mating tester strain matings. A. Matings between isolates of *Ca. macroconidialis* and *Ca. eucalypti*. B. Matings between isolates of *Ca. brachiatica*, *Ca. brassicae*, *Ca. pini* and *Ca. orientalis*. C. Matings between isolates of *Ca. brasiliensis*, *Ca. cerciana*, *Ca. insularis* and *Ca. sulawesiensis*. D. Matings between isolates of *Ca. densa*, *Ca. humicola* and *Ca. spathiphylli*. E. Matings between isolates of *Ca. pauciramosa*, *Ca. pseudoscoparia* and *Ca. scoparia*.



Fig. 7. *Calonectria densa*. A–B. Macroconidiophore of *Ca. densa*. B. Macroconidiophore with lateral stipe extensions. C–D. Fertile branches of macroconidiophore with doliiform to reniform phialides. E–F. Vesicles. G–H. Macroconidia. Scale bars in A–B = 20 μm , C–H = 10 μm .




Fig. 8. *Calonectria eucalypti*. A–F. Teleomorph state of *Ca. eucalypti*. G–L. Anamorph state of *Ca. eucalypti*. A–B. A 10 μm thick vertical section through perithecium. C. Section through perithecium wall. D. Ostiolar region of a perithecium. E. Asci and ascospores F. Ascospores. G–H. Macroconidiophores of *Ca. eucalypti*. I–J. Vesicles. K. Fertile branches of a conidiophore with doliiform to reniiform phialides. L. Macroconidia. Scale bars: A–B = 50 μm , G–H = 20 μm , C–F, I–L = 10 μm .






Fig. 9. *Calonectria humicola*. A–B. Macroconidiophore of *Ca. humicola*. C–D. Fertile branches of macroconidiophore with doliiform to reniform phialides. E–F. Vesicles. G–H. Macroconidia. Scale bars in A–B = 20 μm , C–H = 10 μm .






Fig. 10. *Calonectria orientalis*. A–B. Macroconidiophore of *Ca. orientalis*. C–D. Fertile branches of macroconidiophore with doliiform to reniform phialides. E–F. Vesicles. G–H. Macroconidia. Scale bars in A–B = 20 μm , C–H = 10 μm .






Fig. 11. *Calonectria pini*. A–B. Macroconidiophore of *Ca. pini*. C–D. Fertile branches of macroconidiophore with doliiform to reniform phialides. E–F. Vesicles. G–H. Macroconidia. Scale bars in A–B = 20 μm , C–H = 10 μm .






Fig. 12. *Calonectria pseudoscoparia*. A–B. Macroconidiophore of *Ca. pseudoscoparia*. C–D. Fertile branches of macroconidiophore with doliiform to reniform phialides. E–F. Vesicles. G–H. Macroconidia. Scale bars in A–B = 20 μm , C–H = 10 μm .






Fig. 13. *Calonectria sulawesiensis*. A–B. Macroconidiophore of *Ca. sulawesiensis*. C–D. Fertile branches of macroconidiophore with doliiform to reniform phialides. E–F. Vesicles. G–H. Macroconidia. Scale bars in A–B = 20 μm , C–H = 10 μm .

