

REFERENCE LIST

- Albert, M.J., Bhuiyan, N.A., Talukder, K.A., Faruque, A.S., Nahar, S., Faruque, S.M., Ansaruzzaman, M. and Rahman, M. 1997. Phenotypic and genotypic changes in *Vibrio cholerae* O139 Bengal. *J. Clin. Microbiol.* 45: 2588 – 92.
- Albert, M.J., Islam, D., Nahar, S., Qadri, F., Falklind, S. and Weintraub, A. 1997. Rapid detection of *Vibrio cholerae* O139 Bengal from stool specimens by polymerase chain reaction. *J. Clin. Microbiol.* 35: 1633 – 1635
- Barrett, T.J. and Blake, P.A. 1981. Epidemiological usefulness of changes in hemolytic activity of *Vibrio cholerae* biotype El Tor during the Seventh Pandemic. *J. Clin. Microbiol.* 13: 126 – 129.
- Baumann, P., Furniss, A.L. and Lee, J.V. 1984. Genus 1. *Vibrio*. In: Bergey's manual of systematic bacteriology Vol. 1. Ed. N.R. Krieg, J.G. Holt, R.G.E. Murray, D.J. Brenner, M.P. Bryant, J.W. Moulder, N. Pfennig, P.H.A Sneath, and J.T. Staley. London: Williams and Wilkins. p518 – 538.
- Beltrán, P., Delgado, G., Navarro, A., Trujillo, F., Selander, R.K. and Cravioto A. 1999. Genetic diversity and population structure of *Vibrio cholerae*. *J. Clin. Microbiol.* 37: 581 – 590.
- Bennish, M.L. 1994 Cholera: Pathophysiology, clinical features and treatment. In: *Vibrio cholerae and cholera – molecular to global perspectives*, edited by I.K. Wachsmuth, P.A. Blake and O. Olsvik. Washington: ASM press: 229 – 256.
- Bik, E.M., Gouw, R.D. and Mooi, F.R. 1996. DNA fingerprinting of *Vibrio cholerae* strains with a novel insertion sequence element: a tool to identify epidemic strains. *J. Clin. Microbiol.* 34: 1453 – 1461.

- Blake, P.A. 1994. Historical perspectives on pandemic cholera. In: *Vibrio cholerae and cholera – molecular to global perspectives*, edited by I.K. Wachsmuth, P.A. Blake and O. Olsvik. Washington: ASM press: 293 – 295.
- Byun, R., Elbourne, L.D., Lan, R. and Reeves, P.R. 1999. Evolutionary relationships of pathogenic clones of *Vibrio cholerae* by sequence analysis of four housekeeping genes. *Infect. Immun.* 67: 1116 – 1124.
- Cameron, D.N., Khambarty, F.M., Wachsmuth, I.K., Tauxe, R. and Barrett, T.J. 1994. Molecular characterization of *Vibrio cholerae* O1 strains by pulsed field gel electrophoreses. *J. Clin. Microbiol.* 32: 1685 – 1690.
- Chakraborty, S., Mukhopadhyay, A.K., Bhadra, R.K, Ghosh, A.N., Mitra, R., Shimada, T., Yamasaki, S., Faruque, S.M, Takeda, Y., Colwell R.R. and Nair G.B. 2000. Virulence genes in environmental strains of *Vibrio cholerae*. *Apl. Env. Microbiol.* 66: 4022 – 4028.
- Chattopadhyay, D.J., Sarkar, B.L., Ansari, M.Q., Chakrabarti, B.K., Roy, M.K., Ghosh, A.N. and Pal, S.C. 1993. New phage typing scheme for *Vibrio cholerae* O1 biotype El Tor strains. *J. Clin. Microbiol.* 31: 1579 – 1585.
- Chen, F., Evins, G.M., Cook, W.L., Almeida, R., Hargrett-Bean, N. and Wachsmuth, K. 1991. Genetic diversity among toxigenic and nontoxigenic *Vibrio cholerae* O1 isolated from the Western Hemisphere. *Epidemiol. Infect.* 107: 225 – 233.
- Choudhury, S.R., Bhadra, R.K. and Das, J. 1994. Genome size and restriction fragment length polymorphism analysis of *Vibrio cholerae* strains belonging to different serovars and biotypes. *FEMS Microbiol. Lett.* 115: 329 – 334.

- Chow, K.H., Ng, T.K., Yuen, K.Y. and Yam, C. 2001. Detection of RTX toxin gene in *Vibrio cholerae* by polymerase chain reaction. *J. Clin. Microbiol.* 39: 2594 – 2597.
- Coelho, A., Andrade, J. R., Vicente, A. C. and Salles, C. A. 1995. New variant of *Vibrio cholerae* O1 from clinical isolates in Amazonia. *J. Clin. Microbiol.* 33:114 – 118
- Colwell, R.R. and Huq, A. 1994. *Vibrios* in the environment: Viable but nonculturable *Vibrio cholerae*. In: *Vibrio cholerae and cholera – molecular to global perspectives*, edited by I.K. Wachsmuth, P.A. Blake and O. Olsvik. Washington: ASM press: 117 – 134.
- Colwell, R.R., Hasan, J.A.K., Huq, A., Iloomis, L., Seibeling, R.J., Torres, M., Galvez, S., Islam, S. and Bernstein, D. 1992. Development and evaluation of a rapid, simple, and sensitive monoclonal antibody based co-agglutination test for the direct detection of *Vibrio cholerae* O1. *FEMS Microbiol. Lett.* 97: 215 – 220.
- Comstock, L.E., Johnson, J.A., Michalski, J.M., Morris, J.G.(Jr.) and Kaper, J.B. 1996. Cloning and sequence of a region encoding a surface polysaccharide of *Vibrio cholerae* O139, and characterization of the insertion site in the chromosome of *Vibrio cholerae* O1. *Mol. Microbiol.* 19: 815 – 826.
- Dalsgaard, A., Echeverri, P., Larsen, J.L., Siebeling, R., Serichanelergs, O. and Huss, H.H. 1995. Application of ribotyping for differentiating *Vibrio cholerae* non-O1 isolated from shrimp farms in Thailand. *Appl. Env. Microbiol.* 61: 245 – 251.
- Dalsgaard, A., Forslund, A. and Fussing, V. 1999. Traditional ribotyping shows a higher discrimination than the automated Riboprinter™ system in typing *Vibrio cholerae* O1. *Lett. Appl. Microbiol.* 28: 327 – 333.

- Department of Health. 2004. Cholera. Internet: www.doh.gov.za/cholera.html. Access: 18 October.
- Falklind, S., Stark, M., Albert, M.J., Uhlen, M., Lundberg, J. and Weintraub, A. 1996. Cloning and sequencing of a region of *Vibrio cholerae* O139 Bengal and its use in a polymerase chain reaction based detection. *J. Clin. Microbiol.* 34: 2904 – 2908.
- Farmer III, J.J. and Hickman-Brenner, F.W. 1992. The genera *Vibrio* and *Photobacterium*. In: *The Prokaryotes* (Second edition). Ed. A. Balows, H.G. Truper, M. Dworken, W. Harder and K.-H. Schleifer. New York: Springer-Verlag.
- Faruque, S.M., Asadulghani, Saha, M.N., Abdul Alim, A.R.M., Albert M.J., Nasirul Islam, K.M. and Mekalanos J.J. 1998. Analysis of clinical and environmental strains of non-toxicogenic *Vibrio cholerae* for susceptibility to CTX ϕ : Molecular basis for origination of new strains with epidemic potential. *Infect. Immun.* 66: 5819 – 5825.
- Fassano, A., Baudry, B., Pumplin, D.W., Wasserman, S.S., Tall, B.D., Kelly, J.M. and Kaper, J.B. 1991. *Vibrio cholerae* produces a second enterotoxin which affects intestinal tight junctions. *Proc. Natl. Acad. Sci. USA* 88:5242 – 5246.
- Fields, P.I., Popovic, T., Wachsmuth, K. and Olsvik, O. 1992. Use of the polymerase chain reaction for detection of toxigenic *Vibrio cholerae* O1 strains from the Latin American cholera epidemic. *J. Clin. Microbiol.* 30: 2118-2121.
- Finkelstein, R.A. and LoSpalluto, J.J. 1969. Pathogenesis of experimental cholera: preparation and isolation cholera toxin and cholera toxinogen. *J. Exp. Med.* 130:185 – 202.

- Gustafsson, B. and Holme, T. 1985. Rapid detection of *Vibrio cholerae* O1 by motility inhibition and immunofluorescence with monoclonal antibodies. *Eur. J. Clin. Microbiol.* 4: 291 - 294.
- Halpern, M., Broza, Y.B., Mittler, S., Arakawa, E. and Broza, M. 2004. Chironomid egg masses as a natural reservoir of *Vibrio cholerae* non-O1 and non-O139 in fresh water habitats. *Microbial Ecol.* 47: 341 – 349.
- Holmes, B., Costas, M., Ganner, M., On, S.L.W. and Stevens, M. 1994. Evaluation of Biolog system for identification of some Gram-negative bacteria of clinical importance. *J. Clin. Microbiol.* 32:1970 – 1975.
- Holmes, B., Willcox, W.R. and Lapage, S.P. 1978. Identification of *Enterobacteriaceae* by the API 20 E system. *J. Clin. Pathol.* 31: 22 – 30.
- Honda, T. and Finkelstein, R.A. 1979 Purification and characterization of a hemolysin produced by *Vibrio cholerae* biotype El Tor, another toxic substance produced by cholera *Vibrios*. *Infect. Immun.* 26:1020 – 1027.
- Honda, T., Nishibuchi, M., Miwatani, T. and Kaper, J.B. 1986. Demonstration of a plasmid-borne gene encoding a thermostable direct hemolysin in *Vibrio cholerae* non-O1 strains. *Appl. Env. Microbiol.* 52: 1218 – 1220.
- Hoshino, K., Yamasaki, S., Mukhopadhyay, A.K., Chakraborty, S., Basu, A., Bhattacharya, S.K., Nair, G.B., Shimada, T. and Takeda, Y. 1998. development and evaluation of a multiplex PCR assay for the rapid detection of toxigenic *Vibrio cholerae* O1 and O139. *FEMS Immun. Med. Microbiol.* 20: 201 – 207.

- Huq, A., Colwell, R.R., Chowdhury, M.A., Xu, B., Moniruzzaman, S.M., Islam, M.S., Yunus, M. and Albert, M.J. Coexistence of *Vibrio cholerae* O1 and O139 Bengal in plankton in Bangladesh. 1995. *Lancet* 345: 1249
- Huq, A., Colwell, R.R., Rahman, R., Ali, A., Chowdhury, M.A.R., Parveen, S., Sack, D.A. and Russek-Cohen, E. 1990. Detection of *Vibrio cholerae* O1 in the aquatic environment by fluorescent-monoclonal antibody and culture method. *Appl. Env. Microbiol.* 56: 2370 – 2373.
- Huq, A., Small, E.B., West, P.A., Huq, I., Rahman, R. and Colwell R.R. 1983. Ecological relationship between *Vibrio cholerae* and planktonic crustacean copepods. *Appl. Env. Microbiol.* 45: 275 – 283.
- Huq, A., West, P.A., Small, E.B., Huq, M.I. and Colwell, R.R. 1984. Influence of water temperature, Salinity, and pH on the survival and growth of toxigenic *Vibrio cholerae* serovar O1 associated with live copepods in laboratory microcosms. *Appl. Env. Microbiol.* 48: 420 – 424.
- Ichinose, Y. Yamamoto, K., Nakasone, N., Tanabe, M.J., Takeda T., Miwatane, T. and Iwanaga, M. 1987. Enterotoxicity of El Tor-like hemolysin of non-O1 *Vibrio cholerae*. *Infect. Immun.* 55:1090 – 1093.
- Islam, M.S. 1990. Increased toxin production by *Vibrio cholerae* O1 during survival with a green algae, *Rhizodinium fontanum*, in an artificial aquatic environment. *Microbiol. Immunol.* 34: 557 – 563.
- Janssen, P., Coopman, R., Huys, G., Swings, J., Bleeker, M., Vos, P., Zabeau, M., Kersters, K. 1996. Evaluation of the DNA fingerprinting method AFLP as a new tool in bacterial taxonomy. *Biology*: 142: 1881 – 1893.

- Jesudason, M.V., Thangavelu, C.P. and Lalitha, M.K. 1984. Rapid screening of fecal samples for *Vibrio cholerae* by a coagglutination technique. *J. Clin. Microbiol.* 19: 712 – 713.
- Jiang, S.C., Louis, V., Choopun, N., Sharma, A., Huq, A. and Colwell, R.R. 2000(a). Genetic diversity of *Vibrio cholerae* in Chesapeake Bay determined by amplified fragment length polymorphism fingerprinting. *Appl. Env. Microbiol.* 66:140 – 147.
- Jiang, S.C., Matte, M., Matte, G., Huq, A., Colwell, R.R. 2000(b). Genetic diversity of clinical and environmental isolates of *Vibrio cholerae* determined by amplified fragment length polymorphism fingerprinting. *Appl. Env. Microbiol.* 66: 148 – 153.
- Kaper, J.B., Bradford, H.B., Roberts, N.C. and Falkow, S. 1982. Molecular epidemiology of *Vibrio cholerae* in the U.S. gulf coast. *J. Clin. Microbiol.* 16: 129 – 134.
- Kaper, J.B., Fasano, A. and Trucksis, M. 1994. Toxins of *Vibrio cholerae*. In: *Vibrio cholerae and cholera – molecular to global perspectives*, edited by I.K. Wachsmuth, P.A. Blake and O. Olsvik. Washington: ASM press: 145 – 176.
- Kapley, A., Hermant, T. and Purohit, J. 2001. Detection of the etiological agent for cholera by PCR protocol. *Med. Sci. Monit.* 2001. 7(2): 242 – 245.
- Karaolis, D.K., Johnson, J.A., Bailey, C.C., Boedeker, E.C., Kaper, J.B. and Reeves, P.R. 1998. A *Vibrio cholerae* pathogenicity island associated with epidemic and pandemic strains. *Proc Natl Acad Sci USA* 17;95: 3134-9

- Karunasagar, I., Sugumar, G., Karunasagar, I. and Reilly A. 1995. Rapid detection of *Vibrio cholerae* contamination of seafood by polymerase chain reaction. *Mol. Mar. Biol. Biotechnol.* 4(4): 365 – 368.
- Kay, B.A., Bopp, C.A. and Wells, J.G. 1994. Isolation and identification of *Vibrio cholerae* O1 from fecal specimens. In: *Vibrio cholerae and cholera – molecular to global perspectives*, edited by I.K. Wachsmuth, P.A. Blake and O. Olsvik. Washington: ASM press: 3 – 25.
- Kobayashi, K., Set, K., Akasa, S. and Makano, M. 1990. Detection of toxigenic *Vibrio cholerae* O1 using polymerase chain reaction for amplifying the cholera enterotoxin gene. *J. Jpn. Assoc. Infect. Dis.* 64:1323 – 1329.
- Koch, W.H., Payne, W.L., Wentz, B.A. and Cebula T.A. 1993. rapid polymerase chain reaction method for detection of *Vibrio cholerae* in foods. *Appl. Env. Microbiol.* 59: 556-560.
- Kotetishvili, M., Stine, O.C., Chen, Y., Kreger, A., Sulakvelidze, Sozhamannan, S. and Morris, J.G. (Jr.) 2003. Multilocus sequence typing has better discriminatory ability for typing *Vibrio cholerae* than does pulsed field gel electrophoreses and provides a measure of phylogenetic relatedness. *J. Clin. Microbiol.* 41: 2191 – 2196.
- KZN Health. 2004. Cholera review. Internet: www.kxnhealth.gov.za/cholerareview.pdf Access: 18 October.
- Lan, R. and Reeves, P.R. 2002. Pandemic spread of cholera: Genetic diversity and relationships within the seventh pandemic clone of *Vibrio cholerae* determined by amplified fragment length polymorphism. *J. Clin. Microbiol.* 40: 172 – 181.

- Levine, M.M., Kaper, J.B, Herrington, D., Losonsky, G., Morris, J.G., Clements, M.L., Black, R.E., Tall, B. and Hall, R. 1998. Volunteer studies of deletion mutants of *Vibrio cholerae* O1 prepared by recombinant techniques. *Infect. Immun.* 56: 161 – 167.
- Ling, T.K.W., Liu, Z.K. and Cheng A.F.B. 2003. Evaluation of the VITEK 2 system for rapid identification and susceptibility testing of Gram-negative bacilli from positive blood cultures. *J. Clin. Microbiol.* 41: 4705 – 4707.
- Lyon, W.J. 2001. Taqman PCR for the detection of *Vibrio cholerae* O1, O139, Non-O1, and Non-O139 in pure cultures, raw oysters and synthetic seawater. *Appl. Env. Microbiol.* 67: 4685 – 4693.
- Maiden, M.C., Bygraves, J.A., Feil, E., Morelli, G., Russel, J.E., Urwin, R., Zhang, Q., Zou, J., Zurth, K., Caugant, D.A., Feavers, I.M., Achtman, M. and Spratt, B.G. 1998. Multilocus sequence typing: a portable approach to the identification of clones within populations of pathogenic organisms. *Proc. Natl. Acad. Sci. USA* 95: 3140 – 3145
- Martinez-Govea, A., Ambrosio, J., Gutierrez-Cogco, L. and Flisser A. 2001. Identification and strain differentiation of *Vibrio cholerae* by using antibodies against outer membrane proteins. *Clin. Diag. Lab. Immun.* 8: 768-771.
- Miller, J. and Rhoden, D.L. 1991. Preliminary evaluation of Biolog, a carbon source utilization method for bacterial identification. *J. Clin. Microbiol.* 29: 1143 – 1147.
- Mosley, W.H. and Khan M. 1979. Cholera epidemiology – some environmental aspects. *Prog. Wat. Tech.* 11: 309 – 316.

- Mukhopadhyay, A.K., Chakraborty, S., Takeda, Y., Nair, G.B. and Berg, D.E. 2001. Characterization of VPI pathogenicity island and CTX ϕ prophage in environmental strains of *Vibrio cholerae*. *J. Bacteriol.* 183: 4734 – 4746.
- Nandi, B., Nandy, R.K., mukhopadhyay, S., Nair, G.B., Shimada, T. and Ghose, A.C. 2000. Rapid method for species-specific identification of *Vibrio cholerae* using primers targeted to the gene of the Outer Membrane Protein ompW. *J. Clin. Microbiol.* 38: 4145 – 4151.
- O'Hara, C.M., Rhoden, D.L. and Miller, J.M. 1992. Reevaluation of the API 20E identification system versus conventional biochemicals for identification of members of the family *Enterobacteriaceae*: a new look at an old project. *J. Clin. Microbiol.* 30: 123 – 125.
- Ogawa, A., Kato, J.-I., Watanabe, H., Nair, B.G. and Takeda, T. 1990. Cloning and Nucleotide sequence of a heat-stable enterotoxin gene from *Vibrio cholerae* non-O1 isolated from a patient with traveler's diarrhea. *Infect. Immun.* 58: 3325 – 3329.
- Olsvik, Ø., Whalberg, J., Petterson, B., Uhlen, M., Popovic, T., Wachsmuth, I.K. and Fields, P.I. 1993. Use of automated sequencing of polymerase chain reaction-generated amplicons to identify three types of Cholera Toxin subunit B in *Vibrio cholerae* O1 strains. *J. Clin. Microbiol.* 31: 22 – 25.
- Park, T.S., Ohl, S.L., Lee, E.Y., Park, K.H., Fiqueras, M.J. and chang, C.L. 2003. Misidentification of *Aeromonas veronii* biovar *sobria* as *Vibrio alginolyticus* by the VITEK system. *Lett. Appl. Microbiol.* 37: 349 – 353.
- Pearson G.D.N., Woods, A., Chiang S.L. and Mekalanos J.J. 1993. CTX genetic element encodes a site-specific recombination system and an intestinal colonization factor. *Proc. Natl. Acad. Sci. USA* 90:3750 – 3754.

- Pichel, M., Rivas, M., Chinen, I., Martín, F., Ibarra, C. and Binsztein, N. 2003. Genetic diversity of *Vibrio cholerae* O1 in Argentina and emergence of a new variant. *J. Clin. Microbiol.* 41: 124 – 134.
- Popovic, T., Bopp, C., Olsvik, Ø. and Wachsmuth, K. 1993. Epidemiological application of a standardized ribotype scheme for *Vibrio cholerae* O1. *J. Clin. Microbiol.* 31: 2474 – 2482.
- Rahman, m., Sack, P.A., Mahmood, S., and Hossain, A. 1987. Rapid diagnosis of cholera by coagglutination test using 4-hour fecal enrichment cultures. *J. Clin. Microbiol.* 25: 2204 – 2206.
- Roszak, D.B. and Colwell, R.R. 1987. Survival strategies of bacteria in the natural environment. *Microbiol. Rev.* 51: 365 – 379.
- Rutherford, I., Moody, V., Gavan, T.L., Ayers, L.W. Taylor, D.L. 1977. Comparative study of three methods for identification of *Enterobacteriaceae*. *J. Clin. Microbiol.* 5: 458 – 464.
- Saiki, R.K., Gelfand, D.H., Stoffel S., Scharf, S.J., Higuchi, R., Horn, G.T., Mullis, K.B. and Erlich, H.A. 1988. Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science* 239: 487 – 491
- Sanyal, S.C., Alam, K., Neogi, P.K.B., Huq, M.I. and Al-Mahmud, K.A. 1983. A new cholera toxin. *Lancet* I: 1337.
- Shaffer, N., do Santos, E.S., Andreason, P.-A. and Farmer J.J. III. 1989. Rapid laboratory diagnosis of cholera in the field. *Trans. R. Soc. Trop. Med. Hyg.* 83: 119 – 120.

- Shangkuan, Y.H., Show, Y.S. and Wang, T.M. 1995. Multiplex polymerase chain reaction to detect toxigenic *Vibrio cholerae* and to biotype *Vibrio cholerae* O1. *J. Appl. bacteriol.* 79(3): 264-273.
- Sharma, C., Ghosh, A., Dalsgaard, A., Forslund, A., Ghosh, R. K., Bhattacharya, S. K. and Nair, G. B. 1998. Molecular evidence that a distinct *Vibrio cholerae* O1 biotype El Tor strain in Calcutta may have spread to the African continent. *J. Clin. Microbiol.* 36:843 – 844
- Shirai, H., Nishibuchi, M., Ramamurthy, T., Bhattacharya, S.K., Pal, S.C. and Takeda, Y. 1991. Polymerase chain reaction for the detection of the cholera enterotoxin operon of *Vibrio cholerae*. *J. Clin. Microbiol.* 29: 2517 – 2521.
- Singh, D.V., Isac, S.R. and Colwell, R.R. 2002. development of a hexaplex PCR assay for rapid detection of virulence and regulatory genes in *Vibrio cholerae* and *Vibrio mimicus*. *J. Clin. Microbiol.* 40: 4321 – 4324.
- Singh, D.V., Matte, M.H., Matte, G.R., Jiang, S., Sabeena, F., Shukla, B.N., Sanyal, S.C., Huq, A. and Colwell, R.R. 2001 Molecular analysis of *Vibrio cholerae* O1, O139, non-O1, and non-O139 strains: Clonal relationships between clinical and environmental isolates. *Apl. Env. Microbiol* 67:910 – 921.
- Spanu, T., Sanguinetti, M., Ciccaglione, D., D’Inzeo, T., Romano, L., Leone, F. and Fadda, G. 2003. Use of the VITEK 2 system for the rapid identification of *Staphylococci* from bloodstream infections. *J. Clin. Microbiol.* 41: 4259 – 4263.
- Sprangler, B.D. 1992 Structure and function of cholera toxin and the related *Escherichia coli* heat-labile enterotoxin. *Microbiol. rev.* 56: 622 – 647.
- Stroehler, U.H., Jedani, K.E., Dredge, B.K., Morana, R., Brown, M.H., Karageorges, L.E., Albert, M.J. and Manning, P.A. 1995. Genetic rearrangements

- in the rfb regions of *Vibrio cholerae* O1 and O139. *Proc. Natl. Acad. Sci. USA*. 92: 10374 – 10378.
- Tamplin, M.L., Colwell, R.R., Hall, S., Koqure, K. and Strichartz, G.R. 1987. Sodium channel inhibitors produced by enteropathogenic *Vibrio cholerae* and *Aeromonas hydrophila*. *Lancet* I: 975.
 - Tenover, F. C., Arbeit, R. D., Goering, R.V., Mickelsen, P. A., Murray, B. E., Persing, D. H. and Swaminathan, B. 1995. Interpreting chromosomal DNA restriction patterns produced by pulsed-field gel electrophoresis: criteria for bacterial strain typing. *J. Clin. Microbiol.* 33:2233 – 2239
 - Thal, L.A., Silverman, J., Donabedian, S. and Zervers M.J. 1997. The effect of *Tn 916* insertion on contour-clamped homogeneous electrophoresis patterns of *Enterococcus faecalis*. *J. Clin. Microbiol.* 34: 969 – 972
 - Theron, J., Cilliers, J., Du Preez, M., Brözel, V.S. and Venter.S.N. 2000. Detection of toxigenic *Vibrio cholerae* from environmental water samples by an enrichment broth cultivation-pit-stop semi-nested PCR procedure. *J. Appl. Microbiol.* 89: 539 – 546.
 - Thompson, F.L., Iida, T. and Swings J. 2004. Biodiversity of Vibrios. *Microbiol. Mol. Biol. Rev.* 68:403 – 431
 - Thompson, F.L., Thompson, C.C., Vicente, A.C.P., Theophilo, G.N.D., Hofer, E. and Swings, J. 2003. Genomic diversity of clinical and environmental *Vibrio cholerae* strains isolated in Brazil between 1991 and 2001 as revealed by fluorescent amplified fragment length polymorphism analysis. *J. Clin. Microbiol.* 41: 1946 – 1950.

- Trucksis, M., Galen, J.E., Michalski, J., Fasano, A. and Kaper, J.B. 1993. Accessory cholera enterotoxin (Ace), the third toxin of a *Vibrio cholerae* virulence cassette. *Proc. Natl. Acad. Sci. USA* 90: 5267 – 5271.
- Vos, P., Hogers, R., Bleeker, M., Reijans, M., van de Lee, T., Hornes, M., Frijters, A., Pot, J., Peleman, J., Kuiper, M. and Zabeau, M. 1995. AFLP: a new technique for DNA fingerprinting. *Nucleic Acids Research*. 23(21): 4407 – 4414.
- Wachsmuth, I.K., Bopp, C.A., Fields, P.I. and Camillo, C. 1991. Difference between toxigenic *Vibrio cholerae* O1 from South America and US gulf coast. *Lancet*. 4;337(8749):1097-8.
- Wachsmuth, I.K., Evins, G.M., Fields, P.I., Olsvik, Ø., Popovic, T., Bopp, C.A., Wells, J.G., Carrillo, C. and Blake, P.A. 1993. The molecular epidemiology of cholera in Latin America. *J. Infect. Dis.* 167: 621 – 626.
- Waldor, M.K. and Mekalanos, J.J. 1996 Lysogenic conversion by a filamentous phage encoding cholera toxin. *Science* 272: 1910-1914.
- Weisburg, W.G., Barns, S.M., Pelletier, D.A. and Lane, D.J. 1991. 16S Ribosomal DNA amplification for Phylogenetic study. *J Bacteriol* 173: 697 – 703
- World Health Organization. 2003. *Weekly epidemiological record* 31:269 - 276
- Xu, H.S., Robert, N., Singleton, F.L., Attwell, R.W., Grimes, D.J. and Colwell R.R. 1982. Survival and viability of non-culturable *E. coli* and *Vibrio cholerae* in the estuarine and marine environment. *Microbiol Ecol.* 8: 213 – 223

- Yam, W.C., Lung, M.L. and Ng, M.H. 1991. Restriction fragment length polymorphism analysis of *Vibrio cholerae* strains associated with a cholera outbreak in Hong Kong. *J. Clin. Microbiol.* 29: 1058 – 1059
- Yamai, S., Okitsu, T. and Katsube, Y. 1997. Distribution of serogroups of *Vibrio cholerae* non-O1 non-0139 with specific reference to their ability to produce cholera toxin, and addition of novel serogroups (Japanese, Abstract in English). *Kansenshogaku Zasshi.* 71: 1037 – 45