

# **A taxonomic review of white grubs and leaf chafers (Coleoptera: Scarabaeidae: Melolonthinae) recorded from forestry and agricultural crops in Sub-Saharan Africa**

**James du G. Harrison<sup>1,2,3,4\*</sup> and Michael J. Wingfield<sup>2</sup>**

<sup>1</sup>Department of Zoology & Entomology, University of Pretoria, 0002, South Africa. <sup>2</sup>Forestry & Agriculture Biotechnology Research Institute (FABI), University of Pretoria, 0002, South Africa. <sup>3</sup>School of Animal, Plant & Environmental Sciences (AP&ES), University of the Witwatersrand, 2050, Johannesburg, South Africa. <sup>4</sup>Department of Invertebrates, Ditsong: National Museum of Natural History (formerly Transvaal Museum), Pretoria 0001, South Africa.

## **Abstract**

Integrated Pest Management (IPM) is difficult to implement when one knows little about the pest complex or species causing the damage in an agricultural system. To implement IPM on Sub-Saharan African melolonthine pests access to taxon specific knowledge (their identity) and what is known (their biology) of potential pest species is a crucial step. What is known about Sub-Saharan Africa melolonthine white grubs and chafers has not yet been amalgamated, and this review thus synthesizes all available literature for the Region. The comprehensive nature of the review highlights pest taxon trends within African melolonthines. To facilitate the retrieval of this information for IPM purposes, all relevant taxonomic and biological information is provided for the taxa covered including an on-line supplementary annotated-checklist of taxon, crop, locality and reference(s). Based on the literature reviewed, recommendations are made to promote effective and efficient management of African melolonthine scarab pests. An on-line supplementary appendix provides a list of specialists, useful internet resources, keys, catalogues and sampling methods for the larvae and adults of melolonthine scarab beetles for subsequent morphological or molecular work.

**Keywords:** Africa, southern Africa, Scarabaeidae, Melolonthinae, White Grubs, Chafers, Scarab Pests, Larvae, Agriculture, Identification, Taxonomy.

## **Introduction**

This review aims to provide a checklist of the species of Sub-Saharan African Melolonthinae that have been recorded as pests in agricultural and forestry crops (Table 1 supplementary material),

---

\*Author for correspondence  
Fax: +27 11 717 6494  
E-mail: James.Harrison@wits.ac.za

and draws attention to groups most in need of revision from a pest management perspective. Additionally, it provides comprehensive review and bibliography of all traced literature (especially taxonomic) specific to the melolonthine white grubs and chafers of the Region. It is hoped that this will facilitate white grub research in Sub-Saharan Africa via enhanced access to information, and the reduction of duplicated research efforts. Based on this synthesis, recommendations relevant to white grub and chafer taxonomy in Sub-Saharan Africa are proposed. To constrain the size and scope of the review, it covers only the subfamily Melolonthinae, with the remaining scarab subfamilies that contain species of economic concern being dealt with in forthcoming studies.

The larvae (white grubs) and adults (chafers) of five subfamilies of Scarabaeidae (Aphodiinae, Dynastinae, Cetoniinae, Melolonthinae and Rutelinae) include species that feed on the roots, stems, fruit or foliage of many crops, and therefore may be sporadic forestry and agricultural pests (Table 1 supplementary material). Colloquially these larvae are known as white grubs, cane grubs or curl grubs when they damage agricultural crops in different regions of the World. White grubs are readily identified by their 'C-shaped' bodies and sclerotized head capsules, while the more variable adults generally have an ovoid body shape and lamellate antennae (Richter, 1958, 1966). Related scarab families, for example dung beetles (Scarabaeinae), have similar larvae, but as these groups are beneficial recyclers, they are generally not referred to as white grubs.

Melolonthine scarab beetles have a complete life-cycle (holometabolous); a fertilized female lays eggs, these hatch into the first of three larval stages or instars, the final (or third) instar enters a pre-pupal stage before pupation, from which adults emerge when environmental conditions are conducive (Ritcher, 1958, 1966). The duration of the larval stage can vary from one to three years depending on the environmental conditions and species' life cycle (Ratcliffe, 1991). However, species with two to three year life-cycles often have adults active each year due to an overlapping of generations. The phenology of the adults is unknown for most species, but based on adult phenological data compiled from museum records for *Asthenopholis* (Harrison, 2009) and *Pegylis* (Harrison, 2014b), adults from these genera are present in varying numbers throughout summer each year.

Species distributions are dependent on a variety of factors. Some white grubs have a narrow distribution (stenotopic, e.g., *Asthenopholis subfasciata* (Harrison, 2009) and *Macrophylla* spp.) while others have a wider distribution (eurytopic, e.g., *Pegylis sommeri* (Harrison, 2014b)). Soil type, texture and moisture content can play an important role in the distribution of certain white grub species (Allsopp, *et al.*, 1992; Cherry & Allsopp, 1991; Logan, 1997). Females of some white grub species are flightless, e.g., *Macrophylla pubens* (Fenwick, 1947; Omer-Cooper, *et al.*, 1941, 1941-1942, 1948), and this has implications for control strategies and the geographic extent of an outbreak. The adult activity period can be quite narrow (just a few days after rain) for some species

of chafer (Harrison, pers. obs.) or extend over a longer time period, i.e., weeks and months for *Pegylis sommeri* (see phenology figs in Harrison (2014b)).

### **Pest status of scarabs**

In large numbers the feeding activity of white grubs and chafers in crops reduces yields and facilitates secondary microbial infections through the damaged plant cuticle (Smith, *et al.*, 1995; Miller, *et al.*, 1999). For example, in southern Africa, white grubs have been recorded as sporadic subterranean pests on tree-seedlings, sorghum, sugarcane, pineapples, potatoes and turf grass (see Table 1 supplementary material for a complete list). Additionally, the adult chafers are often defoliators (e.g. *Pegylis* spp.) in forest plantations, fruit orchards, vineyards and rose gardens (Table 1 supplementary material). At least 50 different species of commercially grown plants have records of Melolonthinae being destructive to parts of these plants (Table 1 supplementary material).

### **Introduced alien species**

Most, if not all, African melolonthine scarab pests are endemic to parts of Africa, and no introduced scarab pest species are recorded for South Africa (Picker & Griffiths, 2011). However, as this is a broad and general information source, it by no means rules out the possibility of there being alien scarabs in Sub-Saharan Africa. A possible exception is *Phyllophaga smithi* (Arrow, 1912), which was originally described as *Phytalus smithi* from the Caribbean (Barbados & Trinidad) and introduced into Mauritius (Evans, 2003). This species was recorded (Macdonald, *et al.*, 2003: 74) from Tanzania and mentioned as a sorghum stem-borer in eastern Ethiopia (Tefera, 2004). However, there is no published confirmation of this introduction in the primary literature for sugarcane pests known from Tanzania, but the African endemic *Cochliotis melolonthoides* is well established as a sugarcane pest in Tanzania (Jepson, 1956; Carnegie, 1974a, b). Consequently, the Macdonald *et al.* (2003) and Tefera (2004) records of *P. smithi* occurring on the African mainland may be based on the exotics being confused with an African species of *Schizonycha*, which is quite similar to *Phyllophaga* (= *Phytalus*) Arrow (1912).

### **Conservation of localized endemic species**

Localized endemics are of particular conservation importance. For example, *Asthenopholis subfasciata* can become an important sporadic pest of pineapples in the Eastern Cape of South Africa (Petty, 1976, 1977a,b, 1978, 1982, 1990, 1994, 2001, Petty, *et al.*, 2002). However, it is endemic to South Africa and is also one of only seven known species of *Asthenopholis* (Harrison, 2009). As a localized endemic and part of the country's biodiversity, we need to control it responsibly when it reaches localized pest levels. Another example of a localized endemic that can become a sporadic pest species is *Pseudachloa leonina* on golf greens near Pretoria, South Africa (pers. comm. At Schoeman 2005).

### **Access to information**

Fragmentary, unsynthesized information relating to scarabs as pests in African crops reduces the efficiency of research on economically important species. For example, the paucity of information on *Eucamenta eugeniae*, originally described as a pest of clove (*Eugenia caryophyllata*) from Zanzibar (Arrow, 1932; Andre Moutia, 1941). But a recent paper (Conlong & Mugalula, 2003) omitted these early publications and reported *E. eugeniae* only as a new pest of sugarcane in Uganda.

Information relating to the same insect taxon, but disguised due to an incorrect identification is another problem. For example, the incorrect identification of *Pegylis sommeri* (previously *Hypopholis*) as *Macrophylla ciliata* (Herbst) as a pest of pineapple in South Africa (Petty, 1976, 1977b, 1978, 1990, 2001; Petty, *et al.*, 2002) resulted in the redescription of the larvae of *P. sommeri* (Smith, *et al.*, 1995), which had previously been described (Prins, 1965). Misidentification resulted in a duplication of effort, lack of efficient control strategies, crop destruction, and reduced economic benefits.

### **The importance of having named pests**

The success of an Integrated Pest Management (IPM) program for melolonthine white grubs depends strongly on accurate identification of larvae and adults (Danks, 1988). Species level identification is important because life cycles of white grub species vary from one to three years and susceptibility to insecticides varies for species (as demonstrated in Australian cane grubs; Allsopp, *et al.*, 1995). Correct diagnosis relies upon available taxonomic expertise, published research, trained systematists, and access to museum specimens, voucher specimens and type material. In contrast to more developed countries (e.g. the United States, Australia and New Zealand) southern Africa in our view lags behind by a number of years in scarab taxonomic research on melolonthine white grub identification and research.

In a discussion on white grub identification (Omer-Cooper, *et al.*, 1941; Omer-Cooper, *et al.*, 1941-1942) the authors state that, "... very little work has been done on these insects in South Africa, and with the present state of our knowledge we would not be prepared to say in all cases which are pests and which are not". Later (Sweeney, 1967) added that "Few published papers deal with the morphology and taxonomy of larval Scarabaeoidea in southern Africa, and the larvae of many species are undescribed". However, recent taxonomic work by Ahrens (2007a,b), Conlong & Mugalula (2003), Dittrich-Schröder *et al.* (2009), Goble *et al.* (2012) and Harrison (2004, 2009, 2014a,b, this review) is a positive advancement in melolonthine research in the Region.

### **Morphological identification of melolonthine larvae and adults**

To morphologically describe a scarab larva, the chitinous head capsule is softened in lactic acid or a weak solution of KOH and disarticulated from points of natural articulation of structures (Dittrich-Schröder, *et al.*, 2009). Ritcher (1966) provided numerous illustrations and descriptions of disarticulated scarab head capsules. This requires association of the larva with a named adult and formal description in a published paper. African melolonthines in which larvae have been described and illustrated (Table 2 supplementary material), include a small percentage of the known melolonthine diversity.

Adult identification also relies on the beetle species having been described in a validly published paper. Insect drawers of African Melolonthinae sorted only to subfamily in museum collections are testament to the lack of taxonomists dealing with this diverse group of beetles. Consequently, in many cases one can at best identify a pest melolonthine to genus level only, and even here good taxonomic keys for generic diagnoses (e.g. Péringuey, 1904; Lacroix, 2010) are the exception rather than the rule.

An ideal scenario would be as follows: a taxonomist is presented with larvae and adults of *Pegylis sommeri* and asked to identify these. In this particular instance, the larvae could be identified using Prins (1965) and the adults using Harrison (2014b). Adult specimens of *Asthenopholis* can be identified using Harrison (2009), but the larva of only one species of this genus has been described, i.e., *A. subfasciata* in Smith *et al.* (1995), leaving the larvae of the remaining six species unassociated with their adults via the taxonomic literature.

### **Molecular diagnostic techniques**

A frequently encountered problem with white grub outbreaks is determining the white grub species. Identification is especially important because control action must be immediate to prevent further losses. Molecular DNA barcoding techniques allow for the DNA of grubs or adults to be matched with DNA sequence data (e.g. archived Genbank sequences) and possible retrieval of an identification based on these data. Once a match is retrieved the morphology of the grubs can be described in order to facilitate accurate and efficient identification in the future. This process can be used to develop identification tools. For example, Dittrich-Schröder *et al.* (2009) compiled an identification key to the scarab beetle larvae attacking sugarcane in South Africa. Using molecular methods and associating grubs with adults, tools for identification of chafers were developed for use on the Nepalese fauna (Ahrens, *et al.*, 2007a,b). This research is best done by a collaborating team of molecular biologists, beetle taxonomists and crop specialists.

## **Integrated Pest Management (IPM) options for white grubs and chafers**

IPM of white grubs and chafers is a review topic of its own, and consequently only brief mention is included here. Control options include various forms of chemical control, cultural methods, parasitic Diptera (flies), Hymenoptera (wasps), fungi, and tillage methods. Smit (1964), Annecke & Moran (1982) and Visser (2005, 2009) provide general coverage of IPM methods relevant to the African context for many of the taxa included in this review.

## **Taxonomic review of the Melolonthinae with published records of significance to Sub-Saharan African forestry and agricultural crops**

The world melolonthine fauna is presently divided into 28 extant tribes and 12 subtribes (Smith, 2006), of which six tribes and nine subtribes are known to occur on the African mainland. Sub-Saharan Africa includes taxa from six tribes and six subtribes as presented below. All known Sub-Saharan African Melolonthinae recorded in the literature as having the potential to become sporadic pests in agricultural and forestry crops are reviewed (Table 1). Included in Table 1 are taxonomic details (generic and specific description dates and authors) and thus this information is not repeated in the text. However, taxa not included in the checklist (Table 1) are referred to in the text including their author and publication date.

Also summarized are melolonthine species, crop attacked, life stage involved, country of origin and the source reference (Table 1 supplementary material). Thus, if not specifically mentioned below, the life stage, larva or adult, involved in the damage is included in Table 1 supplementary material.

### **1. Tribe ABLABERINI Blanchard, 1850**

For garden flowers, Annecke & Moran (1982) list adults of *Ablabera pellucida* as feeding on carnations. Adults of *Camenta innocua* appears to occur in most major centers in South Africa (JduGH pers. obs.) and can be found at night feeding on a variety of garden plants, with roses, being one of those most seriously damaged (JduGH pers. obs.). The larvae of what was possibly *C. innocua* caused significant damage to raspberry seedlings in the George and Hermanus areas of South Africa (pers. comm. Tim Sobey Nov. 2012). Oberholzer (1959a,b) described the third instar larvae of *C. innocua* from specimens collected in Johannesburg, South Africa, but he did not indicate any pest status for the species.

In the interests of taxonomic stability, Ahrens (2007a) designated types for various African genera in the Ablaberini, including *Ablabera* Dejean, 1833; *Eucamenta* Péringuey, 1904; *Hybocamenta* Brenske, 1898; and *Paracamenta* Péringuey, 1904. Neither *Ablabera* nor *Camenta* have been recently revised, making species level identification in these genera difficult or impossible, especially bearing in mind the large number of undescribed taxa within the Ablaberini. The syntype

of *C. innocua* is a single female (pers. comm. Dirk Ahrens), making the unequivocal identity of this species currently impossible in the absence of the male genitalia necessary for identification.

## **2. Tribe DIPLLOTAXINI Kirby, 1837**

Peacock (1913), looking at insect pests of cacao or cocoa bean (*Theobroma cacao* L.) in southern Nigeria, recorded *Apogonia nitidula* together with various other scarabs, but included these in a list of taxa with doubtful significance (associated with the crop but not showing signs of being an actual pest). Lepesme & Paulian (1944) recorded the same species on *Coffea robusta* in Gabon, but they provided no further details about the association. In work applying to species from neighbouring South Asia Islam *et al.* (1984) included species of *Apogonia* among the insect pests associated with green gram (*Vigna radiata* L.) from Bangladesh, India. Of the beetles studied by Islam *et al.* (1984), *Apogonia* were responsible for consuming the largest leaf area.

Bezděk (2004a) provided a catalogue of Diplotaxini of the Old World and a detailed revision on the African monotypic genus *Ceratogonia* Kolbe, 1899 (Bezděk, 2004b). He provided synonymic notes for *Apogonia cupreoviridis* Kolbe, 1886 and *A. nigroolivacea* Heyden, 1886 (Bezděk, 2008), and notes on the synonymy and geographic distribution of *Apogonia niponica* Lewis, 1895 (Bezděk, 2009). Lacroix (2008a,b) described new genera and species of Diplotaxini, and Lacroix & Bezděk (2009) proposed replacement names for *Metagonia* Kolbe, 1899 (nec Simon, 1893).

## **3. Tribe HOPLIINI Latreille, 1829**

In South Africa, these beetles are colloquially referred to as monkey beetles (Picker & Midgley 1996). Smit (1964) recorded adult Hopliini burrowing into garden Compositae (Asteraceae), especially marigolds in Pretoria, South Africa. Prins (1965) listed *Monocheilus calcaratus*, which he referred to as the small wattle chafer, as a wattle defoliator. Prins (1965) also described its larva, the only description of an African Hopliini larva (known to us). For garden flowers, Annecke & Moran (1982) recorded *Eriesthis stigmatica* and *Heterochelus connatus* feeding on carnations, and *E. vestita* feeding on dahlias. Swain & Prinsloo (1986) list species of *Heterochelus* feeding on *Acacia mearnsii*, and *Monocheilus calcaratus* feeding on *Acacia decurrens*, *A. mearnsii* and *A. melanoxylon*. Gess (1968), Myburgh *et al.* (1973, 1974), Myburgh & Rust (1975) and Coetzee & Giliomee (1985) investigated insects associated with *Protea* flowers as both pests damaging the flowers and insects likely to be exported with flowers due to their biological associations with the *Protea*. Myburgh & Rust (1975) provided a list of 34 'free-living pests in protea flower-heads', including the following monkey beetles: *Anisonyx nasuus*, *Heterochelus rufimanus*, and a species of *Platycheilus*. Coetzee & Giliomee (1985), in a study specific to *Protea repens* (L.), listed adult *Diaplocheilus longipes* as a floral visitor. Dombrow (2006b) described two new species of *Diaplocheilus* sent abroad with consignments of exported *Protea* from South Africa to Poland and the USA.

Due to the charismatic morphological diversity of the southern African Hopliini, this tribe has received greater taxonomic attention than any other in the Region. Dombrow, in particular, has revised many of the southern African genera (a list of his known papers is provided). These papers included taxonomic revisions of *Eriesthis* (Dombrow, 1997c, 2002b), *Heterochelus* (Dombrow, 2001c) and *Diaplochelus* (Dombrow, 2006b). Schein (1958) studied South African *Heterochelus* and his paper included a key (pp. 256-257) to the genera and subgenera of Heterochelina (= Hopliina).

### **3a. Subtribe HOPLIINA Latreille, 1829**

Swain & Prinsloo (1986) included adults of *Hoplia sordita* as feeding on *Acacia mearnsii*. We are not aware of any other references relating to the biology or pest status of *Hoplia* species in Sub-Saharan Africa, but there is extensive literature on the non-African *Hoplia*. For example, Ansari *et al.* (2008) examined the susceptibility of *Hoplia philanthus* (Füessly) larvae and pupae to entomopathogenic nematodes in Belgium, while Zhang *et al.* (2011) examined the attractive responses of *Hoplia spectabilis* Medvedev to un-baited yellow or white cross-pane funnel traps in China.

### **4. Tribe PACHYDEMINI Burmeister, 1855**

Omer-Cooper *et al.* (1941) noted that southern Africa turf (golf courses) is a human-made habitat often composed of the indigenous grass (*Cynodon dactylon* Pers.), and thus its pests are endemic, opportunistically making use of an abundant food resource. The damage caused by white grubs when they feed on grass roots, completely 'scalps' the roots and can result in death of the grass, especially following a dry period. Larvae of species of *Macrophylla* are potential pests of golf greens in the Eastern Cape of South Africa (Omer-Cooper, *et al.*, 1941).

Omer-Cooper *et al.* (1941, 1941-1942, 1948) provided notes on the biology of species of *Macrophylla*, where the females of at least some of the species appear to be flightless. Fenwick (1947) redescribed the male of *M. pubens* and provided the first description of its flightless female based on specimens collected on the Humewood Golf Green near Port Elizabeth, South Africa. Schoeman (1996) recorded that white grubs in turfgrass are "*usually patchily distributed, unpredictable and capable of doing severe damage before they are detected*". Due to the incorrect identification of *Hypopholis sommeri* (now *Pegylis sommeri*) as *Macrophylla ciliata* and published as *M. ciliata* in Petty's earlier papers, the larva of *M. ciliata* (but actually *P. sommeri*) was described by Smith *et al.* (1995). However, this is actually a redescription of the larva of *P. sommeri* (see Prins, 1965). Consequently, we are not aware of any descriptions of southern African Pachydemini larvae (here we exclude *Sparrmannia flava* covered by Evans (1989) and Scholtz (1988), due to the preferred placement of *Sparrmannia* in the Melolonthini).



Evans (1988a) reviewed the taxonomy and systematics of southern Africa Pachydemini and revised some genera (Evans, 1987a,b,c, 1988b). Lacroix provided descriptions of many new African Pachydemini genera and species in 14 papers (Lacroix, 1997, 1999, 2001, 2003, 2004, 2005a,b, 2006a,b,c,d, 2008e, 2009f, 2011a) and a catalogue (Lacroix, 2007).

### **5. Tribe MELOLONTHINI Leach, 1819**

#### **5a. Subtribe MELOLONTHINA Leach, 1819**

Swain & Prinsloo (1986) included *Psilonychus groendahli* on *Acacia mearnsii* in a list of phytophagous insects on forest trees and shrubs in South Africa. Specimens of *Psilonychus* were collected in pheromone traps set to catch the lepidopterous pest *Helicoverpa armigera* (African Bollworm) from chicory fields in the Eastern Cape of South Africa (Midgley *et al.*, 2008, Midgley pers. comm.). There are eight described species of *Psilonychus* and Harrison (in prep.) plans to revise this genus, making its species and what we know about them accessible to other entomologists.

The adults of three *Sparrmannia* species (*S. acicularis*, *S. flava* and *S. transvaalica*) were recorded in the vicinity of pistachio nut trees in the Prieska area of the Northern Cape of South Africa by Louw (2001). However, only *S. flava* caused economic damage by defoliating trees, thereby forcing them to produce new leaves. It is suspected that this regrowth reduced the ultimate nut yield of trees (Louw, 2001). Haddad *et al.* (2005) and Haddad & Louw (2006) recorded the spider groups present in pistachio orchards in South Africa and noted their potential as biological control agents for pistachio pests. *Sparrmannia* species are caught in spiders' webs (Harrison pers. obs.) and thus one can expect that both large sedentary and orb-weaving spiders would catch and feed on *Sparrmannia* adults. Scholtz (1988) investigated the breeding biology of *S. flava* in the arid Kalahari and noted that, under desert conditions, the life cycle is one year in duration. Larvae are able to maximize feeding after rain and then become dormant until rain allows them to feed again. Quite unique for the melolonthines, or perhaps just unrecorded due to the absence of studies on African arid chafers, is the larval behaviour of foraging at night on the soil surface for antelope dung, which is then taken below the soil surface, rehydrated and fed upon by these larvae. Evans (1989) summarized what is known about the biology of *Sparrmannia*. All species appear to feed on plants as adults, but there is a lack of established host plant records, even from mass swarming observations. Of the 24 known species, two appear to be diurnal while many other species have been collected at light traps at night. Mating swarms of *S. transvaalica* have been observed at dusk, and in the morning and late afternoon (Harrison pers. obs.). Evans (1989) provided a detailed and thorough revision of the 24 known species of *Sparrmannia* and a description of the second instar larvae of *S. flava*.

### **5b. Subtribe SCHIZONYCHINA Burmeister, 1855**

Larvae of *Entyposis impressa* were recorded by Mansfield-Aders (1920) feeding on the roots of caladiums and castor oil plants. Lacroix & Montreuil (2012) revised *Entyposis* and recognized nine species in the genus. Male *Entyposis* are unusual amongst melolonthines in that they have indented dynastine-like pronota including small pronotal horns making it easy to recognize this genus of African melolonthines.

Schmutterer (1964), in an overview of the insect pests on southern Somalian crops, noted larvae of a species of *Schizonycha* feeding on banana plant roots. Büttiker & Bünzli (1957) included *Schizonycha profuga*, *S. citima* and *Hecistopsilus molitor* in their survey of the more important leaf chafers on tobacco in Zimbabwe. Tarr (1958) examined the relationship between feeding by species of *Schizonycha* grubs on Dolichos Bean (*Dolichos lablab*) seedlings, resulting in wilt and often followed by stem blight. They discuss how various chemical seed dressings can reduce losses. In a list of phytophagous insects in South Africa, Swain & Prinsloo (1986) listed *S. fimbriata* as associated with *Acacia mearnsii*. Larvae of *S. fimbriata* caused damage to soya beans in KwaZulu-Natal (Hittersay, 2005). In the Trans Nzoia district of Kenya, changes in the intercropping practices to enhance soil nutrition resulted in an increase in the abundance of root-feeding chafer (*Schizonycha* spp.) grubs (Medvecký, *et al.*, 2006, 2007; Medvecký & Ketterings 2009). Apparently, the enhanced soil nutrition may have benefitted white grubs developing in the soil. In a review of the insect pests associated with yam production and storage, Korada *et al.* (2010) included unnamed species of *Schizonycha* larvae as boring into and feeding on the tubers during the pre-harvest period. Hajek *et al.* (2005) in their 'Catalogue of introductions of pathogens and nematodes for classical biological control of insects and mites' listed *Paenibacillus popilliae* (Dutky) (Bacillaceae) as having been released in 1956 in Kenya for the control of *Schizonycha* spp. In this instance, *P. popilliae* was not recovered after release (Hajek, *et al.*, 2005). *Schizonycha* is the African equivalent to the diverse *Phyllophaga* Harris of the New World. Pope (1960) revised the southern African species of *Schizonycha*, and at that time recorded over 300 described species, with 292 of these being from Africa, and 120 being from southern Africa. Undoubtedly, many undescribed species remain, complicating efforts for routine identification of this important pest containing genus. Lacroix (2010) provided a World checklist of 349 species for *Schizonycha*.

### **5c. Subtribe PEGYLINA Lacroix, 1989**

Harrison (2014a) provides a phylogeny for the Pegylina while Harrison (2014b) provides detailed coverage of the natural history, pest status, larval description, parasites and previous chemical control of South African *Pegylis* species. Lacroix (2008c, d) described new species of *Pegylis* and included all known species of *Pegylis* in his 2010 book.

#### **5d. Subtribe LEUCOPHOLINA Burmeister, 1855**

Lepesme & Paulian (1944) recorded larvae of *Afrolepis pygidialis* feeding on the roots of *Coffea excelsa* in Gabon, provided a brief description of the larvae and pupae of *A. pygidialis*, and a key to the species in the genus. Decelle (1968) proposed *Afrolepis* as a replacement name for *Oligolepis* Brenske, 1903 as the latter generic name was already occupied within the vertebrates. Lacroix (2010) provided redescriptions, a key to species, and maps for the five known species of *Afrolepis*.

Harrison (2009) provides detailed coverage of the natural history and pest status of *Asthenopholis*, revises the genus and recognizes seven species of *Asthenopholis* from these countries: South Africa (5 spp.), Lesotho (1), Swaziland (1), Kenya (1), Tanzania (2) and Uganda (1).

The larvae of *Cochliotis melolonthoides* are considered an important pest of sugarcane in Tanzania (Jepson, 1956). Lacroix (2009a) provided a revision of the four known species of *Cochliotis*, including the description of a new species from Somalia. *Cochliotis* species are known from Kenya, Tanzania and Somalia (Lacroix, 2009a). Jepson (1956) discussed the biology, larvae (including third instar head capsule and rastal pattern illustrations) and adults of *C. melolonthoides*. Additionally, Jepson (1956) included recommendations for the cultural, biological and chemical control of *C. melolonthoides* larvae, which are pests of sugarcane in Tanzania. Hajek *et al.* (2005) listed *Paenibacillus popilliae* (Dutky) (Bacillaceae) as having been released in 1968 in Tanzania for the control of *C. melolonthoides*. It is possible that *P. popilliae* became established, but the presence of an indigenous milky disease confounded conclusive results.

Taylor & Smithers (1959) recorded *Eulepida mashona* as a pest of Ley pasture in Zimbabwe. “*In ley farming, the field is alternately used for grain or other cash crops for a number of years and ‘laid down to ley’, i.e., left fallow, used for growing hay or used for pasture for another number of years. After that period it is again ploughed and used for cash/field crops*” (Wikipedia, 2015). Towards the end of the growing season in Zimbabwe, Wilson (1963, 1972) mentioned that larvae of *E. mashona* can damage maize plants when the cobs begin to ripen in March. In their list of phytophagous insects on forest trees and shrubs, Swain & Prinsloo (1986) included *E. mashona* as associated with *Acacia mearnsii*. Gomez (1988) included *E. mashona* in the list of ‘*indigenous and traditional foods in Zimbabwe*’. From an IPM perspective eating *E. mashona* would reduce high populations of this chafer and also provide rural communities with free nutrients. However, Tagwireyi *et al.* (2000) reported a case of cantharadin poisoning presumably caused by ingestion of a blister beetle (Meloidae) by a four-year-old child in Zimbabwe. The cause of the ingestion is thought to be due to mistaken identification of the edible *Eulepida mashona* with the blister beetle *Mylabris distincta* Thomas. Lacroix (2008f, 2009a,b,c,d,e, 2011b) revised various genera of Leucopholina. Lacroix (2010) covered the 26 known species of *Eulepida* (including descriptions of

three new species); he also provided descriptions, keys, distribution maps and line drawings, including the male genitalia for all 26 species.

### **6. Tribe SERICINI Kirby, 1837**

#### **6a. Subtribe SERICINA Kirby, 1837**

In Nigeria, *Pseudotrochalus concolor* is associated with the following plant species: *Bixa orellana*, *Citrus acida*, *Citrus limonia*, *Gossypium hirsutum*, *Haematoxylon campechianum*, *Quisqualis indica*, *Spondias lutea* and *Zea mais* (Golding, 1927, 1937). Certain genera of African Sericinae (Ahrens, 2007a) are currently being revised (Ahrens pers com).

#### **6b. Subtribe TROCHALINA Brenske, 1898**

In southern Nigeria, Peacock (1913) recorded several species of *Trochalus* together with other scarabs on cacao or cocoa beans (*Theobroma cacao* L.), but he placed these taxa on his list of doubtful significance (associated with the crop but not an actual pest). Hargreaves (1937) recorded *T. carinatus*, *T. gibbus* and *T. pilula* as feeding on avocado, *Cola nitida* and cotton in Sierra Leone. Swain & Prinsloo (1986) included *T. byrrhinus* and *T. fulgidus* in their list of phytophagous insects of forest trees and shrubs in South Africa for *Acacia mearnsii*.

## **Conclusion**

Identification of melolonthine white grubs and adult chafers unlocks published information on these taxa and creates the necessary nomenclatural foundation for research into the biology and ecology of the species. This review provides a foundation for taxonomic research on this agriculturally important pest group. Research is required in all areas relating to African white grub and leaf chafers in order to improve our ability to make species level pest identifications, understand their biology, and implement IPM strategies when required. It is important to conserve these same species as they are also African endemics and an integral part of the continent's rich biodiversity. Recent work by Ahrens, Bezděk, Conlong, Dittrich-Schröder, Goble, Harrison, Lacroix and Way (see reference list) indicates a positive trend to more taxonomic and applied research being undertaken on South African and southern African scarab pests.

## **Recommendations**

Based on this review of the literature from 1889 (Ormerod & Janson, 1889) to the present, the following recommendations can be made to improve the continuity of information relating to African scarab pest taxonomy and associated research in future. These include:

- inclusion of the following key words in paper titles / keyword lists to facilitate electronic literature searchers, i.e., 'white grub, leaf chafer, Scarabaeidae, Melolonthinae, Africa and scarab pest';

- clear indication of the methods used to establish larvae-to-adult conspecificity, when identifications are provided, e.g., adults bred from larvae, larvae and adults found in association with one another, or via molecular matching techniques;
- a clear statement of the person responsible (and their professional affiliation) for taxon identifications, e.g., 'Riaan Stals of the ARC-PPRI<sup>2</sup>'; and
- information on the national museum / research institution in which the vouchers on which the identification is based have been deposited, e.g., 'ARC-PPRI or SANC<sup>3</sup>'.

### Acknowledgements

We extend our gratitude to these people and organizations for their contributions to this paper: Darren J. Mann, Gudrun Dittrich-Schroder, Mike Way, Paul Schoolmeesters, Petr Šípek and Tarryn Goble for providing literature; Riaan Stals and Beth Grobbelaar (ARC-PPRI) for access to their extensive pest related beetle reprint collection; Kevin Balkwill allowed JduGH to attend a number of writing retreats at WITS' Pullen Nature Reserve, where much of the review was written; Irene and Mick Jackson provided funding for the upgrade of facilities at Pullen N.R. making the Jackson Field Station such a pleasant research environment; The WITS Faculty Research Committee and WITS SPARC covered JduGH's research costs. Marcus Byrne (WITS, RSA), Mary Liz Jameson (Wichita State University, USA), Connal Eardley (ARC-PPRI), Simon van Noort (SAMC<sup>4</sup>) and Brett Hurley (FABI<sup>5</sup>) provided valuable comments to an earlier version of the paper; Dirk Ahrens provided his knowledge and insight into the current taxonomy of the Ablaberini, Sericini and Trochalina. Most importantly, JduGH thanks his family and especially Margaret A.C. Harrison for their love, support and patience while completing his PhD.

### References

- Ahrens, D.** (2007a) Type species designations of Afrotropical Ablaberini and Sericini genera (Coleoptera: Scarabaeidae: Melolonthinae). *Zootaxa* **1496**, 53–62.
- Ahrens, D.** (2007b) Taxonomic changes and an updated catalogue for the Palaearctic Sericini (Coleoptera: Scarabaeidae: Melolonthinae). *Zootaxa* **1504**, 1-51.
- Ahrens, D., Monaghan, M.T. & Vogler, A.P.** (2007a) DNA-based taxonomy for associating adults and larvae in multi-species assemblages of chafers (Coleoptera: Scarabaeidae). *Molecular Phylogenetics and Evolution* **44**, 436–449.
- Ahrens, D., Zorn, C., Dhoj, Y.G.C., Keller, S., & Nagel, P.** (2007b) Illustrated key of phytophagous scarabs of Nepal. A guide to white grubs and chafers of the lower central regions (Coleoptera, Scarabaeidae). *Opuscula Biogeographica Basiliensia* **5**, 1–44.

<sup>2</sup> Agricultural Research Council - Plant Protection Research Institute. <http://www.arc.agric.za>

<sup>3</sup> South African National Collection of Insects. <http://www.arc.agric.za/home>

<sup>4</sup> Iziko, South African Museum <http://www.iziko.org.za/museums/south-african-museum>

<sup>5</sup> Forestry and Agricultural Biotechnology Institute <http://www.fabinet.up.ac.za/>

- Allsopp, P.G., Klein, M.G. and McCoy, E.L.** (1992) Effect of soil moisture and soil texture on oviposition by Japanese beetle and rose chafer (Coleoptera: Scarabaeidae). *Journal of Economic Entomology* **85**, 2194-2200.
- Allsopp, P.G., McGill, N.G. and Bull, R.M.** (1995) Use of suSCon Blue against larvae of *Lepidiota picticollis* Lea (Coleoptera: Scarabaeidae) in Australian sugar-cane and the effect of infestation on yield. *Crop Protection* **14**, 69-73.
- Andre Moutia, L.** (1941) The search for parasites of white grubs (Melolonthids) in Zanzibar, Algeria, Morocco and France. *Bulletin of Entomological Research* **31**, 193-208.
- Annecke, D.P. & Moran, V.C.** (1982) *Insects and mites of cultivated plants in South Africa*. Butterworths, Durban, South Africa. 1-383.
- Anonymous** (1992) White Grub Survey Report 1991–1992 Season, Swaziland Sugar Association Extension Services, Swaziland, 10 pp.
- Ansari, M.A., Adhikari, B.N., Ali, F. & Moens, M.** (2008) Susceptibility of *Hoplia philanthus* (Coleoptera: Scarabaeidae) larvae and pupae to entomopathogenic nematodes (Rhabditida: Steinernematidae, Heterorhabditidae). *Biological Control* **47**, 315-321.
- Arrow, G.J.** (1912) XLVII - On a new species of Melolonthine beetle (*Phytalus smithi*) destructive to sugar-cane. *The Annals and Magazine of Natural History* **8**(9), 455-459.
- Arrow, G.J.** (1917) Some insects injurious to Cacao plants in the Belgian Congo. *Bulletin of Entomological Research* **8**, 111-118.
- Arrow, G.J.** (1925) XXIII. A few new species of melolonthine Coleoptera. *The Annals and Magazine of Natural History* **16**(9), 209-221.
- Arrow, G.J.** (1932) XXVII. A few new species of melolonthine Coleoptera. *The Annals and Magazine of Natural History* **9**(10), 209-221.
- Barnes, B.N., Daiber, K.C. & Donaldson, J.M.** (1986) Fruit eating and leaf-eating beetles. pp. 35-42 in Myburgh, A.C. (Editor) 1986. *Crop pests in southern Africa. Vol. 1 Deciduous fruit, grapes and berries*. Bulletin **407**, Plant Protection Research Institute, Pretoria. 1-92.
- Bezděk, A.** (2004a) Revision of the genus *Ceratogonia* Kolbe, 1899. *Annales Zoologici* **54**(4), 797-801.
- Bezděk, A.** (2004b) Catalogue of Diplotaxini of the Old World. *Zootaxa* **463**, 1-90.
- Bezděk, A.** (2008) Synonymical notes on *Apogonia cupreoviridis* and *A. nigroolivacea*. *Annales Zoologici (Warsaw)* **58**(1), 71-77.
- Bezděk, A.** (2009) Synonymical and geographic distribution notes for *Apogonia niponica* Lewis, 1895. *The Coleopterists Bulletin* **63**(4), 438-444.
- Bünzli, G.H. & Büttiker, W.W.** (1960a) Fungous diseases of lamellicorn larvae in southern Rhodesia. *Bulletin of Entomological Research* **50**, 89-96.
- Bünzli, G.H. & Büttiker, W.W.** (1960b) A comparative study of the soil conditions favored by certain melolonthid and tenebrionid pests of tobacco in southern Rhodesia. *Bulletin of Entomological Research* **50**, 259-263.

- Burgeon, L.** (1945a) Les Ablaberini du Congo belge (Coleopt. Scarab. Melolonthinae). *Revue de Zoologie et de Botanique Africaines* **38**(3-4), 212-251.
- Burgeon, L.** (1945b) Les Diplotaxina du Congo belge (Coleopt. Scarab. Melolonthinae). *Revue de Zoologie et de Botanique Africaines* **39**(1), 1-23.
- Burgeon, L.** (1946a) Melolonthini et Pachydemini du Congo belge (suite) (Coleopt. Scarab. Melolonthinae). *Revue de Zoologie et de Botanique Africaines* **39**(3), 230-273.
- Burgeon, L.** (1946b) Melolonthini et Pachydemini du Congo belge (suite). *Revue de Zoologie et de Botanique Africaines* **39**(4), 339-366.
- Büttiker, W.W. & Bünzli, G.H.** (1957) Biological notes on Southern Rhodesian tobacco pests (1): records of host plants of adult chafer beetles. *The Rhodesian Agricultural Journal* **54**(5), 462-472.
- Carnegie, A.J.M.** (1974a) Insects of sugarcane in South Africa. pp. 1-14 in Carnegie, A.J.M., Dick, J. & Harris, R.H.G. 1974. Insects and nematodes of South African sugarcane. *Journal of the Entomological Society of Southern Africa Memoirs* **39**, 1-19.
- Carnegie, A.J.M.** (1974b) Sugarcane white grubs (Scarabaeoidea) and their control in South Africa. *Proceedings of International Society of Sugar Cane Technologists* **15**, 498-512.
- Carnegie, A.J.M.** (1988) White grubs (Scarabaeoidea) continue to cause sporadic damage to sugarcane in South Africa and Swaziland. *Proceedings of the South African Sugar Technologists' Association* (June), 161-163.
- Cherry, R.H. & Allsopp, P.G.** (1991) Soil texture and the distribution of *Antitrogus parvulus* Britton, *Lepidiota crinita* Brenske and *L. negatoria* Blackburn (Coleoptera: Scarabaeidae) in South Queensland sugarcane fields. *Journal of the Australian Entomological Society* **30**, 89-92.
- Coetzee, J.H. & Giliomee, J.H.** (1985) Insects in association with the inflorescence of *Protea repens* (L.) (Proteaceae) and their role in pollination. *Journal of the Entomological Society of Southern Africa* **48**(2), 303-314.
- Colwill, D.J.** (1967) *Pest control in groundnuts*. PANS Manual No. **2**, 24-54.
- Conlong, D.E. & Mugalula, A.M.** (2003) *Idaecamenta eugeniae* (Coleoptera: Scarabaeidae: Melolonthinae): a new species of white grub in Ugandan sugarcane. *Proceedings of the South African Sugar Technologists Association* **77**, 250-255.
- Dalla Torre, K.W. von** (1912a) Fam. Scarabaeidae. Subfam. Melolonthinae. I. Coleopterorum catalogus, vol. 20, pars 45, Scarabaeidae; Melolonthidae I. pp. 1-84.
- Dalla Torre, K.W. von** (1912b) Fam. Scarabaeidae. Subfam. Melolonthinae. II. Coleopterorum catalogus, vol. 20, pars 47, Scarabaeidae; Melolonthidae II. pp. 85-134.
- Dalla Torre, K.W. von** (1912c) Fam. Scarabaeidae. Subfam. Melolonthinae. III. Coleopterorum catalogus, vol. 20, pars 49, Scarabaeidae; Melolonthidae III. pp. 135-290.
- Dalla Torre, K.W. von** (1913) Fam. Scarabaeidae. Subfam. Melolonthinae. IV. Coleopterorum catalogus, vol. 20, pars 50, Scarabaeidae; Melolonthidae IV. pp. 291-450.

- Danks, H.V.** (1988) Systematics in support of entomology. *Annual Review of Entomology* **33**, 271-296.
- de Villiers, E.A., van den Berg, M.A. & Petty, G.T.** (1987) Borers and root feeders. pp. 91-96 in Myburgh, A.C. 1987. *Crop pests in southern Africa. Vol. 2 Citrus and other subtropicals*. Bulletin **411**. Plant Protection Research Institute, Pretoria. 1-118.
- Decelle, J.** (1968) Contribution à l'étude des Melolonthidae d'Afrique: II. Notes sur les Leucopholini et description d'espèces nouvelles. *Revue de Zoologie et de Botanique Africaines* **77**, 62-82.
- Dittrich-Schröder, G. Conlong, D.E., Way, M.J., Harrison, J. du G. & Mitchell, A.** (2009) Identification key to scarabaeid larvae attacking sugarcane in South Africa using DNA barcoding and integrative taxonomy. *Proceedings of the South African Sugar Technologists Association* **82**, 500-524.
- Dombrow, H.** (1997a) Revision of the genus *Denticnema* Schein 1959 (Melolonthinae: Hopliini). *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **1**, 1-21.
- Dombrow, H.** (1997b) Revision of the genus *Pariesthis* Moser 1918 (Melolonthinae: Hopliini). *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **1**, 22-39.
- Dombrow, H.** (1997c) Revision of the genus *Eriesthis* Burmeister (Melolonthinae: Hopliini). *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **1**, 41-115.
- Dombrow, H.** (1997d) New species of Hopliini from the Northern and Western Cape Provinces of the Republic of South Africa (Scarabaeidae: Melolonthinae). *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **1**, 137-159.
- Dombrow, H.** (1997e) Revision of the genera *Chasme* Serville 1825, *Khoina* Péringuey 1902 and *Lepitrichula* Schein 1959 (Melolonthinae: Hopliini). *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **1**, 161-201.
- Dombrow, H.** (1998a) Revision of the genus *Hoplocnemis* Harold 1869 (Melolonthinae: Hopliini). *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **2**, 1-37.
- Dombrow, H.** (1998b) Revision of the genus *Pachycnema* Serville 1825 (Melolonthinae: Hopliini). *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **2**, 39-110.
- Dombrow, H.** (1999a) New species of Hopliini from southern Africa. 8<sup>th</sup> Contribution to the knowledge of African Hopliini (Scarabaeidae: Melolonthinae). *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **3**, 139-144.
- Dombrow, H.** (1999b) Revision of the genus *Clania* Schein, 1956 (Melolonthinae: Hopliini). 9<sup>th</sup> Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **3**, 145-158.
- Dombrow, H.** (1999c) Revision of the genus *Lepisia* Serville, 1825 (Melolonthinae: Hopliini). 10<sup>th</sup> Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **3**, 159-176.



- Dombrow, H.** (1999d) Revision of the genus *Scelophysa* Burmeister, 1844 (Melolonthinae: Hopliini). 11<sup>th</sup> Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **3**, 177-202.
- Dombrow, H.** (2000a) Revision of the genus *Lepithrix* Serville, 1825 (Melolonthinae: Hopliini). 12<sup>th</sup> Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **4**, 93-146.
- Dombrow, H.** (2000b) Two new species of the genus *Hoplocnemis* Harold, 1869 from southern Africa (Melolonthinae: Hopliini). 13<sup>th</sup> Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **4**, 147-152.
- Dombrow, H.** (2001a) Revision of the genus *Gymnoloma* Burmeister 1844 with observations on three new genera (Melolonthinae: Hopliini). 14<sup>th</sup> Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **5**, 107-148.
- Dombrow, H.** (2001b) Two new species of the genus *Lepithrix* Serville from Northern and Western Cape Provinces of the Republic of South Africa (Melolonthinae: Hopliini). 15<sup>th</sup> Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **5**, 427-432.
- Dombrow, H.** (2001c) Three new species of the genus *Heterochelus* Burmeister, 1844 out of Kulzer's 11<sup>th</sup> group from Northern Cape (Melolonthinae: Hopliini). 16<sup>th</sup> Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **5**, 433-440.
- Dombrow, H.** (2002a) *Dentiheterochelus* gen. n. and *Toxocnemis* gen. n., two new genera of Hopliini from the Republic of South Africa (Scarabaeidae: Melolonthinae). 17<sup>th</sup> Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **6**, 103-108.
- Dombrow, H.** (2002b) A new species of the genus *Eriesthis* Burmeister, 1844 from Namibia (Melolonthinae: Hopliini). 18<sup>th</sup> Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **6**, 109-112.
- Dombrow, H.** (2002c) A new species of the genus *Gymnoloma* Burmeister from Western Cape of the Republic of South Africa (Melolonthinae: Hopliini). 19<sup>th</sup> Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **6**, 113-116.
- Dombrow, H.** (2002d) *Amblymelanoplia* a new genus of South African Hopliini (Scarabaeidae: Melolonthinae). 20<sup>th</sup> Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **6**, 117-128.
- Dombrow, H.** (2002e) *Nigroplia* gen. n. and *Mauromecistoplia* gen. n., two new genera of South African Hopliini (Scarabaeidae: Melolonthinae). 21<sup>st</sup> Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **6**, 129-137.

- Dombrow, H.** (2002f) *Ceratochelus* gen.n. and *Paraceratochelus* gen.n., two new genera of South African Hopliini (Scarabaeidae: Melolonthinae). 22<sup>nd</sup> Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **6**, 139-156.
- Dombrow, H.** (2003) *Argoplia* gen.n., *Burmeistoplia* gen.n. and *Dolichiomicroscelis* gen.n. three new genera of South African Hopliini (Scarabaeidae: Melolonthinae). 23<sup>rd</sup> Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **7**, 103-115.
- Dombrow, H.** (2005a) Three new species of the genus *Lepithrix* Serville from the Northern and Western Cape Provinces of the Republic of South Africa (Melolonthinae: Hopliini). 24<sup>th</sup> Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **9**, 117-124.
- Dombrow, H.** (2005b) *Beckhoplia* a new genus of Hopliini from the Northern Cape of the Republic of South Africa (Scarabaeidae: Melolonthinae). 25<sup>th</sup> Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **9**, 125-130.
- Dombrow, H.** (2006a) Two new species of the genus *Knysna* Péringuey from Western Cape of the Republic of South Africa (Melolonthinae: Hopliini). 28<sup>th</sup> Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **10**, 83-88.
- Dombrow, H.** (2006b) Two new species of the genus *Diaplochelus* Burmeister from the Republic of South Africa (Melolonthinae: Hopliini). 29<sup>th</sup> Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **10**, 89-94.
- Dombrow, H.** (2006c) Two new species of the genus *Scelophysa* Burmeister from Western Cape of the Republic of South Africa (Melolonthinae: Hopliini). 30<sup>th</sup> Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **10**, 95-100.
- Dombrow, H.** (2006d) Two new species of the genus *Lepisia* Serville from the Northern and Western Cape of the Republic of South Africa (Melolonthinae: Hopliini). 31<sup>st</sup> Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **10**, 101-106.
- Dombrow, H.** (2007a) Three new species of the genus *Denticnema* Schein 1959 from the Western Cape of the Republic of South Africa (Melolonthinae: Hopliini). 32<sup>nd</sup> Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **11**, 125-132.
- Dombrow, H.** (2007b) Two new species of the genus *Dolichiomicroscelis* Dombrow 2003 from the Western Cape of the Republic of South Africa (Melolonthinae: Hopliini). 33<sup>rd</sup> Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **11**, 133-138.
- Dombrow, H.** (2007c) Eleven new species of the genus *Lepithrix* Serville from the Northern and Western Cape Provinces of the Republic of South Africa (Melolonthinae: Hopliini). 34<sup>th</sup>

- Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **11**, 139-160.
- Dombrow, H.** (2007d) Six new species of the genus *Kobousa* Péringuey from the Western Cape of the Republic of South Africa (Melolonthinae: Hopliini). 35<sup>th</sup> Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **11**, 161-174.
- Dombrow, H.** (2008) *Omocnemus mitzlaffi* sp. n. a new species of the genus *Omocnemus* Schein from Namibia (Melolonthinae: Hopliini). 36<sup>th</sup> Contribution to the knowledge of African Hopliini. *Coleoptera, Schwanfelder Coleopterologische Mitteilungen* **12**, 221-226.
- Elmer, J.L. & Smith, R.G.** (1956) Some pests of Black Wattle in Kenya: with a list of other insects inhabiting the plantations. *East African Agricultural Journal* **21**, 230-246.
- Evans, A.V.** (1987a) A new genus of Melolonthinae from southern Africa. *Journal of the Entomological Society of Southern Africa* **50**(2), 363-370.
- Evans, A.V.** (1987b) Revision of the genus *Aipeiopsis* Péringuey. *Journal of the Entomological Society of Southern Africa* **50**(2), 481-486.
- Evans, A.V.** (1987c) A new species of *Eucyclophylla* Waterhouse. *Journal of the Entomological Society of Southern Africa* **50**(2), 487-491.
- Evans, A.V.** (1988a) *Systematics of the Sub-Saharan Pachydemini (Coleoptera, Melolonthidae, Melolonthinae)*. Doctoral dissertation, Department of Entomology. Pretoria, University of Pretoria, 1-210.
- Evans, A.V.** (1988b) Three new species of *Glyptoglossa* Brenske. *Journal of the Entomological Society of Southern Africa* **51**(1), 87-96.
- Evans, A.V.** (1989) Revision of the genus *Sparrmannia* Laporte (Coleoptera, Melolonthidae, Melolonthinae). *Journal of the Entomological Society of Southern Africa* **52**(1), 11-44.
- Evans, A.V.** (2003) A checklist of the New World chafers (Coleoptera: Scarabaeidae: Melolonthinae). *Zootaxa* **211**, 1-458.
- Evans, A.V. & Smith, A.B.T.** (2009) *An electronic checklist of the New World Chafers (Coleoptera: Scarabaeidae: Melolonthinae). Version 3*. Electronically published, Ottawa, Canada. 353 pp. Available from <http://museum.unl.edu/research/entomology/SSSA/nwmeos.htm> (accessed 15 February 2015).
- Feakin, S.D.** (1973) *Pest control in groundnuts*. PANS Manual No. 2. pp. 1-197.
- Fenwick, E.M.** (1947) On *Macrophylla pubens* Péringuey (Coleoptera, Scarabaeidae, Melolonthinae). *Journal of the Entomological Society of Southern Africa* **10**, 24-30.
- Gess, F.W.** (1968) Insects found on Proteas. *Journal of the Botanical Society of South Africa* **54**, 29-33.
- Goble, T.A., Costet, L., Robene, I., Nibouche, S., Rutherford, R.S., Conlong, D.E. & Hill, M.P.** (2012) *Beauveria brongniartii* on white grubs attacking sugarcane in South Africa. *Journal of Invertebrate Pathology* **111**, 225–236.

- Golding, F.D.** (1927) Notes on the food-plants and habits of some southern Nigerian Insects. *Bulletin of Entomological Research* **17**, 95-99.
- Golding, F.D.** (1937) Further notes on the food plants of Nigerian Insects IV. *Bulletin of Entomological Research* **28**, 5-9.
- Gomez, M.** (1988) A resource inventory of indigenous and traditional foods in Zimbabwe. *Zambezia* **15**(1), 53-73.
- Govender, P.** (2007) Status of seedling establishment pests of *Acacia mearnsii* De Wild. (Mimosaceae) in South Africa. *South African Journal of Science* **103**, 141-147.
- Haddad, C.R., Dippenaar-Schoeman, A.S. & Pekár, S.** (2005) Arboreal spiders (Arachnida: Araneae) in pistachio orchards in South Africa. *African Plant Protection* **11**, 32–41.
- Haddad, C.R. & Louw, S. vd M.** (2006) Phenology, ethology and fecundity of *Heliophanus pistaciae* Wesolowska (Araneae: Salticidae), an agrobiont jumping spider in South African pistachio orchards. *African Plant Protection* **12**, 1–11.
- Hajek, A.E., McManus, M.L., Delalibera Júnior, I.** (2005) *Catalogue of introductions of pathogens and nematodes for classical biological control of insects and mites*. U.S. Department of Agriculture, Forest Service Publication FHTET-2005-05, 1-65.
- Hargreaves, E.** (1937) Some insects and their food plants in Sierra Leone. *Bulletin of Entomological Research* **28**, 505-520.
- Harrison, J.duG.** (2004) Revision of the endemic southern African genus *Rhabdopholis* Burmeister, 1855 (Coleoptera: Scarabaeidae: Melolonthinae). *African Entomology*, **12**(1), 39-54.
- Harrison, J.duG.** (2009) A taxonomic revision of the African leaf chafer genus *Asthenopholis* Brenske, 1898 (Coleoptera: Scarabaeidae: Melolonthinae): a SEM study. *Zootaxa* **2225**, 1-48.
- Harrison, J.duG.** (2014a) A morphological analysis of the subtribe Pegylina Lacroix, 1989 (Scarabaeidae: Melolonthinae: Melolonthini) reconstitutes its generic composition. *African Entomology* **22**(4): 726–741.
- Harrison, J.duG.** (2014b) Review of the South African species of *Pegylis* Erichson, 1847 (Coleoptera: Scarabaeidae: Melolonthinae) commonly known as large wattle chafers. *African Entomology* **22**(4): 685–713.
- Hepburn, G.A.** (1966) A revised list of wattle insects and spiders of southern Africa. *Wattle Research Institute Report* 67-83.
- Hittersay, P.** (2005) White grub damages soya crop. *Farmers Weekly* 1 April. 23.
- Howard, C.W.** (1910) Cockchafers and flower beetles. *Transvaal Agricultural Journal* **6**, 1-4.
- Islam, W., Ahmed, K.N. & Joarder, O.I.** (1984) Timing and extent of damage caused by insect pests of green gram (*Vigna radiata* L.) in Bangladesh. *Crop Protection* **3**(3), 343-348.
- Jepson, W.F.** (1956) The biology and control of the sugar-cane chafer beetles in Tanganyika. *Bulletin of Entomological Research* **47**, 377-397.

- Korada, R.R., Naskar, S.K., & Edison, S.** (2010) Insect pests and their management in yam production and storage: a world review. *International Journal of Pest Management* **56**(4), 337-349.
- Lacroix, M.** (1997) Contribution à la connaissance des Pachydeminae africains. I. Nouveaux genres et nouvelles espèces (Coleoptera, Scarabaeoidea, Melolonthidae). *Revue française d'Entomologie, (N.S.)* **19**(1-2), 51-62.
- Lacroix, M.** (1999) Contribution à la connaissance des Pachydeminae africains. II. Nouveaux genres et nouvelles espèces. Définition des caractères génériques pour la région afrotropicale (Coleoptera, Scarabaeoidea, Melolonthidae). *Coléoptères* **5**(4), 43-83.
- Lacroix, M.** (2001) Pachydeminae de l'Est africain (Kenya et Tanzanie) (Coleoptera, Melolonthidae). *Coléoptères* **7**(13), 173-226.
- Lacroix, M.** (2002) Nouveaux Melolonthidae afrotropicaux (Coleoptera, Scarabaeoidea). *Coléoptères* **8**(9), 153-168.
- Lacroix, M.** (2003) Contribution à la connaissance des Pachydeminae africains, III. Nouveau genre et nouvelles espèces (Coleoptera, Melolonthidae). *Coléoptères* **9**(6), 63-82.
- Lacroix, M.** (2004) Contribution à la connaissance des Pachydeminae africains, IV. Nouvelles espèces (Coleoptera, Melolonthidae). *Coléoptères* **10**(9), 107-118.
- Lacroix, M.** (2005a) Contribution à la connaissance des Pachydeminae africains, V. Nouvelles espèces de *Oedanomerus* Waterhouse (Coleoptera, Melolonthidae). *Coléoptères* **11**(6), 51-65.
- Lacroix, M.** (2005b) Contribution à la connaissance des Pachydeminae africains, VI. Nouveaux genres et nouvelles espèces (Coleoptera, Melolonthidae). *Coléoptères* **11**(22), 273-300.
- Lacroix, M.** (2006a) Contribution à la connaissance des Pachydeminae africains, VII. Le genre *Goniorrhina* Quedenfeldt, 1888 (Coleoptera, Melolonthidae). *Coléoptères* **12**(4), 23-46.
- Lacroix, M.** (2006b) Contribution à la connaissance des Pachydeminae africains, VIII. Nouveaux genres et nouvelles espèces (Coleoptera, Melolonthidae). *Coléoptères* **12**(11), 159-178.
- Lacroix, M.** (2006c) Contribution à la connaissance des Pachydeminae africains, IX. Nouveau genre et nouvelles espèces (Coleoptera, Melolonthidae). *Coléoptères* **12**(14), 203-214.
- Lacroix, M.** (2006d) Contribution à la connaissance des Pachydeminae africains, X. Nouveau genre, mises au point et désignation de lectotypes (Coleoptera, Melolonthidae). *Coléoptères* **12**(18), 235-270.
- Lacroix, M.** (2007) *Pachydeminae du Monde, Genera et Catalogue* (Coleoptera, Melolonthidae). Editions M. Lacroix / Collection Hannetons. 1-450 + 264 plates.
- Lacroix, M.** (2008a) Contribution à la connaissance des Melolonthidae africains, I. Nouveaux genres et nouvelles espèces du complexe *Apogonia* Kirby, 1819 (Coleoptera, Melolonthidae). *Coléoptères* **14**(7), 67-98.
- Lacroix, M.** (2008b) Contribution à la connaissance des Melolonthidae africains, II. Nouvelles espèces d' *Apogonia* Kirby (Coleoptera, Melolonthidae). *Coléoptères* **14**(9), 105-116.

- Lacroix, M.** (2008c) Contribution à la connaissance des Melolonthidae africains, III. Nouvelles espèces de *Pegylis* Erichson, 1848 (Coleoptera, Melolonthidae). *Coléoptères* **14**(12), 133-147.
- Lacroix, M.** (2008d) Nouvelles espèces de *Pegylis* Erichson, 1848, d'Afrique orientale (Coleoptera, Melolonthidae). *Coléoptères* **14**(16), 163-186.
- Lacroix, M.** (2008e) Révision du genre *Aglaphyra* Brenske, 1896, d'Afrique orientale (Coleoptera, Melolonthidae, Pachydeminae). *Coléoptères* **14**(19), 213-228.
- Lacroix, M.** (2008f) *Tanzanilepis badia*, nouveau genre et nouvelle espèce de Leucopholini d'Afrique orientale (Coleoptera, Melolonthidae). *Coléoptères* **14**(20), 229-240.
- Lacroix, M.** (2009a) Révision du genre *Cochliotis* Kolbe (Coleoptera, Melolonthidae, Leucopholini). *Coléoptères* **15**(6), 39-50.
- Lacroix, M.** (2009b) Révision du genre *Lepidomela* Kolbe (Coleoptera, Melolonthidae, Leucopholini). *Coléoptères* **15**(7), 51-62.
- Lacroix, M.** (2009c) Révision des genres *Spaniolepis* Kolbe et *Camerunopholis* Lacroix (Coleoptera, Melolonthidae, Leucopholini). *Coléoptères* **15**(8), 63-78.
- Lacroix, M.** (2009d) Révision du genre *Brachylepis* Kolbe (Coleoptera, Melolonthidae, Leucopholini). *Coléoptères* **15**(19), 153-168.
- Lacroix, M.** (2009e) Révision du genre *Pholidochris* Kolbe (Coleoptera, Melolonthidae, Leucopholini). *Coléoptères* **15**(20), 169-196.
- Lacroix, M.** (2010) *Melolonthinae afrotropicaux, Genera et Catalogue* (Coleoptera, Melolonthidae). Editions M. Lacroix, Paris. 1-277, 140 plates + 52 colour plates.
- Lacroix, M.** (2011) *Kenyailepis montana*, nouveau genre et nouvelle espèce de Leucopholini d'Afrique orientale (Coleoptera, Melolonthidae). *Coléoptères* **17**(2), 13-20.
- Lacroix, M. & Bezděk, A.** (2009) Battle of two replacement names for the genus *Metagonia* Kolbe, 1899 (Coleoptera: Scarabaeidae: Melolonthinae). *Zootaxa* **2237**, 67–68.
- Lacroix, M. (2012. & Montreuil, O.** (2012) Révision du genre *Entyposis* Kolbe, 1894 (Insecta, Coleoptera, Scarabaeoidea, Melolonthidae). *Zoosystema* **34**(4), 721-735.
- Lepesme, P. & Paulian, R.** (1944) Sur deux Mélolonthides récoltés sur Caféier dans l'Ouest africain. *Revue de Zoologie et de Botanique Africaines* **38**(1-2), 99-103.
- Logan, P.** (1997) Effect of soil moisture on oviposition by Childers Canegrub, *Antitrogus parvulus* Britton (Coleoptera: Scarabaeidae). *Australian Journal of Entomology* **36**, 175-178.
- Louw, S. vd M.** (2001) Diversity and flight activity of chafers (Coleoptera: Scarabaeidae) associated with pistachio in South Africa. *Proceedings of the 13<sup>th</sup> Entomological Congress of the Entomological Society of Southern Africa* 41-42.
- Macdonald, I.A.W., Reaser, J.K., Bright, C., Neville, L.E., Howard, G.W., Murphy, S.J. & Preston, G.** (Eds) (2003) *Invasive alien species in southern Africa: National reports & directory of resources*. pp 1-125. Global Invasive Species Programme, Cape Town, South Africa.

- Mansfield-Aders, W.** (1920) Insects injurious to economic crops in the Zanzibar Protectorate. *Bulletin of Entomological Research* **10**(2), 145-155 + plates XIII-XV.
- Medvecky, B.A. & Ketterings, Q.M.** (2009) Incorporation of legume residues does not increase productivity of intercropped beans on smallholder farms in Trans-Nzoia District, Kenya. *Biological Agriculture and Horticulture* **26**(4), 323-335.
- Medvecky, B.M., Ketterings, Q.M. & Nelson, E.B.** (2007) Relationships among soil borne bean seedling diseases, *Lablab purpureus* L. and maize stover residue management, bean insect pests, and soil characteristics in Trans Nzoia District, Kenya. *Applied Soil Ecology* **35**, 107–119.
- Medvecky, B.M., Ketterings, Q.M. & Vermeulen, F.M.** (2006) Bean seedling damage by root-feeding grubs (*Schizonycha* spp.) in Kenya as influenced by planting time, variety, and crop residue management. *Applied Soil Ecology* **34**, 240–249.
- Midgley, J.M., Hill, M.P. & Villet, M.H.** (2008) Baited traps may be an alternative to conventional pesticides in integrated crop management of chicory (Compositae) in South Africa. *Journal of Economic Entomology* **101**(1), 99-106.
- Miller, L.J., Allsopp, P.G., Graham, G.C. and Yeates, D.K.** (1999) Identification of morphologically similar canegrubs (Coleoptera, Scarabaeidae, Melolonthini) using a molecular diagnostic technique. *Australian Journal of Entomology* **38**, 189-196.
- Mitchell, B.L.** (1946) White grub control in tobacco lands. *The Rhodesia Agricultural Journal* **43**(6), 499-504.
- Myburgh, A.C., Rust, D.J. & Starke, L.C.** (1973) Pests of protea cut-flowers. *Journal of the Entomological Society of Southern Africa* **36**(2), 251-255.
- Myburgh, A.C., Starke, L.C. & Rust, D.J.** (1974) Destructive insects in the seed heads of *Protea barbiger* Meisner (Proteaceae). *Journal of the Entomological Society of Southern Africa* **37**(1), 23-29.
- Oberholzer, J.J.** (1959a) A morphological study of some South African Lamellicorn larvae. I - Descriptions of the third instar larvae. *South African Journal of Agricultural Science* **2**, 41-74.
- Oberholzer, J.J.** (1959b) A morphological study of some South African Lamellicorn larvae. II - Comparative morphology. *South African Journal of Agricultural Science* **2**, 75-88.
- Oberprieler, R.G., Donaldson, J.M., Daiber, K.C., Coetzee, J.H., de Villiers, E.A., Rust, D.J. & Myburgh, A.C.** (1990) Flower- and leaf-eating beetles. pp. 21-27 in Myburgh, A.C. 1990. *Crop pests in southern Africa. Vol. 5 Flowers and other ornamentals*. Bulletin **419**. Plant Protection Research Institute, Pretoria. 1-98.
- Omer-Cooper, J., Whitnall, A.B.M. & Fenwick, E.M.** (1941) The problem of South African turf in relation to invertebrate pests. *South African Journal of Science* **37**, 273-284.
- Omer-Cooper, J., Whitnall, A.B.M. & Fenwick, E.M.** (1941-1942) The insect and the golfer. Reprinted from *South African Golf* (June 1941 to August 1942), 1-20.

- Omer-Cooper, J., Whitnall, A.B.M. & Fenwick, E.M.** (1948) Notes on four species of Coleoptera attacking turf in the Eastern Cape Province. *South African Journal of Science* **44**, 125-134.
- Ormerod, E.A. & Janson, O.E.** (1889) *Notes and descriptions of a few injurious farm and fruit insects of South Africa*. Simpkin, Marshall and Co., Stationer's Hall Court, London. 1-116.
- Peacock, A.D.** (1913) Entomological pests and problems of southern Nigeria. *Bulletin of Entomological Research* **4**, 191-220 + plates XXIII-XXVIII.
- Péringuey, L.** (1904) Descriptive catalogue of the Coleoptera of South Africa (Lucanidae and Scarabaeidae). *Transactions of the South African Philosophical Society* **13**, 1-293 + 4 plates.
- Petty, G.J.** (1976) The bionomics and control of pineapple white grubs (Coleoptera: Scarabaeoidea) in the Eastern Cape. *Crop Production* **5**, 55-59.
- Petty, G.J.** (1977a) Beetle pests of pineapples, some biological aspects. *The Citrus and Subtropical Fruit Journal* (Dec.), 4-7.
- Petty, G.J.** (1977b) White-grub rears head in Eastern Cape. *Citrus and Subtropical Fruit Research Institute Information Bulletin* **53**, 1-2.
- Petty, G.J.** (1978) White grubs in pineapples. *Farming in South Africa* **12**, 1-4.
- Petty, G.J.** (1982) Isazophos, a promising new insecticide for control of larvae of fossorial rhizophagous beetles (Coleoptera, Scarabaeoidea) in pineapple fields. *Journal of the Entomological Society of Southern Africa* **45**(1), 33-41.
- Petty, G.J.** (1990) White grubs in pineapples. *Farming in South Africa* **12**, 1-2.
- Petty, G.J.** (1994) White grubs in pineapples, an alternative prevention strategy. *Institute for Tropical and sub-tropical Crops Information Bulletin* **265**(Sept.), 11-12.
- Petty, G.J.** (2001) Order Coleoptera: family Scarabaeidae: white grubs. pp. 247–257 in van den Berg, M.A., de Villiers, E.A. & Joubert, P.H. (Eds) *Pests and Beneficial Arthropods of Tropical and Non-Citrus Subtropical Crops in South Africa*. ARC-Institute for Tropical and Subtropical Crops, Mbombela. pp 1-525.
- Petty, G.J., Stirling, G.R. & Bartholomew, D.P.** (2002) Pests of pineapple. pp. 157–189 in Peña, J.E., Sharp, J.L. & Wysoki, M. (Eds) *Tropical Fruit Pests and Pollinators: biology, economic importance, natural enemies and control*. CABI publishing.
- Picker, M.D. & Griffiths, C.** (2011) *Alien and invasive animals. A South African perspective*. Struik, Cape Town. 1-240.
- Picker, M.D. & Midgley, J.J.** (1996) Pollination by monkey beetles (Coleoptera: Scarabaeidae: Hopliini): flower and colour preferences. *African Entomology* **4**(1), 7–14.
- Pollard, D.G.** (1956) The control of chafer grubs (*Schizonycha* sp., Coleoptera, Melolonthinae) in the Sudan. *Bulletin of Entomological Research* **47**(3), 347-360.
- Pope, R.D.** (1960) A revision of the species of *Schizonycha* Dejean (Col., Melolonthidae) from southern Africa. *Bulletin of the British Museum (Natural History) Entomology* **9**(2), 65-216 + 14 plates.



- Prins, A.J.** (1965) Notes on the biology and morphology of the wattle chafers *Monochelus calcaratus* Burm. (Melolonthidae), *Hypopholis sommeri* Burm. (Melolonthidae), and *Adoretus ictericus* Burm. (Rutelidae) with some references to natural enemies (Coleoptera: Lamellicornia). *Entomology Memoirs* **9**, 1-55.
- Ratcliffe, B.C.** (1991) The scarab beetles of Nebraska. *Bulletin of the University of Nebraska State Museum* **12**, 1-333.
- Ritcher, P.O.** (1938) A field key to Kentucky white grubs. *Journal of the Kansas Entomological Society* **11**(1), 24-27.
- Ritcher, P.O.** (1949) Larvae of Melolonthinae with keys to tribes, genera, and species (Coleoptera: Scarabaeidae). *Kentucky Agricultural Experiment Station Bulletin* **537**, 1-36 + plates i-iv.
- Ritcher, P.O.** (1958) Biology of Scarabaeidae. *Annual Review of Entomology* **3**, 311-334.
- Ritcher, P.O.** (1966) White Grubs and Their Allies, a Study of North American Scarabaeoid Larvae. *Oregon State University Monograph Series No. 4*, 1-219.
- Ritcher, P.O.** (1967) Keys for identifying larvae of Scarabaeoidea (Coleoptera) to the family and subfamily. *California Department of Agriculture, Occasional Papers No. 10*, 1-8.
- Schein, H.** (1958) Bestimmungstabellen der Heterochelides mit ausnahme von *Heterochelus* und *Ischnochelus*. *Annals of the South African Museum* **44**(6), 229-267.
- Schmutterer, H. von** (1964) Zur kenntnis der schädlinge and krankheiten der kulturpflanzen in Südsomalia. *Journal of Pest Science* [no vol.], 102-106.
- Schoeman, A.** (1996) Scarabaeidae as pests of turfgrass in South Africa. in Way, M.J. and du Toit, H.A. (Eds) *White grub working group: pest status, biology and rearing of white grub (Coleoptera: Scarabaeidae) pests in southern Africa*. 6-7 June 1996. *SASRI Technical Report, Mount Edgecombe, KwaZulu-Natal*, 1-23.
- Scholtz, C.H.** (1988) Biology of *Sparrmannia flava* Arrow (Coleoptera, Scarabaeidae, Melolonthinae). *The Coleopterists Bulletin* **42**(1), 57-62.
- Schoolmeesters, P.** (2015) <http://www.datascarabaeoidea.net/> (accessed 21 February 2015).
- Smit, B.** (1964) Chapter 27. Dung beetles, cockchafers, and other beetles. pp. 273-278 in *Insects in southern Africa: How to control them*. Oxford University Press, Cape Town, 1-399.
- Smith, A.B.T.** (2006) A review of the family-group names for the Superfamily Scarabaeoidea (Coleoptera) with corrections to nomenclature and a current classification. *Coleopterists Society Monograph* **5**, 144–204.
- Smith, T.J., Petty, G.J. & Villet, M.H.** (1995) Description and identification of white grubs (Coleoptera, Scarabaeidae) that attack pineapple crops in South Africa. *African Entomology* **3**(2), 153-166.
- Swain, V.M. & Prinsloo, G.L.** (1986) A list of phytophagous insects and mites on forest trees and shrubs in South Africa. *Journal of the Entomological Society of Southern Africa Memoirs* **66**, 1-91.

- Sweeney, C.** (1967) The Scarabaeoidea associated with sugar-cane in Swaziland; an account of preliminary investigations into the bionomics and control, August 1965 –June 1967. *Swaziland Ministry of Agriculture Research Bulletin* **16**, 1-163.
- Tagwireyi, D., Ball, D.E., Loga, P.J. & Moyo, S.** (2000) Case report. Cantharidin poisoning due to blister beetle ingestion. *Toxicon* **38**, 1865-1869.
- Tarr, S.A.J.** (1958) Experiments in the Sudan Gezira on control of wilt of *Dolichos* bean (*Dolichos lablab*) associated with attack by cockchafer grubs (*Schizonycha* sp.). *Annals of Applied Biology* **46**(4), 630-638.
- Taylor, C.E. & Smithers, C.N.** (1959) The biology and control of whitegrubs (*Eulepida mashona* Arrow) in Ley Pastures. *Rhodesia Agricultural Journal* **56**(6), 240-242.
- Tefera, T.** (2004) Farmers' perceptions of sorghum stem-borer and farm management practices in eastern Ethiopia. *International Journal of Pest Management* **50**(1), 35-40.
- van Eeden, C.F., van Rensburg, J.B.J. & de K. van der Linde, T.C.** (1991) A check list of insects occurring in groundnut (*Arachis hypogaea*) fields in the Highveld Region. *Phytophylactica* **23**, 255-259.
- Visser, D.** (2005) *Guide to potato pests and their natural enemies in South Africa*. ARC-Roodeplaat Vegetable and Ornamental Plant Institute, Pretoria. 1-105.
- Visser, D.** (2009) *A complete guide to vegetable pests in South Africa*. ARC-Roodeplaat Vegetable and Ornamental Plant Institute, Pretoria. 1-316.
- Wang, H.** (1997) Chapter 5. Black wattle plantations in South Africa: Protection. pp. 39-53 in *Black Wattle and its Utilisation - Abridged English Edition*. Edited for the Rural Industries Research & Development Corporation by Brown, A.G. & Ho Chin Ko. RIRDC Publication No. 97/72.
- Wikipedia** (2015) [http://en.wikipedia.org/wiki/Ley\\_farming](http://en.wikipedia.org/wiki/Ley_farming). Accessed 13 March 2015.
- Williams, J.R.** (1985) White grubs in sugar cane, Report on a visit to Swaziland 30 July to 4 September 1985. *Sugar Industry Research Institute Réduit Report*. 1-24.
- Wilson, K.J.** (1963) Underground pests of maize in Northern Rhodesia. *Rhodesia Agricultural Journal* **60**(4), 111-114.
- Wilson, K.J.** (1972) A review of soil dwelling insects that have become important pests of agriculture in Rhodesia. *Rhodesia Agricultural Journal* **69**(1), 17-21.
- Zhang, Qing-He, Ma, Jian-Hai, Yang, Qi-Qing, Byers, J.A., Klein, M.G. Zhao, Feng-Yu & Luoe, You-Qing.** (2011) Olfactory and visual responses of the long-legged chafer *Hoplia spectabilis* Medvedev (Coleoptera: Scarabaeidae) in Qinghai Province, China. *Pest Management Science* **67**, 162-169.

## Supplementary material

### Appendix 1. Supplementary information covering useful scarab biology and taxonomy resources.

Queries to JduGH on African scarab pests indicated a need to provide a starting point for non-scarab specialist to available scarab resources, which is included below as supplementary information to the review.

#### Scarab beetle specialists

The 'Scarab Workers World Directory' provides an excellent resource of World scarab taxonomists and biologists. The "*Scarab Beetle Taxonomy Discussion List*" ([http://listserv.unl.edu/...](http://listserv.unl.edu/)) "... is for professionals, amateurs, and students interested in scarab beetle taxonomy. It is useful for posting questions, gathering information, and noting new discoveries regarding scarab systematics, biology, and nomenclature." Thus sending a question to this list provides one with immediate access to many specialists working in this field World Wide.

#### Scarab literature

Paul Schoolmeester's website <http://www.datascaraebaeoidea.net/> provides an alphabetical list of thousands of scarab related papers. Un-copyrighted papers are downloadable, and Paul can be e-mailed for many of the others as he maintains a large PDF reprint collection. Obviously, university or research based professional researchers have access to online search engines, but the advantage of those listed above is that one only requires internet access, rather than library subscriptions to journal databases. Additionally, Schoolmeester's site hosts obscure, gray literature and older literature that is difficult to obtain.

#### Important larval references

A list of scarab papers by Dr. Paul Richter is available at <http://museum.unl.edu/research/entomology/workers/PRitcher.htm> These papers by Richter (1938, 1949, 1958, 1966 & 1967) are equally useful to anyone interested in white grub taxonomy in southern Africa. Richter's work, especially Richter (1966), directs one to much of the earlier scarab larval research including coverage of larval terminology.

#### Catalogues and checklists

Dalla Torre (1912a,b,c & 1913) provided in Junk's *Coleopterorum Catalogus* a now outdated, but nevertheless vital catalogue of melolonthine genera, species and their type localities. Evans (2003) compiled '*A checklist of the New World chafers*', and Smith & Evans (2005) updated this with '*A supplement to the checklist of the New World chafers ... with notes on their tribal classification*'.

Although these last two papers do not cover the African fauna, they provide valuable bibliographies of literature relevant for an understanding of the World melolonthine fauna. An electronic checklist (Evans & Smith 2009) combining these two papers is available from [www.museum.unl.edu/research/ento](http://www.museum.unl.edu/research/ento). Bezděk (2004) catalogued the Diplotaxini of the Old World. Ahrens (2007a, b) designated type species for Afrotropical Ablaberini and Sericini and provided an updated catalogue of the Palaeartic Sericini which includes many North African taxa. Lacroix (2010) covered most of the known African leaf chafer genera and species (adults only) including many useful illustrations, maps and keys to taxa. Lacroix (2010) included a checklist of the described species of *Schizonycha*, and thus Pope (1960) remains the primary source for identification of this difficult and diverse genus, which includes many sporadic pest species (Table 3). Smith's (2006) review of the family-group names for the Superfamily Scarabaeoidea (Coleoptera), which included corrections to its nomenclature and a current classification, provides a high-level classification for the understanding of scarab taxa within a tribal and subtribal classification.

### **Keys to taxa**

Presently all-encompassing keys to the known genera and species of African leaf chafers do not exist. The most recent generic revisionary work should be used for species level keys, for example, Harrison (2009) for *Asthenopholis*, Harrison (2014b) for South African *Pegylis*, and Lacroix (2010) for many other African melolonthines. Additionally, clear habitus photographs can be e-mailed to the first author for a generic or tribal level diagnosis. Péringuey (1904) includes keys to many southern African taxa. Burgeon (1945a, b 1946a, b) provided useful keys to the Ablaberini, Diplotaxini, Melolonthini and Pachydemini, respectively.

### **Collection and preservation of scarab larvae and adults**

Reliable species level identification is based on adult or larval specimens. A few of the better known species, e.g., *P. sommeri*, can be identified from a digital habitus photograph, but this is one of the exceptions. A comparative series of 10 to 20 individuals per morphospecies, ideally including male and female adults should be collected, allowing for analysis of morphological variation within a species. A similar number of larvae will provide sufficient material for their dissection, and subsequent deposition in museum collections. Primary collecting data should include: Country, Province, Locality, latitude and longitude, date, collector's name, and as much other biologically relevant information as possible, e.g., crop, type of damage, extent of damage, etc.

Larvae can be killed by placing them in boiled water for 3-5 mins, which denatures their proteins allowing for better preservation once placed into 70% ethanol. Adults can be frozen, and then kept frozen, or dried, or also placed into 70% ethanol. For molecular vouchers, grub heads can be cut off and preserved in absolute ethanol, while the body can be boiled and then stored in 70%

ethanol. Live adults can be killed in absolute ethanol, which is then replenished 12-24 hours later with new absolute ethanol and then kept in a fridge or freezer before DNA extraction.

**Appendix 1 supplementary material.** Additional references not cited in the review article but referred to in this online appendix.

- Burgeon, L.** (1945a) Les Ablaberini du Congo belge (Coleopt. Scarab. Melolonthinae). *Revue de Zoologie et de Botanique Africaines* **38**(3-4), 212-251.
- Burgeon, L.** (1945b) Les Diplotaxina du Congo belge (Coleopt. Scarab. Melolonthinae). *Revue de Zoologie et de Botanique Africaines* **39**(1), 1-23.
- Burgeon, L.** (1946a) Melolonthini et Pachydemini du Congo belge (suite) (Coleopt. Scarab. Melolonthinae). *Revue de Zoologie et de Botanique Africaines* **39**(3), 230-273.
- Burgeon, L.** (1946b) Melolonthini et Pachydemini du Congo belge (suite). *Revue de Zoologie et de Botanique Africaines* **39**(4), 339-366.
- Dalla Torre, K.W. von** (1912b) Fam. Scarabaeidae. Subfam. Melolonthinae. II. Coleopterorum catalogus, vol. 20, pars 47, Scarabaeidae; Melolonthidae II. pp. 85-134.
- Dalla Torre, K.W. von** (1912c) Fam. Scarabaeidae. Subfam. Melolonthinae. III. Coleopterorum catalogus, vol. 20, pars 49, Scarabaeidae; Melolonthidae III. pp. 135-290.
- Dalla Torre, K.W. von** (1913) Fam. Scarabaeidae. Subfam. Melolonthinae. IV. Coleopterorum catalogus, vol. 20, pars 50, Scarabaeidae; Melolonthidae IV. pp. 291-450.
- Ritcher, P.O.** (1938) A field key to Kentucky white grubs. *Journal of the Kansas Entomological Society* **11**(1), 24-27.
- Ritcher, P.O.** (1949) Larvae of Melolonthinae with keys to tribes, genera, and species (Coleoptera: Scarabaeidae). *Kentucky Agricultural Experiment Station Bulletin* **537**, 1-36 + plates i-iv.
- Ritcher, P.O.** (1967) Keys for identifying larvae of Scarabaeoidea (Coleoptera) to the family and subfamily. *California Department of Agriculture, Occasional Papers* No. **10**, 1-8.
- Schoolmeesters, P.** (2015) <http://www.datascarabaeoidea.net/> (accessed 21 February 2015).

**Table 1 supplementary material.** Checklist of the Sub-Saharan African melolonthine white grub and leaf chafer taxa (Scarabaeidae: Melolonthinae) recorded from agricultural crops, including the life stage involved, larvae (L) or adult (A), locality (if stated), and source reference. Locality abbreviations include: Democratic Republic of the Congo (DRC), KwaZulu-Natal (KZN), South Africa (SA), North (N), South (S), East (E) and West (W). For non-taxonomists, *cf.* is short for ‘confer’ meaning ‘compare with’ and is used when taxonomists have a specimen that is close to a species, but not being absolutely confident they indicate via *cf.* that it might also be another closely related species. Consequently, it is best to regard these as genus level, rather than species level identifications. Taxonomy of the Ablaberini, Sericini and Trochalina courtesy of Dr Dirk Ahrens, including the endnotes for this section of the table.

Taxon	Crop	Life stage/s	Locality	Reference/s
<b>ABLABERINI Blanchard, 1850</b>				
<i>Ablabera</i> Dejean, 1833				
<i>A. pellucida</i> Burmeister, 1855	Carnations	A	Pretoria, SA	Annecke & Moran 1982
<i>A. pellucida</i>	Roses, Carnations, Dahlias etc.	A	SA	Oberprieler <i>et al.</i> , 1990
<i>Camenta</i> Erichson, 1847				
<i>C. innocua</i> Boheman, 1857	Roses	A	Pretoria, SA	Harrison (pers obs)
<i>C. cf. innocua</i>	Raspberry	L	Hermanus, George, SA	Tim Sobey (pers comm 2012)
<i>Camenta sp.</i>	Sugarcane	L	KZN Midlands, SA	Dittrich-Schröder <i>et al.</i> , 2009
<i>Idaecamenta</i> Péringuey, 1904				
<i>I. eugeniae</i> Arrow, 1932	Clove ( <i>Eugenia caryophyllata</i> )	L	Zanzibar	Arrow 1932
<i>I. eugeniae</i>	Clove ( <i>Eugenia caryophyllata</i> )	L	Zanzibar	Andrea Moutia 1941
<i>I. eugeniae</i>	Sugarcane	L	Uganda	Conlong & Mugalula 2003
<i>Paracamenta</i> Péringuey, 1904				
<i>P. suturalis</i> Péringuey, 1904	Wattle	A	Southern Africa	Hepburn 1966
<b>DILOTAXINI Kirby, 1837</b>				
<i>Apogonia</i> Kirby, 1819				
<i>A. nitidula</i> Thomson, 1858	Cacao or cocoa bean ( <i>Theobroma cacao</i> L.)	A	Southern Nigeria	Peacock 1913

Taxon	Crop	Life stage/s	Locality	Reference/s
<i>A. nitidula</i>	<i>Coffea robusta</i>	A	Gabon	Lepesme & Paulian 1944
<b>HOPLIINI Latreille, 1829</b>				
Hopliini <i>sensu lato</i>	Compositae esp. marigolds	A	Pretoria, SA	Smit 1964
<i>Anisonyx</i> Latreille, 1807				
<i>Anisonyx ursus</i> (Fabricius, 1775)	Garden flowers	A	Cape, South Africa	Oberprieler <i>et al.</i> , 1990
<i>Anisonyx nasuus</i> Wiedeman in Germar, 1821	Proteaceae ( <i>Protea</i> )	A	W. & S. Cape, SA	Myburgh & Rust 1975
<i>Congella</i> Péringuey, 1902				
<i>C. valida</i> Péringuey, 1902 <sup>i</sup>	Pineapple	L	Eastern Cape, SA	Smith <i>et al.</i> , 1995
<i>Diaplochelus</i> Burmeister, 1844				
<i>Diaplochelus karnkowskii</i> Dombrow, 2006	<i>Protea</i> sp.	A	W. Cape, SA & Poland <sup>ii</sup>	Dombrow 2006b
<i>Diaplochelus olseni</i> Dombrow, 2006	<i>Protea</i> sp.	A	N. Cape, SA & USA <sup>iii</sup>	Dombrow 2006b
<i>Diaplochelus longipes</i> (Fabricius, 1787)	<i>Protea repens</i> (L.)	A	Cape, SA	Coetzee & Giliomee 1985
<i>Eriesthis</i> Burmeister, 1844				
<i>E. stigmatica</i> (Billberg in Schönherr, 1817)	Maize	A	Fort Beaufort, SA	Omerod & Janson 1889
<i>E. stigmatica</i>	Carnation	A	Pretoria, SA	Annecke & Moran 1982
<i>E. stigmatica</i>	Roses, Carnations, Dahlias etc.	A	SA	Oberprieler <i>et al.</i> , 1990
<i>E. vestita</i> Burmeister, 1844	Dahlia	A	Pretoria, SA	Annecke & Moran 1982
<i>Heterochelus</i> Burmeister, 1844				
<i>H. connatus</i> Burmeister, 1844	Carnation	A	Pretoria, SA	Annecke & Moran 1982
<i>H. connatus</i>	Roses	A	SA	Annecke & Moran 1982
<i>H. rufimanus</i> (Castelnau, 1840)	Proteaceae ( <i>Protea</i> )	A	W. & S. Cape, SA	Myburgh & Rust 1975
<i>Heterochelus</i> sp.	Wattle ( <i>Acacia mearnsii</i> )	A	SA	Annecke & Moran 1982
<i>Heterochelus</i> sp.	Wattle ( <i>Acacia mearnsii</i> )	A	SA	Swain & Prinsloo 1986
<i>Heterochelus</i> sp.	Wattle	A	Southern Africa	Hepburn 1966
<i>Monochelus</i> Serville, 1825				

Taxon	Crop	Life stage/s	Locality	Reference/s
<i>M. calcaratus</i> Burmeister, 1844	Wattle	A	Southern Africa	Hepburn 1966
<i>M. calcaratus</i>	Black wattle	A	SA	Wang 1997
<i>M. calcaratus</i>	Wattle ( <i>Acacia mearnsii</i> )	A	SA	Swain & Prinsloo 1986
<i>Platychelus</i> Burmeister, 1844				
<i>Platychelus</i> sp.	Proteaceae ( <i>Protea</i> )	A	W. & S. Cape, SA	Myburgh & Rust 1975
<b>HOPLIINA Latreille, 1829</b>				
<i>Hoplia</i> Illiger, 1803				
<i>H. coffeae</i> Arrow, 1925	<i>Coffea</i> sp. Shoots & young leaves	A	Kenya	Arrow 1925
<i>H. sordida</i> Burmeister, 1844	Wattle	A	Southern Africa	Hepburn 1966
<i>H. sordida</i>	Wattle ( <i>Acacia mearnsii</i> )	A	SA	Annecke & Moran 1982
<i>H. sordida</i>	Wattle ( <i>Acacia mearnsii</i> )	A	SA	Swain & Prinsloo 1986
<b>PACHYDEMINI Burmeister, 1855</b>				
<i>Aegostheta</i> Dejean, 1833				
<i>A. cf. ciliata</i> Herbst, 1790	Turfgrass	L	Mossel Bay, SA	Omer-Cooper <i>et al.</i> , 1948
<i>Macrophylla</i> Hope, 1837				
<i>Macrophylla</i>	Turfgrass	L	E. Cape, SA	Smit 1964
<i>M. ciliata</i> (Herbst, 1790)	Turfgrass	L	Mossel Bay, SA	Omer-Cooper <i>et al.</i> , 1948
<i>M. ciliata</i> <sup>IV</sup>	Pineapple	L	SA	Annecke & Moran 1982
<i>M. ciliata</i>	Pineapple	L	SA	De Villiers <i>et al.</i> , 1987
<i>M. maritima</i> Blanchard, 1851	Turfgrass	L	Port Alfred, SA	Omer-Cooper <i>et al.</i> , 1948
<i>M. maritima</i>	Lawn grasses	L	SA	Annecke & Moran 1982
<i>M. nigricollis</i> Péringuey, 1904				
<i>M. pubens</i> Péringuey, 1904	Turfgrass	L	Humewood, SA	Omer-Cooper <i>et al.</i> , 1948
<i>M. pubens</i>	Turfgrass	L	Humewood, SA	Fenwick 1947
<i>Pseudachloa</i> Péringuey, 1904				
<i>P. leonina</i> Péringuey, 1904	Turfgrass	A	Pretoria, SA	At Schoeman (pers comm)
<b>MELOLONTHINI Leach, 1819</b>				



Taxon	Crop	Life stage/s	Locality	Reference/s
<i>Sparrmannia</i> Laporte, 1840				
<i>S. acicularis</i> Evans, 1989	Pistachio nut	A	Prieska, N. Cape, SA	Louw 2001
<i>S. flava</i> Arrow, 1917	Pistachio nut	A	Prieska, N. Cape, SA	Louw 2001
<i>S. transvaalica</i> Péringuey, 1904	Pistachio nut	A	Prieska, N. Cape, SA	Louw 2001
<b>SCHIZONYCHINA Burmeister, 1855</b>				
<i>Entyposis</i> Kolbe, 1894				
<i>E. impressa</i> Kolbe, 1894	Castor & caladiums	L	Zanzibar	Mansfield-Aders 1920
<i>E. impressa</i>	Clove	L	Zanzibar	Andre Moutia 1941
<i>E. impressa</i>	Sugarcane	L	Tanzania	Jepson 1956
<i>Hecistopsilus</i> Kolbe, 1894				
<i>H. molitor</i> Kolbe, 1894	Tobacco	A	S. Zimbabwe	Büttiker & Bünzli 1957
<i>Schizonycha</i> Dejean, 1833				
<i>S. affinis</i> Boheman, 1857	Sugarcane	L	South Africa	Carnegie 1974, 1988
<i>S. affinis</i>	Sugarcane	L	SA	Annecke & Moran 1982
<i>S. affinis</i>	Wattle	L	SA	Annecke & Moran 1982
<i>S. affinis</i>	Sugarcane	L	KZN Midlands, SA	Dittrich-Schröder <i>et al.</i> , 2009
<i>S. affinis</i>	Sugarcane	L	KZN Midlands, SA	Goble <i>et al.</i> , 2012
<i>S. cf. carbonaria</i> Boheman, 1857	Wattle	A	Southern Africa	Hepburn 1966
<i>S. citima</i> Péringuey, 1904	Tobacco	A	S. Zimbabwe	Büttiker & Bünzli 1957
<i>S. elegans</i> Péringuey, 1904	Orchards & vineyards	A	Transvaal, SA	Howard 1910
<i>S. fimbriata</i> Brenske, 1898	Wattle ( <i>Acacia mearnsii</i> )	A	SA	Swain & Prinsloo 1986
<i>S. fimbriata</i>	Soya	L	Bergville, KZN, SA	Hittersay 2005
<i>S. infantilis</i> Péringuey, 1904	Lawn grasses	L	SA	Annecke & Moran 1982
<i>S. plausibilis</i> Péringuey, 1904	Roses, Carnations, Dahlias etc.	A	SA	Oberprieler <i>et al.</i> , 1990
<i>S. plausibilis</i>	Lawn grasses	L	SA	Annecke & Moran 1982
<i>S. plausibilis</i>	Roses	A	SA	Annecke & Moran 1982
<i>S. profuga</i> Péringuey, 1904	Tobacco	A	S. Zimbabwe (Zimbabwe)	Büttiker & Bünzli 1957, Bünzli & Büttiker 1960a,b

Taxon	Crop	Life stage/s	Locality	Reference/s
<i>S. vastatrix</i> Paoli, 1934 <sup>v</sup>	Cotton	L	S. Somalia	Paoli 1934, Schmutterer 1964
<i>Schizonycha</i> sp.	Beans ( <i>Phaseolus vulgaris</i> L.)	L	Kenya	Medvecky <i>et al.</i> , 2006, 2007, 2009
<i>Schizonycha</i> sp.	Black wattle	A	SA	Wang 1997
<i>Schizonycha</i> sp. 18	Black wattle	L	Kenya	Elmer & Smith 1956
<i>Schizonycha</i> sp.	Clove	L	Zanzibar	Andre Moutia 1941
<i>Schizonycha</i> sp.	Dolichos Bean ( <i>Dolichos lablab</i> )	L	Sudan	Tarr 1958
<i>Schizonycha</i> sp.	Dura ( <i>Sorghum vulgare</i> )	A+L	Sudan	Pollard 1956
<i>Schizonycha</i> sp.	Grapevines	A	Cape, SA	Barnes <i>et al.</i> , 1986
<i>Schizonycha</i> sp.	Groundnuts ( <i>Arachis hypogaea</i> )	A+L	Potchefstroom, SA	Van Eeden <i>et al.</i> , 1991
<i>Schizonycha</i> sp.	Lubia ( <i>Dolichos lallab</i> )	L	Sudan	Pollard 1956
<i>Schizonycha</i> sp. [as <i>Phyllophaga</i> ] <sup>vi</sup>	Sorghum	A+L	Ethiopia	Tefera 2004
<i>Schizonycha</i> sp.	Tobacco	A	Zimbabwe	Mitchell 1946
<i>Schizonycha</i> sp.	Yams ( <i>Dioscorea</i> sp., family Dioscoreaceae)	L <sup>vii</sup>	Sub Saharan Africa	Korada <i>et al.</i> , 2010
<b>PEGYLINA Lacroix, 1989</b>				
<i>Pegylis</i> Erichson, 1847				
<i>P. sommeri</i> (Burmeister, 1855)	Black wattle	A	SA	Wang 1997
<i>P. sommeri</i>	<i>Eucalyptus camaldulensis</i>	A	SA	Swain & Prinsloo 1986
<i>P. sommeri</i>	<i>Eucalyptus globulus</i>	A	SA	Swain & Prinsloo 1986
<i>P. sommeri</i>	<i>Eucalyptus viminalis</i>	A	SA	Swain & Prinsloo 1986
<i>P. sommeri</i>	Fruit trees	A	SA	Carnegie 1974
<i>P. sommeri</i>	Nestle Factory	A <sup>viii</sup>	Estcourt, SA	Greg Davies pers comm 2005
<i>P. sommeri</i>	<i>Pinus patula</i>	A	Southern Africa	Hepburn 1966
<i>P. sommeri</i>	<i>Pinus patula</i>	A	SA	Swain & Prinsloo 1986
<i>P. sommeri</i>	Roses	A	SA	Annecke & Moran 1982
<i>P. sommeri</i>	Sugarcane	L	KZN Midlands, SA	Goble <i>et al.</i> , 2012
<i>P. sommeri</i>	Sugarcane	L	South Africa	Carnegie 1974
<i>P. sommeri</i>	Turf	L	SA	Carnegie 1974
<i>P. sommeri</i>	Vines	A	Eastern Cape, SA	Omerod & Janson 1889
<i>P. sommeri</i>	Wattle	L+A	Southern Africa	Hepburn 1966
<i>P. sommeri</i>	Wattle ( <i>Acacia mearnsii</i> De Wild Mimosaceae)	L+A	KZN Midlands, SA	Govender 2007

Taxon	Crop	Life stage/s	Locality	Reference/s
<i>P. sommeri</i>	Wattle ( <i>Acacia mearnsii</i> )	A	SA	Swain & Prinsloo 1986
<i>P. sommeri</i>	Pineapple	L	Eastern Cape, SA	Smith <i>et al.</i> , 1995
<i>P. sommeri</i>	Sugarcane	L	SA	Carnegie 1988
<i>P. sommeri</i>	Sugarcane	L	SA	Annecke & Moran 1982
<i>P. sommeri</i>	Pine	A	SA	Annecke & Moran 1982
<i>P. sommeri</i>	Litchis <sup>ix</sup>	L	SA	Annecke & Moran 1982
<i>P. sommeri</i>	Pineapple	L	SA	Annecke & Moran 1982
<i>P. sommeri</i>	Roses, Carnations, Dahlias etc.	A	SA	Oberprieler <i>et al.</i> , 1990
<i>P. sommeri</i> <sup>x</sup>	Pineapple	L	SA	De Villiers <i>et al.</i> , 1987
<i>P. sommeri</i> [as <i>Hypopholis</i> ]	Sugarcane	L	KZN Midlands, SA	Dittrich-Schröder <i>et al.</i> , 2009
<i>P. vittata</i> (Fåhraeus in Boheman, 1857)	Wattle	A	Southern Africa	Hepburn 1966
<i>P. vittata</i>	Wattle ( <i>Acacia mearnsii</i> )	A	SA	Swain & Prinsloo 1986
<i>P. vittata</i>	Sugarcane	L	Swaziland	Sweeney 1967
<b>LEUCOPHOLINA Burmeister, 1855</b>				
<i>Afrolepis</i> Decelle, 1968 <sup>xi</sup>				
<i>A. pygidialis</i> (Brenske, 1903)	<i>Coffea excelsa</i>	L	Cameroon	Lepesme & Paulian 1944
<i>Asthenopholis</i> Brenske, 1898				
<i>A. subfasciata</i> (Blanchard, 1850)	Pineapple	L	Eastern Cape, SA	Smith <i>et al.</i> , 1995
<i>A. subfasciata</i>	Sugarcane	L	SA	Carnegie 1988
<i>A. subfasciata</i> <sup>xii</sup>	Sugarcane	L	Swaziland (Mhlume)	Sweeney 1967
<i>A. minor</i> Brenske, 1898 <sup>xiii</sup>	Sugarcane	L	SA & Swaziland	Harrison 2009
<i>Cochliotis</i> Kolbe, 1894				
<i>C. melolonthoides</i> (Gerstaecker, 1867)	Sugarcane	L	Tanzania	Jepson 1956
<i>C. melolonthoides</i>	Sugarcane	L	Tanzania	Carnegie 1974
<i>Eulepida</i> Kolbe, 1894				
<i>E. mashona</i> (Arrow, 1902) = <i>Lepidiota mashona</i> Arrow, 1902	<i>Arachis hypogaea</i> (groundnut)	L	Africa	Colwill 1967 <sup>xiv</sup> , Feakin 1973 <sup>xiv</sup>
<i>E. mashona</i>	Wattle	L+A	S. Zimbabwe	Hepburn 1966
<i>E. mashona</i>	Black wattle	A	SA	Wang 1997
<i>E. mashona</i>	Wattle ( <i>Acacia mearnsii</i> )	A	SA	Swain & Prinsloo 1986

Taxon	Crop	Life stage/s	Locality	Reference/s
<i>E. mashona</i>	Sugarcane	L	Zimbabwe	Carnegie 1974
<i>E. mashona</i>	Ley pasture ( <i>Chloris gayana</i> (Kunth.))	L	Zimbabwe	Taylor & Smithers 1959
<i>E. mashona</i>	Maize	L	N. Zimbabwe	Wilson 1963, 1972
<i>E. mashona</i>	As a fried or sundried food source	A	Zimbabwe	Gomez 1988
<b>MELOLONTHINA Leach, 1819</b>				
<i>Psilonychus</i> Burmeister, 1855				
<i>P. groendahli</i> (Billberg, 1820)	Wattle	A	Southern Africa	Hepburn 1966
<i>P. groendahli</i>	Wattle ( <i>Acacia mearnsii</i> )	A	SA	Swain & Prinsloo 1986
<i>P. cf. groendahli</i> (Billberg, 1820)	Wattle	A	Southern Africa	Hepburn 1966
<b>SERICINI Kirby, 1837</b>				
<i>Autoserica</i> Brenske, 1897 <sup>xv</sup>				
<i>A. tessellata</i> Péringuey, 1904 <sup>xvi</sup>	Wattle	A	Southern Africa	Hepburn 1966
<i>A. tessellata</i> (Péringuey, 1904)	Wattle ( <i>Acacia mearnsii</i> )	A	SA	Swain & Prinsloo 1986
<i>Aserica</i> Lewis, 1895 <sup>xvii</sup>				
<i>Aserica variegata</i> Arrow, 1917	Cacao young leaves	A	DRC	Arrow 1917
<i>Aserica cf. lucidula</i> (Péringuey, 1904) <sup>xviii</sup>	Black wattle	L	Kenya	Elmer & Smith 1956
<i>Lepiserica</i> Brenske, 1900 <sup>xix</sup>				
<i>L. cf. lucidula</i> (Péringuey, 1892)	Black wattle	L	Kenya	Elmer & Smith 1956
<i>Neoserica</i> Brenske, 1984				
<i>N. carneola</i> Péringuey, 1892 <sup>xx</sup>	Wattle	A	Southern Africa	Hepburn 1966
<i>N. carneola</i>	Wattle ( <i>Acacia mearnsii</i> )	A	SA	Swain & Prinsloo 1986
<i>Pseudotrochalus</i> Quedenfeldt, 1884				
<i>P. concolor</i> Kolbe, 1883	<i>Bixa orellana</i>	A	S. Nigeria	Golding 1937
<i>P. concolor</i>	Cacao	A	DRC	Arrow 1917
<i>P. concolor</i>	<i>Citrus acida</i>	A	S. Nigeria	Golding 1927
<i>P. concolor</i>	<i>Citrus limonia</i>	A	S. Nigeria	Golding 1937

Taxon	Crop	Life stage/s	Locality	Reference/s
<i>P. concolor</i>	<i>Gossypium hirsutum</i>	A	S. Nigeria	Golding 1927
<i>P. concolor</i>	<i>Haematoxylon campechianum</i>	A	S. Nigeria	Golding 1927
<i>P. concolor</i>	<i>Quisqualis indica</i>	A	S. Nigeria	Golding 1927
<i>P. concolor</i>	<i>Spondias lutea</i>	A	S. Nigeria	Golding 1927
<i>P. concolor</i>	<i>Zea mais</i>	A	S. Nigeria	Golding 1927
<i>Triodonta</i> Mulsant, 1842 <sup>xxi</sup>				
<i>T. procera</i> Lansberg, 1886	Cacao	A	DRC	Arrow 1917
<b>TROCHALINA Brenske, 1898</b>				
<i>Trochalus</i> Laporte, 1832				
<i>T. aeruginosus</i> Burmeister, 1855 <sup>xxii</sup>	<i>Hibiscus</i>	A	Swaziland	Sweeney 1967
<i>T. aeruginosus</i>	Roses	A	Swaziland	Sweeney 1967
<i>T. byrrhinus</i> Fåhræus in Boheman, 1858	Wattle ( <i>Acacia mearnsii</i> )	A	SA	Swain & Prinsloo 1986
<i>T. byrrhinus</i>	Wattle	A	Southern Africa	Hepburn 1966
<i>T. carinatus</i> Gyllenhal in Schönherr, 1817				
<i>T. carinatus</i>	Avocado	A	Sierra Leone	Hargreaves 1937
<i>T. fulgidus</i> Fåhræus in Boheman, 1857	Wattle ( <i>Acacia mearnsii</i> )	A	SA	Swain & Prinsloo 1986
<i>T. fulgidus</i>	Wattle	A	Southern Africa	Hepburn 1966
<i>T. gibbus</i> (Fabricius, 1781)	<i>Cola nitida</i>	A	Sierra Leone	Hargreaves 1937
<i>T. gibbus</i>	Cotton	A	Sierra Leone	Hargreaves 1937
<i>T. gibbus</i>	Avocado	A	Sierra Leone	Hargreaves 1937
<i>T. pilula</i> Klug, 1835	<i>Cola nitida</i>	A	Sierra Leone	Hargreaves 1937
<i>T. pilula</i>	Avocado	A	Sierra Leone	Hargreaves 1937
<i>T. politus</i> Moser, 1919	Pineapple	L	Eastern Cape, SA	Smith <i>et al.</i> , 1995
<i>T. politus</i>	Pineapple	L	Eastern Cape, SA	De Villiers <i>et al.</i> , 1987
<i>T. politus</i>	Pineapple	L	SA	Annecke & Moran 1982
<i>Trochalus</i> sp.	Cacao bean ( <i>Theobroma cacao</i> L.)	A	S. Nigeria	Peacock 1913

**Table 1 supplementary material.** Additional explanatory endnotes.

- i. Misidentified as *Adoretus tessulatus* (Rutelinae) in Petty's earlier papers.
- ii. Accidentally exported to Poland with *Protea* flowers.

- iii. Accidentally exported to the USA with *Protea* flowers.
- iv. An incorrect identification has placed this species name in the literature as a pest of pineapple; the correct species identification is *Pegylis sommeri* (see Harrison, 2014b). Petty (1978) provides a clear photograph of *P. sommeri*, but identified as *Macrophylla ciliata*.
- v. Schmutterer (1964) refers to this species as *S. vastatrix* Chiar. [Chiaromonte]. Paul Schoolmeesters (pers. comm.) drew our attention to its first description in Paoli (1934); no mention is made to this species in Pope (1960) or Lacroix (2010).
- vi. To a non-specialist the genera *Phyllophaga* and *Schizonycha* are morphologically similar. The former is not known from Africa, while the latter is (see Pope 1960).
- vii. Larvae feed on both yam roots and tubers.
- viii. Adult beetles attracted to factory security lights at night, and then entered the factory where foodstuffs are being produced and risked being packaged with these foodstuffs.
- ix. Anneck & Moran (1982: 79) list *Macrophylla* as being a pest of Litchis, but if one looks at the geographic distribution of where Litchis were grown at that time (Lowveld and KZN South Coast) it is far more likely to have been a species of *Pegylis*.
- x. Incorrectly referred to as *Macrophylla ciliata* (Herbst, 1790).
- xi. *Afrolepis* Decelle, 1968 (replacement name for preoccupied *Oligolepis* Brenske, 1903).
- xii. Sweeney (1967) includes *A. subfasciata* for Swaziland (Mhlume). After revising *Asthenopholis* Harrison (2009) suggests that the Swaziland (Mhlume) specimens are most likely attributable to *A. minor*.
- xiii. "Williams (1985) records *A. minor* (as *A. subfasciata*) to have a limited distribution in the Nokwane (Mhlume) and S.I.S. (Swaziland Irrigation Scheme) sugarcane farms in Swaziland. Carnegie (1988) mentions *A. minor* and *A. subfasciata* as sugarcane pests in Swaziland and Emoyeni, KwaZulu–Natal. White grubs of *A. subfasciata* (Anonymous 1992) are reported from sugarcane fields from the Mhlume area (Vuvulane) in Swaziland. However, the present study establishes that these are misidentifications. *Asthenopholis subfasciata* does not occur in Swaziland or KwaZulu–Natal (i.e. Port Natal = Durban), save for a few old and questionable records. *Asthenopholis minor* is more likely to be a sporadic pest of sugarcane in these regions." Harrison (2009).
- xiv. Literature not seen in person but quoted in van Eeden *et al.* (1991).
- xv. Currently valid genus name is *Maladera* Mulsant & Rey, 1871 (pers. comm. Dirk Ahrens).

- xvi. Swain & Prinsloo (1986) list this species as *Maladera tessellata* (Péringuey), but we follow Dalla Torre (1912a) and include it here as *Autoserica tessellata* Péringuey, 1904.
- xvii. Currently valid genus name is *Maladera* Mulsant & Rey, 1871 (pers. comm. Dirk Ahrens).
- xviii. This species was originally described in *Serica* (pers. comm. Dirk Ahrens).
- xix. The taxonomic position of *Lepiserica* in relation to *Maladera* requires investigation (pers. comm. Dirk Ahrens).
- xx. Swain & Prinsloo (1986) list this species as *Maladera carneola* (Péringuey), but we follow Dalla Torre (1912a) and include it here as *Neoserica carneola* Péringuey, 1892.
- xxi. Currently valid genus name is *Triodontella* Reitter, 1919 (pers. comm. Dirk Ahrens).
- xxii. Spelt as *T. aerugineus* in Sweeney (1967), but as *T. aeruginosus* in Dalla Torre (1912a).

**Table 1 supplementary material.** Additional references not cited in the review article but referred to in this online table.

- Barnes, B.N., Daiber, K.C. & Donaldson, J.M.** (1986) Fruit eating and leaf-eating beetles. pp. 35-42 in Myburgh, A.C. (Editor) 1986. *Crop pests in southern Africa. Vol. 1 Deciduous fruit, grapes and berries*. Bulletin **407**, Plant Protection Research Institute, Pretoria. 1-92.
- Bünzli, G.H. & Büttiker, W.W.** (1960a) Fungous diseases of lamellicorn larvae in southern Rhodesia. *Bulletin of Entomological Research* **50**, 89-96.
- Bünzli, G.H. & Büttiker, W.W.** (1960b) A comparative study of the soil conditions favored by certain melolonthid and tenebrionid pests of tobacco in southern Rhodesia. *Bulletin of Entomological Research* **50**, 259-263.
- Colwill, D.J.** (1967) *Pest control in groundnuts*. PANS Manual No. **2**, 24-54.
- de Villiers, E.A., van den Berg, M.A. & Petty, G.T.** (1987) Borers and root feeders. pp. 91-96 in Myburgh, A.C. 1987. *Crop pests in southern Africa. Vol. 2 Citrus and other subtropicals*. Bulletin **411**. Plant Protection Research Institute, Pretoria. 1-118.
- Elmer, J.L. & Smith, R.G.** (1956) Some pests of Black Wattle in Kenya: with a list of other insects inhabiting the plantations. *East African Agricultural Journal* **21**, 230-246.
- Feakin, S.D.** (1973) *Pest control in groundnuts*. PANS Manual No. **2**. pp. 1-197.
- Govender, P.** (2007) Status of seedling establishment pests of *Acacia mearnsii* De Wild. (Mimosaceae) in South Africa. *South African Journal of Science* **103**, 141-147.
- Hepburn, G.A.** (1966) A revised list of wattle insects and spiders of southern Africa. *Wattle Research Institute Report* 67-83.

- Howard, C.W.** (1910) Cockchafers and flower beetles. *Transvaal Agricultural Journal* **6**, 1-4.
- Mitchell, B.L.** (1946) White grub control in tobacco lands. *The Rhodesia Agricultural Journal* **43**(6), 499-504.
- Oberprieler, R.G., Donaldson, J.M., Daiber, K.C., Coetzee, J.H., de Villiers, E.A., Rust, D.J. & Myburgh, A.C.** (1990) Flower- and leaf-eating beetles. pp. 21-27 in Myburgh, A.C. 1990. *Crop pests in southern Africa. Vol. 5 Flowers and other ornamentals*. Bulletin **419**. Plant Protection Research Institute, Pretoria. 1-98.
- Pollard, D.G.** (1956) The control of chafer grubs (*Schizonycha* sp., Coleoptera, Melolonthinae) in the Sudan. *Bulletin of Entomological Research* **47**(3), 347-360.
- van Eeden, C.F., van Rensburg, J.B.J. & de K. van der Linde, T.C.** (1991) A check list of insects occurring in groundnut (*Arachis hypogaea*) fields in the Highveld Region. *Phytophylactica* **23**, 255-259.
- Wang, H.** (1997) Chapter 5. Black wattle plantations in South Africa: Protection. pp. 39-53 in *Black Wattle and its Utilisation - Abridged English Edition*. Edited for the Rural Industries Research & Development Corporation by Brown, A.G. & Ho Chin Ko. RIRDC Publication No. 97/72.



**Table 2 supplementary material.** Summary of all African Melolonthinae (pest and non-pest) in which larval instars are described and illustrated. Larval head capsule widths are listed as follows: range (mm), mean (mm), sample size in brackets. All these descriptions include an illustration of the raster (necessary for identification). Locality abbreviations include: South Africa (SA), North (N), South (S), East (E) and West (W). **Table 1** (included in the published *Bulletin of Entomological Research* review paper) includes all appropriate additional taxonomic information.

Taxon	Locality	First instar	Second instar	Third instar	Reference
<i>Asthenopholis cf. minor</i> <sup>i</sup>	Swaziland	-	-	4.8-5.1 (30)	Sweeney 1967
<i>Asthenopholis subfasciata</i>	E. Cape, SA	-	-	5.1-5.8 (10)	Oberholzer 1959a
<i>Asthenopholis subfasciata</i>	E. Cape, SA	-	-	mean 5.6 (5)	Smith <i>et al.</i> , 1995
<i>Camenta innocua</i>	Johannesburg, SA	-	-	3.03-3.87 (10)	Oberholzer 1959a
<i>Cochliotis melolonthoides</i>	Tanzania	-	4.6 (n?)	7.0-7.8 (n?)	Jepson 1956
<i>Congella valida</i> <sup>ii</sup>	E. Cape, SA	1 <sup>st</sup> instar overlooked	2.7-2.8; mean 2.75 (n?)	4.2-4.3; mean 4.25 (n?)	Petty 1977b, 1978
<i>Congella valida</i>	E. Cape, SA	-	mean 2.6 (7)	-	Smith <i>et al.</i> , 1995
<i>Monochelus calcaratus</i>	KwaZulu-Natal, SA	0.68-0.80 (n?)	-	-	Prins 1965
<i>Oligolepis pygidialis</i>	Cameroon	-	-	-	Lepesme & Paulian 1944
<i>Pegylis sommeri</i> <sup>iii</sup>	KwaZulu-Natal, SA	-	-	4.4-5.0 (10)	Oberholzer 1959a
<i>Pegylis sommeri</i> <sup>iv</sup>	KwaZulu-Natal, SA	1.44-1.70 (n?)	1.87 (n?)	-	Prins 1965
<i>Pegylis sommeri</i> <sup>v</sup>	E. Cape, SA	-	-	mean 4.8 (3)	Smith <i>et al.</i> , 1995
<i>Cf. Pegylis vittata</i>	Swaziland	-	-	4.7-5.1 (15)	Sweeney 1967
<i>Sparrmannia flava</i>	N.W. Cape, SA	1.66-1.74; 1.7 (8)	2.8-3.2; 3.0; (9)	4.0-4.3; 4.2 (5)	Scholtz 1988
<i>Sparrmannia flava</i>	N.W. Cape, SA	-	max width 3.7 (3)	-	Evans 1989
<i>Trochalus politus</i>	E. Cape, SA	1.2-1.3; mean 1.25 (n?)	1.8-1.9; mean 1.85 (n?)	2.8-2.9; mean 2.85 (n?)	Petty 1977b, 1978
<i>Trochalus politus</i>	E. Cape, SA	-	-	mean 2.9 (8)	Smith <i>et al.</i> , 1995

**Table 2 supplementary material.** Additional explanatory endnotes.

- i. Identified as *A. subfasciata* in Sweeney (1967), but following Harrison (2009) only *A. minor* is known to occur in Swaziland.

- ii. Originally incorrectly identified as *Adoretus tessulatus* (see Petty 1977b, 1982, 1990), later corrected to *Congella valida* (see Smith *et al.*, 1995, Petty 2001).
- iii. As *Hypopholis sommeri*, now *Pegylis sommeri* (see Harrison, 2014b).
- iv. As *Hypopholis sommeri*, now *Pegylis sommeri* (see Harrison, 2014b).
- v. Due to an incorrect identification these larvae were described as *Macrophylla ciliata*, but following Harrison (2014b) the species involved is *Pegylis sommeri*.