


BMJ Open Impacts of economic inequality on healthcare worker safety at the onset of the COVID-19 pandemic: cross-sectional analysis of a global survey

Sean P Harrigan,¹ Vivian W L Tsang,² Annalee Yassi,¹ Muzimkhulu Zungu,^{3,4} Jerry M Spiegel ¹

To cite: Harrigan SP, Tsang VWL, Yassi A, *et al*. Impacts of economic inequality on healthcare worker safety at the onset of the COVID-19 pandemic: cross-sectional analysis of a global survey. *BMJ Open* 2022;**12**:e064804. doi:10.1136/bmjopen-2022-064804

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2022-064804>).

Received 17 May 2022
Accepted 06 September 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY. Published by BMJ.

¹Global Health Research Program, School of Population and Public Health, The University of British Columbia, Vancouver, British Columbia, Canada

²Faculty of Medicine, The University of British Columbia, Vancouver, British Columbia, Canada

³National Institute for Occupational Health, Johannesburg, Gauteng, South Africa

⁴School of Health Systems and Public Health, University of Pretoria Faculty of Health Sciences, Pretoria, Gauteng, South Africa

Correspondence to
Dr Jerry M Spiegel;
jerry.spiegel@ubc.ca

ABSTRACT

Objectives To assess the extent to which protection of healthcare workers (HCWs) as COVID-19 emerged was associated with economic inequality among and within countries.

Design Cross-sectional analysis of associations of perceptions of workplace risk acceptability and mitigation measure adequacy with indicators of respondents' respective country's economic income level (World Bank assessment) and degree of within-country inequality (Gini index).

Setting A global self-administered online survey.

Participants 4977 HCWs and healthcare delivery stakeholders from 161 countries responded to health and safety risk questions and a subset of 4076 (81.2%) answered mitigation measure questions. The majority (65%) of study participants were female.

Results While the levels of *risk* being experienced at the pandemic's onset were consistently deemed as unacceptable across all groupings, participants from countries with less income inequality were somewhat less likely to report unacceptable levels of risk to HCWs regarding both workplace environment (OR=0.92, p=0.012) and workplace organisational factors (OR=0.93, p=0.017) compared with counterparts in more unequal national settings. In contrast, considerable variation existed in the degree to which *mitigation* measures were considered adequate. Adjusting for other influences through a logistic regression analysis, respondents from lower middle-income and low-income countries were comparatively much more likely to assess both occupational health and safety (OR=10.91, p<0.001) and infection prevention and control (IPC) (OR=6.61, p=0.001) protection measures as inadequate, despite much higher COVID-19 rates in wealthier countries at the time of the survey. Greater within-country income inequality was also associated with perceptions of less adequate IPC measures (OR=0.94, p=0.025). These associations remained significant when accounting for country-level differences in occupational and gender composition of respondents, including specifically when only female care providers, our study's largest and most at-risk subpopulation, were examined.

Conclusions Economic inequality threatens resilience of health systems that rely on health workers working safely to provide needed care during emerging pandemics.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ A major strength of the study is its novel empirical testing of the 'income inequality' hypothesis for a comparative cross-country analysis of a major global health challenge: protection for a workforce central to the provision of healthcare services during a pandemic.
- ⇒ This study is based on a unique global self-administered online survey conducted by a network of occupational health experts coordinated by WHO through a large array of professional networks and social media.
- ⇒ A major limitation of the study is its character as a convenience sample with different compositions by gender and occupation among countries and small sample sizes in some countries; however, access to gender and occupation identifiers of respondents has enabled adaptive strategies to take this into consideration.
- ⇒ The study is exploratory in considering associations with economic inequality, but does not provide a way to consider pathways for this effect, so further research will be needed for this.

INTRODUCTION

As the COVID-19 pandemic emerged, attention was quickly drawn to risks faced by front-line healthcare providers^{1 2}—and the urgent need to strengthen their protection.^{3 4} By September 2020, it was estimated that 10% of global infections had been in healthcare workers (HCWs), and over 7000 had died.^{5 6} Notwithstanding inconsistent reporting, Papoutsis and colleagues, in reviewing the global burden of COVID-19 for HCWs by country,^{6 7} estimated the percentage of HCW cases among the total cases by April 2020 as ranging from less than 1% in Hong Kong and India, to 19% in Spain.

Despite 60 million people employed in the healthcare sector worldwide,^{8 9} a global shortage of HCWs persists and is especially

critical in low and middle-income countries (LMICs),¹⁰ where the greatest global burden of disease exists.¹¹ Risk mitigation is marked by considerable variation,¹² with shortcomings in infrastructure and mitigation programmes contributing to higher burdens of disease and HCW risk in more poorly resourced settings.¹³ The danger that HCWs face of acquiring COVID-19 adds to extensive existing risks in infectious disease endemic states, for example, with tuberculosis in sub-Saharan Africa.¹⁴

While lack of personal protective equipment (PPE) was highlighted early in the COVID-19 pandemic, consideration of broader occupational health and safety (OHS) factors and mitigation measures attracted less initial attention.^{7 8 15} To ascertain the extent of OHS risk exposure and the adequacy of mitigation measures in place to meet the challenge of COVID-19, an ad hoc expert group of WHO, the International Labour Organization (ILO) and the International Commission on Occupational Health prepared and circulated a questionnaire survey to identify HCW perceptions of the most common threats to their health and safety as well as the adequacy of mitigation measures in the emerging pandemic.¹⁶

Further to a preliminary analysis of survey results¹⁶ regarding risk and adequacy of protection, we sought to ascertain the degree to which perceived risk exposure of HCWs and adequacy of mitigation measures is associated with a country's economic characteristics. Considerable attention, after all, has been given to the impact of economic disparity on health,^{17 18} especially in relation to Wilkinson's 'economic inequality hypothesis' suggesting that greater inequality is associated with poorer health.¹⁹ In recognition that 'the traditional exposure-disease framework used in occupational health research is not equipped to address societal contexts in which work is embedded',²⁰ we sought to examine how such driving forces²¹ as a country's economic inequality might be affecting the well-being of HCWs.

A variety of factors have been examined that might have influenced how the onset of the COVID-19 pandemic was experienced in different national settings, including consideration of cultural traits,²² specific government regulations and non-pharmaceutical interventions^{23 24} and political leadership characteristics.²⁵ Our study sought to apply a cross-country perspective to consider the effects of economic inequality, recognising this to be a dimension of considerable relevance in global public health research.

With this focus, we set out to first consider variation in perceptions of the acceptability of work-related risks and the adequacy of mitigation measures that were being experienced by HCWs as COVID-19 emerged; and, second, to determine the extent to which variations were associated with a country's comparative income level and degree of income inequality.

METHODS

Survey development

Shortly after *WHO Ad Hoc Study Group on Health and Safety of Health Workers* was established when the COVID-19 pandemic emerged, it created an online survey aimed at HCWs from all WHO regions globally. In addition to the capture of demographic indicators of respondents, the survey contained 41 questions—17 on health and safety risks and 24 on mitigation measures¹⁶ (online supplemental appendix). Risk questions were grouped into those related to infectious disease transmission, physical work environment, psychological work environment and work organisation. For each risk question, participants were asked, '*Think about the working conditions of health workers in your country, jurisdiction or health facility....; rate the current level of these risks, now during the COVID-19 pandemic*'. Questions regarding mitigation measures were divided into two groups: OHS and infection prevention and control (IPC). Here, participants were similarly asked: '*Think about the working conditions of health workers in your country, jurisdiction or health facility...rate the level of application of these measures according to your knowledge of the real situation now during the COVID-19 pandemic*'.

Patient and public involvement

The participation of health workers (whose well-being is the focus for this study in relation to their assessment of the adequacy of measures to protect them) was indirectly included through the participation of their representatives (unions within the ILO and other professional bodies) who were directly involved in the creation of the research instrument and in the dissemination of the online survey and its initial results.

Study population and inclusion criteria

Participants were recruited by convenience sampling, with dissemination through a large array of professional networks and social media. The survey, self-administered online to enable rapid low-cost recruitment, was available in Arabic, Chinese, English, French, German, Italian, Portuguese, Russian, Spanish and Swahili. A range of HCWs and stakeholders involved with healthcare delivery were invited to participate. In addition to HCWs in direct patient care in both formal and informal settings and in public and private facilities, respondents also included allied health and supporting staff, including OHS and IPC professionals, administration, management, drivers, public health workers, community health workers and others as defined by the International Standard Classification of Occupations 2008. Data collection occurred between 5 May and 25 June 2020. Participant results were excluded if they failed to complete demographic questions or if they failed to provide any responses to the risk and mitigation questions. As the survey was designed to be completed and submitted anonymously, no formal request for signed consent was solicited, with participants' submission itself indicating consent to use the information provided as anonymised aggregated

data. This work was supported by the International Development Research Centre under grant M20-00559 and the Canadian Institutes of Health Research under grant vs1-175519 for the ‘Protecting healthcare workers from COVID-19: a comparative contextualized analysis’ research programme.

Independent variables

Demographic information for individual survey respondents was collected on country, gender and occupation—the latter separated into 13 categories and then grouped into patient care/health services; specialised technical support; clerical support/administration and management; and other. Details about the study sample population composition and demographic characteristics of participants are presented in online supplemental figure S1; Table S1, respectively.

Our research group, drawn from two WHO collaborating centres participating in the survey process, conducted the analysis by consolidating respondents by their home country and then linking this to a WHO geographic region²⁶; a comparative country-level economic classification by World Bank income groups based on the annual Atlas gross national income per-capita estimates^{26 27}; and the country’s Gini index—a measure used for the analysis of income inequality present within a country, with a score of 0 representing perfect equality, and a score of 1 representing complete inequality.²⁸ Data for Gini and economic classification were taken from the World Bank, using the most recent data available. To take account of the variation across regions present during the initial phase of the pandemic, we also considered COVID-19 incidence per million (logarithmic scale) in each country at the time when the survey was completed, as an indicator of the intensity as of a particular date, using values for June 2020 drawn from ‘Our World in Data’ database.²⁹

Dependent variables

Acceptability of workplace risks and adequacy of mitigation measures—the dependent variables in this study—were derived from a factor analysis of individual survey responses, then aggregated to enable subsequent analysis of the effect of country-level characteristics. Factor analysis¹⁶ was used to reduce the 41 survey questions into coherent groupings and principal component analysis with varimax rotation carried out to create factors from each set of workplace health and safety risk exposure questions (corresponding to workplace risk and workplace organisation acceptability) and mitigation measure questions (corresponding to IPC and OHS adequacy); online supplemental table S1 summarises the subject matter covered by the questions consolidated in each factor. Separate factor analyses were run on risk questions and preventive measure questions. Missing values were excluded in a listwise fashion. The rotated component matrix was used to identify factors. To measure scale reliability, Cronbach’s alpha was used for each individual factor. Scores over 0.7 are considered to be acceptable for

internal consistency.³⁰ The results from the factor analysis are outlined further in our preliminary analysis.¹⁶

The questions were administered as a 3-point Likert scale, then converted to a 10-point scale for clearer communication (ie, midpoint of 2 becoming 5). Numerical scores were assigned to each answer to establish a scale for both the risk and mitigation measure factors, with higher scores corresponding to more desirable states. For health and safety risks, a score of 0 was assigned to ‘risk is not acceptable at all’; 5 to ‘risk is acceptable for a short time’; and 10 for ‘risk is negligible’. For mitigation measures, a score of 0 was assigned to ‘does not exist at all’; 5 to ‘exists and offers some protection’; and 10 to ‘exists and offers full protection’. Responses of ‘don’t know/unsure’ were assigned blanks. Factor scores were then calculated to form an individual respondent’s factor score for each of the four groupings, that is, work environment risk acceptability, work organisation risk acceptability, OHS adequacy and IPC adequacy, and then aggregated to generate a mean value for each country’s respondents, so that intercountry comparison could be conducted. The higher the scores, the greater the perceived adequacy of mitigation measures or acceptability of risk deemed as being experienced.

Analysis

The mean country dependent variable factor scores derived from the aggregation of individual participants’ responses served as the basis for considering associations by WHO region, economic classification, Gini coefficients and COVID-19 incidence. Comparisons of survey mean scores were carried out using analysis of variance (ANOVA), with an alpha of 0.05 used to test significance. To compare means for the continuous variable Gini coefficient and COVID-19 incidence scores, we ordinarily divided groups of countries into quartiles by values.

To ensure that intercountry variation was not purely explained by possible gender and occupational compositional differences among a particular country’s respondents, we carefully examined possible sources of discrepancy (online supplemental table S3), using ANOVA to consider effects that could complicate the cross-country comparison of all respondents. To minimise any such effect, we considered different ways to stratify our analysis of the study population, notably by focusing only on those populations that had the most direct workplace experience to personally being ‘at risk’. Noting the presence of gender differences among patient care deliverers, we specifically isolated female respondents, who in fact constituted the largest demographic group of respondents in the study, representing 1998 respondents from 112 countries (n=1968 from 112 countries), the largest subpopulation.

Finally, to measure the effect that the interaction of independent variables had on the likelihood of workplace risks being considered as acceptable in a country setting as COVID-19 was emerging, and workplace protection and control measures being deemed as adequate, we



created and applied a logistic regression model. Preferred outcomes for this analysis were assessed as mean factor scores ≥ 5 , corresponding to assessments that mitigation 'exists and offers some protection' or better; or 'risk is acceptable for a short time' or better. All statistical analyses were done using R and SPSS Statistics software (version 25).^{31 32}

RESULTS

Overall study population and survey responses

There were 4977 participants who responded to health and safety risk questions and a subset of 4076 (81.2%) who answered mitigation measure questions. The majority of study participants were female (65%), reflecting the make-up of the health sector workforce. Most participants were from the European region (35%), followed by the Americas (31%), the Western Pacific region (15%) and Africa (10%); the South East Asian (4%) and Eastern Mediterranean regions (3%) made up the smallest proportion of participants. In total, there were 161 countries represented in the survey. Portugal (n=549, 11%), USA (n=451, 9%), Brazil (n=373, 7%), Canada (n=263, 5%) and China (n=233, 5%) had the most participants. The majority of respondents were from countries of high-economic classification (59%), followed by upper middle (27%), lower middle (10%) and low (4%). Most survey participants worked for a health services employer (61%), followed by government services (15%) and businesses and farms (10%). Those working in academia, professional associations, international organisations and non-government organisations each encompassed less than 10%. Finally, the type of occupation was predominantly patient care/health (56%) services, followed by 29% providing technical services such as IPC or OHS specialists, 7% in administration and 10% identified as working in other sectors (online supplemental table S1).

The largest percentage of countries was in Europe (30%) and over a third of all countries were high-income

countries (high income countries (HICs), 35%). The average Gini index was 37.8 (SD=7.7) and the mean and median COVID-19 incidence rates per million were 1360 and 278, respectively, at the time the survey was conducted.

Table 1 illustrates that considerable variation exists in these variables across the different WHO regions, indicating the distinct characteristics and conditions present at the onset of the pandemic. It is especially noteworthy that case levels had been far greater in high-income country areas at the survey midpoint (1 June 2020). For example, cases per million were 2525 in Europe versus 119 in Africa; 5408 in the USA; and only 138 in India and 97 in Indonesia.

As summarised in table 2 (full table in online supplemental table S4), the majority of respondents designated most of the health and safety risk parameters as 'not acceptable at all'. Circumstances most reported as such included bullying or psychological harassment in the workplace (54%), physical violence and assaults (54%), exposure to blood, bodily fluids and other infectious materials (52%), inadequate sanitation facilities (52%) and sexual harassment (50%). In contrast, areas such as time pressure and high workload (38%), skin damage from PPE (33%) and shift work with night shifts (23%) were deemed to be less of a concern. There were no risk categories in which the most common response was 'risk is negligible'.

Mitigation measures related to the above areas of concern were seen as particularly lacking, with only the category of 'policies for facilities for hand hygiene' designated as 'exists and offers full protection' (full table in online supplemental table S5). For example, despite psychosocial-related risks, including bullying, harassment, physical violence and sexual harassment, ranked consistently high (54%, 54% and 50%, respectively), only 21% indicated that corresponding policies 'exist and offer full protection', with similar dissatisfaction for the adequacy

Table 1 Country characteristics of different WHO regions

Region	Countries (n)	Countries by income classification*				Mean country values		Study population characteristics	
		High	Upper middle	Lower middle	Low	Inequality	COVID-19	Gender	Occupation
						Gini coefficient*	Cases per million†*	Female* (%)	Front-line patient care*(%)
Overall	161	57	42	36	26	37.8	1360	65.5	56.4
AFRO	37	0	6	12	19	43.2	119	44.8	52.1
EMRO	20	6	3	7	4	35.2	2407	39.8	29.4
EURO	48	32	13	2	1	31.8	2525	68.2	64.4
PAHO	30	11	14	4	1	44.8	1135	73.3	46.5
SEARO	9	0	2	6	1	35.0	86	36.7	56.3
WPRO	17	8	4	5	0	37.0	512	70.7	68.8

Full listing of WHO region countries at https://en.wikipedia.org/wiki/List_of_WHO_regions

*P<0.001.

†COVID-19 rates as of June 2020.

AFRO, Africa; EMRO, Eastern Mediterranean; EURO, Europe; PAHO, Americas; SEARO, South-East Asian; WPRO, Western Pacific.

Table 2 Risk acceptability and mitigation adequacy—selected worldwide survey responses

Risk acceptability	Risk is not acceptable at all (%)	Risk is acceptable for a short time (%)	Risk is negligible (%)	Don't know/unsure (%)
Infectious risk work environment				
Exposure to blood, body fluids, respiratory secretions and other potentially infectious materials	52	29	15	4
Inadequate sanitation facilities	52	21	23	4
Skin damage from personal protective equipment and/or frequent hand hygiene	33	46	16	5
Physical work environment				
Crowded workplace	42	36	18	4
Thermal discomfort (cold, heat, humidity)	25	46	24	5
Psychosocial work environment				
Bullying or psychological harassment	54	18	21	7
Sexual harassment	50	10	31	9
Work organisation				
Time pressure, high workload	38	49	10	3
Shift work with night shifts	23	48	21	8
Mitigation measure adequacy	Does not exist at all (%)	Exists and offers some protection (%)	Exists and offers full protection (%)	Don't know/unsure (%)
Infection prevention and control				
IPC policy in the health facility	8	60	28	4
Personal protective equipment (eg, masks, gloves, goggles, gowns) are readily available	8	55	34	3
Training and education of workers about infection prevention and control	11	54	32	3
Facilities for hand hygiene (hand washing and disinfection) are readily available	3	40	54	3
Occupational health and safety				
Occupational health and safety policy and management system in the facility	14	58	22	6
Regular assessment of workplace health and safety risks and controls	22	51	21	6
Engineering controls, such as ventilation, physical barriers, safer devices	19	54	19	8
Prevention of workplace violence and security measures	21	52	21	6
Workplace policies against bullying, psychological and sexual harassment	27	43	21	9

Most cited response highlighted in bold.
IPC, infection prevention and control.

of mitigation measures for other key areas such as IPC policy (28%), availability of PPE (34%), as well as training and education of workers about OHS (21%) and IPC (32%). Only in two mitigation measure areas—availability of facilities for hand hygiene, and policies for postexposure prophylaxis (such as HIV or hepatitis B)—did most participants indicate that measures existed and offered full protection (54% and 42%, respectively). These results show an overwhelming majority of participants indicating that the risks they faced were not acceptable at all and that very few of the corresponding mitigation measures offered adequate protection to HCWs.

Associations with risk exposure acceptability and mitigation measure adequacy

Unacceptable levels of *risk* (ie, factor scores below 5) were consistently reported for both work organisation and

work environment across geographic regions, economic income-level categories, equity classifications and COVID-19 incidence rates, with no statistically significant differences observed within these categories (table 3). However, we observed multiple significant differences in how the adequacy of OHS and especially IPC (overall mean of 4.67) *mitigation measures* was perceived. These apparent associations, observed to be present for all the explanatory factors we examined, drew attention to the need to consider the adjusted effect of each independent variable through the logistic regression analysis that we subsequently conducted.

To understand potential sources of difference that could be attributed to heterogeneous composition of country responses that is encountered in conducting a cross-country comparison such as the one we conducted,

**Table 3** Unadjusted risk acceptability and mitigation adequacy associations

Explanatory variable	Risk acceptability				Mitigation adequacy			
	Work environment		Work organisation		IPC		OHS	
	Mean	P value	Mean	P value	Mean	P value	Mean	P value
Total								
By country means	4.23		4.29		4.67		6.08	
By individuals	3.88		3.87		4.79		6.28	
Region								
AFRO	4.11	0.34	4.17	0.30	3.68	<0.01*	5.31	0.03*
EMRO	4.01		4.25		5.02		6.33	
EURO	4.47		4.24		5.28		6.54	
PAHO	4.03		3.99		4.24		5.92	
SEARO	3.44		4.76		5.11		6.30	
WPRO	4.83		5.03		5.24		6.35	
Economic classification								
High	4.51	0.24	4.62	0.15	5.61	<0.01*	6.99	<0.01*
Upper middle	4.05		4.05		4.85		6.17	
Lower middle	3.78		4.05		3.58		5.15	
Low	4.51		4.27		3.88		5.29	
Gini coefficient								
Q1 (lowest)	4.80	0.11	4.51	0.34	5.26	0.01*	6.64	0.04
Q2	4.10		4.29		4.31		5.81	
Q3	3.90		4.04		4.72		6.20	
Q4	3.98		3.80		3.89		5.55	
COVID-19 incidence rate								
Q1 (lowest)	3.95	0.50	4.09	0.84	4.17	<0.01*	5.64	0.07
Q2	4.18		4.25		4.39		5.95	
Q3	4.50		4.39		4.66		6.10	
Q4	4.19		4.16		5.44		6.62	

Full listing of WHO region countries at https://en.wikipedia.org/wiki/List_of_WHO_regions

*Indicates statistical significance ($p < 0.05$) of differences among the means of country mean values for category; significant values in bold.

AFRO, Africa; EMRO, Eastern Mediterranean; EURO, Europe; IPC, infection prevention and control; OHS, occupational health and safety; PAHO, Americas; Q, quartile; SEARO, South-East Asian; WPRO, Western Pacific.

table 4 presents a summary of the survey's individual-level data to indicate how gender and occupation were associated with respondent perceptions of acceptability and adequacy. Females were somewhat more likely than males to report workplace risks being unacceptable (3.76 vs 4.11; $p < 0.001$), but the strong presence of front-line patient care providers in the gendered health workforce was largely responsible for this, as no statistically significant differences were observed within other occupation groupings (see online supplemental table S3). In fact, patient care providers themselves stood out as being the occupational grouping most critical of workplace risk acceptability as well as OHS and IPC measure adequacy. In contrast, male administrators/managers stood out as the most likely to indicate that acceptable risk exposure and adequate risk mitigation measures were present. This discrepancy is understandable as front-line workers, and women in this occupation grouping, represent those most directly experiencing the impact of the COVID-19 pandemic. However, even in these more extreme

circumstances where differences were observed, the comparative differences in mean scores (that were then aggregated in calculating country mean values) were not large. Moreover, the fact that the African region, where strongest concerns about unacceptable risk and inadequate mitigation were expressed, actually had proportionately fewer female respondents, indicates that even these regional concerns that we observed may well have been under-represented in this unadjusted analysis.

Influence of between-country and within-country income disparities

Table 5 summarises the adjusted comparative effects of income level and income distribution disparity in each country setting while taking into consideration potential influences prompted by differing COVID-19 rates in the initial phase of the pandemic. While there was no difference between *higher and lower income countries* regarding the perception of unacceptable levels of risks in health-care workplaces in all settings, *within-country inequality* was

Table 4 Risk acceptability and mitigation adequacy associations* with gender and occupation

Explanatory variable	n†	Risk acceptability				Mitigation adequacy			
		Work environment		Work organisation		IPC		OHS	
		Mean	P value	Mean	P value	Mean	P value	Mean	P value
Gender‡									
Total	4863	3.88	<0.01§	3.87	0.40	4.79	0.09	6.28	0.07
Female	3220	3.76		3.85		4.74		6.33	
Male	1643	4.11		3.92		4.88		6.19	
Occupation¶									
Total	4916	3.88	0.04	3.87	0.10	4.79	<0.01§	6.28	0.19
Patient care	2792	3.91		3.88		4.63		6.27	
Specialist	1404	3.84		3.80		4.90		6.30	
Admin-manager	327	4.14		4.22		5.50		6.50	
Other	393	3.55		3.82		5.03		6.08	

*This table reports on total respondents in each category, without any consideration for different mixes of gender within different occupations, and different mixes of occupation within genders; online supplemental table S3 provides the results with full occupation and gender breakdowns.

†Total n varies by specific factor; this column refers to n for workplace environment, where response was greatest.

‡Only respondents indicating male or female were included in exploring differences.

§Indicates statistical significance ($p < 0.05$); significant values in bold.

¶Occupation was initially coded with finer detail but then consolidated in these composites for comparative analysis.

IPC, infection prevention and control; OHS, occupational health and safety.

associated with a mildly increased likelihood of *unacceptable levels of risk* with regard to both workplace environment (OR=0.92; $p=0.012$) and workplace organisational (OR=0.93; $p=0.017$) factors.

As was observed in unadjusted bivariate analyses, there was much stronger divergence in perceptions of acceptable *mitigation measures* by both *country income level* and *income inequality*, with an almost sevenfold greater likelihood of IPC measures (OR=6.61; $p=0.001$) being considered adequate in wealthier countries, and over a 10-fold difference in adequacy of OHS measures (OR=10.91; $p < 0.001$), despite the greater intensity of COVID-19 in wealthier countries at the time of the survey. In fact, the counterintuitive positive association that seemed to be present between intensity of COVID-19 and perceptions of adequacy disappeared in our adjusted multivariable analysis. And further to the observed unadjusted effect, higher inequality decreased the likelihood (OR=0.94; $p=0.025$) of deeming IPC measures to be adequate.

Analysis of the more homogeneously constituted population of female patient care provider respondents (online supplemental table S6) further revealed that this group's more critical assessment of risk that we had documented in table 4 especially influenced perceptions of risk acceptability in settings where COVID-19 exposure had intensified. In this regard, workplace organisational factors, which included consideration of the workload being encountered, were substantially more likely to be seen as unacceptable (OR=0.44; $p=0.034$) by female patient care providers in countries with higher COVID-19 presence; a perception reinforced by a further (although less pronounced) effect of in-country income inequality (OR=0.95; $p=0.093$).

As we had observed was the case for all respondents, female care providers in higher income countries were more likely to perceive mitigation measures to be adequate (OHS OR=3.94; $p=0.047$ and IPC OR=11.25; $p=0.004$) than those in more poorly resourced settings, and this was further accompanied by an effect of within-country inequality also contributing some explanatory power (OHS OR=0.92; $p=0.020$).

DISCUSSION

High levels of concern about emerging threats to HCWs were widely published in the first year of the COVID-19 pandemic, providing extensive evidence about morbidity and mortality associated with healthcare work^{33 34} as well as effects on job satisfaction.³⁵ Although meta-analyses have been conducted to synthesise such findings,³⁶ our article provides one of the first worldwide examinations of contextual factors affecting the well-being of HCWs during the COVID-19 pandemic, enabling a comparative cross-country analysis. In doing so, it notably complements studies calling attention to inadequate implementation of OHS and IPC measures, for example, in South Africa,³⁷ as well as a need to consider the influence of structural determinants that affect how risks are experienced in specific health worker exposure contexts.³⁸ The results presented here contribute a theoretical and empirically based understanding of the importance of inequality among and within countries in this regard. This has implications for preparedness for any future pandemic outbreaks.

Our findings clearly demonstrate that there is a strong need for improvements in OHS for HCWs to protect

Table 5 Factors associated with perceived risk acceptability and mitigation adequacy

Explanatory variable (organised by outcome area)	Unadjusted bivariate			Adjusted multivariable model*			
	OR*	95% CI	P value	B	OR*	95% CI	P value
Acceptable WP environmental risk							
Country income level†	1.56	1.08 to 2.27	0.231	0.215	1.24	0.70 to 2.20	0.708
Gini coefficient‡	0.91	0.88 to 0.94	0.005	-0.087	0.92	0.89 to 0.95	0.012
COVID log¶	1.01	0.83 to 1.23	0.965	0.083	1.09	0.78 to 1.51	0.801
Acceptable WP organisation risk							
Country income level†	0.83	0.59 to 1.17	0.587	-0.341	0.71	0.42 to 1.21	0.52
Gini coefficient‡	0.94	0.91 to 0.97	0.028	-0.076	0.93	0.90 to 0.96	0.017
COVID log¶	0.8	0.67 to 0.97	0.243	-0.113	0.89	0.66 to 1.21	0.710
Adequate IPC mitigation							
Country income level†	6.8	1.36 to 34.60	0.006	1.889	6.61	3.68 to 11.88	0.001
Gini coefficient‡	0.93	0.90 to 0.95	0.006	-0.036	0.94	0.91 to 0.96	0.025
COVID log¶	1.85	1.51 to 2.26	0.002	-0.064	0.76	0.55 to 1.03	0.373
Adequate OHS mitigation							
Country income level†	8.91	5.76 to 13.80	<0.001	2.389	10.91	5.63 to 21.12	<0.001
Gini coefficient‡	0.97	0.94 to 0.99	0.183	-0.009	0.99	0.96 to 1.02	0.779
COVID log¶	0.94	0.91 to 0.97	0.028	0.079	1.08	0.77 to 1.52	0.816

OR expressed as Exp(B) value in logistic regression analysis. * $P \leq 0.05$; ** $p < 0.001$. B is coefficient.

Variables where statistical significance is present are shown in bold.

*ORs were calculated by assessing the likelihood (OR) of the presence of a mean score ≥ 5 corresponding to assessments that mitigation 'exists and offers some protection' or better; or level of risk is assessed as 'risk is acceptable for a short time' or better.

†Country income was coded as comparing 'High and Upper-Middle Income' countries versus 'Low and Lower-Middle Income' countries.

‡Gini coefficient was considered in the logistic regression analysis as a continuous variable.

§Indicates statistical significance ($p < 0.05$); also bolded.

¶COVID-19 levels with the log value of the rate of cases per million at the beginning of the survey (taken 1 June 2020); log values to smooth very high levels while taking variation into account.

CI, Confidence Interval; IPC, infection prevention and control; OHS, occupational health and safety; WP, workplace.

against infectious disease transmission and to control the threat of psychosocial risks, a consideration that resonates with studies highlighting the effects on mental health of HCWs as already stressed workplaces with intensifying pressures when pandemics emerge.^{39–43} Widespread concerns about health risks identified in diverse locations such as Ethiopia, Turkey, Italy and Spain in many facets of health work^{44–47} signal a strong rationale for international collaboration in seeking effective technical and policy approaches to best protect HCWs.

Despite a common assessment of unacceptable levels of risk everywhere, our study revealed important differences in the perceived adequacy of protective measures to meet this challenge. Such results point to the need to add explicit attention to OHS measures in WHO's call for better planning healthcare human resources¹⁰ as well as the updating of WHO's *Global Plan of Action for Occupational Health*, considering what this means for HCWs in light of the COVID-19 experience.

While the case prevalence in any one single country clearly influences the intensity of possible healthcare workplace exposure as a global pandemic emerges, HCWs in all countries face the same need for proper PPE, appropriate testing and vaccines as they compete in the same markets and the same supply chains.^{48 49} While there is now appropriate attention focused on the need to address global inequities in vaccine accessibility,⁵⁰ our study highlights other inequities that also call for greater attention. Moreover, our analysis stands out by considering how variation in protecting HCWs may be associated with the presence of contextual social and economic inequities, itself an important social determinant of health that has been prominent in global health research literature. What is of particular relevance here is the vulnerability of HCWs as 'canaries' in a workplace made vulnerable by the emergence of a novel infectious disease,⁵¹ where preparedness to meet a new challenge is critical.

While the presence of unacceptable risk was clearly identified in all countries, it was striking that the strongest concern about inadequate protection of HCWs came not from the HICs hit most intensely by the initial wave of COVID-19 in early 2020, but rather less well-resourced settings that had yet to be as strongly affected. This vividly echoes pre-COVID findings that resource-poor countries have decreased capacities for protecting HCWs^{13 14} even beyond needs for testing and contact tracing, and consistent with studies noting needs for training and PPE for HCWs.⁵² This furthermore mirrors experience in previous pandemics such as Ebola in West Africa where meaningful investments in PPE were shown to be important elements in combating the spread of disease,⁵³ a matter that is now being observed with regard to COVID-19.⁵⁴ Our finding that country income level is strongly associated with greater capacity to provide prevention and mitigation within a health system is thus not surprising.

Previous literature on the effects of income inequality within a society has however been less conclusive, at times

contesting the implications of the Wilkinson's 'economic inequality hypothesis'. In this regard, Blázquez-Fernández and colleagues concluded that income inequality does not significantly reduce health in 'developed' societies⁵⁵ and Mellor and Milyo further argued that there is little support for relation between income inequality and individual or population health after fixed division effects were included.⁵⁶ However, when attention is paid to methodological concerns,¹⁷ strong evidence of the effect of economic inequality has been observed in sub-Saharan African countries.⁵⁷ Looking beyond levels of economic indicators alone, a systematic study of 'welfare regimes' (ie, characterisations of policy orientations dominant in a country at a particular time) has suggested that precarious workers fare better in the context of 'Scandinavian state' policies.⁵⁸ Indeed, countries that recognised COVID-19 as a work-related disease and supported workers with compensation and appropriate absence policies were reported to have reduced mental health stressors, pointing to opportunities for improving HCW well-being.⁵⁹ However, a systematic review of the impact of political economy on health observed substantial gaps in knowledge, calling for 'higher-quality reviews and empirical studies in this area'.⁶⁰

Our study suggests that societies with greater national income equality may well be characterised by policies that are more protective of vulnerable populations such as HCWs, a group whose comparatively high occupational health risk is aggravated by the onset of pandemics. To better understand the pathways and iterative relationships that can explain this, case study examinations would certainly be of value. Moreover, with health worker protection so strategically important to health system functioning during such crises that threaten global health equity, countries known to be highly unequal might accordingly be deemed to be in need of even further technical assistance and attention to ensure that adequate protection is provided to HCWs at risk.

Recognising that appreciation of the contribution of HCWs soared as the COVID-19 pandemic advanced, our observations that economic inequality among and within countries is associated with the degree to which HCWs face unacceptable risk and inadequate protection signal a vital need to promote social justice for those who play such an important role in the care of populations *before* a new pandemic emerges. In light of this, from an analytical perspective, we strongly endorse the call for a new paradigm⁶¹ to better understand how upstream and sociopolitical factors could be 'affecting the nature of work and employment and their impact on the health of workers, the public, and the planet'.⁶² This includes consideration of international cooperation with respect to vaccine supply, and to ensure that less wealthy countries receive technical assistance in establishing protection and mitigation programmes as well as attention to pathways sensitive to the offloading of risks to more marginalised worker populations.



Limitations and further research needs

Cross-country comparative studies such as ours rely on a convenience sample, leading to some countries being over-represented while others were under-represented or non-existent. To address possible concerns about the influence of countries with low respondent counts, we examined this concern by conducting sensitivity analyses, summarised in online supplemental table S7, to consider possible implications, but concluded that this did not warrant a questioning of our findings. Additionally, the classification of countries purely by national income levels leads to designating some countries as high income in settings where national institutions may be minimally developed despite high levels of income earned through high-value exports such as petroleum or in settings of small populations with externally controlled tourism sectors. As such, we developed grouping strategies to allow for a consideration of national contexts where resources could be considered comparatively more or less readily available to protect health workers. Stratification by WHO region was also important because these regions, while large and often heterogenous in nature, do constitute administrative units with an important governance role to play during the emergence of global outbreaks and pandemics.

It should also be acknowledged that differing perceptions of risks and mitigation measures around the world may be influenced by different HCW training and education standards, cultural nuances and institutional expectations. For example, Senthil and colleagues observed that workers in India found a high prevalence of workers unable to identify even immediate risks in an evidently hazardous environment.⁶³ Studies in the Middle East also reported gaps between actual hazards and HCW recognition.^{64 65} Ndejjo and colleagues report similar findings in Uganda and across sub-Saharan Africa.⁶⁶

CONCLUSION

This study adds to the literature on how risks become unevenly distributed, focusing here on country income level but also on within-country income inequality. As noted by Gostin *et al*,⁶⁷ WHO has an important role in supporting LMICs with technical guidance and operational assistance, while simultaneously meeting the needs of high-income countries for information sharing, research coordination and convening authorities, despite lacking both the authority and the resources to mount a more effective response to a global emergency such as this. Our study strongly suggests that international agencies with mandates related to fair trading practices and economic aid have to step up to address the disparities that threaten the healthcare workforce, and ensure that there is sufficient resilience to retain health workers needed for broader delivery of health services. It is also a matter of social justice that they do so.

Acknowledgements The authors would like to thank all the members of WHO occupational health ad hoc committee as well as all WHO occupational health collaborating centres around the world who contributed to the dissemination of the survey.

Contributors SH, AY and JMS conceptualised the research. AY and MZ participated in preparation of the original survey and its use. SH prepared the data and conducted the initial analysis. SH and JMS conducted subsequent data analyses and data interpretation. VWLT conducted literature reviews. SH wrote the initial manuscript draft. SH, AY, VWLT, MZ and JMS reviewed and edited the manuscript. AY and JMS accept full responsibility for the finished work and/or the conduct of the study as guarantors, had access to the data and controlled the decision to publish. All authors read and approved the final manuscript.

Funding This article has received funding from the International Development Research Centre (M20-00559) and the Canadian Institutes of Health Research (VS1-175519).

Competing interests None declared.

Patient and public involvement Patients and/or the public were involved in the design, or conduct, or reporting or dissemination plans of this research. Refer to the Methods section for further details.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants and was approved by Behavioural Research Ethics Board of the University of British Columbia (Ref: H20-01825). As the survey was designed to be completed and submitted anonymously, no formal request for signed consent was solicited, with participants' submission itself indicating consent to use the information provided as anonymised aggregated data.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. Anonymised data that underlie the results reported in this article are available on justified request to the corresponding author.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iD

Jerry M Spiegel <http://orcid.org/0000-0002-4699-3082>

REFERENCES

- 1 International Council of nurses. ICN calls for data on healthcare worker infection rates and deaths. *ICN - International Council of Nurses* <https://www.icn.ch/news/icn-calls-data-healthcare-worker-infection-rates-and-deaths>
- 2 World Health Organization. World Day for Safety and Health at Work: WHO key facts & key messages to support the day. Available: <https://www.who.int/news-room/detail/28-04-2020-who-calls-for-healthy-safe-and-decent-working-conditions-for-all-health-workers-amidst-covid-19-pandemic> [Accessed 12 Aug 2020].
- 3 Talisuna A, Yoti Z, Lee C, *et al*. The urgent need for a global commitment to protect healthcare workers. *BMJ Glob Health* 2020;5:e004077.
- 4 La T. COVID-19: protecting health-care workers. *Lancet* 2020;395:922. doi:10.1016/S0140-6736(20)30644-9
- 5 International A. Amnesty analysis reveals over 7,000 health workers have died from COVID-19. Available: <https://www.amnesty.org/en/latest/news/2020/09/amnesty-analysis-7000-health-workers-have-died-from-covid19/> [Accessed 24 Sep 2020].
- 6 The World Health Organization. World Health organization (who) on Twitter. Available: <https://twitter.com/WHO/status/1284148139797209093> [Accessed 13 Oct 2020].
- 7 Papoutsis E, Giannakoulis VG, Ntella V, *et al*. Global burden of COVID-19 pandemic on healthcare workers. *ERJ Open Res* 2020;6.

- doi:10.1136/23120541.00195-2020. [Epub ahead of print: 06 07 2020].
- 8 World Health Organization. Coronavirus disease 2019 (COVID-19) situation report; 2019.
 - 9 World Health Organization. Global spending on health: a world in transition. who.. Available: <https://www.who.int/publications/i/item/WHO-HIS-HGF-HFWorkingPaper-19.4> [Accessed 12 May 2022].
 - 10 World Health Organization. The World Health Report 2006 - working together for health. Available: <https://apps.who.int/iris/handle/10665/43432> [Accessed 12 May 2022].
 - 11 The world bank. physicians (per 1,000 people). Available: https://data.worldbank.org/indicator/SH.MED.PHYS.ZS?most_recent_value_desc=false [Accessed 24 Sep 2020].
 - 12 McDiarmid MA. Hazards of the health care sector: looking beyond infectious disease. *Ann Glob Health* 2014;80:315–9.
 - 13 Caeiro JP, Garzón MI. Controlling infectious disease outbreaks in low-income and middle-income countries. *Curr Treat Options Infect Dis* 2018;10:55–64.
 - 14 Mossburg S, Agore A, Nkimbeng M, et al. Occupational hazards among healthcare workers in Africa: a systematic review. *Ann Glob Health* 2019;85. doi:10.5334/aogh.2434. [Epub ahead of print: 06 06 2019].
 - 15 Gross JV, Mohren J, Erren TC. COVID-19 and healthcare workers: a rapid systematic review into risks and preventive measures. *BMJ Open* 2021;11:e042270.
 - 16 Harrigan SP, Tsang VWL, Spiegel JM. Health and safety of health workers in the context of COVID-19: a global survey. who collaborating centre, University of British Columbia. Available: http://med-fom-ghrp-spsh.sites.olt.ubc.ca/files/2020/09/WHO_HCW_Survey_COVID-19-report_MASTER.pdf
 - 17 Nowatzki NR. Wealth inequality and health: a political economy perspective. *Int J Health Serv* 2012;42:403–24.
 - 18 Subramanian SV, Kawachi I. Inequality I. Income inequality and health: what have we learned so far? *Epidemiol Rev* 2004;26:78–91.
 - 19 Pickett KE, Wilkinson RG. Income inequality and health: a causal review. *Soc Sci Med* 2015;128:316–26.
 - 20 Fujishiro K, Ahonen EQ, Gimeno Ruiz de Porras D, et al. Sociopolitical values and social institutions: studying work and health equity through the lens of political economy. *SSM Popul Health* 2021;14:100787.
 - 21 Adu PA, Spiegel JM, Yassi A. Towards TB elimination: how are macro-level factors perceived and addressed in policy initiatives in a high burden country? *Global Health* 2021;17:11.
 - 22 Gelfand MJ, Jackson JC, Pan X, et al. The relationship between cultural tightness-looseness and COVID-19 cases and deaths: a global analysis. *Lancet Planet Health* 2021;5:e135–44.
 - 23 Haug N, Geyrhofer L, Londei A, et al. Ranking the effectiveness of worldwide COVID-19 government interventions. *Nat Hum Behav* 2020;4:1303–12.
 - 24 Brauner JM, Mindermann S, Sharma M, et al. Inferring the effectiveness of government interventions against COVID-19. *Science* 2021;371. doi:10.1126/science.abd9338. [Epub ahead of print: 19 02 2021].
 - 25 Rinaldi C, Bekker MPM. A scoping review of populist radical right parties' influence on welfare policy and its implications for population health in Europe. *Int J Health Policy Manag* 2021;10:141–51.
 - 26 World Health Organization. Definition of regional groupings. who. Available: http://www.who.int/healthinfo/global_burden_disease/definition_regions/en/ [Accessed 12 Aug 2020].
 - 27 The world bank. world bank country and lending groups – world bank data help desk. Available: <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups> [Accessed 12 Aug 2020].
 - 28 The world bank. GINI index (world bank estimate). Available: <https://data.worldbank.org/indicator/SI.POV.GINI> [Accessed 12 Aug 2020].
 - 29 Ritchie H, Ortiz-Ospina E, Beltekian D. Coronavirus (COVID-19) Testing - Statistics and Research. Our World in Data. Available: <https://ourworldindata.org/coronavirus-testing> [Accessed 30 Nov 2020].
 - 30 Taber KS. The Use of Cronbach's Alpha When Developing and Reporting Research Instruments in Science Education. *Res Sci Educ* 2018;48:1273–96.
 - 31 R Core Team. R: a language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; 2013. <http://www.R-project.org/>
 - 32 IBM Corp. SPSS; 2018.
 - 33 Mhango M, Dzobo M, Chitungo I, et al. COVID-19 risk factors among health workers: a rapid review. *Saf Health Work* 2020;11:262–5.
 - 34 Bandyopadhyay S, Baticulon RE, Kadhum M, et al. Infection and mortality of healthcare workers worldwide from COVID-19: a systematic review. *BMJ Glob Health* 2020;5:e003097.
 - 35 Afulani PA, Nutor JJ, Agbadi P, et al. Job satisfaction among healthcare workers in Ghana and Kenya during the COVID-19 pandemic: role of perceived preparedness, stress, and burnout. *PLoS Glob Public Health* 2021;1:e0000022.
 - 36 Gómez-Ochoa SA, Franco OH, Rojas LZ, et al. COVID-19 in health-care workers: a living systematic review and meta-analysis of prevalence, risk factors, clinical characteristics, and outcomes. *Am J Epidemiol* 2021;190:161–175.
 - 37 Zungu M, Vuyi K, Mlangeni N, et al. Organizational factors associated with health worker protection during the COVID-19 pandemic in four provinces of South Africa. *BMC Health Serv Res* 2021;21:1080.
 - 38 Jeleff M, Traugott M, Jirovsky-Platter E, et al. Occupational challenges of healthcare workers during the COVID-19 pandemic: a qualitative study. *BMJ Open* 2022;12:e054516.
 - 39 Pappa S, Ntella V, Giannakas T, et al. Prevalence of depression, anxiety, and insomnia among healthcare workers during the COVID-19 pandemic: a systematic review and meta-analysis. *Brian Behav Immun* 2020;88:901–7.
 - 40 Krishnamoorthy Y, Nagarajan R, Saya GK, et al. Prevalence of psychological morbidities among general population, healthcare workers and COVID-19 patients amidst the COVID-19 pandemic: a systematic review and meta-analysis. *Psychiatry Res* 2020;293:113382.
 - 41 Salazar de Pablo G, Vaquerizo-Serrano J, Catalan A, et al. Impact of coronavirus syndromes on physical and mental health of health care workers: systematic review and meta-analysis. *J Affect Disord* 2020;275:48–57.
 - 42 De Brier N, Stroobants S, Vandekerckhove P, et al. Factors affecting mental health of health care workers during coronavirus disease outbreaks (SARS, MERS & COVID-19): A rapid systematic review. *PLoS One* 2020;15:e0244052.
 - 43 Dye TD, Alcantara L, Siddiqi S, et al. Risk of COVID-19-related bullying, harassment and stigma among healthcare workers: an analytical cross-sectional global study. *BMJ Open* 2020;10:e046620.
 - 44 Atnafie SA, Anteneh DA, Yimenu DK, et al. Assessment of exposure risks to COVID-19 among frontline health care workers in Amhara region, Ethiopia: a cross-sectional survey. *PLoS One* 2021;16:e0251000.
 - 45 Arslanca T, Fidan C, Daggez M, et al. Knowledge, preventive behaviors and risk perception of the COVID-19 pandemic: a cross-sectional study in Turkish health care workers. *PLoS One* 2021;16:e0250017.
 - 46 Conti C, Fontanesi L, Lanzara R, et al. Fragile heroes. The psychological impact of the COVID-19 pandemic on health-care workers in Italy. *PLoS One* 2020;15:e0242538.
 - 47 Garralda Fernandez J, Molero Vilches I, Bermejo Rodríguez A, et al. Impact of SARS-CoV-2 pandemic among health care workers in a secondary teaching hospital in Spain. *PLoS One* 2021;16:e0245001.
 - 48 Shone E. More than 850 health and social care workers have died of Covid in England and Wales since the pandemic began. The Scotsman. Available: <https://www.scotsman.com/health/coronavirus/more-than-850-health-and-social-care-workers-have-died-of-covid-in-england-and-wales-since-the-pandemic-began-3114202> [Accessed 10 Jun 2021].
 - 49 Our key findings about US healthcare worker deaths in the pandemic's first year. The Guardian. Available: <https://www.theguardian.com/us-news/ng-interactive/2020/dec/22/lost-on-the-frontline-our-findings-to-date> [Accessed 10 Jun 2021].
 - 50 Kavanagh MM, Gostin LO, Sunder M. Sharing technology and vaccine doses to address global vaccine inequity and end the COVID-19 pandemic. *JAMA* 2021;326:219–20.
 - 51 Lim DW, Htun HL, Wang Y, et al. Healthcare workers as 'canaries' for acute respiratory infections and pathogens during the COVID-19 pandemic. *J Hosp Infect* 2021;112:119–20.
 - 52 McMahon DE, Peters GA, Ivers LC, et al. Global resource shortages during COVID-19: bad news for low-income countries. *PLoS Negl Trop Dis* 2020;14:e0008412.
 - 53 Shears P, O'Dempsey TJD. Ebola virus disease in Africa: epidemiology and nosocomial transmission. *J Hosp Infect* 2015;90:1–9.
 - 54 Kim H, Hegde S, LaFiura C, et al. Access to personal protective equipment in exposed healthcare workers and COVID-19 illness, severity, symptoms and duration: a population-based case-control study in six countries. *BMJ Glob Health* 2021;6:e004611.
 - 55 Blázquez-Fernández C, Cantarero-Prieto D, Pascual-Saez M. Does rising income inequality reduce life expectancy? new evidence for 26 European countries (1995–2014). *Global Economic Review* 2018;47:464–79.
 - 56 Mellor JM, Milyo J. Is exposure to income inequality a public health concern? Lagged effects of income inequality



- on individual and population health. *Health Serv Res* 2003;38:137–51.
- 57 Odusanya IA, Akinlo AE, Obafemi Awolowo University, Nigeria. Income inequality and population health in sub-Saharan Africa: a test of income Inequality-Health hypothesis. *Journal of Population and Social Studies* 2021;29:235–54.
- 58 Kim I-H, Muntaner C, Vahid Shahidi F, *et al.* Welfare states, flexible employment, and health: a critical review. *Health Policy* 2012;104:99–127.
- 59 Sandal A, Yildiz AN. COVID-19 as a recognized work-related disease: the current situation worldwide. *Saf Health Work* 2021;12:136–8.
- 60 McCartney G, Hearty W, Arnot J, *et al.* Impact of political economy on population health: a systematic review of reviews. *Am J Public Health* 2019;109:e1–12.
- 61 Flynn MA, Check P, Steege AL, *et al.* Health equity and a paradigm shift in occupational safety and health. *Int J Environ Res Public Health* 2021;19:349.
- 62 Loewenson R. Rethinking the paradigm and practice of occupational health in a world without Decent work: a perspective from East and southern Africa. *New Solut* 2021;31:107–12.
- 63 Senthil A, Anandh B, Jayachandran P, *et al.* Perception and prevalence of work-related health hazards among health care workers in public health facilities in southern India. *Int J Occup Environ Health* 2015;21:74–81.
- 64 Muralidhar S, Singh PK, Jain RK, *et al.* Needle stick injuries among health care workers in a tertiary care hospital of India. *Indian J Med Res* 2010;131:405–10.
- 65 Kotwal A, Taneja D, Workers HC. Health care workers and universal precautions: perceptions and determinants of non-compliance. *Indian J Community Med* 2010;35:526–8.
- 66 Ndejjo R, Musinguzi G, Yu X, *et al.* Occupational health hazards among healthcare workers in Kampala, Uganda. *J Environ Public Health* 2015;2015:e913741.
- 67 Gostin LO, Moon S, Meier BM. Reimagining global health governance in the age of COVID-19. *Am J Public Health* 2020;110:1615–9.

Appendices

Survey instrument

Health and safety of health workers in COVID-19

Welcome to the survey on health and safety of health workers in COVID-19

Dear colleague,

This survey aims to identify the most common occupational risks for the health and safety of health workers and the measures for their prevention in the context of the ongoing pandemic of Corona Virus Infectious Disease (COVID-19).

In this survey we are interested in the health and safety of all health workers - all people engaged in the promotion, protection or improvement of the health of the population. This includes health workers involved in direct patient care, both formal and informal, in public and private facilities, including traditional medicine, as well as other assisting and supporting staff, including administration, management, ambulance drivers, public health workers, community health workers, and others.

The survey is intended for health workers, managers, and practitioners providing services for protecting the health and safety of workers in health facilities. The results will be used to inform action at all levels for improving the protection of health and safety of health workers in the ongoing response to COVID-19.

The survey has been developed by an international group of experts convened by the World Health Organization and the International Labour Organization and should take approximately 7 minutes to complete.

Your answers are completely confidential, and the data will be processed and analyzed in a way that will not link your answers to your identity.

B. About yourself and your area of work

1. In what country do you usually work?

Standard list of all countries in the world

Other (please specify)

* 2. Your gender?

Male

Female

Other

Prefer not to answer

3. What is your primary area of work? (responses below were randomized)

Administration and clerical support
Allied health professional
Community health worker
Infection prevention and control
Management and human resources
Mental health and psychosocial support
Occupational and environmental health
Patient care (medicine, nursing, midwifery, dentistry)
Pharmacy
Public health
Support staff – cleaner, driver, food worker
Other

4. You work most of the time for: (responses below were randomized)

Academia, research
Business enterprise or farm
Employers' association /hospital federation
Healthcare facility - hospital, primary health-care centre, isolation camp
Local community
National government agency
Other
Professional association
Social care facility (e.g. nursing home, home care)
Sub-national (provincial, district) authority
Trade union

C. Risks for health and safety of health workers

Think about the working conditions of health workers in your country, jurisdiction or health facility - those that you are most familiar with. No workplace is without risk, but some risks are negligible, or acceptable for a short time, and some are not acceptable at all. Below are some common risks for the health and safety of health workers; we are asking you to rate the current level of these risks, now during the COVID-19 pandemic.

5. How would you rate the level of these risks for health workers, now? (randomized)

Questions	Risk is negligible	Risk is acceptable for a short time	Risk is not acceptable at all	Don't know/Unsure
Skin damage from personal protective equipment and/or frequent hand hygiene				
Needle-sticks and sharps injuries				
Inadequate sanitation facilities				
Insufficient access to facilities for personal hygiene, such as, shower and menstrual hygiene				
Exposure to blood, body fluids, respiratory secretions, and other potentially infectious materials				

6. How would you rate the level of these risks for health workers, now? (randomized)

Questions	Risk is negligible	Risk is acceptable for a short time	Risk is not acceptable at all	Don't know/Unsure
Back injury from manual handling of patients and heavy objects				
Hazardous chemicals, drugs, cleaning and disinfection agents				
Slips, trips, and falls				
Crowded workplace				
Thermal discomfort (cold, heat, humidity)				

7. How would you rate the level of these risks for health workers, now? (randomized)

Questions	Risk is negligible	Risk is acceptable for a short time	Risk is not acceptable at all	Don't know/Unsure
Physical violence and assaults				
Bullying or psychological harassment at the workplace				
Sexual harassment				

8. How would you rate the level of these risks for health workers, now? (randomized)

Questions	Risk is negligible	Risk is acceptable for a short time	Risk is not acceptable at all	Don't know/Unsure
Regular long working hours (more than 48 hours a week)				
Time pressure, high workload				
Shift work with night shifts				
Insufficient time-off duty to rest (less than 11 hours between shifts)				

9. How would you rate the level of these risks for health workers, now? (randomized)

Questions	Risk is negligible	Risk is acceptable for a short time	Risk is not acceptable at all	Don't know/Unsure
Skin damage from personal protective equipment and/or frequent hand hygiene				
Needle-sticks and sharps injuries				
Inadequate sanitation facilities				
Insufficient access to facilities for personal hygiene, such as, shower and menstrual hygiene				
Exposure to blood, body fluids, respiratory secretions, and other potentially infectious materials				

D. Preventive measures

There are measures for the prevention of most risks for health and safety at work, but these measures may not be fully implemented and not all workers may benefit from these measures. Think again about the working conditions of health workers in your country, jurisdiction or health facility - those that you are most familiar with. The following questions are about the preventive measures for their health and safety in the real situation, now, during the COVID-19 pandemic.

10. How would you rate the level of application of these measures in the health services according to your knowledge? (randomized)

Questions	Does not exist at all	Exists and offers <u>some</u> protection	Exists and offers <u>full</u> protection	Don't know/Unsure
Policy for infection prevention and control in the health facility				
Processes for triage of patient in place at the emergency room, including early detection and isolation of infectious patients				
Routine assessment of the risk of exposure to body substances or contaminated surfaces before any health care activity and use of appropriate measures for personal protection				
Regular environmental clean-up and disinfection				
Immunization of health workers				

11. How would you rate the level of application of these measures in the health services according to your knowledge? (randomized)

Questions	Does not exist at all	Exists and offers <u>some</u> protection	Exists and offers <u>full</u> protection	Don't know/Unsure
Reporting of incidental exposures to blood, body fluids, or respiratory secretions				
Policies in place for post-exposure prophylaxis, such as, for HIV, Hepatitis B				
Facilities for hand hygiene (hand washing and disinfection) are readily available				
Personal protective equipment, such as masks, gloves, goggles, gowns are readily available				
Training and education of workers about infection prevention and control				

12. How would you rate the level of application of these measures in health services according to your knowledge? (randomized)

Questions	Does not exist at all	Exists and offers <u>some</u> protection	Exists and offers <u>full</u> protection	Don't know/Unsure
Prevention of workplace violence and security measures				
Management of working time, rest and recuperation				
Workplace policies against bullying, psychological and sexual harassment				
Human resource management of safe staffing and workload				
Psycho-social support and counselling				

13. How would you rate the level of application of these measures in the health services according to your knowledge? (randomized)

Questions	Does not exist at all	Exists and offers <u>some</u> protection	Exists and offers <u>full</u> protection	Don't know/Unsure
Occupational safety and health policy and management system in the facility				
Regular assessment of workplace health and safety risks and controls				
Engineering controls, such as ventilation, physical barriers, safer devices				
Ergonomic workplace design and furniture				
Devices for patient handling and lifting of loads				

14. How would you rate the level of application of these measures in the health services according to your knowledge? (randomized)

Questions	Does not exist at all	Exists and offers some protection	Exists and offers full protection	Don't know/Unsure
Regular medical check-ups of health workers				
Medical first aid kits				
Consultations between management and workers regarding health and safety at work				
Training and education of workers about occupational safety and health				

Other (please specify)

Table A 1: Survey options for occupation

Group	Survey responses
Patient care/health services	Patient care (medicine, nursing, midwifery, dentistry) Allied health professional Mental health and psychosocial support Pharmacy Community health worker
Specialized support	Occupational and environmental health Public health Infection prevention and control Support staff – cleaner, driver, food worker
Clerical support/administration and management	Administration and clerical support Management and human resources

Table A 2: Comparison between participants who answered risk and mitigation questions

Variables		Those who responded to risk questions (n=4977)	Those who responded to mitigation measure questions (n=4076)
Countries by region	AFRO	10%	11%
	EMRO	4%	4%
	EURO	35%	36%
	PAHO	31%	31%
	SEARO	3%	3%
	WPRO	15%	16%
Economic Class	High	59%	60%
	Lower-middle	10%	10%
	Upper-middle	27%	26%
	Low	4%	4%
Sex	Male	33%	34%
	Female	65%	65%
	Other / prefer not to answer	1%	1%
Occupation	Patient care/health services	56%	58%
	Specialized support	29%	29%
	Clerical support/administration and management	7%	7%
	Other	8%	7%



Speak up
for health worker safety!



Health and safety of health workers in the context of COVID-19: A global survey

Sean P. Harrigan, Vivian W. L. Tsang, Jerry M. Spiegel, Annalee Yassi

September 15, 2020




a place of mind
THE UNIVERSITY OF BRITISH COLUMBIA

Faculty of Medicine
School of Population and Public Health



WHO Collaborating Centre
for Occupational Health

© Global Health Research Program, School of Population and Public Health,



ONLINE SUPPLEMENTARY INFORMATION

Table S1. Study population demographics

Variables	n	%
Total number of participants	4977	100%
Total who replied to the health and safety risks for health workers questions	4977	100%
Total who replied to the mitigation measures questions	4076	82%
Total number of countries	161	100%
Countries by region		
African region	516	10%
Americas region	1565	31%
Eastern Mediterranean region	221	4%
European region	1757	35%
South East Asian region	158	3%
Western Pacific region	760	15%
Economic Class		
High	2960	59%
Lower-middle	480	10%
Upper-middle	1324	27%
Low	213	4%
Sex		
Male	1654	33%
Female	3259	65%
Other / prefer not to answer	64	1%
Occupation		
Patient care/health services	2805	56%
Specialized support	1426	29%
Clerical support/administration and management	341	7%
Other	405	10%
Type of employer		
Health services	3038	61%
Government services	758	15%
Business and farms	473	10%
Academia	367	7%
Professional associations	177	4%
International organization/NGO and non-for-profit	51	1%
Other	113	2%

Table S2. Summary of Factors and their constituent question areas

Survey area of focus	Factor	Description of question
Health and Safety Risks	Work Environment (Factor 1)	Blood and bodily fluids exposure Skin damage from PPE and hand hygiene Needlesticks and sharps injuries Sanitation facilities Personal hygiene Thermal discomfort Crowded workplace Slips trips & falls Back injury form heavy lifting Chemicals Bullying & harassment Sexual harassment
	Work Organization (Factor 2)	Physical violence and assaults Time pressure Shift work Long working hours Insufficient rest
Mitigation measures	Infection protection and control (Factor 3)	IPC policy Patient triage Standard precautions Cleanup & disinfection Immunization Reporting of blood exposure Policies for post-exposure prophylaxis Hand hygiene PPE IPC training
	Occupational health and safety (Factor 4)	OSH policy Occupational health and safety risk assessment Occupational health and safety engineering controls Ergonomic workplace design and furniture Safe patient handling Violence prevention Management of working time & rest Policy harassment Safe staffing & workload Psychosocial support Medical checkups Medical first aid kits Labour management consultation on OSH Occupational health and safety training

Table S3: Detailed breakdowns in individual level responses by occupation and gender**a. Detailed breakdowns in responses by occupation showing gender differences**

Population characteristic	n ^c	Risk acceptability				Mitigation adequacy			
		Work environment		Work organization		IPC		OHS [f3]	
		mean	p	mean	p	mean	p	mean	P
All Occupations	4916	3.88	0.04	3.87	0.101	4.79	<0.01	6.28	0.19
<i>m</i>		4.11	<0.01	3.92	0.397	4.88	<u>0.089</u>	6.19	<u>0.07</u>
<i>f</i>		3.76		3.85		4.74		6.33	
- Patient Care	2792	3.92	0.001	3.88	0.296	4.63	0.76	6.27	<u>0.092</u>
<i>m</i>		4.19		3.96		4.65		6.15	
<i>f</i>		3.80		3.84		4.62		6.32	
- Specialist	1404	3.84	0.129	3.80	0.669	4.90	0.218	6.30	0.193
<i>m</i>		3.97		3.77		5.00		6.20	
<i>f</i>		3.73		3.83		4.81		6.39	
- Admin-Mgr	327	4.14	<u>0.06</u>	4.22	0.632	5.50	0.512	6.50	0.872
<i>m</i>		4.65		4.33		5.65		6.54	
<i>f</i>		3.92		4.17		5.43		6.48	
- Other	393	3.55	0.205	3.82	0.189	5.03	0.20	6.08	0.89
<i>m</i>		3.82		4.10		5.30		6.11	
<i>f</i>		3.43		3.69		4.90		6.08	

Note: Statistical significance at p<.05 level indicated by bold; statistical significance at p<.10 level indicated by underlining.

Italic font is used for breakdown analysis (i.e. by gender) of population attribute (regular font) being analyzed (i.e. by occupation) and values are right-justified.

Regular font is used for analysis of the population attribute (regular font) being analyzed (i.e. by occupation).

Green colour indicates significant value deemed as associated with less risk (i.e. less unacceptable exposure; more adequate mitigation).

Red colour indicates statistically significant value associated with more risk (i.e. more unacceptable exposure; less adequate mitigation).

b. Detailed breakdowns in responses by occupation showing gender differences

Population characteristic	n ^c	Risk acceptability				Mitigation adequacy			
		Work environment		Work organization		IPC		OHS [f3]	
		mean	p	mean	p	mean	p	mean	P
Total	4863	3.88	<0.01*	3.87	0.4	4.79	<u>0.09</u>	6.28	<u>0.07</u>
<i>Patient Care</i>		3.91		3.88		4.63		6.27	
<i>Specialist</i>		3.84		3.80		4.90		6.30	
<i>Admin-Mgr</i>		4.14		4.22		5.50		6.50	
<i>Other</i>		3.55		3.82		5.03		6.08	
Female	3220	3.76	0.189	3.85	0.273	4.74	<0.01*	6.33	0.288
<i>Patient Care</i>		3.80		3.84		4.62		6.32	
<i>Specialist</i>		3.73		3.83		4.82		6.39	
<i>Admin-Mgr</i>		3.92		4.17		5.43		6.48	
<i>Other</i>		3.43		3.69		4.90		6.07	
Male	1643	4.11	<u>0.089</u>	3.92	0.166	4.88	0.001	6.19	0.527
<i>Patient Care</i>		4.19		3.96		4.65		6.15	
<i>Specialist</i>		3.97		3.77		5.00		6.20	
<i>Admin-Mgr</i>		4.65		4.33		5.65		6.54	
<i>Other</i>		3.82		4.10		5.30		6.11	

Note: Statistical significance at p<.05 level indicated by bold; statistical significance at p<.10 level indicated by underlining.

Italic font is used for breakdown analysis (i.e. by occupation) of population attribute (regular font) being analyzed (i.e. by gender) and values are right-justified.

Regular font is used for analysis of the population attribute (regular font) being analyzed (i.e. by gender).

Green colour indicates significant value deemed as associated with less risk (i.e. less unacceptable exposure; more adequate mitigation).

Red colour indicates statistically significant value associated with more risk (i.e. more unacceptable exposure; less adequate mitigation).

Table S4. Responses to health and safety risk questions

Question	Risk is not acceptable at all	Risk is acceptable for a short time	Risk is negligible	Don't know/unsure
Infectious risks				
Exposure to blood, body fluids, respiratory secretions, and other potentially infectious materials	52%	29%	15%	4%
Skin damage from personal protective equipment and/or frequent hand hygiene	33%	46%	16%	5%
Needle-sticks and sharps injuries	46%	21%	26%	7%
Inadequate sanitation facilities	52%	21%	23%	4%
Insufficient access to facilities for personal hygiene, such as, shower and menstrual hygiene	49%	22%	23%	6%
Physical work environment				
Thermal discomfort (cold, heat, humidity)	25%	46%	24%	5%
Crowded workplace	42%	36%	18%	4%
Slips, trips, and falls	34%	26%	33%	7%
Back injury from manual handling of patients and heavy objects	41%	34%	19%	6%
Hazardous chemicals, drugs, cleaning and disinfection agents	36%	36%	22%	6%
Psychosocial work environment				
Bullying or psychological harassment at the workplace	54%	18%	21%	7%
Sexual harassment	50%	10%	31%	9%
Physical violence and assaults	54%	16%	24%	6%
Work organization				
Time pressure, high workload	38%	49%	10%	3%
Shift work with night shifts	23%	48%	21%	8%
Regular long working hours (more than 48 hours a week)	38%	42%	15%	5%
Insufficient time-off duty to rest (less than 11 hours between shifts)	40%	36%	18%	6%

Note: Most cited response highlighted in bold

Table S5. Responses to mitigation measure questions

Question	Does not exist at all	Exists and offers some protection	Exists and offers full protection	Don't know/ unsure
Infection prevention and control				
IPC policy in the health facility	8%	60%	28%	4%
Patient triage	9%	54%	28%	9%
Standard precautions	15%	53%	25%	7%
Regular environmental clean-up and disinfection	6%	56%	34%	4%
Immunization of health workers	19%	46%	26%	9%
Reporting of incidental exposures to blood, body fluids, or respiratory secretions	10%	50%	32%	8%
Policies in place for post-exposure prophylaxis, such as, for HIV, Hepatitis B	8%	41%	42%	9%
Facilities for hand hygiene (hand washing and disinfection) are readily available	3%	40%	54%	3%
Personal protective equipment, such as masks, gloves, goggles, gowns are readily available	8%	55%	34%	3%
Training and education of workers about infection prevention and control	11%	54%	32%	3%
Occupational safety and health				
Occupational safety and health policy and management system in the facility	14%	58%	22%	6%
Regular assessment of workplace health and safety risks and controls	22%	51%	21%	6%
Engineering controls, such as ventilation, physical barriers, safer devices	19%	54%	19%	8%
Ergonomic workplace design and furniture	33%	48%	12%	7%
Devices for patient handling and lifting of loads	27%	48%	15%	10%
Prevention of workplace violence and security measures	21%	52%	21%	6%
Management of working time, rest and recuperation	20%	55%	19%	6%
Workplace policies against bullying, psychological and sexual harassment	27%	43%	21%	9%
Human resource management of safe staffing and workload	24%	52%	16%	8%
Psycho-social support and counselling	32%	46%	15%	7%
Regular medical check-ups of health workers	33%	43%	18%	6%
Medical first aid kits	15%	48%	30%	7%
Consultations between management and workers regarding health and safety at work	25%	50%	19%	6%
Training and education of workers about occupational safety and health	20%	54%	21%	5%

Note: Most cited response highlighted in bold

Table S6: Comparing analyses of acceptability and adequacy in different stratified populations

	Patient Care Female		Patient Care All		Frontline ^e all		All occupations	
Countries included	112		133		156		161	
Explanatory Variable (organized by outcome area)	OR ^d	p	OR ^d	p	OR ^d	p	OR ^d	p
Acceptable WP Enviro. Risk								
Country Income Level ^a	0.81	0.736	0.63	0.415	1.01	0.981	1.24	0.708
Gini Coefficient ^b	0.98	0.609	0.96	0.1991	0.94	0.068**	0.92	0.012*
COVID-Log ^c	1.12	0.777	1.33	0.398	1.22	0.549	1.09	0.801
Acceptable WP Org. Risk								
Country Income Level ^a	1.30	0.657	1.57	0.446	1.06	0.926	0.71	0.52
Gini Coefficient ^b	0.95	0.093**	0.93	0.024*	0.93	0.056**	0.95	0.017*
COVID-Log ^c	0.44	0.034*	0.66	0.248	0.98	0.951	0.89	0.710
Adequate IPC mitigation								
Country Income Level ^a	11.25	0.004*	7.48	0.006*	6.92	0.001*	6.61	0.001*
Gini Coefficient ^b	0.95	0.220	0.95	0.125	0.95	0.047*	0.94	0.025*
COVID-Log ^c	0.69	0.479	0.7	0.181	0.70	0.264	0.76	0.373
Adequate OHS mitigation								
Country Income Level ^a	3.94	0.047*	5.86	0.004*	18.39	<0.001*	10.91	<0.001*
Gini Coefficient ^b	0.92	0.020*	0.95	0.083**	0.99	0.755	0.99	0.779
COVID-Log ^c	0.79	0.556	0.69	0.281	0.99	0.988	1.08	0.816

^a Country Income was coded as comparing “High and Upper-Middle Income” countries versus “Low and Lower-Middle Income” countries

^b Gini coefficient was considered in the logistic regression analysis as a continuous variable;

^c COVID levels where the log value of the rate of cases per million at the beginning of the survey (taken June 1, 2020); log values to smooth very high levels while taking variation into account

^d Odds Ratios were calculated by assessing the likelihood (OR) of the presence of a mean scores ≥ 5 corresponding to assessments that mitigation “exists and offers some protection” or better; or level of risk is assessed as “risk is acceptable for a short time” or better.

^e includes all patient care and workplace specialist support personnel such as OHS and IPC professionals

* Indicates statistical significance ($p < 0.05$) and also bolded; ** $p < .10$ but $> .05$; indicated in italics and bold

Table S7: Comparing analyses when minimum country sample size provisions are applied

	Patient Care Female		Patient Care Female		Frontline ^e		Frontline ^e	
Minimum country sample	2		all		4		all	
Countries included	80		112		101		161	
Explanatory Variable (organized by outcome area)	OR ^d	p	OR ^d	p	OR ^d	p	OR ^d	p
Acceptable WP Enviro. Risk								
Country Income Level ^a	0.62	0.556	0.81	0.736	1.03	0.967	1.01	0.981
Gini Coefficient ^b	0.97	0.404	0.98	0.609	0.95	0.228	0.94	0.068**
COVID-Log	1.11	0.852	1.12	0.777	0.89	0.797	1.22	0.549
Acceptable WP Org. Risk								
Country Income Level ^a	1.2	0.830	1.30	0.657	1.33	0.736	1.06	0.926
Gini Coefficient ^b	0.97	0.487	0.95	0.093**	0.94	0.230	0.93	0.056**
COVID-Log	0.77	0.617	0.44	0.034*	0.42	0.457	0.98	0.951
Adequate IPC mitigation								
Country Income Level ^a	9.62	0.041*	11.25	0.004*	36.89	0.001*	6.92	0.001*
Gini Coefficient ^b	0.94	0.223	0.95	0.220	0.98	0.647	0.95	0.047*
COVID-Log	0.54	0.433	0.69	0.479	0.48	0.190	0.70	0.264
Adequate OHS mitigation								
Country Income Level ^a	3.00	0.153	3.94	0.047*	5.46	0.020*	18.39	<0.001*
Gini Coefficient ^b	0.94	0.077**	0.92	0.020*	0.94	0.061**	0.99	0.755
COVID-Log	0.67	0.392	0.79	0.556	0.56	0.152	0.99	0.988

^a Country Income was coded as comparing “High and Upper-Middle Income” countries versus “Low and Lower-Middle Income” countries

^b Gini coefficient was considered in the logistic regression analysis as a continuous variable;

^c COVID levels where the log value of the rate of cases per million at the beginning of the survey (taken June 1, 2020); log values to smooth very high levels while taking variation into account

^d Odds Ratios were calculated by assessing the likelihood (OR) of the presence of a mean scores ≥ 5 corresponding to assessments that mitigation “exists and offers some protection” or better; or level of risk is assessed as “risk is acceptable for a short time” or better.

^e includes all patient care and workplace specialist support personnel such as OHS and IPC professionals

* Indicates statistical significance ($p < 0.05$) and also bolded; ** $p < .10$ but $> .05$; indicated in italics and bold