



Author index

AGNAGNA, M.

- Aerobic intestinal flora of wild-caught African dwarf crocodiles *Osteolaemus tetraspis*—F.W. HUCHZERMEYER, M.M. HENTON, J. RILEY and M. AGNAGNA 201–204

AIRE, T.A.

- Ultrastructural study of the luminal surface of the ducts of the epididymis of gallinaceous birds—T.A. AIRE and D. JOSLING 191–199

AITCHISON, H.

- The prevalence of different African horsesickness virus serotypes in the Onderstepoort area near Pretoria, during an outbreak of African horsesickness in South Africa in 1995/1996—C.W. BREMER, G.H. GERDES, H. AITCHISON, I. LOUW, R.R. GREYLING and J. WELGEMOED 65–70

AMANFU, W.

- Bovine borreliosis in Botswana—S.P. SHARMA, W. AMANFU and T.C. LOSHO 221–223

ANJILI, C.O.

- Leishmania major* infections in *Phlebotomus duboscqi* fed on murine models immunized with *L. major* sub-cellular antigens and sandfly gut antigens—P.A. MBATI, C.O. ANJILI, S. ODONGO, P. OGAJA and W. TONU 57–63

ANOSA, V.O.

- In vitro* erythrophagocytosis by cultured macrophages stimulated with extraneous substances and those isolated from the blood, spleen and bone marrow of Boran and N'Dama cattle infected with *Trypanosoma congolense* and *T. vivax*—V.O. TAIWO and V.O. ANOSA 273–287

- Differential expression of surface membrane antigens on bovine monocytes activated with recombinant cytokines and during *Trypanosoma congolense* infection—V.O. TAIWO, V.O. ANOSA and J.O. OLUWANIYI 289–296

BATH, G.F.

- The effect of a natural maedi-visna virus infection on the productivity of South African sheep—B. DUNGU, J. VORSTER, G.F. BATH and D.W. VERWOERD 87–96

BENGIS, R.G.

- The rise and fall of tuberculosis in a free-ranging chacma baboon troop in the Kruger National Park—D.F. KEET, N.P.J. KRIEK, R.G. BENGIS, D.G. GROBLER and A. MICHEL 115–122

BINTA, M.G.

- Helminth parasites of domestic pigeons (*Columba livia domestica*) in Sebele, Gaborone, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, R. NDEBELE and R. PANZIRAH 75–76

BLIGHT, G.W.

- Evaluation of a commercially available ELISA kit for detection of antibodies to *Anaplasma marginale* and *Anaplasma centrale* in cattle in Australia and Zimbabwe—P.M. BOWLES, J.B. MOLLOY, G.W. BLIGHT, S. SINGH and L.G. MABIKACHECHE 83–86

BOOMKER, J.

- A check-list of the nematode parasites of South African serpentes (snakes) and sauria (lizards)—S.F.B.N. HERING-HAGENBECK and J. BOOMKER 1–13
- Parasites of South African wildlife. XVI. Helminths of some antelope species from the Eastern and Western Cape Provinces—J. BOOMKER, I.G. HORAK, R. WATERMEYER and D.G. BOOYSE 31–41
- Selected chemical parameters in the blood and metals in the organs of the Nile crocodile, *Crocodylus niloticus*, in the Kruger National Park—D. SWANEPOEL, J. BOOMKER and N.P.J. KRIEK 141–148
- Molineus cati* n. sp. (Nematoda, Trichostrongylina, Molineoidea), a parasite of feral cats, *Felis catus* Linnaeus, 1758 in South Africa—MARIE-CLAUDE DURETTE-DESSET, J. BOOMKER and F.S. MALAN 173–177
- Studies on the genus *Setaria* Viborg, 1795 in South Africa. I. *Setaria africana* (Yeh, 1959)—R. WATERMEYER, J. BOOMKER and J.F. PUTTERILL 229–234

BOOYSE, D.G.

- Parasites of South African wildlife. XVI. Helminths of some antelope species from the Eastern and Western Cape Provinces—J. BOOMKER, I.G. HORAK, R. WATERMEYER and D.G. BOOYSE 31–41

BOTHA, C.J.

- Investigation of biological samples for monofluoroacetate and *Dichapetalum cymosum* poisoning in southern Africa—P.P. MINNAAR, R.I. McCRINDLE, T.W. NAUDÉ and C.J. BOTHA 27–30

BOWLES, J.

- Evaluation of a commercially available ELISA kit for detection of antibodies to *Anaplasma marginale* and *Anaplasma centrale* in cattle in Australia and Zimbabwe—P.M. BOWLES, J.B. MOLLOY, G.W. BLIGHT, S. SINGH and L.G. MABIKACHECHE 83–86

BRAACK, L.E.O.

- Parasites of domestic and wild animals in South Africa. XXXVIII. Ixodid ticks collected from 23 wild carnivore species—I.G. HORAK, L.E.O. BRAACK, L.J. FOURIE and JANE B. WALKER 239–250

BREMER, C.W.

- The prevalence of different African horsesickness virus serotypes in the Onderstepoort area near Pretoria, during an outbreak of African horsesickness in South Africa in 1995/1996—C.W. BREMER, G.H. GERDES, H. AITCHISON, I. LOUW, R.R. GREYLING and J. WELGEMOED 65–70

CHABO, R.G.

- Helminth parasites of domestic pigeons (*Columba livia domestica*) in Sebele, Gaborone, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, R. NDEBELE and R. PANZIRAH 75–76

CHITAURO, D.

- Pasteurella gallinarum*: Zimbabwean experience of a versatile pathogen—K. MOHAN, F. DZIVA and D. CHITAURO 301–305

COETZEE, L.

- Oribatid mites (Acari, Oribatida) as intermediate hosts of tapeworms of the Family Anoplocephalidae (Cestoda) and the transmission of *Moniezia expansa* cysticercooids in South Africa—R. SCHUSTER, L. COETZEE and J.F. PUTTERILL 49–55

DAKWA, C.

- Fertility in rats immunized with steroid-free bovine follicular fluid—C.R. SEWANI-RUSIKE and C. DAKWA 257–262

DICKS, L.M.T.

- Phage types of *Salmonella enterica* serovar Enteritidis isolated in South Africa from 1991–1995—L. MARÉ, M.L. VANDER WALT and L.M.T. DICKS 129–133

DIPPENAAR, S.M.

- Observations on a lesser-known monogenean, *Udonella myliobati*, from a copepod parasite, *Lepeophtheirus natalensis*, parasitizing the spotted ragged-tooth shark, *Carcharias taurus*, from South African waters—P.A.S. OLIVIER, S.M. DIPPENAAR, L.F. KHALIL and N.M. MOKGALONG 135–140

DU PREEZ, J.H.

- Parameters for the determination and evaluation of heat stress in dairy cattle in South Africa—J.H. DU PREEZ 263–271

DU PLESSIS, L.

- The influence of *Mannheimia haemolytica* A1 seed culture inoculum cell density on the production of leukotoxin in submerged culture supernatant—M.W. ODENDAAL and L. DU PLESSIS 205–216

- DUNGU, B.**
The effect of a natural maedi-visna virus infection on the productivity of South African sheep—B. DUNGU, J. VORSTER, G.F. BATH and D.W. VERWOERD 87–96
- DURETTE-DESSET, MARIE-CLAUDE**
Molineus cati n. sp. (Nematoda, Trichostrongylina, Molineoidea), a parasite of feral cats, *Felis catus* Linnaeus, 1758 in South Africa—MARIE-CLAUDE DURETTE-DESSET, J. BOOMKER and F.S. MALAN 173–177
- DZIVA, F.**
Capsular serogroups of *Pasteurella multocida* isolated from animals in Zimbabwe—F. DZIVA, K. MOHAN and A. PAWANDIWA 225–228
Pasteurella gallinarum: Zimbabwean experience of a versatile pathogen—K. MOHAN, F. DZIVA and D. CHITAURO 301–305
- FOURIE, L.J.**
Parasites of domestic and wild animals in South Africa. XXXVIII. Ixodid ticks collected from 23 wild carnivore species—I.G. HORAK, L.E.O. BRAACK, L.J. FOURIE and JANE B. WALKER 239–250
- GATHUMA, J.M.**
Application of ultrasonography in prevalence studies of hydatid cysts in goats in north-western Turkana, Kenya and Toposaland, southern Sudan—E.M. NJOROGE, P.M.F. MBITHI, J.M. GATHUMA, T.M. WACHIRA, J.K. MAGAMBO and E. ZEYHLE 251–255
- GATHUMBI, P.K.**
Suspected sheep-associated malignant catarrhal fever in a zero-grazed dairy herd in Kenya—C.M. MULEI, P.K. GATHUMBI and P.G. MBUTHIA 43–47
Biochemical and haematological changes in rats administered an aqueous extract of *Prunus africana* stem bark at various dosage levels—P.K. GATHUMBI, J.W. MWANGI, S.M. NJIRO and G.M. MUGERA 123–128
- GERBER, H.M.**
Oral infectivity of cryopreserved *Haemonchus contortus* infective larvae that recovered relatively slowly when thawed after more than 15 years in liquid nitrogen—J.A. VAN WYK and H.M. GERBER 149–152
- GERDES, G.H.**
The prevalence of different African horsesickness virus serotypes in the Onderstepoort area near Pretoria, during an outbreak of African horsesickness in South Africa in 1995/1996—C.W. BREMER, G.H. GERDES, H. AITCHISON, I. LOUW, R.R. GREYLING and J. WELGEMOED 65–70
- GREYLING, R.R.**
The prevalence of different African horsesickness virus serotypes in the Onderstepoort area near Pretoria, during an outbreak of African horsesickness in South Africa in 1995/1996—C.W. BREMER, G.H. GERDES, H. AITCHISON, I. LOUW, R.R. GREYLING and J. WELGEMOED 65–70
- GROBLER, D.G.**
The rise and fall of tuberculosis in a free-ranging chacma baboon troop in the Kruger National Park—D.F. KEET, N.P.J. KRIEK, R.G. BENGIS, D.G. GROBLER and A. MICHEL 115–122
- GROENEWALD, H.B.**
Morphological changes associated with the development of the rumino-reticulum in growing lambs fed different rations—G.E. SWAN and H.B. GROENEWALD 105–114
- HENTON, M.M.**
Aerobic intestinal flora of wild-caught African dwarf crocodiles *Osteolaemus tetraspis*—F.W. HUCHZERMEYER, M.M. HENTON, J. RILEY and M. AGNAGNA 201–204
- HERING-HAGENBECK, S.F.B.N.**
A check-list of the nematode parasites of South African serpentes (snakes) and sauria (lizards)—S.F.B.N. HERING-HAGENBECK and J. BOOMKER 1–13
- HIRUMI, H.**
Suggested dosage rates of melarsoprol in the treatment of mice experimentally infected with *Trypanosoma brucei gambiense*—P.A. MBATI, K. HIRUMI, N. INOUE, N.H. SITUAKIBANZA and H. HIRUMI 71–74
- HIRUMI, K.**
Suggested dosage rates of melarsoprol in the treatment of mice experimentally infected with *Trypanosoma brucei gambiense*—P.A. MBATI, K. HIRUMI, N. INOUE, N.H. SITUAKIBANZA and H. HIRUMI 71–74

HORAK, I.G.	
Parasites of South African wildlife. XVI. Helminths of some antelope species from the Eastern and Western Cape Provinces—J. BOOMKER, I.G. HORAK, R. WATERMEYER and D.G. BOOYSE	31–41
Parasites of domestic and wild animals in South Africa. XXXVIII. Ixodid ticks collected from 23 wild carnivore species— I.G. HORAK, L.E.O. BRAACK, L.J. FOURIE and JANE B. WALKER	239–250
HUCHZERMEYER, F.W.	
Aerobic intestinal flora of wild-caught African dwarf crocodiles <i>Osteolaemus tetraspis</i> —F.W. HUCHZERMEYER, M.M. HENTON, J. RILEY and M. AGNAGNA	201–204
INOUE, N.	
Suggested dosage rates of melarsoprol in the treatment of mice experimentally infected with <i>Trypanosoma brucei gambiense</i> —P.A. MBATI, K. HIRUMI, N. INOUE, N.H. SITUAKIBANZA and H. HIRUMI	71–74
JAMAL, S.	
The distribution of tsetse (Diptera: Glossinidae) and bovine trypanosomosis in the Matutuine District, Maputo Province, Mozambique—I. SIGAUQUE, P. VAN DEN BOSSCHE, M. MOIANA, S. JAMAL and L. NEVES	167–172
JOSLING, D.	
Ultrastructural study of the luminal surface of the ducts of the epididymis of gallinaceous birds—T.A. AIRE and D. JOSLING	191–199
KAGIRA, J.	
The role of parasitic diseases as causes of mortality in cattle in a high potential area of central Kenya: a quantitative analysis—P.W.N. KANYARI and J. KAGIRA	157–161
KANYARI, P.W.N.	
The role of parasitic diseases as causes of mortality in cattle in a high potential area of central Kenya: a quantitative analysis—P.W.N. KANYARI and J. KAGIRA	157–161
KAPPMEIER, KARIN	
A newly developed odour-baited "H trap" for the live collection of <i>Glossina brevipalpis</i> and <i>Glossina austeni</i> (Diptera: Glossinidae) in South Africa—KARIN KAPPMEIER	15–26
Diurnal activity patterns of <i>Glossina brevipalpis</i> and <i>G. austeni</i> (Diptera: Glossinidae) in South Africa, with reference to season and meteorological factors—KARIN KAPPMEIER	179–189
KEET, D.F.	
The rise and fall of tuberculosis in a free-ranging chacma baboon troop in the Kruger National Park—D.F. KEET, N.P.J. KRIEK, R.G. BENGIS, D.G. GROBLER and A. MICHEL	115–122
KHALIL, L.F.	
Observations on a lesser-known monogenean, <i>Udonella myliobati</i> , from a copepod parasite, <i>Lepeophtheirus natalensis</i> , parasitizing the spotted ragged-tooth shark, <i>Carcharias taurus</i> , from South African waters—P.A.S. OLIVIER, S.M. DIP-PENAAR, L.F. KHALIL and N.M. MOKGALONG	135–140
KOKOSHAROV, T.	
Bacterial colonization and endotoxin activity during experimental acute fowl typhoid in chickens—T. KOKOSHAROV .	297–300
KRIEK, N.P.J.	
The rise and fall of tuberculosis in a free-ranging chacma baboon troop in the Kruger National Park—D.F. KEET, N.P.J. KRIEK, R.G. BENGIS, D.G. GROBLER and A. MICHEL	115–122
Selected chemical parameters in the blood and metals in the organs of the Nile crocodile, <i>Crocodylus niloticus</i> , in the Kruger National Park—D. SWANEPOEL, J. BOOMKER and N.P.J. KRIEK	141–148
LEVINS, R.	
Models for regional heartwater epidemiology in a variable environment—R. LEVINS	163–165
LOSHO, T.C.	
Bovine borreliosis in Botswana—S.P. SHARMA, W. AMANFU and T.C. LOSHO	221–223
LOUW, I.	
The prevalence of different African horsesickness virus serotypes in the Onderstepoort area near Pretoria, during an outbreak of African horsesickness in South Africa in 1995/1996—C.W. BREMER, G.H. GERDES, H. AITCHISON, I. LOUW, R.R. GREYLING and J. WELGEMOED	65–70

- MABIKACHECHE, L.G.**
Evaluation of a commercially available ELISA kit for detection of antibodies to *Anaplasma marginale* and *Anaplasma centrale* in cattle in Australia and Zimbabwe—P.M. BOWLES, J.B. MOLLOY, G.W. BLIGHT, S. SINGH and L.G. MABIKACHECHE 83–86
- MAGAMBO, J.K.**
Application of ultrasonography in prevalence studies of hydatid cysts in goats in north-western Turkana, Kenya and Toposaland, southern Sudan—E.M. NJORGE, P.M.F. MBITHI, J.M. GATHUMA, T.M. WACHIRA, J.K. MAGAMBO and E. ZEYHLE 251–255
- MALAN, F.S.**
Molineus cati n. sp. (Nematoda, Trichostrongylina, Molineoidea), a parasite of feral cats, *Felis catus* Linnaeus, 1758 in South Africa—MARIE-CLAUDE DURETTE-DESSET, J. BOOMKER and F.S. MALAN 173–177
- MARÉ, L.**
Phage types of *Salmonella enterica* serovar Enteritidis isolated in South Africa from 1991–1995—L. MARÉ, M.L. VANDER WALT and L.M.T. DICKS 129–133
- MASHEGO, S.N.**
Occurrence of *Neodiplozoon polycotyleus* Paperna, 1973 (Diplozoidae: Monogenea) in cyprinid fish in South Africa—S.N. MASHEGO 153–154
- MBATI, P.A.**
Leishmania major infections in *Phlebotomus duboscqi* fed on murine models immunized with *L. major* sub-cellular antigens and sandfly gut antigens—P.A. MBATI, C.O. ANJILI, S. ODONGO, P. OGAJA and W. TONU 57–63
Suggested dosage rates of melarsoprol in the treatment of mice experimentally infected with *Trypanosoma brucei gambiense*—P.A. MBATI, K. HIRUMI, N. INOUE, N.H. SITUAKIBANZA and H. HIRUMI 71–74
- MBITHI, P.M.F.**
Application of ultrasonography in prevalence studies of hydatid cysts in goats in north-western Turkana, Kenya and Toposaland, southern Sudan—E.M. NJORGE, P.M.F. MBITHI, J.M. GATHUMA, T.M. WACHIRA, J.K. MAGAMBO and E. ZEYHLE 251–255
- MBUTHIA, P.G.**
Suspected sheep-associated malignant catarrhal fever in a zero-grazed dairy herd in Kenya—C.M. MULEI, P.K. GATHUMBI and P.G. MBUTHIA 43–47
- McCRINDLE, R.I.**
Investigation of biological samples for monofluoroacetate and *Dichapetalum cymosum* poisoning in southern Africa—P.P. MINNAAR, R.I. McCRINDLE, T.W. NAUDÉ and C.J. BOTHA 27–30
- MEKONNEN, S.**
Efficacy of flumethrin 1 % pour-on against ticks on cattle under field condition in Ethiopia—S. MEKONNEN 235–237
- MICHEL, A.**
The rise and fall of tuberculosis in a free-ranging chacma baboon troop in the Kruger National Park—D.F. KEET, N.P.J. KRIEK, R.G. BENGIS, D.G. GROBLER and A. MICHEL 115–122
- MINNAAR, P.P.**
Investigation of biological samples for monofluoroacetate and *Dichapetalum cymosum* poisoning in southern Africa—P.P. MINNAAR, R.I. McCRINDLE, T.W. NAUDÉ and C.J. BOTHA 27–30
- MOHAN, K.**
Capsular serogroups of *Pasteurella multocida* isolated from animals in Zimbabwe—F. DZIVA, K. MOHAN and A. PAWANDIWA 225–228
Pasteurella gallinarum: Zimbabwean experience of a versatile pathogen—K. MOHAN, F. DZIVA and D. CHITAURO 301–305
- MOIANA, M.**
The distribution of tsetse (Diptera: Glossinidae) and bovine trypanosomosis in the Matutuine District, Maputo Province, Mozambique—I. SIGAUQUE, P. VAN DEN BOSSCHE, M. MOIANA, S. JAMAL and L. NEVES 167–172
- MOKGALONG, N.M.**
Observations on a lesser-known monogenean, *Udonella myliobati*, from a copepod parasite, *Lepeophtheirus natalensis*, parasitizing the spotted ragged-tooth shark, *Carcharias taurus*, from South African waters—P.A.S. OLIVIER, S.M. DIPPENAAAR, L.F. KHALIL and N.M. MOKGALONG 135–140

MOLLOY, J.B.	
Evaluation of a commercially available ELISA kit for detection of antibodies to <i>Anaplasma marginale</i> and <i>Anaplasma centrale</i> in cattle in Australia and Zimbabwe—P.M. BOWLES, J.B. MOLLOY, G.W. BLIGHT, S. SINGH and L.G. MABIKACHECHE	83–86
MOSALA, K.P.	
The prevalence of internal and external parasites in pigs of different ages and sexes in Southeast District, Botswana—S.J. NSOSO, K.P. MOSALA, R.T. NDEBELE and S.S. RAMABU	217–220
MUGERA, G.M.	
Biochemical and haematological changes in rats administered an aqueous extract of <i>Prunus africana</i> stem bark at various dosage levels—P.K. GATHUMBI, J.W. MWANGI, S.M. NJIRO and G.M. MUGERA	123–128
MÜLDERS, MARIA S.G.	
Stability and partitioning of closantel and rafoxanide in ruminal fluid of sheep—G.E. SWAN, D.W. OLIVER, J. VAN RENSBURG, H.S. STEYN and MARIA S.G. MÜLDERS	97–103
MULEI, C.M.	
Suspected sheep-associated malignant catarrhal fever in a zero-grazed dairy herd in Kenya—C.M. MULEI, P.K. GATHUMBI and P.G. MBUTHIA	43–47
MUSHI, E.Z.	
Helminth parasites of domestic pigeons (<i>Columba livia domestica</i>) in Sebele, Gaborone, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, R. NDEBELE and R. PANZIRAH	75–76
MWANGI, J.W.	
Biochemical and haematological changes in rats administered an aqueous extract of <i>Prunus africana</i> stem bark at various dosage levels—P.K. GATHUMBI, J.W. MWANGI, S.M. NJIRO and G.M. MUGERA	123–128
NAUDÉ, T.W.	
Investigation of biological samples for monofluoroacetate and <i>Dichapetalum cymosum</i> poisoning in southern Africa—P.P. MINNAAR, R.I. McCRINDLE, T.W. NAUDÉ and C.J. BOTHA	27–30
NDEBELE, R.	
Helminth parasites of domestic pigeons (<i>Columba livia domestica</i>) in Sebele, Gaborone, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, R. NDEBELE and R. PANZIRAH	75–76
NDEBELE, R.T.	
The prevalence of internal and external parasites in pigs of different ages and sexes in Southeast District, Botswana—S.J. NSOSO, K.P. MOSALA, R.T. NDEBELE and S.S. RAMABU	217–220
NEVES, L.	
The distribution of tsetse (Diptera: Glossinidae) and bovine trypanosomosis in the Matutuine District, Maputo Province, Mozambique—I. SIGAUQUE, P. VAN DEN BOSSCHE, M. MOIANA, S. JAMAL and L. NEVES	167–172
NJIRO, S.M.	
Biochemical and haematological changes in rats administered an aqueous extract of <i>Prunus africana</i> stem bark at various dosage levels—P.K. GATHUMBI, J.W. MWANGI, S.M. NJIRO and G.M. MUGERA	123–128
NJOROGE, E.M.	
Application of ultrasonography in prevalence studies of hydatid cysts in goats in north-western Turkana, Kenya and Toposaland, southern Sudan—E.M. NJOROGE, P.M.F. MBITHI, J.M. GATHUMA, T.M. WACHIRA, J.K. MAGAMBO and E. ZEYHLE	251–255
NSOSO, S.J.	
The prevalence of internal and external parasites in pigs of different ages and sexes in Southeast District, Botswana—S.J. NSOSO, K.P. MOSALA, R.T. NDEBELE and S.S. RAMABU	217–220
ODENDAAL, M.W.	
The influence of <i>Mannheimia haemolytica</i> A1 seed culture inoculum cell density on the production of leukotoxin in submerged culture supernatant—M.W. ODENDAAL and L. DU PLESSIS	205–216
ODONGO, S.	
<i>Leishmania major</i> infections in <i>Phlebotomus duboscqi</i> fed on murine models immunized with <i>L. major</i> sub-cellular antigens and sandfly gut antigens—P.A. MBATI, C.O. ANJILI, S. ODONGO, P. OGAJA and W. TONU	57–63

- OGAJA, P.**
Leishmania major infections in *Phlebotomus duboscqi* fed on murine models immunized with *L. major* sub-cellular antigens and sandfly gut antigens—P.A. MBATI, C.O. ANJILI, S. ODONGO, P. OGAJA and W. TONU 57–63
- OLIVER, D.W.**
 Stability and partitioning of closantel and rafoxanide in ruminal fluid of sheep—G.E. SWAN, D.W. OLIVER, J. VAN RENSBURG, H.S. STEYN and MARIA S.G. MÜLDERS 97–103
- OLIVIER, P.A.S.**
 Observations on a lesser-known monogenean, *Udonella myliobati*, from a copepod parasite, *Lepeophtheirus natalensis*, parasitizing the spotted ragged-tooth shark, *Carcharias taurus*, from South African waters—P.A.S. OLIVIER, S.M. DIP-PENAAR, L.F. KHALIL and N.M. MOKGALONG 135–140
- OLUWANIYI, J.O.**
 Differential expression of surface membrane antigens on bovine monocytes activated with recombinant cytokines and during *Trypanosoma congolense* infection—V.O. TAIWO, V.O. ANOSA and J.O. OLUWANIYI 289–296
- PANZIRAH, R.**
 Helminth parasites of domestic pigeons (*Columba livia domestica*) in Sebele, Gaborone, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, R. NDEBELE and R. PANZIRAH 75–76
- PAWANDIWA, A.**
 Capsular serogroups of *Pasteurella multocida* isolated from animals in Zimbabwe—F. DZIVA, K. MOHAN and A. PAWANDIWA, A. 225–228
- PUTTERILL, J.F.**
 Oribatid mites (Acari, Oribatida) as intermediate hosts of tapeworms of the Family Anoplocephalidae (Cestoda) and the transmission of *Moniezia expansa* cysticercoids in South Africa—R. SCHUSTER, L. COETZEE and J.F. PUTTERILL 49–55
 Studies on the genus *Setaria* Viborg, 1795 in South Africa. I. *Setaria africana* (Yeh, 1959)—R. WATERMEYER, J. BOOMKER and J.F. PUTTERILL 229–234
- RAMABU, S.S.**
 The prevalence of internal and external parasites in pigs of different ages and sexes in Southeast District, Botswana—S.J. NSOSO, K.P. MOSALA, R.T. NDEBELE and S.S. RAMABU 217–220
- RILEY, J.**
 Aerobic intestinal flora of wild-caught African dwarf crocodiles *Osteolaemus tetraspis*—F.W. HUCHZERMEYER, M.M. HENTON, J. RILEY and M. AGNAGNA 201–204
- SCHUSTER, L.**
 Oribatid mites (Acari, Oribatida) as intermediate hosts of tapeworms of the Family Anoplocephalidae (Cestoda) and the transmission of *Moniezia expansa* cysticercoids in South Africa—R. SCHUSTER, L. COETZEE and J.F. PUTTERILL 49–55
- SEWANI-RUSIKE, C.R.**
 Fertility in rats immunized with steroid-free bovine follicular fluid—C.R. SEWANI-RUSIKE and C. DAKWA 257–262
- SHARMA, S.P.**
 Bovine borreliosis in Botswana—S.P. SHARMA, W. AMANFU and T.C. LOSHO 221–223
- SIGAUQUE, I.**
 The distribution of tsetse (Diptera: Glossinidae) and bovine trypanosomosis in the Matutuine District, Maputo Province, Mozambique—I. SIGAUQUE, P. VAN DEN BOSSCHE, M. MOIANA, S. JAMAL and L. NEVES 167–172
- SINGH, S.**
 Evaluation of a commercially available ELISA kit for detection of antibodies to *Anaplasma marginale* and *Anaplasma centrale* in cattle in Australia and Zimbabwe—P.M. BOWLES, J.B. MOLLOY, G.W. BLIGHT, S. SINGH and L.G. MABIKACHECHE 83–86
- SITUAKIBANZA, N.H.**
 Suggested dosage rates of melarsoprol in the treatment of mice experimentally infected with *Trypanosoma brucei gambiense*—P.A. MBATI, K. HIRUMI, N. INOUE, N.H. SITUAKIBANZA and H. HIRUMI 71–74
- STEYN, H.S.**
 Stability and partitioning of closantel and rafoxanide in ruminal fluid of sheep—G.E. SWAN, D.W. OLIVER, J. VAN RENSBURG, H.S. STEYN and MARIA S.G. MÜLDERS 97–103

- SWAN, G.E.**
 Stability and partitioning of closantel and rafoxanide in ruminal fluid of sheep—G.E. SWAN, D.W. OLIVER, J. VAN RENSBURG, H.S. STEYN and MARIA S.G. MÜLDERS 97–103
 Morphological changes associated with the development of the rumino-reticulum in growing lambs fed different rations—G.E. SWAN and H.B. GROENEWALD 105–114
- SWANEPOEL, J.**
 Selected chemical parameters in the blood and metals in the organs of the Nile crocodile, *Crocodylus niloticus*, in the Kruger National Park—D. SWANEPOEL, J. BOOMKER and N.P.J. KRIEK 141–148
- TAIWO, V.O.**
In vitro erythrophagocytosis by cultured macrophages stimulated with extraneous substances and those isolated from the blood, spleen and bone marrow of Boran and N'Dama cattle infected with *Trypanosoma congolense* and *T. vivax*—V.O. TAIWO and V.O. ANOSA 273–287
 Differential expression of surface membrane antigens on bovine monocytes activated with recombinant cytokines and during *Trypanosoma congolense* infection—V.O. TAIWO, V.O. ANOSA and J.O. OLUWANIYI 289–296
- TONU, W.**
Leishmania major infections in *Phlebotomus duboscqi* fed on murine models immunized with *L. major* sub-cellular antigens and sandfly gut antigens—P.A. MBATI, C.O. ANJILI, S. ODONGO, P. OGAJA and W. TONU 57–63
- VAN DEN BOSSCHE, P.**
 The distribution of tsetse (Diptera: Glossinidae) and bovine trypanosomosis in the Matutuine District, Maputo Province, Mozambique—I. SIGAUQUE, P. VAN DEN BOSSCHE, M. MOIANA, S. JAMAL and L. NEVES 167–172
- VAN DER WALT, M.L.**
 Phage types of *Salmonella enterica* serovar Enteritidis isolated in South Africa from 1991–1995—L. MARÉ, M.L. VAN DER WALT and L.M.T. DICKS 129–133
- VAN RENSBURG, J.**
 Stability and partitioning of closantel and rafoxanide in ruminal fluid of sheep—G.E. SWAN, D.W. OLIVER, J. VAN RENSBURG, H.S. STEYN and MARIA S.G. MÜLDERS 97–103
- VAN WYK, J.A.**
 Oral infectivity of cryopreserved *Haemonchus contortus* infective larvae that recovered relatively slowly when thawed after more than 15 years in liquid nitrogen—J.A. VAN WYK and H.M. GERBER 149–152
- VERWOERD, D.W.**
 The effect of a natural maedi-visna virus infection on the productivity of South African sheep—B. DUNGU, J. VORSTER, G.F. BATH and D.W. VERWOERD 87–96
- VORSTER, J.**
 The effect of a natural maedi-visna virus infection on the productivity of South African sheep—B. DUNGU, J. VORSTER, G.F. BATH and D.W. VERWOERD 87–96
- WACHIRA, T.M.**
 Application of ultrasonography in prevalence studies of hydatid cysts in goats in north-western Turkana, Kenya and Toposaland, southern Sudan—E.M. NJOROGE, P.M.F. MBITHI, J.M. GATHUMA, T.M. WACHIRA, J.K. MAGAMBO and E. ZEYHLE 251–255
- WALKER, JANE B.**
 Parasites of domestic and wild animals in South Africa. XXXVIII. Ixodid ticks collected from 23 wild carnivore species—I.G. HORAK, L.E.O. BRAACK, L.J. FOURIE and JANE B. WALKER 239–250
- WATERMEYER, R.**
 Studies on the genus *Setaria* Viborg, 1795 in South Africa. I. *Setaria africana* (Yeh, 1959)—R. WATERMEYER, J. BOOMKER and J.F. PUTTERILL 229–234
 Parasites of South African wildlife. XVI. Helminths of some antelope species from the Eastern and Western Cape Provinces—J. BOOMKER, I.G. HORAK, R. WATERMEYER and D.G. BOOYSE 31–41
- WELGEMOED, J.**
 The prevalence of different African horsesickness virus serotypes in the Onderstepoort area near Pretoria, during an outbreak of African horsesickness in South Africa in 1995/1996—C.W. BREMER, G.H. GERDES, H. AITCHISON, I. LOUW, R.R. GREYLING and J. WELGEMOED 65–70

ZEYHLE, E.

- Application of ultrasonography in prevalence studies of hydatid cysts in goats in north-western Turkana, Kenya and Toposaland, southern Sudan—E.M. NJOROGI, P.M.F. MBITHI, J.M. GATHUMA, T.M. WACHIRA, J.K. MAGAMBO and E. ZEYHLE 251–255