

# The abundance of *Ixodes ricinus* ticks depends on tree species composition and shrub cover

W. TACK<sup>1\*</sup>, M. MADDER<sup>2,3</sup>, L. BAETEN<sup>1,4</sup>, P. DE FRENNE<sup>1,5</sup> and K. VERHEYEN<sup>1</sup>

<sup>1</sup> *Laboratory of Forestry, Department of Forest and Water Management, Ghent University, Geraardsbergsesteenweg 267, 9090 Melle-Gontrode, Belgium*

<sup>2</sup> *Unit of Vector Biology, Department of Biomedical Sciences, Institute of Tropical Medicine, Nationalestraat 155, 2000 Antwerp, Belgium*

<sup>3</sup> *Department of Veterinary Tropical Diseases, Faculty of Veterinary Science, University of Pretoria, Onderstepoort, 0110, South Africa*

<sup>4</sup> *Terrestrial Ecology Unit, Department of Biology, Ghent University, K. L. Ledeganckstraat 35, 9000 Ghent, Belgium*

<sup>5</sup> *Forest Ecology and Conservation Group, Department of Plant Sciences, University of Cambridge, Downing Street, Cambridge CB2 3EA, UK*

(Received 26 January 2012; revised 14 March 2012; accepted 14 March 2012; first published online 13 April 2012)

## SUMMARY

The mainstream forestry policy in many European countries is to convert coniferous plantations into (semi-natural) deciduous woodlands. However, woodlands are the main habitat for *Ixodes ricinus* ticks. Therefore, assessing to what extent tick abundance and infection with *Borrelia* spirochetes are affected by forest composition and structure is a prerequisite for effective prevention of Lyme borreliosis. We selected a total of 25 pine and oak stands, both with and without an abundant shrub layer, in northern Belgium and estimated tick abundance between April and October 2008–2010. Additionally, the presence of deer beds was used as an indicator of relative deer habitat use. *Borrelia* infections in questing nymphs were determined by polymerase chain reactions. The abundance of larvae, nymphs, and adults was higher in oak stands compared to pine stands and increased with increasing shrub cover, most likely due to differences in habitat use by the ticks' main hosts. Whereas tick abundance was markedly higher in structure-rich oak stands compared to homogeneous pine stands, the *Borrelia* infection rates in nymphs did not differ significantly. Our results indicate that conversion towards structure-rich deciduous forests might create more suitable tick habitats, but we were unable to detect an effect on the infection rate.

Key words: *Ixodes ricinus*, *Borrelia burgdorferi*, tick, roe deer, habitat, forest conversion.

## INTRODUCTION

In recent decades, Lyme borreliosis has become a subject of international concern because of the increasing number of human cases diagnosed each year (World Health Organization, 2004; Bacon *et al.* 2008). The disease is caused by spirochetes belonging to the *Borrelia burgdorferi* sensu lato (s.l.) complex, which is maintained in an enzootic cycle involving mainly ticks of the *Ixodes ricinus* species complex (Acari: Ixodidae) and numerous vertebrates (Piesman and Gern, 2004). Several bird and small mammal species, rodents in particular, serve as hosts for the larval and nymphal stages and are important reservoirs for *Borrelia* spirochetes (Kurtenbach *et al.* 1998; Humair *et al.* 1999; Comstedt *et al.* 2006). Ticks may acquire the spirochetes by feeding on infected hosts, maintain the infection to the subsequent life stages through trans-stadial transmission, and transmit the infection to other hosts or humans during the next bloodmeal.

\* Corresponding author: Laboratory of Forestry, Department of Forest and Water Management, Ghent University, Geraardsbergsesteenweg 267, 9090 Melle-Gontrode, Belgium. Tel: +32 (0) 9 264 90 30. Fax: +32 (0) 9 264 90 92. E-mail: Wesley.Tack@UGent.be

The risk of human infection is typically associated with forested areas, as forests provide ticks with access to a broad range of vertebrate hosts and favourable environmental conditions for tick survival (Gray, 1998). It is commonly assumed that the observed increase in Lyme borreliosis incidence is due to an actual rise in the number of infections and not only to an enhanced surveillance and awareness of the disease. This has been attributed to various factors, particularly to human encroachment into forested areas and habitat modifications, resulting in a closer human contact with ticks and an increase in the abundance and range of tick populations. For instance, the marked increase of deer populations throughout Europe and North America during the latter half of the twentieth century (Gill, 1990; Cederlund *et al.* 1998), largely caused by changes in land use and land cover (e.g., reforestation), has been ascribed a major role in the emergence and spread of Lyme borreliosis (Spielman, 1994; Sood *et al.* 2011). Although incompetent reservoirs for *B. burgdorferi*, deer are the preferred hosts of adult *Ixodes* ticks and play an important role in their reproductive success. Moreover, both tick abundance and infection prevalence in ticks are favoured by forest fragmentation (Allan *et al.* 2003; Brownstein *et al.* 2005; Halos *et al.*

2010; Tack *et al.* 2012), since deer and rodents benefit from the presence of forest edge habitat and abundant ecotonal vegetation (Tufto *et al.* 1996; Saïd and Servanty, 2005; Boyard *et al.* 2008). In Sweden, *I. ricinus* has become more abundant and has expanded its range during the last 3 decades, which is probably caused by changes in climate (increased duration of the vegetation period and milder winters) and increased abundances of deer (Jaenson *et al.* 2012).

In many European countries, the conversion of monospecific coniferous forests into mixed, structure-rich forests dominated by native broad-leaved species has become a major objective of sustainable, multipurpose forest management, with the aim of optimizing the production of various goods and ecosystem services (Olsthoorn *et al.* 1999; Spiecker *et al.* 2004). However, deciduous forests, especially those harbouring significant numbers of cervids, are generally considered to be ideal habitats for *I. ricinus*, which is the most common tick species associated with Lyme borreliosis in Europe (Gray, 1998). By altering the forest composition and structure, these forest management activities involve a large-scale land-use change that might influence the suitability of forests for ticks and, consequently, might influence the epidemiology of tick-borne diseases (i.e., forest conversion creating an ecosystem dysfunction). Yet, there have been relatively few studies addressing the variation in tick abundance between forest types. While it has recently been quantitatively shown that the abundance of *I. ricinus* ticks is higher in oak stands compared to pine stands and increases with increasing shrub cover (Tack *et al.* 2012), little is known on the longer-term temporal variation and on the effects of forest composition and structure on the resulting *Borrelia* infection rate. Here, we selected a total of 25 pine (*Pinus* sp.) and oak (*Quercus* sp.) stands, both with and without a substantial shrub layer, and sampled *I. ricinus* tick populations between April and October in 3 successive years in northern Belgium to describe the spatiotemporal variation in the abundance of larvae, nymphs, and adults and to relate this variation to forest composition and structure. Additionally, habitat use by cervids was determined by counting the number of deer beds. *Borrelia burgdorferi* s.l. spirochete infections in tick nymphs, the life stage predominantly responsible for pathogen transmission, were determined by polymerase chain reactions to assess the potential impact of forest conversion on the infection prevalence.

## MATERIALS AND METHODS

### Study area

The study was conducted at 2 forest sites in the Campine ecoregion in northern Belgium. Forest site

A (51°17' N, 5°12' E) was located near the border with the Netherlands in the municipality Postel and forest site B (51°2' N, 4°58' E) was located approximately 30 km to the south in the municipalities Herselt and Tessenderlo. The climate is sub-atlantic: the mean annual precipitation amounts to 799 mm and is evenly distributed throughout the year, with mean monthly precipitation ranging from 53 mm in March to 79 mm in July. The mean annual temperature is 9.0 °C, with minimum and maximum mean monthly temperatures of 1.4 °C and 16.7 °C in January and July, respectively (Royal Meteorological Institute of Belgium, URL <http://www.kmi.be/>, accessed November 18, 2011). The region's characteristic forests are pine plantations—mainly consisting of Scots pine (*Pinus sylvestris*) and, to a lesser extent, Corsican pine (*P. nigra* subsp. *laricio*)—on nutrient-poor and acidic sandy soils. The pine stands are interspersed with deciduous stands of pedunculate oak (*Quercus robur*), red oak (*Q. rubra*), common beech (*Fagus sylvatica*), silver birch (*Betula pendula*), and downy birch (*B. pubescens*) (Waterinckx and Roelandt, 2001). Most forests were established in the nineteenth and first half of the twentieth century on former heathlands, which once formed an important component of the traditional agricultural system and covered most of the landscape. The then prevailing microclimatic conditions (temperature and moisture) were most likely limiting for tick survival, which is strongly supported by recent studies carried out in heathlands (Estrada-Peña, 2001; Lindström and Jaenson, 2003; Wielinga *et al.* 2006). However, the large-scale afforestation and the subsequent rise in deer populations probably made this region suitable for tick population establishment and survival. Nowadays, the Campine region is known as a hotspot area in Belgium for Lyme borreliosis (Linard *et al.* 2007). Local vertebrate hosts of nymphal and female ticks are large and medium-sized mammals such as roe deer (*Capreolus capreolus*), red fox (*Vulpes vulpes*), European hare (*Lepus europaeus*), European hedgehog (*Erinaceus europaeus*), least weasel (*Mustela nivalis*), European pole cat (*Mustela putorius*), and red squirrel (*Sciurus vulgaris*). Very common small mammalian hosts for larvae include pygmy shrew (*Sorex minutus*), common shrew (*Sorex araneus*), wood mouse (*Apodemus sylvaticus*), bank vole (*Myodes glareolus*), and field vole (*Microtus agrestis*) (Verkem *et al.* 2003; Tack *et al.* unpublished data).

### Forest stand selection

At each forest site, 6 pine stands and 6 oak stands were selected on poor, sandy soils with half of the stands having little or no shrub layer (<15% shrub layer cover in the 1–7 m height class) and the other half having a well-developed shrub layer (>50% cover). An additional oak stand was selected with low shrub

cover at forest site B. In summary, ticks were sampled in 25 forest stands (12 in forest site A and 13 in forest site B) and in 4 distinct forest stand types: pine stands and oak stands, both with and without a substantial shrub layer. The relative contribution of *Pinus* sp. (*P. sylvaticus* or *P. nigra*) or *Quercus* sp. (mainly *Q. robur*) to the total estimated canopy cover of the tree layer (>7 m) was greater than or equal to 80% in each pine and oak stand, respectively. In each forest stand, the percentage cover of the shrub layer (1–7 m) and herb layer (<1 m) was estimated visually. Shrub cover estimates were very comparable between pine and oak stands at both forest sites. The structure-rich oak stands had an average shrub cover of 66.7% at site A and 70.0% at site B, and the pine stands had an average shrub cover of 70.0% at site A and 58.3% at site B. The shrub layer mainly consisted of alder buckthorn (*Frangula alnus*), black cherry (*Prunus serotina*), and rowan (*Sorbus aucuparia*) in the pine stands and alder buckthorn, pedunculate oak, and sycamore (*Acer pseudoplatanus*) in the oak stands. The herbaceous layer was dominated either by wavy hair-grass (*Deschampsia flexuosa*), purple moor-grass (*Molinia caerulea*), broad buckler-fern (*Dryopteris dilatata*), or bilberry (*Vaccinium myrtillus*), providing a comparable blanket contact when drag sampling for ticks (see below). Forest stands with a dense bracken (*Pteridium aquilinum*) understory were avoided because this vegetation can seriously impede tick sampling (Tack *et al.* 2011). Because of the height and rough vegetation surface of bracken, ticks are easily brushed off the blanket, causing tick abundance to be underestimated. However, the sampled vegetation types are representative for the Campine region so we do not expect our sampling procedure to greatly affect the results.

#### Sampling strategy

Tick sampling was carried out between April and October in 2008, 2009, and 2010 for a total of 11 occasions at site A and 12 occasions at site B (12 stands  $\times$  11 occasions + 13 stands  $\times$  12 occasions = 288). Sampling consisted of dragging a white flannel blanket (1  $\times$  1 m<sup>2</sup>) over the herbaceous vegetation and litter. In each forest stand and at each sampling occasion, we performed 6  $\times$  1 min blanket drags (each extending a distance of *ca.* 25 m) at random and recorded the air temperature and relative humidity 3 times at a height of 1.25 m above the soil surface, using a portable digital temperature and relative air humidity meter (DM509, Eijkelkamp Agrisearch Equipment, Giesbeek, The Netherlands). Sampling was always performed on dry (no rain) and non-windy days (<2 Bft) during day time (between 10:00 am and 05:00 pm) when the vegetation was dry. To avoid time of day and changing meteorological conditions as a source of bias, the 4 forest stand types were sampled in random

order on each sampling day. After each transect, larvae, nymphs, and adults were removed from the blanket using forceps and stored in vials containing 70% ethanol for later identification and counting. The ticks were counted and identified morphologically with a stereo-microscope using the identification keys of Hillyard (1996).

Additionally, the number of fecal pellet groups and beds of roe deer were counted at each sampling occasion, along the same transects used for tick sampling. Pellet-group counting is a widely used method for assessing habitat use by deer. In our study, however, the number of pellet groups counted was too small (only 21 pellet groups in total) for proper analysis. Instead, we have used the number of deer beds in each forest stand type to examine differences in habitat selection for bedding sites (Smith *et al.* 1986; B  r   *et al.* 2006). Deer beds were easily detectable in the sandy soil of the study area and were distinguished as oval depressions in the soil or as flattened areas of vegetation, often accompanied by other signs of roe deer (e.g., hoof prints, hair).

#### Identification of *Borrelia* infections

Twenty pooled samples per forest site per year (20 samples  $\times$  2 sites  $\times$  3 years = 120), with each sample consisting of 5 nymphs, were used for further molecular analyses for the presence of *B. burgdorferi* s.l. spirochetes. We did not identify the *Borrelia* genospecies. Instead, only screening up to species level was performed to get an idea of the overall infection prevalence. For each forest site and each year, 10 samples consisted of nymphs collected in pine stands with low shrub cover while the other 10 samples were collected in oak stands with high shrub cover. Potential differences in infection prevalence are most likely to occur between these two contrasting forest stand types. DNA was extracted using the method of Boom *et al.* (1990). This method is based on the lysing and nuclease-inactivating properties of proteinase K, together with the nucleic acid-binding properties of silica particles. A standard PCR amplification was performed in 25  $\mu$ l reaction mixtures containing 5  $\mu$ l of the extracted DNA, 1.65 mM MgCl<sub>2</sub>, 0.2 mM of all 4 dNTPs, 10 pM of 2 primers (BorrSLOspAF/BorrSLOspAR) (Demaerschack *et al.* 1995), 1 Utaq polymerase enzyme (Promega), and 1  $\mu$ l of Yellow SubTM (GENEO Bioproducts, Hamburg, Germany). After a hot start of 10 s at 84  $^{\circ}$ C, an initiation of 4 min at 92  $^{\circ}$ C was performed and followed by a 40-cycle denaturation-hybridization-elongation step (30 s at 92  $^{\circ}$ C, 45 s at 58  $^{\circ}$ C, and 60 s at 72  $^{\circ}$ C). The PCR ended with an extension step of 10 min at 72  $^{\circ}$ C. Then 5  $\mu$ l of each reaction mixture were mixed with 2  $\mu$ l of loading buffer and loaded onto 2% agarose gels (Sigma) to be examined for the presence

of DNA fragments. A 1.5 kb DNA ladder (MBI Fermentas, Lithuania) was loaded on every gel. The samples were run for 20 min at 100 V, stained in ethidium bromide for 30 min, washed under running tap water, and photographed under UV illumination.

### Statistical analysis

Questing tick abundance, expressed as the number of ticks collected per 100 m<sup>2</sup>, was first log<sub>10</sub>(n+1) transformed to approach normality, which was verified using the Kolmogorov-Smirnov test. Subsequently, log-transformed tick abundances were modelled with linear mixed models using the *lmer*-function of the *lme4*-library (Bates *et al.* 2011) in R 2.13.0 (R Development Core Team, 2011). Data for each life stage (larva, nymph, and adult) were analysed separately. Models included tree species (pine *vs* oak), shrub cover (in%), herb cover (in%), year, and all their two-way interactions as fixed effects and forest stand (nested within forest site (A or B)) and sampling occasion as non-nested random-effect terms. To analyse the effects of tree species, shrub cover, year, and all their two-way interactions on the presence of roe deer (scored as 1 or 0 depending on whether deer beds were (1) or were not (0) encountered in the forest stand while dragging), we applied a generalised linear mixed model (GLMM) with similar random-effects structure as above, but with a binomial error distribution and logit link function. Analysis of nymphal infection with *B. burgdorferi* s.l. (pooled samples of nymphs infected (1) or not (0)) were also performed with a GLMM with binomial error distribution and logit link function. This model included forest type (pine stands with low shrub cover *vs* oak stands with high shrub cover), year, and their interaction term as fixed effects and forest stand (nested within forest site) as random-effect term. We always compared all possible models (i.e., build by each combination of the fixed-effects terms) using Akaike's Information Criterion, adjusted for sample size (AIC<sub>C</sub>) (Hurvich and Tsai, 1989). The ΔAIC<sub>C</sub> of a model was then calculated as the difference in AIC<sub>C</sub> value for that model and the model with the lowest AIC<sub>C</sub> value (best fit to the data). Models with ΔAIC<sub>C</sub> ≤ 4 were considered equivalent (Bolker, 2008). To determine the relative importance of the explanatory variables, we used the sum of Akaike weights of the set of all top models (ΔAIC<sub>C</sub> ≤ 4) in which the variable appeared (Burnham and Anderson, 2002). The Akaike weight reflects the weight of evidence in support of a particular model relative to the entire model set, and varies from 0 (no support) to 1 (complete support). Finally, the parameter values of the model with the lowest AIC<sub>C</sub> value were estimated with restricted maximum likelihood estimation.

## RESULTS

In total, 110 770 *I. ricinus* ticks were collected, of which 89 017 were larvae, 18 685 were nymphs, and 3 068 were adults (1 634 males and 1 434 females). During tick collection, the air temperature ranged from 7.1 °C to 31.7 °C and the relative humidity ranged from 28.1% to 92.6%. The mean ± standard error of the number of ticks collected per 100 m<sup>2</sup> was 206.1 ± 20.8 larvae (range 0 to 4 263), 43.3 ± 2.1 nymphs (range 1 to 215), and 7.1 ± 0.4 adults (range 0 to 44). On each sampling occasion, all 3 life stages were active and ticks were found questing in all 25 forest stands studied. In May 2009, a very high number of larvae was collected along a single transect in one of the oak stands with high shrub cover, which resulted in a peak in larval activity in May (Fig. 1a). This high variance in larval abundance was not unexpected and reflects the limited dispersal capability of larvae after emergence from the egg mass, consisting of up to 2 000 eggs (Jongejan, 2001). By considering this single transect as an outlier, questing larvae showed a summer peak (August) each year. Nymphs were active throughout the study period without displaying a clear peak (Fig. 1b). Adult tick abundance peaked in spring (April–May) each year and steadily declined in summer (Fig. 1c). Our data were not suited to study seasonal variation in tick abundance, but our results are in line with those of Gassner *et al.* (2011), who examined the temporal dynamics of *I. ricinus* in a neighbouring country, The Netherlands.

For both larvae and adults, the best model explaining the variation in tick abundance included tree species and shrub cover as explanatory variables (Table 1). For adults, a second closely competing model also included a tree species by year interaction term. The best model for nymphs included tree species, shrub cover, and year, whereas the second best model included only tree species and year (Table 1). Herb cover did not appear in any of the top models. Tree species, on the other hand, was present in all top models of each life stage and was therefore the variable with the highest relative importance in explaining tick abundance (Table 2). The temporal fluctuations in tick abundance were very similar in oak and pine stands, but the mean abundance was consistently higher in the oak stands (Fig. 1a–c; Table 3). Larvae, nymphs, and adults were on average 3.3, 1.6, and 1.5 times more abundant in the oak stands. Shrub cover was also a variable of high relative importance (Table 2) and had a positive effect on tick abundance (Table 3). On each sampling occasion, the mean number of ticks collected was higher in forest stands with high shrub cover compared to stands with low shrub cover. Overall, the number of larvae, nymphs, and adults was 2.1, 1.5, and 1.8 times higher in forest stands with high shrub cover (> 50% cover) (Fig. 2a–c). Hence, mean tick




Fig. 1. Mean number of *Ixodes ricinus* larvae, nymphs, and adults (a–c) and mean number of deer beds (d) in pine and oak stands between May and October in 3 successive years. The results from the 2 forest sites were pooled. Error bars denote the standard error of the mean. Note the difference in values on the y-axis.

abundance was lowest in pine stands with low shrub cover ( $43.3 \pm 8.2$  larvae,  $20.7 \pm 2.0$  nymphs, and  $3.8 \pm 0.5$  adults per  $100 \text{ m}^2$ ) and highest in oak stands with high shrub cover ( $418.9 \pm 72.2$  larvae,  $61.6 \pm 4.9$  nymphs, and  $11.1 \pm 1.0$  adults per  $100 \text{ m}^2$ ) (Fig. 2a–c).

A very similar pattern was observed regarding the number of deer beds we encountered during tick sampling (Figs 1d and 2d). The best model explaining the presence of deer beds included tree species, shrub cover, and year (Table 1), with the first two being the variables with the highest relative importance (Table 2). The probability of encountering deer beds was significantly higher in oak stands ( $n = 288$ ,  $P = 0.006$ ) and in forest stands with high shrub cover ( $n = 288$ ,  $P = 0.015$ ) (Table 3). The mean number of deer beds was 1.6 times higher in forest stands with high shrub cover and twice as high in oak stands, which resulted in 4 times as many deer beds in oak stands with high shrub cover compared to pine stands with low shrub cover.

*Borrelia*-positive nymphs were found each year at both forest sites. The average infection rate with *B. burgdorferi* s.l. was 8.3% (95% confidence interval: 4.8–13.2%) in 2008, 11.3% (7.0–16.9%) in 2009, and

6.2% (3.4–10.7%) in 2010. A similar infection rate was observed at both forest sites in the first 2 years of our study, but the infection rate at site A (1.0%) was considerably lower compared to site B in 2010 (12.9%). No significant difference in infection rate was observed between the homogeneous pine stands and the structure-rich oak stands ( $n = 120$ ,  $P = 0.850$ ). The average infection rate was 8.3% (5.4–12.2%) in the pine stands and 8.7% (5.7–12.8%) in the oak stands.

#### DISCUSSION

Our results show that tree species composition and vertical structure are important variables in explaining tick abundance in forests. The abundance of all 3 life stages was higher in oak stands compared to pine stands, and increased with increasing shrub cover. Interestingly, this pattern was observed at both forest sites and on almost every sampling occasion. So, although some annual and seasonal fluctuation in tick numbers occurred, the mean tick abundance was always lowest in the homogeneous pine stands and almost always highest in the structure-rich oak

Table 1. Model selection statistics for the analyses of effects of tree species (T), shrub layer cover (S), and year (Y) on the abundance of *Ixodes ricinus* larvae, nymphs, and adults and on the presence of deer beds

( $\Delta AIC_C$ : the difference in values of the corrected Akaike Information Criterion ( $AIC_C$ ) between a model and the best model having the lowest  $AIC_C$  value;  $w$ : Akaike weight, indicating relative support for the model.)

Response	Model	D.F.	$\Delta AIC_C$	$w$
Larvae	T+S	7	—	0.630
	T+S+Y	9	2.31	0.198
	T	6	2.60	0.172
Nymphs	T+S+Y	9	—	0.684
	T+Y	8	1.54	0.316
Adults	T+S	7	—	0.490
	T×Y+S	11	1.17	0.273
	T+S+Y	9	2.75	0.124
	T	6	2.94	0.113
Deer beds	T+S+Y	8	—	0.311
	T+S	6	0.42	0.253
	T×S+Y	9	2.04	0.112
	T×S	7	2.43	0.092
	T×Y+S	10	3.00	0.070
	T+Y	7	3.13	0.065
	T	5	3.55	0.053
T+S×Y	10	3.89	0.044	

stands. On average, the abundance of larvae, nymphs, and adults was 9.7, 3.0 and 2.9 times higher in the oak stands with high shrub cover than in the pine stands with no or little shrub cover, while intermediate abundances were recorded in the 2 remaining forest stand types. The observed differences in tick abundance between the forest stand types must not necessarily depend directly on differences in tree species composition or structure, but may rather be caused by differences in activity of host animals. Our observations from deer bed counts indicate that roe deer were more often present in oak stands and in stands with high shrub cover, most likely because of the availability of high-quality forage and shelter. The importance of deer in maintaining tick populations has been stressed in several European studies (Gray *et al.* 1992; Pichon *et al.* 1999; Ruiz-Fons and Gilbert, 2010). Being the most common large mammals in the study area, roe deer are almost certainly the most important hosts for adult ticks and, therefore, their habitat use largely determines the location where engorged female ticks drop off and lay eggs. The immature stages (larvae and nymphs) also feed on large mammals such as roe deer, but they generally feed on small to medium-sized mammals and birds. Rodents, such as bank vole and wood mouse, have been identified by several authors as key hosts for larval ticks (Tälleklint and Jaenson, 1997; Humair *et al.* 1999; Estrada-Peña *et al.* 2005). These rodent species, together with other mammal

Table 2. Relative importance of each explanatory variable, calculated across all top models ( $\Delta AIC_C \leq 4$ , see Table 1) in which the variable appeared

Variable	Larvae	Nymphs	Adults	Deer beds
T	1.000	1.000	1.000	1.000
S	0.828	0.684	0.887	0.882
Y	0.198	1.000	0.397	0.602
T×S	0.000	0.000	0.000	0.204
T×Y	0.000	0.000	0.273	0.070
S×Y	0.000	0.000	0.000	0.044

species such as foxes and hedgehogs, are common in the study area and provide immature ticks the opportunity to successfully obtain a bloodmeal and develop into the next life stage, which explains the relatively high nymphal and adult abundances in our study.

Besides being important hosts for immature ticks, small mammals and birds are also important reservoir hosts for *Borrelia* spirochetes. *Borrelia afzelii* has been associated with mice, voles, and red squirrels, *B. burgdorferi* sensu stricto with red squirrels, and *B. garinii* and *B. valaisiana* mainly with birds (Humair and Gern, 1998; Kurtenbach *et al.* 1998; Humair *et al.* 1999; Hanincová *et al.* 2003). The different genospecies tend to cause distinct clinical manifestations affecting different systems (van Dam *et al.* 1993) and, thus, the vertebrate host composition will determine not only the density of *Borrelia*-infected ticks but also the relative risk of different clinical forms of Lyme borreliosis. We did not identify the *Borrelia* genospecies, which could be considered a shortcoming of our study. However, *B. afzelii* and *B. garinii*, both known to be pathogenic to humans, are the two most common *Borrelia* species in Belgium, the Netherlands, and northern France (Rauter and Hartung, 2005), suggesting that most larvae feed on small rodents and birds in this region. A study carried out in the Netherlands (Gassner *et al.* 2008) showed a significantly higher nymphal abundance and *Borrelia* infection rate in oak plots than in pine plots, which was ascribed to differences in rodent densities. In our study, however, the nymphal infection rate with *Borrelia* varied substantially for the different forest sites and years, but no significant effect was found for forest type. Yet, as the absolute number of ticks was considerably higher in oak stands and stands with an abundant shrub layer, the chance of getting bitten by ticks and acquiring infection is in fact influenced by forest type.

The results of this study have important implications for forest management, as management activities can alter the composition and structure of forests, which could have a profound impact on the epidemiology of tick-borne diseases such as Lyme borreliosis. In response to environmental concerns and changing societal needs, one of the main goals

Table 3. Parameter estimates (P.E.) of the best model (see Table 1) for the abundance of *Ixodes ricinus* larvae, nymphs, and adults and for the presence of deer beds

(A positive effect for tree species means a higher tick abundance or deer presence in oak stands compared to pine stands. A positive effect for the year 2009 or 2010 means a higher tick abundance or deer presence in that year compared to 2008.)

Source of variation	Larvae		Nymphs		Adults		Deer beds	
	P.E.	t-value	P.E.	t-value	P.E.	t-value	P.E.	z-value
Intercept	2.931	8.871	2.330	12.170	1.341	7.270	-0.873	-1.469
Tree species	1.428	5.505	0.604	4.870	0.394	3.589	1.189	2.760
Shrub cover	0.017	4.287	0.008	4.381	0.008	4.725	0.016	2.429
Year 2009			0.588	2.906			-0.594	-0.992
Year 2010			0.873	4.135			-1.450	-2.290




Fig. 2. The effects of tree species and shrub layer cover on the number of *Ixodes ricinus* larvae, nymphs, and adults (a–c) and on the number of deer beds (d) in 3 successive years. Shrub cover estimates were grouped into 2 classes: low (<15%) and high (>50%) cover. The results from the 2 forest sites were pooled. Error bars denote the standard error of the mean. Note the difference in values on the y-axis.

of the forest management policy in many parts of Europe is the conversion of (often coniferous) plantations to semi-natural forest types. To achieve this, large areas of homogeneous coniferous stands are being converted into mixed, structure-rich deciduous stands with oak as one of the main constituents. Our results indicate that this forest type can support higher tick population levels than monospecific plantations.

However, whereas tick abundance was highly affected by tree species and shrub cover, the overall *Borrelia* infection rates in ticks were similar in the 2 contrasting forest types. On the other hand, it is important to note that monospecific pine stands cover most of the area in both forest sites, while oak stands, especially those with an abundant shrub layer, are relatively scarce. Large-scale forest conversion programmes could change the composition and

abundance of wildlife communities to the extent that the relative proportion of reservoir-competent and incompetent hosts changes, thereby influencing not only tick abundance but the infection prevalence in ticks as well. In the past decade, increasing attention has been paid to the role of biodiversity in mediating infection levels and disease, termed the dilution effect (Ostfeld and LoGiudice, 2003). The current study underlines the importance of considering spatial heterogeneity in forest habitat quality when studying tick populations and supports vegetation management as a tool to control tick populations. Relatively simple interventions such as mowing the vegetation and clearing brush along forest trails have been shown to be effective in reducing the local abundance of ticks (Wilson, 1986; Schulze *et al.* 1995). However, further studies will be required in order to fully understand the effects of forest conversion on Lyme borreliosis risk.

#### ACKNOWLEDGEMENTS

The authors owe special thanks to Miguel Lyssens-Danneboom, Kris Ceunen, and Filip Ceunen for assistance with fieldwork and to Margot Vanhellemont and Robert Gruwez for critical reading of the manuscript. We are also grateful to Natuurpunt vzw and the Flemish Nature and Forest Agency (ANB) for the permission to work in the forests.

#### FINANCIAL SUPPORT

This work was supported by IWT-Flanders, the Institute for the Promotion of Innovation through Science and Technology in Flanders.

#### REFERENCES

- Allan, B. F., Keesing, F. and Ostfeld, R. S. (2003). Effect of forest fragmentation on Lyme disease risk. *Conservation Biology* **17**, 267–272. doi: 10.1046/j.1523-1739.2003.01260.x.
- Bacon, R. M., Kugeler, K. J. and Mead, P. S. (2008). Surveillance for Lyme disease—United States, 1992–2006. *Morbidity and Mortality Weekly Report* **57**, 1–9.
- Bates, D., Maechler, M. and Bolker, B. (2011). *lme4: Linear Mixed-Effects Models Using Eigen and S4*. R package version 0.999375-42, URL <http://CRAN.R-project.org/package=lme4/> (accessed November 18, 2011).
- Bíró, Z., Szemethy, L., Katona, K., Heltai, M. and Pető, Z. (2006). Seasonal distribution of red deer (*Cervus elaphus*) in a forest-agriculture habitat in Hungary. *Mammalia* **70**, 70–75. doi: 10.1515/MAMM.2006.016.
- Bolker, B. M. (2008). *Ecological Models and Data*. Princeton University Press, Princeton, NJ, USA.
- Boom, R., Sol, C. J. A., Salimans, M. M. M., Jansen, C. L., Wertheim-van Dillen, P. M. E. and van der Noordaa, J. (1990). Rapid and simple method for purification of nucleic acids. *Journal of Clinical Microbiology* **28**, 495–503.
- Boyard, C., Vourc'h, G. and Barnouin, J. (2008). The relationship between *Ixodes ricinus* and small mammal species at the woodland-pasture interface. *Experimental and Applied Acarology* **44**, 61–76. doi: 10.1007/s10493-008-9132-3.
- Brownstein, J. S., Skelly, D. K., Holford, T. R. and Fish, D. (2005). Forest fragmentation predicts local scale heterogeneity of Lyme disease risk. *Oecologia* **146**, 469–475. doi: 10.1007/s00442-005-0251-9.
- Burnham, K. P. and Anderson, D. R. (2002). *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach*. Springer, New York, USA.
- Cederlund, G., Bergqvist, J., Kjellander, P., Gill, R., Gaillard, J. M., Boisauvert, B., Ballon, P. and Duncan, P. (1998). Managing roe deer and their impact on the environment: maximising the net benefits to society. In *The European Roe deer: The Biology of Success* (ed. Andersen, R., Duncan, P. and Linnell, J. D. C.), pp. 337–372. Scandinavian University Press, Oslo, Norway.
- Comstedt, P., Bergström, S., Olsen, B., Garpmo, U., Marjavaara, L., Mejlon, H., Barbour, A. G. and Bunikis, J. (2006). Migratory passerine birds as reservoirs of Lyme borreliosis in Europe. *Emerging Infectious Diseases* **12**, 1087–1095.
- Demaerschalck, I., Ben Messaoud, A., De Kesel, M., Hoyois, B., Lobet, Y., Hoet, P., Bigaignon, G., Bollen, A. and Godfroid, E. (1995). Simultaneous presence of different *Borrelia burgdorferi* genospecies in biological fluids of Lyme disease patients. *Journal of Clinical Microbiology* **33**, 602–608.
- Estrada-Peña, A. (2001). Distribution, abundance, and habitat preferences of *Ixodes ricinus* (Acari: Ixodidae) in northern Spain. *Journal of Medical Entomology* **38**, 361–370. doi: 10.1603/0022-2585-38.3.361.
- Estrada-Peña, A., Osácar, J. J., Pichon, B. and Gray, J. S. (2005). Hosts and pathogen detection for immature stages of *Ixodes ricinus* (Acari: Ixodidae) in North-Central Spain. *Experimental and Applied Acarology* **37**, 257–268. doi: 10.1007/s10493-005-3271-6.
- Gassner, F., van Vliet, A. J. H., Burgers, S. L. G. E., Jacobs, F., Verbaarschot, P., Hovius, E. K. E., Mulders, S., Verhulst, N. O., van Overbeek, L. S. and Takken, W. (2011). Geographic and temporal variations in population dynamics of *Ixodes ricinus* and associated *Borrelia* infections in The Netherlands. *Vector-Borne and Zoonotic Diseases* **11**, 523–532. doi: 10.1089/vbz.2010.0026.
- Gassner, F., Verbaarschot, P., Smallegange, R. C., Spitzen, J., Van Wieren, S. E. and Takken, W. (2008). Variations in *Ixodes ricinus* density and *Borrelia* infections associated with cattle introduced into a woodland in The Netherlands. *Applied and Environmental Microbiology* **74**, 7138–7144. doi: 10.1128/AEM.00310-08.
- Gill, R. M. A. (1990). *Monitoring the Status of European and North American Cervids*. Global Environment Monitoring System, United Nations Environment Programme, Nairobi, Kenya.
- Gray, J. S. (1998). The ecology of ticks transmitting Lyme borreliosis. *Experimental and Applied Acarology* **22**, 249–258. doi: 10.1023/A:1006070416135.
- Gray, J. S., Kahl, O., Janetzki, C. and Stein, J. (1992). Studies on the ecology of Lyme disease in a deer forest in County Galway, Ireland. *Journal of Medical Entomology* **29**, 915–920.
- Halos, L., Bord, S., Cotté, V., Gasqui, P., Abrial, D., Barnouin, J., Boulouis, H. J., Vayssier-Taussat, M. and Vourc'h, G. (2010). Ecological factors characterizing the prevalence of bacterial tick-borne pathogens in *Ixodes ricinus* ticks in pastures and woodlands. *Applied and Environmental Microbiology* **76**, 4413–4420. doi: 10.1128/AEM.00610-10.
- Hanincová, K., Schäfer, S. M., Etti, S., Sewell, H. S., Taragelová, V., Ziak, D., Labuda, M. and Kurtenbach, K. (2003). Association of *Borrelia afzelii* with rodents in Europe. *Parasitology* **126**, 11–20. doi: 10.1017/S0031182002002548.
- Hillyard, P. D. (1996). *Ticks of North-West Europe*. The Natural History Museum, London, UK.
- Humair, P. F. and Gern, L. (1998). Relationship between *Borrelia burgdorferi* sensu lato species, red squirrels (*Sciurus vulgaris*) and *Ixodes ricinus* in enzootic areas in Switzerland. *Acta Tropica* **69**, 213–227. doi: 10.1016/S0001-706X(97)00126-5.
- Humair, P. F., Rais, O. and Gern, L. (1999). Transmission of *Borrelia afzelii* from *Apodemus* mice and *Clethrionomys* voles to *Ixodes ricinus* ticks: differential transmission pattern and overwintering maintenance. *Parasitology* **118**, 33–42. doi: 10.1017/S0031182098003564.
- Hurvich, C. M. and Tsai, C. L. (1989). Regression and time series model selection in small samples. *Biometrika* **76**, 297–307. doi: 10.1093/biomet/76.2.297.
- Jaenson, T. G. T., Jaenson, D. G. E., Eisen, L., Petersson, E. and Lindgren, E. (2012). Changes in the geographical distribution and abundance of the tick *Ixodes ricinus* during the past 30 years in Sweden. *Parasites & Vectors* **5**, 8. doi: 10.1186/1756-3305-5-8.
- Jongejan, F. (2001). *Teken en door Teken Overgedragen Ziekten*. Stichting Diergeneeskundig Memorandum, Boxtel, The Netherlands.
- Kurtenbach, K., Peacey, M., Rijpkema, S. G. T., Hoodless, A. N., Nuttall, P. A. and Randolph, S. E. (1998). Differential transmission of the genospecies of *Borrelia burgdorferi* sensu lato by game birds and small rodents in England. *Applied and Environmental Microbiology* **64**, 1169–1174.
- Linard, C., Lamarque, P., Heyman, P., Ducoffre, G., Luyasu, V., Tersago, K., Vanwambeke, S. O. and Lambin, E. F. (2007).



- Determinants of the geographic distribution of Puumala virus and Lyme borreliosis infections in Belgium. *International Journal of Health Geographics* **6**, 15. doi: 10.1186/1476-072X-6-15.
- Lindström, A. and Jaenson, T. G. T.** (2003). Distribution of the common tick, *Ixodes ricinus* (Acari: Ixodidae), in different vegetation types in southern Sweden. *Journal of Medical Entomology* **40**, 375–378. doi: 10.1603/0022-2585-40.4.375.
- Olsthoorn, A. F. M., Bartelink, H. H., Gardiner, J. J., Pretzsch, H., Hekhuis, H. J. and Franc, A.** (1999). *Management of Mixed-Species Forest: Silviculture and Economics*. DLO Institute for Forestry and Nature Research (IBN-DLO), Wageningen, The Netherlands.
- Ostfeld, R. S. and LoGiudice, K.** (2003). Community disassembly, biodiversity loss, and the erosion of an ecosystem service. *Ecology* **84**, 1421–1427. doi: 10.1890/02-3125.
- Pichon, B., Mousson, L., Figureau, C., Rodhain, F. and Perez-Eid, C.** (1999). Density of deer in relation to the prevalence of *Borrelia burgdorferi* s.l. in *Ixodes ricinus* nymphs in Rambouillet forest, France. *Experimental and Applied Acarology* **23**, 267–275. doi: 10.1023/A:1006023115617.
- Piesman, J. and Gern, L.** (2004). Lyme borreliosis in Europe and North America. *Parasitology* **129** (Suppl.), S191–S220. doi: 10.1017/S0031182003004694.
- Rauter, C. and Hartung, T.** (2005). Prevalence of *Borrelia burgdorferi* sensu lato genospecies in *Ixodes ricinus* ticks in Europe: a metaanalysis. *Applied and Environmental Microbiology* **71**, 7203–7216. doi: 10.1099-2240/05/7203-7216.
- R Development Core Team** (2011). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org/> (accessed November 18, 2011).
- Ruiz-Fons, F. and Gilbert, L.** (2010). The role of deer as vehicles to move ticks, *Ixodes ricinus*, between contrasting habitats. *International Journal for Parasitology* **40**, 1013–1020. doi: 10.1016/j.ijpara.2010.02.006.
- Said, S. and Servanty, S.** (2005). The influence of landscape structure on female roe deer home-range size. *Landscape Ecology* **20**, 1003–1012. doi: 10.1007/s10980-005-7518-8.
- Schulze, T. L., Jordan, R. A. and Hung, R. W.** (1995). Suppression of subadult *Ixodes scapularis* (Acari: Ixodidae) following removal of leaf litter. *Journal of Medical Entomology* **32**, 730–733.
- Smith, H. D., Oveson, M. C. and Pritchett, C. L.** (1986). Characteristics of mule deer beds. *Great Basin Naturalist* **46**, 542–546.
- Sood, S. K., O'Connell, S. and Weber, K.** (2011). The emergence and epidemiology of Lyme borreliosis in Europe and North America. In *Lyme Borreliosis in Europe and North America: Epidemiology and Clinical Practice* (ed. Sood, S. K.), pp. 1–35. John Wiley & Sons, Inc., New Jersey, USA. doi: 10.1002/9780470933961.ch1.
- Spiecker, H., Hansen, J., Klimo, E., Skovsgaard, J. P., Sterba, H. and von Teuffel, K.** (2004). *Norway Spruce Conversion—Options and Consequences*. Koninklijke Brill, Leiden, The Netherlands.
- Spielman, A.** (1994). The emergence of Lyme disease and human babesiosis in a changing environment. *Annals of the New York Academy of Sciences* **740**, 146–156. doi: 10.1111/j.1749-6632.1994.tb19865.x.
- Tack, W., Madder, M., Baeten, L., Vanhellemont, M., Gruwez, R. and Verheyen, K.** (2012). Local habitat and landscape affect *Ixodes ricinus* tick abundances in forests on poor, sandy soils. *Forest Ecology and Management* **265**, 30–36. doi: 10.1016/j.foreco.2011.10.028.
- Tack, W., Madder, M., De Frenne, P., Vanhellemont, M., Gruwez, R. and Verheyen, K.** (2011). The effects of sampling method and vegetation type on the estimated abundance of *Ixodes ricinus* ticks in forests. *Experimental and Applied Acarology* **54**, 285–292. doi: 10.1007/s10493-011-9444-6.
- Tälleklint, L. and Jaenson, T. G. T.** (1997). Infestation of mammals by *Ixodes ricinus* ticks (Acari: Ixodidae) in south-central Sweden. *Experimental and Applied Acarology* **21**, 755–771. doi: 10.1023/A:1018473122070.
- Tufto, J., Andersen, R. and Linnell, J.** (1996). Habitat use and ecological correlates of home range size in a small cervid: the roe deer. *Journal of Animal Ecology* **65**, 715–724. doi: 10.2307/5670.
- van Dam, A. P., Kuiper, H., Vos, K., Widjojokusumo, A., de Jongh, B. M., Spanjaard, L., Ramselaar, A. C. P., Kramer, M. D. and Dankert, J.** (1993). Different genospecies of *Borrelia burgdorferi* are associated with distinct clinical manifestations of Lyme borreliosis. *Clinical Infectious Diseases* **17**, 708–717. doi: 10.1093/clinids/17.4.708.
- Verkem, S., De Maeseneer, J., Vandendriessche, B., Verbeylen, G. and Yskout, S.** (2003). *Zoogdieren in Vlaanderen. Ecologie en Verspreiding van 1987 tot 2002*. Natuurpunt Studie & JNM-Zoogdierenwerkgroep, Mechelen & Gent, Belgium.
- Waterinckx, M. and Roelandt, B.** (2001). *De Bosinventaris van het Vlaamse Gewest*. Ministerie van de Vlaamse Gemeenschap, Afdeling Bos & Groen, Brussel, Belgium.
- Wielinga, P. R., Gaasenbeek, C., Fonville, M., de Boer, A., de Vries, A., Dimmers, W., Akkerhuis Op Jagers, G., Schouls, L. M., Borgsteede, F. and van der Giessen, J. W. B.** (2006). Longitudinal analysis of tick densities and *Borrelia*, *Anaplasma*, and *Ehrlichia* infections of *Ixodes ricinus* ticks in different habitat areas in The Netherlands. *Applied and Environmental Microbiology* **72**, 7594–7601. doi: 10.1128/AEM.01851-06.
- Wilson, M. L.** (1986). Reduced abundance of adult *Ixodes dammini* (Acari: Ixodidae) following destruction of vegetation. *Journal of Economic Entomology* **79**, 693–696.
- World Health Organization** (2004). *The Vector-borne Human Infections of Europe: Their Distribution and Burden on Public Health*. World Health Organization, Copenhagen Denmark.