

REFERENCES

- Abbas, G., M.J. Arif, and S. Saeed. 2005. Systematic status of a new species of the genus *Phenacoccus* Cockerell (Pseudococcidae), a serious pest of cotton, *Gossypium hirsutum* L. Pakistan Entomologist, 27: 83-84.
- Abo, M.E. and Sy, A.A. 1998. Rice virus diseases: epidemiology and management strategies. Journal of Sustainable Agriculture 11: 113–134.
- Acland, J.D. 1971. East African crops: an introduction to the production of field and plantation crops in Kenya, Tanzania and Uganda. FAO and Longman Group UK; Rome and London.
- Addison, P. 2002. Chemical stem barriers for the control of ants (Hymenoptera: Formicidae) in vineyards. South African Journal of Enology and Viticulture 23, 1–8.
- Adenuga, A.O., 1975. Mutualistic association between ants and some Homoptera, its significance in cacao production. Psyche 82, 24–28.
- African Union 2009. Biological control program for the mango mealybug *Rastrococcus invadens* Williams (Homoptera: Pseudococcidae) in West Africa. Semi-Arid Africa Agricultural Research and Development. Final report. Pp 5-32.
- Agounké and Fischer (1993). Biological control of the mango mealybug (*Rastrococcus invadens*) in Togo. Acta Horticulturae 341: 441-451.
- Agounké, D., Agricola, U, and Bokonon-Ganta, A. 1988. *Rastrococcus invadens* Williams (Hemiptera, Pseudococcidae), a serious exotic pest of fruit trees and other plants in West Africa. Bulletin of Entomological Research 78: 695–702.
- Agounké, D., Agricola, U. and Bokonon-Ganta, A. 1989. *Rastrococcus invadens* Williams (Homoptera: Pseudococcidae), an exotic pest of fruit trees in West Africa. Bulletin du Service de la Protection des Végétaux 11: 17–29.
- Agricola , U. and Fischer, H.U. 1991 Hyperparasitism in two newly introduced parasitoids; *Epidinocarsis lopezi* and *Gyranusoidea tebygi* (Hymenoptera; Encyrtidae) after their establishment in Togo. Bulletin of Entomological Research 81: 127- 132.
- Agricola, U., Agounké, D., Fischer, H.U., Moore, D., 1989. The control of *Rastrococcus invadens* Williams (Homoptera: Pseudococcidae) in Togo by the introduction of *Gyranusoidea tebygi* Noyes (Hymenoptera: Encyrtidae). Bulletin of Entomological Research 79, 671–678.
- Akintola, A.J., and A.T. Ande. 2008. First record of *Phenacoccus solenopsis* Tinsley (Hemiptera: Pseudococcidae) on *Hibiscus rosa-sinensis* in Nigeria. Agricultural Journal, 3:1-3.
- Ali M, 1970. A catalogue of the Oriental Coccoidea. Part IV. Bulletin of Indian Museum 5:71-150.

- Al-Saffar, Z.Y. and J.C. Aldrich. 1998. *Pontania proxima* (Tenthredinidae: Hymenoptera): natural enemies and defensive behaviour against *Pnigalio nemati* (Eulophidae: Hymenoptera). Annals of the Entomological Society of America 91: 858-862.
- Amarasekare, K.G., Chong, J-H., Epsky, N.D., Mannion, C.M., 2008. Effect of temperature on the life history of the mealybug *Paracoccus marginatus* (Hemiptera: Pseudococcidae), Journal of Economic Entomology 101(6), 1798-804.
- Anand, P., Ayub, K. 2000. The effect of five insecticides on *Maconellicoccus hirsutus* (Green) (Homoptera: Pseudococcidae) and its natural enemies *Anagyrus kamali* Moursi (Hymenoptera: Encyrtidae), and *Cryptolaemus montrouzieri* Mulsant and *Scymnus coccivora* Aiyar (Coleoptera: Coccinellidae). International Pest Control 42: 170–173.
- Anderson, R.P., Gomez-Laverde, M. and Peterson, A.T. (2002) Geographical distributions of spiny pocket mice in South America: insights from predictive models. Global Ecology and Biogeography 11, 131–141.
- Anderson, R.P., Lew, D. and Peterson, A.T. (2003) Evaluating predictive models of species' distributions: criteria for selecting optimal models. Ecological Modeling 162, 211–232.
- Arai, T. 2002. Attractiveness of sex pheromone of *Pseudococcus cryptus* Hempel (Homoptera: Pseudococcidae) to adult males in a citrus orchard. Applied Entomology and Zoology 37: 69–72.
- Arif, M.J., Gogi, M.D., Mirza, M., Zia, K., Hafeez, F., 2006. Impacts of plant spacing and abiotic factors on population dynamics of sucking insect pests of cotton. Pakistan Journal of Biological Sciences 9, 1364-1369.
- Austin, M.P., 2002. Spatial prediction of species distribution: an interface between ecological theory and statistical modelling. Ecological Model. 157, 101–118.
- Babu, T.R. and Ramanamurthy, G. 1998. Residual toxicity of pesticides to the adults of *Scymnus coccivora* Aiyar (Coccinellidae: Coleoptera) a predator on mealybugs on grape. Indian Journal of Plant Protection 26: 96–98.
- Bach, C.E. 1991. Direct and indirect interactions between ants (*Pheidole megacephala*), scales (*Coccus viridis*) and plants (*Pluchea indica*). Oecologia, 87: 233-239.
- Baggen, L.R. and Gurr, G.M. 1998. The influence of food on *Copidosoma koehleri* (Hymenoptera: Encyrtidae), and the use of flowering plants as a habitat management tool to enhance biological control of potato moth, *Phthorimaea operculella* (Lepidoptera: Gelechiidae). Biological Control 11, 9–17.
- Barbosa, P., Saunders, J.A. and Waldvogel, M. 1982. Plant-mediated variation in herbivore suitability and parasitoid fitness. Proc . 5th int. Sump. Insect-Plant Relationships, Wageningen, the Netherlands.
- Barker, J.S.F., Toll, G.L., East, P.D., Miranda, M. and Phaff, H.J. 1982. Heterogeneity of the yeast flora in the breeding sites of catophilic *Drosophila*. Canadian Journal of Microbiology 29: 6-14.

- Barratt, B.I.P. and Johnstone, P.D. 2001. Factors affecting parasitism by *Microctonus aethiopoides* (Hymenoptera: Braconidae), and parasitoid development in natural and novel host species. *Bulletin of Entomological Research* 91, 245–253.
- Bartlett, B.R., 1961. The influence of ants upon parasites, predators, and scale insects. *Annals of the Entomological Society of America* 54, 543–551.
- Beentje, H. 1994. Kenya Trees, Shrubs and Lianas. National Museums of Kenya, Nairobi (Kenya), pp. 564–570.
- Bell, V.A., Suckling, D.M., Walker, J.T.S., Millar, J.G., Manning, L.A. and El-Sayed, A.M. 2008. Obscure mealybug pheromone is the kairomone of an introduced parasitoid in New Zealand. *Proc XXIII Int Cong Entomol*, 6–12 July 2008, Durban (Abstract). Available at <http://www.ice2008.org.za/pdf/proceedings.pdf>.
- Bell, V.A., Walker, J.T.S., Suckling, D.M., Manning, L.A., El-Sayed, A.M., Shaw, P.W., Wallis, D.R. and Millar, J.G. 2006. Trapping obscure mealybug (*Pseudococcus viburni*) and its natural enemy *Pseudaphycus maculipennis* (Hymenoptera: Encyrtidae) in apple orchards. *New Zealand Plant Protection* 59: 364.
- Ben-Dov, Y. 1994. A systematic catalogue of the mealybugs of the world (Insecta: Homoptera: Coccoidea: Pseudococcidae and Putoidae) with data on geographical distribution, host plants, biology and economic importance. Intercept Limited, Andover, UK. Pp 435–436.
- Ben-Dov, Y. 2005a. The malvastrum mealybug *Ferrisia malvastra* (Hemiptera: Coccoidea: Pseudococcidae): Distribution, host plants and pest status in Israel. *Phytoparasitica* 33: 154–156.
- Ben-Dov, Y. 2006. Scales in a family/genus query. Available at <http://www.sel.barc.usda.gov/scalecgi/chklist.exe?Family=Pseudococcidae&genus=Accessed> 14 August 2008.
- Berlinger, M.J. and Golberg, A.M. 1978. Effect of the fruit sepals on the citrus mealybug population and on its parasite. *Entomologia Experimentalis et Applicata* 24: 238–243.
- Bess, H.A. 1958. The green scale, *Coccus viridis* (Green) (Homoptera: Coccidae), and ants. *Proceedings of the Hawaiian Entomological Society*, 16: 349–355.
- Bethke, J.A., Redak, R.A. and Schuch U.K. 1998. Melon aphid performance on chrysanthemum as mediated by cultivar, and deferential levels of fertilization and irrigation. *Entomologia Experimentalis et Applicata* 88: 41–47.
- Bhat, A.I., Devasahayam, S., Sarma, Y.R. and Pant, R.P. 2003. Association of a badnavirus in black pepper(*Piper nigrum* L) transmitted by mealybug (*Ferrisia virgata*) in India. *Current Science* 84:1547–1550
- Biassangama, A., Fabres, G. and Moussa, J.B. 1991. The presence in Congo of *Gyranusoidea tebygi* (Hymenoptera: Encyrtidae), parasitoid of *Rastrococcus invadens* (Homoptera: Pseudococcidae). *Bulletin de la Société Entomologique de France* 96: 209–211.
- Blumberg, D. 1997. Parasitoid encapsulation as a defense mechanism in the Coccoidea (Homoptera) and its importance in biological control. *Biological Control* 8: 225–236.

- Blumberg, D., and Van Driesche, R.G. 2001. Encapsulation rates of three encyrtid parasitoids by three mealybug species (Homoptera: Pseudococcidae) found commonly as pests in commercial greenhouses. *Biological Control* 22, 191-199.
- Blumberg, D., Ben-Dov, Y. and Mendel, Z. 1999. The citriculus mealybug, *Pseudococcus cryptus* Hempel, and its natural enemies in Israel: history and present situation. *Entomologica*, Bari 33: 233–242.
- Blumberg, D., Franco, J.C., Suma, P., Russo, A. and Mendel, Z. 2001. Parasitoid encapsulation in mealybugs (Hemiptera: Pseudococcidae) as affected by the host-parasitoid association and super-parasitism. *Boll Zool Agr Bachicolt* 33: 385–395.
- Blumberg, D., Klein, M. and Mendel, Z. 1995. Response by encapsulation of 4 mealybug species (HomopteraPseudococcidae) to parasitization by *Anagyrus pseudococci*. *Phytoparasitica* 23: 157–163.
- Boavida, C. and Neuenschwander, P. 1995. Influence of host plant on the mango mealybug, *Rastrococcus invadens*. *Entomologia Experimentalis et Applicata* 76: 179-188.
- Boavida, C., and Neuenschwander, P. 1995. Population dynamics and life tables of the mango mealybug, *Rastrococcus invadens* Williams, and its introduced natural enemy *Gyranusoidea tebygi* Noyes in Benin. *Biocontrol Science and Technology* 5, 489–508.
- Boavida, C., M. Ahounou, M. Vos, P. Neuenschwander and van Alphen, J.J.M. 1995. Host stage selection and sex allocation by *Gyranusoidea tebygi* (Hymenoptera: Encyrtidae), a parasitoid of the mango mealybug, *Rastrococcus invadens* (Homoptera: Pseudococcidae). *Biological Control* 5: 487-496.
- Bokonon-Ganta, A.H. and Neuenschwander, P. 1995. Impact of the biological control agent *Gyranusoidea tebygi* Noyes (Hymenoptera: Encyrtidae) on the mango mealybug, *Rastrococcus invadens* Williams (Homoptera: Pseudococcidae), in Benin. *Biocontrol Science and Technology* 5, 95–107.
- Bokonon-Ganta, A.H., De Groote H. and Neuenschwander. P. 2002. Socio-economic impact of biological control of mango mealybug in Benin. *Agriculture, Ecosystem and Environment* 93: 367-378.
- Bokonon-Ganta, A.H., Neuenschwander, P., Van alphen, J.J.M, and Vost, M. 1995. Host stage selection and sex allocation by *Anagyrus mangicola* Noyes (Hymenoptera: Encyrtidae) a parasitoid of the mango mealybug, *Rastrococcus invadens* Williams (Hemiptera: Pseudococcidae). *Biological Control*. 5: 479-486.
- Bole, P.V., and Vaghani, Y. 1986. Field guide to common trees of India. Oxford University Press, Bombay, India. 125 pp.
- Bongiorni, S., Mazzuoli, M., Masci, S. and Prantera, G. 2001. Facultative heterochromatization in para-haploid male mealybugs: involvement of a heterochromatin-associated protein. *Development* 128: 3809–3817.

- Boussienguet, J. and Herren, H.R. 1992. Introduction et dynamique de dispersion de la cochenille du mangue, *Rastrococcus invadens* Williams (Hom., Pseudococcidae) au Gabon. Mémoire de la Société Royale Belge d'Entomologie 35 : 363–367.
- Boussienguet, J. and Moulougou, J. 1993. Demographic pressure and host plant choice of *Rastrococcus invadens*, a pest of mango recently introduced into Africa. Bulletin de la Société entomologique de France 98: 139-148.
- Bradley, G.A. 1973. Effect of *Formica obscuripes* (Hymenoptera: Formicidae) on the predatory-prey relationship between *Hyperaspis congreensis* (Coleoptera: Coccinellidae) and *Toumeyella numismaticum* (Homoptera: Coccidae). The Canadian Entomologist, 105: 1113-1118.
- Brandis, S. D. 1907. Indian Trees. London: Constable. 246 pp.
- Briese, D.T. 1982. Damage to saltbush by the coccid *Pulvinaria maskelli* Olliff, and the role played by an attendant ant. Journal of the Australian Entomological Society, 21: 293-294.
- Brito, J.C. and Crespo, E.G. 2002. Distributional analysis of two vipers (*Vipera latastei* and *V. seoanei*) in a potential area of sympatry in the Northwestern Iberian Peninsula. In: Biology of the Vipers, p. 129-138. Schuett, G.W., Hoggren, M., Douglas, M.E., Greene, H.W., Eds, Eagle Mountain Publishing, LC.
- Brown, J.H., Stevens, G.C. and Kaufman, D.M. 1996. The geographic range: size, shape, boundaries, and internal structure. Annual Review of Ecology and Systematics 27, 597–623.
- Browning, H.W. 1994. Early classical biological control on citrus. In: Rosen D, Bennett FD, Capinera JL (eds) Pest management in the subtropics: biological control – a Florida perspective. Intercept, Andover.
- Buckley, R. and Gullan, P. 1991. More aggressive ant species (Hymenoptera: Formicidae) provide better protection for soft scale and mealybugs (Homoptera: Coccidae; Pseudococcidae). Biotropica, 23, 282–286.
- Buckley, R.C. 1987a. Ant-plant-homopteran interactions. Advances in Ecological Research, 16: 53-85.
- Buckley, R.C. 1987b. Interactions involving plants, Homoptera and ants. Annual Review of Ecology and Systematics, 18, 111–135.
- Budenberg, W.J. 1990. Honeydew as a contact kairomone for aphid parasitoids. Entomologia Experimentalis et Applicata 55: 139– 148.
- Burns, D.P. 1973. The foraging and tending behaviour of *Dolichoderus taschenbergi* (Hymenoptera: Formicidae). The Canadian Entomologist, 105: 97-104.
- Butcher, C.F. 1983. Cottony cushion scale, Seychelles scale, and Egyptian fluted scale. South Pacific Commission advisory leaflet no. 16. 4p.
- CABI. 1995. International Institute of Entomology *Rastrococcus iceryoides*. Distribution maps of pests, Series A No. 561, Wallingford, UK: 2 pp.

- CABI. 2000. Crop protection compendium. Global module, 2nd edition. CABI Publishing, Wallingford, UK.
- Calatayud P.A., Duchon S. and Amaze, T.L. 2000. Estimation of carbon and nitrogen modification during water deficiency in leaves of cassava, *Manihot esculenta* Crantz. In: Proceedings of the IVth International Scientific Meeting of Cassava Biotechnology Network, Carvalho L. J. C. B., Thro A. M. & A. D. Vilarinhos (eds), November 03-07, 1998, Salvador, Bahia, Brazil, p. 288-298.
- Calatayud, P.A., Polania, M.A., Seligmann, C.D. and Bellotti, A.C. 2002. Influence of water-stressed cassava on *Phenacoccus herreni* and three associated parasitoids. *Entomologia Experimentalis et Applicata* 102: 163–175.
- Calatayud, P.A., Rahbe, Y., Delobel, B., Khuonghuu, F., Tertuliano, M. and Leru, B. 1994. Influence of secondary compounds in the phloem sap of cassava on expression of antibiosis towards the mealybug *Phenacoccus manihoti*. *Entomologia Experimentalis et Applicata* 72: 47–57.
- Caltagirone, L.E. and Doutt, R.L. 1989. The history of vedalia beetle importation to California and its impact on the development of biological control. *Annual review of entomology* 34: 1-16.
- Cammell, M.E. and Knight, J.D. 1992. Effects of climate change on the population dynamics of crop pests. *Advances in Ecological Research* 22:117–162.
- Carey, J.R. 1993. Applied demography for biologists with special emphasis on insects. Oxford University Press, New York.
- Carlton, J.T. 1996. Pattern, process, and prediction in marine invasion ecology. *Biological Conservation* 78: 97 - 106.
- Carroll, C.R. and Janzen, D.H., 1973. Ecology of foraging by ants. *Annual Review of Ecology and Systematics* 4, 231-57.
- Charles, J.G. 1993. A survey of mealybugs and their natural enemies in horticultural crops in North Island, New Zealand, with implications for biological control. *Biocontrol Science and Technology* 3: 405–418.
- Charles, S., Teppen, B.J., H. Li, Laird, D.A. and Boyd, S.A. 2006. Ex-changeable cation hydration properties strongly influence sorption of nitroaromatic compounds by soils. *Soil Science Society of America Journal*. In press.
- Charleston, K., and D. Murray. 2010. Exotic mealybug species – a major new pest in cotton. Queensland Government. The Beat sheet. Accessed online 17 September 2011 at <http://thebeatsheet.com.au/mealybugs/exotic-mealybug-species-a-major-new-pest-in-cotton/>
- Charnov, E.L., Skinner, S.W. 1985. Complementary approaches to the understanding of parasitoid oviposition decisions. *Environmental Entomology* 14, 383–391.
- Chaudhari, G.B., Bharpoda, T.M., Patel, J.J., Patel, K.I. and Patel, J.R. 1999. Effect of weather on activity of cotton bollworms in middle Gujarat. *Journal of Agrometeorology* 1, 137-142.

- Chih, H.W., Chiu, H.F., Tang, K.S., Chang, F.R, and Wu, Y.C. 2001. Bullatacin, a potent antitumor Annonaceous acetogenin, inhibits proliferation of human hepatocarcinoma cell line 2×2×15 by apoptosis induction; *Life Sciences* 69: 1321–1331.
- Chong JH and Oetting R.D. 2007b. Specificity of *Anagyrus* sp nov nr. *sinope* and *Leptomastix dactylopii* for six mealybug species. *Biocontrol* 52: 289–308.
- Chong, J.H. and Oetting, R.D. 2006. Influence of temperature and mating status on the development and fecundity of the mealybug parasitoid, *Anagyrus* sp nov nr *sinope* Noyes and Menezes (Hymenoptera: Encyrtidae). *Environmental Entomology* 35: 1188–1197.
- Chong, J.H. and Oetting, R.D. 2007. Specificity of *Anagyrus* sp. nov. nr. *sinope* and *Leptomastix dactylopii* for six mealybug species. *BioControl* 52, 289-308.
- Chong, J.H. and Oetting, R.D. 2007a. Intraguild predation and interference by the mealybug predator *Cryptolalemus montrouzieri* on the parasitoid *Leptomastix dactylopii*. *Biocontrol Science and Technology* 17: 933–944.
- Chong, J.H., Oetting, R.D. and Osborne, L.S. 2005. Development of *Diomus austrinus* Gordon (Coleoptera: Coccinellidae) on two mealybug prey species at five constant temperatures. *Biological Control* 33: 39–48.
- Chong, J.H., van Iersel, M.W. and Oetting, R.D. 2004. Effects of elevated carbon dioxide levels and temperature on the life history of the Madeira mealybug (Hemiptera: Pseudococcidae). *Journal of Entomological Science* 39: 387–397.
- Chong, J-H, Oetting, R.D. and Van Iersel, M.W. 2003. Temperature Effects on the Development, Survival, and Reproduction of the Madeira Mealybug, *Phenacoccus madeirensis* Green (Hemiptera: Pseudococcidae), on Chrysanthemum. *Annals of the Entomological Society of America* 96 (4), 539-543.
- Chong, J-H, Roda, A.L. and Mannion, C.M. 2008. Life History of the Mealybug, *Maconellicoccus hirsutus* (Hemiptera: Pseudococcidae), at Constant Temperatures. *Environmental Entomology*. 37(2): 323-332.
- Christopher K. Ruffo, Ann Birnie and Bo Tengnas. 2002. Edible wild plants of Tanzania. *Regional land management unit (RELMA)*. Technical handbook No. 27.
- Cid, M., Pereira, S., Cabaleiro, C., Faoro, F. and Segura, A. 2007. Presence of Grapevine leafroll-associated virus 3 in primary salivary glands of the mealybug vector *Planococcus citri* suggests a circulative transmission mechanism. *European Journal of Plant Pathology* 118: 23–30.
- Clarke, S.R., De Barr, G.L. and Berisford, C.W. 1989. The life history of *Toumeyella pini* (King) (Homoptera: Coccidae) in loblolly pine seed orchards in Georgia. *The Canadian Entomologist*, 121: 853-860.
- Clausen, C.P. (Ed) 1978. Introduced parasites and predators of arthropod pests and weeds: a world review. Washington, DC Agricultural research Service, United States Department of Agriculture, Agricultural Handbook No. 480.

- Clausen, C.P. 1958. Biological control of insect pests. Annual Review of Entomology 3: 291-310.
- Cloyd, R.A. and Dickinson, A. 2006. Effect of Insecticides on Mealybug Destroyer (Coleoptera: Coccinellidae) and Parasitoid *Leptomastix dactylopii* (Hymenoptera: Encyrtidae), Natural Enemies of Citrus Mealybug (Homoptera: Pseudococcidae). Journal of Economic Entomology 99:1596–1604.
- Cochard, R, and Jackes, B.R. 2005. Seed ecology of the invasive tropical tree *Parkinsonia aculeata*. Plant Ecology 180:13-31.
- Colman-Saizarbitoria, T., Gu, Z.M., Zhao, G.X., Zeng, L., Kozlowski, J.F. and McLaughlin, J.L. 1995. Venezenin: a new bioactive Annonaceous acetogenin from the bark of *Xylopia aromaticata*. Journal of Natural Products 58: 532–539.
- Compere, H. 1940. Parasites of the black scale, *Saissetia oleae*, in Africa. Hilgardia, 13: 387-425.
- Conover, W.J. 1999. Practical Nonparametric Statistics. Wiley Series in Probability and Statistics: Applied Probability and Statistics Section. New York: John Wiley and Sons, Inc.
- Copland, M.J.W., Tingle, C.C.D., Saynor, M. and Panis, A. 1985. Biology of glasshouse mealybugs and their predators and parasitoids. In: Hussey, NW, Scopes N (Eds) Biological pest control: the glasshouse experience. Blandford Press, Poole, UK.
- Cox, J. 1989. The mealybug genus *Planococcus* (Homoptera: Pseudococcidae). Bulletin of the British Museum (Natural History), Zoology series (Entomology) 58: 1–78.
- Cox, J.M. and Pearce, M.J. 1983. Wax produced by dermal pores in three species of mealybug (Homoptera, Pseudococcidae). International Journal of Insect Morphology and Embryology 12: 235–248.
- CPC (2002) Crop Protection Compendium Database, CAB Int'l. Wallingford, UK.
- Cranshaw, W., Jevremovic, Z., Sclar, D.C. and Mannix, L. 2000. Observations on the biology and control of the hawthorn (two-circuli) mealybug, *Phenacoccus dearnessi* (King). The Journal of Arboriculture 26: 225–229.
- Cross, A.E., Moore, D. 1992. Developmental studies on *Anagyrus mangicola* (Hymenoptera: Encyrtidae) a parasitoid of the mealybug *Rastrococcus invadens* (Homoptera: Pseudococcidae). Bulletin of Entomological Research 82, 307 – 312.
- Cudjoe, A.R., Neuenschwander, P. and Copland, M.J.W. 1993. Interference by ants in biological control of the cassava mealybug *Phenacoccus manihoti* (Hemiptera: Pseudococcidae) in Ghana. Bulletin of Entomological Research, 83, 15–22.
- Daane, K.M, Bentley, W.J. and Weber, E.A. 2004a. Vine mealybug: a formidable pest spreads throughout California vineyards. Prac Winery Vineyard Ma 3:35–40.
- Daane, K.M, Malakar-Kuenen, R.D, and V.M. Walton. 2004b. Temperature-dependent development of *Anagyrus pseudococci* (Hymenoptera: Encyrtidae) as a parasitoid of the

- vine mealybug, *Planococcus ficus* (Homoptera: Pseudococcidae). Biological Control. 31:123–132.
- Daane, K.M., Bentley, W.J., Walton, V.M., Malakar-Kuenen, R., Millar, J.G., Ingels, C.A., Weber, E.A. and Gispert, C. 2006a. New controls investigated for vine mealybug. California Agriculture 60: 31–38.
- Daane, K.M., Sime, K.R., Fallon, J. and Cooper, M.L. 2007. Impacts of Argentine ants on mealybugs and their natural enemies in California's coastal vineyards. Ecological Entomology 32, 583–596.
- Daane, K.M., Sime, K.R., Hogg, B.N., Bianchi, M.L., Cooper, M.L., Rust, M.K. and Klotz, J.H. 2006b. Effects of liquid insecticide baits on Argentine ants in California's coastal vineyards. Crop Protection 25: 592–603.
- Das, G.M. 1959. Observations on the association of ants with coccids of tea. Bulletin of Entomological Research, 50: 437.
- Davies, A.P., Ceballo, F.A. and Walter, G.H. 2004. Is the potential of *Coccidoxyenoides perminutus*, a mealybug parasitoid, limited by climatic or nutritional factors? Biological Control 31: 181–188.
- Davis, A.J., Lawton, J.H., Shorrocks, B. and Jenkinson, L.S. 1998. Individualistic species responses invalidate simple physiological models of community dynamics under global environmental change. Journal of Animal Ecology 67, 600–612.
- De Meyer, M., Robertson, M.P., Peterson, A.T. and Mansell, M.W. 2008. Ecological niches and potential geographical distributions of Mediterranean fruit fly (*Ceratitis capitata*) and Natal fruit fly (*Ceratitis rosa*). Journal of Biogeography 35, 270–281.
- de Souza, J.C., Reis, P.R., Ribeiro, J.A., Santa Cecília, L.V.C. and Silva, R.A. 2007. Chemical control of the coffee root mealybug *Dysmicoccus texensis* (Tinsley, 1900) in coffee plants (*Coffea arabica* L.). Coffee Sciences 2:29–37.
- DeBach, P. 1949. Population Studies of the Long-Tailed Mealybug and Its Natural Enemies on Citrus Trees in Southern California, Ecology 30(1), 14-25
- DeBach, P., Fleschner, C.A. and Dietrick, E.J. 1951. A biological check method for evaluating the effectiveness of entomophagous insects. Journal of Economic Entomology, 44, 763–766.
- Degen, A.A. and Gersani, M. 1989. Environmental effects on activity and honeydew collection by the weaver ant *Polyrhachis simplex* (Hymenoptera, Formicidae) when attending the mealybug *Trabutina* sp (Homoptera, Pseudococcidae). Journal of Zoology, 218: 421–432.
- Delalibera, I., Humber, R.A., Bento, J.M.S. and DeMatos, A.P. 1997. First record of the entomopathogenic fungus *Neozygites fumosa* on the cassava mealybug *Phenacoccus herreni*. Journal of Invertebrate Pathology, 69: 276–278.
- Del-Claro, K. and Oliveira, P.S. 2000. Conditional outcomes in a Neotropical treehopper–ant association: temporal and species-specific variation in ant protection and homopteran fecundity. Oecologia 124, 156–165.

- Derzelle, S., Ngo S., Turlin, E., Duchaud, E., Namane, A., Kunst, F., Danchin, A., Bertin, P. and Charles J.F. 2004. AstR-AstS, a new two-component signal transduction system, mediates swarming, adaptation to stationary phase and phenotypic variation in *Photobradybus luminescens*. *Journal of the Society for General Microbiology* 150, 897-910.
- Dhawan, A.K., Kamaldeep, S.A. and Sarika, S. 2009. Distribution of mealybug, *Phenacoccus solenopsis* Tinsley in cotton with relation to weather factors in South-Western districts of Punjab. *Journal of Entomological Research* 33(1), 59-63.
- Dikshith, T.S.S. 1966. Studies on the cytology of coccids. II Cytology of *Rastrococcus iceryoides* (Green) (Pseudococcidae: Coccoidea). *Cytologia*, 31: 117-124.
- Duan, J.J. and Messing, R.H. 2000. Effects of host substrate and vibration cues on ovipositor probing behaviour in two larval parasitoids of tephritid fruit flies. *Journal of Insect Behaviour* 13 (2), 175-186.
- Dufour, B. 1991. Place and importance of different insect species in the ecology of Cssv (Cocoa swollen shoot virus) in Togo. *Café Cacao Thé* 35: 197–204.
- Dunkelblum, E., Ben-Dov, Y., Goldschmidt, Z., Wolk, J.L. and Somekh, L. 1987. Synthesis and field bioassay of some analogs of sex pheromone of citrus mealybug, *Planococcus citri* (Risso). *Journal of Chemical Ecology*, 13: 863–871.
- Eisner, T. and Silberglied, R.E. 1988. A chrysopid larva that cloaks itself in mealybug wax. *Psyche* 95:15–20.
- Ekesi, S., Nderitu, P.W. and Chang, C.L. 2007. Adaptation to and small-scale rearing of invasive fruit fly *Bactrocera invadens* (Diptera: Tephritidae) on artificial diet. *Annals of Entomological Society of America* 100: 562–567.
- Elith, J., Graham, C.H. and Anderson, R.P. 2006. Novel methods to improve prediction of species' distributions from occurrence data. *Ecography*, 29, 129–151.
- Entomological society of Nigeria, 1991. Newsletter 10, pg 14.
- Entwistle, P.F. and Longworth, J.F. 1963. Relationships between cacao viruses and their vectors – feeding behaviour of three mealybug (Homoptera : Pseudococcidae) species. *Annals of Applied Biology* 52: 387–391.
- EON, 2001. Encyclopedia of the Nations: Tanzania-Climate. <http://www.nationsencyclopedia.com/Africa/Tanzania-CLIMATE.html>.
- Fabres, G. and Le Rü, B. 1988. Plant-insect relationships studies to improve cassava mealybug regulation methods. Proceedings of the Seventh Symposium of The International Society for Tropical Root Crops organized by INRA, 1-6 July 1985, GosierGuadeloupe, Editions INRA, p. 563-577.
- Fang, X.P., Rieser, M.J., Gu, Z.M., Zhao, G.X, and McLaughlin, J.L. 1993. Annonaceous acetogenins: an update review. *Phytochemical Analysis* 4: 27–48.
- FAO. 2004. The Food and Agriculture Organisation database (FAOSTAT). Published on the internet; <http://apps.fao.org> [accessed 18 February 2011].

- Ferris, G.F. 1954. Report upon scale insects collected in China (Homoptera: Coccoidea) Part V. Microentomology, 19: 51-66.
- Fielding, A.H. and Bell, J.F. 1997. A review of methods for the assessment of prediction errors in conservation presence/ absence models. Environmental Conservation 24, 38–49.
- Flaherty, D.L., Peacock, W.L., Bettiga, L. and Leavitt, G.M. 1982. Chemicals losing effect against grape mealybug. California Agriculture 36: 15-16.
- Flanders, S.E. 1943. The Argentine ant versus the parasites of the black scale. The California Citrograph, March. pp. 117–137.
- Flanders, S.E. 1945. Coincident infestations of *Aonidiella citrina* and *Coccus hesperidium*, a result of ant activity. Journal of Economic Entomology, 38, 711–712.
- Flanders, S.E. 1951. The role of the ant in the biological control of homopterous insects. The Canadian Entomologist, 83: 93-98.
- Flanders, S.E. 1958. The role of the ants in the biological control of scale insects in California. Proceedings of the X International Congress of Entomology, 4, 579–584, Montreal, Canada.
- Fletcher, B.S. 1989. Temperature-development rate relationships of immature stages and adult of tephritid fruit flies. pp. 273–289 in Robinson, A.S. & Hooper, G. (Eds) Fruit Flies: Their Biology, Natural Enemies and Control. Amsterdam, The Netherlands, Elsevier.
- Floridata. 2001. Online information: *Parkinsonia aculeata*. Floridata.com L.C. Available: http://www.floridata.com/ref/p/park_acu.cfm. 3 p (Accessed: February 10, 2011).
- Foldi, I. 1983. Structure and functions of the integumentary glands of mealybugs Pseudococcidae and of their secretions. Annales de la Société Entomologique de France 19: 155–166.
- Folkina, M.Y. 1978. A case of mass extermination of the San Jose scale *Quadraspidiotus perniciosus* by an ant *Crematogaster subdentata*. Journal of Zoology 57: 301.
- Franco, J.C., Silva, E.B. and Carvalho, J.P. 2000. Mealybugs (Hemiptera, Pseudococcidae) associated with citrus in Portugal. ISA Press, Lisbon.
- Franco, J.C., Silva, E.B., Cortegano, E., Campos, L., Branco, M., Zada, A. and Mendel, Z. 2008c. Kairomonal response of the parasitoid *Anagyrus* spec. nov near *pseudococci* to the sex pheromone of the vine mealybug. Entomologia Experimentalis and Applicata 126: 122–130.
- Franco, J.C., Suma, P., Silva E.B., Blumberg, D. and Mendel, Z. 2004b. Management strategies of mealybug pests of citrus in Mediterranean countries. Phytoparasitica 32 (5), 507-522.
- Franco, J.C., Zada, A. and Mendel, Z. 2009. Novel Approaches for the Management of Mealybug Pests. In: I. Ishaaya and A.R. Horowitz (eds.), Biorational Control of Arthropod Pests. Pp 233-278.

- Fu Castillo, A.A., Miranda Blanco, J.L., Osorio Acosta, G. and Martinez Carrillo, J.L. 2004. Chemical control of mealybug *Planococcus ficus* Signoret (Homoptera: Pseudococcidae) in table grapes. Agricultural Technology of Mexico 30: 101–105.
- Fujimoto, Y., Eguchi, T., Kakinuma, K., Ikekawa, N., Sahai, M, and Gupta, Y. K. 1998. Squamocin, a new cytotoxic bistetra hydrofuran containing acetogenin from *Annona squamosa*. Chemical & Pharmaceutical Bulletin (Tokyo) 36: 4802–4806.
- Gado, A and Neuenschwander, P. 1993. Report on a survey from 16 to 30-6-93 for the mango mealybug *Rastrococcus iceryoides* and its natural enemies in Tanzania. IITA internal report, pp. 8.
- Galán Saúco, V. 2002. Mango: Emerging tropical export crop. Chronica Horticulturae 42, 4: 14–17.
- Galán Saúco, V. 2004. Mango production and world market: Current situation and future prospects. Proceedings of the VIIth International Symposium on Mango, Editors Pinto, A.C.Q. et al. Acta Horticulturae 645: 107-116.
- Garcia-Valente, F., Ortega-Arenas, L.D., Gonzalez-Hernandez, H., Villanueva- Jimenez, J.A., Lopez-Collado, J., Gonzalez-Hernandez, A., and Arredondo-Bernal, H.C. 2009. Natural and induced parasitism of *Anagyrus kamali* against pink hibiscus mealybug on teak shoots in Bahia de Banderas, Nayarit. *Agrociencia*, 43(7), 729-738.
- Gevrey, M. and Worner, S.P. 2006. Prediction of global distribution of insect pest species in relation to climate by using an ecological informatics method. Journal of Economic Entomology 99:979–986.
- Gibernau, M. and Dejean, A. 2001. Ant protection of a Heteroptera trophobiont against a parasitoid wasp. Oecologia 126, 53–57.
- Gillani, W.A. and Copland, M.J.W. 1999. Defensive behaviour of the longtailed mealybug *Pseudococcus longispinus* (Targioni Tozzetti) (Hemiptera: Pseudococcidae) against the brown lacewing *Symppherobius fallax* Navas (Neuroptera: Hemerobiidae). Entomologica, Bari 33:279–285.
- Giordanengo, P. and Nenon, J.P. 1990. Melanization and encapsulation of eggs and larvae of *Epidinocarsis lopezi* by its host *Phenacoccus manihoti* – effects of superparasitism and egg- laying patterns. Entomologia Experimentalis and Applicata 56: 155–163.
- Godfray, H.C.J. 1994. Parasitoids: behaviour and evolutionary ecology. Princeton University Press, Princeton, NJ, USA.
- Gonzalez, R.H., Jorge Poblete, G. and Gerardo-Barria, P. 2001. The tree fruit mealybug in Chile, *Pseudococcus viburni* (Signoret), (Homoptera: Pseudococcidae). Review Frutic 22: 17–26.
- Goolsby, J.A., Alan A. K. and Meyerdirk, D.E. 2002. Seasonal Phenology and Natural Enemies of *Maconellicoccus hirsutus* (Hemiptera: Pseudococcidae) in Australia. The Florida Entomologist, 85(3): 494-498.
- Goolsby, J.A., DeBarro, P.J., Kirk, A.A., Sutherst, R.W., Canas, L., Ciomperlik, M.A., Ellsworth, P.C., Gould, J.R., Hartley, D.M., Hoelmer, K.A., Naranjo, S.E., Rose, M.,

- Roltsch, W.J., Ruiz, R.A., Pickett, C.H. and Vacek, D.C. 2005. Post-release evaluation of biological control of *Bemisia tabaci* biotype “B” in the USA and the development of predictive tools to guide introductions for other countries. *Biological Control* 32:70–77.
- Grafton-Cardwell, E.E. and Reagan, C.A. 1999. Southern fire ant insecticide efficacy trial, 1997. *Arthropod. Management Tests*, 24, 92.
- Greathead, D.J. 1971. A review of biological control in the Ethiopian region. Technical Bulletin of the Commonwealth Institute of Biological Control, no. 5. Commonwealth Agricultural Bureau, Farnham Royal, Slough.
- Greathead, D.J. 2003. Historical overview of biological control in Africa. In “Biological Control in IPM Systems in Africa” (P. Neuenschwander, C. Borgemeister and J. Langewald, Eds.), CABI Publishing, Wallingford, U.K, pp. 1-26.
- Greathead, D.R. 1986. Parasitoids in classical biological control. In J. Waage and D. Greathead (eds.), *Insect Parasitoids*, pp. 289-318. Academic Press, New York. Pp 383.
- Griesbach, J. 2003. Mango growing in Kenya. Nairobi: World Agroforestry Centre (ICRAF). ISBN 92 9059 149 8.
- Griesbach, J. 1992. A guide to propagation and cultivation of fruit trees in Kenya. Schriftreihe der GTZ, No. 230. Eschborn, Germany: GTZ. ISBN: 3 88085 482 3.
- Grinnell, J. 1917. Field tests of theories concerning distributional control. *American Naturalist* 51, 115–128.
- Grinnell, J. 1924. Geography and evolution. *Ecology* 5, 225–229.
- Gross, S., Biraty, Y. and Gal, S. 2001. Using powdery and microcapsular prepares to decimate ant populations on citrus trees. *Alon Hanotea* 55: 219–221.
- Gross, S., Gefen, D., Rotman, N., Tadmor, U., Zemer, B., Gotlib, A. and Gefen, Y. 2000. Chemical control of the spherical mealybug (*Nipaecoccus viridis*) (Newstead) in citrus. *Alon Hanotea* 54: 234– 240.
- Grout, T.G. and Stephen, P. 2005. Use of an inexpensive technique to compare systemic insecticides applied through drip irrigation systems in citrus. *Africa Entomology*, 13: 353–358.
- Gullan, P. and Martin, J.H. 2003. Sternorrhyncha (jumping plant lice, whiteflies, aphids, and scale insects). In: Resh VH, Cardé RT (Eds) *Encyclopedia of insects*. Academic, Amsterdam.
- Gullan, P.J. and Kosztarab, M. 1997. Adaptations in scale insects. *Annual Review of Entomology* 42: 23–50.
- Guo, Q.F., Qian, H., Ricklefs, R.E. and Xi, W.M. 2006. Distributions of exotic plants in eastern Asia and North America. *Ecology Letters* 9:827–834.
- Gutierrez, A.P., Daane, K.M., Ponti, L., Walton, V.M. and Ellis, C.K. 2008a. Prospective evaluation of the biological control of vine mealybug: refuge effects and climate. *Journal of Applied Ecology*, 45: 524–536.

- Gutierrez, A.P., Neuenschwander, P, and Vanalphen, J.J.M. 1993. Factors affecting biological control of cassava mealybug by exotic parasitoids – a ratio-dependent supply-demand driven model. *Journal of Applied Ecology*, 30:706–721.
- Gutierrez, A.P., Ponti, L., d'Oultremont, T. and Ellis, C.K. 2008b. Climate change effects on poikilotherm tritrophic interactions. *Climate Change* 87: S167–S192.
- Hala, N., Quilici S., Gnago, A.J., N'Depo, O.R. N'D a Adopo A. Kouassi, P. and Allou, K. 2004. "Status of fruit flies (Diptera: Tephritidae) in Côte d'Ivoire and implications for mango Exports". *Fruits of Economic importance: from basic to applied knowledge, proceedings of the 7th International Symposium on fruits flies of economic importance*, 10-15 September 2006, Salvador, Brazil: 233-239.
- Hall, D.G., Roda, A., Lapointe, S.L. and Hibbard, K. 2008. Phenology of *Maconellicoccus hirsutus* (Hemiptera: Pseudococcidae) in Florida based on attraction of adult males to pheromone traps. *Florida Entomologist*, 91: 305–310.
- Hall, R.W. and Ehler, L.E. 1979. Rate of establishment of natural enemies in classical biological control. *Bulletin of the Entomological Society of America* 25:280–282.
- Hall, W.J. 1945. The identity of a mealybug vector of swollen shoot-virus disease of cacao in West Africa. *Bulletin of Entomological Research*, 36: 305–313.
- Hance, T., van Baaren, J., Vernon, P. and Boivin, G. 2007. Impact of extreme temperatures on parasitoids in a climate change perspective. *Annual Review of Entomology* 52:107–126.
- Hanemann, L.P. 2006. An Assessment of the Export Competitiveness of the Banana, Mango and Mango Pulp Sectors in the State of Tamil Nadu. Paper for the World Bank, Abt Associates, Inc.
- Hanks, L.M. and Sadof, C.S. 1990. The effect of ants on nymphal survivorship of *Coccus viridis* (Homoptera: Coccoidea). *Biotropica*, 22: 210-213.
- Hardin, M.R., Benrey, B., Coll, M., Lamp, W.O., Roderick, G.K. and Barbosa, P. 1995. Arthropod pest resurgence: an overview of potential mechanisms. *Crop Protection*, 14: 3–18.
- Harris, S.C. 1982. Nitrogen fixation by tropical woody legumes: potential source of soil enrichment. In: J.P. Roskoski, J. Monano, C. van Kessel, G. Castilleja, and P.H. Graham, eds. *Biological nitrogen fixation technology for tropical agriculture*. Centro International de Agricultura Tropical. Cali, Colombia. p. 447 - 454.
- Hassell, M.P. 1978. *The Dynamics of Arthropod predator-Prey Systems*. Princeton University Press, Princeton, N. J. 237pp.
- Hassell, M.P. and Waage, J.K. 1984. Host-parasitoid population interactions. *Annual Review Entomology*, 29: 89–114.
- Haukioja, E. and Neuvonen, S. 1985. The relationship between size and reproductive potential in male and female *Epirita autumnata* (Lep., Geometridae). *Ecological Entomology* 10: 267-270.

- Hayat, M. 1998. Aphelinidae of India (Hymenoptera: Chalcidoidea): A taxonomic revision. *Memoirs on Entomology, International*. Vol. 13: 1-422.
- HCA. 2010. Horticultural Council of Africa. Published on the internet; <http://fpeak.org/hca.html> [accessed 18 February 2011].
- Hcidari, M. and Jahan, M. 2000. A study of ovipositional behaviour of *Anagyrus pseudococci* a parasitoid of mealybugs. *Journal of Agriculture, Science and Technology* 2, 49-53.
- Helms, K.R. and Vinson, S.B. 2002. Widespread association of the invasive ant *Solenopsis invicta* with an invasive mealybug. *Ecology* 83:2425–2438.
- Hernandez, P.A., Graham, C.H., Master, L.L. and Albert, D.L. 2006. The effect of sample size and species characteristics on performance of different species distribution modeling methods. *Ecography* 29, 773–785.
- Heu, R.A., Fukada, M.T., Conant, P., 2007. Papaya mealybug, *Paracoccus marginatus* Williams and Granara de Willink (Hemiptera: Pseudococcidae). New Pest Advisory. 4(3). Department of Agriculture, State of Hawaii, Honolulu, HI.
- Herren, H.R., Neuenschwander, P., Henessey, R.D. and Hammond, W.N.O. 1987. Introduction and dispersal of *Epidinocarsis lopezi* (Hymenoptera: Encyrtidae), an exotic parasitoid of the cassava mealybug *Phenacoccus manihoti* (Homoptera: Pseudococcidae), in Africa. *Agriculture, Ecosystems and Environment* 19, 131–144.
- Higgins, S.I., Richardson, D.M., Cowling, R.M. and Trinder-Smith, T.H. 1999. Predicting the landscape scale distribution of alien plants and their threat to plant diversity. *Conservation Biology*. 13: 303 - 313.
- Ho, C.T. and Khoo, K.C. 1997. Partners in biological control of cocoa pests: Mutualism between *Dolichoderus thoracicus* (Hymenoptera: Formicidae) and *Cataenococcus hispidus* (Hemiptera: Pseudococcidae). *Bulletin Entomology Research*, 87: 461–470.
- Hodges, G. and Hodges, A. 2004. New invasive species of mealybugs, *Palmicitor lumpurensis* and *Chaetococcus bambusae* (Hemiptera: Coccoidea: Pseudococcidae), on bamboo in Florida. *Florida Entomologist* 87: 396–397.
- Hodgson, C., Abbas, G., Arif, M.J., Saeed, S. and Krar, H. 2008. *Phenacoccus solenopsis* Tinsley (Sternorrhyncha: Coccoidea: Pseudococcidae), an invasive mealybug damaging cotton in Pakistan and India, with a discussion on seasonal morphological variation. *Zootaxa* 1913: 1–35.
- Hoelmer, K.A. and Kirk, A.A. 2005. Selecting arthropod biological control agents against arthropod pests: can the science be improved to decrease the risk of releasing ineffective agents? *Biological Control* 34:255–264.
- Hoffmann, M.H. 2001. The distribution of *Senecio vulgaris*: capacity of climatic range models for predicting adventitious ranges. *Flora* 196: 395 - 403.
- Hogendorp, B.K., Cloyd, R.A. and Swiader, J.M. 2006. Effect of nitrogen fertility on reproduction and development of citrus mealybug, *Planococcus citri* Riso (Homoptera:

- Pseudococcidae), Feeding on two colors of Coleus, *Solenostemon scutellarioides* L. Codd, Environmental Entomology 35(2): 201-211.
- Høgh-Jensen, H., Myaka, F.A., Sakala, W.D., Kamalongo, D., Ngwira, A., Vesterager, J.M., Odgaard, R. and Adu-Gyamfi, J.J. 2007. Yields and qualities of pigeon pea varieties grown under smallholder farmers' conditions in Eastern and Southern Africa. African Journal of Agricultural Research 2(6): 269-278.
- Hölldobler, B. and Edward, O.W. 1978. The multiple recruitment systems of the African weaver ant *Oecophylla longinoda* (Latreille) (Hymenoptera: Formicidae). Behavioral Ecology Sociobiology, 3: 19-60.
- Hölldobler, B. and Wilson, E.O. 1990. The ants. Cambridge, Massachusetts (MA): The Belknap Press of Harvard University Press. 732 pp.
- Holler, C., Borgemeister, C., Haardt, H. and Powell, W. 1993. The relationship between primary parasitoids and hyperparasitoids of cereal aphids: an analysis of field data. Journal of Animal Ecology 62, 12-21.
- Horton, J.R. 1918. The Argentine ant in relation to citrus groves. Bulletin No. 647. Government Printing Office, Washington, USA.
- Hulspas-Jordaan, P.M. and van Lenteren, J.C. 1978. The relationship between host plant leaf structure and parasitisation efficiency of the parasitic wasp *Encarsia formosa* Gahan (Hymenoptera: Aphelinidae). Mededelingen van de Faculteit Landbouwwetenschappen, Rijksuniversiteit Gent 43, 431–440.
- Hulting, F.L., Orr, D.B. and Obrycki, J.J. 1990. Computer program for calculation and statistical comparison of the intrinsic rate of increase and associated life table parameters. Florida Entomologist 73, 602–612.
- ICIPE. 2006. Development and Implementation of a Sustainable IPM Program for Major Mango Pests and Opportunity for Improving Market and Processing Information in Sub-Saharan Africa. Proposal submitted to the German Ministry for Economic Cooperation and Development (BMZ).
- Inkerman, P.A., Ashbolt, N.J., Carver, M. and Williams, D.J. 1986. Observations on the pink sugarcane mealybug, *Saccharicoccus sacchari* (Cockerell), in Australia (Homoptera: Pseudococcidae). Proc. 19th Congr. Int. Soc. Sugar Cane Technol. (Jakarta, Indonesia) 1:612-616.
- Iperti, G. 1999. Biodiversity of predaceous coccinellidae in relation to bioindication and economic importance. Agriculture, Ecosystems & Environment 74:323–342.
- IPNI. 2005. International Plant Name Index. – Published on the Internet <http://www.ipni.org/index.html>. Viewed February 2, 2011.
- Irulandi, S., Kumar, P.K.V., Seetharama, H.G. and Sreedharan, K. 2001. Bioefficacy of neem formulations alone and in combination with synthetic insecticide against mealybug, *Planococcus citri* (Risso) on *Coffea*. Journal of Coffee Research 29: 56–60.

- Islam, K.S. and Copland, M.J.W. 1997. Host preference and progeny sex ratio in a solitary koinobiont mealybug endoparasitoids, *Anagyrus pseudococci* (Girault) in response to its host stage. *Biocontrol Science and Technology* 7, 449–456.
- Islam, K.S. and Copland, M.J.W. 2000. Influence of egg load and oviposition time interval on the host discrimination and offspring survival of *Anagyrus pseudococci* (Hymenoptera: Encyrtidae), a solitary endoparasitoid of citrus mealybug, *Planococcus citri* (Hemiptera: Pseudococcidae). *Bulletin of Entomological Research* 90: 69-75.
- Itioka, T. and Inoue, T. 1996. The role of predators and attendant ants in regulation and persistence of a population of the Citrus Mealybug *Pseudococcus citriculus* in a Satsuma orange orchard. *Applied Entomology and Zoology*, 31, 195–202.
- Ivbijaro, M. F., Udensi, N., Ukwela U.M. and Anno-Nyako, F.V. 1991. "Geographical distribution and host range in Nigeria of the mango mealy bug, *Rastrococcus invadens* Williams, a serious exotic pest of horticulture and other crops". *Insect Science and Its Application* 13: 411-416.
- Ivbijaro, M.F. and Udensi, N. 1988. "A preliminary survey of Lagos, Ogun and Oyo State for the incidence of attack of mangoes and other plants by a complex of mealybug and fungi", a study report submitted through the National Horticultural Research Institute, Ibadan, to the Federal Department of Agriculture, Abuja, September 19, 8pp.
- James, D.G. 2004. Effect of buprofezin on survival of immature stages of *Harmonia axyridis*, *Stethorus punctum pictipes* (Coleoptera: Coccinellidae), *Orius tristis* (Hemiptera: Anthocoridae), and *Geocoris* spp. (Hemiptera: Geocoridae). *Journal Economic Entomology* 97: 900–904.
- James, D.G., Stevens, M.M., and O'Malley, K.J. 1998. Prolonged exclusion of foraging ants (Hymenoptera: Formicidae) from citrus trees using controlled-release chlorpyrifos trunk bands. *International Journal of Pest Management*, 44, 65–69.
- James, D.G., Stevens, M.M., O' Malley, K.J. and Heffer, R. 1996. Activity of a hydramethylnon-based bait against the citrus ant pests, *Iridomyrmex rufoniger* gp. spp. and *I. purpureus*. *Plant Protection Quart*, 11, 122–125.
- Jervis, M.A. and Copland, M.J.W. 1996. The life cycle. pp. 376–394 in Jervis, M.A. & Kidd, N. (Eds) *Insect natural enemies: practical approaches to their study and evaluation*. London, Chapman and Hall.
- Jiggins, C., Majerus, M.E.N. and Gough, U. 1993. Ant defense of colonies of *Aphis fabae* Scolopi (Hemiptera: Aphididae), against predation by lady birds. *British Journal of Natural History* 6, 129–137.
- Joubert, C.J. 1943. The introduction into the Union of South Africa of some natural enemies of mealybugs. *Journal of Entomological Society of Southern Africa* 6: 131-136.
- Joubert, P.H., Daneel, M.S. and Grove, T. 2000. Progress towards integrated pest management (IPM) on mangoes in South Africa. *Acta Horticulturae No. 509*, 811-817.

- Joyce, A.L., M.S. Hoddle, T.S. Bellows and Gonzalez, D. 2001. Oviposition behaviour of *Coccidoxyenoides peregrinus*, a parasitoid of *Planococcus ficus*. *Entomologia Experimentalis and Applicata* 98: 49-57.
- Kairo, M.T.K. and Murphy, S.T. 1999. Host age choice for oviposition in *Pauesia juniperorum* (Hymenoptera: Braconidae, Aphidiinae) and its effect on the parasitoid's biology and host population growth. *Biocontrol Science and Technology*, 9: 475-486.
- Kairo, M.T.K., Pollard, G.V., Peterkin, D.D. and Lopez, V.F. 2000. Biological control of the hibiscus mealybug, *Maconellicoccus hirsutus* Green (Hemiptera: Pseudococcidae) in the Caribbean. *Integrated Pest Management Reviews* 5: 241-54.
- Kairo, M.T.K., Pollard, G.V., Peterkin, D.D. and Lopez, V.F. 2000. Biological control of the hibiscus mealybug, *Maconellicoccus hirsutus* Green (Hemiptera: Pseudococcidae) in the Caribbean. *Integrated Pest Management Reviews* 5, 241-54.
- Karamaouna, F. and Copland, M.J.W. 2000. Host suitability, quality and host size preference of *Leptomastix epona* and *Pseudaphycus flavidulus*, two endoparasitoids of the mealybug *Pseudococcus viburni*, and host size effect on parasitoid sex ratio and clutch size. *Entomologia Experimentalis and Applicata* 96: 149 – 158.
- Khosla, S., Mendiratta, G. and Brahmachari, V. 2006. Genomic imprinting in the mealybugs. *Cytogenetic and Genome Research* 113: 41–52.
- Kim, K.C. 1993. Insect pests and evolution. In: Kim KC, McPheron BA (Eds) *Evolution of insect pests: patterns of variation*. Wiley, New York.
- Kiritani, K. 2006. Predicting impacts of global warming on population dynamics and distribution of arthropods in Japan. *Population Ecology* 48:5–12.
- Klingauf, F.A. 1987. Feeding, adaptation and excretion. In: *Aphids Their Biology, Natural Enemies and Control*, Vol. A. Ed. By Minka, AK, Harrewijn P. Amsterdam, The Netherlands: Elsevier: pp. 225–253.
- Klotz, J.H., Greenberg, L. and Venn, E.C. 1998. Liquid boric acid bait for control of the Argentine ant (Hymenoptera: Formicidae). *Journal of Economic Entomology*, 91, 910–914.
- Klotz, J.H., Vail, K.M. and Williams, D.F. 1997. Toxicity of a boric acid– sucrose water bait to *Solenopsis invicta* (Hymenoptera:Formicidae). *Journal of Economic Entomology*, 90, 488–491.
- Kontodimas, D.C., Eliopoulos, P.A., Stathas, G.J. and Economou, L.P. 2004. Comparative temperature-dependent development of *Nephush includens* (Kirsch) and *Nephush bisignatus* (Bohemian) (Coleoptera: Coccinellidae) preying on *Planococcus citri* (Risso) (Homoptera: Pseudococcidae): evaluation of a linear and various nonlinear models using specific criteria. *Environmental Entomology* 33: 1–11.
- Koricheva, J., Larsson, S. and Aukioja, E.H. 1998. Insect performance on experimentally stressed woody plants: a meta-analysis. *Annual Review of Entomology* 43: 195-216.
- Kosztarab, M. and Kozár, F. 1988. Scale insects of Central Europe. Dr. W. Junk Publishers, Dordrecht.

- Kottek, M., Grieser, J., Beck, C., Rudolf, B. and Rubel, F. 2006. World map of the Koppen-Geiger climate classification updated. *Meteorologische Zeitschrift* 15, 259–263.
- Kouamé, K.I., and Mackauer, M. 1991. Influence of aphid size, age and behaviour on host choice by the parasitoid wasp *Ephedrus californicus*: a test of host-size models. *Oecologia*, 88, 197–203.
- Kriegler P.J. and Whitehead V.B. 1962. Notes on the biology and control of *Crematogaster peringueyi* var. *anguistor*, Arnold on grape vines (Hymenoptera: Formicidae). *Journal of Entomological Society of Southern Africa* 25: 287-290.
- Kriticos, D.J. and Randall, R.P. 2001. A comparison of systems to analyze potential weed distributions. In: Groves, R.H., Panetta, F.D., Virtue, J.G. (Eds.), *Weed Risk Assessment*. CSIRO Publishing, Melbourne, pp. 61–79.
- Kubiriba, J., Legg, J.P., Tushemereirwe, W. and Adipala, E. 2001. Vector transmission of Banana streak virus in the screenhouse in Uganda. *Annals of Applied Biology*, 139: 37–43.
- Kumashiro, B.R., R.A. Heu, G.M. Nishida, and Beardsley, J.W. 2001. New state records of immigrant insects in the Hawaiian Islands for the year 1999. *Proceedings of the Hawaiian Entomological Society*, 35: 170-184.
- Lacey, E.P. 1998. What is an adaptive environmentally induced parental effect? In: Mousseau TA, Fox CW (eds) *Maternal effects as adaptations*. Oxford University Press, New York, USA, pp. 54–66.
- Lach, L., 2003. Invasive ants: unwanted partners in ant–plant interactions? *Annals of the Missouri Botanical Garden* 90, 91–108.
- Larrain, P.S. 1999. Effect of chemigation and painted applications of imidacloprid (ConfidorReg.) upon *Pseudococcus viburni* (Signoret) (Homoptera: Pseudococcidae) populations in table grapes. *Agric Tec Santiago* 59: 13–25.
- Larraín, S.P. 2002. Insect and mite pest incidence on sweet pepinos, *Solanum muricatum* (Ait.) cultivated in the IV Region, Chile. *Agriculture T'ecnica*, 62, 15–26.
- Le Rü B., Izquierdo Y., Biassangama A. and Kiyindou, A. 1991. Variations d'abondance et facteurs de régulation de la cochenille du manioc *Phenacoccus manihoti* (Hom. Pseudococcidae) cinq ans après l'introduction d'*Epidinocarsis lopezi* (Hym. Encyrtidae) au Congo en 1982. *Entomophaga* 36: 499-511.
- Legner, E.F. and Bellows, T.S. 1999. Exploration for natural enemies. In. Bellows, T.S., Fisher, T.W. (Eds.), *Handbook of Biological Control*. Academics Press, San Diego, pp 87-101.
- Leru, B. 1986. Epizootiology of the entomophthoraceous fungus *Neozygites fumosa* in a population of the cassava mealybug, *Phenacoccus manihoti* (Hom, Pseudococcidae). *Entomophaga* 31: 79–89.
- Leru, B. and Tertuliano, M. 1993. Tolerance of different host-plants to the cassava mealybug *Phenacoccus manihoti* Matile-Ferrero (Homoptera, Pseudococcidae). *International Journal of Pest Management* 39: 379–384.

- Leru, B., Diangana, J.P. and Beringar, N. 1994. Effects of nitrogen and calcium on the level of resistance of cassava to the mealybug *P. manihoti*. Insect Science and its Application 15: 87–96.
- Liu, F., Li, W., Wang, Y. and Lin, J. 1995. Convenient syntheses of some analogs of the sex pheromone of citrus mealybug, *Planococcus citri* (Risso). Synthetic Communications 25: 3837–3843.
- Lockhart, B.E.L., Autrey, L.J.C. and Comstock, J.C. 1992. Partial purification and serology of Sugarcane mild mosaic virus, a mealybug-transmitted closterolike virus. Phytopathology 82: 691–695.
- López, O.P., Hénaut, Y., Cancino, J., Lambin, M., Cruz-López L., and Rojas, J.C. 2009. Is host size an indicator of quality in the mass-reared parasitoid *Diachasmimorpha longicaudata* (Hymenoptera: Braconidae). Florida Entomologist 92(3), 441-449.
- Luck, R., P.S. Messenger and Barbieri, J.F. 1981. The influence of hyperparasitism on the performance of biological control agents. In: D. Rosen (ed), *The role of hyperparasitism in biological control: a symposium*. Division of Agricultural Sciences, University of California Publications, 4103. pp. 34–42.
- Luck, R.F. and Forster, L.D. 2003. Quality of augmentative biological control agents: a historical perspective and lessons learned from evaluating *Trichogramma*. In: van Lenteren JC (Ed) Quality control and production of biological control agents: theory and testing procedures. CABI, Wallingford Gutierrez et al. (2008a).
- Luhanga, W.W., and Gwinner, J. 1993. Mango mealybug (*Rastrococcus iceryoides*) on *Mangifera indica* in Malawi. FAO Plant Protection Bulletin 41(2), 125-126.
- Lunderstadt, J. 1998. Impact of external factors on the population dynamics of beech scale (*Cryptococcus fagisuga*) (Homoptera: Pseudococcidae) in beech (*Fagus sylvatica*) stands during the latency stage. Journal of Applied Entomology, 122: 319–322.
- Lux, S.A, Copeland R.S, White I.M, Manrakhan A and Billah M.K. 2003. A new invasive fruit fly species from the *Bactrocera dorsalis* (Hendel) group detected in East Africa. Insect Science and its Application, 23(4), 355-361.
- Madulu, R.B., and Chalamila, B.N. 2007. Potential and constraints of fruit trees in Coast region in Tanzania: A case study of Mkuranga and Bagamoyo districts. African Crop Science Conference Proceedings. Vol. 8. Pp. 1399-1401.
- Maia, A.H.N., Luiz, A.J.B. and Campanhola, C. 2000. Statistical inference on associated fertility life table parameters using Jackknife technique: computational aspects. Journal of Economic Entomology 93: 511–518.
- Malsch, A.K.F., Kaufmann, E., Heckroth, H.P., Williams, D.J., Maryati, M. and Maschwitz, U. 2001. Continuous transfer of subterranean mealybugs (Hemiptera, Pseudococcidae) by *Pseudolasius* spp. (Hymenoptera: Formicidae) during colony fission? Insectes Sociaux 48: 333–341.

- Maneepun, S. and Yunchalad, M. 2004. Developing processed mango products for international markets. Proceedings of the VIIth International Symposium on Mango, Editors Pinto, A.C.Q. et al. *Acta Horticulturae* 645: 93–105.
- Mani and Krishnamoorthy, A. 1998b. Regulation of *Rastrococcus iceryoides* (Green) on guava. *Insect Environment* 4(3): 71.
- Marohasy, J. 1997. Acceptability and suitability of seven plant species for the mealybug *Phenacoccus parvus*. *Entomologia Experimentalis et Applicata* 84: 239 - 246.
- Marohasy, J. 2003. Acceptability and suitability of seven plant species for the mealybug *Phenacoccus parvus*. *Entomologia Experimentalis et Applicata* 84: 239–246.
- Maron, J.L., Vila', M., Bommarco, R., Elmendorf, S. and Beardsley, P. 2004. Rapid evolution of an invasive plant. *Ecological Monographs* 74: 261 - 280.
- Martin, N.A. and Workman, P.J. 1999. Efficacy of insecticides for long-tailed mealybug control. In: Proceedings of the Fifty Second New Zealand Plant Protection Conference, Auckland Airport Centre, Auckland, New Zealand, 10–12 August 1999. Wine Industry Association, Western Australia.
- Martinez-Ferrer, M.T., Grafton-Cardwell, E.E. and Shorey, H.H. 2003. Disruption of parasitism of the Californian red scale (Homoptera: Diaspididae) by three ant species (Hymenoptera: Formicidae). *Biological Control*, 26, 279–286.
- Martinez-Meyer, E., Peterson, A.T. and Hargrove, W.W. 2004. Ecological niches as stable distributional constraints on mammal species, with implications for Pleistocene extinctions and climate change projections for biodiversity. *Global Ecology and Biogeography* 13, 305–314.
- Matile-Ferrero, D. 1977. New species infesting cassava in Equatorial Africa, *Phenacoccus manihoti* n. sp. (Homoptera-Coccoidea-Pseudococcidae). *Annales de la Société Entomologique de France* 13:145–152
- Matokot, L., Reyd, G., Malonga, P. and Ru, B.I. 1992. Population dynamics of *Rastrococcus invadens* (Homoptera: Pseudococcidae) in the Congo; influence of accidental introduction of the Asiatic parasitoid *Gyranusoidea tebygi* (Hymenoptera: Encyrtidae). *Entomophaga* 37: 123–140.
- May, R.M. and Hassell, M.P. 1981. The dynamics of multiparasitoids-host interactions. *American Naturalist*, 117: 234–261.
- McDougall, S.J. and Mills, N.J. 1997. The influence of hosts, temperature and food sources on the longevity of *Trichogramma platneri*. *Entomologia Experimentalis et Applicata* 83: 195 – 203.
- McKenzie, H.L. 1967. Mealybugs of California with taxonomy, biology and control of North American species (Homoptera: Coccoidea: Pseudococcidae) University of California Press, Berkeley, LA, 534 pp.
- McLeod, P., Diaz, J., Vasquez, L. and Johnson, D.T. 2002. Within-plant distribution and sampling of mealybugs in plantain var. FHIA 21. *Tropical Agriculture* 79: 150–153.

- Mehta, R. and George, J. 2003 Processed Food Products Exports from India: An Exploration with SPS Regime.
- Mendel, Z., Gross, S., Steinberg, S., Cohen, M. and Blumberg, D. 1999. Trials for the control of the citrus mealybug in citrus orchards by augmentative release of two encyrtid parasitoids. *Entomologica*, Bari 33: 251–265.
- Messenger, P.S. and van den Bosch, R. 1971. The adaptability of introduced biological control agents. In. *Biological Control*, (ed.) C.B. Huffaker, Plenum Press, New York. Pp 68-92.
- Meyer, J.S., Ingersoll, C.G., McDonald, L.L. and Boyce, M.S. 1986. Estimating uncertainty in population growth rates: Jackknife vs. bootstrap techniques. *Ecology* 67:1156–1166.
- Meyerdirk, D.E., French, J.V. and Hart, W.G. 1982. Effect of pesticide residues on the natural enemies of citrus mealybug. *Environmental Entomology* 11: 134–136.
- Meyerdirk, D.E., Muniappan, R., Warkentin, R., Bamba, J, and Reddy, G.V.P. 2004. Biological control of the papaya mealybug, *Paracoccus marginatus* (Hemiptera: Pseudococcidae) in Guam. *P. P. Q.* 19(3): 110-114.
- Meyerdirk, D.E., Warkentin, R., Attavian, B., Gersabeck, E., Francis, A., Adams M. and Francis, G. 1998. Biological control of the pink hibiscus mealybug project manual. United States Department of Agriculture, Washington, DC.
- Mgocheki, N. and Addison, P. 2009. Interference of ants (Hymenoptera: Formicidae) with biological control of the vine mealybug *Planococcus ficus* (Signoret) (Hemiptera: Pseudococcidae). *Biological Control* 49, 180–185.
- Millar, J.G., Daane, K.M., McElfresh, J.S., Moreira, J.A., Malakar-Kuenen, R., Guillen, M. and Bentley, W.J. 2002. Development and optimization of methods for using sex pheromone for monitoring the mealybug *Planococcus ficus* (Homoptera: Pseudococcidae) in California vineyards. *Journal of Economic Entomology* 95:706–714.
- Miller, D.R., Miller, G.L. and Watson, G.W. 2002. Invasive species of mealybugs (Hemiptera: Pseudococcidae) and their threat to U.S. agriculture. *Proceedings of the Entomological Society of Washington*, 104: 825-836.
- Missouri Botanical Garden - W³ Tropicos [Web Database.]. 2006. St. Louis, Missouri. Available from <http://mobot.mobot.org/W3T/Search/vast.html> (accessed 05 April 2011).
- Mittler, T.E. and Douglas, A.E. 2003. Honeydew. In: Resh VH, Cardé RT (Eds) *Encyclopedia of insects*. Academic, Amsterdam.
- Mohamed, S.A., Overholt, W.A., Wharton, R.A., Lux, S.A. and Eltoum, E.M. 2003. Host specificity of *Psyllalia cosyrae* (Hymenoptera: Braconidae) and the effect of different host species on parasitoid fitness. *Biological Control* 28, 155-163.
- Mohamed, S.A., Wharton, R.A., von Me'rey, G., and Schulthess, F. 2006. ‘Acceptance and Suitability of Different Host Stages of *Ceratitis capitata* (Wiedemann) (Diptera: Tephritidae) and Seven Other Tephritid Fruit Fly Species to *Tetrastichus giffardii* Silvestri (Hymenoptera: Eulophidae)’, *Biological Control*, 39: 262-271.

- Moore, D. 1988. Agents used for biological control of mealybugs (Pseudococcidae). *Biocontrol News and Information* 9:209–225.
- Moore, D. 2004. Biological control of *Rastrococcus invadens*. *Biocontrol News and Information* 25: 17–27.
- Moore, D. 2004. Biological control of *Rastrococcus invadens*. *Biological Control News and Information*. 25: 17-27.
- Moore, D. and Cross, A.E. 1992. Competition between primary parasitoids, *Gyranusoidea tebygi* Noyes and *Anagyrus magicola* Noyes attacking the mealybug *Rastrococcus invadens* Williams and the influence of an hyperparasitoid. *Chartocarus hyalipennis* Hayat. *Biocontrol Science and Technology*, 2: 225-234.
- Moreno, D.S., Haney, P.B. and Luck, R.F. 1987. Chlorpyrifos and diazinon as barriers to Argentine ant (Hymenoptera: Formicidae) foraging on citrus trees. *Journal of Economic Entomology*, 80: 208–214.
- Morishita, M. 2005. Effect of bark-scraping, dormant spray of petroleum oil and applying pesticide in late spring on density of Japanese mealybug, *Planococcus kraunhiae* (Kuwana), in persimmon. *Annual Report of The Kansai Plant Protection Society* 47: 123–124.
- Morris, R.J., Müller, C.B. and Godfray, H.C.J. 2001. Field experiments testing for apparent competition between primary parasitoids mediated by secondary parasitoids. *Journal of Animal Ecology* 70, 301–309.
- Morton, J. 1987. Fruits of warm climates, Miami, Florida. Accessed online at [http://www.hort.purdue.edu/newcrop/morton/sweet orange](http://www.hort.purdue.edu/newcrop/morton/sweet%20orange).
- Moussa, J.B. and Matile-Ferrero, D. 1988. Sur la présence du nouveau ravageur, *Rastrococcus invadens* Williams, 1986 en République populaire du Congo [Hemiptera, Coccoidea, Pseudococcidae]. *Bulletin de la Société Entomologique de France* 93: 1–2.
- Mukherjee, N.G. 1919. Handbook of Sericulture. Bengal Secretariate Press, Calcutta, pp. 121-127.
- Muniappan, R., B.M. Shepard, G.W. Watson, G.R. Carner, D. Sartiami, A. Rauf, and Hammig, M.D. 2008. First report of the papaya mealybug, *Paracoccus marginatus* (Hemiptera: Pseudococcidae), in Indonesia and India. *Journal of Agricultural and Urban Entomology*, 25:37-40.
- Muniappan, R., Meyerdirk, D.E., Sengebau, F.M., Berringer, D.D. and Reddy, G.V.P. 2006. Classical biological control of the papaya mealybug, *Paracoccus marginatus* (Hemiptera: Pseudococcidae) in the Republic of Palau. *Florida Entomologist* 89: 212–217.
- Muniappan, R., Shepard, B.M., Watson, G.W., Carner, G.R., Rauf, A., Sartiami, D., Hidayat, P., Afun, J.V.K., Goergen, G. and Ziaur Rahman, A.K.M. 2011. New Records of Invasive Insects (Hemiptera: Sternorrhyncha) in Southeast Asia and West Africa. *Journal of Agricultural and Urban Entomology* 26(4):167-174.

- Muthukrishnan, N., Manoharan, T., Thevan, P.S.T. and Anbu, S. 2005. Evaluation of buprofezin for the management of grape mealybug, *Maconellicoccus hirsutus* (Green). *Journal of Entomological Research* 29: 339–344.
- Myburgh, A.C., and Siebert, M.W. 1964. Experiments on parathion-resistant mealybugs. *Deciduous Fruit Grower* 14: 190-193.
- Myburgh, A.C., Swart, P.L. and Urban, A.J. 1986. Mealybugs and Australian bugs. In: Myburgh, A.C. (Ed.) *Crop Pests in Southern Africa Vol. 1. Deciduous Fruit, Grapes and Berries*. 43–47. Plant Protection Research Institute, Department of Agriculture and Water Supply, Pretoria.
- Nakahira, K. and Arakawa, R. 2006. Development and reproduction of an exotic pest mealybug, *Phenacoccus solani* (Homoptera: Pseudococcidae) at three constant temperatures. *Applied Entomology and Zoology* 41: 573–575.
- Narai, Y. and Murai, T. 2002. Individual rearing of the Japanese mealybug, *Planococcus kraunhiae* (Kuwana) (Homoptera: Pseudococcidae) on a germinated broad bean seeds. *Applied Entomology and Zoology* 37: 295-298.
- Narasimham, A.U. and Chacko, M.J. 1988. *Rastrococcus* spp. (Hemiptera: Pseudococcidae) and their natural enemies in India as potential biocontrol agents for *R. invadens* Williams. *Bulletin of Entomological Research*, 78(4): 703-708.
- Narasimham, A.U. and Chacko, M.J. 1991. The distribution of some *Rastrococcus* spp. (Homoptera: Pseudococcidae) on mango in India. *Bulletin of Entomological Research*, 81(4): 445-448.
- Narasimham, A.U. and Sankaran, T. 1985. Some mealybugs and their natural enemies in India. *Proceedings of the National Seminar on Entomophagous Insects, Calicut*, 29-36.
- NAS 2002. Predicting invasions of non indigenous plants and plant pests. National Academy of Sciences, Washington, D.C.
- Nassar, N.M.A. 2007. Cassava genetic resources and their utilization for breeding of the crop. *Genetics and Molecular Research* 6: 1151–1168.
- Nelson-Rees, W.A. 1960. A study of sex predetermination in the mealy bug *Planococcus citri* (Risso). *Journal of Experimental Zoology* 144: 111–137.
- Nestel, D., Cohen, H., Saphir, N., Klein, M. and Mendel, Z. 1995. Spatial distribution of scale insects – comparative study using Taylor's power-law. *Environmental Entomology* 24: 506–512.
- Neuenschwander, P. 1993. Human interactions in classical biological control of cassava and mango mealybugs on subsistence farms in tropical Africa. In: *Crop Protection Strategies for subsistence Farmers*. Altieri, M. A. (Ed.), pp. 143-177.
- Neuenschwander, P. 1996. Evaluating the efficacy of biological control of three exotic homopteran pests in tropical Africa. *Entomophaga* 41: 405–424.

- Neuenschwander, P. 2001. Biological control of the cassava mealybug in Africa: A review. *Biological Control*. 21:214–229.
- Neuenschwander, P. 2003. Biological control of cassava and mango mealybugs. In *Biological Control in IPM Systems in Africa*. Neuenschwander, P., Borgemeister, C and Langewald, J. (Editors). CABI Publishing in association with the ACP-EU Technical Centre for Agricultural and Rural Cooperation (CTA) and the Swiss Agency for Development and Cooperation (SDC), pp. 45–59. ISBN: 0-85199-639-6.
- Neuenschwander, P. and Ajuonu, O. 1993. Report on a survey from 22-2 to 20-3-93 for cassava mealybug and its natural enemies and the impact of biological control on cassava growth and yield, in Tanzania. IITA internal report, 10 pp.
- Neuenschwander, P., Boavida, C., Bokonon-Ganta, A., Gado, A. and Herren, H.R. 1994. Establishment and spread of *Gyranusoidea tebygi* Noyes and *Anagyrus mangicola* Noyes (Hymenoptera: Encyrtidae), two biological control agents released against the mango mealybug *Rastrococcus invadens* Williams (Homoptera: Pseudococcidae) in Africa. *Biocontrol Science and Technology* 4: 61–69.
- Nicholas, J.A. and Inkerman, P.A. 1989. Acetic Acid Bacterial Biota of the Pink Sugar Cane Mealybug, *Saccharococcus sacchari*, and Its Environs. *Applied and environmental microbiology*, 56(3): 707-712.
- Nixon, G.E.J. 1951. The Association of Ants with Aphids and Coccids. Commonwealth Institute of Entomology, London.
- Nordlund, D.A., Lewis, W.J. and Altieri, M.A. 1988. Influences of plant produced allelochemicals on the host and prey selection behavior of entomophagous insects. In “Novel Aspects of Insect– Plant Interactions” (P. Barbosa and D. Letourneau, Eds.), pp. 65–90. Wiley, New York.
- Norgaard, R.B. 1988. The biological control of cassava mealybug in Africa. *American Journal of Agricultural Economics* 72: 366–371.
- Normark, B.B. 2003. The evolution of alternative genetic systems in insects. *Annual Review of Entomology* 48: 397–423.
- Novak, H. 1994. The influence of ant attendance on larval parasitism in hawthorn psyllids (Homoptera: Psyllidae). *Oecologia*, 99, 72–78.
- Noyes, J.S. and Hayat, M. 1994. Oriental Mealybug Parasitoids of the Anageryrini (Hymenoptera: Encyrtidae). CAB International, The Natural History Museum, London, 544 p.
- Noyes, J.S. 1988. *Gyranusoidea tebygi* sp. n. (Hymenoptera: Encyrtidae), a parasitoid of *Rastrococcus* (Hemiptera: Pseudococcidae) on mango in India. *Bulletin of Entomological Research* 78, 313–316.
- Nur, U. 1990. Heterochromatization and euchromatization of whole genomes in scale insects (Coccoidea, Homoptera) Development. (Suppl.) 1: 29–34.

- Nyambo, B. and Verschoor, R. 2005. Partnership for market access: towards a sustainable market-oriented horticultural sector in Tanzania. Wageningen UR Position Paper.
- Oberlies, N.H., Chang, C.J., and McLaughlin, J.L. 1997. Structure activity relationships of diverse Annonaceous acetogenins against multidrug resistant human mammary adenocarcionma (MCF-7/Adr) cells. *Journal of Medical Chemistry* 40: 2102–2106.
- Opp, S.B. and Luck R.F. 1986. Effect of host size on selected fitness components of *Aphytis melinus* and *A. lingnanensis* (Hymenoptera: Aphelinidae). *Annals of the Entomological Society of America* 79, 700-704.
- Orwa C., A. Mutua, R. Kindt, R. Jamnadass and Simons, A. 2009. Agroforestry Database:a tree reference and selection guide version 4.0 (<http://www.worldagroforestry.org/af/treedb/>)
- Osman, M., Bin, S. and Chettanachitara, C. 1989. Postharvest insects and other pests of rambutan, pp. 57 – 60. In P. F. Lam and S. Kosiyanchinda (Eds.). *Rambutan: Fruit development, Postharvest Physiology and Marketing in ASEAN*. ASEAN Food Handling Bureau, Kuala Lumpur, Malaysia.
- Ott, R.L. and Longnecker, M. 2010. *An Introduction to Statistical Methods and Data Analysis*, 6th ed.; Brooks/Cole, Cengage Learning: Belmont, CA, USA.
- Overholt, W.A., Ngi-Song, A.J., Omwega, C.O., Kimani-Njogu, S.W., Mbapila, J.C., Sallam, M.N.S. and Ofomata, V.C. 1997. A review of the introduction and establishment of *Cotesia xavipes* Cameron (Hymenoptera: Braconidae) in East Africa for biological control of cereal stemborers. *Insect Science and its Application* 17, 79–95.
- Panis, A. 1969. Observations faunistiques et biologiques sur quelques Pseudococcidae (Homoptera, Coccoidea) vivant dans le midi de la France. *Annales de Zoologie Ecologie Animale* 1: 211–244.
- Pascal, J.P. and Ramesh, B.R. 1987, 1997. A field key to the trees and lianas of the evergreen forests of the Western Ghats (India). Institut Français de Pondicherry, Sri Aurobindo Ashram Press, Pondicherry, India. 236 pp. [Describes 502 species, including 39 lianas. Illustrations for 400 species. Most of the illustrations are excellent leaf prints obtained by placing a thin paper on dry leaves and darkening them with a pencil. Glossary.]
- Persad, A. and Khan, A. 2007. Effects of four host plants on biological parameters of *Maconellicoccus hirsutus* green (Homoptera: Pseudococcidae) and efficacy of *Anagyrus kamali* Moursi (Hymenoptera: Encyrtidae). *Journal of Plant Protection Research* 47, 35-42.
- Peterson, A.T. 2003. Predicting the geography of species' invasions via ecological niche modeling. *Quarterly Review Biology* 78, 419–433.
- Peterson, A.T. 2005. Predicting potential geographic distributions of invading species. *Current Science* 89, 9.
- Peterson, A.T. and Nakazawa, Y. 2008. Environmental data sets matter in ecological niche modeling: an example with *Solenopsis invicta* and *Solenopsis richteri*. *Global Ecology and Biogeography* 17, 135–144.

- Peterson, A.T. and Shaw, J.J. 2003. Lutzomyia vectors for cutaneous leishmaniasis in southern Brazil: ecological niche models, predicted geographic distributions, and climate change effects. *International Journal of Parasitology* 33: 919-931.
- Peterson, A.T. and Vieglais, D.A. 2001. Predicting species invasions using ecological niche modeling. *BioScience* 51, 363– 371.
- Peterson, A.T., Papes, M. and Eaton, M. 2007. Transferability and model evaluation in ecological niche modeling: a comparison of GARP and Maxent. *Ecography* 30, 550–560.
- Peterson, A.T., Papes, M. and Soberon, J. 2008. Rethinking receiver operating characteristic analysis applications in ecological niche modeling. *Ecological Modeling* 213, 63–72.
- Peterson, A.T., Soberon, J. and Sanchez-Cordero, V. 1999. Conservatism of ecological niches in evolutionary time. *Science* 285, 1265–1267.
- Phillips, S.J., Anderson, R.P. and Schapire, R.E. 2006. Maximum entropy modeling of species geographic distributions. *Ecological Modeling* 190, 231–259.
- Pierce, N.E. and Mead, P.S. 1981. Parasitoids as selective agents in the symbiosis between Lycaenid butterfly larvae and ants. *Science* 211, 1185–1187.
- Pieterse, W., Muller, D.L. and Jansen van Vuuren, B. 2010. A Molecular Identification Approach for Five Species of Mealybug (Hemiptera: Pseudococcidae) on Citrus Fruit Exported from South Africa. *African Entomology*, 18(1): 23-28.
- Pillai, S.R.M., Gopi, K.C., Salarkhan, A.M. and Pankajam, S. 1991. Insect pest problems of *Albizia lebbek* (L.) Benth. in nurseries and young plantations. *Indian Journal of Forestry* 14(4), 253-260.
- Pitan, O.O., T.A. Akinlosotu, and Odebiyi, J.A. 2000. Impact of *Gyranusoidea tebygi* Noyes (Hymenoptera: Encyrtidae) on the mango mealybug, *Rastrococcus invadens* (Homoptera: Pseudococcidae) in Nigeria. *Bioccontrol Science and Technology* 10: 245–254.
- Pitan, O.O.R., Mwansat, G., Akinyemi, S.O.S., Adebayo, O.S. and Akinlosotu, T.A. 2002. Effect of mango mealybug and sooty mould attack on mango and the impact of the released *Gyranusoidea tebygi* Noyes on yield. *Fruits (Paris)* 57, 105–113.
- Powell, W. and Pickett, J.A. 2003. Manipulation of parasitoids for aphid pest management: progress and prospects. *Pest Management Science* 59: 149–155.
- Pramanik, A. and Ghose, S.K. 1991. Observations on the biology of the mango mealybug, *Rastrococcus iceryoides* (Green) (Pseudococcidae, Homoptera). *Annals of Entomology*, 9(1): 25-30.
- Prasad, K., Divakar, B.N., Hegde, N.K. and Ganigara, B.S. 1998. Nature of damage and efficacy of insecticides against mealybug, *Ferrisia virgata* (Ckll.) on black pepper cuttings. *Pest Management In Horticultural Ecosystems*, 4:52–53
- Price, P.W., Bouton, C.E., Gross, P., McPheron, B.A., Thompson, J.N., and Weis, A.E. 1980. Interactions among three trophic levels: Influence of plants on interactions between insect herbivores and natural enemies. *Annual Review of Ecology and Systematics* 11, 41–65.

- Rae, D.J. and Jones, R.E. 1992. Influence of host nitrogen levels on development, survival, size and population dynamics of sugarcane mealybug, *Saccharicoccus saccari* (Cockerell) (Homoptera: Pseudococcidae). Australian Journal of Zoology, 40: 327-342.
- Raj A.X.S. 2008. An economic analysis of mango pulp agro based industry in Krishnagiri District, Tamil Nadu. Political Economy Journal of India Jan-June 2008.
- Ramachandran, S. and Ayyar, T.V.R. 1934. Host plant index of Indo-Ceylonese coccidae. Imperial Council Agricultural Research Miscellaneous Bulletin No. 4.
- Ramadan, M.M., Wong, T.T.Y. and Messing, R.H. 1995. Reproductive Biology of *Biosteres vandenboschi* (Hymenoptera: Braconidae), a parasitoid of early-instar oriental fruit fly. Annals of the Entomological Society of America 88, 189-195.
- Ranjan, R. 2006. Economic impacts of pink hibiscus mealybug in Florida and the United States. Stochastic Environmental Research and Risk Assessment, 20: 353–362.
- Rawat, R.R. and Jakhmola, S.S. 1970. Bionomics of the mango-coccid (*Rastrococcus iceryoides* Green; Homoptera, Coccidae). Indian Journal of Agricultural Sciences, 40: 140-144.
- Raxworthy, C.J., Martinez-Meyer, E., Horning, N., Nussbaum, R.A., Schneider, G.E., Ortega-Huerta, M.A. and Peterson, A.T. 2003. Predicting distributions of known and unknown reptile species in Madagascar. Nature 426, 837–841.
- Rice, N., Martinez-Meyer, E. and Peterson, A.T. 2003. Ecological niche differentiation in the phelocoma jays: a phylogenetic perspective. Biological Journal of the Linnean Society 80, 369–383.
- Richardson, D.M. and van Wilgen, B.M. 2004. Invasive alien plants in South Africa: how well do we understand the ecological impacts? South African Journal of Science 100, 45–52.
- Roltsch, W.J., Meyerdirk, D.E., Warkentin, R., Andress, E.R. and Carrera, K. 2006. Classical biological control of the pink hibiscus mealybug, *Maconellicoccus hirsutus* (Green), in southern California. Biological Control 37: 155–166.
- Rosen, D. 1981. A new species of *Pseudaphycus* (Hym, Encyrtidae), with notes on the *Angelicus* group. Entomophaga 26:251–263.
- Rotundo, G. and Tremblay, E. 1975. Sull' attractività delle femmine vergini di due specie de Pseudococcidi (Homoptera: Coccoidea) per un Imenottero parassita (HymenopteraChalcidoidea). Bollettino del Laboratorio di Entomologia Agraria Filippo Silvestri, Portici 32: 172–179.
- RTA. 2008. Fruits and Vegetables (East and Southern Africa). Round Table Africa Presentation MSM. Published on the internet; <http://www.roundtableafrica.net/File/presentations> [accessed 18 February 2011].
- Ruberson, J.R., Tauber, M.J. and Tauber, C.A. 1989. Intraspecific variability in hymenopteran parasitoids: comparative studies of two biotypes of the egg parasitoid *Edovum puttleri* (Hymenoptera: Eulophidae). Journal of the Kansas Entomological Society 62, 189-202.

- Sadof, C.S., Neal, J.J. and Cloyd, R.A. 2003. Effect of variegation on stem exudates of coleus and life history characteristics of citrus mealybug (Hemiptera: Pseudococcidae). *Environmental Entomology* 32: 463–469.
- Sagarra, L. A, and Vincent, C. 1999. Influence of host stage on oviposition, development, sex ratio and survival of *Anagyrus kamali* Moursi (Hymenoptera: Encyrtidae), a parasitoid of the hibiscus mealybug *Maconellicoccus hirsutus* Green (Homoptera: Pseudococcidae). *Biological Control*. 15:51–56.
- Sagarra, L.A. and Peterkin, D.D. 1999. "Invasion of the Caribbean by the hibiscus mealybug, *Maconellicoccus hirsutus* Green [Homoptera: Pseudococcidae]", *Phytoprotection*, 80(2): 103-113.
- Sagarra, L.A., Peterkin, D.D., Vincent, C. and Stewart, R.K. 2000. Immune response of the hibiscus mealybug, *Maconellicoccus hirsutus* Green (Homoptera: Pseudococcidae), to oviposition of the parasitoid *Anagyrus kamali* Moursi (Hymenoptera: Encyrtidae). *Journal of Insect Physiology* 46: 647–653.
- Sagarra, L.A., Vincent, C. and Stewart, R.K. 2001. Body size as an indicator of parasitoid quality in male and female *Anagyrus kamali* (Hymenoptera: Encyrtidae). *Bulletin of Entomological Research* 91, 363-367.
- Sagarra, L.A., Vincent, C. and Stewart, R.K. 2001. Suitability of nine mealybug species (Homoptera: Pseudococcidae) as hosts for the parasitoid *Anagyrus kamali* (Hymenoptera: Encyrtidae). *Florida Entomologist* 84: 112–116.
- Sahoo, S.K. and Ghosh, A.B. 2000. Biology of the mealybug *Rastrococcus invadens* Williams (Pseudococcidae: Hemiptera). *Journal of Environment and Ecology* 18: 752– 756.
- Salt, G. 1940. Experimental studies in insect parasitism. VII. The effects of different hosts on the parasite *Trichogramma evanescens* westw. (Hymenoptera: Chalcidoidea). *Proceeding of the Royal society of London, Series A General Entomology* 15: 81-95.
- Samways, M.J. and Grech, N.M. 1986. Assessment of the fungus *Cladosporium oxysporum* (Berk and Curt) as a potential biocontrol agent against certain homoptera. *Agriculture, Ecosystems and Environment* 15: 231–239.
- Samways, M.J. and Tate, B.A. 1984. Evaluation of several trunk barriers used to prevent the movement of the pugnacious ant (*Anoplolepis custodiens* (Smith)) into citrus trees. *Citrus and Subtropical Fruit Journal* 608, 9–12.
- Samways, M.J., Nel, M. and Prins, A.J. 1982. Ants (Hymenoptera: Formicidae) foraging in citrus trees and attending honey-producing Homoptera. *Phytophylactica* 14:155–157
- Sánchez, L. 2008. Sex-determining mechanisms in insects. *International Journal of Developmental Biology* 52: 837–856.
- SAS Institute. 2010. SAS/STAT Users Guide: Statistics, version 9.1.3, SAS Institute, Cary, NC.
- Sazo, L., Pizarro, E. and Araya, J.E. 2006. Effect of the form of application of imidacloprid on control of the long-tailed mealybug *Pseudococcus longispinus* (Targioni & Tozzetti) on

- avocado and its impact on *Neoseiulus californicus* (McGregor) in Chile. Boletín de Sanidad Vegetal Plagas 32: 483–490.
- Scott, J.M., Heglund, P.J., Morrison, M.L., Haufler, J.P., Raphael, M.G., Wall, W.A., Samson, F.B. (eds). 2002. Predicting species occurrences: issues of accuracy and scale. Island Press, Washington, DC, pp 868.
- Serrano, M.S. and Lapointe, S.L. 2002. Evaluation of host plants and a meridic diet for rearing *Maconellicoccus hirsutus* (Hemiptera: Pseudococcidae) and its parasitoid *Anagyrus kamali* (Hymenoptera: Encyrtidae). Florida Entomologist 85: 417- 425.
- Sether, D.M. and Hu, J.S. 2002a. Closterovirus infection and mealybug exposure are necessary for the development of mealybug wilt of pineapple disease. Phytopathology 92: 928–935.
- Sether, D.M. and Hu, J.S. 2002b. Yield impact and spread of pineapple mealybug wilt associated virus-2 and mealybug wilt of pineapple in Hawaii. Plant Diseases 86: 867–874.
- Sether, D.M., Melzer, M.J., Bustos, J., Zee, F. and Hu, J.S. 2005. Diversity and mealybug transmissibility of ampeloviruses in pineapple. Plant Diseases 89: 450–456.
- Sforza, R. 2008a. Les cochenilles sur la vigne. In “*Les ravageurs de la vigne*” pp. 188–210. Eds Feret, Bordeaux, p. 389.
- Sforza, R. 2008b. Espèces invasives en viticulture. Phytoma-La défense des végétaux. 619: 24–29.
- Sforza, R., Boudon-Padieu, E. and Greif, C. 2003. New mealybug species vectoring Grapevine leafroll- associated viruses-1 and -3 (GLRaV-1 and-3). European Journal of Plant Pathology 109: 975–981.
- Shafqat, S., Munir, A., Mushtaq, A. and Kwon, Y.J. 2007. Insecticidal control of the mealybug *Phenacoccus gossypiphilous* (Hemiptera: Pseudococcidae), a new pest of cotton in Pakistan. Entomological Research 37: 76–80.
- Shanahan, M. and Compton, S.G. 2000. Ant Predation Suppresses Populations of the Scale Insect *Aulacaspis marina* in Natural Mangrove Forests. Biotropica 32(4a): 764-768.
- Shorey, H.H., Gaston, L.K., Gerber, R.G., Phillips, P.A., Wood, D.L. 1992. Disruption of foraging by argentine ants *Iridomyrmex humilis* (Mayr) (Hymenoptera: Formicidae), in citrus trees through the use of semiochemicals and related chemicals. Journal of Chemical Ecology, 18, 2131–2142.
- Shorey, H.H., Gaston, L.K., Gerber, R.G., Sisk, C.B. and Wood, D.L. 1993. Disruption of foraging by *Formica aerata* (Hymenoptera: Formicidae) through the use of semiochemicals and related chemicals. Environmental Entomology, 22, 920–924.
- Shree, M.P. and Boraiah, G. 1988. Incidence of turka and bacterial blight on mulberry plants in Germplasm Bank. Current Science 57 (22): 111-113.
- Shrewsbury, P.M., Bejleri, K. and Lea-Cox, J.D. 2004. Integrating cultural management practices and biological control to suppress citrus mealybug. In: proceeding of XXVI IHC

- Protected cultivation 2002: in search of structures, systems and plant Materials for sustainable greenhouse production. *Acta Horticulture*. 633, ISHS.
- Silva, E.B. and Mexia, A. 1999a. Histological studies on the stylet pathway, feeding sites and nature of feeding damage by *Planococcus citri* (Risso) (Homoptera: Pseudococcidae) in sweet orange. *Entomologica*, Bari 33:347–350.
- Silva, E.B. and Mexia, A. 1999b. The pest complex *Cryptoblabes gnidiella* (Millière) (Lepidoptera: Pyralidae) and *Planococcus citri* (Risso) (Homoptera: Pseudococcidae) on sweet orange groves (*Citrus sinensis* (L.) Osbeck) in Portugal: interspecific association. *Boletín de Sanidad Vegetal Plagas* 25: 89–98.
- Silverman, J. and Brightwell, R.J. 2008. The Argentine ant: challenges in managing an invasive unicolonial pest. *Annual Review of Entomology* 53:231–252.
- Singh, L.B. 1968. The mango. Botany, Cultivation and Utilization. World Crop Books. Leonard Hill, London.
- Sisk, C.B., Shorey,H.H.,Gerber,R.G. and Gaston, L.K. 1996. Semio- chemicals that disrupt foraging by the Argentine ant (Hymenoptera: Formicidae): laboratory bioassays. *Journal of Economic Entomology*, 89, 381–385.
- Skov, F. 2000. Potential plant distribution mapping based on climatic similarity. *Taxon* 49: 503 - 515.
- Slansky, J.E. and Rodriguez, J.G. 1987b. Nutritional ecology of insects, mites, spiders, and related invertebrates: an overview. In: Slansky Jr E & Rodriguez JG (eds), *Nutritional Ecology of Insects, Mites, Spiders, and Related Invertebrates*. John Wiley & Sons, New York, pp. 1-69.
- Smit, B. and Bishop, H.J. 1934. A study of the citrus mealybug and its association with ants in the Eastern Province. *Science Bulletin No. 125*. The Government Printer, Pretoria.
- Soberon, J. and Peterson, A.T. 2005. Interpretation of models of fundamental ecological niches and species' distributional areas. *Biodiversity Informatics* 2, 1–10.
- Sokal, R.R. and Rohlf, F.J. 1981. *Biometry: The principles and Practices of Statistics in Biological Research*, second ed. Freeman, New York, NY.
- Sopow, S.L. and Quiring, D.T. 1998. Body size of spruce-galling adelgids is positively related to realized fecundity in nature. *Ecological Entomology* 23: 467-479.
- Souissi, R. 1999. The influence of the host plant of the cassava mealybug *Phenacoccus manihoti* on the plant and host preferences of its parasitoid *Apoanagyrus lopezi*. *Biological Control* 15(1), 64-70.
- Souissi, R. and Le Ru, B.1997a. Effect of host plant on fecundity and development of *Apoanagyrus lopezi*, an endoparasitoid of the cassava mealybugs *Phenacoccus manihoti*. *Entomologia Experimentalis et Applicata* 82: 235–238.

- Souissi, R. and Le Rü, B. 1997a. Effect of host plant on fecundity and development of *Apoanagyrus lopezi*, an endoparasitoid of the cassava mealybugs *Phenacoccus manihoti*. *Entomologia Experimentalis et Applicata* 82, 235–238.
- Souissi, R. and Le Rü, B. 1997b. Influence of host plant on the ecobiological characteristics of *Epidinocarsis lopezi* (Hymenoptera: Encyrtidae), parasitoid of the cassava mealybug. *Bulletin of Entomological Research* 87, 1-7.
- Souissi, R. and Le Rü, B. 1998a. Influence of host plant of the cassava mealybug *Phenacoccus manihoti* (Hemiptera: Pseudococcidae) on biological characteristics of its parasitoid *Apoanagyrus lopezi* (Hymenoptera: Encyrtidae). *Bulletin of Entomological Research* 87, 1–7.
- Souissi, R., Ne'non, J.P. and Le Ru, B. 1998b. Olfactory responses of the parasitoid *Apoanagyrus lopezi* (Hymenoptera: Encyrtidae) to odor of plants, mealybugs, and plant-mealybug complexes. *Journal of Chemical Ecology* 1, 37–48.
- Sriharan, T.P., Samsom, M.V. and Krishnaswami, S. 1979. Studies on tukra disease of mulberry. *India journal of Sericulture* 18: 78.
- Stechmann DH, Volk W and Stary' P. 1996. Ant-attendance as a critical factor in the biological control of the banana aphid *Pentalonia nigronervosa* Coq. (Hom. Aphididae) in Oceania. *Journal of Applied Entomology* 120:119–123.
- Steyn, J.J. 1954. The pugnacious ant, *Anoplolepis custodiens* (Smith), and its relation to the control of the citrus scale at Letaba. *Memoirs of the Entomological Society of South Africa* No. 3. Pretoria.
- Stiling, P. 1990. Calculation the establishment rates of parasitoids in classical biological control. *American Entomologist* 36:225–230.
- Stiling, P. 1993. Why do natural enemies fail in classical biological control programs? *American Entomologist* 39:31–37.
- Stockwell, D.R.B. and Noble, I.R. 1992. Induction of sets of rules from animal distribution data: a robust and informative method of analysis. *Mathematics and Computers in Simulation* 33: 385 - 390.
- Stockwell, D.R.B. and Peterson, A.T. 2002. Effects of sample size on accuracy of species distribution models. *Ecological Modeling* 148, 1–13.
- Streito, J.C. and Martinez, M. 2005. Nouveaux ravageurs, 41 espèces depuis 2000. Phytoma, La défense des végétaux. 586: 16–20.
- Strickland, A.H. 1950. The entomology of swollen shoot of cacao. I.—The insect species involved, with notes on their biology. *Bulletin of Entomological Research*, 41: 725-748.
- Stuart, R.J., Polavarapu, S., Lewis, E.E. and Gaugler, R. 1997. Differential susceptibility of *Dysmicoccus vaccinii* (Homoptera: Pseudococcidae) to entomopathogenic nematodes (Rhabditida: Heterorhabditidae and Steinernematidae) *Journal of Economic Entomology* 90: 925–932.

- Sugonyayev, E.S. 1995. Ants nesting on living plants in the tropics as refuges for soft-scale insects (Homoptera, Coccoidea, protecting them from attacks of chalcidoid parasites (Hymenoptera, Chalcidoidea). Entomological Review, 75: 120-127.
- Sullivan, D.J. 1987. Insect hyperparasitism. Annual Review of Entomology, 32: 49-70.
- Suresh, S., Kavitha, P.C. 2008. Seasonal incidence of economically important coccid pests in Tamil Nadu. Branco M, Franco JC & Hodgson C (eds) (2008) Proceedings of XI International Symposium on Scale Insect Studies, Oeiras (Portugal), 24-27 September 2007. ISA Press, Lisbon, 322 p.
- Sutherst, R.W. 2003. Prediction of species geographical ranges. Journal of Biogeography 30:805–816.
- Swirski, E., Izhar, Y., Wysoki, M., Gurevitz, E. and Greenberg, S. 1980. Integrated control of the long-tailed mealybug, *Pseudococcus longispinus* (Homoptera: Pseudococcidae), in avocado plantations in Israel. Entomophaga 25: 415–426.
- Takabayashi, J., M. Dicke and Posthumus, M.A. 1991. Variation in composition of predator-attracting allelochemicals emitted by herbivore-infested plants: relative influence of plant and herbivore. Chemoecology 2: 1–6.
- Takahashi, S., M. Hajika, J. Takabayashi and Fukui, M. 1990. Oviposition stimulants in the coccoid cuticular waxes of *Aphytis yanonensis* De Back & Rosen. Journal of Chemical Ecology 16: 1657–1665.
- Tandon, P.L. and Lal, B. 1978. The mango coccid, *Rastrococcus iceryoides* Green (Homoptera: Coccoidea) and its natural enemies. Current Science 47(13):467- 468.
- Tandon, P.L. and Srivastava, R.P. 1980. New records of parasites and predators of important insect pests of mango. Entomology 5(3), 243-244.
- Terra, W.R. and Ferreira, C. 2003. Digestive system. In: Resh VH, Cardé RT (Eds) Encyclopedia of insects. Academic Press, Amsterdam.
- Tertuliano, M. and Lero, B. 1992. Interaction between cassava mealybugs (*Phenacoccus manihoti*) and their host plants – amino-acid and sugar contents of sap. Entomologia Experimentalis et Applicata 64: 1–9.
- Tertuliano, M., Dossougbe, S. and Lero, B. 1993. Antixenotic and antibiotic components of resistance to the cassava mealybug *Phenacoccus manihoti* (Homoptera: Pseudococcidae) in various host- plants. Insect Science and its Application 14: 657–665.
- Tharanathan, R.N., Yashoda, H.M. and Prabha, T.N. 2006 Mango (*Mangifera indica* L.), “The king of fruits”—An overview. Food Reviews International 22: 95–123.
- Tho, Y.P. 1978. Living in harmony. Nature Malaysiana, 3: 34-39.
- Thomson, K.G., Dietzgen, R.G., Thomas, J.E. and Teakle, D.S. 1996. Detection of pineapple bacilliform virus using the polymerase chain reaction. Annals of Applied Biology 129: 57–69.

- Tingle, C.C.D., and Copland, M.J.W. 1989. Progeny production and adult longevity of the mealybug parasitoids *Anagyrus pseudococci*, *Leptomastix dactylopii*, and *Leptomastidea abnormis* [Hymenoptera: Encyrtidae] in relation to temperature. *Entomophaga* 34: 111 - 120.
- Tobih, F.O., Omoloye, A.A., Ivbijaro, M.F. and Enobakhare, D.A. 2002. Effects of field infestation by *Rastrococcus invadens* Williams (Hemiptera: Pseudococcidae) on the morphology and nutritional status of mango fruits, *Mangifera indica* L. *Crop Protection* 21, 757–761.
- Tonkyn, D.W. and Whitcomb, R.F. 1987. Feeding strategies and the guild concept among vascular feeding insects and microorganisms. In: Harris KF (Ed) Current topics in vector research, Vol. 4. Springer-Verlag, New York.
- Tsai, C., Chan, J., Fernandez, L., Bosco, D., Daane, K.M. and Rodrigo, R.P. 2008. Transmission of grapevine leafroll-associated virus 3 by the vine mealybug (*Planococcus ficus*). *Phytopathology* 98: S159.
- Tuda, M., Matsumoto, T., Itioka, T., Ishida, N., Takanashi, M., Ashihara, W., Kohyama, M. and Takagi, M. 2006. Climatic and intertrophic effects detected in 10-year population dynamics of biological control of the arrowhead scale by two parasitoids in southwestern Japan. *Population Ecology* 48:59–70.
- Turlings, T.C.J., Tumlinson, J.H., Lewis, W.J. 1991. Larval damaged plants: Sources of volatile synomones that guide the parasitoid *Cotesia marginiventris* to the micro-habitat of its hosts. *Entomologia Experimentalis et Applicata* 58, 75–82.
- Ukwela, M.U. 2009. Influence of hyperparasitoids on the exotic parasitoid *Gyranusoidea tetygi noyes* (Hymenoptera: Encyrtidae) of mango mealybug on mango varieties in guinea savanna ecology. *African Journal of Science and Technology (AJST)*, 10 (1): 16 – 23.
- Unwin, D.M. and Corbet, S.A. 1991. Insects, Plants and microclimate. Richmond Publishing, Surrey.
- Vacek, D.C. 1982. Interactions between microorganisms and Catophilic *Drosophila* in Australia, p.175-190. In J.S.F. Barker and W.T. Starmer (ed.), Ecological genetics and evolution. The cactus-yeast-Drosophila model system. Academic Press, Inc., Sydney, Australia.
- Van Alphen, J.J.M. and M.A. Jervis. 1996. Foraging behaviour, Pp. 1-62. In M.A. Jervis and N.A.C. Kidd [eds.], Insect natural enemies: practical approaches to their study and evaluation. Chapman and Hall, London, UK.
- van Alphen, J.J.M., Vet, L.E.M. 1986. An evolutionary approach to host finding and selection. In Insect Parasitoids 13th Symposium of the Royal Entomological Society of London, 18-19 September 1985, London (ed. J. K. Waage and D. J. Greathead), pp. 23-61. London: Academic Press.
- van Lenteren, J.C. 1986. Evaluation, mass production, quality control and release of entomophagous insects. In: J.M. Franz (ed), Biological Plant and Health Protection. Series Progress in Zoology 32, Fischer, Stuttgart. pp. 31–56.

- Van Lenteren, J.C. 2006. How not to evaluate augmentative biological control. *Biological Control* 39: 115–118.
- van Lenteren, J.C. and Noldus, L.P.J.J. 1990. Whitefly–plant relationships: behavioural and ecological aspects. In: Gerling D (ed) *White-flies: bionomics pest status and management*. Intercept Ltd, Andover, Hants, pp. 47–90
- van Lenteren, J.C., Bigler, F., Burgio, G., Hokkanen, H.M.T., Thomas, M.B. 2002. Risks of importation and release of exotic biological control agents: how to determine host specificity? *IOBC/ WPRS Bulletin*, 25 (1), 281-284.
- Van Mele, P., Vayssières, J.-F., Van Tellingen, E. and Vrolijks, J. 2007. Effects of an African weaver ant, *Oecophylla longinoda*, in controlling mango fruit flies (Diptera: Tephritidae) in Benin. *Journal of Economic Entomology* 100: 695-701.
- Van Mele, P., Vayssières, J.-F., Van Tellingen, E., Vrolijks, J. 2007. Effects of an African weaver ant, *Oecophylla longinoda*, in controlling mango fruit flies (Diptera: Tephritidae) in Benin. *Journal of Economic Entomology* 100, 695-701.
- Van Nouhuys, S., and Hanski, I. 2000. Apparent competition between parasitoids mediated by a shared hyperparasitoid. *Ecology Letters*. 3, 82–84.
- Van Veen, F.J.F., Rajkumar, A., Müller, C.B. and Godfray, H.C.J. 2001. Increased reproduction by pea aphids in the presence of secondary parasitoids. *Ecological Entomology* 26, 425–429.
- Varela, AM, Seif, A and Nyambo, B. 2006. A guide to IPM in mango production in Kenya. The International Centre of Insect Physiology and Ecology. Second edition. ISBN 92 9064 17-6 2.
- Vargas, R.I., Chang, H.B.C., Komura, M. and Kawamoto, D. 1987. Mortality, stadal duration, and weight loss in three species of mass-reared fruit fly pupae (Diptera: Tephritidae) held with and without vermiculite at selected relative humidities. *Journal of Economic Entomology* 80, 972– 974.
- Vasundhara, M., Siddappaji, C. and Kotikal, Y.K. 1990. New record of downey snow line mealybug on *Jasminum rigidum* and its predators. *Current Research - University of Agricultural Sciences (Bangalore)*, 19(6): 99-100.
- Vera, M.T., Rodriguez, R., Segura, D.F., Cladera, J.L. and Sutherst, R.W. 2002. Potential geographical distribution of the Mediterranean fruit fly, *Ceratitis capitata* (Diptera: Tephritidae), with emphasis on Argentina and Australia. *Environmental Entomology* 31, 1009–1022.
- Vet, L.E.M. and Dicke, M. 1992. The ecology of infochemical use by natural enemies of herbivores in a tritrophic context. *Annual Review of Entomology* 37, 141–172.
- Vinson, S.B. 1976. Host location by insect parasitoids. *Annual Review of Entomology* 21, 109–134.
- Vinson, S.B. and Iwantsch, G.F. 1980. Host regulation by insect parasitoids. *Quarterly Review of Biology* 55, 143 – 165.

- Vinson, S.B. and Scarborough, T.A. 1991. Interaction between *Solenopsis invicta* (Hymenoptera: Formicidae), *Rhopalosiphum maidis* (Homoptera: Aphididae) and the parasitoid *Lysiphlebus testaceipes* Cresson (Hymenoptera: Aphidiidae). Annals of the Entomological Society of America, 84, 158 – 164.
- Vinson, S.B. and Williams, H.J. 1991. Host selection behavior of *Campoplexis sonorensis*: A model system. Biological Control 1, 107–117.
- Vitullo, J., Wang, S.F., Zhang, A.J., Mannion, C. and Bergh, J.C. 2007. Comparison of sex pheromone traps for monitoring pink hibiscus mealybug (Hemiptera: Pseudococcidae). Journal Economic Entomology 100: 405–410.
- Vögele, J.M., Agounké, D. and Moore, D. 1991. Biological control of the fruit tree mealybug *Rastrococcus invadens* Williams in Togo: a preliminary sociological and economic evaluation. Tropical Pest Management 37: 379–382.
- Völkl, W. 1995. Behavioral and morphological adaptations of the coccinellid, *Platynaspis luteorubra*, for exploiting ant-attended resources (Coleoptera: Coccinellidae). Journal of Insect Behaviour 8, 653-670.
- Völkl, W. and Mackauer, M. 2000. Oviposition behaviour of aphidiine wasps (Hymenoptera: Braconidae, Aphidiinae): morphological adaptations and evolutionary trends. The Canadian Entomologist 132: 197-212.
- Waage, J.K. 1986. Family planning in parasitoids: adaptive patterns of progeny sex allocation. pp. 63–95 in Waage, J. & Greathead, D. (Eds) Insect parasitoids. London, Academic Press.
- Waage, J.K. and Ng, S.M. 1984. The reproductive strategy of a parasitic wasp. 1. Optimal progeny and sex allocation in *Trichogramma evanescence*. Journal of Animal Ecology 53, 401–416.
- Walker, J.T.S., Charles, J.G., Froud, K.J. and Connolly, P. 2004. Leafroll virus in vineyards: modelling the spread and economic impact. Horticulture and Food Research Institute of New Zealand Ltd, Mt Albert, NZ.
- Walton, V.M. and Pringle, K.L. 1999. Effects of pesticides used on table grapes on the mealybug parasitoid *Coccidoxyenoides peregrinus* (Timberlake) (Hymenoptera: Encyrtidae). South African Journal of Enology and Viticulture 20: 31-34.
- Walton, V.M. and Pringle, K.L. 2005. Developmental biology of vine mealybug, *Planococcus ficus* (Signoret) (Homoptera: Pseudococcidae), and its parasitoid *Coccidoxyenoides perminutus* (Timberlake) (Hymenoptera: Encyrtidae). African Entomology 13: 143–147.
- Walton, V.M., Daane, K.M. and Pringle, K.L. 2004. Monitoring *Planococcus ficus* in South African vine-yards with sex pheromone-baited traps. Crop Protection 23: 1089–1096.
- Walton, V.M., Daane, K.M., Bentley, W.J., Millar, J.G., Larsen, T.E. and Malakar-Kuenen, R. 2006. Pheromone-based mating disruption of *Planococcus ficus* (Hemiptera: Pseudococcidae) in California vineyards. Journal of Economic Entomology 99: 1280–1290.
- Wang, X.G. and Keller, M.A. 2002. A comparisons of the host-searching efficiency of two larval parasitoids of *Plutella xylostella*. Ecological Entomology 27: 105-114.

- Ward, P.S. 1991. Phylogenetic analysis of pseudomyrmecine ants associated with domatia-bearing plants. In: C.R. Huxley and D.F. Cutler (Editors), *Ant-Plant Interactions*. Oxford University Press, Oxford, pp. 335-352.
- Waterhouse, D.F. 1993. Biological control: Pacific prospects-Supplement 2. ACIAR monograph no. 20. 138 p.
- Watson, G.W. and Kubiriba, J. 2005. Identification of mealybugs (Hemiptera: Pseudococcidae) on banana and plantain in Africa. *African Entomology* 13: 35–47.
- Watson, G.W. and Kubiriba, J. 2005. Identification of mealybugs (Hemiptera: Pseudococcidae) on banana and plantain in Africa. *African Entomologist* 13 (1): 35-47.
- Way, M.J. 1954. Studies on the association of the ant *Oecophylla longinoda* (Latr.) (Formicidae) with the scale insect *Saissetia zanzibarensis* Williams (Coccidae). *Bulletin of Entomological Research* 45, 113–134.
- Way, M.J. 1954b. Studies on the association of the ant *Oecophylla longinoda* (Latr.) (Formicidae) with the scale insect *Saissetia zanzibarensis* Williams (Coccidae). *Bulletin of Entomological Research*, 45: 113-134.
- Way, M.J. 1963. Mutualism between ants and honeydew-producing Homoptera. *Annual Review of Entomology*, 8: 307-344.
- Way, M.J. and Khoo, K.C. 1992. Role of ants in pest management. *Annual Review of Entomology*, 37: 479-503.
- Weeds of National Significance. 2011. Weed Management Guide. Department of the Environment and Heritage and the CRC for Australian Weed Management. <http://www.deh.gov.au/biodiversity/invasive/publications/p-aculeata.html> (accessed February 01th, 2011).
- Welk, E., Schubert, K. and Hoffmann, M.H. 2002. Present and potential distribution of invasive mustard (*Alliaria petiolata*) in North America. *Divers. Distributions* 8, 219–233.
- Wessels and Wolfgang. 1997. An Ever Closer Fusion? A Dynamic Macropolitical View on Integration Processes, in: *Journal of Common Market Studies*, 35(2): 267-299.
- Wharton, T.N. and Kriticos, D.J. 2004. The fundamental and realized niche of the Monterey pine aphid, *Essigella californica* (Essig) (Hemiptera: Aphididae): implications for managing softwood plantations in Australia. *Diversity and Distributions* 10, 253–262.
- Wheeler, W.M. 1910. Ants. Their structure, Development and Behaviour. Columbia University Press, New York and London, 663 pp.
- Whiley, A.W. 1993. Environmental effects on phenology and physiology of mango-a review. *Acta Horticulturae* 341, 168-176.
- Whitehead, V.B. 1957. A study of the predators and parasites of *Planococcus citri* (Risso) (Homoptera: Pseudococcidae) on vines in the Western Cape Province, South Africa. MSc Thesis, Rhodes University, Grahamstown.

- Wiens, J.J. and Graham, C.H. 2005. Niche conservatism: integrating evolution, ecology, and conservation biology. *Annual Review of Ecology, Systematics and Evolution* 36, 519–539.
- Williams, D.J. 1986. *Rastrococcus invadens* sp. n. (Hemiptera : Pseudococcidae) introduced from the Oriental Region to West Africa and causing damage to mango, citrus and other trees. *Bulletin of entomological Research* 76: 695-699.
- Williams, D.J. 1989. The mealybug genus *Rastrococcus* Ferris (Hemiptera: Pseudococcidae). *Journal of Systematic Entomology* 14(4): 433-486.
- Williams, D.J. 2004. Mealybugs of southern Asia. The Natural History Museum, London, UK, Southdene SDN. BHD, Kuala Lumpur, Malaysia, 896 pp.
- Williams, D.J. and Granara de Willink, M.C. 1992. Mealybugs of Central and South America. CABI, Wallingford.
- Willink, E. and Moore, D. 1988. Aspects of the biology of *Rastrococcus invadens* Williams (Hemiptera: Pseudococcidae), a pest of fruit crops in West Africa, and one of its primary parasitoids, *Gyranusoidea tebygi* Noyes (Hymenoptera: Encyrtidae). *Bulletin of Entomological Research* 78: 709–715.
- Winotai, A., G. Goergen, M. Tamò, and Neuenschwander, P. 2010. Cassava mealybug has reached Asia. *Biocontrol News and Information*, 31: 10N-11N.
- Wysoki, M., Izhar, Y., Swirski, E., Gurevitz, E. and Greenberg, S. 1977. Susceptibility of avocado varieties to long-tailed mealybug, *Pseudococcus longispinus* (Targioni Tozzetti) (Homoptera: Pseudococcidae), and a survey of its host plants in Israel. *Phytoparasitica* 5: 140–148.
- Yang, J. and Sadof, C.S. 1995. Variegation in *Coleus blumei* and the life history of citrus mealybug (Homoptera: Pseudococcidae). *Environmental Entomology* 24: 1650 -1655.
- Yang, J.S. and Sadof, C.S. 1997. Variation in the life history of the citrus mealybug parasitoid *Leptomastix dactylopii* (Hymenoptera: Encyrtidae) on three varieties of *Coleus blumei*. *Environmental Entomology* 26: 978–982.
- Yaninek, J.S., Onzo, A. and Ojo, B. 1993. Continent-wide releases of neotropical phytoseiids against the exotic cassava green mite in Africa. *Experimentalis and Applied Acarology*. 17: 145-160.
- Yom-Tov, Y. and Kadmon, R. 1998. Analysis of the distribution of insectivorous bats in Israel. *Divers. Distributions* 4, 63–70.
- Yusuf, S.A. and Salau, A.S. 2007. Forecasting Mango and Citrus Production in Nigeria: A Trend Analysis MPRA Paper No. 2691, Munich Personal RePEc Archive. Available on line at <http://mpra.ub.uni-muenchen.de/2691/>.
- Zada, A., Dunkelblum, E., Assael, F., Franco, J.C., Silva, E.B., Protasov, A. and Mendel, Z. 2008. Attraction of *Planococcus ficus* males to racemic and chiral pheromone baits: flight activity and bait longevity. *Journal of Applied Entomology* 132:480–489

- Zaka, S.M., S. Saeed, S.A. Bukhari, and E. Baksh. 2006. Mealybug, *Phenacoccus solenopsis* (Homoptera: Pseudococcidae): a novel pest of cotton in Pakistan. Proceedings of 34th Pakistan(SAARC) Countries Science conference, p. 32. University of Veterinary and Animal Sciences, Lahore, Pakistan.
- Zalba, S.M., Sonaglioni, M.I., Compagnoni, C.A. and Belenguer, C.J. 2000. Using a habitat model to assess the risk of invasion by an exotic plant. *Biological Conservation* 93: 203 - 208.
- Zaman, Z. and Maiti, B. 1994. Insects and mites infesting seedlings of mango in West Bengal. *Environmental Ecology* 12(3), 734-736.
- Zar, J.H. 2009. *Biostatistical Analysis*, 4th ed.; Prentice Hall: Upper Saddle River, NJ, USA.
- Zeddies, J., Schaab, R.P., Neuenschwander, P. and Herren, H.R. 2001. Economics of biological control of cassava mealybug in Africa. *Agricultural Economics* 24: 209–219.
- Zhang, A.J., Wang, S.F., Vitullo, J., Roda, A., Mannion, C. and Bergh, J.C. 2006. Olfactory discrimination among sex pheromone stereoisomers: Chirality recognition by pink hibiscus mealybug males. *Chemical Senses* 31: 621–626.
- Zorloni, A., Prati, S., Bianco, P.A. and Belli, G. 2006. Transmission of Grapevine virus A and Grapevine leafroll-associated virus 3 by *Heliothrips bohemicus*. *Journal of Plant Pathology* 88: 325–328.