

References

- Abe, M., Abe, K., Kuroda, M., and Arai, S. (1992) Corn kernel cystein protease inhibitor as a novel cystatin superfamily member of plant origin. *Eur. J. Biochem.*, 209: 933-937.
- Abe, M., Abe, K., Domoto, C., and Arai, S. (1995) Two distinct species of corn cystatin in corn kernels. *Biosci. Biotech. Biochem.*, 59: 756-758.
- Abe, M., Domoto, C., Watanabe, H., Abe, K. and Arai, S. (1996) Structural organization of the gene encoding corn cystatin. *Biosci. Biotech. Biochem.*, 60: 1173-1175.
- Adam, Z. and Clarke, A.K. (2002) Cutting edge of chloroplast proteolysis. *Trends Plant Sci.*, 7: 451-456.
- Ainsworth, E.A., Davey, P.A., Bernacchi, C.J., Dermody, O.C., Heaton, E.A., Moore, D.J., Morgan, P.B., Naidu, S.L., Yoo Ra, H-S., Zhu, X-G., Peters, C., Long, S.P. (2002) A meta-analysis of elevated [CO₂] effects on soybean (*Glycine max*) physiology, growth and yield. *Glob. Change Biol.*, 8: 695-709.
- Ainsworth, E.A. and Long, S.P. (2005) What have we learned from 15 years of free-air CO₂ enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO₂. *New Phytol.*, 165: 351-371.
- Akbar, M., Khush, G.S., and Hillerislambers, D. (1986) Genetics of salt tolerance in rice, in: Rice Genetics, International Rice Research Institute, Manila, Philippines, pp. 399-409.
- Allen, D.J. and Ort, D.R. (2001) Impacts of chilling temperatures on photosynthesis in warm-climate plants. *Trends Plant Sci.*, 6: 36-42.
- Alonso, J.M. and Granell, A. (1995) A putative vacuolar processing protease is regulated by ethylene and also during fruit ripening in *citrus* fruit. *Plant Physiol.*, 109: 541-547.

Altpeter, F.; Diaz, I.; McAuslane, H.; Gaddour, K.; Carbonero, P. and Vasil, I.K. (1999). Increased insect resistance in transgenic wheat stably expressing trypsin inhibitor CMe. Mol. Breeding, 5:53-63.

Altschul, S.F., Gish, W., Miller, W., Myers, E.W., Lipman, D.J. (1990) Basic local alignment search tool. J. Mol. Biol., 215: 403-410

Amme, S., Rutten, T., Melzer, M., Sonsmann, G., Vissers, J.P.C., Schlesier, B., Mock, H-P. (2005) A proteome approach defines protective functions of tobacco leaf trichomes. Proteomics, 5: 2508-2518.

Anderson, L.J., Maherli, H., Johnson, H.B., Polley, H.W., Jackson, R.B. (2001) Gas exchange and photosynthetic acclimation over sub-ambient to elevated CO₂ in a C₃-C₄ grassland. Glob. Change Biol., 7: 693–707.

Antão, C.M. and Malcata, F.X. (2005) Plant serine proteases: biochemical, physiological and molecular features. Plant Physiol. Bioch., 43: 637-650.

Aono, M., Saji, H., Sakamoto, A., Tanaka, K., Kondo, N., Tanaka, K. (1995) Paraquat tolerance of transgenic *Nicotiana tabacum* with enhanced activities of glutathione reductase and superoxide dismutase. Plant Cell Physiol., 36: 1687-1691.

Arai, S., Abe, K. (2000) Cystatin-based control of insects with special reference to the efficacy of oryzacystatin. In *Recombinant Protease Inhibitors in Plants*; Michaud, D., Ed.; Landes: Georgetown, TX; pp. 27-42.

Aravind, L. and Koonin, E.V. (2002) Classification of the caspase-hemoglobinase fold: detection of new families and implications for the origin of the eukaryotic separins. Proteins, 46: 355-367.

Arena, C., Vitale, L., Virzo de Santo, A. (2005) Photosynthetic response of *Quercus ilex* L. plants grown on compost and exposed to increasing photon flux densities and elevated CO₂. Photosynthetica, 43: 615-619.

Babu, R.M., Sajeena, A., Seetharaman, K., and Reddy, M.S. (2003) Advances in genetically engineered (transgenic) plants in pest management – an over view. Crop Prot., 22: 1071-1086.

Baena-González, E., Rolland, F., Thevelein, J.M., Sheen, J. (2007) A central integrator of transcription networks in plant stress and energy signalling. Nature, 448: 938-943.

Baginsky, S., Siddique, A., Gruissem, W. (2004) Proteome analysis of tobacco bright yellow-2 (BY-2) cell culture plastids as a model for undifferentiated heterotrophic plastids. J. Proteome Res., 3: 1128 – 1137.

Barrett, A.J. (1980) Fluorimetric assays for cathepsin B and cathepsin H with methylcoumarylamide substrates. Biochem. J., 187: 909-912.

Barret, A.J., Rawlings, N.D., Cavies, M.E. Machleidt, W., Salvesen, G. and Turk, V. (1986). Cysteine protease inhibitors of the cystatin super family. In: Protease inhibitors (Barrett, A.J. and Salvesen, G., eds.) pp. 515-569. Elsevier. Amsterdam

Barta, E., Pintar, A., Pongor, S. (2002) Repeats with variations: accelerated evolution of the *Pin2* family of proteinase inhibitors. Trends Genet., 18: 600-603.

Basset, G., Raymond, P., Malek, L. and Brouquisse, R. (2002) Changes in the expression and the enzymatic properties of the 20S proteosome in sugar-starved maize roots. Evidence for an in vivo oxidation of the proteosome. Plant Physiol., 128: 1149-1162.

Bazzaz, F.A. (1990) The response of natural ecosystems to the rising global CO₂ levels. Annu. Rev. Ecol. Syst., 21: 167-196.

Beers, E.P., Woffenden, B.J., Zhao, C. (2000) Plant proteolytic enzymes: possible roles during programmed cell death. Plant Mol. Biol., 44: 399-415.

Belenghi, B., Acconcia, F., Trovato, M., Perazzolli, M., Bocedi, A., Polticelli, F., Ascenzi, P., and Delledonne, M. (2003) AtCYS1, a cystatin from *Arabidopsis thaliana*, suppresses hypersensitive cell death. Eur. J. Biochem., 270: 2593-2604.

Belenghi, B., Romero-Puertas, M.C., Vercammen, D., Brackenier, A., Inzé, D., Delledonne, M., Van Breusegem, F. (2007) Metacaspase activity of *Arabidopsis thaliana* is regulated by S-nitrosylation of a critical cysteine residue. J. Biol. Chem., 282: 1352-1358.

Benchekroun, A., Michaud, D., Nguyen-Quoc, B., Overney, S., Desjardins, Y., Yelle, S. (1995) Synthesis of active oryzacystatin I in transgenic potato plants. Plant Cell Rep., 14: 585-588.

Benson, A.A. and Calvin, M. (1950) Carbon dioxide fixation by green plants. Ann. Rev. Plant Physio., 1: 25–40.

Berger, D. and Altmann, T. (2000). A subtilisin-like serine protease involved in the regulation of stomatal density and distribution in *Arabidopsis thaliana*. Gene Dev., 14: 1119-1131.

Besford, R.T. (1990) The Greenhouse Effect: Acclimation of tomato plants growing in high CO₂, relative changes in Calvin cycle enzymes. Journal of Plant Physiol., 136: 458-463.

Beyene, G., Foyer, C.H. and Kunert, K.J. (2006) Two new cysteine proteases with specific expression patterns in mature and senescent tobacco (*Nicotina tabacum* L.) leaves. J. Exp. Bot., 57: 1431-1443.

Bhalerao, R., Keskitalo, J., Sterky, F., Erlandsson, R., Bjorkbacka, H., Birve, S.J., Karlsson, J., Gardestrom, P., Gustafsson, P., Lundeberg, J. and Jansson, S. (2003) Gene expression in Autumn leaves. Plant Physiol., 131: 430-442.

Bindi, M., Hacour, A., Vandermeiren, K., Craigon, J., Ojanperä, K., Selldén, G., Högy, P., Finn, J., Fibbi, L. (2002) Chlorophyll concentration of potatoes grown under elevated carbon dioxide and/or ozone concentrations. *Eur. J. Agron.*, 17: 319-335.

Bleecker, A.B. and Patterson, S.E. (1997) Last exit: senescence, abscission, and meristem arrest in *Arabidopsis*. *Plant Cell*, 7: 1169-1179.

Boese, S.R., Wolfe, D.W., Melkonian, J.J. (1997) Elevated CO₂ mitigates chilling-induced water stress and photosynthetic reduction during chilling. *Plant Cell Environ.*, 20, 625-632.

Boetsch, J., Chin, J., Ling, M., Croxdale, J. (1996) Elevated carbon dioxide affects the patterning of subsidiary cells in *Tradescantia* stomatal complexes. *J. Exp. Bot.*, 47: 925-931.

Bolstad, B.M., Irizarry, R.A., Astrand, M., Speed, T.P. (2003) A comparison of normalization methods for high density oligonucleotide array data based on bias and variance. *Bioinformatics*, 19: 185-193.

Bolter, C.J. (1993) Methyl jasmonate induces papain inhibitor(s) in tomato leaves. *Plant Physiol.*, 103: 1347-1353.

Botella, M. A., Xu, Y., Prabha, T.N., Zhao, Y., Narasimhan, M.L., Wilson, K.A., Nielsen, S.S., Bressan, R.A., Hasegawa, P.M. (1996) Differential expression of soybean cysteine protease inhibitor genes during development and in response to wounding and methyl jasmonate. *Plant Physiol.*, 112: 1201-1210.

Boulter, D (1993) Insect pest control by copying nature using genetically engineered crops. *Biochemistry* 34: 1453-1466.

Bowes, G. (1991) Growth at elevated CO₂: photosynthetic responses mediated through rubisco. *Plant Cell Environ.*, 14: 795-806.

Bowes, G. (1993) Facing the inevitable: plants and increasing atmospheric CO₂. *Ann. Rev. Plant Physio. and Plant Mol. Biol.*, 44: 309-332.

Bowler, C. and Fluhr, R. (2000) The role of calcium and activated oxygens as signals for controlling cross-tolerance. *Trends Plant Sci.*, 5: 241-246.

Bozhkov, P.V., Suarez, M.F., Filonova, L.H., Daniel, G., Zamyatnin, A.A., Jr., Rodriguez-Nieto, S., Zhivotovsky, B., Smertenko, A. (2005) Cysteine protease mcII-Pa executes programmed cell death during plant embryogenesis. *Proc. Natl. Acad. Sci. U. S. A.*, 102: 14463-14468.

Bradford, M.M. (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein dye-binding. *Anal. Biochem.*, 72: 248–254.

Budai-Hadrian, O., Davydov, O., Wasternack, C., Fluhr, R. (2006) SAS1 Serpin is at a node of cross-talk between jasmonic and salicylic acid pathways in *Arabidopsis*. *Israel J. Plant Sci.*, 54: 55 – 86.

Bushnell, T.P., Bushnell, D., Jagendorf, A.T. (1993) A purified zinc protease of pea chloroplast, EP1, degrades the large subunit of ribulose -1,5-bisphosphate carboxylase/oxygenase. *Plant Physiol.*, 103: 585-591.

Callis, J. (1995) Regulation of protein degradation. *Plant Cell*, 7: 845–857.

Callis, J. and Vierstra, R.D. (2000) Protein degradation in signalling. *Curr. Opin. Plant Biol.*, 3: 381-386.

Campbell, W.J., Allen, L.H., Bowes, G. (1988) Effects of CO₂ concentration on rubisco activity, amount, and photosynthesis in soybean leaves. *Plant Physiol.*, 88: 1310-1316.

Carlson, R.W. and Bazzaz, F.A. (1981) Photosynthetic and growth response to fumigation with SO₂ at elevated CO₂ for C₃ and C₄ plants. *Oecologia*, 54: 50-54.

Cercos, M., Urbez, C., Carbonell, J. (2003) A serine carboxypeptidase gene (PsCP), expressed in early steps of reproductive and vegetative development in *Pisum sativum*, is induced by gibberellins. *Plant Mol. Biol.*, 51: 165-174.

Chan, P.H. and Wildman, S.G. (1972) Chloroplast DNA codes for the primary structure of the large subunit of fraction I protein. *Biochim. Biophys. Acta*, 277: 677-680.

Chaves, M.M. and Pereira, J.M. (1992) Water stress, CO₂ and climate change. *J. Exp. Bot.*, 43: 1131-1139.

Chen, T.H.H. and Murata, N. (2002) Enhancement of tolerance of abiotic stress by metabolic engineering of betaines and other compatible solutes. *Curr. Opin. Plant Biol.*, 5: 250-257.

Cheng, S-H., Moore, B.D., Seemann, J.R. (1998) Effects of short- and long-term treatments at elevated CO₂ on the expression of Rubisco genes and leaf carbohydrate accumulation in *Arabidopsis thaliana* (L.) Henyh. *Plant Physiol.*, 116: 715-723.

Chenna, R., Sugawara, H., Koike, T., Lopez, R., Gibson, T.J., Higgins, D.G., Thompson, J.D. (2003) Multiple sequence alignment with the Clustal series of programs. *Nucleic Acids Res.*, 31: 3497-3500.

Cheong, Y.H., Chang, H.-S., Gupta, R., Wang, X., Zhu, T., Luan, S. (2002) Transcriptional profiling reveals novel interactions between wounding, pathogen, abiotic stress, and hormonal responses in *Arabidopsis*. *Plant Physiol.*, 129: 1-17.

Chiang, C.-H., Wang, J.J., Jan, F.-J., Yeh, S.-D., and Gonsalves, D. (2001) Comparative reactions of recombinant papaya ringspot viruses with chimeric coat protein (CP) genes and wild-type viruses on CP-transgenic papaya. *J. Gen. Virol.*, 82: 2827-2836.

Chiba, A., Ishida, H., Nishizawa, N.K., Makino, A., Mae, T. (2003) Exclusion of ribulose-1,5-bisphosphate carboxylase/oxygenase from chloroplasts by specific bodies in naturally senescing leaves of wheat. *Plant Cell Physiol.*, 44: 914-921.

Chinnusamy, V., Schumaker, K., Zhu, J.-K. (2004) Molecular genetic perspectives on cross-talk and specificity in abiotic stress signalling in plants. *J. Exp. Bot.*, 55: 225-236.

Chrispeels, M.J. and Herman, E.M. (2000) Endoplasmic reticulum-derived compartments function in storage and as mediators of vacuolar remodelling via a new type of organelle, precursor protease vesicles. *Plant Physiol.*, 123: 1227-1233.

Christou, P., Capell, T., Kohli, A., Gatehouse, J.A., Gatehouse, A.M.R. (2006) Recent developments and future prospects in insect pest control in transgenic crops. *Trends Plant Sci.*, 11: 302-308.

Clemente, T.E., LaVallee, B.J., Howe, A.R., Conner-Ward, D., Rozman, R.J., Hunter, P.E., Broyles, D.L., Kasten, D.S., Hinchee, M.A. (2000) Progeny analysis of glyphosate selected transgenic soybeans derived from *Agrobacterium*-mediated transformation. *Crop Sci.*, 40: 797-803.

Colfalonieri, M., Allegro, G., Balestrazzi, A., Fogher, C., and Delledonne, M. (1998) Regeneration of *Populus nigra* transgenic plants expressing a Kunitz protease inhibitor (Kti3) gene. *Mol. Breed.*, 4: 137-145.

Conley, M.M., Kimball, B.A., Brooks, T.J., Pinter, Jr., P.J., Hunsaker, D.J., Wall, G.W., Adam, N.R., LaMorte, R.L., Matthias, A.D., Thompson, T.L., Leavitt, S.W., Ottman, M.J., Cousins, A.B., Triggs, J.M. (2001). CO₂ enrichment increases water-use efficiency in sorghum. *New Phytol.*, 151, 407-412.

Cooper, B., Eckert, D., Andon, N.L., Yates, J.R., Haynes, P.A. (2003) Investigative proteomics: identification of an unknown plant virus from infected plants using mass spectrometry. *J. Am. Soc. Mass Spectrom.*, 14: 736–741.

Coupe, S.A., Palmer, B.G., Lake, J.A., Overy, S.A., Oxborough, K., Woodward, F.I., Gray, J.E., Quick, W.P. (2006) Systemic signalling of environmental cues in *Arabidopsis* leaves. *J. Exp. Bot.*, 57: 329-341.

Cousins, A.B., Adam, N.R., Wall, G.W., Kimball, B.A., Pinter, Jr., P.J., Leavitt, S.W., LaMorte, R.L., Matthias, A.D., Ottman, M.J., Thompson, T.L., Webber, A.N. (2001) Reduced photorespiration and increased energy-use efficiency in young CO₂-enriched sorghum leaves. *New Phytol.*, 150: 275-284.

Crafts-Brandner, S.J. and Salvucci, M.E. (2000) Rubisco activase constrains the photosynthetic potential of leaves at high temperature and CO₂. *P. Natl. Acad. Sci. USA*, 97: 13430-13435.

Croxdale, J. (1998) Stomatal patterning in monocotyledons: *Tradescantia* as a model system. *J. Exp. Bot.*, 49: 279–292.

Dal Degan, F., Rocher, A., Cameron-Mills, V., Von Wettstein, D. (1994) The expression of serine carboxypeptidases during maturation and germination of the barley grain. *P. Natl. Acad. Sci. USA*, 91: 8209–8213.

Dani, V., Simon, W.J., Duranti, M., Croy, R.R.D. (2005) Changes in the tobacco leaf apoplast proteome in response to salt stress. *Proteomics*, 5: 737-745.

Dean, C., Dunsmuir, P., Bedbrook, J. (1989) Structure, evolution, and regulation of *rbcS* genes in higher plants. *Ann. Rev. Plant Phys.*, 40: 415-439.

De Block, M., Verduyn, C., De Brouwer, D., Cornelissen, M. (2005) Poly(ADP-ribose) polymerase in plants affects energy homeostasis, cell death and stress tolerance. *Plant J.*, 41: 95-106.

De Las Rivas, J., Andersson, B., Barber, J. (1992) Two sites of primary degradation of the D1-protein induced by acceptor or donor side photo-inhibition in photosystem II core complexes. *FEBS Lett.*, 301: 246-52.

De Leo, F., Bonadé-Bottino, M., Ceci, L.R., Gallerani, R., Jouanin, L. (2001) Effects of a mustard trypsin inhibitor expressed in different plants on three lepidopteran pests. *Insect Biochem. Molec.*, 31: 593-602.

Del Pozo, O. and Lam, E. (1998) Caspases and programmed cell death in the hypersensitive response of plants to pathogens. *Curr. Biol.*, 8: 1129–1132.

Deng, X. and Woodward, F.I. (1998) The growth and yield responses of *Fragaria ananassa* to elevated CO₂ and N supply. *Ann Bot – London*, 81: 67-71.

Desimone, M., Henke, A., Wagner, E. (1996) Oxidative stress induces partial degradation of the large subunit of ribulose 1,5- bisphosphate carboxylase/oxygenase in isolated chloroplasts of barley. *Plant Physiol.*, 111: 789-796

Diop, N.N., Kidric, M., Repellin, A., Gareil, M., d'Arcy-Lameta, A., Pham Thi, A.T. and Zuijly-Fodil, Y. (2004) A multicystatin is induced by drought-stress in cowpea (*Vigna unguiculata* (L.) Walp.) leaves. *FEBS Lett.*, 577: 545-550.

Domínguez, F. and Cejudo, F.J. (1998) Germination-related genes encoding proteolytic enzymes are expressed in the nucellus of developing wheat grains. *Plant J.*, 15: 569-574.

Domínguez, F., González, M., Cejudo, F.J. (2002) A germination-related gene encoding a serine carboxypeptidase is expressed during the differentiation of the vascular tissue in wheat grains and seedlings. *Planta*, 215: 727-734.

Drake, B.G., Gonzalez-Meler, M.A., Long, S.P. (1997) More efficient plants: A consequence of rising atmospheric CO₂? *Ann. Rev. Plant Phys.*, 48, 609-639.

Driscoll, S.P., Prins, A., Olmos, E., Kunert, K.J., Foyer, C.H. (2006). Specification of adaxial and abaxial stomata, epidermal structure and photosynthesis to CO₂ enrichment in maize leaves. *J. Exp. Bot.*, 57: 381-390.

Dreher, K. and Callis, J. (2007) Ubiquitin, hormones and biotic stress in plants. *Ann. Bot.-London*, 99: 787-822.

Duan, X.L., Li, X.G., Xue, Q.Z., Aboelsaad, M., Xu, D.P., and Wu, R. (1996) Transgenic rice plants harboring an introduced potato protease inhibitor II gene are insect resistant. *Nat. Biotechnol.*, 14: 494-498.

Eason, J.R., Ryan, D.J., Pinkney, T.T., and O'Donoghue, E.M. (2002) Programmed cell death during flower senescence: isolation and characterization of cysteine proteases from *Sandersonia aurantiaca*. *Funct. Plant Biol.*, 29: 1055–1064.

Edmonds, H.S., Gatehouse, L.N., Hilder, V.A., Gatehouse, J.A. (1996) The inhibitory effects of the cysteine protein protease inhibitor, oryzacystatin, on digestive proteases and on larval survival and development of the southern corn rootworm (*Diabrotica undecimpunctata howardi*). *Entomol. Exp. Appl.*, 78: 83-94.

Edwards, G. and Walker, D. (1983) C₃, C₄: mechanisms, and cellular and environmental regulation of photosynthesis. Blackwell Science Publishers, London, UK.

Elbaz, M., Avni, A., Weil, M. (2002) Constitutive caspase-like machinery executes programmed cell death in plant cells. *Cell Death Differ.*, 9: 726-733.

Ellis, R.J. (1981). Chloroplast proteins: synthesis, transport and assembly. *Ann. Rev. Plant Physio.* 32, 111-137.

Emanuelsson, O., Nielsen, H., Brunak, R., Von Heijne, G. (2000) Predicting subcellular localization of proteins based on their N-terminal amino acid sequence. *J. Mol. Biol.*, 300: 1005-1016.

Fan, S-G. and Wu, G-J. (2005) Characteristics of plant protease inhibitors and their applications in combating phytophagous insects. *Bot. Bull. Acad. Sinica*, 46: 273-292.

Farrar, J.F. and Williams, M.L. (1991) The effects of increased atmospheric carbon-dioxide and temperature on carbon partitioning, source-sink relations and respiration. *Plant Cell Environ.*, 14: 819-830.

Feller, U. Anders, I., Mae, T. (2007) Rubiscoolytics: fate of Rubisco after its enzymatic function in a cell is terminated. *J. Exp. Bot.*, doi:10.1093/jxb/erm242

Ferris, R., Long, L., Bunn, S.M., Robinson, K.M., Bradshaw, H.D., Rae, A.M., Taylor, G. (2002) Leaf stomatal and epidermal cell development: Identification of putative quantitative trait loci in relation to elevated carbon dioxide concentration in poplar. *Tree Physiol.*, 22: 633–640.

Ferry, N., Edwards, M.G., Gatehouse, J.A., Gatehouse, A.M.R. (2004) Plant-insect interactions: molecular approaches to insect resistance. *Curr. Opin. Biotech.*, 15: 155-161.

Finkelstein, R.R. and Gibson, S.I. (2002) ABA and sugar interactions regulating development: cross-talk or voices in a crowd? *Curr. Opin. Plant Biol.*, 5: 26–32.

Finkelstein, R.R. and Lynch, T.J. (2000) Abscisic acid inhibition of radicle emergence but not seedling growth is suppressed by sugars. *Plant Physiol.*, 122: 1179–1186.

Fisher, A. and Feller, U. (1994) Senescence and protein degradation in leaf segments of young winter wheat: influence of leaf age. *J. Exp. Bot.*, 45: 103-109.

Fischhoff, D.A., Bowdish, K.S., Perlak, F.J., Marrone, P.G., McCormick, S.H., Niedermeyer, J.G., Dean, D.A., Kusano-Kretzmer, K., Mayer, E.J., Rochester, D.E., Rogers, S.G., Fraley, R.T. (1987) Insect tolerant transgenic tomato plants. *Bio/Technol.*, 5: 807–813.

Flowers, T.J., and Yeo, A.R. (1995) Breeding for salinity resistance in crop plants: where next? *Aust. J. Plant Physiol.*, 22: 875-884.

Flowers, T.J. (2004) Improving crop salt tolerance. *J. Exp. Bot.*, 55: 307-319.

Foyer, C.H. and Noctor, G. (2003) Redox sensing and signaling associated with reactive oxygen in chloroplasts, peroxisomes and mitochondria. *Physiol. Plantarum*, 119: 355–364.

Foyer, C.H. and Noctor, G. (2007) Shape-shifters building bridges? Stromules, matrixules and metabolite channelling in photorespiration. *Trends Plant Sci.*, 12: 381-383.

Franceschetti, M., Perry, B., Thompson, B., Hanfrey, C., Michael, A.J. (2004) Expression proteomics identifies biochemical adaptations and defense responses in transgenic plants with perturbed polyamine metabolism. FEBS Lett, 576: 477-480.

Frugis, G. and Chua, N.-H. (2002) Ubiquitin-mediated proteolysis in plant hormone signal transduction. Trends Cell Biol., 12: 308-311.

Fuentes-Prior, P. and Salvesen, G.S. (2004) The protein structures that shape caspase activity, specificity, activation and inhibition. Biochem. J., 384: 201–232.

Fujita, M., Fujita, Y., Noutoshi, Y., Takahashi, F., Narusaka, Y., Yamaguchi-Shinozaki, K., Shinozaki, K. (2006) Crosstalk between abiotic and biotic stress responses: a current view from the points of convergence in the stress signaling networks. Curr. Opin. Plant Biol., 9: 436-442.

Furbank, R.T. and Taylor, W.C. (1995) Regulation of photosynthesis in C3 and C4 Plants: A molecular approach. Plant Cell, 7: 797-807.

Gaddour K, Vicente-Carbajosa J, Lara P, Isabel-Lamonedo I, Díaz I, Carbonero P. (2001) A constitutive cystatin-encoding gene from barley (*Icy*) responds differentially to abiotic stimuli. Plant Mol. Biol., 45: 599–608.

Garcia-Ferris, C. and Moreno, J. (1994) Oxidative modification and breakdown of ribulose-1,5-bisphosphate carboxylase oxygenase induced in *Euglena gracilis* by nitrogen starvation. Planta, 193: 208-215.

Gasteiger, E., Hoogland, C., Gattiker, A., Duvaud, S., Wilkins,M.R., Appel, R.D., Bairoch, A. (2005) In: The Proteomics Protocols Handbook (Walker, J.M., ed) pp. 571-607. Humana Press.

Gepstein, S., Sabehi, G., Carp, M.-J., Hajouj, T., Nesher, M.F.O., Yariv, I., Dor, C., Bassani, M. (2003) Large-scale identification of leaf senescence-associated genes. Plant J., 36: 629–642.

Gesch, R.W., Boote, K.J., Vu, J.C.V., Allen, H. Jr., Bowes, G. (1998) Changes in growth CO₂ result in rapid adjustments of ribulose-1,5-bisphosphate carboxylase/oxygenase small subunit gene expression in expanding and mature leaves of rice. *Plant Physiol.*, 118: 521-529.

Ghannoum, O., Von Caemmerer, S., Barlow, E.W.R., Conroy, J.P. (1997) The effect of CO₂ enrichment and irradiance on the growth, morphology and gas-exchange of a C₃ (*Panicum laxum*) and a C₄ (*Panicum antidotale*) grass. *Aust. J. Plant Physiol.*, 24: 227–237.

Ghannoum, O., Von Caemmerer, S., Conroy, J.P. (2001) Plant water use efficiency of 17 Australian NAD-ME and NADP-ME C₄ grasses at ambient and elevated CO₂ partial pressure. *Austr. J. Plant Physiol.*, 28: 1207–1217.

Gietl, C. and Schmid, M. (2001) Ricinosomes: an organelle for developmentally regulated programmed cell death in senescing plant tissues. *Naturwissenschaften*, 88: 49–58.

Gish, W. and States, D. J. (1993). Identification of protein coding regions by database similarity search. *Nat. Genet.*, 3: 266-72.

Giri, A.P., Wünsche, H., Mitra, S., Zavala, J.A., Muck, A., Svatoš, A., Baldwin, I.T. (2006) Molecular interactions between the specialist herbivore *Manduca sexta* (Lepidoptera, Sphingidae) and its natural host, *Nicotiana attenuata*. VII. Changes in the plant's proteome. *Plant Physiol.*, 142: 1621-1641.

Glickman, M.H. (2000) Getting in and out of the proteasome. *Semin. Cell Dev. Biol.*, 11: 149-158.

Gong, Z., Lee, H., Xiong, L., Jagendorf, A., Stevenson, B., Zhu, J.-K. (2002) RNA helicase-like protein as an early regulator of transcription factors for plant chilling and freezing tolerance. *P. Natl. Acad. Sci. USA*, 99: 11507-11512.

Granat, S.J., Wilson, K.A., Tan-Wilson, A.L. (2003) New serine carboxypeptidases in mung bean seedling cotyledons. *J. Exp. Bot.*, 160: 1263–1266.

Granell A, Cercos M, Carbonell J. (1998) Plant cysteine proteases in germination and senescence. In: *The Handbook of Proteolytic Enzymes* (Barrett, A.J., Rawlings, N.D. and Woessner, J.F., eds), pp. 578-583. San Diego: Academic Press.

Gray, J.E., Holroyd, G.H., Van Der Lee, F.M., Bahrami, A.R., Sijmons, P.C., Woodward, F.I., Schuch, W., Hetherington, A.M. (2000) The HIC signalling pathway links CO₂ perception to stomatal development. *Nature*, 408: 713-716.

Grbić, V. (2003) SAG2 and SAG12 protein expression in senescing *Arabidopsis* plants. *Physiol. Plantarum*, 119:263–269.

Greer, D.H., Laing, W.A., Campbell, B.D. (1995) Photosynthetic responses of thirteen pasture species to elevated CO₂ and temperature. *Aust. J. Plant Physiol.*, 22: 713–722.

Griffin, K.L. and Seemann, J.R. (1996) Plants, CO₂ and photosynthesis in the 21st century. *Chem. Biol.*, 3: 245-254.

Grime, J.P. (1989) Whole-plant responses to stress in natural and agricultural systems. In: H. Jones, T.J. Flowers, M.B. Jones (Eds.), *Plants Under Stress*, Cambridge University Press, Cambridge. Pp. 31-46.

Grover, A., Sahi, C., Sanan, N., and Grover, A. (1999) Taming abiotic stresses in plants through genetic engineering: current strategies and perspective. *Plant Sci.*, 143: 101-111.

Grover, A., Agarwal, M., Katiyar-Agarwal, S., Sahi, C., Agarwal, S. (2000) Production of high temperature tolerant transgenic plants through manipulation of membrane lipids. *Curr. Sci. India*, 79: 557-559.

Grover, A., Aggarwal, P.K., Kapoor, A., Katiyar-Agarwal, S., Agarwal, M., Chandramouli, A. (2003) Addressing abiotic stresses in agriculture through transgenic technology. *Curr. Sci. India*, 84: 355-367.

Gruis, D.F., Selinger, D.A., Curran, J.M., Jung, R. (2002) Redundant proteolytic mechanisms process seed storage proteins in the absence of seed-type members of the vacuolar processing enzyme family of cysteine proteases. *Plant Cell*, Vol. 14, 2863-2882.

Gunning, B.E.S. (2005) Plastid stromules: video microscopy of their outgrowth, retraction, tensioning, anchoring, branching, bridging, and tip-shedding. *Protoplasma*, 225: 33-42.

Gutiérrez-Campos, R., Torres-Acosta, J.A., Pérez-Martínez, J.D.J. and Gómez-Lim, M.A. (2001) Pleiotropic effects in transgenic tobacco plants expressing oryzacystatin I gene. *HortScience*, 36: 118–119.

Guy, C.L., Huber, J.L.A., Huber, S.C. (1992). Sucrose phosphate synthase and sucrose accumulation at low temperature. *Plant Physiol.*, 100: 502-508.

Hara-Nishimura, I., Inoue, K., Nishimura, M. (1991) A unique vacuolar processing enzyme responsible for conversion of several proprotein precursors into the mature forms. *FEBS Lett.*, 294: 89-93.

Hara-Nishimura, I., Takeuchi, Y., Nishimura, M. (1993) Molecular characterization of a vacuolar processing enzyme related to a putative cysteine protease of *Schistosoma mansoni*. *Plant Cell*, 5: 1651-1659.

Hara-Nishimura, I., Kinoshita, T., Hiraiwa, N., Nishimura, M. (1998) Vacuolar processing enzymes in protein-storage vacuoles and lytic vacuoles. *J. Plant Physiol.*, 152: 668-674.

Harrak, H., Azelmat, S., Baker, E.N. and Tabaeizadeh, Z. (2001) Isolation and characterization of gene encoding a drought-induced cysteine protease in tomato (*Lycopersicon esculentum*). *Genome*, 44: 368-374.

Hatch, M.D. (1992) C₄ Photosynthesis: An unlikely process full of surprises. *Plant Cell Physiol.*, 33: 333-342.

Hayashi, H., Alia Mustardy, L., Deshnium, P., Ida, M., Murata, N. (1997) Transformation of *Arabidopsis thaliana* with the codA gene for choline oxidase: accumulation of glycinebetaine and enhanced tolerance to salt and cold stress. *Plant J.*, 12: 133-142.

Hayashi, Y., Yamada, K., Shimada, T., Matsushima, R., Nishizawa, N.K., Nishimura, M., Hara-Nishimura, I. (2001). A protease-storing body that prepares for cell death or stresses in the epidermal cells of *Arabidopsis*. *Plant Cell Physiol.*, 42: 894-899.

Heagle, A.S., Miller, J.E., Sherrill, D.E., Rawlings, J.O. (1993) Effects of ozone and carbon dioxide mixtures on two clones of white clover/ *New Phytol.*, 123: 751-762.

Hellman, H. and Estelle, M. (2002) Plant development: regulation by protein degradation. *Science*, 297: 793-797.

Hensel, L.L., Grbic, V., Baumgarten, D.A., Bleecker, A.B. (1993) Developmental and age-related processes that influence the longevity and senescence of photosynthetic tissues in *Arabidopsis* *Plant Cell*, 5: 553-64.

Herman, E. and Schmidt, M. (2004) Endoplasmic reticulum to vacuole trafficking of endoplasmic reticulum bodies provides an alternative pathway for protein transfer to the vacuole. *Plant Physiol.*, 136: 3440-3446.

Herrick, J.D. and Thomas, R.B. (2003) Leaf senescence and late-season net photosynthesis of sun and shade leaves of overstorysweetgum (*Liquidambar styraciflua*) grown in elevated and ambient carbon dioxide concentrations. *Tree Physiol.*, 23: 109-118.

Hetherington, A.M. and Woodward, F.I. (2003) The role of stomata in sensing and driving environmental change. *Nature*, 424: 901-908.

Hilder, V.A., Gatehouse, A.M.R., Sheerman, S.E., Baker, R.F., and Boulter, D. (1987) A novel mechanism of insect resistance engineered into tobacco. *Nature*, 330: 160-163.

Hilder, V.A. and Boulter, D. (1999) Genetic engineering of crop plants for insect resistance – a critical review. *Crop Prot.*, 18: 177-191.

Hiraiwa, N., Takeuchi, Y., Nishimura, M. and Hara-Nishimura, I. (1993) A vacuolar processing enzyme in maturing and germinating seed: its distribution and associated changes during development. *Plant Cell Physiol.*, 34: 1197-1204.

Hochstrasser, M. (1995) Ubiquitin, proteasomes, and the regulation of intracellular protein degradation. *Curr. Opin. Cell Biol.*, 7: 215–223.

Hoeberichts, F.A., Ten Have, A., Woltering, E.J. (2003) A tomato metacaspase gene is upregulated during programmed cell death in *Botrytis cinerea*-infected leaves. *Planta*, 217: 517-522.

Hopkins, M., Taylor, C., Liu, Z., Ma, F., McNamara, L., Wang, T.-W., Thompson, J.E. (2007) Regulation and execution of molecular disassembly and catabolism during senescence. *New Phytol.*, 175: 201–214.

Horsch, R.B., Fry, J.E., Hoffman, N.L., Wallroth, M., Eichholtz, D., Rogers, S.G., and Frailey, R.T. (1985) A simple and general method for transferring genes into plants. *Science*, 227: 1229-1231.

Hörtensteiner, S. and Feller, U. (2002) Nitrogen metabolism and remobilization during senescence. *J. Exp. Bot.*, 53: 927-937.

Hsiao, T.C. and Jackson, R.B. (1999) Interactive effects of water stress and elevated CO₂ on growth, photosynthesis, and water use efficiency. In: Luo, Y. and Mooney, H.A. (eds). *Carbon Dioxide and Environmental Stress*. Academic Press. Pp. 3-26.

Huber, S.C. and Huber, J.L. (1992) Role of sucrose-phosphate synthase in sucrose metabolism in leaves. *Plant Physiol.*, 4: 1275-1278.

Huber, S.C. and Huber, J.L. (1996) Role and regulation of sucrose-phosphate synthase in higher plants. *Annu. Rev. Plant Phys.*, 47: 431-444.

Huffaker, R.C. (1990) Proteolytic activity during senescence of plants. *New Phytol.*, 116: 199–231.

Humbeck, K., Quast, S. and Krupinska, K. (1996) Functional and molecular changes in the photosynthetic apparatus during senescence of flag leaves from field-grown barley plants. *Plant Cell Environ.*, 19: 337–344.

Hymus, G.J., Baker, N.R., Long, S.P. (2001) Growth in elevated CO₂ can both increase and decrease photochemistry and photoinhibition of photosynthesis in a predictable manner. *dactylis glomerata* grown in two levels of nitrogen nutrition *Plant Physiol.*, 127: 1204–1211.

Iba, K. (2002) Acclimative response to temperature stress in higher plants: Approaches of gene engineering for temperature tolerance. *Annu. Rev. Plant Biol.*, 53: 225-245.

Idso, S.B., Kimball, B.A., Pettit, G.R., Garner, L.C., Pettit, G.R., Backhaus, R.A. (2000) Effects of atmospheric CO₂ enrichment on the growth and development of *Hymenocallis littoralis* (Amaryllidaceae) and the concentrations of several antineoplastic and antiviral constituents of its bulbs. *Am. J. Bot.*, 87: 769-773.

Inoue, K., Motozaki, A., Takeuchi, Y., Nishimura, M. and Hara-Nishimura, I. (1995) Molecular characterization of proteins in protein-body membrane that disappear most rapidly during transformation of protein bodies into vacuoles. *Plant J.*, 7: 235-243.

Irie, K., Hosoyama, H., Akeuchi, T., Iwabuchi, K., Watanabe, H., Abe, M., Abe, K., Arai, S. (1996) Transgenic rice established to express corn cystatin exhibits strong inhibitory activity against insect gut proteases. *Plant Mol. Biol.*, 30: 149-157.

Irving, L.J. and Robinson, D. (2006) A dynamic model of RuBisCO turnover in cereal leaves. *New Phytol.*, 169: 493-504.

Ishida, H., Nishimori, Y., Sugisawa, M., Makino, A., Mae, T. (1997) The large subunit of ribulose-1,5-bisphosphate carboxylase/oxygen- ase is fragmented into 37-kDa and 16-kDa

polypeptides by active oxygen in the lysates of chloroplasts from primary leaves of wheat. Plant Cell Physiol., 38: 471-479.

Ishida, H., Shimizu, S., Makino, A., Mae, T. (1998) Light-dependent fragmentation of the large subunit of ribulose-1,5-bisphosphate carboxylase/oxygenase in chloroplasts isolated from wheat leaves. Planta, 204: 305-309.

Ishida, H., Yoshimoto, K., Reisen, D., Makino, A., Ohsumi, Y., Hanson, M., Mae, T. (2007) Visualisation of RuBisCO-containing bodies derived from chloroplasts in living cells of Arabidopsis. Photosynthesis Res., 91:275-276.

Jablonski, L.M., Wang, X., Curtis, P.S. (2002) Plant reproduction under elevated CO₂ conditions: a meta-analysis of reports on 79 crop and wild species. New Phytol., 156: 9–26.

Jacob, J., Greitner, C., Drake, B.G. (1995) Acclimation of photosynthesis in relation to Rubisco and non-structural carbohydrate contents and in situ carboxylase activity in *Scirpus olneyi* grown at elevated CO₂ in the field. Plant Cell Environ., 18: 875-884.

Jain, R.K., and Selvaraj, G. (1997) Molecular genetic improvement of salt tolerance in plants. Biotechnol. Annu. Rev., 3: 245-267.

Jang, J.C. and Sheen, J. (1994) Sugar sensing in higher plants. Plant Cell, 6: 1665–79.

Janska, H. (2005) ATP-dependent proteases in plant mitochondria: what do we know about them today? Physiol. Plant, 123: 399–405.

Jiang, M. and Zhang, J. (2003) Cross-talk between calcium and reactive oxygen species originated from NADPH oxidase in abscisic acid-induced antioxidant defence in leaves of maize seedlings. Plant Cell Environ., 26: 929-939.

Johnson, R., and Ryan, C.A. (1990). Wound-inducible potato inhibitor II genes: Enhancement of expression by sucrose. Plant Mol. Biol., 14: 527-536.

Johansson, E., Olsson, O., Nystrom, T. (2004) Progression and specificity of protein oxidation in the life cycle of *Arabidopsis thaliana*. *J. Biol. Chem.*, 279: 22204-22208.

Jones, A.M. and Dangl, J.L. (1996) Logjam at the Styx: programmed cell death in plants. *Trends Plant Sci.*, 1: 114-119.

Jones, M.G.K., Outlaw, Jr. W.H., Lowry, O.H. (1977). Enzymic assay of 10^{-7} to 10^{-14} moles of sucrose in plant tissues. *Plant Physiol.*, 60: 379-383.

Juarez, M.T., Twigg, R.W., Timmerman, S.M.C.P. (2004) Specification of adaxial cell fate during maize leaf development. *Development*, 131: 4533–4544.

Kardailsky, I.V. and Brewin, N.J. (1996) Expression of cysteine protease genes in pea nodule development and senescence. *Mol. Plant. Microbe Interact.*, 8: 689-695.

Kato, H. and Minamikawa, T. (1996) Identification and characterization of a rice cysteine endopeptidase that digests glutelin. *Eur. J. Biochem.*, 239: 310-316.

Kavi Kishor, P.B., Hong, Z., Miao, G., Hu, C.A., and Verma, D.P.S. (1995) Overexpression of Δ -pirroline-5-carboxylate synthetase increases proline production and confers osmotolerance in transgenic plants. *Plant Physiol* 108: 1387-1394.

Keith, C.S., Hoang, D.O., Barrett, B.M., Feigelman, B., Nelson, M.C., Thai, H., and Baysdorfer, C. (1993) Partial sequence analysis of 130 randomly selected maize cDNA clones. *Plant Physiol.*, 101:329-332.

Keys, A.J. and Parry, M.A.J. (1990) Ribulose bisphosphate carboxylase/oxygenase and carbonic anhydrase. *Meth. Plant Biochem.*, 3: 1-15.

Khan, S., Andralojc, P.J., Lea, P.J. and Parry, M.A.J. (1999) 2-carboxy-D-arabinitol 1-phosphate protects ribulose 1,5-bisphosphate carboxylase/oxygenase against proteolytic breakdown. *Eur. J. Biochem.*, 266: 840–847.

Khanna-Chopra, R., and Sinha, S.K. (1998) Prospects of success of biotechnological approaches for improving tolerance to drought stress in crop plants. Curr. Sci., 74: 25-34.

Kingston-Smith, A.H. and Foyer, C.H. (2000) Overexpression of Mn-superoxide dismutase in maize leaves leads to increased monodehydroascorbate reductase, dehydroascorbate reductase and glutathione reductase activities. J. Exp. Bot., 51: 1867-1877.

Kinoshita, T., Nishimura, M., Hara-Nishimura, I. (1995) Homologues of a vacuolar processing enzyme that are expressed in different organs in *Arabidopsis thaliana*. Plant Mol. Biol., 29: 81-89.

Kinoshita, T., Yamada, K., Hiraiwa, N., Kondo, M., Nishimura, M., Hara-Nishimura, I. (1999) Vacuolar processing enzyme is up-regulated in the lytic vacuoles of vegetative tissues during senescence and under various stressed conditions. Plant J., 19: 43-53.

Kloetzel, P. M. (2001) Antigen processing by the proteasome. Nat. Rev. Mol. Cell Biol., 2: 179-187.

Knight, H. and Knight, M.R. (2001) Abiotic stress signalling pathways: specificity and cross-talk. Trends Plant Sci., 6: 262-267.

Kobayashi, T., Kobayashi, E., Sato, S., Hotta, Y., Miyajima, N., Tanaka, A. and Tabata, S. (1994) Characterization of cDNAs induced in meiotic prophase in lily microsporocytes. DNA Res., 1: 15-26.

Koiwa, K., Shade, R.E., Zhu-Salzman, K., Subramanian, L., Murdock, L.L., Nielsen, S.S., Bressan, R.A., Hasegawa, P.M. (1998) Phage display selection can differentiate insecticidal activity of soybean cystatins. Plant J., 14: 371-379.

Koizumi, M., Yamaguchi-Shinozaki, K., Tsuji, H. and Shinozaki, K. (1993) Structure and expression of two genes that encode distinct drought-inducible cysteine proteases in *Arabidopsis thaliana*. Gene, 29: 175-182.

Kondo H, Abe K, Nishimura I, Watanabe H, Emori Y, Arai S. (1990) Two distinct cystatin species in rice seeds with different specificities against cysteine proteases. *J. Biol. Chem.*, 265: 15832–15837.

Kondo, H., Abe, K., Emori, Y. and Arai, S. (1991) Gene organization of oryzacystatin-II, a new cystatin super family member of plant origin, is closely related to that of oryzacystatin-I but different from those of the animal cystatins. *FEBS Lett.* 278: 87-90.

Konno, K., Hirayama, C., Nakamura, M., Tateishi, K., Tamura, Y., Hattori, M., Kohno, K. (2004) Papain protects papaya trees from herbivorous insects: role of cysteine proteases in latex. *Plant J.*, 37:370–378.

Kreps, J.A., Wu, Y., Chang, H.S., Zhu, T., Wang, X., Harper, J.F. (2002) Transcriptome changes for *Arabidopsis* in response to salt, osmotic, and cold stress. *Plant Physiol.*, 130: 2129–2141.

Kruger, N.J. (1990) Carbohydrate synthesis and degradation. In *Plant Physiology, Biochemistry and Molecular Biology* (Dennis, D.T. and Turpin, D.M., eds), pp. 59–76. Longman, Harlow, UK.

Krüger, J., Thomas, C.M., Golstein, C., Dixon, M.S., Smoker, M., Tang, S., Mulder, L., Jones, J.D.G. (2002) A tomato cysteine protease required for *Cf-2*-dependent disease resistance and suppression of autonecrosis. *Science*, 296: 744–747.

Krupinska, K. (2006) Fate and Activities of Plastids During Leaf Senescence. In: *Advances in Photosynthesis and Respiration*. Wise, R.R. and Hoober, J.K. (eds), vol. 23: The Structure and Function of Plastids. Springer, Netherlands. Pp. 433-449.

Kuroda, M., Ishimoto, M., Suzuki, K., Kondo, H., Abe, K., Kitamura, K., Arai, S. (1996) Oryzacystatins exhibit growth-inhibitory and lethal effects on different species of bean insect pests, *Callosobruchus chinensis* (Coleoptera) and *Riptortus clavatus* (Hemiptera). *Biosci. Biotech. Bioch.*, 60: 209-212.

Kuroda M, Kiyosaki T, Matsumoto I, Misaka T, Arai S, Abe K. (2001) Molecular cloning, characterization, and expression of wheat cystatins. *Biosci. Biotech. Bioch.*, 65: 22–28.

Lai, L.B., Nadeau, J.A., Lucas, J., Lee, E.-K., Nakagawa, T., Zhao, L., Geisler, M., Sack, F.D. (2005) The Arabidopsis R2R3 MYB proteins FOURLIPS andMYB88 restrict divisions late in the stomatal cell lineage. *Plant Cell*, 17: 2754–2767.

Lake, J.A., Quick, W.P., Beerling, D.J., Woodward, F.I. (2001) Plant development-signals from mature to new leaves. *Nature*, 411: 154-154.

Lake, J.A., Woodward, F.I., Quick, W.P. (2002) Long distance CO₂ signalling in plants. *J. Exp. Bot.*, 53: 183-193.

Langridge, P., Paltridge, N., Fincher, G. (2006) Functional genomics of abiotic stress tolerance in cereals. *Brief. Funct. Genomic. Proteomic.*, 4: 343-354.

Larkin, J.C., Marks, M.D., Nadeau, J., Sack, F. (1997) Epidermal cell fate and patterning in leaves. *Plant Cell*, 9: 1109–1120.

Laukens, K., Deckers, P., Esmans, E., Van Onckelen, H., Witters, E. (2004) Construction of a two-dimensional gel electrophoresis protein database for the *Nicotiana tabacum* cv. Bright Yellow-2 cell suspension culture. *Proteomics*, 4: 720-727.

Lawson, T., Craigon, J., Tulloch, A-M., Black, C.R., Colls, J.J., Landon, G. (2001) Photosynthetic responses to elevated CO₂ and O₃ in field-grown potato (*Solanum tuberosum*). *J. Plant Physiol.*, 158, 309–323.

Leakey, A.D.B., Bernacchi, C.J., Dohleman, F.G., Ort, D.R., Long, S.P. (2004) Will photosynthesis of maize (*Zea mays*) in the US Corn Belt increase in future [CO₂] rich atmospheres? An analysis of diurnal courses of CO₂ uptake under free-air concentration enrichment (FACE). *Glob. Change Biol.*, 10: 951–962.

Leakey, A.D.B., Uribelarrea, M., Ainsworth, E.A., Naidu, S.L., Rogers, A., Ort, D.R., Long, S.P. (2006) Photosynthesis, productivity, and yield of maize are not affected by

open-air elevation of CO₂ concentration in the absence of drought. *Plant Physiol.*, 140: 779–790.

Le Cain, D.R. and Morgan, J.A. (1998) Growth, gas exchange, leaf nitrogen and carbohydrate concentrations in NAD-ME and NADP-ME C₄ grasses grown in elevated CO₂. *Physiol. Plantarum*, 102: 297–306.

Lee, J.S. and Ellis, B.E. (2007) *Arabidopsis* MAPK Phosphatase 2 (MKP2) positively regulates oxidative stress tolerance and inactivates the MPK3 and MPK6 MAPKs. *J. Biol. Chem.*, 282: 25020–25029.

León, P. and Sheen, J. (2003) Sugar and hormone connections. *Trends Plant Sci.*, 8: 110–116.

Leple, J.C., Bonadebottino, M., Augustin, S., Pilate, G., Letan, V.D. Delplanque, A. Cornu D., Jouanin, L. (1995) Toxicity to *Chrysomela tremulae* (Coleoptera, Chrysomelidae) of transgenic poplars expressing a cysteine protease inhibitor, *Mol. Breeding*, 1: 319–328.

Lewis, C.E., Noctor, G., Causton, D., Foyer, C.H. (2000) Regulation of assimilate partitioning in leaves. *Aust. J. Plant Physiol.*, 27: 507–517.

Lewis, C.E., Peratoner, G., Cairns, A.J., Causton, D.R., Foyer, C.H. (1999) Acclimation of the summer annual species, *Lolium temulentum*, to CO₂ enrichment. *Planta*, 210: 104–114.

Li, Z., Sommer, A., Dingermann, T., Noe, C.R. (1996) Molecular cloning and sequence analysis of a cDNA encoding a cysteine protease inhibitor from Sorghum bicolor seedlings. *Mol. Gen. Genet.*, 251: 499–502.

Li, J., Lease, K.A., Tax, F.E., Walker, J.C. (2001) BRS1, a serine carboxypeptidase, regulates BRI1 signaling in *Arabidopsis thaliana*. *P. Natl. Acad. Sci. USA*, 98: 5916–5921.

Liang, C., Brookhart, G., Feng, G.H., Reeck, G.R., Kramer, K.J. (1991) Inhibition of digestive proteases of stored grain coleopteran by oryzacystatin, a cysteine proteinase inhibitor from rice seed. FEBS Lett., 278: 139-142.

Lichtenhaller, H.K. and Wellburn, A.R. (1983) Determination of total carotenoids and chlorophylls a and b of leaf extracts in different solvents. Biochem. Soc. T., 11: 591-592.

Lin, J., Jach, M. E., Ceulemans, R. (2001) Stomatal density and needle anatomy of Scots pine (*Pinus sylvestris*) are affected by elevated CO₂. New Phytol., 150: 665-674.

Lin, J.S. and Wu, S.H. (2004) Molecular events in senescing *Arabidopsis* leaves. Plant J., 39: 612-628.

Ling, J., Kojima, T., Shiraiwa, M., Takahara, H. (2003) Cloning of two cysteine proteases genes, CysP1 and CysP2, from soybean cotyledons by cDNA representational difference analysis. Biochim Biophys Acta, 1627: 129-139.

Linthorst, H.J.M., Vanderdoes, C., Brederode, F.T. and Bol, J.F. (1993) Circadian expression and induction by wounding of tobacco genes for cysteine protease. Plant Mol. Biol., 21: 685-694.

Lipke, H., Fraenkel, G.S., Liener, I.E. (1954) Effects of soybean inhibitors on growth of *Tribolium confusum*. J. Agr. Food Chem., 2: 410-415.

Liu, Q., Kasuga, M., Sakuma, Y., Abe, H., Miura, S., Yamaguchi-Shinozaki, K., and Shinozaki, K. (1998) Two transcription factors, DREB1 and DREB2, with an EREBP/AP2 DNA binding domain separate two cellular signal transduction pathways in drought- and low-temperature-responsive gene expression, respectively, in *Arabidopsis*. Plant Cell, 10: 1391-1406.

Liu, J. and Zhu, J-K (1998) A calcium sensor homolog required for plant salt tolerance. Science, 280: 1943-1945.

Lohman, K.N., Gan, S., John, M.C., Amasino, R.M. (1994) Molecular analysis of natural leaf senescence in *Arabidopsis thaliana*. *Physiol. Plantarum*, 92: 322-328.

Long, S.P., Ainsworth, E.A., Alistair, R., Ort, D.R. (2004) Rising atmospheric carbon dioxide: plants FACE the future. *Ann Rev Plant Biol.*, 55: 591-628.

Long, S.P. and Drake, B.G. (1992) Photosynthetic CO₂ assimilation and rising atmospheric CO₂ concentrations. In: Baker NR and Thomas H (eds) *Crop Photosynthesis: Spatial and Temporal Determinants*, pp 69–95. Elsevier Science Publishers, Amsterdam.

Long, S.P., Zhu, X.-G., Naidu, S.L., Ort, D.R. (2006) Can improvement in photosynthesis increase crop yields? *Plant Cell Environ.*, 29: 315-330.

Lorimer, G.H. (1981) The carboxylation and oxygenation of ribulose-1,5-bisphosphate: The primary events in photosynthesis and photorespiration. *Ann. Rev. Plant Physio.*, 32: 349-382.

Ludewig, F. and Sonnewald, U. (2000) High CO₂-mediated down-regulation of photosynthetic gene transcripts is caused by accelerated leaf senescence rather than sugar accumulation. *FEBS Lett.*, 479: 19-24.

MacIntosh, S.C., Kishore, G.M., Perlak, F.J., Marrone, P.G., Stone, T.B., Sims, S.R., Fuchs, R.L. (1990) Potentiation of *Bacillus thuringiensis* insecticidal activity by serine protease inhibitors. *J. Agr. Food Chem.*, 38: 50-58.

Mae, T., Kai, N., Makino, A. and Ohira, K. (1984) Relationship between ribulose bisphosphate carboxylase content and chloroplast number in naturally senescent primary leaves of wheat. *Plant Cell Physiol.*, 25: 333-336.

Maestre, F.T., Valladres, F., Reynolds, J.F. (2005) Is the change of plant-plant interactions with abiotic stress predictable? A meta-analysis of field results in arid environments. *J. Ecol.*, 93: 748-757.

Majeau, N. and Coleman, J.R. (1996) Effect of CO₂ concentration on carbonic anhydrase and ribulose-1,5-carboxylase/oxygenase expression in pea. *Plant Physiol.*, 112: 569-574.

Makino, A., Sakuma, H., Sudo, E., Mae, T. (2003) Differences between maize and rice in N-use efficiency for photosynthesis and protein allocation. *Plant Cell Physiol.*, 44:952–956.

Manzara, T. and Gruisse, W. (1988) Organization and expression of the genes encoding ribulose-1,5-bisphosphate carboxylase in higher plants. *Photosynth. Res.*, 16: 117-139.

Margis R., Reis E.M., Villeret V. 1998. Structural and phylogenetic relationships among plant and animal cystatins. *Arch. Biochem. Biophysics*, 359: 24–30.

Marín-Navarro, J. and Moreno, J. (2006) Cysteines 449 and 459 modulate the reduction–oxidation conformational changes of ribulose 1·5-bisphosphate carboxylase/oxygenase and the translocation of the enzyme to membranes during stress *Plant Cell Environ.*, 29: 898–908.

Maroco, J.P., Edwards, G.E., Ku, M.S.B. (1999) Photosynthetic acclimation of maize to growth under elevated levels of carbon dioxide. *Planta*, 210: 115-125.

Martin, C. and Glover, B.J. (2007) Functional aspects of cell patterning in aerial epidermis. *Curr. Opin. Plant Biol.*, 10: 70–82.

Martínez, M., Rubio-Somoza, I., Carbonero, P., and Díaz, I. (2002) A cathepsin B-like cysteine protease gene from *Hordeum vulgare* (gene *CatB*) induced by GA in aleurone cells is under circadian control in leaves. *J. Exp. Bot.*, 54: 951-959.

Masle, J. (2000) The effects of elevated CO₂ concentrations on cell division rates, growth patterns, and blade anatomy in young wheat plants are modulated by factors related to leaf position, vernalization, and genotype. *Plant Physiol.*, 122: 1399–1415.

Masoud, S.A., Johnson, L.B., White, F.F. and Reeck, G.R. (1993) Expression of a cysteine protease inhibitor (oryzacystatin-I) in transgenic tobacco plants. *Plant Mol. Biol.*, 21: 655–663.

Matros, A., Amme, S., Kettig, B., Buck-Sorlin, G-H., Sonnewald, U., Mock, H-P. (2006) Growth at elevated CO₂ concentrations leads to modified profiles of secondary metabolites in tobacco cv. SamsunNN and to increased resistance against infection with *potato virus Y*. *Plant Cell Environ.*, 29: 126-137.

Matsumoto, I., Watanabe, H., Abe, K., Arai, S., Emori, Y. (1995) A putative digestive cysteine protease from *Drosophila melanogaster* is predominantly expressed in the embryonic and larval midgut. *Eur. J. Biochem.*, 227: 582-587.

Matsumoto, I., Emori, Y., Abe, K., Arai, S. (1997) Characterization of a gene family encoding cysteine proteases of *Sitophilus zeamais* (maize weevil) and analysis of the protein distribution in various tissues including alimentary tracts and germ cells. *J. Biochem.*, 121: 464-476.

Matsumoto, I., Abe, K., Arai, S., Emori, Y. (1998) Functional expression and enzymatic properties of two *Sitophilus zeamais* cysteine proteases showing different antolytic processing profiles in vitro. *J. Biochem.*, 123: 693-700.

Mayers, P.R., Kearns, P., McIntyre, K. E., Eastwood, J. A. (2002) Case study: canola tolerant of Roundup herbicide. An assessment of its substantial equivalence compared to non-modified canola. Genetically modified crops: assessing safety. Atherton, K. T. (ed). Taylor & Francis

McCouch, S. (2005) Diversifying selection in plant breeding. *PLoS Biology*, 2: e347.

McGrath, M.E. (1999) The lysosomal cysteine proteases. *Annu. Rev. Biophys. Biomol. Struct.*, 28: 181– 204.

McKee, I.F. and Woodward, F.I. (1994) CO₂ enrichment responses of wheat: interactions with temperature, nitrate and phosphate. *New Phytol.*, 127: 447–453.

McKersie, B.D., Bowley, S.R., Jones, K.S. (1999) Winter survival of transgenic alfalfa overexpressing superoxide dismutase. *Plant Physiol.*, 119: 839–848.

McKersie, B.D., Murnaghan, J., Jones, K.S., Bowley, S.R. (2000) Iron-superoxide dismutase expression in transgenic alfalfa increases winter survival without a detectable increase in photosynthetic oxidative stress tolerance. *Plant Physiol.*, 122: 1427-1438.

Mello, M.O., Tanaka, A.S., Silva-Filho, M.C. (2003) Molecular evolution of Bowman–Birk type proteinase inhibitors in flowering plants. *Mol. Phylogenet. Evol.*, 27: 103-112.

Michaud, D., Faye, L., Yelle, S. (1993a) Electrophoretic analysis of plant cysteine and serine proteases using gelatin-containing polyacrylamide gels and class-specific protease inhibitors. *Electrophoresis*, 14: 94-98.

Michaud, D., Nguyen-Quoc, B., Yelle, S. (1993b) Selective inhibition of Colorado potato beetle cathepsin H by oryzacystatins I and II. *FEBS Lett.*, 331: 173-176.

Michaud, D., Cantin, L., Vrain, T.C. (1995) Carboxy-terminal truncation of oryzacystatin II by oryzacystatin-insensitive insect digestive proteases. *Arch. Biochem. Biophys.*, 322: 469-472.

Mickel, C.E. and Standish, J. (1974) Susceptibility of processed soy flour and soy grits in storage to attack by *Tribolium castaneum*. University of Minnesota Agricultural Experimental Station Technical Bulletin: 178: 1-20.

Millenium Ecosystem Assessment 2005. Ecosystems and human well-being: Synthesis. Island Press, Washington, D.C., U.S.A. <http://www.maweb.org/proxy/Document.356.aspx>

Miller, J.E., Heagle, A.S., Pursley, W.A. (1998) Influence of ozone stress on soybean response to carbon dioxide enrichment: II. Biomass and development. *Crop Sci.*, 38: 122-128.

Minamikawa, T., Toyooka, K., Okamoto, T., Hara-Nishimura, I. and Nishimura, M. (2001) Degradation of ribulose-bisphosphate carboxylase by vacuolar enzymes of senescing French bean leaves: immunocytochemical and ultrastructural observations. *Protoplasma*, 218: 144–153.

Mitsuhashi, W., Crafts-Brandner, S.J., Feller, U. (1992) Ribulose-1,5-bisphosphate carboxylase/oxygenase degradation in isolated pea chloroplasts incubated in the light or in the dark. *J. Plant Physiol.*, 139: 653-658.

Mitsuhashi, W. and Oaks, A. (1994) Development of endopeptidase activities in maize (*Zea mays L.*) endosperms. *Plant Physiol.*, 104: 401-407.

Miyazawa, S.I., Livingston, N.J., Turpin, D.H. (2006) Stomatal development in new leaves is related to the stomatal conductance of mature leaves in poplar (*Populus trichocarpa × P. deltoides*). *J. Exp. Bot.*, 57: 373–380.

Mochizuki, A., Nishizawa, Y., Onodera, H., Tabei, Y., Toki, S., Habu, Y., Ugaki, M., and Ohashi, Y. (1999) Transgenic rice plants expressing trypsin inhibitor are resistant against rice stem borers, *Chilo suppressalis*. *Entomol. Exp. Appl.*, 93: 173-178.

Moons, A., Bauw, G., Prinsen, E., Montagu, M.V., and Straeten, D.V.D. (1995) Molecular and physiological responses to abscisic acid and salts in the roots and salt-sensitive and salt-tolerant indica rice varieties. *Plant Physiol.*, 107: 177-186.

Moore, B.D., Cheng, S-H., Rice, J., Seemann, J.R. (1998) Sucrose cycling, rubisco expression and prediction of photosynthetic acclimation to elevated atmospheric CO₂. *Plant Cell Environ.*, 21: 905-915.

Moore, B.D., Cheng, S-H., Sims, D., Seemann, J.R. (1999) The biochemical and molecular basis for photosynthetic acclimation to elevated atmospheric CO₂. *Plant Cell Environ.*, 22: 567-582.

Moore, B., Zhou, L., Rolland, F., Hall, Q., Cheng, W.H., Liu, Y.-X., Hwang, I., Jones, T., Sheen, J. (2003) Role of the *Arabidopsis* glucose sensor HXK1 in nutrient, light, and hormonal signaling. *Science*, 300: 332–36.

Morgan, J.M. (1984) Osmoregulation and water stress in higher plants. *Annu. Rev. Plant Physiol.*, 299-319.

Mott, K.A., Cardon, Z.G., Berry, J.A. (1993) Assymetric patchy stomatal closure for two surfaces of *Xanthium strumarium* L. leaves at low humidity. *Plant Cell Environ.*, 16: 25–34.

Mott, K.A., Gibson, A.C., O'Leary, J.W. (1982) The adaptive significance of amphistomatic leaves. *Plant Cell Environ.*, 5: 455–460.

Mott, K.A. and Michaelson, O. (1991) Amphistomaty as an adaptation to high light intensity in *Ambrosia cordifolia* (compositae). *Am. J. Bot.*, 78: 76–79.

Mott, K.A. and O'Leary, J.W. (1984) Stomatal behaviour and CO₂ exchange characteristics in amphistomatous leaves. *PlantPhysiol.*, 74: 47–51.

Moura, D.S., Bergey, D.R., Ryan, C.A. (2001) Characterization and localization of a wound-inducible type I serine-carboxypeptidase from leaves of tomato plants (*Lycopersicon esculentum* Mill.) *Planta*, 212: 222-230.

Mulholland, B.J., Craigon, J., Black, C.R., Colls, J.J., Atherton, J., Landon, G. (1997) Impact of elevated atmospheric CO₂ and O₃ on gas exchange and chlorophyll content in spring wheat (*Triticum aestivum* L.). *J. Exp. Bot.*, 48: 1853-1863.

Murata, N., Ishizaki-Nishizawa, O., Higashi, S., Hayashi, H., Tasaka, Y., Nishida, I. (1992) Genetically engineered alteration in the chilling sensitivity of plants. *Nature*, 356: 710-713.

Nagata, K., Kudo, N., Abe, K., Arai, S., Tanokura, M. (2000) Three-dimensional solution structure of oryzacystatin-I, a cysteine protease inhibitor of the rice, *Oryza sativa* L. *japonica*. Biochemistry-US, 39: 14753-14760.

Nakagawa, T. and Yuan, J. (2000) Cross-talk between two cysteine protease families: activation of caspase-12 by calpain in apoptosis. J. Cell Biol., 150: 887-894.

Nelson, J.M., Lane, B., Freeling, M. (2002) Expression of a mutant maize gene in ventral leaf epidermis is sufficient to signal a switch of the leaf's dorsoventral axis. Development, 129: 4581–4589.

Nguyen, H.T., Babu, R.C., and Blum, A. (1997) Breeding for drought resistance in rice: physiology and molecular genetics considerations. Crop Sci., 37: 1426-1434.

Nicklin, M.J.H. and Barrett, A.J. (1984). Inhibition of cysteine proteases and dipeptidyl peptidase I by egg-white cystatin. Biochem. J., 223: 245-253.

Nie, G., Hendrix, D.L., Webber, A.N., Kimball, B.A., Long, S.P. (1995) Increased accumulation of carbohydrates and decreased photosynthetic gene transcript levels in wheat grown at an elevated CO₂ concentration in the field. Plant Physiol., 108: 975-983.

Novitskaya, L., Trevanion, S., Driscoll, S.P., Foyer, C.H., Noctor, G. (2002) How does photorespiration modulate leaf amino acid contents? A dual approach through modelling and metabolite analysis. Plant Cell Environ., 25: 821–836.

Nowak, R. S., Ellsworth, D. S., Smith, S. D (2004) Functional responses of plants to elevated atmospheric CO₂ - do photosynthetic and productivity data from FACE experiments support early predictions? New Phytol., 162: 253-280.

Okamoto, T., Shimada, T., Hara-Nishimura, I., Nishimura, M., Minamikawa, T. (2003) C-terminal KDEL sequence of a KDEL-tailed cysteine protease (sulphydryl-endopeptidase) is involved in formation of KDEL vesicle and in efficient vacuolar transport of sulphydryl-endopeptidase. Plant Physiol., 132: 1892–1900.

Ono, K., Hashimoto, H. and Katoh, S. (1995) Changes in the number and size of chloroplasts during senescence of primary leaves of wheat grown under different conditions. *Plant Cell Physiol.*, 36: 9-17.

Padgette, S.R., N.B. Taylor, D.L. Nida, M.R. Bailey, J. MacDonald, L.R. Holden, and Fuchs, R.L. (1996) The composition of glyphosate-tolerant soybean seeds is equivalent to that of conventional soybeans. *J. Nutr.*, 126:702-716.

Pannetier, C., Gibaud, M., Couzi, P., Letan, V., Mazier, M., Tourneur, J., Hau, B. (1997) Introduction of new traits into cotton through genetic engineering: insect resistance as an example. *Euphytica* 96: 163-166.

Pareek, A., Singla, S.L., and Khush, A.K. (1997) Short term salinity and high temperature stress-associated ultrastructural alterations in young leaf cells of *Oryza sativa* L. *Ann. Bot.*, 80: 629-639.

Parry, M.A.J., Andralojc, P.J., Parmar, S., Keys, A.J., Habash, D., Paul, M.J., Alred, R., Quick, W.P., and Servaites, J.C. (1997) Regulation of Rubisco by inhibitors in the light. *Plant Cell Environ.*, 20: 528–534.

Paul, M.J. and Pellny, T.K. (2003) Carbon metabolite feedback regulation of leaf photosynthesis and development. *J. Exp. Bot.*, 54: 539-547.

Paul, M. and Stitt, M. (1993). Effects of nitrogen and phosphate deficiencies on levels of carbohydrates, respiratory enzymes, and metabolites in seedlings of tobacco, and their response to exogenous sucrose. *Plant Cell Environ.*, 16: 1047-1057.

Pechan, T., Ye, L., Chang, Y.-M, Mitra, A., Lin, L., Davis, F.M. Williams, W.P., Luthe, D.S. (2000) A unique 33-kD cysteine protease accumulates in response to larval feeding in maize genotypes resistant to fall armyworm and other Lepidoptera. *Plant Cell*, 12: 1031–1041.

Peet, M.M., Huber, S.C., Patterson, D.T. (1986) Acclimation to high carbon dioxide in monoecious cucumbers (*Cucumis sativus*): II. Carbon exchange rates, enzyme activities, and starch and nutrient concentrations. *Plant Physiol.*, 80: 63–67.

Pellegrineschi, A., Reynolds, M., Pacheco, M., Brito, R.M., Almeraya, R., Yamaguchi-Shinozaki, K., Hoisington, D. (2004) Stress-induced expression in wheat of the *Arabidopsis thaliana* DREB1A gene delays water stress symptoms under greenhouse conditions. *Genome*, 47: 493–500.

Penuelas, J. and Matamala, R. (1990) Changes in N and S leaf content, stomatal density and specific leaf area of 14 plant species during the last three centuries of CO₂ increase. *J. Exp. Bot.*, 41: 1119–1124.

Perlak, F.J., Deaton, R.W., Armstrong, T.A., Fuchs, R.L., Sims, S.R., Greenplate, J.T., Fischhoff, D.A. (1990) Insect resistant cotton plants. *Bio/Technol.*, 8: 939–943.

Perlak, F.J., Stone, T.B., Muskopf, Y.M., Petersen, L.J., Parker, G.B., McPherson, S.A., Wyman, J., Love, S., Reed, G., Biever (1993) Genetically improved potatoes: protection from damage by Colorado potato beetles. *Plant Mol. Biol.*, 22: 313–321.

Pernas, M., Sánchez-Monge, R., Salcedo, G. (2000) Biotic and abiotic stress can induce cystatin expression in chestnut. *FEBS Lett.*, 467: 206–210.

Peterson, L.W., Kleinkopf, G.E., Huffaker, R.C. (1973) Evidence for lack of turnover of ribulose-1,5-diphosphate carboxylase in barley leaves. *Plant Physiol.*, 51: 1042–1045.

Pickart, C.M. (2001) Mechanisms underlying ubiquitination. *Annu. Rev. Biochem.*, 70: 503–533.

Pingali, P.L. (ed.). 2001. CIMMYT 1999–2000 World maize facts and trends. meeting world maize needs: technological opportunities and priorities for the public sector. Mexico, D.F.: CIMMYT

Pino, M.-T., Skinner, J.S., Park, E.-J., Jeknić, Z., Hayes, P.M., Tomashow, M.F. Chen, T.H.H. (2007) Use of a stress inducible promoter to drive ectopic *AtCBF* expression improves potato freezing tolerance while minimizing negative effects on tuber yield. *Plant Biotech. J.*, 5: 591-604.

Pons, L. (2003) Pumping iron into western Africa's corn. *Agricultural Research (USDA)*.
<http://www.ars.usda.gov/is/AR/archive/apr03/iron0403.pdf>

Pontius, J.U., Wagner, L., Schuler, G.D. (2003) UniGene: a unified view of the transcriptome. In: *The NCBI Handbook*. Bethesda (MD): National Center for Biotechnology Information. 11pp.

Poorter, H. and Navas, M.-L. (2003) Plant growth and competition at elevated CO₂: on winners, losers and functional groups. *New Phytol.*, 157: 175–198.

Porter, M.A. and Grodzinski, B. (1984) Acclimation to high CO₂ in bean. Carbonic anhydrase and ribulose bisphosphate carboxylase. *Plant Physiol.*, 74: 413-416.

Portis, A.R. Jr. (2003) Rubisco activase – Rubisco's catalytic chaperone. *Photosynth. Res.*, 75: 11-27.

Puntervoll, P., Linding, R., Gemund, C., Chabanis-Davidson, S., Mattingsdal, M., Cameron, S., Martin, D. M., Ausiello, G., Brannetti, B., Costantini, A. Ferrè, F., Maselli, V., Via, A., Cesareni, G., Diella, F., Superti-Furga, G., Wyrwicz, L., Ramu, C., McGuigan, C., Gudavalli, R., Letunic, I., Bork, P., Rychlewski, L., Küster, B., Helmer-Citterich, M., Hunter, W.N., Aasland, R., Gibson, T.J. (2003). ELM server: A new resource for investigating short functional sites in modular eukaryotic proteins. *Nucleic Acids Res.*, 31: 3625-3630.

Quirino, B.F., Normanly, J. and Amasino, R.M. (1999) Diverse range of gene activity during *Arabidopsis thaliana* leaf senescence includes pathogen-independent induction of defense-related genes. *Plant Mol. Biol.*, 40: 267-278.

Raines, C.A., Horsnell, P.R., Holder, C., Lloyd, J.C. (1992) *Arabidopsis thaliana* carbonic anhydrase: cDNA sequence and effect of CO₂ on mRNA levels. *Plant Mol. Biol.*, 20:1143–1148.

Ramagli, L.S. (1999) Quantifying protein in 2-D PAGE solubilization buffers. In: *Methods in Molecular Biology*, 112: 99-103, Link, A.J. (ed), Humana Press Inc, Totowa, NJ.

Reichard, U., Cole, G.T., Hill, T.W., Ruchel, R., Monod, M. Molecular characterization and influence on fungal development of ALP2, a novel serine proteinase from *Aspergillus fumigatus* (2000) *Int. J. Med. Microbiol.*, 290: 549-558.

Repelin, A., Baga, M., Jauhar, P.P., Chibbar, R.N. (2001) Genetic enrichment of cereal crops via alien gene transfer: New challenges. *Plant Cell Tiss. Org.*, 64: 159-183.

Rintamäki, E., Keys, A.J. and Parry, M.A.J. (1988) Comparison of the specific activity of ribulose-1,5-bisphosphate carboxylase-oxygenase from some C3 and C4 plants. *Physiol. Plant.*, 74: 326–331.

Rivard, D., Anguenot, R., Brunelle, F., Le, V.Q., Vézina, L-P., Trépanier, S., Michaud, D. (2006). An in-built protease inhibitor system for the protection of recombinant proteins recovered from transgenic plants. *Plant Biotech. J.*, 4, 359-368.

Robertson, E.J. and Leech, R.M. (1995) Significant changes in cell and chloroplast development in young wheat leaves (*Triticum aestivum* cv. Hereward) grown in elevated CO₂. *Plant Physiol.*, 107: 63–71.

Robertson, E.J., Williams, M., Harwood, J.L., Lindsay, J.G., Leaver, C.J., Leech, R.M. (1995) Mitochondria increase three-fold and mitochondrial proteins and lipid change dramatically in post-meristematic cells in young wheat leaves grown at elevated CO₂. *Plant Physiol.*, 108: 469–474.

Rodermel, S. (1999) Subunit control of Rubisco biosynthesis – a relic of an endosymbiotic past? *Photosynth. Res.*, 59: 105-123.

Rogers, A., Ellsworth, D.S., Humphries, S.W. (2001) Possible explanation of the disparity between the *in vitro* and *in vivo* measurements of Rubisco activity: a study in loblolly pine grown in elevated $p\text{CO}_2$. *J. Exp. Bot.*, 52: 1555-1561.

Rogers, H.H., Runion, G.B., Krupa, S.V. (1994) Plant responses to atmospheric CO_2 enrichment with emphasis on roots and the rhizosphere. *Environ. Pollution*, 83: 155-189.

Roitsch, T., Balibrea, M.E., Hofmann, M., Proels, R., Sinha, A. K. (2003) Extracellular invertase: key metabolic enzyme and PR protein. *J. Exp. Bot.*, 54: 513-524.

Rojo, E., Martín, R., Carter, C., Zouhar, J., Pan, S., Plotnikova, J., Jin, H., Paneque, M., Sánchez-Serrano, J.J., Baker, B., Ausubel, F.M., Raikhel, N.V. (2004) VPE \square exhibits a caspase-like activity that contributes to defense against pathogens. *Curr. Biol.*, 14: 1897-1906.

Romeis, J., Meissle, M., Bigler, F. (2006) Transgenic crops expressing *Bacillus thuringiensis* toxins and biological control. *Nat. Biotechnol.*, 24: 63-71.

Romero, C., Belles, J.M., Vayá, J.L., Serrano, R., Culiañez-Maciá, F.A. (1997) Expression of the yeast trealose-6-phosphate synthase gene in transgenic tobacco plants: pleiotropic phenotypes include drought tolerance. *Planta* 201: 293-297.

Roulin, S. and Feller, U. (1998) Light-independent degradation of stromal proteins in intact chloroplasts isolated from *Pisum sativum* L. leaves: requirement for divalent cations. *Planta*, 205: 297-304.

Rouquié, D., Peltier, J.B., Marquis-Mansion, M., Tournaire, C., Doumas, P., Rossignol, M. (1997) Construction of a directory of tobacco plasma membrane proteins by combined two-dimensional gel electrophoresis and protein sequencing. *Electrophoresis*, 18: 654-660.

Rowland-Bamford, A.J., Baker, J.T., Allen, L.H.J., Bowes, G. (1991) Acclimation of rice to changing atmospheric carbon dioxide concentration. *Plant Cell Environ.*, 14: 577-583.

Ryan, C.A. (1989) Insect-induced chemical signals regulating natural plant protection responses. In: Denno, R.F. and McClure, M.S., eds. *Variable plants and herbivores in natural and managed systems*. New York, Academic Press, pp.43-60.

Ryan, C.A. (1990) Protease inhibitors in plants: genes for improving defenses against insects and pathogens. *Annu. Rev. Phytopathol.*, 28: 425-449.

Sage, R.F., Pearcy, R.W., Seemann, J.R. (1987) The nitrogen use efficiency of C₃ and C₄ plants. III. Leaf nitrogen effects on the activity of carboxylating enzymes in *Chenopodium album* (L.) and *Amaranthus retroflexus* (L.). *Plant Physiol.*, 85: 355–359.

Sage, R.F., Sharkey, T.D., Seeman, J.R. (1988) The *in vivo* response of the ribulose-1,5-bisphosphate carboxylase activation state and the pool sizes of photosynthetic metabolites to elevated CO₂ in *Phaseolus vulgaris* L. *Planta*, 174: 407-16.

Sage, R.F., Sharkey, T.D., Seeman, J.R. (1989) Acclimation of photosynthesis to elevated CO₂ in five C₃ species. *Plant Physiol.*, 89, 590-596.

Sage, R.F. (1994) Acclimation of photosynthesis to increasing atmospheric CO₂: The gas exchange perspective. *Photosynth. Res.*, 39: 351-368.

Sage, R. F. (2004) The evolution of C₄ photosynthesis. *New Phytol.*, Vol. 161: 341-370.

Sakamoto, W. (2006) Protein degradation machineries in plastids. *Annu Rev. Plant Biol.*, 57: 599-621.

Salisbury, E.J. (1927) On the causes and ecological significance on stomatal frequency with special reference to woodland flora. *Philos. T. R. Soc. B*, 216: 1–65.

Salvucci, M.E. and Ogren, W.L. (1996) The mechanism of Rubisco activase: Insights from studies of the properties and structure of the enzyme. *Photosynth. Res.*, 47: 1-11.

Sambrook, J., Fritsch, E.F., Maniatis, T. (1989) Molecular Cloning, 2nd edn. Cold Spring Harbor Laboratory Press, New York.

Sanmartín, M., Jaroszewski, L., Raikhel, N.V., Rojo, E. (2005) Caspases. Regulating death since the origin of life. *Plant Physiol.*, 137: 841-847.

Schaffer, M.A. and Fischer, R.L. (1988) Analysis of mRNAs that accumulates in response to low temperature identifies a thiol protease gene in tomato. *Plant Physiol.*, 87: 431-436.

Schaller, A. (2004) A cut above the rest: the regulatory function of plant proteases. *Planta*, 220: 183-197.

Schmid, M., Simpson, D, and Gietl, C. (1999) Programmed cell death in castor bean endosperm is associated with the accumulation and release of a cysteine endopeptidase from ricinosomes. *P. Natl. Acad. Sci. USA*, 96: 14159–14164.

Schmidt, G.W. and Mishkind, M.L. (1983) Rapid degradation of unassembled ribulose 1,5-biphosphate carboxylase small subunits in chloroplasts. *P. Natl. Acad. Sci. USA*, 80: 2632–2636.

Schwanz, P. and Polle, A. (2001) Differential stress responses of antioxidative systems to drought in pendunculate oak (*Quercus robur*) and maritime pine (*Pinus pinaster*) grown under high CO₂ concentrations. *J. Exp. Bot.*, 52: 133-143.

Sharma, H.C., Sharma, K.K., Crouch, J.H. (2004) Genetic transformation of crops for insect resistance: potential and limitations. *Crit. Rev. Plant Sci.*, 23: 47-72.

Shen, B., Jensen, R.G., and Bohnert, H.J. (1997) Increased resistance to oxidative stress in transgenic plants by targeting mannitol biosynthesis to chloroplasts. *Plant Physiol.*, 113: 1177-1183.

Shi, Y. (2002) Mechanisms of caspase activation and inhibition during apoptosis. *Mol. Cell*, 9: 459-470.

Shimada, T., Hiraiwa, N., Nishimura, M. and Hara-Nishimura, I. (1994) Vacuolar processing enzyme of soybean that converts proproteins to the corresponding mature forms. *Plant Cell Physiol.*, 35: 713-718.

Shinozaki, K. and Yamaguchi-Shinozaki, K. (2000) Molecular responses to dehydration and low temperature: differences and cross-talk between two stress signaling pathways. *Curr. Opin. Plant Biol.*, 3: 217-223.

Sicher, R.C., Kremer, D.F., Rodermel, S.R. (1994) Photosynthetic acclimation to elevated CO₂ occurs in transformed tobacco with decreased ribulose-1,5-bisphosphate carboxylase/oxygenase content. *Plant Physiol.*, 104: 409–415.

Sims, D.A. and Gamon, J.A. (2002) Relationships between leaf pigment content and spectral reflectance across a wide range of species, leaf structures and developmental stages. *Remote Sens. Environ.*, 81: 337-354.

Smart, D.R., Chatterton, N.J., Bugbee, B. (1994) The influence of elevated CO₂ on non-structural carbohydrate distribution and fructan accumulation in wheat canopies. *Plant Cell Environ.*, 17: 435–442.

Soares, A.S., Driscoll, S.P., Olmos, E., Harbinson, J., Arrabaça, M.C., Foyer, C.H. (2008) Adaxial/abaxial specification in the regulation of photosynthesis and stomatal opening with respect to light orientation and growth with CO₂ enrichment in the C₄ species Paspalum dilatatum. *New Phytol.*, 177: 186-198.

Solomon, M., Belenghi, B., Delledonne, M., Menachem, E., Levine, A. (1999) The involvement of cysteine proteases and protease inhibitor genes in the regulation of programmed cell death in plants. *Plant Cell*, 11: 431-443.

Spreitzer, R.J. and Salvucci, M.E. (2002) Rubisco: structure, regulatory interactions, and possibilities for a better enzyme. *Annu. Rev. Plant Biol.*, 53: 449-475.

Stitt, M. (1991) Rising CO₂ levels and their potential significance for carbon flow in photosynthetic cells. *Plant Cell Environ.*, 14: 741-762.

Strauss, S.H., Knowe, S.A., and Jenkins, J.J. (1997) Benefits and risk of transgenic, Roundup Ready® cottonwoods. *J. For.*, 95: 12–19.

Sugiyama, T., Mizuno, M., Hayashi, M (1984) Partitioning of nitrogen among ribulose-1,5-bisphosphate carboxylase/oxygenase, phosphoenolpyruvate carboxylase, and pyruvate orthophosphate dikinase as related to biomass productivity in maize seedlings. *Plant Physiol.*, 75:665–669.

Sullivan, J.A., Shirasu, K., Deng, X.W. (2003) The diverse roles of ubiquitin and the 26S proteasome in the life of plants. *Nat. Rev. Genet.*, 4: 948-958.

Sun, J., Gibson, K.M., Kuirats, O., Okita, T.W., Edwards, G.E. (2002) Interactions of nitrate and CO₂ enrichment on growth, carbohydrates, and Rubisco in *Arabidopsis* starch mutants. Significance of starch and hexose. *Plant Physiol.*, 130: 1573–1583.

Swidzinski, J.A., Sweetlove, L.J. and Leaver, C.J. (2002) A custom microarray analysis of gene expression during programmed cell death in *Arabidopsis thaliana*. *Plant J.*, 30: 431–446.

Tajima, H. and Koizumi, N. (2006) Induction of BiP by sugar independent of a *cis*-element for the unfolded protein response in *Arabidopsis thaliana*. *Biochem. Biophys. Res. Co.*, 346: 926-930.

Tanaka, H., Onouchi, H., Kondo, M., Hara-Nishimura, I., Nishimura, M., Machida, C., Machida, Y. (2001) A subtilisin-like serine protease is required for epidermal surface formation in *Arabidopsis* embryos and juvenile plants. *Development*, 128: 4681-4689.

Tarczynski, M.C., Jensen, R.G., and Bohnert, H.J. (1993) Stress protection of transgenic tobacco by production of osmolyte mannitol. *Science*, 259: 508-510.

Taylor, G., Ranasinghe, S., Bosac, C., Gardner, S.D.L., Ferris, R. (1994) Elevated CO₂ and plant growth: cellular mechanisms and responses of whole plants. *J. Exp. Bot.*, 45: 1761–1774.

Taylor,, G., Tallis, M.J., Giardina, C.P.Percy, K.E., Miglietta, F., Gupta, P.S., Gioli, B., Calfapietra, C., Gielen, B., Kubiske, M.E., Scarascia-Mugnozza, G.E., Kets, K., Long, S.P., Karnosky, D.F. (2008) Future atmospheric CO₂ leads to delayed autumnal senescence. *Global Change Biol.*, 14: 264–275.

Tissue, D.T., Thomas, R.B., Strain, B.R. (1993) Long-term effects of elevated CO₂ and nutrients on photosynthesis and rubisco in loblolly pine seedlings. *Plant Cell Environ.*, 16: 859-865.

Toyooka, K., Okamoto, T., Minamikawa, T. (2000) Mass transport of proform of a KDEL-tailed cysteine protease (SH-EP) to protein storage vacuoles by endoplasmic reticulum–derived vesicle is involved in protein mobilization in germinating seeds. *J. Cell Biol.*, 148: 453-464.

Trobacher, C.P., Senatore, A., Greenwood, J.S. (2006) Masterminds or minions? Cysteine proteinases in plant programmed cell death. *Can. J. Bot.*, 84: 651-667.

Tuba, Z., Szente, K., Kock, J. (1994) Response of photosynthesis, stomatal conductance, water use efficiency and production to long-term elevated CO₂ in winter wheat. *J. Plant Physiol.*, 144: 661–668.

Turk, V., Turk, B., Turk, D. (2001) Lysosomal cysteine proteases: facts and opportunities. *EMBO J.* 20: 4629–4633.

Ueda, T., Seo, S., Ohashi, Y. and Hashimoto, J. (2000) Circadian and senescence-enhanced expression of a tobacco cysteine protease gene. *Plant Mol. Biol.*, 44: 649–657.

Uren, A.G., O'Rourke, K., Aravind, L., Pisabarro, M.T., Seshagiri, S., Koonin, E.V., Dixit, V.M. (2000) Identification of paracaspases and metacaspases two ancient families of caspase-like proteins, one of which plays a key role in MALT lymphoma. *Mol. Cell*, 6: 961-967.

Urwin, P.E., Atkinson, H.J., Waller, D.A., McPherson, M.J. (1995) Engineered oryzacystatin-I expressed in transgenic hairy roots confers resistance to *Globodera pallida*. Plant J., 8: 121-131.

Urwin, P.E., Lilley, C.J., McPherson, M.J., Atkinson, H.J. (1997) Resistance to both cyst and rootnot nematodes conferred by transgenic *Arabidopsis* expressing a modified plant cystatin. Plant J. 12: 455-461.

Ussuf, K.K., Laxmi, N.H., Mitra, R. (2001) Protease inhibitors: Plant-derived genes of insecticidal protein for developing insect-resistant transgenic plants. Curr. Sci., 80: 847-853.

Vain, P., Worland, B., Clarke, M.C., Richard, G., Beavis, M., Liu, H., Kohli, A., Leech, M., Snape, J., Christou, P. (1998) Expression of an engineered cysteine protease inhibitor (Oryzacystatin-I delta D86) for nematode resistance in transgenic rice plants. Theor. Appl. Genet., 96: 266-271.

Van Breusegem, F., Slooten, L., Stassart, J.-M., Moens, T., Boterman, J., Van Montagu, M., Inzé, D. (1999) Overproduction of *Arabidopsis thaliana* FeSOD confers oxidative stress tolerance to transgenic maize. Plant Cell Physiol., 40: 515-523.

Vanderauwera, S., De Block, M., Van de Steene, N., Van de Cotte, B., Metzlaff, M., Van Breusegem, F. (2007) Silencing of poly(ADP-ribose) polymerase in plants alters abiotic stress signal transduction. P. Natl. Acad. Sci. USA, 104: 15150-15155.

Van der Vyver, C., Schneidereit, J., Driscoll, S., Turner, J., Kunert, K., and Foyer, C.H. (2003) Oryzacystatin I expression in transformed tobacco produces a conditional growth phenotype and enhances chilling tolerance. Plant Biotech. J., 1: 101-112.

Van Doorn, W.G. and Woltering, E.J. (2005) Many ways to exit? Cell death categories in plants. Trends Plant Sci., 10: 117-122.

Van Heerden, P.D.R., Krüger, G.G.J., Loveland, J.E., Parry, M.A.J., Foyer, C.H. (2003) Dark chilling imposes metabolic restrictions on photosynthesis in soybean. *Plant Cell Environ.*, 26: 323–337.

Van Oosten, J-J. and Besford, R.T. (1996) Acclimation of photosynthesis to elevated CO₂ through feedback regulation of gene expression: Climate of opinion. *Photosynth. Res.*, 48: 353-365.

Van Oosten, J-J., Wilkins, D., Besford, R.T. (1994) Regulation of the expression of photosynthetic nuclear genes by high CO₂ is mimicked by carbohydrates: a mechanism for the acclimation of photosynthesis to high CO₂. *Plant Cell Environ.*, 17: 913-923.

Vercammen, D., Belenghi, B., Van de Cotte, B., Beunens, T., Gavigan, J.-A., De Rycke, R., Brackenier, A., Inzé, D., Hariies, J.L., Van Breusegem, F. (2006) Serpin 1 of *Arabidopsis thaliana* is a suicide inhibitor for metacaspase 9. *J. Mol. Biol.*, 364: 625-636.

Vernon, L.P. (1960). Spectrophotometer determination of chlorophylls and pheophytins in plant extracts. *Anal. Chem.* 32: 1144–1150.

Vernon, D.M., Tarczynski, M.C., Jensen, R.G., Bohnert, H.J. (1993) Cyclitol production in transgenic tobacco. *Plant J.*, 4: 199-205.

Vierstra, R.D. (1996) Proteolysis in plants: mechanisms and functions. *Plant Mol. Biol.*, 32: 275-302.

Vierstra, R.D. (2003) The ubiquitin/26S proteasome pathway, the complex last chapter in the life of many plant proteins. *Trends Plant Sci.*, 8, 135-142.

Vinocur, B. and Altman, A. (2005) Recent advances in engineering plant tolerance to abiotic stress: achievements and limitations. *Curr. Opin. Biotech.*, 16: 123-132.

Virgin, I., Salter, A.H., Ghanotakis, D.F., Andersson, B. (1991) Light-induced D1 protein degradation is catalyzed by a serine type protease. *FEBS Lett.*, 281: 125 128.

Vitale, A. and Pedrazzini, E. (2005) Recombinant pharmaceuticals from plants: the plant endomembrane system as bioreactor. *Mol. Interv.*, 5: 216-225.

Voges, D., Zwickl, P., Baumeister, W. (1999) The 26S proteasome: A molecular machine designed for controlled proteolysis. *Annu. Rev. Biochem.*, 68: 1015-1068.

Von Caemmerer, S. and Farquhar, G.D. (1981) Some relationships between the biochemistry of photosynthesis and the gas exchange of leaves. *Planta*, 153: 376-387.

Von Caemmerer, S., Ghannoum, O., Conroy, J.P., Clark, H., Newton, P.C.D. (2001) Photosynthetic responses of temperate species to free air CO₂ enrichment (FACE) in a grazed New Zealand pasture. *Aust. J. Plant Physiol.*, 28: 439–450.

Vu, C.V., Allen, L.H. Jr., Bowes, G. (1983) Effects of light and elevated atmospheric CO₂ on the ribulose bisphosphate carboxylase activity and ribulose bisphosphate level of soybean leaves. *Plant Physiol.*, 73: 729-734.

Vu, J.C.V., Allen Jr., L.H., Bowes, G. (1989) Leaf ultrastructure, carbohydrates and protein of soybeans grown under CO₂ enrichment. *Environ. Exp. Bot.*, 29: 141–147.

Wagstaff, C., Leverentz, M.K., Griffiths, G., Thomas, B., Chanasut, U., Stead, A.D., Rogers, H.J. (2002) Cysteine protease gene expression and proteolytic activity during senescence of *Alstroemeria* petals. *J. Exp. Bot.*, 53: 233-240.

Walker-Simmons, M. and Ryan, C.A. (1980) Isolation and properties of carboxypeptidase from leaves of wounded tomato plants. *Phytochemistry*, 19: 43-47.

Walting, J.R. and Press, M.C. (1997) How is the relationship between the C₄ cereal *Sorghum bicolor* and the C₃ root hemi-parasites *Striga hermonthica* and *Striga asiatica* affected by elevated CO₂? *Plant Cell Environ.*, 20: 1292–1300.

Wand, S.J.E., Midgley, G.F., Jones, M.H., Curtis, P.S. (1999) Responses of wild C₄ and C₃ grass (*Poaceae*) species to elevated CO₂ concentration: a meta-analytic test of current theories and perceptions. *Glob. Change Biol.*, 5: 723–741.

Wand, S.J.E., Midgley, G.F., Stock, W.D. (2001) Growth responses to elevated CO₂ in NADP-ME, NAD-ME and PCK C₄ grasses and a C₃ grass from South Africa. Aust. J. Plant Physiol., 28: 13–25.

Ward J.K., Tissue D.T., Thomas R.B., Strain B.R. (1999) Comparative responses of model C₃ and C₄ plants to drought in low and elevated CO₂. Glob. Change Biol., 5: 857-867.

Watanabe, N. and Lam, E. (2004) Recent advance in the study of caspase-like proteases and Bax inhibitor-1 in plants: their possible roles as regulator of programmed cell death.

Watling, J.R., Press, M.C., Quick, W.P. (2000) Elevated CO₂ induces biochemical and ultrastructural changes in leaves of the C₄ cereal sorghum. Plant Physiol., 123: 1143–1152.

Webber, A.N., Nie, G.-Y., Long, S.P. (1994) Acclimation of photosynthetic proteins to rising atmospheric CO₂. Photosynth. Res., 39: 413-425.

Werneke, J.M., Chatfield, J.M. and Ogren, W.L. (1989) Alternative mRNA splicing generates the two ribulosebisphosphate carboxylase/oxygenase activase polypeptides in spinach and Arabidopsis. Plant Cell, 1: 815-825.

Winder, T.L., Anderson, J.C., Spalding, M.H. (1992) Translational regulation of the large and small subunits of ribulose bisphosphate carboxylase/oxygenase during induction of the CO₂ concentrating mechanism in *Chlamydomonas reinhardtii*. Plant Physiol. 98: 1409–1414.

Wingler, A., Purdy, S., MacLean, J.A., Pourtau, N. (2006) The role of sugars in integrating environmental signals during the regulation of leaf senescence. J. Exp. Bot., 57: 391-399.

Winzeler, M., Dubois, D., Nosberger, J. (1990) Absence of fructan degradation during fructan accumulation in wheat stems. J. Plant Physiol., 136: 324–329.

Woltering, E.J., Van der Bent, A., Hoeberichts, F.A. (2002) Do plant caspases exist? *Plant Physiol.*, 130: 1764-1769.

Woodward, F.I. (1987) Stomatal numbers are sensitive to increases in CO₂ from pre-industrial levels. *Nature*, 327: 617-18.

Woodward, F.I. (2002) Potential impacts of global elevated CO₂ concentrations on plants. *Curr. Opin. Plant Biol.*, 5: 207–211.

Woodward, F.I. and Kelly, C.K. (1995) The effect of CO₂ concentration on stomatal density. *New Phytol.*, 131, 311-327.

Woodward, F. I J., Lake, A, Quick, W. P. (2002) Stomatal development and CO₂: ecological consequences. *New Phytol.*, 153: 477–484.

Wullschleger, S. D., Norby, R. J. Gunderson, C. A. (1992) Growth and maintenance respiration in leaves of *liriodendron tulipifera* L. exposed to long-term carbon dioxide enrichment in the field. *New Phytol.*, 121: 515-523.

Xiong, Y., Contento, A.L., Nguyen, P.Q., Bassham, D.C. (2007) Degradation of oxidised proteins by autophagy during oxidative stress in *Arabidopsis*. *Plant Physiol.*, 143: 291–299.

Xu, X-F. and Chye, L-M. (1999) Expression of cysteine protease during developmental events associated with programmed cell death in brinjal. *Plant J.*, 17: 321-328.

Yamada, T., Ohta, H., Shinohara, A., Iwamatsu, A., Shimada, H., Tsuchiya, T., Masuda, T., and Takamiya, K. (2000) A cysteine protease from maize isolated in a complex with cystatin. *Plant Cell Physiol.* 41: 185-191.

Yamauchi, M., Aguilar, A.M., and Vaughan, D.A. (1993) Seshu, Rice (*Oryza sativa* L.) germplasm suitable for direct sowing under flooded soil surface. *Euphytica*, 67: 177-184.

Yancey, P.H., Clark, M.E., Hand, S.C., Bowlus, R.D., and Somero, G.N. (1982) Living with water stress: evolution of osmolyte systems. *Science*, 217: 1214-1222.

Yang, D.-H., Webster, J., Adam, Z., Lindahl, M., Andersson, B. (1998) Induction of acclimative proteolysis of the light-harvesting chlorophyll a/b protein of photosystem ii in response to elevated light intensities. *Plant Physiol.*, 118: 827-834.

Ye, G.-N., Hajdukiewicz, P.T.J., Broyles, D., Rodriguez, D., Xu, C.W., Nehra, N., Staub, J.M. (2001) Plastid-expressed 5-enolpyruvylshikimate-3-phosphate synthase genes provide high level glyphosate tolerance in tobacco. *Plant J.*, 25: 261–270.

Yoshida, T. and Minamikawa, T. (1996) Successive amino-terminal proteolysis of the large subunit of ribulose 1,5-bisphosphate carboxylase/oxygenase by vacuolar enzymes from french bean leaves. *Eur. J. Biochem.*, 238: 317–324.

Zhang, N., Kallis, R.P., Ewy, R.G., Portis, Jr., A.R. (2002) Light modulation of RuBisCO in Arabidopsis requires a capacity for redox regulation of the larger RuBisCO activase isoform. *P. Natl. Acad. Sci. USA*, 99: 3330-3334.

Zhang, H. and Nobel, P.S. (1996) Photosynthesis and carbohydrate partitioning for the C₃ desert shrub *Encelia farinosa* under current and doubled CO₂ concentrations. *Plant Physiol.*, 110: 1361-1366.

Zhao, Y., Botella, M.A., Subramanian, L., Niu, X., Nielsen, S.S., Bressan, R.A. Hasegawa, P.M. (1996). Two wound-inducible soybean cysteine protease inhibitors have greater insect digestive protease inhibitory activities than a constitutive homologue. *Plant Physiol.*, 111: 1299- 1306.

Zhu, J., Goldstein, G., Bartholomew, D.P. (1999) Gas exchange and carbon isotope composition of Ananas comosus in response to elevated CO₂ and temperature. *Plant Cell Environ.*, 22: 999-1007.



Zimmermann, P., Hirsch-Hoffmann, M., Hennig, L., Gruissem, W. (2004) GENEVESTIGATOR. Arabidopsis microarray database and analysis toolbox. Plant Physiol., 136: 2621-2632.

Ziska, L.H. and Bunce, J.A. (1999) Effect of elevated carbon dioxide concentration at night on the growth and gas exchange of selected C4 species. Aust. J. Plant Physiol., 26: 71-77.