



## References

- Alsaadawi, I. S., 2001. Allelopathic influence of decomposing wheat residues in agroecosystem. *Journal of Crop Production*. 4, 185-196.
- Alsaadawi, I.S., Al-Ekelle, M.H.S. & Al-Hamzawi, M.K., 2005. Allelopathic potential of *Sorghum bicolor* L. (Moench) genotypes against weeds. *Proceedings of the 4th World Congress on Allelopathy*, August 2005, Wagga Wagga, Australia. 254-257.
- Alsaadawi, I. S., Zwain, K. H. Y. & Shahata, H.A., 1998. Allelopathic inhibition of growth of rice by wheat residues. *Allelopathy Journal*. 5, 163-169.
- An, M., Pratley, J.J.E. & Haig, T.T., 2001. Phytotoxicity of *Vulpia* residues: IV. Dynamics of allelochemicals during decomposition of *Vulpia* residues and their corresponding phytotoxicity. *Journal of Chemical Ecology*. 27, 395-409.
- Ash, G.J., Raman, R., & Crump, N.S., 2003. An investigation of genetic variation in *Carthamus lanatus* in New South Wales, Australia, using inter-simple sequence repeats (ISSR) analysis. *Weed Research* 43, 208-213.
- Ashton, F.M. & Monaco, T.J., 1991. *Weed Science Principles and Practices*. John Wiley & Sons, New York.
- Bacilio-Jimenez M, Aguilar-Flores S, Ventura-Zapata E, Perez-Campos E, Bouquelet S & Zenteno E (2003) Chemical characterization of root exudates from rice (*Oryza sativa*) and their effects on the chemotactic response of endophytic bacteria. *Plant and Soil* 249, 271-277.
- Baghestani, A., Lemieux, C., Leroux, G.D., Baziramakenga, R. & Simard, R.R., 1999. Determination of allelochemicals in spring cereal cultivars of different competitiveness. *Weed Science* 47, 498-504.
- Bais, H.P., Vepachedu, R., Gilroy, S., Callaway, R.M. & Vivanco, J.M., 2003. Allelopathy and exotic plant invasion: from molecules and genes to species interactions.

Science 301, 1377-1380.

Balfourier, F., Imbert, C. & Charmet, G., 2000. Evidence for phylogeographic structure in *Lolium* species related to the spread of agriculture in Europe: a cpDNA study. *Theoretical and Applied Genetics* 101, 131-138.

Barberi, P., 2002. Weed management in organic agriculture: are we addressing the right issues? *Weed Research* 42, 177-193.

Bastiaans, L., 2008. Focus on ecological weed management: what is hindering adoption. *Weed Research* 48, 481-491.

Batish, D.R., Singh, H.P., Kohli, R.K. & Kaur, S., 2001. Crop allelopathy and its role in ecological agriculture. *Journal of Crop Production* 4, 121-161.

Batish, D.R., Tung, P., Singh, H.P. & Kohli, R.K., 2002. Phytotoxicity of sunflower residues against some summer season crops. *Journal of Agronomy and Crop Science* 188, 19-24.

Baucom, R.S., 2009. A herbicide defense trait that is distinct from resistance: The evolutionary ecology and genomics of herbicide tolerance. In: Stewart, C.N., Jr. (ed.), *Weedy and Invasive Plant Genomics*. Wiley-Blackwell, Iowa, USA.

Belz, R.G., 2004. Evaluation of allelopathic traits in *Triticum* L. spp. and *Secale cereale* L. PhD Thesis, University of Hohenheim, Stuttgart, Germany.

Belz, R.G., 2007. Allelopathy in crop/weed interactions – an update. *Pest Management Science* 63, 308-326.

Belz, R.G., 2008. Stimulation versus inhibition-bioactivity of parthenin, a phytochemical from *Parthenium hysterophorus* L. *Dose response* 6, 80-96.

Belz, R.G., van der Laan, M., Reinhardt, C.F. & Hurle, K., 2009. Soil degradation of parthenin—does it contradict the role of allelopathy in the invasive weed

*Parthenium hysterophorus* L.? *Journal of Chemical Ecology* 35, 1137-1150.

Ben-Hammouda, M., Ghorbal, H., Kreme, R.J. & Oueslati, O., 2001. Allelopathic effects of barley extracts on germination and seedling growth of bread and durum wheats. *Agronomie* 21, 65-71.

Bertholdsson, N.O., 2004. Variation in allelopathic activity over 100 years of barley selection and breeding. *Weed Research* 44, 78-86.

Bertholdsson, N.O., 2005. Early vigour and allelopathy – two useful traits for enhanced barley and wheat competitiveness against weeds. *Weed Research* 45, 94-102.

Bertin, C., Yang, X. & Weston, L.A., 2003. The role of root exudates and allelochemicals in the rhizosphere. *Plant and Soil* 256, 67-83.

Bhatia, R.K., Gill, H.S., Bhandari, S.C. & Khurana, A.S., 1984. Allelopathic interaction of some tropical weeds. *Indian Journal of Weed Science* 16, 182-189.

Blum, U., 1996. Allelopathic interactions involving phenolic acids. *Journal of Nematology* 28, 259-267.

Bouwmeester, H.J. & Karssen, C.M., 1993. Seasonal periodicity in germination of seeds of *Chenopodium album* L. *Annals of Botany* 72, 463-473.

Bosendorf, O., Auge, H., Lafuma, L., Roger, W.E., Siemann, E. & Prati, D., 2005. Phenotypic and genetic differentiation between native and introduced plant populations. *Oecologia* 144, 1-11.

Breland, T.A., 1996. In: Physiology of stressed crops. Volume III. The Stress of Allelochemicals 2005. U.S. Gupta (ed.), University of Georgia, Science Publishers, Inc., Enfield (NH), USA.

Brecke, B.J., Shilling, D.G., 1996. Effect of crop species, tillage, and rye (*Secale cereale*) mulch on sicklepod (*Senna obtusifolia*). *Weed Science* 44, 133-136.

- Burgess, L.W., Backhouse, D., Summerell, B.A. & Swan, L.J., 2001. Crown rot of wheat. *In: Summerell, B.A., Leslie, J.F., Backhouse, D., Bryden, W.L., Burgess, L.W. (eds.), Fusarium – Paul Nelson Memorial Symposium, American Phytopathological Society, St Paul, Minnesota, USA.*
- Calabrese, E.J., 2007. Biological stress response terminology: integrating the concepts of adaptive and preconditioning stress within a hormetic dose-response framework. *Toxicology Applied in Pharmacology* 222, 122-128.
- Callaway, R.M. & Aschehoug, E., 2000. Invasive plants versus their new and old neighbors: a mechanism for exotic invasion. *Science* 290, 521-523.
- Caussanel, J.P., 1979. Effects non compétitifs entre le chenopode blanc (*Chenopodium album* L.) de le maïs (INRA 258). *Weed Research* 19, 123-135.
- Caussanel, J.P., Martin, J., Annelle, D., Peiento, M., Vallet, J.K. & Kurnej, K., 1977. Inhibition of growth of young radicles of maize by exudation in culture solutions and extracts of ground roots of *Chenopodium album* L. *In: A M Grodzinsky (ed.) Interaction of plants and microorganisms in phytocenoses. Naukova Dumka* (In Russian, English summary).
- Chao, W.S., Horvath, D.P., Anderson, J.V. & Foley, M., 2005. Potential model weeds to study genomics, ecology, and physiology in the 21<sup>st</sup> century. *Weed Science* 53, 927-937.
- Charmet, G. & Balfourier, F., 1994. Isozyme variation and species relationships in the genus *Lolium* L. (ryegrasses, Graminaceae). *Theoretical and Applied Genetics* 87, 641-649.
- Chou, C.H., 1999. Roles of allelopathy in plant biodiversity and sustainable agriculture. *Critical Reviews in Plant Science* 18, 609-636.
- Chung, I-M., Seigler, D., Miller, D.A., & Kyung, S.H., 2000. Autotoxic compounds from

fresh alfalfa leaf extracts: identification and biological activity. *Journal of Chemical Ecology* 26, 315-27.

Conklin, A.E., Erich, M.S . & Liebman, M., 2002. Effects of red clover (*Trifolium pratense*) green manure and compost soil amendments on wild mustard (*Brassica kaber*) growth and incidence of disease. *Plant and Soil* 238, 245-256.

Cousens, R., 1985. A simple model relating yield loss to weed density. *Annals of Applied Biology* 107, 239-252.

Dabney, S.M., Schreiber, J.D., Rothrock, C.S. & Johnson, J.R., 1996. Cover crops affect sorghum seedling growth. *Agronomy Journal* 88, 961-970.

Daldon, B.R., Blum, U. & Weed, S.B., 1983. Allelopathic substances in ecosystems: Effectiveness of sterile soil components in altering recovery of ferulic acid. *Journal of Chemical Ecology* 9, 1185-1201.

Dinelli, G., Bonetti, A., Lucchese, C., Catizone, P., Bravin, F. & Zanin, G., 2002. Taxonomic evaluation of Italian populations of *Lolium* spp resistant and susceptible to diclofop-methyl. *Weed Research* 42, 156-165.

Duke, S.O., Dayan, F.E., Romagni, J.G. & Rimando, A.M., 2000. Natural products as sources of herbicides: current status and future trends. *Weed Research* 40, 99-111.

Dyck, E. & Liebman, M., 1994. Soil fertility management as a factor in weed control: the effect of crimson clover residue, synthetic nitrogen fertilizer, and their interaction on emergence and early growth of lambsquarters and sweet corn. *Plant and Soil* 167, 227-237.

Efthimiadou, A.P., Karkanis, A.C., Bilalis, D.J. & Efthimiadis, P., 2009. Review: the phenomenon of crop-weed competition; a problem or a key for sustainable weed management? *Journal of Food, Agriculture & Environment* 7, 861-868.

Einhellig, F.A., 1996. Interactions involving allelopathy in cropping systems. *Agronomy Journal* 88, 886-893.

Ehrenfeld, J.G., 2006. A potential novel source of information for screening and monitoring the impact of exotic plants on ecosystems. *Biological Invasions* 8, 1511-1521.

El-Shatnawi, M.K.J. & Makhadmeh, I.M., 2001. Ecophysiology of the plant-rhizosphere system. *Journal of Agronomy and Crop Science* 187, 1-9.

Fay, P.K. & Duke W.B., 1977. An assessment of allelopathic potential in *Avena* germplasm. *Weed Science* 25, 224-228.

Fourie, J.C., 2005. Cover crop management in the vineyards of the Lower Orange River region, South Africa: 1. Performance of grass and broadleaf species. *South African Journal of Enology and Viticulture* 26, 140-146.

Fourie, J.C., Louw, P.J.E. & Agenbag, G.A., 2005. Cover crop management in a Sauvignon blanc/Ramsey vineyard in the semiarid Olifants River Valley, South Africa. 1. Effect of management practices on selected grass and broadleaf species. *South African Journal of Enology and Viticulture* 26, 131-139.

Frigo, M.J., Mangolin, C.A., Oliveira, R.S. Jr. & Machado, M.P.S., 2009. Esterase polymorphism for analysis of genetic diversity and structure of wild poinsettia (*Euphorbia heterophylla*) populations. *Weed Science* 57, 54-60.

Gallandt, E.R., Molloy, T., Lynch, R.P. & Drummond, F.A., 2005. Effect of cover-cropping systems on invertebrate seed predation. *Weed Science* 53, 69-76.

Garland, J.L. & Mills, A.L., 1991. Classification and characterization of heterotrophic microbial communities on the basis of patterns of community level sole-carbon-source utilization. *Applied and Environmental Microbiology* 57, 2351-2359.

Gill, G.P., Wilcox, P.L., Whittaker, D.J., Winz, R.A., Bickerstaff, P., Echt, C.E., Kent, J.,

- Humphreys, M.O., Elborough, K.M. & Gardner, R.C., 2006. A framework linkage map of perennial ryegrass based on SSR markers. *Genome* 49, 354-364.
- Gu, Y., Li, H.B. & Kong, C.H. 2008a. Allelopathic potential of barnyard grass on rice and soil microbes in paddy. *Allelopathy Journal* 21, 389-396.
- Gu, Y., Wang, P. & Kong, C.H., 2008b. Effects of rice allelochemicals on the microbial community of flooded paddy soil. *Allelopathy Journal* 21, 299-310.
- Guenzi, W.D., Kehr, W.R. & McCalla, T.M., 1964. Water soluble phytotoxic substances in alfalfa forage: Variation with variety, cutting, year, and stage of growth. *Agronomy Journal* 56, 499-500.
- Gupta, U.S., 2005. Physiology of stressed crops. Volume III. The Stress of Allelochemicals, University of Georgia, Science Publishers, Inc., Enfield (NH), USA.
- Hall, M.H. & Henderlong, P.R., 1989. Alfalfa autotoxic fraction characterization and initial separation. *Crop Science* 29, 425-28.
- Henson, I.E., 1970. The effects of light, potassium nitrate and temperature on the germination of *Chenopodium album* L. *Weed Research* 10, 27-39.
- Heap, I., 2010. The International Survey of Herbicide Resistant Weeds ([www.weedscience.com](http://www.weedscience.com)).
- Hierro, J.L. & Callaway, R.M., 2003. Allelopathy and exotic plant invasion. *Plant and Soil* 256, 29-39.
- Hirata, M., Cai, H., Inoue, M., Yuyama, N., Miura, Y., Komatsu, T., Takamizo, T. & Fujimori, M., 2006. Development of simple sequence repeat (SSR) markers and construction of an SSR-based linkage map in Italian ryegrass (*Lolium multiflorum* Lam.). *Theoretical and Applied Genetics* 113, 270-9.

Hoffman, M.L., Weston, L.A., Snyder, J.C. & Regnier, E.E., 1996. Allelopathic influence of germinating seeds and seedlings of cover crops on weed species. *Weed Science* 44, 579-584.

Holm, L.G., Plucknett, D.L., Pancho, J.V. & Herberger, J.P., 1977. The World's Worst Weeds: Distribution and Biology. Hawaii, Honolulu; University Press.

Hoult, A.H.C. & Lovett, J.V., 1993. Biologically active secondary metabolites of barley. III. A method for identification and quantification of hordenine and gramine in barley by high-performance liquid chromatography. *Journal of Chemical Ecology* 19, 2245-2254.

Huang, H.C. & Chou, C.H., 2005. Impact of Plant Disease Biocontrol and Allelopathy on Biodiversity and Agricultural Sustainability. *Plant Pathology Bulletin* 14, 1-12.

Inderjit., 2005. Soil microorganisms: an important determinant of allelopathic activity. *Plant and Soil* 274, 227-236.

Jasieniuk, M. & Maxwell, B.D., 2001. Plant diversity: new insights from molecular biology and genomics technologies. *Weed Science* 49, 257-265.

John, J.A., & M.H. Quenouille. 1977. Experiments: Design and Analysis, 2<sup>nd</sup> Edition, The Garden City Press Limited, Letchworth, Hertfordshire, UK, 296 pp.

Jones, S., Dupal, P., Dumsday, L., Hughes, J. & Forster, W., 2002. An SSR-based genetic linkage map for perennial ryegrass (*Lolium perenne L.*). *Theoretical and Applied Genetics* 105, 577-584.

Jones, E., Jessop, R.S., Sindel., B.M. & Hoult. A., 1999. Utilising crop residues to control weeds. p. 373-376. *in:* Bishop, A. Boersma, M. and Barnes, C.D. (ed.), Proceedings of the 12<sup>th</sup> Australian Weeds Conference. Tasmanian Weeds Society, Devonport, Australia.

Kato-Noguchi, H., 2000. Assessment of the allelopathic potential of extracts of *Evolvulus*

*alsinoides*. *Weed Research* 40, 343-350.

Kato-Noguchi H, Salam MA & Kobayashi T (2009) A quick seeding test for allelopathic potential of Bangladesh rice cultivars. *Plant Production Science* 12, 47-49.

Kazinczi, G., Horváth, J. & Takács, A.P., 2005. Plant-plant and plant-virus interactions. Lectures and papers presented at the 7th Slovenian Conference on Plant Protection, Zreče, Slovenia, 8-10 March.

Khanh, T.D., Chung, M.I., Xuan, T.D. & Tawata, S., 2005. The exploitation of crop allelopathy in sustainable agricultural production. *Journal of Agronomy & Crop Science* 191, 172-184.

Kong, C.H., 2008. Rice allelopathy. *Allelopathy Journal* 21, 261-274.

Kong, C.H., Wang, P., Zhao, H., Xu, X.H. & Zhu, Y.D., 2008. Impact of allelochemical exuded from allelopathic rice on soil microbial community. *Soil Biology & Biochemistry* 40, 1862-1869.

Kruidhof, H.M., 2008. Cover crop-based ecological weed management: exploration and optimization. PhD Thesis, Wageningen University, Wageningen, The Netherlands. 156 pp.

Kumar, V., Brainard, D.C. & Bellinder, R.R., 2008. Suppression of Powell amaranth (*Amaranthus powellii*), shepherd's-purse (*Capsella bursa-pastoris*), and corn chamomile (*Anthemis arvensis*) by buckwheat residues: role of nitrogen and fungal pathogens. *Weed Science* 56, 271-280.

Kumar, V., Brainard, D.C. & Bellinder, R.R. 2009. Suppression of Powell amaranth (*Amaranthus powellii*) by buckwheat residues: role of allelopathy. *Weed Science* 57, 66-73.

Lamprecht, S.C., Marasas, W.F.O., Hardy, M.B. & Calitz, F.J., 2006. Effect of crop rotation on crown rot and the incidence of *Fusarium pseudograminearum* in wheat in the

Western Cape, South Africa. *Australasian Plant Pathology* 35, 419-426.

Lee, C.E., 2002. Evolutionary genetics of invasive species. *Trends in Ecology and Evolution* 17, 386-391.

[Lehle, F.R.](#), [Frans, R.](#) & [McClelland, M.](#), 1983. Allelopathic potential of Hope white lupine (*Lupinus albus*) herbage and herbage extracts. *Weed Science* 31, 513-519.

[Leflaive, J.](#) & [Ten-Hage, L.](#), 2007. Algal and cyanobacterial secondary metabolites in freshwaters: a comparison of allelopathic compounds and toxins. *Freshwater Biology* 52, 199-214.

Levene, H. 1960: Robust test in the equality of variance IN: Contr. Prob. Stat., I. Olkin (Ed), 278-292. Palo Alto, CA, USA: Stanford University Press.

Levitt, J., Lovett, J.V. & Garlick, P.R., 1984. *Datura stramonium* allelochemicals longevity in soil, and ultrastructural effects on root tip cells of *Helianthus annuus* L. *New Phytologist* 97, 213-218.

Liebl, R.A., Simmons, F.W., Wax, L.M. & Stoller, E.W., 1992. Effect of rye (*Secale cereale*) mulch on weed control and soil moisture in soybean (*Glycine max*). *Weed Technology* 6, 838-846.

Liebman, M. & Davis, A.S., 2000. Integration of soil, crop and weed management in low-extemal-input farming systems. *Weed Research* 40, 27-47.

Liebman, M. & Mohler, C.L., 2001. Weeds and the soil environment. p. 210-268. In: Ecological Management of Agricultural Weeds, M. Liebman, C.L. Mohler and C.P. Staver (eds.), Cambridge University Press, Cambridge.

Lipin'ska, H. & Lipin'ski, W., 2009. Initial growth of *Phleum pratense* under the influence of leaf water extracts from selected grass species and the same extracts improved with MgSO<sub>4</sub>.7H<sub>2</sub>O. *Journal of Elementology* 14, 101-110.

[Lin W-X.](#), [Xiong, J.](#), [Zhou, J-J.](#), [Qiu, L.](#), [Shen, L-H.](#), [Li, Z-F.](#), [Chen, H.](#), [Hao, H-R.](#), [Chen, T.](#), [Lin, R-Y.](#), [He, H-B.](#) & [Liang, Y-Y.](#), 2007. Research status and its perspective on the properties of rhizosphere biology mediated by allelopathic plants. [Zhongguo Shengtai Nongye Xuebao / Chinese Journal of Eco-Agriculture](#) 15, 1-8.

Locken, L.J. & Kelsey RG., 1987. Cnicin concentrations in *Centaurea maculosa*, spotted knapweed. *Biochemical Systems Ecology* 15, 313-320.

Lopes, M.S., Bernardi, L.E., Farencena, I.R. & Lamb, P.R., 1987. Effect of extracts of weeds on germination and initial growth of rice (*Oryza sativa* L. cv BR-IRGA 409). *Lavoura Arrozeira* 40, 22-23.

Lovett, J.V., 1983. Self-defense chemicals of plants. *Proceedings of the 10<sup>th</sup> International Congress on Plant Protection*. p 838.

Lovett, J.V. & Hoult, A.H.C., 1995. Allelopathy and self-defense in barley. In: Allelopathy: Organisms, processes and applications, K.M.N. Inderjit, & F.A.E. Dakshini (eds.), American Chemical Society Symposium Series 582, 170-183.

Lovett, J.V., Ryunyu, M.Y. & Liu, D.L., 1989. Allelopathy, chemical communication, and plant defense. *Journal of Chemical Ecology* 15, 1193-1201.

Machado, S., 2007. Allelopathic potential of various plant species on downy brome: implications for weed control in wheat production. *Agronomy Journal* 99, 127-132.

Madhou, P.A., Wells, A., Pang, E.C.K. & Stevenson, T.W., 2005. Genetic variation in populations of Western Australian wild radish. *Australian Journal of Agricultural Research* 56, 1079-1087.

Mamolos, A.P. & Kalburjji, K.L., 2001. Significance of allelopathy in crop rotation. *Journal of Crop Production* 4, 197-218.

Manici, L.M., Caputo, F. & Babini, V., 2004. Effect of green manure on *Pythium* spp. population and microbial communities in intensive cropping systems. *Plant and*

*Soil* 263, 133-142.

McCalla, T.M. & Norstadt, F.A., 1974. Toxicity problems in mulch tillage. p. 29. In: Physiology of stressed crops. Volume III. The Stress of Allelochemicals, 2005. U.S. Gupta (ed.), University of Georgia, Science Publishers, Inc., Enfield (NH), USA.

Mian, M.A., Saha, M.C., Hopkins, A.A. & Wang, Z.Y., 2005. Use of tall fescue EST-SSR markers in phylogenetic analysis of cool-season forage grasses. *Genome* 48, 637-647.

Mohler, C.L. & Teasdale, J.R., 1993. Response of weed emergence to rate of *Vicia villosa* Roth and *Secale cereale* L. residue. *Weed Research* 33, 487-499.

Molisch, H. 1937. Der Einfluss einer Pflanze auf die andere- Allelopathie. Gustav Fischer, Jena.

Monaghan, N.M., 1980. The biology and control of *Lolium rigidum* as a weed of wheat. *Weed Research*. 20, 117-121.

Moodie, M., Finch, R.P. & Marshall, G., 1997. Analysis of genetic variation in wild mustard (*Sinapis arvensis*) using molecular markers. *Weed Science* 45, 102-107.

Mwaja, V.N., Masiunas, J.B. & Weston, L.A., 1995. Effects of fertility on biomass, phytotoxicity, and allelochemical content of cereal rye. *Journal of Chemical Ecology* 21, 81-96.

Narwal, S. S., 1994. Allelopathy in Crop Production. Scientific Publishers, Jodhpur, India.

[Oberan, L.V.](#), [Djurdjevic, L.](#), [Mitrovic, M.](#), [Pavlovic, P.](#) & [Kostic, O.](#), 2008. Allelopathic interactions between the soil microorganisms and dominant plants in *Orno-Quercetum virginiana* forest on Avala Mt. (Serbia). [Allelopathy Journal](#) 22, 167-180.

Oerke, E.C., 2006 Crop losses to pests. *The Journal of Agricultural Science* 144, 31-43.

O'Hanlon, P.C., Peakall, R. & Briese, D.T., 2000. A review of new PCR-based genetic markers and their utility to weed ecology. *Weed Research* 40, 239-254.

Olofsdotter, M., 2001. Getting closer to breeding for competitive ability and the role of allelopathy – an example from rice (*Oryza sativa*). *Weed Technology* 15, 798-806.

Olofsdotter, M., Jensen, L.B. & Courtois, B., 2002. Improving crop competitive ability using allelopathy – an example from rice. *Plant Breeding* 121, 1-9.

Olofsdotter, M., Navarez, D. & Moody, K., 1995. Allelopathic potential in rice (*Oryza sativa* L.) germplasm. *Annals of Applied Biology* 127, 543-560.

Olofsdotter, M., Navarez, D., Rebulanan, M. & Streibig, J.C., 1999. Weed-suppressing rice cultivars – does allelopathy play a role? *Weed Research* 39, 441-454.

Ott, R.L. 1998. An introduction to statistical methods and data analysis. 807-837. Belmont, CA, USA: Duxbury Press.

Pacheco, A. & Pohlan, J., 2007. Intercropping aromatic plants with coffee (*Coffea arabica* L.) under greenhouse and field conditions. *Revista Colombiana de Ciencias Hortícolas* 1, 94-102.

Pieterse, P.J. & Cairns, A.L.P., 2008. Current status of herbicide resistance in South Africa. Combined Congress 21-24 January 2008, Grahamstown, South Africa.

Putnam, A. R. & DeFrank, J., 1983. Use of phyto toxic plant residues for selective weed control. *Crop Protection* 2, 173-181.

Purvis, C.E., 1990. Differential Response of Wheat to Retained Crop Stubbles. I. Effect of Stubble Type and Degree of Decomposition. *Australian Journal of Agricultural Research* 41, 225-42.

Qasem, J.R., 1994. Allelopathic effect of whitetop (*Lepidium draba*) and wheat and barley. *Allelopathy Journal* 1, 29-40.

Qasem, J.R. & Foy, C.L., 2001. Weed allelopathy, its ecological impacts and future prospect: a review. *Journal of Crop Production* 4, 43-120.

Qasem, J.R. & Hill, T.A., 1989. Possible role of allelopathy in the competition between tomato, *Senecio vulgaris* L. and *Chenopodium album* L. *Weed Research* 29, 349-356.

Qasem, J.R., 1994. Allelopathic effect of whitetop (*Lepidium draba*) and wheat and barley. *Allelopathy Journal* 1, 29-40.

Reberg Horton, S.C., Burton, J.D. & Danehower, D.A., 2005. Changes over time in the allelochemical content of ten cultivars of rye (*Secale cereale* L.). *Journal of Chemical Ecology* 31, 179-193.

Reinhardt, C.F., Meissner, R. & Labuschagne, N., 1994. Allelopathic interaction of *Chenopodium album* L. and certain crop species. *South African Journal of Plant and Soil* 11, 45-49.

Reinhardt, C.F., Meissner, R. & Van Wyk, L.J., 1997. Allelopathic effects of *Chenopodium album* L. and *Chenopodium polyspermum* L. on another weed and two crop species. *South African Journal of Plant and Soil* 14, 165-168.

Rice, E.L., 1984. Allelopathy. 2<sup>nd</sup> edn. Academic Press, Orlando, Florida, USA.

Ridenhour, W.M. & Callaway, R.M., 2001. The relative importance of allelopathy in interference: The effects of an invasive weed on a native bunchgrass. *Oecologia* 126, 444-450.

Saha, M.C., Mian, M.A., Eujayl, I., Zwonitzer, J.C., Wang, L. & May, G.D., 2004. Tall fescue EST-SSR markers with transferability across several grass species. *Theoretical and Applied Genetics* 109, 783-91.

Saha, M.C., Mian, R., Zwoniter, J.C., Chekovskiy, K. & Hopkins, A.A., 2005. An SSR- and

AFLP-based genetic linkage map of tall fescue (*Festuca arundinacea* Schreb.).  
*Theoretical and Applied Genetics* 110, 323-336.

[San Emeterio, L.](#), [Arroyo, A.](#) & [Canals, R.M.](#), 2004. Allelopathic potential of *Lolium rigidum* Gaud. on the early growth of three associated pasture species. [\*Grass and Forage Science\*](#) 59, 107-112.

Sarika, P., Pandey, N. & Rao, P.B. 2008. Response of certain weed species extracts on germination and seedling growth of wheat (*Triticum aestivum* L.). *Environment and Ecology* 26, 2061-2066.

Schumacher, W.J., Thill, D.C. & Lee, G.A., 1983. Allelopathic potential of wild oat (*Avena fatua*) on spring wheat (*Triticum aestivum*) growth. *Journal of Chemical Ecology* 9, 1235-1245.

Seal, A.N., Pratley, J.E. & Haig, T.J., 2005. Evaluation of rice varieties for allelopathic effects on Australian rice weeds – linking laboratory to field. Proceedings of the 4<sup>th</sup> World Congress on Allelopathy, August 2005, Wagga Wagga, Australia. 164-167.

Senda, T., Kubo, N., Hirai, M. & Tominaga, T., 2004. Development of microsatellite markers and their effectiveness in *Lolium temulentum*. *Weed Research* 44, 136-141.

Senda, T., Saito, M., Ohsako, T., & Tominaga, T., 2005. Analysis of *Lolium temulentum* geographical differentiation by microsatellite and AFLP markers. *Weed Research* 45, 18-26.

Shapiro, S.S., & Wilk, M.B., 1965. An analysis of variance test for normality (complete samples). *Biometry* 52, 591-611.

Singh, H.P., Batish, D.R. & Kohli, R.K., 2001. Allelopathy in agroecosystems: an overview. *Journal of Crop Production* 4, 1-41.

Smith, M.W., Wolf, M.E., Cheary, B.S. & Carroll, B.L., 2001. Allelopathy of bermudagrass,

tall fescue, redroot pigweed, and cutleaf evening primrose on pecan. *HortScience* 36, 1047-1048.

Soil Classification Working Group., 1991. Soil Classification – A Taxonomic System for South Africa. Soil and Irrigation Research Institute, Department of Agricultural Development, Pretoria.

Sozeri, S. & Ayhan, A., 1998. Effect of stubble on wheat seed germination and seedling growth. p. 29. In: Physiology of stressed crops. Volume III. The Stress of Allelochemicals 2005. U.S. Gupta (ed.), University of Georgia, Science Publishers, Inc., Enfield (NH), USA.

Spahillari, M., Hammer, K., Gladis, T. & Diederichsen, A., 1999. Weeds as part of agrobiodiversity. *Outlook on Agriculture* 28, 227-232.

Stefureac, T.I. & Fratilesque-Sesan, T., 1979. Contribution to the study of the reciprocal action on the seeds of plant during germination. *Biologie Vegetala* 31, 55-61.

Studer, B., Boller, B., Hermann, D., Bauer, E., Posset, U.K., Widmer, F. & Kolliker, R., 2006. Genetic mapping reveals a single major QTL for bacterial wilt resistance in Italian ryegrass (*Lolium multiflorum* Lam.). *Theoretical and Applied Genetics* 11, 661-71.

Teasdale, J.R. & Mohler, C.L., 1993. Light transmittance, soil temperature, and soil moisture under residue of hairy vetch and rye. *Agronomy Journal* 85, 673-680.

Teasdale, J.R. & Pillai, P., 2005. Contribution of ammonium to stimulation of smooth pigweed (*Amaranthus hybridus* L.) germination by extracts of hairy vetch (*Vicia villosa* Roth) residue. *Weed Biology and Management* 5, 19-25.

Tranel, P.J. & Trucco, F., 2009. 21st-Century Weed Science: A call for *Amaranthus* genomics p. 53-71. In: Stewart, C.N., Jr. (ed.), *Weedy and Invasive Plant Genomics*. Wiley-Blackwell, Iowa, USA.

Treier, U.A., Broennimann, O., Normand, S., Guisan, A., Schaffner, U., Steinger, T. & Muller-Scharer, H., 2009. Shift in cytotype frequency and niche space in the invasive plant *Centaurea maculosa*. *Ecology* 90, 1366-1377.

Uppar, D.S., Nalini, A.S., Hiremath, S.M. & Kamatar, M.Y., 1993. Allelopathic effects of weeds on germination and vigour index of wheat. University of Agricultural Sciences Bangalore, *Current Research* 22, 47-48.

Vilarino, M.P.D., Mareggiani, G., Grass, M.Y., Leicach, S.R. & Ravetta, D.A. 2005. Post-damage alkaloid concentration in sweet and bitter lupine varieties and its effect on subsequent herbivory. *Journal of Applied Entomology* 129, 233-238.

Vyvyan, J.R., 2002. Allelochemicals as leads for new herbicides and agrochemicals. *Tetrahedron* 58, 1631-1646

Wardle, D.A. & Nilsson, MC., 1997. Microbe-plant competition, allelopathy and arctic plants. *Oecologia* 109, 291-293.

Weston, L.A., 1996. Utilization of allelopathy for weed management in agroeco-systems. *Agronomy Journal* 88, 860-866.

Weston, L.A., 2005. History and Current Trends in the Use of Allelopathy for Weed Management. Proceedings of the 4<sup>th</sup> World Congress on Allelopathy, Wagga Wagga, Australia. 15-21.

Weston, L.A. & Duke, S.O. 2003. Weed and crop allelopathy. *Critical Reviews in Plant Sciences* 22, 367-389.

Whittaker, R.H. & Feeney, P.P., 1971. Allelochemics: chemical interactions between species. *Science* 171, 757-770.

Wink, M., 1983. Inhibition of seed germination by quinolizidine alkaloids. Aspects of allelopathy in *Lupinus albus* L. *Planta* 158, 365-368.

[Wu, H.W.](#), [Pratley, J.](#) & [Haig, T.](#), 2003. Phytotoxic effects of wheat extracts on a herbicide-resistant biotype of annual ryegrass (*Lolium rigidum*). *Journal of Agricultural and Food Chemistry* 51, 4610-4616.

Wu, H.W., Pratley, J., Lemerle, D. & Haig, T., 2001. Allelopathy in wheat (*Triticum aestivum*). *Annals of Applied Biology* 139, 1-9.

Wu, H.W., Pratley, J., Lemerle, D. & Haig, T., 1999. Crop cultivars with allelopathic capability. *Weed Research* 39, 171-180.

Wu, H.W., Pratley, J., Lemerle, D. & Haig, T. 2000b. Laboratory screening for allelopathic potential of wheat (*Triticum aestivum*) accessions against annual ryegrass (*Lolium rigidum*). *Australian Journal of Agricultural Research* 51, 259-266.

[Wu, H.W.](#), [Haig, T.](#), [Pratley, J.](#), [Lemerle, D.](#) & [An, M.](#), 2000a. Distribution and exudation of allelochemicals in wheat *Triticum aestivum*. *Journal of Chemical Ecology* 26, 2141-2154.

[Xu, Y-F.](#), [Bi, Y-F.](#) & [Jin, J-W.](#), 2008. Allelopathic effects of *Cajanus cajan* L. on the seed germination and seedling growth of 6 shrub & grass species. *Journal of Yunnan Agricultural University* 23, 375-380.

Xuan, T.D., Tawata, S., Khanh, T.D. & Chung, M.I., 2005. Decomposition of allelopathic plants in soil. *Journal of Agronomy & Crop Science* 191, 162-171.

Xuan, T.D. & Tsuzuki, E., 2002. Varietal differences in allelopathic potential of alfalfa. *Journal of Agronomy & Crop Science* 188, 2-7.

Yuan, J.S., Tranel, P.J. & Stewart, C.N. Jr., 2007. Non-target-site herbicide resistance: a family business. *Trends in Plant Science* 12, 6-13.