

## 5. References

- Altschul, S.F., T.L. Madden, A.A. Schäffer, J. Zhang, Z. Zhang, W. Miller & D.J. Lipman** (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research* 25: 3389-3402.
- Bent, A.F., B.N. Kunkel, D. Dahlbeck, K.L. Brown, R.L. Schmidt, J. Giraudat, J.L. Leung & B.J. Staskawicz** (1994). *RPS2* of *Arabidopsis thaliana*: A leucine-rich repeat class of plant disease resistance genes. *Science* 265: 1856-1860.
- Blattner, F.R., G. Plunkett III, C.A. Bloch, N.T. Perna, V. Burland, M. Riley, J. Collado-Vides, J.D. Glasner, C.K. Rode, G.F. Mayhew, J. Gregor, N.W. Davis, H.A. Kirkpatrick, M.A. Goeden, D.J. Rose, B. Mau & Y. Shao** (1997). The complete genome sequence of *Escherichia coli* K-12. *Science* 277: 1453-1474.
- Cannon, S.B., H. Zhu, A.M. Baumgarten, R. Spangler, G. May, D.R. Cook & N. D. Young** (2002). Diversity, distribution, and ancient taxonomic relationships within the TIR and non-TIR NBS-LRR resistance gene subfamilies. *Journal of Molecular Evolution* 54: 548-562.
- Chomczynski, P. & N. Sacchi** (1987). Single-step method of RNA isolation by acid guanidium thiocyanate-phenol-chloroform extraction. *Analytical Biochemistry* 162: 156-159.
- Deng, Z., S. Huang, P. Ling, C. Chen, C. Yu, C.A. Weber, G.A. Moore & F.G. Gmitter Jr.** (2000). Cloning and characterization of NBS-LRR class resistance-gene candidate sequences in citrus. *Theoretical and Applied Genetics* 101: 814-822.
- Dixon, M.S., K. Hatzixanthis, D.A. Jones, K. Harrison & J.D.G. Jones** (1998). The tomato Cf-5 disease resistance gene and six homologs show pronounced allelic variation in leucine-rich repeat copy number. *Plant Cell* 10: 1915-1925.
- Edwards, K., C. Johnstone & C. Thompson** (1991). A simple and rapid method for the preparation of plant genomic DNA for PCR analysis. *Nucleic Acids Research* 19: 1349.
- Ellis, J., P. Dodds & T. Pryor** (2000). Structure, function and evolution of plant disease resistance genes. *Current Opinion in Plant Biology* 3: 278-284.

**Garcia-Mas, J., H. Van Leeuwen, A. Monfort, M.C. De Vincente, P. Puigdomènech & P. Arús** (2001). Cloning and mapping of resistance gene homologues in melon. *Plant Science* 161: 165-172.

**Jackson, A.O. & C.B. Taylor** (1996). Plant-microbe interactions: Life and death at the interface. *The Plant Cell* 8: 1651-1668.

**Jones, D.A. & J.D.G. Jones** (1997). The role of leucine-rich repeat proteins in plant defenses. *Advances in Botanical Research* 24: 89-167.

**Jones, D.A., C.M. Thomas, K.E. Hammond-Kosack, P.J. Balint-Kurti & J.D.G. Jones** (1994). Isolation of the tomato *Cf-9* gene for resistance to *Cladosporium fulvum* by transposon tagging. *Science* 266: 789-793.

**Lagudah, E.S., O. Moullet & R. Appels** (1997). Map-based cloning of a gene sequence encoding a nucleotide-binding domain and a leucine-rich region at the *Cre3* nematode resistance locus of wheat. *Genome* 40: 659-665.

**Leister, D., A. Ballvora, F. Salamini & C. Gebhardt** (1996). A PCR-based approach for isolating pathogen resistance genes from potato with potential for wide application in plants. *Nature Genetics* 14: 421-429.

**Leister, D., J. Kurth, D.D. Laurie, M. Yano, T. Sasaki, K. Devos, A. Graner & P. Schulze-Lefert** (1998). Rapid reorganization of resistance gene homologues in cereal genomes. *Proceedings of the National Academy of Science of the United States of America*. 95: 370-375.

**Martin, G.B., S.H. Brommonschenkel, J. Chunwongse, A. Frary, M.W. Ganal, R. Spivey, T. Wu, E.D. Earle & S.D. Tanksley** (1993). Map-based cloning of a protein kinase gene conferring disease resistance in tomato. *Science* 262: 1432-1436.

**Meyers, B.C., D.B. Chin, K.A. Shen, S. Sivaramakrishnan, D.O. Lavelle, Z. Zhang & R.W. Michelmore** (1998). The major resistance gene cluster in lettuce is highly duplicated and spans several megabases. *Plant Cell* 10: 1817-1832.

**Myburg, A.A., M. Cawood, B.D. Wingfield & A-M. Botha** (1998). Development of RAPD and SCAR markers linked to the Russian wheat aphid resistance gene *Dn2* in wheat. *Theoretical and Applied Genetics* 96: 1162-1169.

**Ochman, H., M.M. Medhora, D. Garza & D.L. Hartl** (1990). Amplification of flanking sequences by inverse PCR. In: *PCR protocols, a guide to methods and applications*. Eds. M.A. Innis, D.H. Gelfand, J.J. Sninsky & T.J. White. Academic Press, London, United Kingdom, pp 219-227.

**Pan, Q., Y-S. Liu, O. Budai-Hadrian, M. Sela, L. Carmel-Goren, D. Zamir & R. Fluhr** (2000). Comparative genetics of nucleotide binding site-leucine rich repeat resistance gene homologues in the genomes of two dicotyledons: tomato and Arabidopsis. *Genetics* 155: 309-322.

**Ritter, C. & J.L. Dangl** (1996). Interference between two specific pathogen recognition events mediated by distinct plant disease resistance genes. *Plant Cell* 8: 251-257.

**Sambrook, J., E.F. Fritsch & T. Maniatis** (1989). *Molecular cloning: A laboratory manual*, Second Edition. Cold Spring Harbour Laboratory, NY.

**Sandhu, D. & K.S. Gill** (2002). Gene-containing regions of wheat and the other grass genomes. *Plant Physiology* 128: 803-811.

**Seah, S., K. Sivasithamparam, A. Karakousis & E.S. Lagudah** (1998). Cloning and characterization of a family of disease resistance gene analogs from wheat and barley. *Theoretical and Applied Genetics* 97: 937-945.

**Siebert, P.D., A. Chenchik, D.E. Kellogg, K.A. Lukyanov & S.A. Lukyanov** (1995). An improved PCR method for walking in uncloned genomic DNA. *Nucleic Acids Research* 23: 1087-1088.

**Whitham, S., S.P. Dinesh-Kumar, D. Choi, R. Hehl, C. Corr & B. Baker** (1994). The product of the tobacco mosaic virus resistance gene N: Similarity to Toll and the interleukin-1 receptor. *Cell* 78: 1101-1115.

**Yu, Y.G., G.R. Buss & M.A.S. Maroof (1996).** Isolation of a superfamily of candidate disease-resistance genes in soybean based on a conserved nucleotide-binding site. Proceedings of the National Academy of Science of the United States of America 93: 11751-11756.



## CHAPTER V

### Summary/ Opsomming



## Summary

Infestation by the Russian wheat aphid, *Diuraphis noxia* (Mordvilko), has caused large-scale damage to small-grain crops since its introduction into South Africa in 1978 and many other countries world-wide. The extreme damage caused by *D. noxia* in wheat (*Triticum aestivum*) and barley (*Hordeum vulgare*) has resulted in a concerted effort by scientists to investigate the mechanisms and genes involved in the resistance response. This is done in an effort to understand and ultimately use the knowledge of resistance responses to produce aphid resistant crops. The aim of the present study was to use suppression subtractive hybridization (SSH) and degenerate oligonucleotide primers to amplify cDNA of differentially expressed genes and nucleotide-binding site (NBS)-containing sequences respectively. This served as a source for the identification and characterization of disease-resistance genes in the wheat line PI 294994 after infestation with the Russian wheat aphid. SSH is an effective approach for studying the genetic nature of many biological processes by identifying differentially expressed genes. The sequences that were obtained in this study did indeed show some similarity to the underlying mechanisms involved in lesion formation and disease resistance. At the amino acid level similarities were identified with polypeptides such as lymphoma receptors. At the nucleotide level similarities were identified to enzymes involved in fibrinogen lysis and the chloroplast regulatory system. A heat shock protein gene was also identified, which may possibly play a role in the induction of the resistance response during aphid infestation. The use of PCR with degenerate oligonucleotide primers, designed from the NBS region of cloned disease resistance genes, has led to the cloning of resistance sequences in various plant species. In this study oligonucleotide primers designed from conserved motifs in the NBS domain were used to clone disease resistance gene homologues from the wheat line PI 294994. The sequences obtained showed a homology to an NBS region in lettuce, which was linked to a resistance protein candidate gene. In the wheat line PI 294994 the cloned NBS sequence was expressed at varying levels in different wheat tissues, which requires further study. However, when the full-length cDNA sequence of the NBS-containing gene was sequenced, there was no significant homology to any disease resistance genes at the nucleotide level. At the amino acid level there was homology to a kinase type resistance protein. The degenerate NBS

PCR approach was very successful in producing a number of NBS regions, which may be used for further study even if they are not necessarily involved in disease resistance. These results have aided the process of identifying novel aphid induced transcripts after infestation and has contributed to the growing base of knowledge about the underlying mechanisms involved in lesion formation and resistance responses to insects.

## Opsomming

Sedert die eerste waarneming van Russiese koringluis (*Diuraphis noxia* Mordvilko) in Suid Afrika in 1974, is gootskaalse verliese by klein-graas gewasse gerapporteer. Dit is ook die geval vir ander lande. Die ernstige skade wat deur *D. noxia* op koring (*Triticum aestivum*) en gort (*Hordeum vulgare*) veroorsaak word, het tot 'n gesamentlike poging deur wetenskaplikes gelei om die meganismes en gene betrokke by die weerstandsreaksie te ondersoek. Die poging lei tot 'n beter begrip van die meganisme en gene betrokke by die RWA weerstandsreaksie wat uiteindelik gebruik kan word om nog meer luisweerstandbiedende gewasse te kweek. Die doelwit van die huidige studie was om subtraksie-suppressie-hibridisasie (SSH) en gedegenerereerde oligonukleotiedvoorvoeders te gebruik om komplimentêre-DNA (cDNA) van differensieël uitgedrukte gene en nukleotiedbindingsetels- (NBS) -bevattende volgordes onderskeidelik te amplifiseer. Dit is gedoen om siekteweerstandsgene in die koringlyn PI 294994, na infestering met die Russiese koringluis, te identifiseer en te karakteriseer. SSH is 'n effektiewe nadering om die genetiese aard van baie biologiese prosesse te bestudeer deur die identifisering van differensieël uitgedrukte gene. Die volgordes wat in hierdie studie verkry was, het wel sommige ooreenkomste met die onderliggende meganismes gewys en was dan wel funksioneel in letsselformasie en siekteweerstand. Op aminosuurvlak is ooreenkomste geïdentifiseer met onder andere limfoomreseptors. Terwyl ooreenkomste met ensieme wat by fibrinogeenoplossing en die chloroplasreguleeringsstelsel betrokke is, op nukleotiedvlak geïdentifiseer is. 'n Hitte-skok-proteïen is ook geïdentifiseer wat dalk 'n rol in die induksie van die weerstandsreaksie gedurende plantluisinfestering mag speel. Die gebruik van die polimerasekettingreaksie (PCR) met gedegenerereerde oligonukleotiedvoorvoeders van die NBS-gebied, het tot die klonering van weerstandvolgordes in verskeie plantsoorte gelei. In dié studie is oligonukleotiedvoorvoeders vanaf gekonserveerde motiewe in die NBS-gebied ontwerp en gebruik om die siekteweerstandgeen-homoloë van die koringlyn PI 294994 te kloner. Die volgordes wat verkry is, het homologie met 'n NBS-gebied in blaarslaai getoon wat aan 'n weerstandproteïenkandidaatgeen gekoppel is. In die koringlyn PI 294994 is die gekloneerde NBS-volgorde in variërende vlakke, in verskillende koringweefsel uitgedruk. Hierdie aspek regverdig verdere studie. Die volledige cDNA-basisvolgorde van die NBS-bevattendegeen het geen betekenisvolle



homologie aan enige siekteweerstandsgene op nukleotiedvlak getoon nie. Op aminosuurvlak was daar wel homologie met 'n kinase-tipe weerstandsproteïen. Die gedegeneerde NBS-PCR-nadering was baie suksesvol om 'n aantal NBS-bevattende fragmente te produseer, al hou dit nie noodwendig met siekteweerstand verband nie. Hierdie fragment kan vir verdere studie gebruik word. Hierdie resultate het die proses van identifikasie van nuwe plantluis-geïnduseerde transkripte na infestasië geondersteun en bygedra tot die groeiende basis van kennis oor die onderliggende meganismes wat by letsselformasie en die weerstandsreaksie teen insekte betrokke is.