

**THE EFFECTS OF AN *AD LIBITUM* VERSUS A RESTRICTED PLANE
OF NUTRITION ON THE PERFORMANCE AND
LONGEVITY OF DAIRY CATTLE**

by

CASPARUS MAREE

Submitted to the Faculty of Agriculture
(Department Animal Production)
University of Pretoria

in partial fulfilment of the requirements
for the degree of
D.Sc. (Agric)

**PRETORIA
MAY, 1975**

INDEX

	Page
ACKNOWLEDGEMENTS	iv
LIST OF TABLES	v
LIST OF FIGURES	vi
CHAPTER I : INTRODUCTION	1
1. Paradox in the distribution of food	1
1.1 The physiological regulation of food intake	3
1.2 Some effects of the nutritional level on the production and longevity of animals	5
1.3 The reproductive performance of cattle on a high level of feeding	10
1.4 Intensification and the erosion of dairy cows	14
CHAPTER II : MATERIAL AND PROCEDURE	16
2.1 The calves, their management and development until conception	16
2.1.1 Examination of genitalia and insemination	17
2.1.2 Level of feeding before insemination	18
2.1.3 Observations on body temperature, heart and respiratory rates	19
2.1.4 Observations after the first calving	19
2.1.4.1 Procedures and regrouping after first calving	20
2.1.4.2 Rations	20
2.1.4.3 Body mass	22
2.1.4.4 Body temperature, heart and respiratory rates	22
2.1.4.5 Cyclic and ovarian activity during first and second lactation	22
2.1.4.6 Inseminations	23
2.1.4.7 Observations at calving	23
2.2 Post partum observations	23
2.2.1 Clinical	23
2.2.2 Histological	24
2.2.2.1 Uterine biopsies	24
2.2.2.2. Examination of biopsy sections	24
2.3 Blood serum analysis	25
2.3.1 Clinical observations and treatment	26
2.3.2 Final insemination and slaughtering	27
CHAPTER III : RESULTS AND DISCUSSION	28
3.1 Ovarian activity	28
3.1.1 Ovulation and ovulatory failure	28
3.1.2 Cycle length	37
3.1.3 The time of ovulation from the onset of estrus	41
3.1.4 Ovulation and estrus performance	45
3.1.5 Follicular and corpus luteum development	46
3.1.6 The respective roles of right and left ovaries	51
3.1.7 Quiescent estrus	57

		Page
3.1.8	Conception in relation to ovulatory performance .	59
3.1.9	Body mass	65
3.1.10	Temperature, heart and respiratory rates	67
3.1.11	Laminitis and arthritis	70
3.1.12	Milk production	79
3.1.13	The process of parturition	81
3.1.13.1	The first calving	82
3.1.13.2	The second calving	82
3.1.13.3	The duration of parturition and uterine involution	96
3.1.13.4	Uterine inertia	104
3.2	Histological examination of the endometrium	107
3.2.1	The morphology of the non pregnant uterine wall..	108
3.2.1.1	The epithelium	109
3.2.1.2	The lamina propria	110
3.2.1.3	The stratum compactum	110
3.2.1.4	The caruncles	110
3.2.1.5	The stratum spongiosum	110
3.2.1.6	The uterine glands	110
3.2.1.7	Wandering cells	111
3.2.1.7.1	Neutrophiles	111
3.2.1.7.2	Lymphocytes	112
3.2.1.7.3	Eosinophiles	112
3.2.1.7.4	Plasma cells	113
3.2.1.7.5	Histiocytes	114
3.2.1.7.6	Mast cells	114
3.2.2	Involutional changes in the bovine uterus after parturition	114
3.2.3	Cyclic changes in the endometrium	116
3.2.4	Uterine biopsies as a method of examination of the endometrium	116
3.2.4.1	The cell population of the endometrium in the HIGH and LOW feeding groups	117
3.2.4.2	Neutrophiles	119
3.2.4.3	Plasma cells	119
3.2.4.4	Lymphocytes	120
3.2.4.5	Eosinophiles	120
3.2.4.6	Mast cells	121
3.2.4.7	Histiocytes	121
3.2.4.8	Mobile cells in the surface epithelium	122
3.2.4.9	Cell types during the follicular and luteal phases of the cycle	122
3.2.4.10	Special features of cells in the uterine wall	124
3.2.4.11	Epithelial cell height and nuclei	125
3.2.4.12	The endometrial glands	127
3.2.4.12.1	Wandering cells	128
3.2.4.12.2	Glandular debris	128
3.2.4.12.3	Glandular nuclei	129
3.2.4.12.4	Glandular cell height	129
3.2.4.12.5	Glandular lumina	129
3.2.4.13	Lymph follicles in the endometrium	129
3.2.4.14	Subepithelial haemorrhage	130
3.2.5	Discussion	130

ACKNOWLEDGEMENTS

Without the generous assistance of several people and institutions, this investigation would not have been possible. I am grateful in particular to –

Prof. J.C. Bonsma, my promotor and after his retirement my co-promotor, who through his wide experience rendered me great assistance

Prof. G.N. Louw, my promotor, for his valued association with this investigation

Messrs. National Chemical Products, Germiston, Transvaal for facilities at their experimental farm during the initial stages of this work and for making available trial animals for the entire duration of this investigation

Prof. W. Gernecke for aid with histological techniques and interpretations

Prof. K. van der Walt and Mr. W.A.J. Walzl for clinical pathological analysis

Messrs. R.J. de Kock and A. McLaren for biochemical analysis

Dr. P. Boyazoglu for mineral determinations

Mr. S. Eloff for the preparation of histological sections

Mr. P. van Loggerenberg for the programming and statistical analysis of data

Messrs. H. Swanepoel, P. Bester, Vincent Maledimo and A. Mahlatje for the management and care of experimental animals, particularly Vincent for long hours of faithful duty over many years.

Financial assistance by the Department Agricultural Technical Services made this investigation possible.

LIST OF TABLES

Table		Page
1(a)	Ovulation record of 32 heifers during all cycles from the onset of puberty until conception	29
1(b)	Ovulation record of cows kept on a low feeding level after their first calving	30
1(c)	Ovulation record of cows kept on a high feeding level after their first calving	30
2	The incidence of ovulatory failure at estrus following on cycles of abnormal length	39
3	The time of ovulation from the onset of estrus	42
4	The number of inseminations received per animal and record of ovulations at three breeding periods : (a) as heifers (b) after first calving (c) after second calving	60
5	The duration of parturition, time to involution of the uterus, time to development of the first CL and incidence of quiescent estrus after calving in cows on a low and a high nutritional level	96
6	Birth mass of calves and maternal mass at time of birth in cows on low and high feeding levels	97
7	Ease of calving of cows on high and low feeding levels in relation to the mass of calves at birth expressed as a percentage of the maternal body mass at birth	98
8	Mobile cells in the endometrium during estrus and the luteal phase in cows on a high and a low feeding level	124
9	Blood serum values of cows on a high and a low feeding level before and after the onset of treatment	137
10	Clinical pathological values of blood serum of cows on a high and a low level of feeding during 24 months	138
11	The mineral content of liver specimens of cows kept on a high and a low feeding level	140
12	Post parturient complications, laminitis and arthritis recorded in cows on a high feeding level, together with clinical symptoms and post mortal lesions at slaughtering	143

LIST OF FIGURES

Figure		Page
1(a)	Ovulatory performance of Frieslands dairy animals before and after calving	33
1(b)	Ovulatory pattern before and after calving and on high and low feeding levels	34
2(a)	Differences in cycle length before and after first calving and between high and low feeding groups after calving	35
2(b)	Distribution of cycles of normal and abnormal cycle length – as adults – as heifers	36
3(a)	The time of ovulation from the onset of first estrus as heifers and after first calving	43
3(b)	The time of ovulation from the onset of estrus in cattle on a low and a high level of feeding after first calving	44
4(a)	Numerical index value of follicular activity in both ovaries on each day of the cycle in cows on a high and a low level of feeding	53
4(b)	Numerical index value of follicular activity on each day of the cycle in the ovulating and non-ovulating ovaries respectively in cows on a high and a low level of feeding	54
5	Monthly average body mass of cows on a high (H) and a low (L) level of feeding from July 1969 to July 1971	66
6	Temperature, respiratory and heart rates of cows on high and low nutritional levels	68
7	Decrease in the size of the uterus after parturition as determined by rectal palpation and expressed according to a numerical index for the HIGH and the LOW groups respectively	100
8(a)–(f)	The cell population of the bovine endometrium at weekly intervals from 7 to 72 days after calving in cows on a high and a low level of feeding respectively	118
9	The number of neutrophils, lymphocytes and histiocytes per microscopical field in bovine endometrial epithelium at weekly intervals in cows on high and low feeding levels	123
10	Cellular height and vesicularity of nuclei in endometrial epithelium during the post calving period (a) High feeding level (b) Low feeding level	126