

CHAPTER SIX

REFERENCES

- Abbaszadegan, M., Huber, H.S., Gerba, C.P. and Pepper, I.L. 1993. Detection of enteroviruses in groundwater with the polymerase chain reaction. *Appl. Environ. Microbiol.* 59: 1318-1324.
- Alberts, M.J. 1994. *Vibrio cholerae* O139 Bengal. *J. Clin. Microbiol.* 32: 2345-2349.
- Alvarez, A.J., Hernandez-Delgado, E.A. and Toranzos, G.A. 1993. Advantages and disadvantages of traditional and molecular techniques applied to the detection of pathogens in waters. *Wat. Sci. Tech.* 27: 253 - 256.
- Amann, R. I., Ludwig, W., and Schleifer, K.-H. 1995. Phylogenetic identification and *in situ* detection of individual microbial cells without cultivation. *Microbiol. Rev.* 59: 143-169.
- American Public Health Association. 1998. Standard Methods for the Examination of Water and Wastewater, American Public Health Association, Washington D.C.
- Anderson, S.A., Dewis, G.D. and M.N. Pearson. 1995. Use of gene probes for the detection of quiescent enteric bacteria in marine and fresh waters. *Wat. Sci. Tech.* 31: 291-298.
- Anon. 1996. Waterborne pathogens kill 10M-20M people a year. *World Water Environ. Engin.* June 6.
- Arnheim, N. and Erlich, H. 1992. Polymerase chain reaction strategy. *Ann. Rev. Biochem.* 61: 131-156.

Arroyo, G. and Arroyo, J.A. 1995. Efficiency of different enrichment and isolation procedures for the detection of *Salmonella* serotypes in edible offal. *J. Appl. Bact.* 79: 360-367.

Arroyo, G. and Arroyo, J.A. 1996. Selective action of inhibitors used in different culture media on competitive microflora of *Salmonella*. *J. Appl. Bact.* 78:281-289.

Barret, T J. and Feeley, J.C. 1994. Serologic diagnosis of *Vibrio cholerae* O1 infections, pp 135-141. *In: Vibrio cholerae and cholera: Molecular Methods to Global Perspectives*, I.K. Wachsmuths, P.S.A. Blake, Ø. Olsvik (eds.), American Society for Microbiology, Washington, D.C.

Baumann, P. and Schubert R.H.W. 1984. Family II. Vibrionaceae, pp. 516-550. *In: Bergey's Manual of Systematic Bacteriology*, N. R. Krieg and J.G. Holt (eds.), Vol.1, Williams & Wilkins Co., Baltimore.

Bej, A.K. and Mahbubani, M.H. 1992. Applications of the polymerase chain reaction in environmental microbiology. *PCR Methods and Applications* 1: 151-159.

Bej, AK., Mahbubani, M.H., Miller, R., DiCesare, J.L., Haff, L. and Atlas, R M. 1990. Multiplex PCR amplification and immobilized capture probes for detection of bacterial pathogens and indicators in water. *Mol. Cell. Probes* 4: 353-365.

Bej, A.K., Mahbubani, M.H., Dicesare, J.L. and Atlas, R.M. 1991a. Polymerase chain reaction-gene probe detection of microorganisms by using filter-concentrated samples. *Appl. Environ. Microbiol.* 57: 3529-3534.

Bej, A.K., Steffan, R.J., DiCesare, L., Haff, L. and Atlas, R.M. 1991b. Detection of *Escherichia coli* and *Shigella* spp. in water using the polymerase chain reaction and gene probes for *uid*. *Appl. Environ. Microbiol.* 57: 1013-1017.

Bennish, M.L. 1991. Potentially lethal complications of shigellosis. *Rev. Infect. Diseases* 13: Suppl. 4, S319-324.

Blostein, J. 1991. Shigellosis from swimming in a park pond in Michigan. *Public Health Report* 106: 317-322.

Bobb, C.A., Summer, J.W., Morris, G.K. and Wells, J.G. 1981. Isolation of *Legionella* spp. from environmental water samples by low-pH treatment and use of a selective medium. *J. Clin. Microbiol.* 13: 714-719.

Bode, G., Mauch, F. and Malfertheiner, P. 1993. The coccoid forms of *Helicobacter pylori*: Criteria for their viability. *Epidemiol. Infect.* 111: 483-490.

Bottger, E.C., Jurs, M., Barrett, T., Wachsmuth, K., Metzger, S. and Bitter-Suermann, D. 1987. Qualitative and quantitative determination of enterobacterial common antigen (ECA) with monoclonal antibodies. *Clin. Microbiol.* 25: 377-382.

Brugha, R., Vipond, I.B., Evans, M.R., Sandifer, Q.D., Roberts, R.J., Salmon, R.L., Caul, E.O. and Mukerjee, A.K. 1999. A community outbreak of food-borne small round-structured virus gastroenteritis caused by a contaminated water supply. *Epidemiol. Infect.* 122: 145-54.

Buswell, C.M., Herlihy, Y.M., Lawrence, L.M., McGuiggan, J.T.M., Marsh, P.D., Keevil, C.W. and Leach, S.A. 1998. Extended survival and persistence of *Campylobacter* spp. in water and aquatic biofilms and their detection by immunofluorescent-antibody and r-RNA staining. *Appl. Environ. Microbiol.* 64: 733-741.

Byrd, J.J., Xu, H-S. and Colwell, R.R. 1991. Viable but nonculturable bacteria in drinking-water. *Appl. Environ. Microbiol.* 57:875-78.

Cappelier, J.M. and Federighi, M. 1998. Demonstration of viable but nonculturable state *Campylobacter jejuni*. *Rev. Med. Vet.* 149: 319-326.

Carter, A.M., Pacha, R.E., Clark, G.W. and Williams, E.A. 1987. Seasonal occurrence of *Campylobacter* spp. in surface waters and their correlation with standard indicator bacteria. *Appl. Environ. Microbiol.* 53: 523-526.

Cash, R.A., Music, S.I., Libonati, J.P., Snyder, M.J., Wenzel, R.P. and Hornick, R.B. 1974. Response of man to infection with *Vibrio cholerae*. Clinical, serologic and bacteriologic responses to a known inoculum. *J. Infect. Dis.* 129: 45-52.

Catalan, V., Moreno, F.G., Vila, M.J. and Apraiz, D. 1997. Detection of *Legionella pneumophila* in wastewater by nested polymerase chain reaction. *Res. Microbiol.* 148: 71-78.

Cerdà-Cuéllar, M., Jofre, J. and Blanch, A.R. 2000. A selective medium and a specific probe for detection of *Vibrio vulnificus*. *Appl. Environ. Microbiol.* 66: 855-859.

Cheasty, T. and Rowe, B. 1983. Antigenic relationships between the Enteroinvasive *Escherichia coli* O antigens O28ac, O124, O136, O143, O144, O152 and O164 and *Shigella* O antigens. *J. Clin. Microbiol.* 17: 681-684.

Chopra, M., Wilkinson, D. and Stirling, S. 1997. Epidemic shigella dysentery in children in northern KwaZulu-Natal. *SAMJ* 87: 49-51.

Colwell, R.R., Hasan, J.A.K., Huq, A., Loomis, L., Seibeling, R.J., Torres, M., Galvez, S., Islam, S. and Bernstein, D. 1992. Development and evaluation of a rapid, simple, sensitive, monoclonal antibody-based co-agglutination test for direct detection of *Vibrio cholerae* O1. *FEMS Microbiol. Lett.* 97: 215-220.

Colwell, R.R., Brayton, P.R., Grimes, D.J., Roszak, D.B., Huq, S.A. and Palmer, L.M. 1985. Viable but non-culturable *Vibrio cholerae* and related pathogens in the environment: Implications for release of genetically engineered microorganisms. *Bio. Technol.* 3: 817-820.

Colwell, R.R., Tamplin, M.L., Brayton, P.R., Tavgens, A.L., Tall, B.D., Herrington, D., Levine, M.M., Hall, S., Huq, A. and Sack, D.A. 1990. Environmental aspects of *Vibrio cholerae* in transmission of cholera, pp. 327-343. *In: Advances in Research on Cholera and Related Areas*, R.B. Sack and R. Zinnaki (eds.), 7th Edition, KTK Scientific Publishers, Tokyo.

Covert, T.C., Rodgers, M.R., Reyes, A.L. and Stelma, G.N. 1999. Occurrence of nontuberculous mycobacteria in environmental samples. *Appl. Environ. Microbiol.* 65: 2492-2496.

Craun, G. 1988. Surface Water Supplies and Health. *J. AWWA* 80: 40-52.

Dalsgaard, A., Frimodt-Moller, N., Bruun, B., Hoi, L. and Larsen, J.L. 1996. Clinical manifestations and epidemiology of *Vibrio vulnificus* infections in Denmark. *Eur. J. Clin. Microbiol. Infect. Dis.* 15: 227-231.

Davies, J. 1994. Inactivation of antibiotics and the dissemination of resistance genes. *Science.* 264: 375-382.

De Silva, D.G.H., Candy, D.C.A., Mendis, L.N., Charts, H. and Rowe, B. 1992. Serological diagnosis of infection by *Shigella dysenteriae*-1 in patients with bacillary dysentery. *J. Infect.* 25: 273-278.

De Ryck, R., Struelens, M.J. and Serruys, E. 1994. Rapid biochemical screening for *Salmonella*, *Shigella*, *Yersinia* and *Aeromonas* isolates from stool specimens. *J. Clin. Microbiol.* 32: 1583-1585.

Degrange, V. and Bardin, R. 1995. Detection and counting of *Nitrobacter* populations in soil by PCR. *Appl. Environ. Microbiol.* 61: 2093-2098.

DePaola, A. and Hwang, G. 1995. Effect of dilution, incubation time, and temperature of enrichment on cultural and PCR detection of *Vibrio cholerae* obtained from the oyster

Crassostrea virginica. *Mol. Cell. Probes* 9: 75-81.

Diviacco, S., Norio, P., Zentilin, L., Menzo, S., Clementi, M., Biamonti, G., Riva, S., Falaschi, A. and Giacca, M. 1992. A novel procedure for quantitative polymerase chain reaction by co-amplification of competitive templates. *Gene* 122: 313-320.

Dixon, B. 1998. Viable but not culturable. *ASM NEWS*. 64: 372-373.

Donovan, T.J. and van Netten, P. 1995. Culture media for the enumeration of pathogenic *Vibrio* species in foods and environmental samples. *Int. J. Food Microbiol.* 26: 77-91.

Dorman, C.J. and Porter, M.E. 1998. The *Shigella* virulence gene regulatory cascade: A paradigm of bacterial gene control mechanisms. *Molec. Microbiol.* 29: 677-684.

Dorsch, M., Ashbolt, N.J., Cox, P.T. and Goodman, A.E. 1994. Rapid identification of *Aeromonas* species using 16S rRNA-targeted oligonucleotide primers: A molecular approach based on screening of environmental isolates. *J. Appl. Bact.* 77: 722-726.

DuPont, H L. 1990. *Shigella* species (Bacillary dysentery), pp 1716-1722. *In: Principles and Practice of Infectious Diseases*, G.L. Mandell, R.G Douglas, Jr, and J. E. Bennett. (eds.), 3rd Edition, Churchill, Livingston Inc., New York.

Dupray, E., Caprais, M.P., Derrien, A. and Fach, P. 1997. *Salmonella* DNA persistence in natural seawaters using PCR analysis. *J. Appl. Microbiol.* 82: 507-510.

Engels, D., Madaras, T., Nyandwi, S. and Murray, J. 1995. Epidemic dysentery caused by *Shigella dysenteriae* type 1. A sentinel site surveillance of antimicrobial resistance patterns in Burundi. *Bull. WHO.* 73: 787-791.

Enroth, H. and Engstrand, L. 1995. Immunomagnetic separation and PCR for detection of *Helicobacter pylori* in water and stool specimens. *J. Clin. Microbiol.* 33: 2162-2165.

- Ewing, W.H. and Lindberg, A. A. 1984. Serology of *Shigella*. *Meth. Microbiol.* 14: 113-142.
- Falkind, S., Stark, M.J.A., Uhlén, M., Lundeberg, J. and Weintraub, A. 1996. Cloning and sequencing of a region of *Vibrio cholerae* O139 Bengal and its use in PCR-based detection. *J. Clin. Microbiol.* 34: 2904-2908.
- Falkow, S. and Mekalanos, J. 1990. The enteric bacilli and Vibrios, pp. 561-587. *In: Microbiology*, B.D. Davis, T. Dulbecco, H.N. Eisen and H.S. Ginsberg (eds.), 4th Edition., J.B. Lippincott Company, Philadelphia.
- Farmer III, J.J. and Hickman-Brenner, F.W. 1992. The Genera *Vibrio* and *Photobacterium*, pp. 2952-3011. *In: The Prokaryotes*, A. Balows, H.G. Trüper, M. Dworkin, W. Harder and K-H. Schleifer (eds.), Springer-Verlag, New York.
- Faude, U.C. and Höffle, M.G. 1997. Development and application of monoclonal antibodies for *in situ* detection of indigenous bacterial strains in aquatic ecosystems. *Appl. Environ. Microbiol.* 63: 4534-4542.
- Feldman, R.A. 1992. Transmission of cholera: modes of spread and vehicles. *Microbiol. Dig.* 9: 37-41.
- Fields, P.I., Popovic, T., Wachsmuth, K. and Olsvik, Ø. 1992. Use of the polymerase chain reaction for detection of toxigenic *Vibrio cholerae* O1 strains from the Latin American cholera epidemic. *J. Clin. Microbiol.* 30: 2118-2121.
- Finkelstein, R.A. 1988. Cholera, the cholera enterotoxins, and the cholera enterotoxin-related enterotoxin family, pp. 85-102. *In: Immunological and Molecular Genetic Analysis of Bacterial Pathogens*, P. Owen and T.J. Foster, (eds.), Elsevier Science, Amsterdam.
- Fleming, C.A., Caron, D., Gunn, J.E., Horine, M.S., Matyas, B.T. and Barry, M.A. 2000.

An outbreak of *Shigella sonnei* associated with a recreational spray fountain. *Am. J. Pub. Hlth.* 90:1641-1642.

Ford, T.E. 1999. Microbiological safety of drinking water: United States and global perspectives. *Environ. Hlth. Perspect.* 107: 191-206.

Frahm, E. and Obst, U. 1995. Comparison of two rapid tests for the detection of *Legionellaceae* in water: A microbiological-immunological method and a commercial gene-probing testing. *Wat. Sci. Tech.* 31: 403-406.

Frankel, G., Giron, J.A., Valmassoi, J. and Schoolink, G.K. 1989. Multi-gene amplification: Simultaneous detection of three virulence genes in diarrhoeal stool. *Mol. Microbiol.* 3: 1729-1734.

Furness, B.W., Beach, M.J. and Roberts, J.M. 2000. *Mor. Mortal. Wkly. Rep. CDC Surveill. Sum.* 11: 1-13.

Furtado, C., Adak, G.K., Stuart, J.M., Wall, P.G., Evans, H.S. and Casemor, D.P. 1998. Outbreaks of waterborne infectious intestinal disease in England and Wales. 1992-1995. *Epidemiol. Infect.* 121:109-19.

Garret, C.T., Ferreira-Centeno, A. and Nasim, S. 1993. Molecular diagnostics: Issues of utilization, regulation and organization. *Clinica Chimica Acta.* 217: 85-103.

Gerhardt, P. and Drew, S. 1994. Liquid culture, pp. 224-244. *In: Methods for General and Molecular Bacteriology*, H. Gerhardt, R.G.E. Murray, W.A. Wood and N.R. Krieg (eds.), American Society for Microbiology, Washington D.C.

Glass, R.I., Libel, M. and Brandling-Bennett, A.D. 1992. Epidemic cholera in the Americas. *Science*. 265:1524-1525.

Gonzalez, A. T., Torres, J., Dvorsky, E. and Toranzos, G. A. 1995. Modified culture methods for the detection of *Vibrio* spp. from estuarine waters. *Wat. Sci. Tech.* 31: 283-290.

Goodwin, C.S., Blincow, E.D., Warren, J.R., Waters, T.E., Sanderson, C.R. and L. Easton. 1985. Evaluation of cultural techniques for isolating *Campylobacter pyloridis* from endoscopic biopsies of gastric mucosa. *J. Clin. Pathol.* 38:1127-1131.

Grabow, W.O.K. 1996. Waterborne diseases: Update on water quality assessment and control. *Wat. SA.* 2: 193-202.

Graczyk, T.K., Cranfield, M. R. and Fayer, R. 1997. Recovery of waterborne oocysts of *Cryptosporidium* from water samples by the membrane-filter dissolution method. *Parasitol. Res.* 83: 121-125.

Greenough III, W.B. 1985. *Vibrio cholerae*, p. 1209-1218. In: Principles and Practice of Infectious Diseases, G.L. Mandell, R.G. Douglas and J. E. Bennet, (eds.), John Wiley & Sons, New York.

Guidolin, A. and Manning, P.A. 1987. Genetics of *Vibrio cholerae* and its bacteriophages. *Microbiol. Rev.* 51: 285-298.

Hafliger, D., Hubner, P. and Luthy, J. 2000. Outbreak of viral gastroenteritis due to sewage-contaminated drinking water. *Int. J. Food Microbiol.* 10: 123-132.

Hale, T.L. 1991. Genetic basis of virulence in *Shigella* species. *Microbiol. Rev.* 55: 206-224.

Hartman, A.B., Venkatesan, M., Oaks, E.V. and van Buysse, J.M. 1990. Sequence and molecular characterization of multicopy invasion plasmid antigen gene, *ipaH*, of *Shigella flexneri*. *J. Bact.* 172: 1905-1915.

Hasan, J.A.K., Huq, A., Tamplin, R.J., Siebeling, R.J. and Colwell, R.R. 1994. A novel kit for rapid detection of *Vibrio cholerae* O1. *J. Clin. Microbiol.* 32: 249-252.

Hayes, E.B., Matte, T.D. and O'Brein, T.R. 1989. Large community outbreak of cryptosporidiosis due to contamination of a filtered public water supply. *N. Engl. J. Med.* 320:1372-1376.

Hazen, T.C. and Jimenez, L. 1988. Enumeration and identification of bacteria from environmental samples using nucleic acid probes. *Microbiol. Sci.* 5: 340-343.

Hegarty, J.P., Dowd, M.T. and Baker, K.H. 1999. Occurrence of *Helicobacter pylori* in surface water in the United States. *J. Appl. Microbiol.* 87: 697-701.

Hellard, M. E., Sinclair, M.I., Streeton, C.L. and Fairly, C.K. 1997. Drinking water and microbial pathogens-issues and challenges for the year 2000. *J. Public. Hlth. Med.* 19: 129-131.

Hermans, P.W.M., Schuitema, A.P.J., Van Soolingen, D., Verstynen, C.P.H., Bik, E.M., Thole, J.E.R., Kolk, A.H.J. and Van Embden, J.D.A. 1990. Specific detection of *Mycobacterium tuberculosis* complex strains by polymerase chain reaction. *J. Clin. Microbiol.* 28: 1204-1213.

Høi, L., Dalsgaard, I. and Dalsgaard, A. 1998. Improved isolation of *Vibrio vulnificus* from seawater and sediment with Cellobiose-Colistin agar. *Appl. Environ. Microbiol.* 64: 1721-1724.

Huang, H., Garcia, M.M., Brooks, B. W., Nielsen, K. and Ng, S-P. 1999. Evaluation of culture enrichment procedures for use with *Salmonella* detection immunoassay. *Int. J. Food Microbiol.* 51: 85-94.

Huang, P.W., Laborde, D., Land, V.R., Matson, D.O., Smith, A.W. and Jiang, X. 2000. Concentration and detection of caliciviruses in water samples by reverse transcription-PCR. *Appl. Environ. Microbiol.* 66: 4383-4388.

Islam, M.S., Hasan, M.K., Miah, M.A., Sur, G.C., Felsenstein, M., Venkatesan, M., Sack, R.B. and Albert, M.J. 1993a. Use of the polymerase chain reaction and fluorescent-antibody methods for detecting viable and nonculturable *Shigella dysenteriae* Type 1 in laboratory microcosms. *Appl. Environ. Microbiol.* 59: 536-540.

Islam, M S., Hossain, M Z., Khan, S I., Felsenstein, A., Sack, R.B. and Albert, M.J. 2001. Detection of non-culturable *Shigella dysenteriae* 1 from artificially contaminated volunteers fingers using fluorescent antibody and PCR techniques. www.icddrb.org/journal/jddr152/JDDR152-65.html

Islam, D. and Lindberg, A.A. 1992. Detection of *Shigella dysenteriae* Type 1 and *Shigella flexneri* in feces by immunomagnetic isolation and polymerase chain reaction. *J. Clin. Microbiol.* 30: 2801-2806.

Islam, D., Tzipori, S. and Lindberg, A.A. 1993b. Rapid detection of *Shigella dysenteriae* and *Shigella flexneri* in faeces by an immunomagnetic assay with monoclonal antibodies. *Eur. J. Clin. Microbiol. Infect. Dis.* 12: 25-32.

Ito, T., Kuwahara, S. and Yokota, T. 1983. Automatic and manual latex agglutination tests for measurement of cholera toxin and heat-labile enterotoxin of *Escherichia coli*. *J. Clin. Microbiol.* 17: 7-12.

Jackson, C.J., Fox, A.J. and Jones, D.M. 1996. A novel polymerase chain reaction assay for the detection and speciation of thermophilic *Campylobacter* spp. *J. Appl. Bact.* 81: 467-473.

Jaykus, L.A., Hermard, M.T. and Sobsey, M.D. 1994. Human enteric pathogenic viruses. *In: Environmental Indicators and Shellfish Safety*, pp. 92-153. C.R. Hackney and M.D. Peirson, (eds.), Chapman and Hall, New York.

Josephson, K.L., Gerba, C.P. and Pepper, I.L. 1993. Polymerase chain reaction of nonviable bacterial pathogens. *Appl. Environ. Microbiol.* 59: 3513 - 3515.

Juck, D., Ingraham, J., Prévost, M., Coallier, J. and Greer, C. 1996. Nested PCR protocol for the rapid detection of *Escherichia coli* in potable water. *Can. J. Microbiol.* 42: 862-866.

June, G.A., Sherrod, P.S., Imaguana, R.M., Andrews, W.H. and Hammack, T.S. 1993. Effectiveness of the Bacteriological Analytical Manual culture method for the recovery of *Shigella sonnei* from selected foods. *J. AOAC Int.* 76: 1240-1248.

Kahn, A.S., Moe, C.L., Glass, R.I., Monroe, S.S., Estes, M.K., Chapman, L.E., Jiang, X., Humphrey, C., Pon, E., Iskander, J.K. and Schönberger, L.B. 1994. Norwalk virus-associated gastroenteritis traced to ice consumption aboard a cruise ship in Hawaii: Comparison and application of molecular method-based assays. *J. Clin. Microbiol.* 32: 318-322.

Kang, D.H. and Siragusa, G.R. 1999. Agar underlay method for recovery of sublethally heat-injured bacteria. *Appl. Environ. Microbiol.* 65: 5334-5337.

Kaper, J.B., Fasano, A. and Trucksis, M. 1994. Toxins of *Vibrio cholerae*, pp. 145-176. In: *Vibrio cholerae and Cholera: Molecular to Global Perspectives*, I. K. Wachsmuth, P.A. Blake and Ø. Olsvik (eds.), American Society for Microbiology, Washington, D.C.

Kaper, J.B., Moseley, S.L. and Falkow, S. 1981. Molecular characterization of environmental and non-toxigenic strains of *Vibrio cholerae*. *Infect. Immun.* 32: 661-667.

Kaplan, J.E., Gary, G.W., Baron, R.C., Singh, N., Schönberger, L.B., Feldman, R. and Greenberg, H.B. 1982. Epidemiology of Norwalk gastroenteritis and the role of Norwalk virus in outbreaks of acute nonbacterial gastroenteritis. *Ann. Int. Med.* 96: 756-761.

Kapperud, G., Vardund, T., Skjerve, E., Hornes, E. and Michaelsen, T.E. 1993. Detection of pathogenic *Yersinia enterocolitica* in foods and water by immunomagnetic separation, nested polymerase chain reactions and colorimetric detection of amplified DNA. *Appl. Environ. Microbiol.*, 59: 2938-2944.

Karaolis, D.K.R., Somara, S., Maneval Jr., D.R., Johnson, J.A. and Kaper, J.B. 1999. A bacteriophage encoding a pathogenicity island, type-IV pilus and phage receptor in cholera. *Nature* 399: 375-379.

Kay, B.A., Bopp, C.A. and Wells, J.G. 1994. Isolation and identification of *Vibrio cholerae* O1 from fecal specimens, pp. 3-25. In: *Vibrio and Cholera: Molecular to Global Perspectives*, I. K. Wachsmuth, P.A. Blake and Ø. Olsvik (eds.), American Society for Microbiology, Washington, D.C.

Keasler, S.P. and Hall, R.H. 1993. Detecting and biotyping *Vibrio cholerae* O1 with

multiplex polymerase chain reaction. *Lancet* 341: 1661.

Kfir, R. and Genthe, B. 1993. Advantages and disadvantages of the use of immunodetection techniques for the enumeration of microorganisms and toxins in water. *Wat. Sci. Tech.* 27: 243-248.

King, C. A. and Heyning, W. E. 1973. Deactivation of cholera toxin by sialidase-resistant monosialosylganglioside. *J. Infect. Dis.* 127: 639-647.

Knight, I.T., Di Ruggiero, J. and Colwell, R.R. 1991. Direct detection of enteropathogenic bacteria in estuarine water using nucleic acid probes. *Wat. Sci. Tech.* 24 : 261-266.

Kobayashi, K., Set, K., Akasaka, S. and Makino, M. 1990. Detection of toxigenic *Vibrio cholerae* O1 using polymerase chain reaction for amplifying the cholera enterotoxin gene. *J. Jap. Ass. Infect. Dis.* 64:1323-1329.

Koch, W.H., Payne, W.L., Wentz, B.A. and Cebula, T.A. 1993. Rapid polymerase chain reaction method for detection of *Vibrio cholerae* in foods. *Appl. Environ. Microbiol.* 59: 556-560.

Koenraad, P.M.F.J., Giesendorf, B.A.J., Henkens, M.H.C., Beumer, R.R. and Quint, W.G.V. 1995. Methods for the detection of *Campylobacter* in sewage: Evaluation of efficacy of enrichment and isolation media, applicability of polymerase chain reaction and Latex agglutination assay. *J. Microbiol. Meth.* 23: 309-320.

Kramer, M.H., Sorhabe, F.E., Goldstein, S.T., Dalley, E., Wahlquist, S.P. and Herwaldt, B.L. 1998. First reported outbreak in the United States of cryptosporidiosis associated with a recreational lake. *Clin. Infect. Dis.* 26: 27-33

Kreader, C.A. 1996. Relief on amplification inhibition in PCR with bovine serum albumin or T4 gene 32 protein. *Appl. Environ. Microbiol.* 62: 1102-1106.

Kukkula, M., Maunula, L., Silvennoinen, E. and van Bonsdorff, C.H. 1999. Outbreak of viral gastroenteritis due to drinking water contaminated by Norwalk-like viruses. *J. Infect. Dis.* 180: 1771-1776.

LeChevalier, M.W., Norton, W.D. and Lee, R.G. 1991. Occurrence of *Giardia* and *Cryptosporidium* spp. in surface water supplies. *Appl. Environ. Microbiol.* 57: 2610-2616.

Leser, T.D., Boye, M. and Hendriksen, N B. 1995. Survival and activity of *Pseudomonas* sp. strain B13 (FR1) in a marine microcosm determined by quantitative PCR and an rRNA-targeting probe and its effect on the indigenous bacterioplankton. *Appl. Environ. Microbiol.* 61: 1201-1207.

Lewis, C.W. and Mak, J.L. 1989. Comparison of membrane filtration and autoanalysis Colilert presence-absence techniques for analysis of total coliforms and *Escherichia coli* in drinking water samples. *Appl. Environ. Microbiol.* 55: 3091-3094.

Lewis, M.J. 1997. Shigella, (Bacillary dysentery), pp 262-266. *In: Medical Microbiology. A guide to Microbial Infections: Pathogenesis, Immunity, Laboratory diagnosis and Control*, D. Greenwood, R.C.B. Slack and J.F. Peutherer (eds.), Churchill, Livingston, New York.

Lindberg, A.A., Cam, P.D., Chan, N., Phu, L.K., Trach, D.D., Lindberg, C. and Ekwall, E. 1991. Shigellosis in Vietnam: Seroepidemiological studies with use of lipopolysaccharide antigens in enzyme-immuno assays. *Rev. Infect. Dis.* 13 (Suppl. 4), S231-S237.

Lindqvist, R. 1999. Detection of *Shigella* spp. in food with a nested PCR method-sensitivity and performance compared with a conventional culture method. *J. Appl. Microbiol.* 86: 971-978.

MacKenzie, W.R., Hoxie, N.J., Proctor, M.E., Gradus, M.S., Blair, K.A., Peterson, D.E., Kazmierczak, J.J., Addiss, D.G., Fox, K.R., Rose, J.B. and Davis, J.P. 1994. A massive outbreak in Milwaukee of *Cryptosporidium* infection transmitted through the public water supply. *N. Eng. J. Med.* 331: 161-167.

MacKenzie, W.R., Kazmierczak, J.J. and Davis, J.P. 1995. An outbreak of cryptosporidiosis associated with a resort swimming pool. *Epidemiol. Infect.* 115: 545-553.

Maguire, H.C., Seng C., Chambers S., Cheasty, T., Double, G., Soltanpoor N. and Morse, D. 1998. *Shigella* outbreak in a school associated with eating canteen food and person to person spread. *Commun. Dis. Public. Hlth.* 1: 279-280.

Mahbubani, M.H., Bej, A.K., Miller, R., Haff, L.D. and Atlas, R.M. 1990. Detection of *Legionella* with polymerase chain reaction and gene probe methods. *Mol. Cell. Probes* 4: 175-187.

Maiwald, M., Kissel, K., Srimuang, S., von Knebel Doeberitz, M. and Sontag, H.G. 1995. Comparison of polymerase chain reaction and conventional culture for the detection of legionellas in hospital water samples. *J. Appl. Bact.* 76: 216-225.

Mayer, C.L. and Palmer, C.J. 1996. Evaluation of PCR, nested PCR and fluorescent antibodies for detection of *Giardia* and *Cryptosporidium* species in wastewater. *Appl. Environ. Microbiol.* 62:2081-2085.

McAnulty, J.M., Fleming, D.W. and Gonzalez, A.H. 1994. A community-wide outbreak of cryptosporidiosis associated with swimming at a wave pool. *JAMA* 272: 1597-1600.

McDonald, I.R., Kenna, E.M. and Murrell, J.C. 1995. Detection of methanotrophic bacteria in environmental samples with the PCR. *Appl. Environ. Microbiol.* 61:116-121.

McDougald, D., Rice, S.A., Weichert, D. and Kjelleberg, S. 1998. Nonculturability: Adaptation or devilitation? *FEMS Microbiol. Ecol.* 25: 1-9.

McGinnis, M. J. and Foege, W.H. 1993. Actual causes of death in the United States. *JAMA* 270: 2207-2212.

McKay, A.M. 1992. Viable but nonculturable forms of potentially pathogenic bacteria in water. *Lett. Appl. Microbiol.* 14: 129 -135.

Mekalanos, J.J., Swartz, D.J. and Pearson, G.D.N. 1983. Cholera toxin genes: Nucleotide sequence, deletion analysis and vaccine development. *Nature* 306: 551-557.

Mekalanos, J.J., Rubin, E.J. and Waldor, M.K. 1997. Cholera: Molecular basis for emergence and pathogenesis. *FEMS Immun. Med. Microbiol.* 18: 241-248.

Morinigo, M.A., Martinez-Manzanares, E., Munoz, A., Cornax, R., Romero, P. and Borrego, J.J. 1989. Evaluation of different plating media used in the isolation of salmonellae from environmental samples. *J. Appl. Bact.* 66: 353-360.

Morris, J.G. and Potter, M. 1997. Emergence of new pathogens as a function of changes in host susceptibility. *Emerg. Infect. Dis.* 3: 435-441.

Morris, J. 1994. Non-group O1 *V. cholerae* strains not associated with epidemic disease, pp. 103-115. *In: Vibrio and Cholera: Molecular to Global perspectives*, I. K. Wachsmuth, P.A. Blake and Ø. Olsvik (eds.), American Society for Microbiology, Washington, D.C.

Morse, S.S. 1995. Factors in the emergence of infectious disease. *Emerg. Infect. Dis.* 1: 5-10.

Nabeth, P., Vasset, B., Guerin, P., Doppler, B. and Tectonidis, M. 1997. Health situation of refugees in eastern Zaire. *Lancet.* 349: 1031-1032.

Naik, S.R., Aggarwal, R., Salunke, P.N. and Mehrotra N.N. 1992. A large waterborne viral hepatitis E epidemic in Kanpur, India. *Bull. W.H.O.* 70: 597-604.

Nair, G.B., Sarkar, B.L., De, S.P., Chakraborty, M.K., Bhadra, R.K. and Pal, S.C. 1988. Ecology of *V. cholerae* in the fresh water environs of Calcutta. *Microb. Ecol.* 15: 203-215.

Nair, G.B., Bag, P.K., Shimada, T., Ramamurthy, T., Takeda, T., Yamamoto, S., Kurazono, H. and Takeda, Y. 1995. Evaluation of DNA probes for specific detection of *Vibrio cholerae* O139 Bengal. *J. Clin. Microbiol.* 33: 2186-2187.

National Status Report on Cholera Epidemic in South Africa. 2001. www.sandmc.pwv.gov.za/ndmc/DevChol/Cholrep/Summary/select.asp.

Neu, H.C. 1992. The crisis in antibiotic resistance. *Science* 257: 1064-1072.

Nikaido, H. and Vaara, M. 1984. Outer membrane, pp. 7-22. *In: Escherichia coli and*

Salmonella typhimurium, F.C. Neidhardt, J.L. Ingraham, K. Brooks Low, B. Magasanik, M. Schaechter, H.E. Umbarger (eds.), American Society for Microbiology, Washington, D.C.

Nilsson, L., Oliver, J.D. and Kjelleberg, S. 1991. Resuscitation of *Vibrio vulnificus* from the viable but nonculturable state. *J. Bact.* 173: 5054-5059.

Noah, C. W., Poteet, S.S., Lister, M.M., Roderick, C.N., Smith, D.B., Colvert, R.M., Holland, M.A. and Cerniglia, C.E. 1996. Production and characterization of monoclonal antibodies to detect *Vibrio cholerae* serogroup O1 in a rapid enzyme-linked immunosorbent assay. *J. Food Protec.* 59: 1153-1157.

Olsvik, Ø., Popovic, T. and Fields, P. I. 1993. PCR detection of toxin genes in strains of *Vibrio cholerae* O1, pp 266-276. *In: Diagnostic Molecular Microbiology. Principles and Applications*, D.H. Persing, T.F. Smith, F. C. Tenover and T. J. White, (eds.), American Society for Microbiology, Washington, D.C.

Olsvik, Ø., Popovic, T., Skjerve, E., Hornes, E., Ugelstad, J. and Uhlen, M. 1994. Magnetic separation techniques in diagnostic microbiology. *Clin. Microbiol. Rev.* 7: 43-54.

Oyofa, B.A. and Rollins, D. M. 1993. Efficacy of filter types for detecting *Campylobacter jejuni* and *Campylobacter coli* in environmental water samples by polymerase chain reaction. *Appl. Environ. Microbiol.* 59: 4090-4095.

Palmer, C.J., Tsai, Y.-L., Paszko-Kolva, C., Mayer, C. and Sangermano, L.R. 1993. Detection of *Legionella* species in sewage and ocean water by polymerase chain reaction, direct fluorescent-antibody, and plate culture methods. *Appl. Environ. Microbiol.* 59: 3618-3624.

Payment, P. 1993. Viruses: Prevalence of Disease, Levels, and Sources, pp. 99-144. *In: Safety in Water Disinfection: Balancing Chemical & Microbial Risks*, G.F. Craun, (ed.), ILSI Press, Washington, D.C.

Payment, P. and Franco, E. 1993. *Clostridium perfringens* and somatic coliphages as indicators of the efficiency of drinking water treatment for viruses and protozoan cysts. *Appl. Environ. Microbiol.* 59: 2418-2424.

Pepper, I.L., Josephson, K.L., Bailey, R.L. and Gerba, C.P. 1991. Detection of bacterial pathogens in water: comparison of culturable methodology with polymerase chain reaction technology. Proceedings of the Water Quality Technological Conference, 10-14 November 1991, Orlando, Fla., American Water Works Association, Denver, Colo., pp.101-107.

Pergram, G.C., Rollins, N. and Esprey, Q. 1998. Estimating the costs of diarrhoea and epidemic dysentery in KwaZulu-Natal and South Africa. *Wat. S.A.* 24: 11-20.

Pillai, S.D., Josephson, L., Bailey, R.L., Gerba, C.P. and Pepper, I.L. 1991. Rapid method for processing soil samples for polymerase chain reaction amplification of specific gene sequences. *Appl. Environ. Microbiol.* 57: 2283-2286.

Popovic, T., Fields, P.I., Olsvik, Ø., Wells, J.G., Evins, G.M., Cameron, D.N., Farmer III, J.J., Bopp, C.A., Wachsmuth, K., Sack, R.B., Albert, M.J., Nair, G.B., Shimada, T. and Feeley, J.C. 1995. Molecular subtyping of toxigenic *Vibrio cholerae* O139 causing epidemic cholera in India and Bangladesh, 1992-1993. *J. Infect. Dis.* 171: 122-127.

Prier, R. and Solnick, J.V. 2000. Foodborne and waterborne infectious diseases. *Postgrad. Medicine* 107: 245-255.

Pyle, B.H., Broadway, S.C. and McFeters G.A. 1995. A rapid, direct method for enumerating respiring enterohemorrhagic *Escherichia coli* O157:H7 in water. *Appl. Environ. Microbiol.* 61: 2614-2619.

Quadri, F., Hasan, J.A.K., Hossain, J., Chowdhury, A., Begum, Y.A., Azim, T., Loomis, L., Sack, R.B. and Albert, M.J. 1995. Evaluation of monoclonal antibody-based kit Bengal SMART for rapid detection of *Vibrio cholerae* O139 synonym Bengal in stool samples. *J. Clin. Microbiol.* 33: 732-734.

Quadri, F., Chowdury, A., Hossain, J., Chowdury, K., Azim, T., Shimada, T., Islam, N.K.M., Sack, R.B. and Albert, M.J. 1994. Development and evaluation of rapid monoclonal antibody based coagulation test for direct detection of *Vibrio cholerae* O139 synonym Bengal in stool samples. *J. Clin. Microbiol.* 32: 1589-1590.

Rafii, F., Holland, M.A., Hill, W.E. and Cerniglia, C.E. 1995. Survival of *Shigella flexneri* on vegetables and detection by polymerase chain reaction. *J. Food Protec.* 58: 727-732.

Rafii, F. and Lunsford, P. 1997. Survival and detection of *Shigella flexneri* in vegetables and commercially prepared salads. *J. AOAC Int.* 80:1191-1197.

Ramamurthy, T., Pal, A., Bag, P.K., Bhattacharya, S.K., Nair, G.B., Kurozano, H., Yamasaki, S., Shirai, H., Takeda, T., Uesaka, Y., Horigome, K. and Takeda, Y. 1993. Detection of cholera toxin gene in stool specimens by polymerase chain reaction: Comparison with bead enzyme-linked immunosorbent assay and culture method for laboratory diagnosis of cholera. *J. Clin. Microbiol.* 31: 3068-3070.

Rehnstam, A.S., Norqvist, A., Wolf-Watz, H. and Hagstrom, A. 1989. Identification of *Vibro anoillarum* in fish by using partial 16S rRNA sequences and a specific 16S RNA oligonucleotide probe. *Appl. Environ. Microbiol.* 55: 1907-1910.

Reissbrodt, R., Heier, H., Tschäpe, H., Kingsley, R. A. and Williams, P.H. 2000. Resuscitation by ferrioxamine E of stressed *Salmonella enterica* serovar Typhimurium from soil and water microcosms. *Appl. Environ. Microbiol.* 66: 4128-4130.

Ridgway, J.W. 1984. Bacterial recovery from water, sewage, and sewage effluents, pp.373. *In: The Revival of Injured Microbes*, M.H.E. Andrew and A.D. Russel, (eds.), Academic Press, Orlando.

Romaniuk, P.J. and Trust, T.J. 1987. Identification of *Campylobacter* species by Southern hybridization of genomic DNA using an oligonucleotide probe for 16S rRNA genes. *FEMS Microbiol. Lett.* 43: 331-335.

Rose, J. 1993. Enteric Waterborne Protozoa: Hazard and Exposure Assessment, pp. 115-126. *In: Safety of Water Disinfection: Balancing Chemical and Microbial Risks*, G. Craun, (ed.), ILSI Press, Washington, D.C.

Rose, J.B. 1988. Occurrence and significance of cryptosporidium in water. *J. AWWA* 80: 53-58.

Rose, J.B., Gerba, C.P. and Jakubowski, W. 1991. Survey of potable water supplies for *Cryptosporidium* and *Giardia*. *Environ. Sci. Technol.* 25:1393-1400.

Rosenberg, M.L., Hazlet, K.K., Schaefer, J., Wells, J.G. and Pruneda, R.C. 1976. Shigellosis from swimming. *J. Am. Med. Ass.* 236:1849-1852.

Rossen, L., Norskov, P., Holmstrom, K. and Rasmussen, O.F. 1992. Inhibition of PCR by components of food samples, microbial diagnostic assays and DNA-extraction solutions. *Int. J. Food Microbiol.* 17: 37-45.

Roszak, D.B. and Colwell, R.R. 1987. Survival strategies of bacteria in the natural environment. *Microbiol. Rev.* 51:365-379.

Rowe, B. and Gross, R.J. 1984. Facultatively anaerobic gram negative rods. Genus II. *Shigella*, pp. 423-427. In: Bergey's Manual of Systematic Bacteriology, N.R. Krieg and J.G. Holt (eds.), Williams & Wilkins, Baltimore.

Saïd, B., Smith, H.R., Scotland, S.M. and Rowe, B. 1995. Detection and differentiation of the gene for toxin co-regulated pili (*tcpA*) in *V. cholerae* non-O1 using the polymerase chain reaction. *FEMS Microbiol. Lett.* 125: 205-210.

Saiki, R.K., Gelfand, D.H., Stoffel, S., Scharf, S.J., Higuchi, R., Horn, G.T., Mullis, K.B. and Erlich, H.A. 1988. Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science* 239: 487-491.

Sakazaki, R. and Balows, A. 1981. The genera *Vibrio*, *Pleisiomonas*, and *Aeromonas*, pp. 1272-1301. In: The Prokaryotes. Vol. 2, M.P. Starr, H. Stolp, H.G. Trüper, A. Balows and H.G. Schlegel (eds.), Springer-Verlag, New York.

Salyers, A.A. and Whitt, D.D. 1994. Bacterial Pathogenesis. A Molecular Approach. A.S.M.Press, Washington DC.

Sambrook, J., Fritsch, E.F. and Maniatis, T. 1989. Molecular cloning: A Laboratory Manual. Cold Spring Harbor Laboratory Press. Cold Spring Harbor.

Samonis, G., Elting, L., Skoulika, E., Maraki, S. and Tselentis, Y. 1994. An outbreak of diarrhoeal disease attributed to *Shigella sonnei*. *Epidemiol Infect.* 112: 235-245.

Sandery, M., Stinear, T. and Kaucner, C. 1996. Detection of pathogenic *Yersinia enterocolitica* in environmental waters by PCR. *J. Appl. Bact.* 80: 327-332.

Sansonetti, P.J., Kopecko, D.J. and Formal, S.B. 1982. Involvement of a plasmid in the invasive ability of *Shigella flexneri*. *Infect. Immun.* 35: 852-860.

Sansonetti, P.J. 1999. *Shigella* plays dangerous games. *ASM News.* 65: 661-617.

Sasaki, K., Tajiri, Y., Sata, M., Fujii, Y., Matsubara, F., Zhao, M., Shimizu, S., Toyonaga, A. and Tanikawa, K. 1999. *Helicobacter pylori* in the natural environment. *Scand. J. Infect. Dis.* 31: 275-279.

Schleifer, K-H., Ludwig, W., Kraus, J. and Festl, H. 1985. Cloned ribosomal ribonucleic acid genes from *Pseudomonas aeruginosa* as probes for conserved deoxyribonucleic acid sequences. *Int. J. Syst. Bacteriol.* 35:231-236.

Schwab, K.J., de Leon, R., and Sobsey, M.D. 1996. Immunoaffinity concentration and purification of waterborne enteric viruses for detection by reverse transcriptase PCR. *Appl. Environ. Microbiol.* 62: 2086-2094.

Sellwood, J. and Dadswell, J.V. 1991. Human viruses and water, p. 29. *In: Current Topics in Clinical Virology*, P. Morgan-Capner, (ed.), The Laverham Press, England.

Sethabutr, O., Venkatesan, M., Murphy, G.S., Eampokalap, B., Hoge, C.W. and Echeverria, P. 1993. Detection of *Shigellae* and enteroinvasive *Escherichia coli* by amplification of the invasion plasmid antigen H DNA sequence in patients with dysentery. *J. Infect. Dis.* 167: 458-461.

- Shangkuan, Y.H., Show, Y.S. and Wang, T.M. 1995. Multiplex polymerase chain reaction to detect toxigenic *Vibrio cholerae* and to biotype *Vibrio cholerae* O1. *J. Appl. Bact.* 79: 264-273.
- Shawar, R.M., El-Zaatari, F.A.K., Nataraj, A. and Clarridge, J.E. 1993. Detection of *Mycobacterium tuberculosis* in clinical samples by two-step polymerase chain reaction and nonisotopic hybridization methods. *J. Clin. Microbiol.* 31: 61-65.
- Shears, P. 1996. Review: *Shigella* infections. *Annals of Tropical Medicine and Parasitology.* 90: 105-114.
- Shepherd, K. M. and Wyn-Jones, A. P. 1995. Evaluation of different filtration techniques for the concentration of *Cryptosporidium oocysts* from water. *Wat. Sci. Tech.* 31: 425-429.
- Shimada, T., Sakazaki, R., Fujimura, S., Niwani, K., Mishina, M. and Takizawa, K. 1990. A new selective differential agar medium for the isolation of *Vibrio cholerae* O1: PMT (polymixin mannose-tellurite agar). *Jap. J. Med. Sci. Biol.* 43: 37-41.
- Shirai, H., Nishibuchi, M., Ramamurthy, T., Bhattacharya, S.K., Pal, S.C. and Takeda, Y. 1991. Polymerase chain reaction for detection of the cholera enterotoxin operon of *Vibrio cholerae*. *J. Clin. Microbiol.* 29: 2517-2521.
- Siddique, A.K., Salam, A., Islam, M.S., Akram, K., Majumdar, R.N., Zaman, K., Fronczak, N. and Laston, S. 1995. Why treatment centers failed to prevent cholera deaths among Rwandan refugees in Goma, Zaire. *Lancet.* 345: 359-361.
- Simmons, D.A.R. and Romanowska, E. 1987. Structure and biology of *Shigella flexneri* O antigens. *J. Med. Microbiol.* 23: 289-302.

Sinigalliano, C.D., Kuhn, D.N. and Jones, R.D. 1995. Amplification of the *amoA* gene from diverse species of ammonium-oxidizing bacteria and from an indigenous bacterial population from seawater. *Appl. Environ. Microbiol.* 61: 2702-2706.

Smith, H.V., Robertson, I.J. and Ongerthm, J.E. 1995. Cryptosporidiosis and giardiasis: The impact of waterborne transmission. *J. Water Supply Res. Technol. Aqua.* 44: 258-274.

Sobsey, M.D. 1994. Molecular methods to detect viruses in environmental samples, pp. 387-400. *In: Rapid Methods and Automation in Microbiology and Immunology*, R.C. Spencer, E.P. Wright and S.W.B. Newsme (eds.), Intercept Ltd., Andover, England.

Sorvillo, F.J., Fujioka, K., Nahlen, B., Tormey, M.P., Keababjian, R. and Mascola, L. 1992. Swimming-associated cryptosporidiosis. *Am. J. Public Health* 82: 742-744.

Spangler, B.D. 1992. Structure and function of cholera toxin and related *Escherichia coli* heat-labile enterotoxin. *Microbiol. Rev.* 56: 622-647.

Starnbach, M.N., Falkow, S. and Tompkins, L.S. 1989. Species-specific detection of *Legionella pneumophila* in water by DNA amplification and hybridization. *J. Clin. Microbiol.* 27: 1257-1261.

Steffen, R.J. and Atlas, R.M. 1991. Polymerase chain reaction: Applications in environmental microbiology. *Annu. Rev. Microbiol.* 45: 137-161.

Stone, G.G., Oberst, R.D., Hays, M.P., McVey, S. and Chengappa, M.M. 1994. Detection of *Salmonella* serovars from clinical samples by enrichment broth cultivation-PCR procedure. *J. Clin. Microbiol.* 32: 1742-1749.

Strachan, T. and Read, A.P. 1996. PCR-based DNA cloning and DNA analysis, pp. 129-145. *In: Human Molecular Genetics*. BIOS Scientific Publishers Ltd., Oxford, United Kingdom.

Tauxe, R.Y. and Blake, P.A. 1992. Epidemic cholera in Latin America. *JAMA* 267: 1388-1390.

Tauxe, R., Seminario, L., Tapia, A.R. and Libel, M. 1994. The Latin American epidemic, pp. 321-344. *In: Vibrio cholerae and Cholera*, I.K. Wachsmuth, P.A. Blake and Ø. Olsvik, (eds.), American Society for Microbiology Press. Washington D.C.

Taylor, D.N., Bodhidatta, L. and Echeverria, P. 1991. Epidemiologic aspects shigellosis and other causes of dysentery in Thailand. *Rev. Infect. Dis.* 13:(supl. 4) S226-S230.

Tenover, F.C. and Unger, E.R. 1993. Nucleic acid probes for detection and identification of infectious agents, pp. 3-25. *In: Diagnostic Molecular Microbiology. Principles and Applications*. D. H. Persing, T.F. Smith, F.C. Tenover, T.J. White, (eds)., American Society for Microbiology Press, Washington D.C.

Tibbetts, J. 2000. Water world 2000. *Environ. Hlth. Perspect.* 108: no. 2.

Toranzos, G.A., Alvarez, A.J. and Dvorsky, E.A. 1993. Application of the polymerase chain reaction to the detection of pathogens in water. *Wat. Sci. Tech.* 27: 207-210.

Toro, A., Gonzales, N., Torres, J., Dvorsky, E. and Toranzos, G.A. 1995. Modified culture methods for detection of *Vibrio* spp. from estuarine water. *Wat. Sci. Tech.* 31: 283-290.

Toze, S. 1999. PCR and the detection of microbial pathogens in water and wastewater. *Wat. Res.* 33: 3545-3556.

Tsai, Y.L. and Olson, B.H. 1992. Rapid method for separation of bacterial DNA from humic substances for polymerase chain reaction. *Appl. Environ. Microbiol.* 58: 2291-2295.

U.S. EPA. 1992. Guidelines for water reuse. United States Environmental Protection Agency, Washington D.C.

Vantarakis, A., Komninou, G., Venieri, D. and Papapetropoulou, M. 2000. Development of a multiplex PCR detection of *Salmonella* spp. and *Shigella* spp. in mussels. *Lett. Appl. Microbiol.* 31: 105-109.

Varela, P., Rivas, M., Binsztein, N., Cremona, M.L., Herrmann, P., Burrone, O., Ugalde, R.A. and Frasch, A.C.C. 1993. Identification of toxigenic *Vibrio cholerae* from the Argentine outbreak by PCR for *ctxA1* and *ctxA2-B*. *FEMS Microbiol. Lett.* 315: 74-76.

Venkatesan, M.M., Buysse, J.M. and Hartman, A.B. 1991. Sequence variation in two *ipaH* genes of *Shigella flexneri* 5 and homology to the LRG-like family of proteins. *Mol. Microbiol.* 5: 2435-2445.

Venkatesan, M., Buysse, J.M., Vandendries, E. and Kopecko, D.J. 1988. Development and testing of invasion-associated DNA probes for detection of *Shigella* spp. and enteroinvasive *Escherichia coli*. *J. Clin. Microbiol.* 26: 261-266.

Venkatesan, M.M., Buysse, J.M. and Kopecko, D.J. 1989. Use of *Shigella flexneri ipaC* and *ipaH* gene sequences for the general identification of *Shigella* spp. and enteroinvasive *Escherichia coli*. *J. Clin. Microbiol.* 27: 2687-2691.

Vickers, R.M., Stout, J.E. and Yu, V.L. 1990. Failure of a diagnostic monoclonal immunofluorescent reagent to detect *Legionella pneumophila* in environmental samples. *Appl. Environ. Microbiol.*, 56: 2912-2914.

Vilaginès, P., Sarrette, B., Champsaur, H., Hugues, B., Doubrou, S., Joret, J.C., Laveran, F., Lesne, J., Paquin, J.L., Delattre, H.M., Oger, C., Alame, J., Grateloup, I., Perrollet, H., Serceau, R., Sinègre, F. and Vilaginès, R. 1997. Round robin investigation of glass wool method for poliovirus recovery from drinking water and sea water. *Wat. Sci.Tech.* 35: 445-560.

Waage, A.S., Vardund, T., Lund, V. and Kapperud, G. 1999. Detection of small numbers of *Campylobacter jejuni* and *Campylobacter coli* cells in environmental water, sewage, and food samples by a seminested PCR. *Appl. Environ. Microbiol.* 65: 1636-1643.

Waldor, K.M. and Mekalanos, J.J. 1996. Lysogenic conversion by a filamentous phage encoding cholera toxin. *Science* 272: 1920-1914.

Wand, H., Laht, T., Peters, M.M., Becker, P.M., Stottmeister, U. and Heinaru, A. 1997. Monitoring of biodegradative *Pseudomonas putida* strains in aquatic environments using molecular techniques. *Microb. Ecol.* 33: 124-133.

Warner, J.M. and Oliver, J.D. 1998. Randomly amplified polymorphic DNA analysis of starved and viable but not culturable *Vibrio vulnificus* cells. *Appl. Environ. Microbiol.* 64: 3025-3028.

Watanabe, H. and Timmis, K.N. 1984. A small plasmid in *Shigella dysenteriae* 1 specifies one or more functions essential for O antigen production and bacterial virulence. *Infect Immun.* 43: 391-396.

Way, J.S., Josephson, K.L., Pillai, S.D., Abbazadegan, M., Gerba, C.P. and Pepper, I.L. 1993. Specific detection of *Salmonella* spp. by multiplex polymerase chain reaction. *Appl. Environ. Microbiol.* 59: 1473-1479.

Weaver, J.W. and Rowe, M.T. 1997. Effect of non-target cells on the sensitivity of the PCR for *Escherichia coli* O157:H7. *Lett. Appl. Microbiol.* 25:109-112 .

Whittam, T.S. 1998. Evolution of *Escherichia coli* O157:H7 and other shiga toxin-producing *E. coli* strains, pp.195-209. In: *Escherichia coli* O157:H7 and other shiga-toxin-producing *E. coli* strains, J.B. Kaper and A.D. O'Brien, (eds.), American Society Microbiology Press, Washington D.C.

WHO, 1997. Report on global surveillance of epidemic prone infectious diseases at the end of 1997. www.who.int/emc-documents/surveillance/docs/whocdsrisr2001.html

Wilde, J., Eiden, J. and Yolken, R. 1990. Removal of inhibitory substances from human fecal specimens for detection of group A rotaviruses by reverse transcriptase and polymerase chain reactions. *J. Clin. Microbiol.* 28: 1300-1307.

Wilson, I.G. 1997. Inhibition and facilitation of nucleic acid amplification. *Appl. Environ. Microbiol.* 63: 3741-3751.

Wright, A.C., Miceli, G.A., Landry, W.L., Christy, J.B., Watkins, W.D. and Morris, J.G. 1993. Rapid identification of *Vibrio vulnificus* on nonselective media with an alkaline phosphatase-labeled oligonucleotide probe. *Appl. Environ. Microbiol.* 59: 541-546.

Yavzori, M., Cohen, D., Wasserlauf, R., Ambar, R., Rechavi, G. and Ashkenazi, S. 1994. Identification of *Shigella* species by DNA amplification of different loci of the *Shigella* virulence plasmid. *Eur.J. Clin. Microbiol. Infect. Dis.* 13: 232-237.

Papers published and congress contributions during the course of this study

Papers published:

Detection of toxigenic *Vibrio cholerae* from environmental water samples by an enrichment broth cultivation-pit-stop semi-nested PCR procedure. 2000. Theron, J., Celliers, J., du Preez, M., Brözel, V.S. and Venter, S.N. *J. Appl. Microbiol.* 89:539-546.

A sensitive seminested PCR method for the detection of *Shigella* in spiked environmental water samples. 2001. Theron, J., Morar, D., du Preez, M., Brözel, V.S. and Venter, S.N. *Wat. Res.* 35:869-874.

Contributions to conference proceedings:

Detection of small numbers of toxigenic *Vibrio cholerae* from environmental water samples by a novel polymerase chain reaction procedure. Cilliers, J., du Preez, M., Venter, S.N., Brözel, V.S. and Theron, J. WISA Biennial Conference and Exhibition. Sun City 28May-1 June 2000.

Direct polymerase chain reaction detection of entero-invasive *Shigella* spp. and *E. coli* in environmental water samples. Morar, D., du Preez, M., Theron, J., Brözel, V.S. and Venter, S.N. WISA Biennial Conference and Exhibition. Sun City 28May-1 June 2000.

Rapid detection of *Shigella flexneri* from environmental water samples by an enrichment broth cultivation-PCR procedure. Morar, D., du Preez, M., Theron, J., Brözel, V.S. and Venter, S.N. Bio Y2k Conference, Grahamstown. 23-28 January 2000.

Rapid compact polymerase chain reaction assay for detection of toxigenic *Vibrio cholerae* in environmental water samples. Cilliers, J., du Preez, M., Venter, S.N., Brözel, V.S. and Theron, J. Bio Y2k Conference, Grahamstown. 23-28 January 2000.