

REFERENCES

- Ahmed J & Bonham CD 1982. Optimal allocation in multivariate double sampling for biomass estimates. *J. Range Manage.* 35 : 777 – 779.
- Alipayo D Valdez R Holecheck JL & Cardinas M 1992. Evaluation of microhistological analysis for determining ruminant diet botanical composition. *J. Range Manage.* 45: 71 – 76.
- Anthony RG & Smith NS 1974. Comparisons of rumen and fecal analysis to describe deer diets. *J. Wildl. Manage.* 38 : 534 – 540.
- AOAC 1996. Official Methods of Analysis. *Association of Official Analytical Chemists. Washington DC.*
- APRU 1971. Bone meal and salt supplementation of Tswana steers in Kweneng District of Botswana. Technical Report. *Animal Production Research Division. Ministry of Agriculture. Gaborone. Botswana. Pp. 1 –8.*
- APRU 1977. Animal Production and Range Research Unit Annual Report. *Dept.Agric. Research. Ministry of Agriculture. Botswana*
- APRU 1980. Beef production and range management in Botswana. *Animal Production Research Division. Ministry of Agriculture. Gaborone. Botswana. Pp. 26 – 31.*
- APRU 1985. Animal Production and Range Research Unit Annual Report. *Dept.Agric. Research. Ministry of Agriculture. Botswana*
- Aucamp AJ 1981. Animal production off veld. *Proc. Grassl. Soc. sth Afr.* 16 : 17 – 18.
- Aucamp AJ Danckwerts JE Teague WR & Venter JJ 1983. The role of Acacia karroo in the False Thornveld of the Eastern Cape. *Proc. Grassl. Soc. sth Afr.* 18 : 151 – 154.
- Aucamp AJ Danckwerts JE & Venter JJ 1984. The production potential of an *Acacia karroo* community utilized by cattle and goats. *J. Grassl. Soc. sth. Afr.* 1 : 29 – 32
- Barbour MG Burk JH & Pitts WD 1980. Terrestrial Plant Ecology. *The Benjamin / Cummings Publishing Company, Inc.*
- Barnes DL & McNeill L 1978. Rainfall variability and animal production in the semi –arid savanna of Southern Africa. *Proc. Grassl. Soc. Sth. Afr.* 13 : 59 – 63.
- Barnes DL Odendaal JJ & Beukes BH 1982. Use of dry – weight – rank method of botanical analysis in Eastern Transvaal Highveld. *Proc. Grassl. Soc. sth. Afr.* 17 : 79 – 82.

- Bawden MG & Stobbs RA 1963. The land resources of Eastern Bechuanaland Protectorate. *Directorate of Overseas Survey. Tolworth, Surrey, England.*
- Bergstrom R & Skarpe C 1985. Characteristics of the Kalahari sand in Western Botswana. *J. of Arid Environ. 11 : 147 – 164.*
- Blair Rain A & Yalala AM. 1972. The Central and Southern States, Botswana. *Land Resources Study no. 11. Resources Division. ODA. UK.*
- Bouttom WT Tieszen LL & Imbamba SK 1988. Seasonal changes in nutrition of East African grassland vegetation. *Afr. J. Ecol. 26 : 103 – 115.*
- Brady NC 1974. The nature and properties of soils. 8th Ed. *Macmillan Publishing Co. Inc. NY.*
- Bransby DI 1981. The value of veld and pasture as animal feed. In: Tainton NM (ed.) Veld and Pasture management in South Africa. *Shuter and Shooter. Pietermaritzburg.*
- Bransby DI Matches AG & Krause GF 1977. Disc meter for rapid estimation of herbage yield in grazing trials. *Agron. J. 69 : 393 – 396.*
- Bransby DI & Tainton NM 1977. The disc meter. Possible application in grazing management. *Proc. Grassl. Soc. Sth. Afr. 12 : 393 – 396.*
- Buckley R Wasson R & Gubb A 1967. Phosphorus and potassium status of arid dune field soil in central Australia and Southern Africa, and biogeographic implications. *J. of Arid Environ. 13 : 211 – 216.*
- Central Statistics Office 1986. HIES. Household income and survey. *Min. of Finance & Dev. Planning. Botswana*
- Church DC 1986. Livestock feeds and feeding. (2nd Edition). *Prentice – Hall, New Jersey.*
- Cook CW 1964. Collecting forage samples representative of ingested materials of grazing animals for nutritional studies. *J. Animal Sc. 23 : 265 – 270.*
- Cook CW Kothmann M & LE Harris 1965. Effects of range condition and utilization on nutritive intake of sheep on summer ranges. *J. Range Manage. 18: 69 – 73.*
- Cook CE & Stubbendieck J 1986. Range Research Basic Problems and Techniques. *Am. Soc. for Range Mange.*
- Cooper SM 1982. Feeding behavior of goats and impala. *Proc. Grassl. Soc. sth. Afr. 17 : 117 – 121.*
- Cooper WM & Owen – Smith N 1985. Condensed tannins deter feeding by browsing ruminants in South African savanna. *Oecologia. 67 : 142 – 146.*

- Coughenour MB 1985. Graminoid response to grazing by large herbivores : adaptation, exaptations and interacting processes. *Ann. Miss. Bot. Gard.* 72 : 852 – 863.
- Daines T 1976. Grazing sequence pattern and species selection by cattle in the Dohne Sandveld M.Sc. Thesis. *Rhodes University*.
- Dean S Burkhardt JW & Meewig RO 1981. Estimating twig and foliage biomass of sagebrush, bitterbrush and rabbitbrush in the Great Plains. *J. Range Manage.* 4 : 224 - 227.
- Dearden BL Pegan RE & Hansen RM 1975. Precision of microhistological estimates of ruminants food habits. *J. Wildl. Manage.* 39 : 402 - 407.
- D.H.V. Consulting Engineers 1980. Countrywide Animal and Range Assessment Project. Final Report. *Ministry of Agriculture. Botswana*
- Drewes RH 1984. Smuts Finger grass. Farming on South Africa. *Dept. Agric. Pretoria Bulletin no. 115.*
- Du Toit PF 1968. A preliminary report on the effect of Acacia karroo competition on the composition and yield of sweet grassveld. *Proc. Grassl. Soc. sth Afr.* 3 : 147 – 149.
- Du Toit PF 1972. The goat in bush – grass communities. *Proc. Grassl. Soc. sth Afr.* 7 : 44 – 50.
- Dyksterhuis EJ 1949. Condition and management of rangeland on quantitative ecology. *J. Range Manage.* 2 : 104 – 112.
- Ellison L 1960. Influence of grazing on plant succession of rangelands. *Bot. Rev.* 26 : 71 – 78.
- Ernst WHO & Tolsma DJ 1989. Mineral nutrients in some Botswana savanna types. *British Ecol. Society. Special publication no. 9. Blackwell Scientific Publ.*
- FAO 1990. Explanatory notes on soil map of Republic of Botswana. *Soil mapping and advisory services. Botswana.*
- FAO 1991. Explanatory notes on vegetation of Republic Botswana. *Soil Mapping and advisory services. Botswana*
- Ferreira NA & Bigalke RC 1987. Food selection by grey rhebuck in the Orange Free State. *S. Afr. J. Wildl. Res.* 17 : 123 - 127.
- Field D 1977. A handbook of range ecology for range management in Botswana. *Land Utilization Division. Ministry of Agriculture. Botswana.*

- Foran BD, Tainton NM & Booysen P de V 1978. The development of a method for assessing veld condition in three grassveld types in Natal. *Proc. Grassl. Soc. sth. Afr.* 13 : 27 - 33.
- Fourie JH Engels EA & De Bruyn HL 1986. The influence of stocking rate and grazing system on the crud protein content and digestibility of *Tarchonanthus* veld in Northen Cape. *J. Grassl. Soc. South. Afr.* 3 : 62 – 69.
- Fourie JH Redelinghuys GJ & Opperman DPJ 1984. Influence of stocking rate and grazing system on plant basal cover and botanical composition of veld in the Northern Cape. *J. Grassl. Soc. sth. Afr.* 4 : 20 - 24.
- Free JC Hansen RM & Sims PL 1970. Estimating dry weights of food plants in feces of herbivores. *J. Range Manage.* 23 : 300 - 303.
- Fusco M Holecheck JL Tembo A Alipayo D & Cardenas M 1995. Grazing influence on water point vegetation in the Chihuahuan desert. *J. Range Manage.* 48 : 32 – 88.
- Gammon DM & Roberts BR 1978. Patterns of deloliation during continous and rotational grazing of the Matopo sandveld of Rhodesia. 1. Selectivity of grazing. 2. Severity of grazing. 3. Frequency of defoliation. *Rhod. J. Agric Res.* 16 : 117 - 126.
- Georgiadis NJ & McNaughton SJ 1990. Element and fibre contents of savanna grasses : Variation with grazing , soil type, season and species. *J. Appl. Ecol.* 27 : 623 – 634.
- Grossman D Holden PL & Collinson RFH 1999. Veld management on the game ranch. In: Veld management in South Africa (Edited by NM. Tainton). *Univ. of Natal Press, Scottsville.*
- Halls LK & Harlow RF 1971. Weight - length relations on flowering dogwood twigs. *J. Range Manage.* 24 : 236 - 237.
- Hanan NP Prevost Y Diouf A & Diallo O 1991. Assessment of desertification around deep wells in the Sahel using satellite imagery. *J. Appl. Ecol.* 28 : 173 – 186.
- Hanley TA 1982. The nutritional basis for food selection by ungulates. *J. Range Manage.* 35 : 146 – 151.
- Hansen RM Foppe TM Gilbert MB Clark RC & Renolds HB 1978. The microhistological analyses of feces as an estimate of herbivore dietary. *Composition Analysis Laboratory. Fort Collins. Colorado.*

- Hansen RM & Martin PS 1973. Ungulant diets in lower Grand Canyon. *J. Range Manage.* 26 : 380 – 381.
- Hardy MB & Mantis MT 1986. Grazing dynamics in sour grassveld. *S. Afr. J. Sci.* 82 : 566 – 572.
- Heady HF 1964. Palatability of herbage and animal preference. *J. Range Manage.* 17 : 76 – 82.
- Heithold DV Kiesling HE & Beck RF 1980. Nutritive value of forage on continuous and seasonally – grazed semi-arid range. *Abstr. 72nd Annual Meet. Amer. Soc.*
- Hercus BH 1960. Plant cuticle as an aid to determine the diet of grazing animal. *Proc. Intern. Grassl. Cong.* 8 : 443 – 447.
- Hilder EJ 1966. Distribution of excreta by sheep at pasture. *Proc. of Tenth Int. Grassland Congress.* pp. 977 – 979.
- Hobson FO & de Ridder CH 1991. Quantifying indices of potential competitiveness and browse production of single - stemmed Acacia karroo trees in South - East Africa. *J. Grassl. Soc. Sth. Afr.* 8 : 1 - 5.
- Holechek JL & Gross BD 1982. Training needed to quantify diets from fragmented range plants. *J. Range Manage.* 35 : 644 – 648.
- Holechek JL Pieper RD & Herbel CH 1998. Range Management. Principles and Practices. Third edition. *Prentice Hall, INC. New Jersey.*
- Holecheck JL Vavra M & Pieper RD 1982. Botanical composition of range herbivore diets. A Review. *J. Range Manage.* 35 : 309 - 315.
- Humphrey RR 1962. Range Ecology. *Ronald Press Co. NY.* pp. 193 - 194.
- Hyder DN 1969. The impact of domestic animals on structure of grassland ecosystems. In. “The grassland Ecosystem – a preliminary synthesis.” *Colorado State Univ. Range Sci. Dept Science series no. 2* pp. 243 – 260.
- Johnson MK & Pearson HA 1981. Esophageal, fecal and exclosure estimates of cattle diets on a longleaf pine - bluestem range. *J. Range Manage.* 34 : 232 – 235.
- Jordaan JJ Rethman NFG & Pieterse PA 1991. A technique to quantify available above – ground photomass of grass tufts in defoliation studies. *J. Grassl. Soc. Sth. Afr.* 8 : 189 – 192.

- Kelly RD 1977. The significance of the woody component of semi – arid savanna vegetation in relation to meat production . *Proc. Grassl. Soc. sth Afr. 12* : 105 – 108
- Krebs C J 1989. Ecology. The experimental analysis of distribution and abundance. Third Edition. *Harper & Row, Publishers, NY. Pp. 521.*
- Kreuter UP & Tainton NM 1989. The effects of continuous and rotational grazing of sourveld on the quality of herbage selected by Simmentaler heifers. *J. Grassl. Soc. South. Afr. 6* : 236 – 240.
- Krueger WC 1972. Evaluating animal forage preferences. *J. Range Manage. 25* : 271 - 275.
- Krueger WC Laycock WC & Price DA 1974. Relationships of taste, smell, sight and touch to forage selection. *J. Range Manage. 27* : 257 – 262.
- Lange RT 1969. The Piosphere. Sheep track and dung patterns. *J. of Range Manage. 22* ; 396 – 400.
- Laycock WC Buchanan H & Krueger WC 1972. Three methods of determining diet utilization and trampling damage on sheep range. *J. Range Manage. 25* : 352 – 356.
- Le Houérou H N 1980. The role of browse in the management of natural grazing land. In: International Symposium on Browse in Africa. Edited by H N Le Houérou. *International Livestock Center for Africa.*
- Leistner OA 1967. The plant ecology of the Southern Kalahari. *Memoir of Botanical Survey of South Africa.*
- Liversidge R 1970. Identification of grazed grass using epidermal characters. *Proc. Grassl. Soc sth Afr. 5* : 153 – 165.
- Louw GN. 1969 The nutritive value of natural grazing in South Africa. *Proc. Sth. Afr. Anim. Prod. 8* : 57 - 61
- Lyon JL 1970. Length and weight - diameter relations of serviceberry twigs. *J. Wildl.Manage. 34* : 456 - 460.
- Martin SC & Ward DE 1970. Rotating access to water improves semi desert cattle range near water. *J. Range Manage. 23* : 22 – 26.
- McInnis ML 1977. A comparison of four methods used in determining the diet of large herbivore.MSc Thesis. Oregon State Univ. Cited in: Holechek J. et al. 1982.

- Botanical composition of range herbivore diets. A Review. *J.Range Manage* 35 : 309 – 315.
- McInnis ML Vavra M & Krueger WC 1983. A comparison of four methods used to determine the diet of large herbivore. *J. Range Manage.* 36 : 302 – 306.
- McKay AD. 1968. Range production in Botswana. *E. Afr. For. J.* 34 : 178 – 193.
- McNaughton SJ 1990. Mineral nutrient and spatial concentration of African ungulants. *Nature.* 334 : 343 – 345.
- Morisita M 1959. Measuring of interspecific association and similarity between communities. Cited In: H. Wolda 1981. Similarity indices, sample size and diversity. *Oecologia.* 50 : 296 – 302
- Mueller – Dombis D. & Ellenberg H 1974. Aims and methods of vegetation ecology. *John Wiley & Sons, NY.*
- Omphile U 1997. Seasonal diet selection and quality of large ungulates in Chobe National Park, Botswana. Ph.D thesis. *Univ.of Wyoming. Laramie. USA.*
- O'Reagain PJ & Mentis MT 1988. Seasonal changes in the quality of diet selected by cattle grazing the Natal Sourveld. *J. Grassl. Soc. sth Afr.* 5 : 48 – 51.
- O'Reagain PJ & Mentis MT 1989. The effect of plant structure on the acceptability of different grass species to cattle. *J. Grassl. Soc. sth. Afr.* 6 : 163 – 170.
- O'Reagain PJ & Mentis MT 1990. The effect of veld condition on the quality of diet selected by cattle grazing the Natal sour sandveld. *J. Grassl. Soc. South. Afr.* 7 : 190 – 195.
- Owen – Smith N. 1999. Animal factor in veld management In: Veld management in South Africa. (Edited by N. Taiton). *Univ. of Natal Press. Scottsville.* pp118 – 124.
- Owen – Smit N & Novellie P 1982. What should a clever ungulate eat? *Amer. Nat.* 19 : 151 – 178.
- Palgrave KC 1977. Trees of Southern Africa. *Struik Publishing Co. Cape Town.*
- Pechanec JF & Pickford GD 1937. A weight estimate method for determination of range or pasture production. Cited in: Pieper RD 1978. Measurement techniques for herbaceous and shrubby vegetatin. *Dept. of Animal & Range Sci. New Mexico State Univ. Las Cruces. NM.*

- Perkins JS 1991. The impact of borehole dependent cattle grazing on environment and society of the Kalahari sandveld, Western Central District. Botswana. Ph. D Thesis. *Univ. of Sheffield UK.*
- Pienaar JP Tainton NM Van Ryssen JBJ & Swiegers JP 1993. Factors affecting the voluntary feed intake of sheep grazing Pennisetum clandestinum pastures :Observation in the animal. *Afr. J. Range For. Sci. 10 : 145 – 150.*
- Pieper RD 1978. Measurement techniques for herbaceous and shrubby vegetation. *Dept. of Animal & Range Sci. New Mexico State Univ. Las Cruces. New Mexico.*
- Pieper RD Montoya RJ & Groce VL 1971. Site characteristics of pinyon – juniper and blue grama ranges in southern – central New Mexico. *Bull. 573 New Mexico Agric. Exp. Sta.*
- Pike JG 1971. Rainfall over Botswana. Proceedings of the conference on sustained production from semi – arid areas. *Botswana Notes and Records 1 : 69 – 76.*
- Pratchett D 1978. Effects of bush clearing on grasslands in Botswana. *Proc. First Int. Grassl. Congr. Pp 667 – 670.*
- Pratchett D Capper BG Light DE Miller MD Rutherford AS Rennie TW Buck NG & Trail JC 1977. Factors limiting liveweight gain of beef cattle on rangeland in Botswana. *J. Range Manage. 30 : 442 – 445.*
- Pumphrey FW 1980. Precipitation, temperature and herbage relationships for pine woodlands site in Northern Oregon. *J. Range Manage. 33 : 307 – 310.*
- Roos JH Rethman NFG & Kotze GD 1973. Preliminary results on species selection by animals on sour grassland. *Proc. Grassl. Soc. sth. Afr. 8 : 77 – 81.*
- Rosiere RE Beck RF & Wallace JD 1975. Cattle diets on semi – arid rangeland: botanical composition. *J. Range Manage. 28 : 89 – 93.*
- Rutherford MC 1979. Plant based techniques for determining available browse and browse utilization: A Review. *Botanical Review. Vol. 45 No. 2.*
- Sanders KD Dahl BE & Scott G. 1980. Bite counts, vs. fecal analysis for range animal diets. *J. Range Manage. 32 : 146 – 149.*
- S.A.S. 1985. SAS User's Guide. Basics. Vision 5 Edition. *SAS Institute INC. Carey, N C.*
- S.A.S. 1990. SAS User's Guide. Basic. Vision 6 Edition. *SAS Institute INC. Carey, N C.*

- Scotcher JSB 1979. A review of faecal analysis techniques for determining the diet of wild grazing herbivores. *Proc. Grassl. Soc. sth. Afr.* 14 : 131 – 136.
- Shackleton CM & Mentis MT 1992. Seasonal changes in nutrient content under three defoliation treatments in two coastal grassland communities of Transkei. *J. Grassl. Soc. South. Afr.* 9 : 30 –37.
- Siderius W 1972. Soils of Eastern and Northern Botswana. *Tech.Document no.3 UNDP (SF). FAO project on surveys and Training for the Dev. of Water and Resources and Agric Prod.Botswana.*
- Skarpe C 1983. Cattle grazing and the ecology of the Western Kalahari. In: Which Way Botswana's Wildlife? Proceedings of the Symposium of the Kalahari Conservation Society. *Kalahari Conservation Society. Botswana. Pp 21 – 24.*
- .Smit GN 1989. Quantitative description of woody plant community : Part 1. An approach. *J. Grassl. soc. sth. Afr.* 6 : 186 - 191.
- Smit GN 1994. The influence of intensity of tree thinning on mopani veld. Ph D. Thesis. *University Pretoria. Dept. of Plant Prod. and Soil Sci.* pp.7 - 22.
- Smit GN 1996. BECVOL : Biomass Estimates from Canopy Volume. (Version 2) – User guide.Unpublished manual. *Univ. of Orange Free State. Bloemfontain.*
- Smit GN & Rethman NFG 2000. The influence tree thinning on soil water in a semi – arid savanna of Southern Africa. *J. Arid Environment.* 44 : 41 – 49.
- Smith AD & Shandruk IJ 1979. Comprison of fecal, rumen and utilization methods for assessing pronghorn diets. *J. Range Manage.* 32 : 275 – 279.
- Sparks DR & Malechek JC 1968. Estimating percentage dry weight in diets using a microscope technique. *J. Range Manage.* 21 : 264 - 265.
- Stebbins GL 1981. Co-evolution of grasses and herbivores. *Ann. Miss. Bot. Gard.* 68 : 75-86.
- Steel RGD & Torrie JH 1980. Principles and Procedures of Statistics.*McGrawHill Books. NY*
- Stewart DRM 1967. Analysis of plant epidermis in faeces. A technique for studying the foodpreferences of grazing herbivores. *J. Appl. Ecol.* 4 : 83 - 111.
- Stewart DRM & Stewart J 1970. Food preferences data by faecal analysis for African plain ungulates. *Zool. Afr.* 15 : 155 – 129.
- Storr GM 1961. Microscopic analysis of feces, a procedure for ascertaining the diets of herbivore mammals. *Austr. J. Biol. Sci.* 14 : 157 - 164.

- Stuart – Hill GC 1985. Competitive interactions between *Acacia karroo* and grass sward in the False Thornveld of the Eastern Cape. MSc Thesis. *Univ. of Natal. Peiternaritzberg*
- Stuart – Hill GC 1987. The influence of an *Acacia karroo* tree on grass production in its vicinity. *J. Grassl. Soc. Sth Afr.* 4 : 83 – 88.
- Stuart –Hill GC Aucamp AJ Le Roux CJG & Teague WR 1983. Towards a method of assessing the veld condition of the Valley Bushveld of the Eastern Cape. *J. Grassl. Soc. Sth Afr.* 3 : 19 – 24.
- Stuart – Hill GC & Tainton NM 1988. Browse and forage production in Eastern Cape Thornveld in response to tree size and defoliation frequency. *J. Grassl. Soc. sth Afr.* 5 :42 – 47.
- Sweet RJ & Mphinyane WN 1986. Preliminary observations on the ability of goats to control post – burning regrowth in *Acacia nigrescens/Combretum apiculatum* savanna in Botswana. *J. Grassl. Soc. sth Afr.* 3 : 79 – 84.
- Tadmor NH Brieghet A Noy Meir I Benjamin RW & Eyal E 1975. An evaluation of the calibrated weight – estimate method for measuring production of annual vegetation. *J. Range Manage.* 28 : 65 – 69.
- Teague WR 1983. Expected response of *Acacia karroo* Hayne to moisture stress and defoliation. *Proc. Grassl. sth Afr.* 18 : 147 – 150.
- Teague WR & Smit GN 1992. Relations between woody and herbaceous components and the effects of bush-clearing in Southern Africa. *J. Grassl. Soc. sth. Afr.* 9: 60 – 71.
- Teague WR Trollope WSW & Aucamp AJ 1981. Veld management in the semi – arid bush – grass communities of Eastern Cape. *J. Grassl Soc. sth Afr.* 16 : 23 – 28.
- Theron EP & P de V Booysen 1966. Palatability in grasses. *Proc. Grassl. Soc. sth Afr.* 1 : 111 – 120.
- Thrash I. 1998. Impact of water provision on herbaceous vegetation in Kruger National Park, South Africa. *J. Arid Environ.* 38 : 437 – 450.
- Todd JW & Hansen RM 1973. Plant fragments in feces of pronghorn as indicator of food habits. *J. Wildl. Manage.* 37 : 363 – 366.
- Tolsma DJ Ernst WH & Verwey RA 1987a. Nutrients in soil and vegetation around two artificial water points in Eastern Botswana. *J. Appl. Ecol.* 24 : 991 – 1000.

- Tolsma DJ Ernst WH Verweij RA & Vooijs R 1987b. Seasonal variation of nutrient concentration in semi-arid savanna ecosystem in Botswana. *J. Appl. Ecol.* 75: 755-60.
- Van der Merwe CR 1949. South African soil types. *Commonwealth Bureau of Soil Sci. Tech. Comm. No. 46..*
- Van Dyne GM & Heady HF 1965. Dietary chemical composition of cattle and sheep grazing in common on dry annual range. *J. Range Manage.* 18 : 78 – 86.
- Vavra M Rice RW & Hansen RM 1978. A comparison of esophageal fistula and fecal material to determine steer diets. *J. Range Manage.* 31 : 11 – 13.
- Van Dyne GM & Torell DF 1964. Developing and use of esophageal fistula : A Review. *J. Range Manage.* 17 : 7 – 17.
- Van Rooyen N Bredenkamp GJ Theron GK Bothma J Du P & LeRiche EAN 1994. Vegetational gradients around artificial watering points in the Kalahari Gemsbok National Park. *J. Arid Environ.* 26 : 349 – 361.
- Walker BH Ludwig D Holling CS & Peterman RM 1981. Stability of semi – arid grazing systems. *J. of Ecol.* 69 : 473 – 498.
- Weaver JE & Clements FE 1938 Plant Ecology. *Mc Graw –Hill, NY.*
- Westoby M 1974. Analysis of diet selection by large herbivore. *Amer. Nat.* 108 : 290 – 304.
- Weare PR & Yalala A 1971. Provisional vegetation map of Botswana. *Ministry of Agriculture. Botswana. Pp 18.*
- Williams OB 1969. An approved technique for identification of plant fragments in herbivore feces. *J. Range Manage.* 22 : 51 – 52.
- Wolda H 1981. Similarity indices, sample size and diversity. *Oecologia.* 50 : 296 – 302.
- Zumer Linde M 1976. Botswana. Can Desert Encroachment be Stopped? A study with emphasis on Africa. (Edited by A. Rapp, HN Lehouerou & B Landholm. pp171 – 187). *Ecological Bulletins No. 24. Swedish National Science. Research Council. Stockholm.*