

## References

- Abdulrazak, S.A., Muinga, R.W., Thorpe, W. and Orskov, E.R., 1997. Supplementation with *Gliricidia sepium* and *Leucaena leucocephala* on voluntary food intake, digestibility, rumen fermentation and live weight of crossbred steers offered *Zea mays* stover. *Livest. Prod. Sci.* 49, 53-62.
- Abdulrazak, S.A., Fujihara, T., Ondiek, J.K. and Orskov, E., 2000. Nutritive evaluation of some *Acacia* tree leaves from Kenya. *Anim. Feed Sci. Technol.* 85, 89-98.
- Abou El Nasr, H.M., Kandil, H.M., El Kerdawy, A., Dawlat, Khamis, H.S. and El-Shaer, H.M., 1996. Value of processed saltbush and acacia shrubs as sheep fodders under the arid conditions of Egypt. *Small. Rum. Res.* 24: 15-20.
- AFRC Agricultural Food and Research Council, 1993. Energy and Protein requirements of ruminants. An advisory manual prepared by the AFRC Technical committee on responses to nutrients. CAB International, Wallingford, UK.
- Aganga, A.A., Mtheho, J.K. and Tshwenyane, S., 2003. *Atriplex nummularia* (Old man saltbush): A potential forage crop for arid regions of Botswana. *Pakistan J. Nutr.* 2 (2): 72-75.
- Agricultural Research Council (ARC), 1984. The nutrient requirements of ruminant livestock (Suppl. 1). Commonwealth Agricultural Bureaux, Slough, United Kingdom.
- Ahn, J.H., Elliot, R. and Minson, D.J., 1988. Quality assessment of the fodder legume *Cassia rotundifolia*. *Trop. Grassl.* 22 (2): 63-67.
- Aitchison, E., Gill, M., France, J. and Dhanoa, M.S., 1986. Comparison of methods to describe the kinetics of digestion and passage of fibre in sheep. *J. Sci. Food & Agric.* 37: 1065-1072.
- Antoniewicz, A.M., Heinemann, W.W. and Hanks, E.M., 1980. The effect of changes in intestinal flow of nucleic acids on allantoin excretion in sheep urine. *J. Agric. Sci. Camb.* 95: 395-400.

AOAC, 2000. Official Methods of analysis (17<sup>th</sup> Ed.) Volume I. Association of Official Analytical Chemists, Inc., Maryland, USA.

Armstrong, G. and Gibbs, L., 2000. Handbook of useful trees and shrubs for rural areas of the winter rainfall region. *Dept. of Forest science, University of Stellenbosch* ([gvwyk@maties.sun.ac.za](mailto:gvwyk@maties.sun.ac.za)).

Aronsen, J.A., Pasternak, D. & Danon, A., 1985. Introduction and first evaluation of 120 halophytes. *Proc. Int. Res. & Dev. Conf. Tuscon, Arizona*.1985.

Bailey, E.M. Jr., 1985. Myopathies associated with *Cassia* spp. In "Plant Toxicology." *Proceedings of the Australia-USA Poisonous plants symposium*, Brisbane, Australia, pp395-400.

Barry, T.N. and Duncan, S.J., 1984. The role of condensed tannins in the nutritional value of *Lotus pendiculatus* for sheep-voluntary intake. *Br. J. Nutr.* 51: 485-491.

Barry, T.N. and Manley, T.R., 1986. Interrelationships between the concentration of total condensed tannin, free condensed tannin and lignin in lotus species and their possible consequences in ruminant nutrition. *J. Sci. Food & Agric.* 37: 248-254.

Bekker, J.J., 1995. Variation between lucerne cultivars in protein degradation and utilization by sheep. M.Sc. (Agric) dissertation, University of Pretoria, Pretoria, South Africa.

Benjamin, R.W., Lavie, Y., Forti, M., Barkai, D., Yonatan, R. and Hefetz, Y., 1995. Annual regrowth and edible biomass of two species of *Atriplex* and of *Cassia sturtii* after browsing. *J. Arid. Environm.* 29: 63-84.

Bonsi, M.L.K., Osuji, P.O. and Tuah, A.K., 1995. Effect of supplementing teff straw with different levels of *Leucaena* or *Sesbania* leaves on the degradability of teff straw, *Sesbania*, *Leucaena*, *Tagasaste* and *Vernonia* and on certain rumen and blood metabolites in Ethiopian Menz sheep. *Anim. Feed Sci. Technol.* 52, 101-129.

- Bouزيد, S.M. and Papanastasis, V.P., 1996. Effects of seeding rate and fertilizer on the establishment and growth of *Atriplex halimus* and *Medicago arborea*. *J. Arid Environm.s* 33: 109-115.
- Broderick, G.A. and Kang, J.H., 1980. Automated simultaneous determination of ammonia and total amino acids in ruminal fluid and *in vitro* media. *J. Dairy Sci.* 63: pp 64.
- Chen, X.B., Orskov, E.R. and Hovell, F.D., 1990. Excretion of purine derivatives by ruminants: endogenous excretion, differences between cattle and sheep. *Brit. J. Nutr.*, 63: 121-129.
- Chen, X. B., Chen, Y.K., Franklin, M.F., Orskov, E.R. and Shand, W.J., 1992. The effect of feed intake and body weight on purine derivative excretion and microbial protein supply in sheep. *J Anim Sci.*, 70:1534-1542.
- Chen, X.B. and Gomes, M.J., 1992. Estimation of microbial protein supply to sheep and cattle based on urinary excretion of purine derivatives- An overview of the technical details. *International feed resources unit. Rowett research institute, Bucksburn, Aberdeen, UK.* pp. 1-21.
- Chriyaa, A., Moore, K.J. and Waller, S.S., 1997. Intake, digestion and nitrogen balance of sheep fed shrub foliage and medic pods as a supplement to wheat straw. *Anim. Feed Sci. Technol.* 65, 183-196.
- Clark, J.H., Klusmeyer, T.H. and Cameron, M.R., 1992. Microbial protein synthesis and flows of nitrogen fractions to the duodenum of dairy cows. *J. Dairy Sci.* 75: pp 2304.
- Crampton, E.W., Donefer, E. and Lloyd, L.E., 1960. A nutritive value index for forages. *J. Anim. Sci.* 19: 538 – 544.
- Dean, W.R.J. and MacDonald, I.A.W., 1994. Historical stocking rates of domestic livestock as a measure of semi-arid and arid rangeland degradation in the Cape Province. *S. A. J. Arid Environm.* 26: 281-298.

De Kock, G.C., 1980. Drought resistant fodder shrubs in South Africa. Ed. Le Houerou, H.N., *Browse in Africa, the current state of knowledge*. pp 399-408. Paris: UNESCO.

De Visser, H., Van Der Togt, P.L., Huisert, H. and Tamminga, S., 1992. Structural and non-structural carbohydrates in concentrate supplements of silage-based dairy cow rations. 2. Rumen degradation, fermentation and kinetics. *Neth. J. Agric. Sci.* 40: 431-445.

Dewhurst, R.J. and Webster, A.J.F., 1988. Effects of manipulating rumen fermentation and outflow rate in sheep on microbial protein yield as estimated from allantoin excretion. *Anim. Prod.* 46: 490 (Abstract).

Dewhurst, R.J. and Webster, A.J.F., 1992. Effects of diet, level of intake, sodium bicarbonate and monensin on urinary allantoin excretion in sheep. *Br. J. Nutr.* 67:345-353.

Dicko, M.S. and Sangare, M., 1984. Feeding behaviour of domestic ruminants in Sahelian zone. *Proc. 2<sup>nd</sup> Int. Rangelands Congr. Adelaide, Australia* pp. 388-390.

Djouvinov, D.S. and Todorov, N.A., 1994. Influence of dry matter intake and passage rate on microbial protein synthesis in the rumen of sheep and its estimation by cannulation and a non-invasive method. *Anim. Feed Sci. Technol.* 48: 289-304.

Faucon, P., 2001. Desert Tropicals.

[http://www.desert-tropicals.com/plants/Fabaceae/Cassia\\_sturtii.html](http://www.desert-tropicals.com/plants/Fabaceae/Cassia_sturtii.html)

Firkins, J.L., Allen, M.S., Oldick, B.S. and St-Pierre, N.R., 1998. Modelling ruminal digestibility of carbohydrates and microbial protein flow to the duodenum. *J. Dairy. Sci.* 81:3350–3369.

Fonesca, A.J.M., Dias-da-silva, A.A. and Orskov, E.R., 1998. *In sacco* degradation characteristics as predictors of digestibility and voluntary intake of roughages by mature ewes. *Anim. Feed Sci. Technol.* 72: 205-219.

Fujihara, T., Orskov, E.R., Reeds, P.J. and Kyle, D.J., 1987. The effect of protein infusion on urinary excretion of purine derivatives in ruminants nourished by intragastric nutrition. *J. Agric. Sci.* 109: 7-12.

Goering, H.K. and Van Soest, P.J., 1970. Forage Fiber Analyses (Apparatus, Reagents, Procedures, and Some Applications). Agric. Handbook No. 379. USDA-ARS, Washington, DC.

Gomes, M.J., Hovell, F.D. DeB. and Chen, X.B., 1994. The effect of starch supplementation on microbial protein supply in sheep. *Anim. Feed Sci. Technol.* 49: 277-286.

Grovm, W.L. and Williams, V.J., 1977. Rate of passage of digesta in sheep. 6. The effect of level of food intake on mathematical predictions of the kinetics of digesta in the reticulorumen and intestines. *Brit. J. Nutr.* 38: 425-436.

Gurbuz, Y., 2007. Determination of nutritive value of leaves of several *Vitis vinifera* varieties as a source of alternative feedstuff for sheep using *in vitro* and *in situ* measurements. *Small. Rum. Res.* 71: 59-66.

Herbert, C.D., Flory, W., Seger, C. and Blanchard, R.E., 1983. Preliminary isolation of a myodegenerative toxic principle from *Cassia occidentalis*. *Am. J. Vet. Res.:* 44:1370-1374.

Hoffman, T.M. and Cowling, R.M., 1990. Desertification in the lower Sundays river valley, S. A. *J. Arid Environm.* 19: 105-117.

Hovell, F.D.D. and Greenhalgh, J.F.D., 1978. The utilization of diets containing acetate, propionate or butyrate salts by growing lambs. *Brit. J. Nutr.* 40, 171-182.

Huhtanen, P., 1988. The effects of barley, unmolassed sugar beet pulp and molasses supplements on organic matter, nitrogen and fibre digestion in the rumen of cattle given a silage diet. *Anim. Feed Sci. Technol.* 20 (4): 259 – 278.

Huhtanen, P. and Khalili, H., 1991. Sucrose supplements in cattle given grass silage based diet. 3. Rumen pool size and digestion kinetics. *Anim. Feed Sci. Technol.* 33: 275 – 287.

Ikhimioya, I., Isah. O.A., Ikhatua, U.J. and Bamikole, M.A., 2005. Rumen degradability of dry matter and crude protein in tree leaves and crop residues of humid Nigeria. *Pak. J. Nutr.* 4 (5): 313-320.

- Kaitho, R. J., Umunna, N. M., Nsahlai, I. V., Tamminga, S. and Van Bruchen, J., 1998. Utilization of browse supplements with varying tannin levels by Ethiopian Menz sheep, 1. Intake, digestibility and live weight changes. *Agroforestry Systems* 39: 145-159.
- Kallah, Muh.S., Bale, J.O., Abdullahi, U.S., Muhammad, I.R. and Lawal, R., 2000. Nutrient composition of native forbs of semi-arid and dry sub-humid savannas of Nigeria. *Anim. Feed Sci. Technol.* 84: 137-145.
- Kanani, J., Lukefahr, S.D. and Stanko, R.L., 2006. Evaluation of tropical forage legumes (*Medicago sativa*, *Dolchis lablab*, *Leucaena leucocephala* and *Desmanthus bicontortus*) for growing goats. *Small. Rum. Res.* 65: 1-7.
- Khalili, H., 1993. Supplementation of grass hay with molasses in crossbred (*Bos Taurus* x *Bos indicus*) non-lactating cows: Effect of level of molasses on feed intake, digestion, rumen fermentation and rumen digesta pool size. *Anim. Feed Sci. Technol.* 41: 23-38.
- Khazaal, K. and Orskov, E.R., 1994. The *in vitro* gas production technique: an investigation on its potential use with insoluble polyvinylpyrrolidone for assessment of phenolic-related antinutritive factors in browse species. *Anim. Feed Sci. Technol.* 47: 305-320.
- Komlong, M.K., Barber, D.G. and McNiell, D.M., 2001. Post ruminal protein supply and N retention of weaner sheep fed on a basal diet of lucerne hay (*Medicago sativa*) with increasing levels of quebracho tannins. *Anim. Feed. Sci. Technol.* 92: 59 -72.
- Leng, R.A., 1997. Tree foliage in ruminant nutrition. Food and Agriculture Organization of the United Nations Rome. <http://www.fao.org/docrep/003/w7448e/W7448E00.htm>
- Le Roux, P.M., Kotze, C.D., Nel, G.P. and Glen, H.F., 1994. Bossieveld. Grazing plants of the Karoo and Karoo-like areas. Bulletin 428. Published by the Dept. Agric., Pretoria, South Africa
- Lovegrove, B., 1993. The living deserts of Southern Africa. Published by Fernwood press (Pty) Ltd. South Africa pp. 224.

Macvicar, C.N., De Villiers, J.N., Loxton, R.F., Verster, E., Lamprechts, J.J.N., Merryweather, F.R., Le Roux, J., Van Rooyen, J.A. and Harmse, H.J., 1977. Soil classification: A binomial system for SA. Pretoria, 1<sup>st</sup> ed. Soil & Irrig. Res. Inst. Dept. Agric. Tech. Services.

Mathers, J.C. and Miller E.L., 1981. Quantitative studies of food protein degradation and the energetic efficiency of microbial protein synthesis in the rumen of sheep given chopped lucerne and rolled barley. *Br. J. Nutr.* 45: 587 – 604.

McAllan, A.B. and Smith, R.H., 1973. Degradation of nucleic acids derivatives by rumen bacteria *in vitro*. *Br. J. Nutr.* 29: 467-474.

McDonald, P., Edwards, R.A., Greenhalgh, J.F.D. and Morgan, C.A., 1996. Animal Nutrition, 5<sup>th</sup> edition, Longman Scientific & Technical, Essex, UK, 607pp.

McDonald, P., Edwards, R.A., Greenhalgh, J.F.D. and Morgan, C.A., 2002. Animal Nutrition 6<sup>th</sup> edition Published by Pearson Education Ltd, Edinburgh Gate, Harlow, Essex, UK, 693pp.

McMahon, L.R., McAllister, T.A., Berg, B.P., Majak, W., Acharya, S.N., Popp, J.D., Coulman, B. E., Wang, Y., and Cheng, K.J., 2000. A review of the effects of forage condensed tannins on ruminal fermentation and bloat in grazing cattle. *Can. J. Plant. Sci.* 80: 469–485.

McSweeney, C.S., Palmer, B., McNeill, D.M. and Krause, D.O., 2001. Microbial interactions with tannins: nutritional consequences for ruminants. *Anim. Feed Sci. Technol.* 91, 83-93.

Melaku, S., Peters, K.J. and Tegegne, A., 2004. Microbial nitrogen supply, nitrogen retention and rumen function in Menz sheep supplemented with dried leaves of multipurpose trees, their mixtures or wheat bran. *Small Rum. Res.* 52: 25-36.

Mercer, H.D., Neal, F.C., Himes J.A. and Edds, G.T., 1967. *Cassia occidentalis* toxicosis in cattle. *J. Am. Vet. Med. Assoc.* 151:735- 741.

Mertens, D.R., 1993. Rate and extent of digestion. pp 13–51 *in* Quantitative Aspects of Ruminant Digestion and Metabolism. Ed. Forbes, J. M. & France, J. CAB Int., Wallingford, United Kingdom.

Mgheni, D.M., 2000. Prediction of intake of tropical forages in dairy cattle based on studies from fibre digestion kinetics. PhD Dissertation, Sokoine University of Agriculture.

Milford, R. and Minson, D.J., 1965(a). Intake of tropical forage species. *Proc. Int. Grassl. Congr.* 9, 815-822.

Milford, R. and Minson, D.J., 1965(b). The relation between the crude protein content and the digestible crude protein content of tropical pasture plants. *J. British. Grassland. Soc.* 20:177-179.

Minson, D.J., 1990. Forages in Ruminant Nutrition, Academic press, San Diego, California, pp. 208 – 229 & 316 – 381.

Minson, D.J., 1994. Nutritional value of tropical legumes in grazing and feeding systems. *Forage legumes for energy efficient animal production. Proceedings of a trilateral workshop held in Palmerson North, New Zealand, April 30-May 4.* pp. 192-196.

Misra, A.K., Mishra, A.S., Tripathi, M.K., Chaturvedi, O.H., Vaithiyanathan, S., Prasad, R. and Jakhmola, R.C., 2006. Intake, digestion and microbial protein synthesis in sheep on hay supplemented with prickly pear cactus [*Opuntia ficus-indica* (L.) Mill.] with or without groundnut meal. *Small. Rum. Res.* 63: 125-134.

Moya-Rodriguez, J.G., Ramirez, R.G. and Foroughbakhch, R., 2002. Seasonal changes in cell wall digestion of eight browse species from northeastern Mexico. *Livestock research for rural development.* 14 (1). <http://www.cipav.org.co/lrrd/lrrd12/1/moya141.htm>

Mupangwa, J.F., Ngongoni, N.T., Topps, J.H., Acamovic, T., Hamudikuwanda, H. and Ndlovu, L.R., 2000. Dry matter intake, apparent digestibility and excretion of purine derivatives in sheep fed tropical legume hay. *Small. Rum. Res.* 36: 261 – 268.

Ndlovu, L.R. and Buchanan-Smith, J.G., 1985. Utilization of poor quality roughages by sheep: Effects of alfalfa supplementation on ruminal parameters, fibre digestion and rate of passage from the rumen. *Can. J. Anim. Sci.* 65: 693-703.



Nelson, C.J. and Moser, L.E., 1994. Plant factors affecting forage quality. Forage quality, evaluation and utilization. Ed. Fahey, G.C. pp 115 – 154.

Ngwa, A.T., Nsahlai, I.V. and Iji, P.A., 2002. Effect of supplementing veld hay with dry meal or silage from pods of *Acacia sieberiana* with or without wheat bran on voluntary intake, digestibility, excretion of purine derivatives, nitrogen utilization, and weight gain in South African Merino sheep. *Livest. Prod. Sci.* 77, 253-264.

NRC, 2007. The nutrient requirements of small ruminants (sheep, goats, cervids and new world camelids). *The National Academies press, Washington D.C.* pp. 246-247.

O'Hara, P.J., Pierce, K.R. and Reid, W.K., 1969. Degenerative myopathy associated with ingestion of *Cassia occidentalis*: clinical and pathologic features of the experimentally induced disease. *Am. J. Vet. Res.*; 30:2173-2180.

Ørskov, E.R. and McDonald, I., 1979. The estimation of protein degradability in the rumen from incubation measurements weighted according to rate of passage. *J. Agric. Sci. Camb.* 92:499–503.

Ørskov, E.R., Hovell, F.D.DeB. and Mould, F., 1980. The use of nylon bag technique for the evaluation of feedstuffs. *Trop. Anim. Prod.* 5: 195-213.

Ørskov, E.R., 1994. Recent advances in understanding of microbial transformation in ruminants. *Livest. Prod. Sci.* 39, 53-60.

Osuji, P.O., Nsahlai, I.V. and Khalili, H., 1993 (a). *Feed evaluation*. ILCA Manual 5. ILCA (International Livestock Centre for Africa), Addis Ababa, Ethiopia. 40 pp.

Osuji, P.O., Sibanda, S., Nsahlai, I.V., 1993 (b). Supplementation of maize stover for Ethiopian Menz sheep: effects of cotton seed cake, noug (*Guizotia abyssinica*) or sunflower cake with or without maize on intake, growth, apparent digestibility, nitrogen balance and excretion of purine derivatives. *Anim. Prod.* 57: 429-436.

Pasternak, D., Aronson, J.A., Ben-Dov, J., Forti, M., Mendlinger, S., Nerd, A. and Sitton, D., 1986. Development of new arid zone crops for the Negev Desert of Israel. *J. Arid Environ.* 11:37-59.

Paterson, J.A., 1994. The impact of forage quality and supplementation regimen on ruminant animal intake and performance. Forage quality, evaluation and utilization. Ed. Fahey, G.C. pp 59 – 114.

Pentz, E.I., 1969. Adaptation of the Rimini-Schryver reaction for the measurement of allantoin in urine to the autoanalyzer: Allantoin and taurine excretion following neutral titration. *Anal. Biochem.*, 27, 333-342.

Puchala, R. and Kulasek, G.W., 1992. Estimation of microbial protein flow from the rumen of sheep using microbial nucleic acid and urinary excretion of purine derivatives. *Can. J. Anim. Sci.* 72: 821-830.

Ramirez, R.G., 1998. Nutrient digestion and nitrogen utilization by goats fed native shrubs *Celtis pallid*, *Leucophullum texanum* and *Porlieria angustifolia*. *Small Rum. Res.* 28: 47-51.

Rethman, N.F.G. & Van Niekerk, W.A., 1991. The potential of oldman saltbush as forage for the winter/spring period in sour grassveld areas. *Proc. IVth Int. Rangeland. Congr.* Montpellier, France.

Robertson, J.B. and Van Soest, P.J., 1981. The analysis of dietary fibre in food. James, W.P.T and Theander, O. Dekker, eds, New York.

Robinson, P.H., Tamminga, S. and Van Vuuren, A.M., 1987. Influence of declining level of feed intake and varying the proportion of starch in the concentrate on rumen ingesta quantity, composition and kinetics of ingesta turnover in dairy cows. *Livest. Prod. Sci.* 17: 37-62.

Salawu, M.B., 1997. The nutritive value of the leguminous browse *Caliandra calothyrsus* and the role of condensed tannins in ruminant feeds. PH.D Thesis, University of Aberdeen.

Samuels, M.L., 1989. Statistics for the life sciences. Collier MacMillan Publishers, London.

Scholtz, G.D.J., van der Merwe, H.J. and Tylutki, T.P., 2009. The nutritive value of South African *Medicago Sativa* L. hay. *S. Afr. J. Anim. Sci.* 39 (Supplement 1), 179 – 182.

Snyder, L.J.U., Luginbuhl, J-M., Mueller, J.P., Conrad, A.P. and Turner, K.E., 2007. Intake, digestibility and nitrogen utilization of *Robinia pseudoacacia* foliage fed to growing goat wethers. *Small Rum. Res.* 71: 179-193.

Snyman, H.A., 2003. Revegetation of bare patches in a semi-arid rangeland of South Africa: an evaluation of various techniques. *J. Arid Environm.* 55: pp. 417-432.

Snyman, L.D., 2006. Qualitative characteristics of selected *Atriplex nummularia* (Hatfield select). M.Sc. (Agric) dissertation. Department of Animal and Wildlife sciences. University of Pretoria, Pretoria, South Africa.

Sparks, C.F., 2003. Interspecies variation in nutritive value of certain drought tolerant fodder shrubs. M.Sc. dissertation. Department of Animal and Wildlife Sciences. University of Pretoria, Pretoria, South Africa.

Suliman, H.B. and Shommein, A.M., 1986. Toxic effect of the roasted and unroasted beans of *Cassia occidentalis* in goats. *Vet. Hum. Toxicol.*; 28:6-11.

Statistical Analysis Systems, 2005. SAS User's Guide: Statistics Version 8. SAS Institute Inc. Cary, NC., USA.

Tamminga, S. and Van Vuuren, A.M., 1988. Formation and utilization of end products of lignocellulose degradation in ruminants. *Anim. Feed. Sci. Technol.* 21: 141-159.

Tamminga, S., Robinson, P.H., Vogt, M. and Boer, H., 1989. Rumen ingesta kinetics of cell wall components in dairy cows. *Anim. Feed Sci. Technol.* 25: 89-98.

Tebot, I., Britos, A., Godeau, J.M. and Cirio, A., 2002. Microbial protein production determined by urinary allantoin and renal urea sparing in normal and low protein fed Corriedale sheep. *Vet. Res.* 33: 101-106.

Technicon Auto Analyzer II. 1976. Industrial method 334-74. Technicon Industrial Systems, Tarrytown, NY.

Thomson, S., 2002. Canavine toxicity: Is Sutherlandia a health herb or potent (ial) poison? *Gaia research institute. January 2002.* ([director@gaiaresearch.co.za](mailto:director@gaiaresearch.co.za))

Tilley, J.M.A. and Terry, R.A., 1963. A two stage technique for *in vitro* digestion of forage crops. *J. Br. Grassland Soc.* 18: 104-111.

Tolera, A. and Sundstol, F., 2000. Supplementation of graded levels of *Desmodium intortum* hay to sheep feeding on maize stover harvested at three stages of maturity. 2. Rumen fermentation and nitrogen metabolism. *Anim. Feed Sci. Technol.* 85, 239-257.

Underwood, E.J., and Suttle, N.F., 1999. The mineral nutrition of livestock. 3<sup>rd</sup> ed. CABI publishing. CAB International, Wallingford, Oxon, UK, pp 614.

Van Breda, P.A.B. and Barnard, S.A., 1991. Veld plants of the winter rainfall region. Department of Agriculture Bulletin, 422: 1 –211.

Van Houtert, M.F.J., 1993. The production and metabolism of volatile fatty acids by ruminants fed roughages: A Review. *Anim. Feed Sci. Technol.* 43: 189-225.

Van Niekerk, W.A., 1997. Inname en partiële verteerbaarheid van 'n aantal weidingsgewasse deur skape en die gebruik van enkele kwaliteitsparameters om inname te voorspel. PhD thesis. University of Pretoria. South Africa.

Van Niekerk, W.A., Sparks, C.F., Rethman, N.F.G. and Coertze, R.J., 2004 (a). Qualitative characteristics of some *Atriplex* species and *Cassia sturtii* at two sites in South Africa. *S. Afr. J. Anim. Sci.* 34 (Supplement 1), 108 – 110.

Van Niekerk, W.A., Sparks, C.F., Rethman, N.F.G. and Coertze, R.J., 2004 (b). Mineral composition of certain *Atriplex* species and *Cassia sturtii*. *S. Afr. J. Anim. Sci.* 34 (Supplement 1), 105 – 107.

Van Soest, P.J., 1965. Symposium on factors influencing the voluntary intake of herbage by ruminants: voluntary intake in relation to chemical composition and digestibility. *J. Anim. Sci.* 24, 834-843.

Van Soest, P.J., 1982. Nutritional ecology of the ruminant. O and B books, Corvallis, Oregon, USA.

Van Soest, P.J., 1984. Chemical procedures for estimating nutritive value. Section 3. In: Laboratory evaluation of fibrous feeds. Ed. Knipfel, J.E. Swift current research station, Sadkatchewan.

Van Soest, P.J., and Wine, R.H., 1967. Use of detergents in the analysis of fibrous feeds. IV. Determination of plant cell wall components. *J. Assoc. Off. Anal. Chem.* 50: pp 50.

Ventura, M.R, Castanon, J.I.R., Pieltain, M.C. and Flores, M.P., 2004. Nutritive value of forage shrubs: *Bituminaria bituminosa*, *Rumex lunaria*, *Acacia salincina*, *Cassia sturtii* and *Adenocarpus foliosus*. *Small Rum. Res.* 52, 1-2, pp. 13-18.

Walker, B.H., 1980. A review of browse and its role in livestock production in Southern Africa. Browse in Africa, the current state of knowledge. Ed. Le Houerou, H.N. ICLA Addis Ababa, Ethiopia.

Watson, M.C., O'Leary, W.O. and Glenn, E.P., 1987. Evaluation of *Atriplex lentiformis* (Torr.) S. Wats. and *Atriplex nummularia* Lindl. as irrigated forage crops. *J. Arid Environ.* 13: 293 – 303.

Webb E.C., 1994. Synthesis of long chain fatty acids in ruminants and their effects on meat quality. Chapter 3, page 58-59 (with modifications). PhD thesis. University of Pretoria, South Africa.

Wilcock, T.E., Van Niekerk, W.A., Rethman, N.F.G. and Coertze, R.J., 2004. A comparison of *Cassia sturtii*, *Tripteris sinuatum* and *Sutherlandia microphylla*: three fodder shrubs applicable to revegetation of degraded rangeland in the Northern Cape Province. *S. Afr. J. Anim. Sci.* 34 (Supplement 1), 114 – 116.