

REFERENCES

- Agricultural and Food Research Council., 1992. Nutritive requirements of ruminant animals: Protein. AFRC Technical Committee on Response to Nutrients. Report No. 9. Nutr. Abstr. Rev. (Series B), 62:787-835.
- Agricultural Research Council., 1980. The nutrient requirements of ruminant livestock. Commonwealth Agriculture Bureaux, Famham Royal. Slough. UK. P 126-127.
- Allaway, W. H., 1973. Selenium in the food chain. Cornell Vet., 63: 151-170.
- Ammerman, C.B., Waldroup P.W., Arrington, L.R, Shirley, R.L. and Harms, R.H., 1966. Nutrient digestibility by ruminants of poultry litter containing dried citrus pulp. J. Agric. Food Chem., 14: 279 – 284.
- Angus, K.W., Suttle, N.F., Munro, C.S. and Field, A.C., 1978. Adverse effects on health of including high levels of dried poultry waste in the diets of lambs. J. Comp. Pathol., 88: 449-466.
- Anonymous., 1960. Dorp het plaas geword. Landbouweekblad, Des 6; 66.
- Anonymous., 1967. Standard methods for the examination of dairy products- Microbial and chemical (12th ed.) Amer. Public Health Assoc. Enc. New York.
- Anonymous., 1974. Clinical laboratory. Medico-chemical investigation methods (11th edn).E Merck, Darmstadt.
- Association of Official Analytical Chemists., 1975. Official methods of analysis. Association of Official Analytical Chemists, Washington, DC, 12th edn.
- Association of Official Analytical Chemists., 1990. Official Methods of Analysis. A. O. A. C. Washington, DC,15th Edn.

- Banton, M.I., Nicholson, S. S., Jowett, P.L.H., Brantley, M.B. and Boudreaux, C.L., 1987. Copper toxicosis in cattle fed chicken litter. *J. Am. Vet. Med. Ass.*, 191: 827-828.
- Bare, L.N., Wiseman, R.F. and Abbott, O.J., 1964. Effects of dietary antibiotics and uric acid on the growth of chicks. *J. Nutr.* 83:27- 34.
- Bastianello, S.S., Fourie, N., Prozesky, L., Nel, P.W. and Kellermann, T.S., 1995. Cardiomyopathy of ruminants induced by the litter of poultry fed on rations containing the ionophore antibiotic, maduramicin. II. Macropathology and histopathology. *Onderstepoort J. Vet. Res.*, 62: 5-18.
- Bastianello S .S., McGregor, H. L., Penrith, M.L., Fourie, N., 1996. Achronic cardiomyopathy in feedlot cattle attributed to toxic levels of salinomycin in the feed. *J. SA Veter. Assoc.*, 67: 38-41.
- Bath, I.H. and Rook, J.A.F., 1965. The evaluation of cattle foods and diets in terms of the ruminal concentration of volatile fatty acids. II Roughages and succulents. *J. Agric. Sci.*, 64:67-75.
- Belasco, I.J., 1954. New nitrogen feed compounds for ruminants a laboratory evaluation. *J. Anim. Sci.*, 13: 601-610.
- Ben-Ghedalia, D., Miron, J., Est, Y. and Yosef, E., 1988. SO₂ treatment for converting straw into a concentrate-like feed: A growth study with lambs. *Anim. Fd Sci. Technol.*, 19: 219-229.
- Ben-Ghedalia, D., Miron, J. and Yosefu, E., 1996. Apparent digestibility of minerals by lactating cows from a total mixed ration supplemented with poultry litter. *J. Dairy Sci.*, 79: 454-458.

- Benne, E.J., 1970. A compilation of all samples of poultry waste analysed by E. J. Benne. Mich. Agr. Exp. Sta. Res. Rep., 117: pp 49.
- Bhattacharya, A.N. and Fontenot, J.P., 1965. Utilization of different levels of poultry litter nitrogen by sheep. J. Anim. Sci., 24: 1174-1178.
- Bhattacharya, A.N. and Fontenot, J.P., 1966. Protein and energy value of peanut hull and wood shaving poultry litters. J. Anim. Sci., 25: 367-371.
- Bhattacharya, A.N. and Taylor, J.C., 1975. Recycling animal waste as feedstuff: A review. J. Anim. Sci., 41: 1438-1457.
- Bishop, E.J.B., Wilke, P.I., Nash, W.J., Nell, J.A.G., MacDonald, D.A., Compaan, J.P., Grobler, J. and Kingman, E.R., 1971. Poultry manure as a livestock feed (Part 2). Farming in S. Afr., 46: 49-53.
- Borchers, R., 1977. Allantoin determination. Anal. Biochem., 79:612-613.
- Borland International., 1993. User's guide. Borland Quattro Pro for Windows. Version 5.0 Scotts Valley CA 95067-0001.
- Borgioli, E. and Tocchini, M., 1969. Sterilised poultry litter for feeding young cattle. Aliment. Anim., 13:263. Cited in Nutri. Abstr. Rev., 40:3871 (1970).
- Bosman, S.W., 1973. Chicken litter in fattening rations for cattle and sheep. S. Afr. J. Anim. Sci., 3:57- 61.
- Bredon, R.M., Stewart, P.G. and Dugmore, T.J., 1987. A manual on the nutritive and chemical composition of commonly used South African farm feeds. Natal Region Department of Agric. And Water Supply, Pretoria. RSA.
- Brosh, A., Aharoni, Y., Levy, D. and Holzer, Z., 1998. Effects of source and content of ash in poultry litter used in diets for beef cattle. J. Agric. Sci. Camb., 131: 87- 95.

- Brugman, H.H., Dickey, H.C., Plummer, B.E. and Poulton, B.R., 1964. Nutritive value of poultry litter. *J. Anim. Sci.*, 23: 869. (Abstract).
- Brugman, H.H., Dickey, H.C., Plummer, B.E., Goaster, J., Heitam, R.N. and Taka, M.R.Y., 1968. Drug residue in lamb carcasses fed poultry litter. *J. Anim. Sci.*, 27:1132 (abstr.).
- Bull, L.S. and Reid, J.T., 1971. Nutritive value of chicken manure for cattle. *Livestock Waste Management and Pollution Abatement. Proc. Int. Symp. Livestock Wastes*, Columbus, Ohio, pp. 297- 300.
- Calvert, C.C., 1971. Faecal residues from feed additives- Poultry. In : Animal Waste Reuse-- Nutritive Value and Potential Problems, from Feed Additives. A Review. Prepared by Animal Science Research Division and Agricultural Research Service of the USA Department of Agriculture. pp.14-45.
- Calvert, C.C. and Smith, L.W., 1972. Arsenic in milk and blood of cows fed organic arsenic compounds. *J. Dairy.*, 55:706 (abstr.).
- Calvert, C.C., 1974. Feed additive residues in animal manure processed for feed. *Feedstuffs*, 45 (No. 17):32.
- Carter, T.A. and Poore, M., 1998. Deep stacking broiler litter as a feed for beef cattle. North Carolina Cooperative Extension Service, North Carolina State University, Raleigh, North Carolina (Electronic pub. No. DRO- 49).
- Casey, N.H., Van Niekerk, W.A. and Spreeth, E.B., 1988. Fatty acid composition of subcutaneous and kidney fat of sheep grazed on eight different pastures. *Meat Sci.*, 23: 55-63.
- Caswell, L.F., Webb, K.E. and Fontenot, J.P., 1977. Fermentation, nitrogen utilization, digestibility and palatability of broiler litter ensiled with high moisture corn grain. *J. Anim. Sci.*, 44:803-813.

- Caswell, L.F., Fontenot, J. P. and Webb, K.E., 1978. Fermentation and utilization of broiler litter ensiled at different moisture levels. *J. Anim. Sci.*, 46: 547-561.
- Chaudry, S.M., Fontenot, J.P., Naseer, Z. and Ali, C. S., 1996. Nutritive value of deep stacked and ensiled broiler litter for sheep. *Anim. Feed Sci. Technol.*, 57:165-173.
- Chen, X. B., Meija, A. T., Kyle, D. J., Orskov, E. R., 1995. Evaluation of the use of purine derivative:creatinine ratio in spot urine and plasma samples as an index of microbial protein supply in ruminants: studies in sheep. *J. Agric. Sci., Cambridge* 125:137-143.
- Chester-Jones, H., Fontenot, J.P., Dana, G.R. and Hovatter, M.D., 1972. Performance of growing and finishing cattle fed deep-stacked and ensiled broiler litter. *J. Anim. Sci.*, 55 : 59 (abstract).
- Crickenberger, R.G. and Goode, L., 1998. Guidelines for feeding broiler litter to cattle. Water Quality and Waste Management. North Carolina Cooperative Extension Servive. Publication Number AG-61.
- Cullison, A.E., McCampbell, H.C., Cunningham, A.C., Lowery, R.S., Warren, E.P., McLendon, B.D. and Sherwood, D.H., 1976. Use of poultry manures in steer finishing rations. *J. Anim. Sci.*, 42: 219 - 228.
- Deshck, A., Abo-Shehada, M., Allonby, E., Givens, D.I. and Hill, R., 1998. Assessment of the nutritive value for ruminants of poultry litter. *Anim. Feed Sci. Technol.*, 73: 29-35.
- De Villiers, S., Van der Walt, J. G. and Procos, J., 1977. An accurate, sensitive and producible method for the estimation of free fatty acids. *Onderstepoort J. of Vet. Res.*, 44:169-172.

- Doctorian, D.S. and Evers, G.W., 1998. Utilizing broiler litter as a protein and mineral supplement for beef cows. Texas Agric. Extension Service/ The Texas A & M University System /College Station, Texas. 77843-2468.
- Downes, A.M. and McDonald, I.W., 1964. The chromium⁵¹ complex of ethylene diaminetetraacetic acid as a soluble rumen marker. Brit. J. Nutr., 18:153-162.
- Drake, C.L., McClure, W.H. and Fontenot, J.P., 1965. Effects of levels and kinds of broiler litter for fattening steers. J. Anim. Sci., 24: 879 (abstr.).
- Duncan, W.R.H., Ørskov, E. R., Fraser, C. and Garton, G.A., 1974. Effect of processing of dietary barley and supplementary cobalt and cyanocobalamin on the fatty acid composition of lamb triglycerides, with special reference to branched-chain components. Brit. J. Nutr., 32:71-75.
- Ehlig, C.F. Hogue, D. E., Allaway, W.H. and Hamm, D.J., 1967. Fate of selenium from selenite or selenomethionine with or without vitamin E, in lambs. J. Nutr., 92: 121-126.
- Eiteman, M.A., Gordillo, R.M., Cabrera, M.I. and Fresnius, J., 1994. Anal. Chem. 346:680 - 683.
- El-hag, M.G. and El-hag, G.A., 1981. Effect of supplementing groundnut hulls with dry excreta or cotton seed cake on the performance of Sudan desert sheep. J. Anim. Sci., 51 (suppl.1):133.
- Elmund, G.K., Morrison, S.M., Grant, D.W. and Nevins, M.P., 1980. Role of excreted chlortetracycline on modifying the decomposition process of feedlot waste. In: Digestive Physiology and Metabolism in Ruminants. Ed: Ruckebusch, Y. & Thivend, P. (1971) pp 641-662.

- El-Sabban, F.F., Bratzler, J.W., Long, T.A., Frear, D.E.H. and Gentry, R.F., 1970.
Value of processed poultry waste as a feed for ruminants. *J. Anim. Sci.*, 31: 107-111.
- Erb, R.E., Brown, C.J., Callahan, C.J., Moeller, N.J., Hill, D. L. and Cunningham, M.D., 1976. Dietary urea for dairy cattle. II. Effect on functional traits. *J. Dairy Sci.*, 59: 659-668.
- Essig, H.W., 1975. Recycling nutrients for livestock. *Feedstuffs.*, 47:35.
- Evans, E., Moran, E.T., McLeod, G.K. and Turner, E.M., 1978. Laying hen excreta as a ruminant feed stuff. II. Preservation and acceptability of wet excreta by sheep. *J. Anim. Sci.*, 46:527-534.
- Faichney, G.J., 1975. The use of markers to partition digestion within the gastrointestinal tract of ruminants. New England Press, Armidale, Australia, pp. 277-291.
- Ferrel, C.L. and Jenkins, T.G., 1985. Cow type and nutritional environment: Nutritional aspects. *J. Anim. Sci.*, 61:725- 732.
- Fisher, C., Wise, D. and Filmer, D.G., 1972. The effect of copper on the growth of broilers and the interaction of copper with zinc and iron. 4th World Poultry Congr., Madrid 2: 759-764.
- Flegal, C.J. and Zindel, H.C., 1970. The utilization of poultry waste as a feedstuff for growing chicks. Research Report No. 117, Michigan State Agricultural Experiment Station, East Lansing, Michigan, p. 21.
- Fluharty, F.L. and McClure, K.E., 1997. Effects of dietary intake and protein concentration on performance and visceral organ mass in lambs. *J. Anim. Sci.*, 75:604-610.

- Fontenot, J.P., McClure, W.H., Kelly, R.F. and Litton, G.W., 1963. The value of poultry litter as feedstuffs for fattening beef steers. *Va. Agric. Exp.*
- Fontenot, J. P., Bhattacharya, A.N., Drake, C. L. and McClure, W.H., 1966. Value of broiler litter as feed for ruminants. *Proc. Natl. Symp. Anim. Waste Management*, pp. 105-108.
- Fontenot, J.P., Webb, K.E., Harmon, B.W., Tucker, R.E. and Moore, W.E.C., 1971. Studies of processing nutritional value and palatability of broiler litter for ruminants. *Livestock Waste Management and Pollution Abatement. Proc. Int. Symp. Livestock Waste*, Columbus, Ohio, p. 391.
- Fontenot, J.P. and Webb, K.E., 1974. Poultry waste as feed stuffs for ruminants. *Fed. Proc.*, 33: 1936-1937.
- Fontenot, J.P. and Webb, K.E., 1975. Health aspects of recycling animal waste by feeding. *J. Anim. Sci.*, 40: 1267.
- Fontenot, J. P. and Jurubescu, V., 1980. Processing animal waste by feeding to ruminants. In: *Digestive Physiology and Metabolism in Ruminants*. Y. Ruckebusch and P. Thivend, Eds. Avi Publishing Co., Westport, Connecticut.
- Fontenot, J.P., 1991. Recycling animal waste by feeding to enhance environmental quality. *Prof. Anim. Scientist.*, 7: 1-8.
- Fourie, N., Bastianello, S.S., Prozesky, L., Nel, P.W. and Kellerman, T.S., 1991. Cardiomyopathy of ruminants induced by the litter of poultry fed on rations containing the ionophore antibiotic, maduramicin. Epidemiology, clinical signs and clinical pathology. *Onderstepoort J. Vet. Res.*, 58: 291-296.

- Galmez, J., Santisteban, R., Haardt, C., Crempien, C., Villalta, I. and Terrell, D., 1970. Performance of ewes and lambs fed broiler litter. *J. Anim. Sci.*, 31: 241 (abstr.).
- Ganther, H.E., Vadhanavikit, S. and Ip, C., 1990. Selenium methylation, demethylation and biological activity in the rat. *FASEB J.*, No. 3, Part 1, A372.
- Goering, H. K. and Smith, L. W., 1977. Composition of corn plant ensiled with excreta or nitrogen supplements and its effects on growing wethers. *J. Anim. Sci.*, 44: 452 –461.
- Goetsch, A.L., Anthony, N.B., Woodley, M.A. and Tabler, G.T., 1998. Chemical constituents in broiler litter in two areas of a production unit after different numbers of growing periods. *Bioresource Tech.*, 65: 151-157.
- Golab, T., Barton, S. J. and Scroggs, R. E., 1973. Colometric method for monensin. *J. Assoc. Off. Anal. Chem.*, 56: 171-173.
- Griel, L.C., Kradel, D.C. and Wickersham, E.W., 1969. Abortion in cattle associated with the feeding of poultry litter. *Cornell Vet. J.*, 59: 226-235.
- Grimaud, P. and Doreau, M., 1995. Effect of extended underfeeding on digestion and nitrogen balance in nonlactating cows. *J. Anim. Sci.*, 73: 211-219.
- Hadjipanayiotou, M., 1984. The use of poultry litter as feed in Cyprus. *World Animal Review*, 49: 32-38.
- Hakkarainen, J., 1993. Bioavailability of selenium. *Norwegian J. Agric. Sci. Suppl.*, 11: 21-35.
- Harmon, B.W., Fontenot, J.P. and Webb, K. E., 1974. Effect of processing method of broiler litter on nitrogen utilisation by lambs. *J. Anim. Sci.*, 7:117-120.

- Harmon, B.W., Fontenot, J.P. and Webb, K. E., 1975. Ensiled broiler litter and corn forage. I. Fermentation characteristics. *J. Anim. Sci.*, 40: 144-155.
- Harris, L.E. 1970. Nutrition research techniques for domestic and wild animals. Vol.1. Compiled by E.R. Davie. Animal Nutrition Subdirectorate, Animal Dairy Science Research Institute. Irene. SA.
- Harrison, J.H. and Conrad, H.R., 1984. Effect of dietary calcium on selenium absorption by the nonlactating dairy cow. *J. Dairy Sci.*, 67: 1860-1864.
- Harrison, T.J., Van Ryssen, J.B.J. and Barrowman, P.R., 1987. The influence of breed and dietary molybdenum on the concentration of copper in tissues of sheep. *S. Afr. J. Anim. Sci.*, 17: 104-110.
- Hartmann, F. and Van Ryssen, J.B.J., 1997. Metabolism of selenium and copper in sheep with and without sodium bicarbonate supplementation. *J. Agric. Sci. Camb.*, 128: 357-364.
- Henry, P.R., Echevarria, M.G., Ammerman, C.B.C. and Rao, P.V., 1988. Estimation of the relative biological availability of inorganic selenium sources for ruminants using tissue uptake of selenium. *J. Anim. Sci.*, 66: 2306-2312.
- Hertelandy, F., Taylor, T.G., Mathur, R.S. and Common, R.H., 1965. Isolation of estradio-17 β from hens' urine and its characterization as the crystalline 3-methyl ether. *Can. J. Biochem.*, 43:1379.
- Hill, C.H., 1974. Reversal of selenium toxicity in chicks by mercury, copper, and cadmium. *J. Nutr.*, 104: 593-598.
- Hodgetts, B., 1971. The effects of including dried poultry wastes in the feed of laying hens. Livestock Waste Management and Pollution Abatement. Proc. Int. Symp. Livestock Waste, Columbus, Ohio, p. 311.

- Hovatter, M.D., Sheehan, W., Dana, G.R., Fontenot, J.P., Webb, K.E. and Lamm, W.D., 1979. Different levels of ensiled and deep stacked broiler litter for growing cattle. Va. Polytech. Inst. And State Univ. Res. Div. Rep., 175:77-83.
- Howell, J. M. and Gooneratne, S. R., 1987. The pathology of copper toxicity in animals. In: Howell, J. McC, Path. F. R. C., Gawthorne, J. M. (eds) . Copper in animals and man. Vol 2 . CRC Press, Boca Raton, Florida: pp 53-78
- Hunter, R.A. and Siebert, B.D., 1986. The effect of genotype, age, pregnancy, lactation and rumen characteristics on voluntary intake of roughage diets by cattle. Aust. J. Agric. Res., 37: 549-560.
- Ilian, M.A., Razzaque, M.A. and Salman, A.J., 1988. Unconventional feeds for sheep: Effects on performance and meat quality and composition. Biological Wastes 24, 115-125.
- Jacobs, G.J. L. and Leibholz, J., 1978. The digestion of dry matter, organic matter and nitrogen in calves fed diets containing broiler litter. J. Agric. Sci., Camb., 90: 367-372.
- Jacob, J.P., Kunkle, W.E., Tervola, R.S., Miles, R.D. and Mather, F.B., 1998. Broiler litter, Part 1: A feed ingredient for ruminants.
<http://hammock.Ifas.ufl.edu/fairs/ps001>
- Jakhmola, R.C., Kundu, S.S., Punj, M.L., Singh, K. Kamra, D.N. and Singh, R., 1988. Animal excreta as ruminant feed: Scope and limitations under Indian conditions. Anim. Feed Sci. Technol., 19: 1-23.

Jordan, E.R. and Swanson, L.V., 1979. Effect of crude protein on reproductive efficiency, serum total protein and albumin in the high producing dairy cow. *J. Dairy Sci.*, 48: 1154 (Abstract).

Kaneko, J. J., 1989. Clinical Biochemistry of Domestic Animals (4th edn). Academic Press, New York.

Kargaard, J. and Van Niekerk, B. D. H., 1977. Incorporation of DPW, urea and fish meal with varying molasses levels in cattle feedlot rations. *S. Afr. J. Anim. Sci.*, 7: 117-125.

Kargaard, J. and Van Niekerk, B.D.H., 1978. The use of DPW at varying levels in cattle finishing rations. *S. Afr. J. Anim. Sci.*, 8:143-147.

Kitching, J.P., 1986. The uses and dangers of poultry litter in feeding in cattle. *S. Afr. Vet. Assoc. Biannual Congr.*, August, 1986.

Knight, E.F., McCaskey, T.A., Anthony, W.B. and Walters, J.C., 1977. Microbial population changes and fermentation characteristics of ensiled bovine manure-blended rations. *J. Dairy Sci.*, 60:416-423.

Koenig, K.M., Rode, L.M., Cohen,R.D.H. and Buckley, W.T., 1997. Effect of diet and chemical form of selenium on selenium metabolism in sheep. *J. Anim. Sci.*, 75:817-827.

Koh, T.S. and Benson, T. H., 1983. Critical re-appraisal of fluorometric method for determination selenium in biological materials. *J. Assoc. Off. Anal. Chem.*, 66:918-926.

Kumanov, S., Jankor, B. and Paliev, H., 1969. Use of deep litter from broiler production as a feed:- peleted or as meal. *Nutr. Abstr. Rev.*, 40: 3872.

- Kwak, W., Fontenot, J.P. and Herbein, J.H., 1998. Effect of processing method on ruminal solubility and degradability of broiler litter. *Bioresource Tech.*, 66: 13-18.
- Langlands, J.P., Bowles, J.E., Donald, G.E. and Smith, A.J., 1986. Selenium excretion in sheep. *Aust. J. Agric. Res.*, 37:201-209.
- Lloyd, L.E., McDonald, B.E. and Crampton, E.W., 1978. Chapter18. Fundamentals of Nutrition. (2nd Ed.) , W.H. Freeman and Co. San Francisco. Page 264.
- Lober, U., Eisengarten, H.J. and Flachowsky, G., 1992. A field study on the influence of urea on microbial decontamination and digestibility of broiler litter. *Bioresource Technol.*, 41:135-138.
- Lovett, J., Messer, W. and Read, R., 1971. The microflora of southern Ohio poultry litter. *Poult. Sci.*, 50:746-751.
- Lowry, M., 1992. In: Byford-Jones, C., 1992. Is broiler litter in feed really dangerous? *Farmer's weekly*, (January, 1992) pp 10-11.
- Long, T.A., Bratzler, J.W. and Frear, D.E.H., 1969. The value of hydrolyzed and dried poultry waste as a feed for ruminant animals. In: *Animal Waste Management*. Cornell University, Ithaca, NY, pp 98-104.
- Malik, S. and Bhattacharya, A.N., 1971. Nitrogen and energy utilization from different kinds of poultry litter in sheep. Proc. 10th Int. Congr. Anim. Prod. p. 250.
- Manyuchi, B., Smith, T. and Mikayiri, S., 1992. The use of poultry litter in ruminant diets: 1. Poultry litter and cotton seed meal as supplements for weaner steers grazing natural pasture during the dry season or sheep fed natural pasture hay in pens. *Zimbabwe J. Agric. Res.*, 30: 91-103.

- Manyuchi, B., Smith, T. and Mikayiri, S., 1992. The use of poultry litter in ruminant diets : 2. Poultry litter and cotton seed meal as sources of nitrogen in feedlot diets for cattle and sheep. Zimbabwe J. Agric. Res., 30: 105-115.
- Marais, J.P. and Everwell, T.K., 1983. The use of trichloroacetic acid as precipitant for the determination of “true protein” in animal feeds. S. Afr. J. Anim. Sci., 13: 138 -139.
- McCaskey, T.A. and Anthony, W.B., 1979. Human and animal aspects of feeding livestock excreta. J. Anim. Sci., 48: 163-177.
- McDonald, P., Edwards, R.A. and Greenhalgh, F.D., 1988. Animal Nutrition. (4th edn), Longman group UK. Harlow, UK, pp 219 – 230.
- McInnes, P., Austin, P.J. and Jenkins, D.L., 1968. The value of poultry litter and wheat mixture in the drought feeding of weaner sheep. Aust. J. Exp. Agric. Anim. Husb., 8:401-404.
- McClure, W.H. and Fontenot, J.P., 1987. Poultry litter in corn silage can be used to finish steers. Feedstuffs., 59:12-13.
- McConnell, K.P. and Carpenter, D.M., 1971. Interrelationship between selenium and specific trace elements. Proc. Soc. Exp. Biol. Med., 137: 996-1001.
- Melton, S.L., 1990. Effects of feeds on flavour of red meat: a review. J. Anim. Sci., 68:4421-4435.
- Messer, J.W., Lovett, Y., Murthy, G.K., Wehby, A.J., Schafer, M.C. and Read, R.B., 1971. An assessment of some public health problems resulting from feeding poultry litter to animals. Poult. Sci., 50: 874-881.

- Mills, E.W., Comerford, J.W., Hollender, R., Harpster, H.W., House, B. and Henning, W.R., 1992. Meat composition and palatability of Holstein and beef steers as influenced by forage type and protein source. *J. Anim. Sci.*, 70:2446-2451.
- Ministry of Agriculture, Fisheries and food., 1986. The Analysis of Agricultural Materials., 3rd ed., Her Majesty's Stationery Office, London, pp. 219-221.
- Moksnes, K., Norheim, G., 1983. Selenium and glutathione peroxidase levels in lambs receiving feed supplemented with sodium selenite or selenomethionine. *Acta Vet. Scand.*, 24:45-58.
- Morgan, P.J.K., Pienaar, J.P., Clark, R.A., 1976. Animal based methods of determining herbage intake and quality under grazing conditions. In: Proc. 5th Congr. Grassl. Soc., vol. 11 pp. 73-78.
- Muirhead, S., 1996. Adding hay to broiler litter diet helps protect cows from milk fever. *Feedstuffs*, 68(No. 2):10.
- National Research Council., 1985. Nutrient Requirements of Sheep (6th edn). National Academy Press, Washington DC.
- Nel, P.W., 1989. Toksikologiese effekte by hoendermis aan vee. SAVDP-Kongres.
- Ngongoni, N.T. and Manyuchi, B., 1993. A note on the flow of nitrogen to the abomasum in ewes given a basal diet of milled star-grass hay supplemented with graded levels of deep litter poultry manure. Zimbabwe *J. Agric. Res.*, 31: 135-140.
- Noland, P.R., Ford, B.F. and Ray, M.I., 1955. The use of ground chicken litter as a source of nitrogen for gestating and lactating ewes and fattening steers. *J. Anim. Sci.*, 14: 860-865.

- Ogonowski, K., Barnard, Marie-Luise and Giesecke, W.H., 1984. Bacteriological findings regarding the hygienic safety of poultry litter intended as an ingredient of feeds for ruminants. *Onderstepoort J. Vet. Res.*, 51: 249-252.
- Offer, J. and Offer, N.W., 1994. Distillery by-products for cattle. *Feed Mix* 2, 14-17.
- Ojowi, M., McKinnon, J.J., Mustafa, A. and Christensen, D.A., 1997. Evaluation of wheat- based wet distillers grains for feedlot cattle. *Can. J. Anim. Sci.*, 17:447-454.
- Oldfield, J.E., Burau, R., Moller, G., Ohlendorf, H.M., Ullrey, D., 1994. Risks and benefits of selenium in agriculture. Issue Paper Supplement 3, Council for Agricultural Science and Technology, Ames, IA.
- Olson, K.J., Fontenot, J.P. and Failla, M.L., 1984. Influence of molybdenum and sulfate supplementation and withdrawal of diets containing higher cooper broiler litter on tissue cooper levels in ewes. *J. Anim. Sci.*, 59: 210.
(Abstract)
- Oltjen, R. R., Silyter, L.L., Kozak, A.S. and Williams, E.E., 1968. Evaluation of urea, biuret, urea-phosphate and uric acid as NPN sources for cattle. *J. Nutr.*, 92: 193-202.
- Oosthuizen, S.A., 1979. Which factors have to be considered when feeding poultry manure? *Vleis/Meat*: 12-13.
- Ørskov, E.R., Fraser, C. and Gordon, J.G., 1974. Effects of processing of cereals on rumen fermentation, digestibility, rumination time and firmness of subcutaneous fat in lambs. *Brit. J. Nutr.*, 32:59-69.

- Ørskov, E.R. and Miller, E.I., 1988. Chapter 5. Protein evaluation in ruminants. Feed Science, (Editor- E.R. Ørskov) Elsevier Service, Amsterdam. pp 124-135.
- Park, K.K., Goetsch, A.L., Patil, A.R., Kouakou, B. and Johnson, Z.B., 1995. Composition and in vitro digestibility of fibrous substrates placed in deep-stacked broiler litter. *Anim. Feed Sci. Technol.*, 54:159-174.
- Patil, A.R., Goetsch, A.L., Kouakou, B., Galloway, D.L., and Johnson, Z. B., 1995b. Nutritive value of deep-stacked and composted broiler litter for growing cattle. *Prof. Anim. Sci.*, 11:100-105.
- Patil, A.R., Goetsch, A.L., Kouakou, B., Galloway Sr, D.L., Forester, Jr, L.A. and Park, K.K., 1995a. Effects of corn vs corn plus wheat in forage- based diets containing broiler litter on feed intake, ruminal digesta characteristics and digestion in cattle. *Anim. Feed Sci. Technol.*, 55:87-103.
- Perez, J.F., Balcells, J., Guada, J.A. and Castrillo, C., 1996. Determination of rumen microbial-nitrogen production in sheep: a comparison of urinary purine excretion with methods using N and purine bases as makers of microbial-nitrogen entering the duodenum. *Brit. J. Nutr.*, 75:699-709.
- Phelps, A., 1990. Coccidiostats a health risk when broiler litter is fed. *Feedstuffs*, 62 (39):13.
- Pienaar, J. P., Roux, C. Z. and Cronje, P.B., 1989. Comparison of invivo and in sacco methods to estimate mean retention time of fermentable organic matter in the rumen. *S. Afr. J. Anim. Sci.*, 19:71-75.
- Pitzen, D., 1994. The trouble with iron. Resolved by pro-oxidant: anti-oxidant balancing of dairy rations. *Feed Int.* (August, 1994). Page 3.

- Polin, D., Varghese, S., Neff, M., Gomez, M., Flegal, C.J. and Zindel, H., 1971. The metabolizable energy value of dried poultry waste. Res. Rep. No.152. Michigan Agr. Exp. Sta., East Lansing. p. 32.
- Polman, Y., Newmark, H., Kaim, M. and Kaufmann, W., 1981. Performance, rumen, and blood metabolites in high yielding cows fed varying protein percents and protected soybean. J. Dairy Sci., 64: 759-768.
- Pope, A.L., Moir, R.J., Somers, M., Underwood, E.J. and White, C.E., 1979. The effects of sulphur on Se absorption and retention in sheep. J. Nutri., 109: 1448-1455.
- Pugh, D.G., Rankins, D.L., Eason, J.T., Wenzel, J.G.W. and Spano, J.S., 1994. The effect of feeding broiler litter on the serum calcium, phosphorus and magnesium concentration of beef brood cows. Vet. Clin. Nutr., 1: 18-22.
- Puls, R., 1994. Mineral Levels in Animal Health: Diagnostic Data, Trinity Western University Press, Clearbook, British Colombia, Canada, pp. 250-252.
- Rankins, D.L., Eason, J. T., McCaskey, T.A., Stephenson, A.H. and Floyd, J.G., 1993. Nutritional and toxicological evaluation of three deep-stacking methods for the processing of broiler litter as a foodstuff for beef cattle. Anim. Prod., 56: 321-326.
- Robertson, J.B. and Van Soest, P.J., 1981. The detergent system of analysis and its application to human foods. In: W. P. T. James and O. Theander (Eds), The Analysis of Dietary Fibre in Food. Marcel Dekker, New York. pp. 123-158.

Rossi, J.E., Goetsch, A.L., Patil, A.R., Kouakou, B., Park, K.K., Wang, Z.S..

Galloway, D.L. and Johnson, Z.B., 1996. Effects of forage level in broiler litter-based diets on feed intake, digestibility and particulate passage rate in Holstein steers at different live weights. *Anim. Feed Sci. Technol.*, 62:163-177.

Rossi, J.E., Goetsch, A.L. and Galloway, D.L., 1998. Intake and digestion by Holstein steers consuming different particle size fractions of broiler litter. *Anim. Feed Sci. Technol.*, 71: 145- 156.

Rude, B.J. and Rankins, D.L., 1997. Mineral status in beef cows fed broiler litter diets with cation-anion differences or supplemented with hay. *J. Anim. Sci.*, 75: 727-735.

Ruffin, B.G., and McCaskey, T.A., 1990. Broiler litter can serve as feed ingredient for beef cattle. *Feedstuffs*, 62(15):13-17.

Ruffin, B.G. and McCaskey, T.A., 1998. Feeding broiler litter to beef cattle. Circular ANR-557, <http://gallus.tamu.edu/waste.bfcattle.html>

Rule, D.C., Busboon, J.R. and Kercher, C.J., 1994. Effect of dietary canola on fatty acid composition of bovine adipose tissue, muscle, kidney and liver. *J. Anim. Sci.*, 72: 2735-2744.

Sankari, S., 1993. Methods for evaluation of selenium status. *Norwegian J. Agric. Sci. Suppl.* 11, 51-56. Statistical Analytical Analysis System, 1994. SAS User's Guide:Statistics, Version 6. SAS Institute Inc. Cary, N C, USA.

Shand, P.J., McKinnon, J.J. and Christensen, D.A., 1998. Eating quality of beef from animals fed brewers grain and wheat-based wet distillers' grains. *Can. J. Anim. Sci.*, 78:143-146.

- Shihabi, Z.K., Hindsdale, M.E. and Bleyer, A.J., 1995. Xanthine analysis in biological fluids by capillary electrophoresis. *J. Chromatogr.*, 669:163-169.
- Shuller, M.L., 1987. Utilization of farm waste for food. In: Silanikove, N., Holzer, Z., Cohen, D., Benjamin, R., Gutman, M. and Meltzer, A., 1976. Interrelationship between metabolism of tritiated water, sodium and dry matter intake, by beef cows fed poultry litter and wheat straw in free choice. *Comp. Biochem. Physiol.*, 88A, pp. 113-118.
- Siddons, R.C., Nolan, J.V., Beever, D.E. and MacRae, J.C., 1985. Nitrogen digestion and metabolism in sheep consuming diets containing contrasting forms and levels of nitrogen. *Brit. J. Nutr.*, 54:175 -187.
- Silanikove, N., Holzer, Z., Cohen, D., Benjamin, R., Gutman, M. and Meltzer, A., 1987. Interrelationship between metabolism of tritiated water, ²²sodium and dry matter intake by beef cows fed poultry litter and wheat straw in free choice. *Comp. Biochem. Physiol.* 88A, 113-118.
- Silanikove, N. and Tiomkin, D., 1992. Toxicity induced by poultry litter consumption: effect on measurements reflecting liver function in beef cows. *Anim. Prod.*, 54:203-209.
- Silva, L.A., Van Horn, H.H., Olaloka, E.A., Wilcox, C.J. and Harris, S., 1976. Complete rations for dairy cattle : VII: Dried poultry waste for lactating cows. *J. Dairy Sci.*, 59: 2071-2076.
- Smith, L.W., Goering, H.K. and Gordon, C.H., 1971. Nutritive evaluation of untreated and chemically treated dairy cattle waste. *Proc. Int. Symp. Livestock Wastes*, ASAE Publ., 314.
- Smith, L.W. and Calvert, C.C., 1976. Dehydrated broiler excreta versus soyabean meal as nitrogen supplements for sheep. *J. Anim. Sci.*, 43:1286-1292.

- Smith, L.W. and Calvert, C.C., 1976. Dehydrated poultry excreta versus soybean meal as nitrogen supplements for sheep. *J. Anim. Sci.*, 43: 1286-1292.
- Smith, O.B., McLeod, G.K., Mowat, D.N., Fox, C.A. and Moran, E.T., 1978. Performance and health of calves fed wet caged layer excreta as a protein supplement. *J. Anim. Sci.*, 47: 833-842.
- Smith, L.W. and Wheeler, W.E., 1979. Nutritional and economic value of animal excreta. *J. Anim. Sci.*, 48: 144-156.
- Smith, L.W., Calvert, C.C. and Cross, R.H., 1979. Dehydrated poultry excreta vs. cottonseed meal as nitrogen supplements for Holstein steers. *J. Anim. Sci.*, 48: 633-640.
- Southwell, B.L., Hale, O.M. and McCormick, W.C., 1958. Poultry house litter as protein supplement in steer fattening rations. *Ga. Agr. Exp. Sta. Mimeo. Series N. S.* 55.
- Statistical Analysis System, 1994. SAS User's Guide; Version 6. SAS Inst. Inc. Cary, North Carolina, USA.
- Stowe, H.D. and Herdt, T.H., 1992. Clinical assessment of selenium status of livestock. *J. Anim. Sci.*, 70:3928-3933.
- Symonds, H.W., Mather, D.L. and Collis, K.A., 1981. The maximum capacity of the liver of the adult dairy cow to metabolize ammonia. *Br. J. Nutr.*, 46:481-486.
- Tagari, H., Levy, D., Holzer, Z. and Illan, D., 1976. Poultry litter for intensive beef production. *Anim. Prod.*, 23:317-320.
- Tietz, N. W., 1976. Fundamentals of clinical chemistry (2nd edn). W B Saunders, Philadelphia.

- Tilley, J. M. A. and Terry, R.A., 1964. A two-stage technique for the in vitro digestion of forage crops. *J.Br Grassld. Soc.*, 18:104-111.
- Tinnimit, P., Yu, Y., McGuffy, K. and Thomas, J.W., 1972. Dried animal waste as a protein supplement for sheep. *J. Anim., Sci.*35: 431- 438.
- Titgemeyer, E.C., 1997. Design and interpretation of nutrient digestion studies. *J. Anim. Sci.*, 75:2235-2247.
- Ulrey, D. E., 1992. Basis for regulation of selenium supplements in animal diets. *J. Anim. Sci.*, 70:3922-3927.
- Valdivia, R., Ammerman, C.B., Henry, P.R., Feaster, J.P. and Wilcox, C.J., 1982. Effect of dietary aluminum and phosphorus on performance, phosphorus utilization and tissue mineral composition in sheep. *J. Anim. Sci.*, 7:402-410.
- Van der Lugt, J. J., Henton, M. M. and Steyn, B. G., 1996. Type C botulism in sheep associated with the feeding of poultry litter. *J. S. Afr. Vet. Assoc.*, 67:2-7.
- Van der Merwe, H.J., Pretorius, P.S. and du Toit, J.E.S., 1975. Benutting van rantsoene met battery-hoender-mis as proteienbron. *S. Afr. Tydskr. Veek.*, 5:17 – 21.
- Van der Westhuizen, A. A. and Hugo, J.M., 1972. Die benutting van batteryhoendermis in oorwinteringsrant-soene van jong vleisbeeste. *S. Afr. Tydskr. Veek.*, 2:13.
- Van Ryssen, J.B.J., Channon, P. and Stielau, W.J., 1977. Mineral and nitrogen in poultry manure. *S. Afr. J. Anim. Sci.*,7: 195-199.
- Van Ryssen, J.B.J. and Jagoe, H.M., 1982. Retention of trace elements in the livers of sheep fed poultry manure as a ration component. *S. Afr. J. Anim. Sci.*, 11:273-278.

Van Ryssen, J. B. J., 1988. Poultry litter as a feedstuff for ruminants. Proc. Symp. Natal Branch, SASAS.

Van Ryssen, J.B.J., 1991. Effect of monensin and its metabolites in broiler litter on sheep consuming the broiler litter. J. S. Afr. Vet. Assoc., 62: 94-99.

Van Ryssen, J.B.J., Van Malsen, S. and Verbeek, A.A., 1993. Mineral composition of poultry manure in South Africa with reference to the farm feed act. S. Afr. J. Anim. Sci., 23:54-57.

Van Ryssen, J.BJ., Van Malsen, P.M.S. and Hartmann, F., 1998. Contribution of dietary sulphur to the interaction between selenium and copper in sheep. J. Agric. Sci., Camb., 130: 107-114.

Van Soest, J.P., 1982. Nutritional Ecology of the Ruminant. Cornell Univ. Press. Ithaca. NY.

Van Soest, P.J., 1994. Nutritional Ecology of the Ruminant. 2nd edn. Cornell Univ. Press. Ithaca.NY.

Vest, L. and Dyer, J., 1993. Many factors affect broiler litter's mineral composition. Poultry Digest (September, 1993).

Ways, P and Hanahan, D.J., 1964. Characterization and quantification of red cell lipids in normal man. J. Lipid Res., 5:318-328.

Webb, E.C., 1992. The influence of dietary energy levels on subcutaneous fatty acid profiles and meat quality of sheep. Master of Science (Agriculture), Thesis. University of Pretoria.

Webb, E.C., Casey, N.H. and Van Niekerk, W.A., 1994a. Fatty Acids in the subcutaneous adipose tissue of intensively fed South African Mutton Merino and Dorper wethers. Meat Sci., 38:123-131.

Webb, E.C., Bosman, M.J.C. and Casey, N. H., 1994b. Dietary influences on subcutaneous fatty acid profiles and sensory characteristics of Dorper and South African Mutton Merino wethers. *S. Afr. J. Food Sci. Nutr.*, 6:45-50.

Webb, E.C., Bosman, M.J. and Casey, N.H., 1997. Influence of dietary presentation on the composition of fatty acids and sensory characteristics of meat from wethers. *S. Afr. J. Food Sci. Nutr.*, 9:69-76.

Webb, K.E., Phillips, W.A., Libke, K.G., Harmon, B.W. and Fontenot, J.P., 1973. Different levels of broiler litter in ewe rations. *J. Anim. Sci.*, 36 : 218.(abstr.).

Westing, T.W., Fontenot, J.P., McClure, W.H., Kelly, R.F. and Webb, K.E., 1985. Mineral element profiles of animal wastes and edible tissues from cattle fed animal wastes. *J. Anim. Sci.*, 61: 670-681.

Westlake, K. and Dutton, M.F., 1985. The incidence of mycotoxins in litter, feed and livers of chickens in Natal. *S. A. J. Anim. Sci.*, 15:175 .

Wright, P. L., Bell, M. C., 1966. Comparative metabolism of selenium and tellurium in sheep and swine. *Am. J. Physiol.*, 211: 6-10.

Wuethrich, H.U., 1978. Recycling the manure of cattle, pigs and poultry by feeding to animals. Seminar for the Feed Industry, Cesme, Turkey, Roche Information Service.

Yan, T., Offer, N.W. and Roberts, D.J., 1996. The effect of dietary nitrogen sources and levels on rumen fermentation, nutrient degradation and digestion and rumen microbial activity by wether sheep given a high level of molasses. *Anim. Sci.*, 63:123-131.

Zindel, H.C. and Flegal, C.J., 1971. "Economics of dried poultry waste (DPW) as a feed ingredient or a fertilizer." In: Poultry Pollution: Research results, research report 152, Agric. Exper. Sta., Michigan State Univ., East Lansing, Michigan, November (1971).

Zinn, R.A. and Owens, F.N., 1986. A rapid procedure for purine measurement and its use for estimating net ruminal protein synthesis. *Can. J. Anim. Sci.*, 66: 157-166.

Zinn, R.A., Barajas, R., Montano, M. and Shen, Y., 1996. Protein and energy values of dehydrated poultry excreta in diets for feedlot cattle. *J. Anim. Sci.*, 74: 2331-2335.