

REFERENCES

- Aguirre, G., Bellon, M.R. and Smale, M. 1998.** A regional analysis of maize biological diversity in Southeastern Guanajuato, Mexico. Pp. 22. CIMMYT Economics Working Paper 98-06. Mexico.
- Ahn, S. and Tanksley, S.D. 1993.** Comparative linkage maps of the rice and maize genomes. *Proc. Natl. Acad. Sci. USA* **90**: 7980-7984.
- Ajmone-Marsan, P., Castiglioni, P., Fusari, F., Kuiper, M. and Motto, M. 1998.** Genetic diversity and its relationship to hybrid performance in maize as revealed by RFLP and AFLP markers. *Theor. Appl. Genet.* **96**:219-227.
- Alika, J.E., Aken'ova, M.E. and Fatokun, C.A. 1993.** Variation among maize (*Zea mays* L.) accessions of Bendel State, Nigeria. Multivariate analysis of agronomic data. *Euphytica* **66**: 65-71.
- Appa Rao, S., Mengesha, M.H., Gopal Reddy, V. and Prasada Rao, K.E. 1998.** Collecting and evaluation of sorghum germplasm from Rwanda. *Plant Genet. Res. Newslett.* **114**: 26–28.
- Arias, C.A.A., deSouza, C.L. and Takeda, C. 1999.** Path coefficient analysis of ear weight in different types of progeny in maize. *Maydica* **44**: 251-262.
- Ayana, A. and Bekele, E. 2000.** Geographical patterns of morphological variation in sorghum (*Sorghum bicolor* (L) Moench) germplasm from Ethiopia and Eritrea: Quantitative characters. *Euphytica* **115**: 91-104.
- Barbosa-Neto, J.F., Hernandez, C.M., O'Donoghue, L.S. and Sorrells, M.E. 1997.** Precision of genetic relationship estimates based on molecular markers. *Euphytica* **98**: 59-67.

- Beaumont, M.A., Ibrahim, K.M., Boursot, P. and Bruford, M.W. 1998.** Measuring genetic distance. Pp.315-325. In A. Karp *et al.* (ed.) Molecular tools for screening biodiversity. Chapman and Hall, London.
- Bennetzen, J.L. and Devos, K. 2002.** Plant synteny, colinearity and genome evolution. Encyclopedia of Life Sciences. Pp.1-5. Macmillan Publishers Ltd.
- Bogyo, T.P., Porceddu, E. and Perrino, P. 1990.** Analysis of sampling strategies for collecting genetic material. *Econ. Bot.* **34**: 11-86.
- Bouchez, A., Hospital, F., Causse, M., Gallais, A. and Charcosset, A. 2002.** Marker-assisted introgression of favorable alleles at quantitative trait loci between maize elite lines. *Genetics* **162**: 1945-1959.
- Brown, A.H.D. 1989a.** The case for core collections. Pp. 136-156. In A.H.D. Brown *et al.* (ed.), The use of plant genetic resources. Cambridge University press, Cambridge.
- Brown, A.H.D. 1989b.** Core collections: A practical approach to genetic resources management. *Genome* **31**: 818-824.
- Buckler, E.S. and Thornsberry, J.M. 2002.** Plant molecular diversity and application to genomics. *Curr. Opin .Plant Biol.* **5**: 107-111.
- Buckler, E.S., Thornsberry, J.M. and Kresovich, S. 2001.** Molecular diversity, structure and domestication of grasses. *Genet. Res.* **77**: 213-218.
- Burr, B., Burr, F.A., Thompson, K.H., Albertsen, M.C. and Stuber, C.W. 1988.** Gene mapping and recombinant inbreds in maize. *Genetics* **118**: 519- 526.
- Burstin, J. and Charcosset, A. 1997.** Relationships between phenotypic and marker distance: Theoretical and experimental investigation. *Heredity* **78**: 477-483.
- Causse, M., Santoni, S., Damerval, C., Maurice, A., Charcosset, A., Deatrick, J.**

- and de Vienne, D. 1996.** A composite map of expressed sequences in maize. *Genome* **39**: 418-432.
- CIMMYT, 2000.** International Maize and Wheat Improvement Center (CIMMYT). CGIAR Research areas of research: Maize (*Zea mays* L.) <http://www.cgiar.org/areas/maize.htm>.
- Coe, E. and Neuffer, M.G. 1993.** Genetic loci and linkage map of corn (*Zea mays* L.) (2n = 20), Pp. 6157-6189. In S.J. O'Brien (ed.), Genetic Maps: Locus maps of complex genomes, 6th edition, Cold Spring Harbor Press, Cold Spring Harbor, NY.
- Coe, E., Hancock, D., Kowaleski, S. and Polacco, M. 1995.** Gene list and working maps. *Maize Genet. Coop. Newslett.* **69**: 191-267.
- Crossa, J., Hernandez, C.M., Bretting, P., Eberhart, S.A. and Taba, S. 1993.** Statistical genetic considerations for maintaining germplasm collections. *Theor. Appl. Genet.* **86**: 673-678.
- CSA. 1998.** Central Statistical Authority (CSA): Agricultural sample survey 1997/98. Report on farm management practices of 'belg' season . Vol. **III**. *Statistical bulletins 171*. Addis Ababa, Ethiopia.
- CSA. 2001.** Central Statistical Authority (CSA): Agricultural sample survey 2000/2001 on area production of major crops, private peasant holdings 'meher' season. Vol. **III**. *Statistical bulletins 190*. Addis Ababa. Ethiopia.
- Davis, G.L., McMullen, M.D., Baysdorfer, C., Musket, T., Grant, D., Staebell, M., Xu, G., Polacco, M., Koster, L., Melia-Hancock, S., Houchines K., Chao, S. and Coe, E. 1999.** A maize map standard with sequenced core markers, grass genome reference points and 932 expressed sequenced tagged

sits (ESTs) in a 1736-locus map. *Genetics* **152**: 1137-1172.

Demeke, T., Sasikumar, B., Hucl, P. and Chibbar, R.N. 1997. Random amplified polymorphic DNA (RAPD) in cereal improvement. *Maydica* **42**: 133-142.

Doebley, J.F. 1990. Molecular evidence for gene flow among *Zea* species. *Bio Science* **40**: 443- 448.

Doebley, J.F. and Iltis, H.H. 1980. Taxonomy of *Zea* (Gramineae): I. A subgeneric classification with a key to taxa. *Am. J. Bot.* **67**: 982-993.

Doebley, J.F., Goodman, M.M. and Stuber C.W. 1984. Isoenzymatic variation in *Zea* (Gramineae). *Systematic Bot.* **9**: 203-218.

Dubreuil, P., Dufour, P., Krejet, E., Causse, M., de Vienne, D., Gallais, A. and Charcosset, A. 1996. Organization of RFLP diversity among inbred lines of maize representing the most significant heterotic groups. *Crop Sci.* **36**: 790-799.

EARO. 2000. Ethiopian Agricultural Research Organization (EARO), Research strategy plan for maize. Addis Ababa, Ethiopia.

Edwards, M.D., Stuber, C.W. and Wendel, J.F. 1987. Molecular-marker facilitated investigations of quantitative traits in maize. I. Numbers, genomic distribution and types of gene action. *Genetics* **116**: 113-125.

Enoki, A., Sato, H. and Koinuma, K. 2002. SSR analysis of genetic diversity among maize inbred lines adapted to cold regions of Japan. *Theor. Appl. Gene.* **104**: 1270-1277.

Eubanks, M. 2000. International application published under the patent cooperation treaty (PCT)WO00/07434. Novel genetic materials for transmission into maize.

- Evans, L.T. and Evans, T.L.T. 1993.** Crop evolution, adaptation, and yield. Cambridge University Press; New York.
- Everitt, B. 1980.** Cluster analysis. 2nd edition, Halstead Press, New York.
- Excoffier, L., Smouse, P.E. and Quattro, J.M. 1992.** Analysis of molecular variance inferred from metric distances among DNA haplotypes: Application to human mitochondrial DNA restriction data. *Genetics* **131**: 479-491.
- Eyre-Walker, A., Gaut, R.L., Hilton, H. and Gaut, B.S. 1998.** Investigation of the bottleneck leading to the domestication of maize. *Proc. Natl. Acad. Sci. USA* **95**: 4441-4446.
- Falconer, D.S. and Mackay, T.F.C. 1996.** Introduction to quantitative genetics. 4th edition, Longman Ltd. Group, London.
- FAPRI. 2003.** U.S. and World Agriculture Outlook. Food and Agricultural Policy Research Institute. Iowa State University. Ames, Ia. U.S. (<http://www.fapri.iastate.edu/outlook2003/>).
- Felsenstein, J. 1993.** PHYLIP (Phylogeny Inference Package) version 3.5c. Distributed by the author, department of genetics, University of Washington, Seattle, Washington.
- Flint-Garcia, S.A., Thornsberry, J.M. and Buckler, E.S. 2003.** Structure of linkage disequilibrium in plants. *Annu. Rev. Plant Biol.* **54**: 357-374.
- Gardiner, J.M., Coe, E.H., Melia-Hancock, S., Hoisington, D.A. and Chao, S. 1993.** Development of a core RFLP map in maize using an immortalized F₂ population. *Genetics* **134**: 917-930.
- Gaut, B.S. 2001.** Patterns of chromosomal duplication in maize and their implications for comparative maps of the grasses. *Gen. Research* **11**: 55-56.

- Gaut, B.S., d'Ennequin, M.T., Peek, A.S. and Sawkins, M.C. 2000.** Maize as a model for the evolution of plant nuclear genomes. *Proc. Natl. Acad. Sci. USA* **97**: 7008-7015.
- Gauthier, P., Gouesnard, B., Dallard, J., Redaelli, R., Rebourg, C., Charcosset, A. and Boyat, A. 2002.** RFLP diversity and relationships among traditional European maize populations. *Theor. Appl. Genet.* **105**: 91-99.
- Georges, M., Nielsen, D., Mackinnon, M., Mishra, A., Okimoto, R., Pasquino, A.T., Sargeant, L.S., Sorensen, A., Steele, M.R., Zao, X., Womack, J.E. and Hoeschle, A. 1995.** Mapping quantitative traits loci controlling milk production in dairy cattle by exploiting progeny testing. *Genetics* **139**: 907-920.
- Grossniklaus, U., Koltunow, A. and van-Lookeren-Campagne, M. 1998.** A bright future for apomixis. *Trends Plant Sci.* **3**: 415-416.
- Guthridge, K.M., Dupal, M.P., Kolliker, R., Jones, E.S., Smith, K.F. and Forster, J.W. 2001.** AFLP analysis of genetic diversity within and between populations of perennial ryegrass (*Lolium perenne* L.). *Euphytica* **122**: 191-201.
- Hafnagel, H.P. 1961.** Agriculture in Ethiopia. Food and Agricultural Organization of the United Nations, Rome, Italy.
- Hair, J.R., Anderson, R.E., Tatham, R.L. and Black, W.C. 1995.** Multivariate data analysis with readings. 4th edition, Prentice-Hall, Englewood Cliffs, NJ.
- Hake, S. and Walbot, V. 1980.** The genome of *Zea mays*, its organization and homology to related grasses. *Chromosoma* **79**: 251-270.
- Hallauer, A.R. and Miranda, J.B. 1988.** Quantitative genetics in maize breeding.

Iowa State Univ. Press, Ames, Iowa.

Harjes, C.E., Smith, M.E., McCouch, S.R. and Tanksley, S.D. 1999. Advanced backcross QTL analysis and introgression of perennial teosinte (*Zea diploperennis*) alleles to maize. *Plant and Animal Genome VII* Conference, January 17-21, San Diego, California.

Helentjaris, T., Weber, D. and Wright, S. 1988. Identification of genomic locations of duplicate nucleotide sequences in maize by analysis of restriction fragment length polymorphisms. *Genetics* **118**: 353-363.

Helentjaris, T., King, G., Slocum, M., Siedenstrang, C. and Wegman, S. 1985. Restriction fragment polymorphisms as probes for plant diversity and their development as tools for applied plant breeding. *Plant Mol. Biol.* **5**: 109-118.

Helentjaris, T., Slocum, M., Wright, S., Schaefer, A. and Nienhuis, J. 1986. Construction of genetic linkage maps in maize and tomato using restriction fragment length polymorphisms. *Theor. Appl. Genet.* **72**: 761-769.

Henandez, Z.E. 1985. Maize and man in the greater Southwest. *Econ. Bot.* **39**: 416-430.

Henry, A.M. and Damerval, C. 1997. High rates of polymorphism and recombination at the opaque-2 locus in cultivated maize. *Mol. Gen. Genet.* **256**: 147-157.

Hoisington, D.A. and Ribaut, J. M. 1998. What can biotechnology offer Africa? - Concrete examples. Six Eastern and Southern Africa Regional Maize Conference. Pp. 12-17, Addis Ababa, Ethiopia.

Hoisington, D.A., Khairallah, M., Reeves, T., Ribaut, J.M., Skovmand, B., Taba, S. and Warburton M. 1999. Plant genetic resources: What can they

contribute toward increased crop productivity? *Proc. Natl. Acad. Sci. USA* **96**: 5937-5943.

Hulbert, S.H., Richter, T.E., Axtell, J.D. and Bennetzen, J.L. 1990. Genetic mapping and characterization of sorghum and related crops by means of maize DNA probes. *Proc. Natl. Acad. Sci. USA* **87**: 4251-4255.

Jaccard, P. 1908. Nouvelles recherches sur la distribution florale. *Bull. Soc. Vaudoise Sci. Natl.* **44**: 223-270.

Jerry, H. 2000. Number Cruncher Statistical Systems (NCSS). Statistical software, Utah.

Johnson, A.R. and Wichern, D.W. 1992. Applied multivariate statistical analysis. 3rd edition, Prentice-Hall, Englewood Cliffs, NJ.

Jones, C.J., Edwards, K.J., Castiglioni, S., Winfield, M.O., Sala, F., Van de Wiel, C., Bredemeijer, G., Vosman, B., Matths, M., Daly, A., Brettschneider, R., Bettini, P., Buitti, M., Maestri, E., Malcevski, A., Marmirdi, N., Aert, R., Volkaert, G., Rueda, J., Linacero, R., Vazquez, A. and Karp, A. 1997. Reproducibility testing of RAPD, AFLP, and SSR markers in plants by a network of European laboratories. *Mol. Breed.* **3**: 381-389.

Jones, N., Ougham, H. and Thomas, H. 1997. Markers and mapping: we are all geneticists now. *New Phytologist* **137**: 165-177.

Karp, A., Seberg, O. and Buiatti, M. 1996. Molecular techniques in the assessment of botanical diversity. *Ann. Bot.* **78**: 143-149.

Karp, A., Edwards, K., Bruford, M., Vosman, B., Morgante, M., Seberg, O., Kremer, A., Boursot, P., Arctander, P., Tautz, D. and Hewitt, G. 1997. Newer molecular technologies for biodiversity evaluation: Opportunities and

challenges. *Nature Biotechnol.* **15**: 625-628.

Koester, R., Sisco, P. and Stuber, C. 1993. Identification of quantitative trait loci controlling days to flowering and plant height in two near-isogenic lines of maize. *Crop Sci.* **33**: 1209–1216.

Kolliker, R., Jones, E.S., Jahufer, M.Z.Z. and Forster, J.W. 2001. Bulked AFLP analysis for the assessment of genetic diversity in white clover (*Trifolium repens* L.). *Euphytica* **121**: 305-315.

Kumar, L.S. 1999. DNA markers in plant improvement: An overview. *Biotechnology advances* **17**: 143-182.

Kumar, M.V.N. and Kumar, S.S. 2000. Studies on character association and path coefficient for grain yield and oil content in maize (*Zea mays* L.). *Ann. Agric. Res.* **21**: 73-78.

Leblanc, O., Grimanelli, D., González-de-León, D. and Savidan, Y. 1995. Detection of the apomictic mode of reproduction in maize-Tripsacum hybrids using maize RFLP markers. *Theor. Appl. Genet.* **90**: 1198-1203.

Lee, M., Sharopova, N., Beavis, W.D., Grant, D., Katt, M., Blair, D.L. and Hallauer, A.R. 2002. Expanding the genetic map of maize with intermated B73 x MO17 (IBM) population. *Plant Mol. Bio.* **48**: 453-461.

Levinson, G. and Gutman, G.A. 1987. Slipped-strand mispairing: A major mechanism for DNA sequence evolution. *Mol. Biol. Evol.* **4**: 203-221.

Lin, Y.R., Schertz, K.F. and Paterson, A.H. 1995. Comparative analysis of QTLs affecting plant height and maturity across the Poaceae, in reference to interspecific sorghum population. *Genetics* **141**: 391- 411.

Link, W.C., Dickens, C., Singh, M., Schwall, M. and Melchinger, A.E. 1995.

Genetic diversity in European and Mediterranean fababean germplasm revealed by RAPD markers. *Theor. Appl. Genet.* **90**: 27-32.

Lubberstedt, T., Melchinger, A.E., Duñle, C., Vuylsteke, M. and Kuiper, M. 2000. Relationships among early European maize inbreds: IV. Genetic diversity revealed with AFLP markers and comparison with RFLP, RAPD, and pedigree data. *Crop Sci.* **40**: 783-791.

Lucchin, M., Baraccia, G. and Parrini, P. 2003. Characterization of flint maize (*Zea mays* L. convar. Mays) Italian landraces: Morpho-phenological and agronomic traits. *Genet. Res. Crop Evol.* **50**: 315-327.

Malecot, G. 1948. Les mathematiques de l'heredite. Masson and Cie Paris.

Manly, B.F.J. 1991. Randomization and Monte Carlo methods in biology. Chapman and Hall, New York.

Mantel, N. 1967. The detection of disease clustering and a generalized regression approach. *Cancer Res.* **27**: 209-220.

Martin, J.M., Blake, T.K. and Hockett, E.A. 1991. Diversity among North American spring barley cultivars based on coefficients of parentage. *Crop Sci.* **31**: 1131-1137.

Matsuoka, Y., Mitchell, S.E., Kresovich, S., Goodman, M. and Dobeley J. 2002. Microsatellite in *Zea* –variability, patterns of mutations and use for evolutionary studies. *Theor. Appl. Genet.* **104**: 436-450.

Matsuoka, Y., Vigouroux, Y., Goodman, M.M., Sanchez, J.G., Buckler, E. and Doebley, J. 2001. A single domestication for maize shown by multilocus microsatellite genotyping. *Proc. Natl. Acad. Sci. USA* **99**: 6080-6084.

Melchinger, A.E. 1993. Use of RFLP markers for analysis of genetic relationships

among breeding materials and prediction of hybrid performance. Pp. 621-627. In D.R. Buxton *et al.* (ed.) First international crop science congress. CSSA, Madison, WI.

Melchinger, A.E., Gumber, R.K., Leipert, R.B. and Vuysteke, M. 1998.

Predication of testcross means and variance among F_3 progenies of F_1 crosses from testcross means and genetic distances of their parents in maize. *Theor. Appl. Genet.* **96**: 503-512.

Messmer, M.M., Melchinger, A.E., Herrmann, R.G. and Boppenmaier, J. 1993.

Relationship among early European maize inbreds: Comparison of pedigree and RFLP data. *Crop Sci.* **33**: 944-950.

Michelmore, R.W., Paran, I. and Kessli, R.V. 1991. Identification of markers linked

to disease resistance genes by bulked segregant analysis: A rapid method to detect markers in specific genomic regions using segregenat populations. *Proc. Natl. Acad. Sci. USA* **88**: 9828-9832.

Mitchell, S.E., Kresovich, S., Jester, C.A., Henandez, C.J. and Szewe-McFadden,

A.K. 1997. Application of multiplex PCR and fluorescence-based, semi-automated allele sizing technology for genotyping plant genetic resources. *Crop Sci.* **37**: 617-624.

Moeller, D.A. and Schaal, B.A. 1999. Genetic relationships among Native American

maize accessions of the Great Plains assessed by RAPDS. *Theor. Appl. Genet.* **99**:1061-1067.

Mohammadi, S.A., Prasanna, B.M., Sudan, C. and Singh, N.N. 2002. A

microsatellite marker based study of chromosomal regions and gene effects on yield and yield components in maize. *Cellu. Mol. Biol. Lett.***7**: 599-606.

- Moreau, L., Lemarié, S., Charcosset, A. and Gallais, A. 2000.** Economic efficiency of one cycle of marker-assisted selection. *Crop Sci.* **40**: 329-337.
- Morgante, M., Hanafey, M. and Powell, W. 2002.** Microsatellites are preferentially associated with non-repetitive DNA in plant genomics. *Nat. Genet.* **30**:194-200.
- Mulatu, K., Bogale, G., Tolessa, B., Desalegn, Y. and Afeta A. 1992.** Maize production trends and research in Ethiopia. In: B. Tolessa and J.R. Ransom (eds.). Proceeding of the first national maize workshop in Ethiopia. Addis Ababa, Ethiopia.
- Myburg, A.A., Remington, D.L., O'Malley, D.M., Sederoff, R.R. and Whetten, R.W. 2001.** High-throughput AFLP analysis using infrared dye-labeled primers and an automated DNA sequencer. *BioTechniques* **30**: 348-357.
- Nei, M. and Li, W.H. 1979.** Mathematical model for studying genetic variation in terms of restriction endonucleases. *Proc. Natl. Acad. Sci. USA* **76**: 5269-5273.
- Neter, J., Wasserman, W., and Kutner, M.H. 1990.** Applied linear statistical models. 3rd edition, Irwin, Inc., Homewood, Illinois.
- NMSA. 2000.** National Meteorology Service Agency (NMSA), monthly weather report, Feb. 1980 to Dec. 2000. Addis Ababa, Ethiopia.
- Nuel, G., Baril C.P. and Robin, S. 2000.** Varietal distinctness assisted by molecular markers: A methodological approach. *Acta. Hort.* **546**: 65-71.
- Ohta, T. and Kimura, M. 1973.** A model of mutation appropriate to estimate the number of electrophoretically detectable alleles in a finite population. *Genet. Res.* **22**:201-204.

- Pejic, I., Ajmone-Marson, P., Morgante, M., Kozumplick, V., Castiglioni, P., Tarmino, G. and Motto, M. 1998.** Comparative analysis of genetic similarity among maize inbred line detected by RFLPs, RAPDs, and AFLPs. *Theor. Appl. Genet.* **97**: 1248-1255.
- Pinto, L.R., Vieira, M.L.C., de Souza, C.L.P. and de Souza, A.P. 2003.** Genetic diversity assessed by microastellites in tropical maize populations submitted to a high-intensity reciprocal recurrent selection. *Euphytica* **134**: 277-286.
- Powell, W., Morgante, M., Andre, C., Hanafey, M., Vogel, J., Tingey, S. and Rafalski, A. 1996.** The comparison of RFLP, RAPD, AFLP and SSR (microsatellite) markers for germplasm analysis. *Mol. Breeding* **2**: 225-238.
- Quarrie, S.A., Lazic-Jancic, V., Kovacevic, D., Steed, A. and Pekic, S. 1999.** Bulk segregant analysis with molecular markers and its use for improving drought resistance in maize. *J. Exp. Botany* **50**: 1299-1306.
- Ragot, M., Gay, G., Muller, J.P. and Durovray, J. 2000.** Efficient selection for adaptation to the environment through QTL mapping and manipulation in maize. Pp. 128-130. In: Ribaut, J.M. and Poland, D. (eds.) Molecular approaches for the genetic improvement of cereals for stable production in water-limited environments. A strategic planning workshop held at CIMMYT, Et Batan, Mexico, June 21-25, 1999.
- Rebourg, C., Gouesnard, B. and Charcosset, A. 2001.** Large-scale molecular analysis of traditional European maize populations: Relationships with morphological variation. *Heredity* **86**: 574-587.
- Reif, J.C., Melchinger, A.E., Xia, X.C., Warburton, M.L., Hoisington, D.A., Vasal, S.K., Beck, D., Bohan, M. and Frisch. 2003.** Use of SSRs for

establishing heterotic groups in subtropical maize. *Theor. Appl. Genet.* **107**: 947-957.

Remington, D. L., Thornsberry, J. M., Matsuoka, Y., Wilson, L. M., Whitt, S. R., Doebley, J. F., Kresovitch, S., Goodman, M. M. and Buckler, E. S. 2001. Structure of linkage disequilibrium and phenotypic associations in the maize genome. *Proc. Natl. Acad. Sci. USA* **98**: 11479-11484.

Ribaut, J.M. and Hoisington, D. 1998. Marker-assisted selection: new tools and strategies. *Trends Plant Sci.* **31**: 236-239.

Ribaut, J.M., Edmeades, G., Perotti, E. and Hoisington, D. 2002. QTL analyses, MAS results, and perspectives for drought-tolerance improvement in tropical maize. Pp. 131-136. In: Ribaut, J.M. and Poland, D. (eds.) Molecular approaches for the genetic improvement of cereals for stable production in water-limited environments. A strategic planning workshop held at CIMMYT, Et Batan, Mexico, June 21-25, 1999.

Riek, J.De., Calsyn, E., Everaet, I., van Bockstaele, E. and De Loose, M. 2001. AFLP based alternative for the assessment of distinctness, uniformity and stability in sugar beet varieties. *Theor. Appl. Genet.* **103**: 1254-1265.

Roldan-Ruiz, L., Van Eeuwijk, F.A., Gilliland, T.J., Dubreuil, P., Dillmann, C., Lallemand, J., De Loose, M. and Baril, C.P. 2001. A comparative study of molecular and morphological methods of describing relationships between perennial ryegrass (*Lolium perenne* L.) varieties. *Theor. Appl. Genet.* **103**: 1138-1150.

Saghai-Marouf, M.A., Biyashev, R.M., Yang, G.P., Zhang, Q. and Allard, R.W. 1994. Extraordinarily polymorphic microsatellite DNA in barely: Species

diversity, chromosomal locations and population dynamics. *Proc. Natl. Acad. Sci. USA* **91**: 5466-5470.

SAS. 1993. SAS/STAT users'. Version 6.09. SAS Institute Inc carry, NC.

Sasakawa-Global2000. 2002. Agricultural project in Ethiopia: Activities and output. An assessment report, Sasakawa-Global 2000, Addis Ababa, Ethiopia.

Savidan, Y., Grimanelli, D., Perotti, E. and Leblanc, O. 1998. International application published under the patent cooperation treaty (PCT) WO98/36090. Means for identifying nucleotide sequences involved in apomixis.

Sax, K. 1923. The association of size difference with seed-coat pattern and pigmentation in *Phaseolus Volgaris*. *Genetics* **8**: 552-560.

Senior, M.L. and Heun, M. 1993. Mapping maize microsatellites and polymerase chain reaction conformation of the targeted repeats using a CT primer. *Genome* **36**: 884-889.

Senior, M.L., Murphy, J.P., Goodman, M.M. and Stuber, C.W. 1998. Utility of SSRs for determining genetic similarities and relationships in maize using an agarose gel system. *Crop Sci.* **38**: 1088-1098.

Senior, M.L., Chin, E.C.L., Lee, M., Smith, J.S.C. and Stuber, C.W. 1996. Simple sequence repeat markers developed from maize sequences found in the GENE BANK database: Map construction. *Crop Sci.* **36**: 1676-1683.

Sharopova, N., McMullen, M.D., Schulz, L., Schroeder, S., Sancock-Villeda, H., Gardiner, J., Bergstrom, D., Houchins, K., Melia-Hancock, S., Musket, T., Duru, N., Polacco, M., Edwards, K., Ruff, T., Register, J.C., Brouwer, C., Thompson, R., Velasco, R., Chin, E., Lee, M., Woodman-

- Clikeman, W., Long, M. J., Liscum, E., Cone, K., Davis, G. and JrCoe, E.H. 2002.** Development and mapping of SSR markers for maize. *Plant Mol. Biol.* **48**: 463-481.
- Singh, R.K. and Chaudhary, B.D. 1977.** Biometrical methods in quantitative genetics analysis. Kalyani Publishers, New Delhi.
- Skinner, D.Z., Bauchan, G.R., Auricht, G. and Huges S. 1999.** A method for the efficient management and utilization of large germplasm collections. *Crop Sci.* **39**: 1237-1242.
- Smith, J.S.C. and Smith, O.S. 1992.** Fingerprinting crop varieties. *Adv. Agron.* **47**: 85-140.
- Sneath, P.H.A. and Sokal, R.R. 1973.** Numerical taxonomy. WH Freeman, San Francisco.
- Snedecor, G.W. and Cochran, W.G. 1980.** Statistical Methods. 7th edition, Iowa State Univ. Press, Ames.
- Sokal, R.R. and Michener, C.D. 1958.** A statistical method for evaluating systematic relationships. *Univ. Kansas Sci. Bull.* **38**: 1409-1438.
- Souza, E. and Sorrells, M.E. 1991.** Relationships among 70 American oat germplasm. I. Cluster analysis using quantitative characters. *Crop Sci.* **31**: 599-605.
- Stanton, M.A., Stewart, J.M., Percival, A.E. and Wendel, J.F. 1994.** Morphological diversity and relationships in A-genome cottons, *Gossypium arboreum* and *G. herbaceum*. *Crop Sci.* **34**: 519-527.
- Statistical for Windows. 1995.** Stat Soft. Inc., Computer program manual. Tulsa. OK: Stat Soft. Inc., 2325 East 13th Street Tulsa.

- Stuber C.W. 1989.** Marker-based selection for quantitative traits. Pp.31-49:In: Proc. XII Congr. EUCARPIA. Parey, Berlin, Hamburg.
- Stuber, C.W., Goodman, M.M. and Moll, R.H. 1982.** Improvement of yield and ear number resulting from selection at allozyme loci in a maize population. *Crop Sci.* 22:737-740.
- Stuber, C.W., Polacco, M. and Senior, M.L. 1999.** Synergy of empirical breeding, marker-assisted selection, and genomics to increase crop yield potential. *Crop Sci.* 39: 1571-1583.
- Taba, S., Diaz, J., Franco, J. and Crossa, J. 1998.** Evaluation of Caribbean maize accessions to develop a core subset. *Crop Sci.* 38: 1378-1386.
- Tanksley, S.D. and McCouch, S.R. 1997.** Seed banks and molecular maps: unlocking genetic potential from the wild. *Science* 277: 1063–1066.
- Tanksley, S.D., Grandillo, S., Fulton, T.M., Zamir, D., Eshed, T., Petiard, V., Lopez, J. and Beck-Bunn, T. 1996.** Advance backcross QTL analysis in a cross between an elite processing line of tomato and its wild relative *L. pimpinellifolium*. *Theor. Appl. Genet.* 92: 213-224.
- Tato, K. 1964.** Rainfall in Ethiopia. *Eth. Geog. J.* 2: 28-36.
- Tautz, D. 1989.** Hypervariability of simple sequences as a general source for polymorphic DNA markers. *Nucleic Acids Res.* 17: 6463-6471.
- Thormann, C.E., Ferreira, M.E., Camargo, L.E.A., Tivang, J.G. and Osborn, T.C. 1994.** Comparison of RFLP and RAPD markers to estimating genetic relationships within and among cruciferous species. *Theor. Appl. Genet.* 88: 973-980.
- Thornsberry, J.M., Goodman, M.M., Doebley, J., Kresovich S., Nielsen, D. and**

- Edward, S.E. 2001.** *Dwarf8* polymorphisms associate with variation in flowering time. *Nat. Genet.* **28**: 286-289.
- Tuberosa, R., Salvi, S. and Phillips, S. 1998.** Bulk segregant analysis confirms the importance of the region near *umc89a* for days to pollen shed in maize. *Maize Gene. Newsl.* **72**: 71-72.
- Tuberosa, R., Salvi, S., Sanguineti, M.C., Landi, P., Maccaferri, M. and Conti, S. 2002.** Mapping QTLs regulating morph-physiological traits and yield: Case studies, shortcomings and perspectives in drought-stressed maize. *Annals Botany* **89**: 941-963.
- Twumasi-Afryie, S., Habatmu, Z., Kass, Z., Bayisa, Y. and Sewagegne, T. 2001.** Development and improvement of highland maize in Ethiopia. pp.1-12. 2nd national maize workshop of Ethiopia, Addis Ababa, Ethiopia.
- Uptmoor, R., Wenzel, W., Friedt, W., Donaldson, G., Ayisi, K. and Ordon, F. 2003.** Comparative analysis of the genetic relatedness of *Sorghum bicolor* accessions from Southern Africa by RAPDs, AFLPs and SSRs. *Theor. Appl. Genet.* **106**: 1316-1325.
- Van Eeuwijk, F.A. and Baril, C.P. 2001.** Conceptual and statistical issues related to the use of molecular markers for distinctness and essential derivation. *Acta Hort.* **546**: 35-53.
- Vavilov, N.I. 1951.** Phylogeographic basis of plant breeding: The origin, variation, immunity and breeding of cultivated plants. *Chron. Bot.* **13**: 13-54.
- Veldboom, L.R., Lee, M. and Woodman, W.L. 1994.** Molecular marker-facilitated studies in an elite maize populations: Linkage analysis and determination of QTL for morphological traits. *Theor. Appl. Genet.* **88**: 7-16.

- Virk, P.S., Ford-Lioyed, B., Jackson, M.T., Pooni, H.S., Clemeno, T.P. and Newbury, H.J. 1996.** Predicting quantitative variation within rice germplasm using molecular markers. *Heredity* **76**: 296-304.
- Vos, P., Hogers, R., Bleeker, M., Reijans, M., VanDe Lee, T., Hornes, M., Frijters, A., Pot, J., Poleman, J., Kuiper, M. and Zabeau, M. 1995.** AFLP®: a new technique for DNA fingerprinting. *Nucleic Acids Res.* **23**: 4407-4414.
- Vuylsteke, M., Kuiper, M. and Stam, P. 2000a.** Chromosomal regions involved in hybrid performance and heterosis: Their AFLP-based identification and practical use in prediction models. *Heredity* **85**: 208-218.
- Vuylsteke, M., Mank, R., Brugmans, B., Stam, P. and Kuiper, M. 2000b.** Further characterization of AFLP data as a tool in genetic diversity assessments among maize (*Zea mays* L.) inbred lines. *Mol. Breed.* **6**: 265-276.
- Vuylsteke, M., Mank, R., Antonise, R., Bastiaans, E., Senior, M.L., Stuber, C.W., Melchinger, A.E., Lubberstedt, T., Xia, X.C., Stam, P., Zabeau, M. and Kuiper M. 1999.** Two high-density AFLP linkage maps of *Zea mays* L.: Analysis of distribution of AFLP markers. *Theor. Appl. Genet.* **99**: 921-935.
- Wang, R.L., Stec, A., Hey, J., Lukens, L. and Doebley, J. 1999.** The limits of selection during maize domestication. *Nature* **398**: 236-239.
- Warburton, M.L. and Crossa, J. 2000.** Data analysis in CIMMYT applied biotechnology center for fingerprinting and genetic diversity studies. CIMMYT, Mexico.
- Warburton, M.L., Zianchun, X., Crossa, J., Franco, J., Melchinger, A.E., Frisch, M., Bohn, M. and Hoisington, D. 2002.** Genetic characterization of

CIMMYT inbred maize lines and open pollinated populations using large scale fingerprinting methods. *Crop Sci.* **42**: 1832-1840.

Ward, J.H. 1963. Hierarchical grouping to optimize an objective function. *Am. Stat. Assoc. J.* **56**: 236-244.

Waugh, R., Bonar, N., Baird, E., Thomas, B., Graner, A. and Hayes, P. 1997. Homology of AFLP products in three mapping populations of barely. *Mol. Gen. Genet.* **255**: 311-321.

Welsh, J. and McClell, M. 1990. Fingerprinting genomes using PCR with arbitrary primers. *Nucleic Acids Res.* **18**:7213-7218.

Wendel, J.F. 2000. Genome evolution in polyploids. *Plant Mol. Biol.* **42**: 225-249.

Wiley, E.O. 1981. Phylogenetics: The theory and practice of phylogenetics and systematics. John Wiley, New York.

Williams, J.G.K., Kubelick, A.R., Livak, K.J., Rafalski J.A. and Tingey, S.V. 1990. DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Res.* **18**: 6531-6535.

Wilson, R.N., Heckman, J.W. and Somerville, C.R. 1992. Gibberellin is required for flowering in *Arabidopsis-thaliana* under short days. *Plant Physiol.* **100**: 403-408.