

# First Report of *bla*<sub>IMP-14</sub> on a Plasmid Harboring Multiple Drug Resistance Genes in *Escherichia coli* Sequence Type 131

Nicole Stoesser,<sup>a</sup> Anna E. Sheppard,<sup>a</sup> Gisele Peirano,<sup>b,c</sup> Robert P. Sebra,<sup>d</sup> Tarah Lynch,<sup>c</sup> Luke W. Anson,<sup>a</sup> Andrew Kasarskis,<sup>d</sup> Mary R. Motyl,<sup>e</sup> Derrick W. Crook,<sup>a</sup> Johann D. Pitout<sup>b,c,f,g,h</sup>

Modernizing Medical Microbiology Consortium, Nuffield Department of Medicine, John Radcliffe Hospital, University of Oxford, Oxford, United Kingdom<sup>a</sup>; Division of Microbiology, Calgary Laboratory Services, Calgary, Alberta, Canada<sup>b</sup>; Department of Pathology and Laboratory Medicine, University of Calgary, Calgary, Alberta, Canada<sup>c</sup>; Icahn Institute and Department of Genetics and Genomic Sciences, Icahn School of Medicine, Mount Sinai, New York, New York, USA<sup>d</sup>; Clinical Microbiology, Merck and Co. Inc., Rahway, New Jersey, USA<sup>e</sup>; Department of Microbiology, Immunology, and Infectious Diseases, University of Calgary, Calgary, Alberta, Canada<sup>f</sup>; Snyder Institute for Chronic Diseases, University of Calgary, Calgary, Alberta, Canada<sup>g</sup>; Department of Medical Microbiology, University of Pretoria, Pretoria, South Africa<sup>h</sup>

**The *bla*<sub>IMP-14</sub> carbapenem resistance gene has largely previously been observed in *Pseudomonas aeruginosa* and *Acinetobacter* spp. As part of global surveillance and sequencing of carbapenem-resistant *Escherichia coli*, we identified a sequence type 131 strain harboring *bla*<sub>IMP-14</sub> within a class 1 integron, itself nested within an ~54-kb multidrug resistance region on an epidemic IncA/C<sub>2</sub> plasmid. The emergence of *bla*<sub>IMP-14</sub> in this context in the ST131 lineage is of potential clinical concern.**

The emergence of carbapenemases in clinically prevalent *Escherichia coli* lineages such as sequence type 131 (ST131) is a major problem for the management of patients infected with these strains (1, 2). Globally, five major transmissible carbapenemase enzymes predominate, represented by the KPC, OXA-48-like, NDM, VIM, and IMP families (2, 3).

An IMP metallo-beta-lactamase enzyme (IMP-1) was first detected in Japan in *Pseudomonas aeruginosa* in the late 1980s (4); since then, 52 genetically diverse *bla*<sub>IMP</sub> gene variants 738 to 747 bp in length have been identified (5). Most *bla*<sub>IMP</sub> variants have been isolated from either *Pseudomonas* or *Acinetobacter* spp. and demonstrate a degree of geographic structuring (6); however, some, such as *bla*<sub>IMP-4</sub> and *bla*<sub>IMP-8</sub>, have emerged successfully in members of the family *Enterobacteriaceae* and are distributed over wider geographic regions (6, 7). Associations of *bla*<sub>IMP</sub> with *E. coli* ST131 have, to date, been restricted to *bla*<sub>IMP-4</sub> and *bla*<sub>IMP-8</sub> in Taiwan, China, and Australia (8–11). As part of the Merck Study for Monitoring Antimicrobial Resistance Trends (SMART) (1), we identified an IMP-14-producing ST131 *E. coli* isolate, Ecol<sub>732</sub>, that was isolated in Bangkok, Thailand, in 2012 and was sequenced in order to ascertain the genetic structures associated with this IMP variant in ST131.

Ecol<sub>732</sub> was obtained from the urine of a hospitalized elderly male with a lower urinary tract infection. The MICs of ampicillin-sulbactam, piperacillin-tazobactam, ceftazidime, cefepime, ertapenem, imipenem, amikacin, and ciprofloxacin were determined with microdilution panels prepared at International Health Management Associates, Inc. (Schaumburg, IL, USA), in accordance with 2015 CLSI guidelines. It tested non-susceptible (i.e., either intermediate or resistant) to the above-mentioned agents. The MICs of colistin and tigecycline (determined by E tests) were 0.12 and 1 mg/liter, respectively.

DNA (chromosomal plus plasmid) was extracted from pure overnight subcultures of the isolate for both PacBio (long-read) sequencing and Illumina MiSeq (short-read) sequencing with the Qiagen Genomic-tip 100/G kit and the QIAamp DNA minikit (catalogue numbers 10243 and 51304; Qiagen, Valencia, CA), respectively. Preliminary *de novo* assembly of PacBio reads with HGAP3 was performed; resulting contigs were annotated with Prokka (12) and then trimmed on the basis of sequence/anno-

tation overlaps in Geneious (version 9.04) (13). One-hundred-fifty-base paired-end MiSeq reads for each of the isolates were trimmed with Trimmomatic (version 0.35) (14) and then mapped to the corresponding PacBio assemblies with BWA mem (version 0.7.9a-r786) (15). Read pileups were inspected to confirm the structural integrity of the contigs and correct any small errors in the assembled contigs. Unmapped MiSeq reads were assembled with A5MiSeq (16) in order to identify any small plasmids (<7 kb) that may have been filtered out during the size selection process implemented as part of PacBio library preparation. Additional annotation focused on resistance genes and insertion sequences was performed with reference to the ResFinder (17), PlasmidFinder (18), and ISFinder (19) databases.

The Ecol<sub>732</sub> genome consists of a 5,009,900-bp chromosome and six plasmids, five of which could be fully resolved. These included pEC732-IMP14 (186,826 bp, IncA/C<sub>2</sub>), pEC732\_2 (129,154 bp, IncFII/FIA/FIB/col), pEC732\_3 (82,588 bp, IncB/O/K/Z), pEC732\_4 (4,072 bp, untyped), and pEC732\_5 (1,549 bp, untyped). A partial sixth *mob* plasmid fragment was also present (4,204 bp, untyped). The *bla*<sub>IMP-14</sub> sequence in pEC732-IMP14 differed from the reference AY553332 by a single synonymous substitution (A249C), resulting in the same amino acid sequence. It was located within a 3,791-bp class 1 integron, In687 [*intI1-bla*<sub>IMP-14</sub>-*aac*(6')-*qacE* $\Delta$ -*sulI*]. This integron is almost identical (single nucleotide difference, A925G) to that in *Achromobacter*

Received 19 April 2016 Returned for modification 11 May 2016

Accepted 25 May 2016

Accepted manuscript posted online 31 May 2016

Citation Stoesser N, Sheppard AE, Peirano G, Sebra RP, Lynch T, Anson LW, Kasarskis A, Motyl MR, Crook DW, Pitout JD. 2016. First report of *bla*<sub>IMP-14</sub> on a plasmid harboring multiple drug resistance genes in *Escherichia coli* sequence type 131. *Antimicrob Agents Chemother* 60:5068–5071. doi:10.1128/AAC.00840-16.

Address correspondence to Nicole Stoesser, nicole.stoesser@ndm.ox.ac.uk.

Copyright © 2016 Stoesser et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).




FIG 1 Plasmid pEC732-IMP14 with a detailed view of the ARI-A-like 54-kb resistance region (bottom).


*xylosoxidans* strain R4, which was cultured from a urine sample, also in Thailand (GenBank accession number KJ406505).

The backbone of pEC732-IMP14 was highly similar to that of prototype type 1 IncA/C<sub>2</sub> plasmid pRMH760 (RefSeq database accession no. NC\_023898; from a *Klebsiella pneumoniae* strain) and other type 1 IncA/C<sub>2</sub> plasmids (recently reviewed in reference 20), almost all of which also include a specific region designated ARI-A that contains a variable array of resistance genes and is located 1,711 bp upstream of *rhs* (20). Similarly, in pEC732-IMP14, the *bla*<sub>IMP-14</sub>-harboring integron was part of a much larger, 54,454-bp ARI-A-like region containing antimicrobial, heavy metal, and biocide resistance genes (Fig. 1), including those encoding resistance to beta-lactams (*bla*<sub>OXA-10</sub>, *bla*<sub>IMP-14</sub>), macrolides (drug efflux), rifampin (*arr-2*), sulfonamides (*sul1*), aminoglycosides [*aadA1*, *aadB*, *aph*(3')-IVa, *aac*(6')] chloramphenicol (*cmlA7*), chromate (*srpC*), mercury (*mer* operon), and quaternary ammonium compounds (*qac*). Some of these were part of a second, novel, integron designated In1286 (*int1*-*qacH*-*aadB*-*arr-2*-*cmlA7*-*bla*<sub>OXA-10</sub>-*aadA1*).

An alignment of pEC732-IMP14, prototype IncA/C<sub>2</sub> type 1 plasmid pRMH760, and the only publicly available type 1 IncA/C<sub>2</sub> sequence from Thailand, pR148 (RefSeq database accession no.

NC\_019380, from *Aeromonas hydrophila* [21]), demonstrates the genetic similarity of these plasmids (Fig. 2). All three sequences were >99% similar in the 1- to 86,573-bp region and in the ~27.5-kb region downstream of ARI-A (Fig. 2). Differences in pEC732-IMP14 include a region of clustered single-nucleotide variants suggestive of a recombination event (region, 3,100 to 8,000 bp) and the acquisition of two integrase subunits (regions, 86,573 to 89,203 and 90,138 to 200,167 bp; Fig. 2). Interestingly, the pR148-containing *A. hydrophila* strain was identified on a Thai tilapia fish farm that had successively used several antimicrobial classes (21).

To date, *bla*<sub>IMP-14</sub> has not been described in *E. coli*, to our knowledge, and has largely previously been reported in *P. aeruginosa* and *Acinetobacter baumannii* strains by several hospital centers in Thailand, in some cases as part of clonal outbreaks (22–25). Although *bla*<sub>IMP-14</sub> is similarly associated with class 1 integrons in these cases, as in pEC732-IMP14, the wider plasmid contexts and sequences of these integrons in *P. aeruginosa* and *A. baumannii* strains have not been investigated. It is, however, conceivable that the *bla*<sub>IMP-14</sub>-harboring integron observed in pEC732-IMP14 and *A. xylosoxidans* strain R4 has been exchanged more widely with *Pseudomonas* and *Acinetobacter* spp. in Thailand. Class 1 integrons



**FIG 2** Alignment of pEC732-IMP14, pR148, and pRMH760 demonstrating mean percent pairwise identity over alignment columns (green, 100%; olive, 30 to 70%; red <30%; no color, 0%). Vertical black lines in the sequence bars represent differences between the nucleotide sequences with respect to the pEC732-IMP14 reference; thin horizontal black lines represent deleted regions. The *repA* gene, variable ARI-A resistance region, and *rhs* gene are indicated, as are the region of clustered single-nucleotide variants downstream of the *repA* region and two additional mobile genetic elements present in pEC732-IMP14 and absent from the other two sequences.

have been linked with the recent successful spread and expansion of another metallo-beta-lactamase, *bla*<sub>VIM</sub>, in IncA/C<sub>2</sub> plasmids in members of the family *Enterobacteriaceae* in Greece (26) and *bla*<sub>VIM</sub> and *bla*<sub>IMP</sub> in Spain (27).

The presence of the extensively drug-resistant region observed here on an epidemic IncA/C<sub>2</sub> plasmid in an *E. coli* ST131 strain from Thailand is therefore of concern and may represent wider, regional, horizontal dissemination of *bla*<sub>IMP-14</sub> mediated by mobile genetic elements across bacterial families. The homology of pEC732-IMP14 with an *A. hydrophila* plasmid found on a fish farm and the presence of *bla*<sub>IMP</sub>-harboring plasmids in *E. coli* in other environmental (28) and animal sampling frames (29) suggest that the transmission network for IMP-positive *E. coli* may extend beyond the health care setting. Broad surveillance and control measures that are targeted at both community and health care contexts may be required to monitor and limit *bla*<sub>IMP</sub> dissemination.

**Nucleotide sequence accession numbers.** Complete sequence data for Ecol\_732 have been deposited in GenBank under BioProject number PRJNA316786. The accession numbers of the sequences are CP015138 (chromosome), CP015139 (pEC732-IMP14), CP015140 (pEC732\_2), CP015141 (pEC732\_3), CP015142 (pEC732\_4), CP015143 (pEC732\_5), and CP015144 (pEC732\_6 [partial sequence only]).

#### ACKNOWLEDGMENTS

We are grateful for the support of the Modernizing Medical Microbiology Informatics Group. We also thank Daryl Hoban and the Merck SMART surveillance team.

N.S. is currently funded through a Public Health England/University of Oxford clinical lectureship; the sequencing work was also partly funded through a previous Wellcome Trust doctoral research fellowship (099423/Z/12/Z). Additional funding support was provided by a research grant from Calgary Laboratory Services (10006465) and by the Health Innovation Challenge Fund (a parallel funding partnership between the Well-

come Trust [WT098615/Z/12/Z] and the Department of Health [grant HICF-T5-358]). This research was supported by the National Institute for Health Research (NIHR) Oxford Biomedical Research Centre Programme.

The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication. The views expressed are those of the authors and not necessarily those of the National Health Service, the NIHR, or the Department of Health.

#### FUNDING INFORMATION

This work, including the efforts of Nicole Stoesser, Anna E. Sheppard, Luke W. Anson, and Derrick W. Crook, was funded by The Wellcome Trust (099423/Z/12/Z and WT098615/Z/12/Z). This work, including the efforts of Nicole Stoesser, Anna E. Sheppard, Luke W. Anson, and Derrick W. Crook, was funded by The United Kingdom Department of Health (DH) (HICF-T5-358). This work, including the efforts of Gisele Peirano and Johann D. Pitout, was funded by Calgary Laboratory Services (CLS) (10006465).

#### REFERENCES

1. Peirano G, Bradford PA, Kazmierczak KM, Badal RE, Hackel M, Hoban DJ, Pitout JD. 2014. Global incidence of carbapenemase-producing *Escherichia coli* ST131. *Emerg Infect Dis* 20:1928–1931. <http://dx.doi.org/10.3201/eid2011.141388>.
2. Mathers AJ, Peirano G, Pitout JD. 2015. The role of epidemic resistance plasmids and international high-risk clones in the spread of multidrug-resistant *Enterobacteriaceae*. *Clin Microbiol Rev* 28:565–591. <http://dx.doi.org/10.1128/CMR.00116-14>.
3. Nordmann P, Naas T, Poirel L. 2011. Global spread of carbapenemase-producing *Enterobacteriaceae*. *Emerg Infect Dis* 17:1791–1798. <http://dx.doi.org/10.3201/eid1710.110655>.
4. Watanabe M, Iyobe S, Inoue M, Mitsuhashi S. 1991. Transferable imipenem resistance in *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 35:147–151. <http://dx.doi.org/10.1128/AAC.35.1.147>.
5. Bush K, Palzkill T, Jacoby G. 2015.  $\beta$ -Lactamase classification and amino acid sequences for TEM, SHV and OXA extended-spectrum and inhibitor resistant enzymes. <http://www.lahey.org/studies/>. Accessed 1 March 2016.
6. Cornaglia G, Giamarellou H, Rossolini GM. 2011. Metallo-beta-lactamases: a last frontier for beta-lactams? *Lancet Infect Dis* 11:381–393. [http://dx.doi.org/10.1016/S1473-3099\(11\)70056-1](http://dx.doi.org/10.1016/S1473-3099(11)70056-1).

7. Sidjabat HE, Townell N, Nimmo GR, George NM, Robson J, Vohra R, Davis L, Heney C, Paterson DL. 2015. Dominance of IMP-4-producing *Enterobacter cloacae* among carbapenemase-producing *Enterobacteriaceae* in Australia. *Antimicrob Agents Chemother* 59:4059–4066. <http://dx.doi.org/10.1128/AAC.04378-14>.
8. Ho PL, Cheung YY, Wang Y, Lo WU, Lai EL, Chow KH, Cheng VC. 2016. Characterization of carbapenem-resistant *Escherichia coli* and *Klebsiella pneumoniae* from a healthcare region in Hong Kong. *Eur J Clin Microbiol Infect Dis* 35:379–385. <http://dx.doi.org/10.1007/s10096-015-2550-3>.
9. Yan JJ, Tsai LH, Wu JJ. 2012. Emergence of the IMP-8 metallo-beta-lactamase in the *Escherichia coli* ST131 clone in Taiwan. *Int J Antimicrob Agents* 40:281–282. <http://dx.doi.org/10.1016/j.ijantimicag.2012.05.011>.
10. Zhang F, Zhu D, Xie L, Guo X, Ni Y, Sun J. 2015. Molecular epidemiology of carbapenemase-producing *Escherichia coli* and the prevalence of ST131 subclone H30 in Shanghai, China. *Eur J Clin Microbiol Infect Dis* 34:1263–1269. <http://dx.doi.org/10.1007/s10096-015-2356-3>.
11. Sidjabat HE, Robson J, Paterson DL. 2015. Draft genome sequences of two IMP-4-producing *Escherichia coli* sequence type 131 isolates in Australia. *Genome Announc* 3:e00983-15. <http://dx.doi.org/10.1128/genomeA.00983-15>.
12. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <http://dx.doi.org/10.1093/bioinformatics/btu153>.
13. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton A, Markowitz S, Duran C, Thierer T, Ashton B, Metjies P, Drummond A. 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28:1647–1649. <http://dx.doi.org/10.1093/bioinformatics/bts199>.
14. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <http://dx.doi.org/10.1093/bioinformatics/btu170>.
15. Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25:1754–1760. <http://dx.doi.org/10.1093/bioinformatics/btp324>.
16. Coil D, Jospin G, Darling AE. 2015. A5-miseq: an updated pipeline to assemble microbial genomes from Illumina MiSeq data. *Bioinformatics* 31:587–589. <http://dx.doi.org/10.1093/bioinformatics/btu661>.
17. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 67:2640–2644. <http://dx.doi.org/10.1093/jac/dks261>.
18. Carattoli A, Zankari E, Garcia-Fernandez A, Voldby Larsen M, Lund O, Villa L, Moller Aarestrup F, Hasman H. 2014. *In silico* detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. *Antimicrob Agents Chemother* 58:3895–3903. <http://dx.doi.org/10.1128/AAC.02412-14>.
19. Siguier P, Perochon J, Lestrade L, Mahillon J, Chandler M. 2006. ISfinder: the reference centre for bacterial insertion sequences. *Nucleic Acids Res* 34:D32–36. <http://dx.doi.org/10.1093/nar/gkj014>.
20. Harmer CJ, Hall RM. 2014. pRMH760, a precursor of A/C(2) plasmids carrying bla<sub>CMY</sub> and bla<sub>NDM</sub> genes. *Microb Drug Resist* 20:416–423. <http://dx.doi.org/10.1089/mdr.2014.0012>.
21. Del Castillo CS, Hikima J, Jang HB, Nho SW, Jung TS, Wongtavatchai J, Kondo H, Hirono I, Takeyama H, Aoki T. 2013. Comparative sequence analysis of a multidrug-resistant plasmid from *Aeromonas hydrophila*. *Antimicrob Agents Chemother* 57:120–129. <http://dx.doi.org/10.1128/AAC.01239-12>.
22. Khuntayaporn P, Montakantikul P, Santanirand P, Kiratisin P, Chomnawang MT. 2013. Molecular investigation of carbapenem resistance among multidrug-resistant *Pseudomonas aeruginosa* isolated clinically in Thailand. *Microbiol Immunol* 57:170–178. <http://dx.doi.org/10.1111/1348-0421.12021>.
23. Piyakul C, Tiyawisutrisri R, Boonbumrung K. 2012. Emergence of metallo-beta-lactamase IMP-14 and VIM-2 in *Pseudomonas aeruginosa* clinical isolates from a tertiary-level hospital in Thailand. *Epidemiol Infect* 140:539–541. <http://dx.doi.org/10.1017/S0950268811001294>.
24. Kansakar P, Dorji D, Chongtrakool P, Mingmongkolchai S, Mokmake B, Dubbs P. 2011. Local dissemination of multidrug-resistant *Acinetobacter baumannii* clones in a Thai hospital. *Microb Drug Resist* 17:109–119. <http://dx.doi.org/10.1089/mdr.2010.0062>.
25. Samuelsen O, Toleman MA, Sundsfjord A, Rydberg J, Leegaard TM, Walder M, Lia A, Ranheim TE, Rajendra Y, Hermansen NO, Walsh TR, Giske CG. 2010. Molecular epidemiology of metallo-beta-lactamase-producing *Pseudomonas aeruginosa* isolates from Norway and Sweden shows import of international clones and local clonal expansion. *Antimicrob Agents Chemother* 54:346–352. <http://dx.doi.org/10.1128/AAC.00824-09>.
26. Papagiannitsis CC, Dolejska M, Izdebski R, Giakkoupi P, Skalova A, Chudejova K, Dobiasova H, Vatopoulos AC, Derde LP, Bonten MJ, Gniadkowski M, Hrabak J. 2016. Characterisation of IncA/C2 plasmids carrying an In416-like integron with the bla<sub>VIM-19</sub> gene from *Klebsiella pneumoniae* ST383 of Greek origin. *Int J Antimicrob Agents* 47:158–162. <http://dx.doi.org/10.1016/j.ijantimicag.2015.12.001>.
27. Zamorano L, Miro E, Juan C, Gomez L, Bou G, Gonzalez-Lopez JJ, Martinez-Martinez L, Aracil B, Conejo MC, Oliver A, Navarro F. 2015. Mobile genetic elements related to the diffusion of plasmid-mediated AmpC beta-lactamases or carbapenemases from *Enterobacteriaceae*: findings from a multicenter study in Spain. *Antimicrob Agents Chemother* 59:5260–5266. <http://dx.doi.org/10.1128/AAC.00562-15>.
28. Kieffer N, Poirel L, Bessa LJ, Barbosa-Vasconcelos A, da Costa PM, Nordmann P. 2016. VIM-1, VIM-34, and IMP-8 carbapenemase-producing *Escherichia coli* strains recovered from a Portuguese river. *Antimicrob Agents Chemother* 60:2585–2586. <http://dx.doi.org/10.1128/AAC.02632-15>.
29. Dolejska M, Masarikova M, Dobiasova H, Jamborova I, Karpiskova R, Havlicek M, Carlile N, Priddel D, Cizek A, Literak I. 2016. High prevalence of Salmonella and IMP-4-producing *Enterobacteriaceae* in the silver gull on Five Islands, Australia. *J Antimicrob Chemother* 71:63–70. <http://dx.doi.org/10.1093/jac/dkv306>.