Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehponline@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Levels and Determinants of DDT and DDE Exposure in the

VHEMBE Cohort

Fraser W. Gaspar, Jonathan Chevrier, Lesliam Quirós-Alcalá, Jonah M. Lipsitt, Dana Barr, Nina Holland, Riana Bornman, and Brenda Eskenazi

Table of Contents

- 1. Quantification of *o*,*p*' and *p*,*p*'-DDT and DDE Isomers
- 2. Quality Control Results for VHEMBE Serum Analysis
- 3. Additional Results
 - **Table S1.** Propensity score distributions for estimating E(A|W) for assessing positivity violations.
 - **Table S2.** Algorithm weights used by SuperLearner to estimate E(Y|A,W)
 - **Table S3.** Algorithm weights used by SuperLearner to estimate E(A|W).
 - **Table S4.** p,p'-DDT and p,p'-DDE concentrations (ng/g-lipid) by selected characteristics in VHEMBE participants, Limpopo, South Africa.
 - **Table S5.** Comparison of VHEMBE lipid-adjusted serum concentrations with previously reported blood concentrations in adults living in IRS communities and pregnant women living in the United States
 - **Figure S1.** Directed Acyclical Graph (DAG) used to conceptualize the relationship between exposures and outcomes.
 - **Figure S2.** Spatial distribution of p,p'-DDE concentrations in relation to Tshilidzini Hospital

References