

Local antagonism and resource partitioning between two invasive pine plantation pests

Mesfin Wondafrash^{1,2,3}, Bernard Slippers^{2,4}, Brett P. Hurley^{1,2}, Jeff Garnas^{2,5}

Correspondence: Mesfin Wondafrash, Department of Zoology and Entomology, Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria, 0002, South Africa, email address: Mesfin.Gossa@fabi.up.ac.za , cell phone: +27 (61) 242 7403

Keywords: facilitation; wood borer; phloem-feeding insect; *Pinus*; woodwasp; *Pissodes*; *Sirex noctilio*; plantation pests

Running title: Interaction between two forest insect pests

¹Department of Zoology and Entomology, University of Pretoria, Pretoria, 0002, South Africa

²Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria, 0002, South Africa

³School of Plant Sciences, Haramaya University, Dire Dawa, P.O. Box 38, Ethiopia

⁴Department of Biochemistry, Genetics and Microbiology, Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria, 0002, South Africa

⁵Department of Natural Resources and the Environment, University of New Hampshire, Durham, NH 03824, USA

Abstract

1. The woodwasp *Sirex noctilio* Fabricius, native to Eurasia and North Africa, regularly co-occurs with a North American origin *Pissodes* sp. on *Pinus* trees in South Africa. The nature of this co-occurrence and potential impacts on either of the species is unknown.
2. Using structured sampling, we studied the pattern and degree of co-occurrence of *S. noctilio* and *Pissodes* sp. in six sites in *P. patula* plantations in South Africa to better understand the potential for interactions and population-level feedbacks. We compared density, adult body size and within-tree distribution of both insect species in trees where they co-occurred or occurred singly.
3. *Sirex noctilio* and *Pissodes* sp. co-occurred on $68.0 \pm 3.8\%$ of infested trees (range: 55-80%). Both insect species were more abundant in trees where they occurred alone relative to co-inhabited trees.
4. The within-tree distribution of *S. noctilio* did not differ in the presence of *Pissodes* sp.; however, in the presence of *S. noctilio*, *Pissodes* sp. were more frequent in the bottom than the middle tree section where *S. noctilio* was at higher density and this could be an indication of resource partitioning. Body size of *S. noctilio* and *Pissodes* sp. was not influenced by co-occurrence.
5. Our results suggest broad overlap for both insects in preference for stressed trees in similar states of decline, and moderate antagonism between the species leading to lower emergence of both species in shared trees. The influence of *S. noctilio* on *Pissodes* sp. population densities may still be a net positive at a landscape scale as *Pissodes* sp. prefers declining or recently dead trees which are readily created by *S. noctilio* via attacks on healthy trees.

Keywords: facilitation; wood borer; phloem-feeding insect; *Pinus*; woodwasp; *Pissodes*; *Sirex noctilio*; plantation pests

Introduction

Biological invasion is occurring at an unprecedented rate as a result of increasing global trade and transport, threatening native biota and community integrity in invaded regions (Mack *et al.*, 2000; Sala *et al.*, 2000; Stein *et al.*, 2000; Simberloff, 2011; Schaffner *et al.*, 2013). The likelihood of interaction among sympatric invaders increases with introduced species richness, with potential consequences for their ecology and evolution as well as for invasive impacts (Garnas *et al.*, 2016). These interactions may be direct or indirect, symmetric or asymmetric, strong or weak, pairwise or diffuse, novel or co-evolved in the native range (Denno *et al.*, 1995; Wingfield *et al.*, 2016). Additionally, species interactions may result in positive or negative feedbacks depending in part on whether they are fundamentally antagonistic (e.g., predatory, parasitic, or competitive), mutualistic or facilitative (Hawkins, 2005; Heimpel *et al.*, 2010; Hajek *et al.*, 2016).

Both facilitative and antagonistic interactions are well known in forest insect systems (Dewey *et al.*, 1974; Barbosa & Wagner, 1988). For example, primary tree-killing bark beetles such as the Southern pine beetle *Dendroctonus frontalis* Zimmermann facilitate a suite of secondary colonizers that take advantage of weakened tree defences following mass attack (Hofstetter *et al.*, 2006). Similarly, co-occurrence with other native bark beetles facilitates persistence of low-density endemic populations in the Mountain pine beetle *Dendroctonus ponderosae* Hopkins (Smith *et al.*, 2011). Facilitation of certain species via interactions with shared hosts may be particularly dramatic where invasive species are concerned (Rodriguez, 2006). For example, invaders such as the Emerald ash borer *Agrilus planipennis* Fairmaire provide a short term

pulse in resource availability for saproxylic insects as well as native predators, including woodpeckers (Klooster *et al.*, 2018), and may have facilitated range expansion in at least one other buprestid beetle in Europe (Orlova-Bienkowskaja, 2015). In some cases, the cascade of facilitative interactions among invaders can be quite large (e.g., up to eleven species; Heimpel *et al.* (2010)). Antagonistic interactions among invasive insects are also common. For example, in the eastern United States, introduced Hemlock woolly adelgid *Adelges tsugae* (Annand), and Elongate hemlock scale *Fiorinia externa* Ferris limit one another's populations on eastern hemlock by approximately 30% in mixed versus single species treatments (Preisser *et al.*, 2008; Preisser & Elkinton, 2008). Negative interactions between *F. externa* and the introduced circular hemlock scale *Nuculapsis tsugae* (Marlatt) have also been reported, where the density of the latter was reduced by 74% in the presence of the former (McClure, 1981).

In some cases, an aggressive invader may arrive that alters ecological conditions and/or feedbacks for already-established invaders. Such is the case with the weevil *Pissodes* sp. (Curculionidae, Coleoptera) and the European woodwasp *Sirex noctilio* Fabricius (Siricidae, Hymenoptera), which have both become established in *Pinus* plantations in South Africa, as well as in other parts of the world (Webb, 1974; Hurley *et al.*, 2007; Slippers *et al.*, 2015). *Pissodes* sp. was first detected in South Africa in 1942 and identified as the deodar weevil *Pissodes nemorensis* Germar, a native to North America (Webb, 1974; EPPO, 1999). Recent research, however, has shown that it is not the deodar weevil but rather an unrecognized species of the *P. strobi* complex of North American origin (Wondafrash *et al.*, 2016). As with other members of the genus, *Pissodes* sp. often breeds on the bole of declining or recently dead trees, although it has also been observed attacking terminal leaders of young healthy trees (Gebeyehu & Wingfield, 2003). Published research on flight phenology and voltinism is lacking for this weevil in South Africa. Adults and larvae of the weevil are present throughout the year

however, suggesting multiple, overlapping generations per year. The weevil has also been shown to at least incidentally carry the pitch canker fungus *Fusarium circinatum* Nirenberg & O'Donnell (Coutinho *et al.*, 2007; Mitchell *et al.*, 2011), which causes a disease of major importance in South Africa and other parts of the world.

Sirex noctilio is native to Eurasia and North Africa and was first detected in South Africa in 1994 (Tribe, 1995). The woodwasp has become one of the most damaging insect pests of *Pinus* in South Africa and other Southern Hemisphere countries (Hurley *et al.*, 2007; Slippers *et al.*, 2015). Female *S. noctilio* drills into the sapwood and oviposits while also inoculating trees with spores of an obligate symbiotic fungus, *Amylostereum areolatum* (Chaillet ex Fries) Boidin, and a phytotoxic mucus/venom. The *A. areolatum* and the phytotoxic mucus act synergistically in order to suppress the resistance of infested trees and provide suitable hosts for the larvae (Gaut, 1969; Bordeaux *et al.*, 2014). *Amylostereum areolatum* serves as important source of nutrition for the larvae and is also involved in xylem digestion during larval development, probably by contributing cellulolytic enzymes and acting as an external gut (Thompson *et al.*, 2014). The woodwasp completes its larval development within the xylem of coniferous trees. Eggs are placed within the sapwood where larvae form tunnels and later pupate and emerge as adults. In general, the woodwasp develops in 1-3 years, but in most regions the majority of the emergence occurs after one year (Morgan, 1968). In South Africa, adults emerge from October to January within the summer rainfall area, including KwaZulu Natal. Most trees start declining and die by late summer (February). Flight period and tree decline phenology are approximately one month later in the winter rainfall areas of the Eastern and Western Cape (Ryan & Hurley, 2012). Extensive tree mortality due to *S. noctilio* occurs in parts of the invasive range, especially where trees were exposed to other biotic and abiotic stress factors (Neumann & Minko, 1981).

In South Africa, *Pissodes* sp. and *S. noctilio* are often observed co-occurring on trees. Trees attacked by *S. noctilio* are generally suitable hosts of *Pissodes* sp. since both the woodwasps and the weevils prefer stressed or declining trees (Gebeyehu & Wingfield, 2003; Ryan & Hurley, 2012). While preferences for the exact stage of decline are not currently known for either *S. noctilio* or *Pissodes* sp., there are some indications that the woodwasp attacks trees in earlier stages of decline relative to the weevil. Based on literature and field observations, the woodwasp appears to prefer moderately stressed trees that are alive and standing at the time of attack (Bordeaux & Dean, 2012; Ryan & Hurley, 2012). In contrast, the weevil larvae and pupae are primarily found in severely declining or recently dead trees, including downed trees (Gebeyehu & Wingfield, 2003; Wondafrash *et al.*, 2016). The degree to which *S. noctilio*'s attack behavior provides breeding habitat that would have otherwise been unavailable to *Pissodes* sp. is not known. Facilitation of *Pissodes* sp. by *S. noctilio* seems likely, however, given that woodwasps attack healthy trees that are later co-colonized by weevils as the trees decline and die. The presence of *Pissodes* sp. and *S. noctilio* in South Africa provides an opportunity to study patterns of co-occurrence and possible consequences of direct or indirect interactions therein.

Direct interaction through competition for resources is not expected between *S. noctilio* and *Pissodes* sp. since they utilize different parts/tissues of the tree. *Sirex noctilio* develops in the sapwood (Ryan & Hurley, 2012), while most members of the genus *Pissodes*, including the weevil in South Africa are limited to the inner bark and the phloem, and only excavate superficial galleries in the outer surface of the xylem. These species could interact indirectly, however, through the induction of tree defences (e.g., traumatic resin duct formation or change in resin quantity or quality), tree physiological state (e.g., xylem water content, tree vigour) and/or rate of tree decline/death with effects on nutritional quality or suitability of the resource.

For example, primary bark beetles exhaust resin via mass attack, which make trees more accessible to subsequent bark or phloem feeders (Ryan *et al.*, 2015).

The co-occurrence of *S. noctilio* and *Pissodes* sp. could influence the growth and survival of *A. areolatum*, the obligate fungal symbiont of *S. noctilio*. Previous studies have reported on possible outcomes of interactions between *A. areolatum* and ophiostomatoid fungi associated with bark- and wood-boring insects such as *P. nemorensis* (Hurley *et al.*, 2007; Ryan *et al.*, 2011; Ryan *et al.*, 2012; Yousuf *et al.*, 2014; Haavik *et al.*, 2015). However, the outcome of the interaction between *A. areolatum* and other insect-associated fungi could depend on their respective growth rates and phenologies as well as microhabitat, nutritive and microclimatic optima. Disruption of the growth and survival of *A. areolatum* could impact the development of *S. noctilio* with potential consequences for survival or fecundity.

Here, we examine the patterns and consequences of co-occurrence of the two forest insect pests, *Pissodes* sp. and *S. noctilio* in their invasive range in South Africa. The objectives of the study were to (1) determine the degree and pattern of co-occurrence of *S. noctilio* and *Pissodes* sp.; (2) assess positive or negative consequences of co-occurrence on body size and emergence; and (3) examine evidence for resource partitioning in co-colonized trees in the form of differential emergence as a function of bole height.

Materials and methods

Study sites and host plant

This study was conducted in pine plantations in the central South African province of Mpumalanga (Table 1). Plantations in Mpumalanga were selected because of the high incidence of *S. noctilio* and *Pissodes* sp. in this region in recent years. Six sites across three *Pinus patula* plantations (in Lothair, Rooihogte and Torburnlea) were selected based on their

history of higher *S. noctilio* and *Pissodes* sp. infestations in the previous years. The study was conducted in 16 to 17 year-old stands. Only trees with adult emergence holes and pupal chambers were included in the study as these are clear indicators of infestation and easier to identify. Standing trees infested with *Sirex* and/or *Pissodes* sp. including those attacked in previous years were included in the study. In cases where emergence holes were not visible on the lower bole, resin droplets were used as indicators of *Sirex* infestation higher up in the trees. Such trees were felled and closely checked for emergence holes. Considering older trees raises the possibility of measuring consecutive rather than concurrent colonization (i.e., that *Pissodes* sp. may have arrived after *S. noctilio* had left, or *vice-versa*). It is highly unlikely that *S. noctilio* could have arrived after *Pissodes* sp. had come and gone, given the former's strong preference for live trees (Bordeaux & Dean, 2012; Ryan & Hurley, 2012). There may be some cases where *Pissodes* sp. arrived too late to influence *S. noctilio* development, or in some cases even after woodwasps have emerged, however. Based on preference of *Pissodes* sp. for stressed or recently dead trees (Gebeyehu & Wingfield, 2003) and the rapid rate of decline of woodwasp-infested trees, serial utilization of the resource is likely to be rare.

Study on degree and pattern of co-occurrence

In each of the sites, two transects were sampled along two randomly selected rows of trees. *Pissodes* sp. and/or *S. noctilio*-infested trees were noted and each transect was continued until obtaining 100 infested trees. Here, infested trees include all infestation types (*Pissodes* sp. only (P), *Sirex noctilio* only (S), and *Sirex noctilio* and *Pissodes* sp. (S+P)). Diameter at Breast Height (DBH) of each of the infested trees was measured. Emergence holes for the two species are quite similar, so bark was removed with a drawknife to confirm species identity by assessing the presence or absence of *Pissodes* sp. pupal chambers, diagnostic for the insect (Fig. 1A & B). At the time of our assessment, most insects had already emerged from the tree,

Table 1. Study sites, GPS coordinates and infestation levels of *Sirex noctilio* and *Pissodes* sp. in *Pinus patula* plantations in Mpumalanga, March to May 2015.

Site	Latitude	Longitude	Age of trees (year)	<i>Sirex</i> infestation (%)	<i>Pissodes</i> infestation (%)
Rooihogte C3	S 26° 44.005'	E 30° 18.041'	16	12.2	11.6
Rooihogte N56	S 26° 38.031'	E 30° 20.046'	17	16.8	11.2
Lothair U27	S 26° 23.079'	E 30° 37.437'	16	13.1	11.1
Lothair W17	S 26° 25.521'	E 30° 37.628'	16	9.3	11.0
Lothair H2	S 26° 32.452'	E 30° 41.492'	16	7.7	6.9
Torburnlea T6a	S 26° 12.215'	E 30° 35.054'	17	14.2	11.0
Average				12.2 ± 1.4	10.5 ± 0.7




Figure 1. A. *Pissodes* sp. pupal chambers, B. *Sirex noctilio* emergence holes, C. Logs from mature *Pinus patula* trees infested with *Pissodes* sp. and/or *Sirex noctilio*, D. *Pissodes* sp. pupal chamber and *S. noctilio* emergence holes co-occurring on the same tree.

180x140mm (300 x 300 DPI)

with only a few cases where (typically dead) insects remained. In cases where pupal chambers and emergence holes were absent from the bottom section of dead trees and identification of infestation types were difficult, trees were felled and thoroughly examined for pupal chambers of *Pissodes* sp. (Fig. 1A) and/or emergence holes of *S. noctilio* (Fig. 1B).

Effect of co-occurrence on insect density and body size

Five trees per site of each infestation type (P, S and S+P) were randomly selected from the transects. In total, 90 trees (i.e. 30 per infestation type) were felled and each of them were divided into three equal sections (bottom, middle, top) based on the total tree height. Two logs per tree (80 cm each), one from the bottom and the other from the middle tree section, where symptoms of infestation were observed, were sampled (Fig. 1C). We selected only the bottom and middle tree sections as *Pissodes* sp. is rarely found in top sections (Wondafrash M, personal observation). Length and circumference of each log was measured for the purpose of standardizing estimates of insect density by wood volume. Bark was removed from logs with a drawknife in order to differentiate between *Pissodes* sp. and *S. noctilio* infestations. Pupal chambers of *Pissodes* sp. (Figs. 1A & D) and/or emergence holes of *S. noctilio* (Figs. 1B & D) were then counted with the objectives of comparing emergence densities and within-tree spatial distribution of each species in co-inhabited (S+P) versus *S. noctilio* (S) or *Pissodes* sp. (P) trees.

Emergence hole diameter correlates strongly with body size in *S. noctilio* (Madden, 1981) and was used as a proxy for body size in this study. We measured the diameters of 10 randomly selected *S. noctilio* emergence holes from each log sample collected from S and S+P trees. Prior literature on the correlation between emergence hole or pupal chamber size and body size is lacking for *Pissodes* sp. To establish the strength of this relationship, we determined the

Pearson correlation coefficient (r) between adult body size, the width and length of the pupal chamber, and the diameter of the adult emergence hole in a preliminary rearing study. Logs from *Pissodes*-infested trees were kept in emergence cages and frequently monitored for the emerging adults. Fifty adults were each stored in a separate vial in 96% ethanol and were given unique codes that matched them to each pupal chamber. In some cases, multiple adults emerged between checks of the emergence cage and could not be definitively matched to a single new emergence hole. These adults/holes were excluded from further analysis. Width and length of the corresponding pupal chambers, diameter of emergence holes were measured and measurements were made on five adult morphological characters (Williams & Langor, 2002) using STATA version 13 (StataCorp, 2013). All five characters were found to be strongly correlated with adult emergence hole diameter and the length and width of pupal chambers (r ranged from 0.73 to 0.97; see Table 2 for details and a list of characters assessed). Ultimately, pupal chamber width was selected as the sole proxy for adult size due to ease of measurement and high correlation coefficient (r ranged from 0.87 to 0.97). We then measured the width of 10 pupal chambers from each P and S+P logs only in the bottom tree section. Measurements of *Pissodes* sp. size were not obtained from the middle tree section due to limited number of pupal chambers, particularly in S+P trees.

Data analysis

Co-occurrence of *S. noctilio* and *Pissodes* sp. was compared to random using a Monte Carlo randomization procedure where expected frequencies of co-occurrence were calculated given the observed densities of *S. noctilio* and *Pissodes* sp. trees at each site. The Monte Carlo procedure was necessary as we did not collect data on uncolonized trees and therefore did not have a full contingency table to work with, in which case a Chi-squared or Fisher's exact test would have been appropriate. The model was run for 10,000 iterations per site and coded in R

Table 2. Correlation coefficient among five morphometric characters and diameter of emergence hole and width and length of pupal chamber of *Pissodes* sp. in South Africa.

	EL	EB	PL	PW	SL	EH	WPC	LPC
EL	1.000							
EB	0.973	1.000						
PL	0.904	0.926	1.000					
PW	0.911	0.945	0.803	1.000				
SL	0.884	0.847	0.771	0.770	1.000			
EH	0.945	0.912	0.843	0.860	0.822	1.000		
WPC	0.965	0.933	0.874	0.872	0.879	0.936	1.000	
LPC	0.861	0.822	0.807	0.731	0.811	0.818	0.872	1.000

^{EL} length of elytra along mid line, ^{EB} width of elytra at base, ^{PL} length of pronotum on midline, ^{PW} width of pronotum at widest point, ^{SL} length of snout between apex and eye margin at mid-height, ^{EH} adult emergence hole, ^{WPC} width of pupal chamber and ^{LPC} length of pupal chamber

(R Core Team, 2016). The density of each insect species in trees of different infestation types (infestation types being S and S+P for *Sirex*, and P and S+P for *Pissodes* sp.) was standardized and expressed as the number of insects per unit volume (insects dm⁻³) of the log. *Pissodes* sp. being a phloem feeder, phloem surface area estimates could be considered to be a more appropriate denominator in the estimation of weevil density. However, wood volume estimate was preferred to maintain consistency across the two species and make comparisons on their within-tree distribution. Further, the relevant comparisons (Fig. 3 & 4) are not across but rather within species (i.e., the co-occurring v. the occurring-singly comparison for *Sirex* and *Pissodes*). Additionally, scaling by surface area did not qualitatively influence our results within the range of tree sizes we considered (data not shown). Wood volume was calculated using the equation $V = \pi r^2 h$ assuming the boles as perfect cylinders, where V is wood volume, r is radius and h is length of the log. The effect of co-occurrence on density and adult body size of *S. noctilio* and *Pissodes* sp. was determined using a General Linear Models (GLM) approach using SAS software version 9.3 (SAS, 2011). We performed separate models using density and size of *S. noctilio* and *Pissodes* sp. as the response variables. Response variables were square-root transformed to meet model assumptions of normality and homogeneity of variance.

Analyses were made separately for the bottom and middle tree section as there were no interactions detected between tree section and infestation type. To examine changes in within-tree colonization patterns across trees with different densities of insects, we compared the slopes corresponding to the proportion of insects in the middle section (where a slope of 0.5 corresponds to equal distribution in the middle versus the bottom, < 0.5 to higher densities in the bottom, and > 0.5 to higher densities in the middle). The 95% confidence intervals (CIs) of linear regression slopes were calculated for all the infestation categories based on the density

of insects in middle versus middle plus bottom tree section. The confidence intervals were then checked if they encompass 0.5 or not.

Results

Degree and pattern of co-occurrence

The degree of *S. noctilio* and *Pissodes* sp. co-occurrence ranged from 55-80% depending on study site. Overall, *S. noctilio* and *Pissodes* sp. were found co-occurring on $68.0 \pm 3.8\%$ of the infested trees. The remaining $32.0 \pm 3.8\%$ of trees were infested by a single species, $21.8 \pm 4.6\%$ by *S. noctilio* and the remaining $10.2 \pm 3.2\%$ by *Pissodes* sp. This degree of co-occurrence was indistinguishable from random based on the relative frequencies of each species in each study site, with *p*-values ranging from 0.08 to 0.35 (Fig. 2). A mean DBH of 12.8 ± 0.3 cm was recorded from all infested trees. The DBH of S trees (11.9 ± 0.6 cm) was significantly smaller than S+P trees (13.8 ± 0.5 cm), while that of P trees (12.7 ± 0.5) was not different from either (based on Tukey's HSD, full model: $F_{2, 135} = 3.4, p = 0.04$).

Insect density

A total of 90 trees were felled (i.e. 180 logs – 90 from each of the bottom and middle tree section) to determine the density of *S. noctilio* and/or *Pissodes* sp. In total we counted 4,196 *S. noctilio* and 3,354 *Pissodes* sp. emergence holes across the bottom and middle sections of the 90 trees. On average 24.6 ± 3.7 and 45.3 ± 4.8 *S. noctilio* emergence holes and 38.5 ± 4.4 and 20.8 ± 4.4 *Pissodes* sp. emergence holes were counted per bottom versus middle tree sections.

Sirex noctilio density was significantly lower in S+P trees compared to the density in S trees both in bottom (Fig. 3A; $F_{1, 48} = 13.3, p = 0.0007$) and middle (Fig. 3A; $F_{1, 48} = 21.4, p < 0.0001$) tree section. This effect was considerable, with 63.2% and 57.1% fewer emergence holes in the bottom and middle tree sections, respectively. There was significant though




Figure 2. Observed and expected frequencies of *Sirex noctilio* and *Pissodes* sp. co-occurrence on *Pinus patula* trees in Mpumalanga, South Africa. The histograms indicate the expected number of trees that would host both *S. noctilio* and *Pissodes* sp. within the subset of 100 attacked trees (within each site) under the assumption of randomness. The vertical black lines are the observed count of S+P trees. P-value is the proportion of expected values (of 10000) that were lower than the observed. As all vertical lines fall near the middle of the random distribution, no divergence from random was detected.

229x177mm (300 x 300 DPI)

moderate site effect on *S. noctilio* density in the middle ($F_{5, 48} = 2.6; p = 0.03$), but not for bottom tree section ($F_{5, 48} = 0.9; p = 0.31$). No site \times infestation type interactions were observed in either log section.

The density of *Pissodes* sp. in S+P trees was significantly lower compared to its density in P trees in middle tree section (Fig. 3B; $F_{1, 48} = 18.6, p < 0.0001$). Density of *Pissodes* sp. in the middle section of S+P trees was 62.9% lower than the corresponding density in P trees. The density in bottom tree section was not affected significantly (Fig. 3B; $F_{1, 48} = 1.8, p = 0.18$). There was significant though moderate site effect on *Pissodes* sp. density in the bottom ($F_{5, 48} = 3.1, p = 0.02$), but not in middle tree section ($F_{5, 48} = 0.6, p = 0.71$). No site \times infestation type interactions were observed in either log section.

Insect body size

Body size of *Sirex noctilio* in S+P trees did not differ significantly from the size in S trees either in the bottom ($F_{1, 43} = 0.5, p = 0.47$) or the middle ($F_{1, 46} = 0.7, p = 0.39$) tree section. However, there was a general trend of larger *S. noctilio* in S+P trees compared to S trees. Size of *S. noctilio* in S+P trees was 13.8% and 13.5% greater in middle and bottom tree section, respectively, compared to the size in S trees. The effect of infestation type (S and S+P) on *S. noctilio* size remained consistent across sites both in the bottom (infestation type \times site $F_{5, 43} = 2.0, p = 0.09$) and middle (infestation type \times site $F_{5, 46} = 1.8, p = 0.13$) tree section. There was no significant difference in *Pissodes* sp. body size in S+P versus P trees ($F_{1, 43} = 0.3, p = 0.58$) in bottom tree section. The effect of infestation type on body size of *Pissodes* sp. remained consistent throughout the study sites (infestation type \times site $F_{5, 43} = 0.7, p = 0.61$).




Figure 3. Mean density of A. *Sirex noctilio* emergence holes and B. *Pissodes* sp. pupal chambers in bottom and middle tree sections of co-inhabited versus *S. noctilio* or *Pissodes* sp. only trees. Error bars indicate standard error and means and different letters indicate significant differences at $p \leq 0.05$. The means were not compared between the bottom and middle tree sections, as indicated by the faint vertical lines.

136x200mm (150 x 150 DPI)




Figure 4. Within tree spatial distribution of *S. noctilio* and *Pissodes* sp. in *Pinus patula* trees in South Africa. Distribution of *S. noctilio* in S+P trees (light blue line), *S. noctilio* in S trees (orange line), *Pissodes* sp. in S+P trees (grey line), *Pissodes* sp. in P trees (yellow line), Slope = 0.5 (green line) and Slope = 1.0 (black line). A slope of 0.5 corresponds to equal distribution of insects in the middle versus the bottom tree section, < 0.5 to higher densities in the bottom, > 0.5 to higher densities in the middle, and a slope of 1.0 to all insects in the middle. M – Middle, B – Bottom.

159x117mm (150 x 150 DPI)

Within-tree distribution of insects

The within tree spatial distribution of *Pissodes* sp. varied between S+P and P trees (Fig. 4). In P trees, *Pissodes* sp. was equally abundant in bottom versus middle tree section, relative to the total insect density in each section (slope = 0.50; CI [0.44, 0.55]), while in S+P trees, it was more abundant in bottom tree section than in the middle (slope = 0.24; CI [0.14, 0.34]). In contrast, the within tree spatial distribution of *S. noctilio* did not vary between S+P and S trees. *Sirex noctilio* was more abundant in middle tree section both in S+P (slope = 0.74; CI [0.66, 0.83]) and S trees (slope = 0.71; CI [0.65, 0.76]) compared to bottom tree section.

Discussion

This is the first study on the pattern of co-occurrence and potential interactions between *S. noctilio* and *Pissodes* sp. in their invasive range. Largely by virtue of their preference for the same host trees in similar physiological conditions, *S. noctilio* and *Pissodes* sp. co-occur together on over two-thirds ($68.0 \pm 3.8\%$) of dead trees. In this sense, it can be said that these species are strongly aggregated on similar resources. When comparing actual co-occurrence rates with expectations (i.e., under the simplifying assumption that tree preferences are equivalent), we found no evidence for a departure from a random pattern of co-utilization of trees within the region we assessed (mid-bole or below). In other words, given the total number of trees that were attacked by each species and assuming that their preferences for trees overlap completely, the predicted number of shared and unshared trees does not differ significantly from random chance. This does not preclude facilitation or antagonism, however, and it is possible that both are occurring at different spatiotemporal scales. This may seem counter-intuitive, but it is perfectly plausible to hypothesize that, for example, many or all of the wasp-killed trees with weevil emergence holes would have been unavailable to *Pissodes* sp. in the

absence of *S. noctilio* – in fact this seems highly likely. Thus, it stands to reason that populations of *Pissodes* sp. have likely been augmented (facilitated) by the presence of *S. noctilio* at a landscape scale, even if they are not preferentially selecting or avoiding *Sirex*-colonized trees within stands that contain suitable trees with and without the woodwasps.

Distinguishing patterns arising from positive or negative interactions between species from those that emerge as a consequence of sometimes subtle variation in preference or suitability of host trees is not a trivial task and would require manipulative experiments, which are highly challenging in this system. However, examining within-tree spatial pattern for evidence of resource partitioning can shed some light on their existence, direction, or strength. Interestingly, the density and the within-tree spatial distribution of *S. noctilio* and *Pissodes* sp. in S+P trees were significantly different from S or P trees respectively. Vertical colonization patterns for *Pissodes* sp. appeared to shift lower in the tree in the presence of *S. noctilio* when total insect density per section was accounted for (Fig. 4), while 70-75% of *S. noctilio* colonization was in the mid-bole, irrespective of co-occurrence with *Pissodes* sp. This is an indication of resource partitioning where the within-tree distribution of *Pissodes* sp. is influenced by the occurrence of *S. noctilio*.

Our results on body size contrast with results from Ryan *et al.* (2012) who reported larger body size of *S. noctilio* (particularly for males) in beetle-infested trees in Canada, whereas we found no impacts of *Pissodes* sp. on *S. noctilio* body size. Ryan *et al.* (2012) accounted this to the possible accelerated drying of sapwood in the presence of Ophiostomatoid fungi (fungal associates of many bark beetles and other wood-inhabiting insects; Chow & Obermajer, 2007) and faster growth of *A. areolatum* in drier wood (Coutts & Dolezal, 1965) that might have ultimately improved larval nutrition in beetle-positive trees. The incongruence between this

study and Ryan *et al.* (2012) could be due to differences in the identity and diversity of insect species co-occurring with *S. noctilio* in Canada and South Africa. In South Africa, *Pissodes* sp. is the most frequently co-occurring species with *S. noctilio*. *Orthotomicus erosus*, *Hylastes angustatus* and *Hylurges ligniperda* are also introduced pine-feeding insects (Garnas *et al.*, 2016), but are rare in mature *P. patula* plantations and were not detected in this study. However, in Canada, Ryan *et al.* (2012) reported that five subcortical beetles species (*Tomicus piniperda*, *Ips grandicollis*, *Gnathotricus materiarius*, *Monochamus carolinensis* and *P. nemorensis*) commonly co-occur with *S. noctilio*.

The invasion of *S. noctilio* in South Africa has significantly increased *Pinus* tree mortality, particularly under outbreak conditions (Hurley *et al.*, 2007; Hurley *et al.*, 2012). Ninety percent of dead trees included in the present study were found infested with *S. noctilio*. The majority (> 75%) of these *S. noctilio* infested trees were found co-inhabited by *Pissodes* sp., suggesting a high degree of overlap in preference and/or performance of *Pissodes* sp. on declining trees in a similar physiological condition as utilized by *S. noctilio*. This could also be a result of facilitation of one insect species by the other. Preference of *Pissodes* sp. and *S. noctilio* for declining/recently dead trees is well documented (Hopkins, 1911; Gebeyehu & Wingfield, 2003; Wermelinger & Thomsen, 2012), though the precise physiological state of trees preferred for oviposition or that are optimal for larval development may differ among insects specializing on declining or stressed trees. In line with this, mixed (negative or neutral) co-colonization patterns were observed between *S. noctilio* and bark and wood boring beetles at tree and stand level in North America, which the authors attributed to differences in tree preferences by different insect species (Foelker *et al.*, 2018). Interestingly, *S. noctilio* co-occurred non-randomly with *Tomicus* sp. and with cerambycids in Spanish pine, though not with *Pissodes* sp. (Ayres *et al.*, 2014).

While patterns of co-occurrence cannot strictly be interpreted as proof of competition or facilitation, it seems likely that *S. noctilio* in South Africa has increased availability of suitable trees for *Pissodes* sp. by attacking and weakening trees that would otherwise be too vigorous and therefore unavailable to the weevil. This is based on other evidence such as apparently lower densities of *Pissodes* sp. observed in areas where the density of *S. noctilio* is low and where far fewer dead and dying trees are available. This feedback is likely complex and variable in time and space. For example, in sites/years where *S. noctilio* is at "endemic" or low levels and is principally utilizing stressed and declining trees that would have been available to *Pissodes* sp. anyway, the direction of its effect on *Pissodes* sp. may be slightly negative due to competition for resources. At higher densities however, *S. noctilio* is capable of killing healthier trees (Spradbery, 1973; Ryan & Hurley, 2012; Lantschner & Corley, 2015) which almost certainly creates additional habitat for *Pissodes* sp. This would imply that while the influence of *S. noctilio* on *Pissodes* sp. populations is negative at the tree scale, the influence at the landscape level could be positive or negative depending on population or density of the woodwasp. Overall, we suspect that there may be a net positive effect of *S. noctilio* on *Pissodes* sp., though more research is needed to explore this question further. Based on the current study, it appears that the effect of *Pissodes* sp. on *S. noctilio* may be negligible in the bole. That said, in pine plantations where *Pissodes* sp. feeds on terminal leaders of young trees resulting in double leaders, this could represent a positive feedback if these double leader trees are more susceptible to *S. noctilio*.

The lower density of *S. noctilio* and *Pissodes* sp. in the sampled sections of S+P trees compared to the density in the sampled sections of S or P trees is a potential indicator of negative interactions. This pattern could emerge from reduced preference for attacked trees by *S. noctilio* and *Pissodes* sp., though this seems unlikely for *S. noctilio* given the observation that wasps

typically arrive first. Alternatively, larval mortality, perhaps as a function of excessive bole drying or poor performance of fungal symbiont of *S. noctilio*, could be elevated in the presence of *Pissodes* sp. Previous studies reported potential interactions between *S. noctilio* and the co-occurring community of insects, where co-occurring insects and their associated fungi in North America were suggested as possible factors that regulate *S. noctilio* populations (Ryan *et al.*, 2012; Haavik *et al.*, 2015). These studies were conducted in the Northern Hemisphere where *Pinus* species are native and support a diverse, potentially co-evolved community of insects, including numerous other subcortical phytophagous insects (wood borers and bark beetles) (Ryan *et al.*, 2012; Haavik *et al.*, 2015). Though body size appears unaffected, *S. noctilio* density in South Africa does appear to respond in a similar way to *Pissodes* sp. as compared with co-occurring beetles in the Northern Hemisphere.

Differences in the within-tree spatial distribution of *Pissodes* sp. in S+P versus P trees is another potential indicator of interactions between *S. noctilio* and *Pissodes* sp. A higher proportion of *Pissodes* sp. were present in the bottom tree section of S+P trees as compared to P trees, an indication that *Pissodes* sp. shifts away from the middle tree section (*S. noctilio*'s preferred habitat) (Morgan & Stewart, 1966; Neumann *et al.*, 1982; Ryan *et al.*, 2012) when co-occurring with *S. noctilio*. This suggests partitioning of a common resource among these two co-occurring insects. Previous studies have suggested partitioning of a common resource as a mechanism of minimizing interspecific competition among bark beetle species (Paine *et al.*, 1981; Grunwald, 1986; Flamm *et al.*, 1987). For example, avoidance of oviposition site was observed among four co-occurring insect species on grapevine, namely the European grapevine moth *Lobesia botrana*, the grape berry moth *Eupoecilia ambiguella*, the codling moth *Cydia pomonella* and the oriental fruit moth *C. molesta* (Thiéry & Gabel, 1993). In contrast to *Pissodes* sp., the within-tree spatial distribution of *S. noctilio* was not influenced by

the presence of *Pissodes* sp. The majority of *S. noctilio* emergence holes were in the middle tree section, both in S+P trees and S trees. This is consistent with previous reports in Canada (Ryan *et al.*, 2012), South Africa (Hurley *et al.*, 2008), and New Zealand, Australia and Brazil (Morgan & Stewart, 1966; Neumann *et al.*, 1982; Ryan *et al.*, 2012).

Studies on patterns of co-occurrence of invasive species can shed light on the nature and potential interaction among them, which can drive dynamics and which can also change over time as environments or management regimes change, or as new species arrive. In the present study, we detected high levels of co-occurrence of *S. noctilio* and *Pissodes* sp. and potential indications of both positive (landscape scale) and negative (tree scale) interactions between the two species in their invasive range in South Africa. These interactions will need to be confirmed through manipulative experiments, but if present could have potential implications for the management of these species.

References

- Ayres, M. P., Pena, R., Lombardo, J. A. & Lombardero, M. J. (2014) Host use patterns by the European woodwasp, *Sirex noctilio*, in its native and invaded range. *PLoS One*, **9**, e90321.
- Barbosa, P. & Wagner, M.R. (1988) Introduction to forest and shade tree insects. Academic Press, Inc., San Diego, USA.
- Bordeaux, J.M. & Dean, J.F.D. (2012) Susceptibility and response of pines to *Sirex noctilio*. *The Sirex woodwasp and its fungal symbiont* (ed. by B. Slippers, P. de Groot & M.J. Wingfield). Springer, New York, pp. 31-50.
- Bordeaux, J.M., Lorenz, W.W., Johnson, D., Badgett, M.J., Glushka, J., Orlando, R. & Dean, J.F. (2014) Noctilisin, a venom glycopeptide of *Sirex noctilio* (Hymenoptera:

- Siricidae), causes needle wilt and defense gene responses in pines. *Journal of Economic Entomology*, **107**, 1931-1945.
- Chow, S. & Obermajer, A. (2007) Moisture and blue stain distribution in mountain pine beetle infested lodgepole pine trees and industrial implications. *Wood Science Technology*, **41**, 3–16.
- Coutinho, T.A., Steenkamp, E.T., Mongwaketsi, K., Wilmot, M. & Wingfield, M.J. (2007) First outbreak of pitch canker in a South African pine plantation. *Australasian Plant Pathology*, **36**, 256-261.
- Coutts, M.P. & Dolezal, J.E. (1969) Emplacement of fungal spores by the woodwasp, *Sirex noctilio*, during oviposition. *Forest Science*, **4**, 412–416.
- Denno, R.F., McClure, M.S. & Ott, J.R. (1995) Interspecific interaction in phytophagous insects: competition re-examined and resurrected. *Annual Review of Entomology*, **40**, 297-331.
- Dewey, J.E., Ciesla, W.M. & Meyer, H.E. (1974) Insect defoliation as a predisposing agent to a bark beetle outbreak in eastern Montana. *Environmental Entomology*, **3**, 722-722.
- EPPO (1999) European and Mediterranean Plant Protection Organization (EPPO) Reporting Service, No. 4, Paris, France. <http://archives.eppo.int/EPPOreporting/1999/Rse-9904.pdf> [accessed on 10 July 2016].
- Flamm, R.O., Wagner, T.L., Cook, S.P., Pulley, P.E., Coulson, R.N. & Mcardle, T.M. (1987) Host colonization by cohabiting *Dendroctonus frontalis*, *Ips avulsus*, and *I. calligraphus* (Coleoptera: Scolytidae). *Environmental Entomology*, **16**, 390-399.
- Foelker, C.J., Parry, D. & Fierke, M.K. (2018) Biotic resistance and the spatiotemporal distribution of an invading woodwasp, *Sirex noctilio*. *Biological Invasions*, **20**: 1991-2003.

- Garnas, J., Roux, J., Hurley, B.P., Slippers, B., & Wingfield, M.J. (2016) Insects and diseases of Mediterranean forests: A South African perspective. *Insects and Diseases of Mediterranean Forest Systems* (ed. by T.D. Paine and F. Lieutier), pp. 397-430, Springer International Publishing, Switzerland.
- Gaut, I.P. (1969) Identity of the fungal symbiont of *Sirex noctilio*. *Australian Journal of Biological Sciences*, **22**, 905-914.
- Gebeyehu, S. & Wingfield, M.J. (2003) Pine weevil *Pissodes nemorensis*: threat to South African pine plantations and options for control: review article. *South African Journal of Science*, **99**, 531-536.
- Grunwald, M. (1986) Ecological segregation of bark beetles (Coleoptera, Scolytidae) of spruce. *Journal of Applied Entomology*, **101**, 176-187.
- Haavik, L.J., Dodds, K.J. & Allison, J.D. (2015) Do native insects and associated fungi limit non-native woodwasp, *Sirex noctilio*, survival in a newly invaded environment? *PLoS ONE*, **10**, e0138516.
- Hajek, A.E., Hurley, B.P., Kenis, M., Garnas, J.R., Bush, S.J., Wingfield, M.J., van Lenteren, J.C. & Cock, M.J.W. (2016) Exotic biological control agents: A solution or contribution to arthropod invasions? *Biological Invasions*, **18**, 953-969.
- Hawkins, B.A. (2005) *Pattern and process in host-parasitoid interactions*. Cambridge University Press, Cambridge, pp. 151-168.
- Heimpel, G.E., Frelich, L.E., Landis, D.A., Hopper, K.R., Hoelmer, K.A., Sezen, Z., Asplen, M.K. & Wu, K. (2010) European buckthorn and Asian soybean aphid as components of an extensive invasional meltdown in North America. *Biological Invasions*, **12**, 2913-2931.

- Hofstetter, R.W., Cronin, J.T., Klepzig, K.D., Moser J.C. & Ayres, M.P. (2006). Antagonisms, mutualisms and commensalisms affect outbreak dynamics of the southern pine beetle. *Oecologia* **147**: 679 - 691.
- Hopkins, A.D. (1911) Contributions toward a monograph of the bark weevils of the genus *Pissodes*. Technical series, No. 17, Part 1, US Department of Agriculture. Washington, DC, USA.
- Hurley, B.P., Croft, P., Verleur, M., Wingfield, M.J. & Slippers, B. (2012) The control of the *Sirex* woodwasp in diverse environments: the South African experience. *The Sirex woodwasp and its fungal symbiont* (ed. by B. Slippers, P. de Groot and M.J. Wingfield). Springer, New York, pp. 247-264.
- Hurley, B.P., Slippers, B., Croft, P.K., Hatting, H.J., van der Linde, M., Morris, A.R., Dyer, C. & Wingfield, M.J. (2008) Factors influencing parasitism of *Sirex noctilio* (Hymenoptera: Siricidae) by the nematode *Deladenus siricidicola* (Nematoda: Neotylenchidae) in summer rainfall areas of South Africa. *Biological Control*, **45**, 450-459.
- Hurley, B.P., Slippers, B. & Wingfield, M.J. (2007) A comparison of control results for the alien invasive woodwasp, *Sirex noctilio*, in the Southern Hemisphere. *Agricultural and Forest Entomology*, **9**, 159-171.
- Klooster, W.S., Gandhi, K.J.K, Long, L.C., Perry, K.I., Rice, K.B. & Herms, D.A. (2018) Ecological Impacts of Emerald Ash Borer in Forests at the Epicenter of the Invasion in North America. *Forests*, **9**: 250, doi:10.3390/f9050250
- Lantschner, M.V. & Corley, J.C. (2015) Spatial pattern of attacks of the invasive woodwasp *Sirex noctilio*, at landscape and stand scales. *PLoS ONE*, **10**, e0127099.

- Mack, R.N., Simberloff, D., Lonsdale, M.W., Evans, H., Clout, M. & Bazzaz, F.A. (2000) Biotic invasions: causes, epidemiology, global consequences, and control. *Ecological Applications*, **10**, 689-710.
- Madden, J.L. (1981) Egg and larval development in the woodwasp, *Sirex noctilio* F. *Australian Journal of Zoology*, **29**, 493-506.
- McClure, M.S. (1981) Effects of voltinism, interspecific competition and parasitism on the population dynamics of the hemlock scales, *Fiorinia externa* and *Tsugaspidotus tsugae* (Homoptera, Diaspididae). *Ecological Entomology*, **6**, 47-54.
- Mitchell, R.G., Steenkamp, E.T., Coutinho, T.A. & Wingfield, M.J. (2011) The pitch canker fungus, *Fusarium circinatum*: implications for South African forestry. *Southern Forests: a Journal of Forest Science*, **73**, 1-13.
- Morgan, F.D. (1968) Bionomics of Siricidae. *Annual Review of Entomology*, **13**: 239 – 256.
- Morgan, F.D. & Stewart, N.C. (1966) The biology and behaviour of the wood-wasp *Sirex noctilio* F. in New Zealand. *Transactions and Proceedings of the Royal Society of New Zealand*, **7**, 195-204.
- Neumann, F., Harris, J., Kassaby, F. & Minko, G. (1982) An improved technique for early detection and control of the *Sirex* woodwasp in radiata pine plantations. *Australian Forestry*, **45**, 117-124.
- Neumann, F. & Minko, G. (1981) The *sirex* wood wasp in Australian radiata pine plantations. *Australian Forestry*, **44**, 46-63.
- Orlova- Bienkowskaja, M.J. (2015). Cascading ecological effects caused by the establishment of the emerald ash borer *Agrilus planipennis* (Coleoptera: Buprestidae) in European Russia. *European Journal of Entomology*, **112**: 778 – 789.

- Paine, T.D., Birch, M.C. & Švihra, P. (1981) Niche breadth and resource partitioning by four sympatric species of bark beetles (Coleoptera: Scolytidae). *Oecologia*, **48**, 1-6.
- Preisser, E.L. & Elkinton, J.S. (2008) Exploitative competition between invasive herbivores benefits a native host plant. *Ecology*, **89**, 2671-2677.
- Preisser, E.L., Lodge, A.G., Orwig, D.A. & Elkinton, J.S. (2008) Range expansion and population dynamics of co-occurring invasive herbivores. *Biological Invasions*, **10**, 201-213.
- R (2016) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/> (Version 3.3.2, downloaded on 31 October 2017).
- Rodriguez, L.F. (2006) Can invasive species facilitate native species? Evidence of how, when, and why these impacts occur. *Biological Invasions*, **8**: 927 – 939.
- Ryan, K., de Groot, P. & Smith, S.M. (2012) Evidence of interaction between *Sirex noctilio* and other species inhabiting the bole of *Pinus*. *Agricultural and Forest Entomology*, **14**, 187-195.
- Ryan, K. & Hurley, B.P. (2012) Life history and biology of *Sirex noctilio*. *The Sirex woodwasp and its fungal symbiont* (ed. by B. Slippers, P. de Groot & M.J. Wingfield). Springer, New York, pp. 15-30.
- Ryan, K., Moncalvo, J-M., de Groot, P. & Smith, S.M. (2011) Interactions between the fungal symbiont of *Sirex noctilio* (Hymenoptera: Siricidae) and two bark beetle-vectored fungi. *The Canadian Entomologist*, **143**, 224-235.
- Ryan, M.G., Sapes, G., Sala, A. & Hood, S.M. (2015) Tree physiology and bark beetles. *New Phytologist*, **205**, 955-957.

- Sala, O.E., Chapin, F.S., Armesto, J.J., Berlow, E., Bloomfield, J., Dirzo, R., Huber-Sanwald, E., Huenneke, L.F., Jackson, R.B. & Kinzig, A. (2000) Global biodiversity scenarios for the year 2100. *Science*, **287**, 1770-1774.
- SAS Institute Inc. (2011) SAS[®] 9.3 System Options: Reference, Second Edition. Cary, NC: SAS Institute Inc.
- Schaffner, F., Medlock, J.M. & Bortel, W.V. (2013) Public health significance of invasive mosquitoes in Europe. *Clinical Microbiology and Infection*, **19**, 685-692.
- Simberloff, D. (2011) How common are invasion-induced ecosystem impacts? *Biological Invasions*, **13**, 1255-1268.
- Slippers, B., Hurley, B.P. & Wingfield, M.J. (2015) Sirex woodwasp: A model for evolving management paradigms of invasive forest pests. *Annual Review of Entomology*, **60**, 601-619.
- Smith, G.D., Carroll, A.L. & Lindgren, B.S. (2011) Facilitation in bark beetles: endemic mountain pine beetle gets a helping hand. *Agricultural and Forest Entomology*, **13**: 37-43.
- Spradbery, J.P. (1973) A comparative study of the phytotoxic effects of siricid woodwasps on conifers. *Annals of Applied Biology*, **75**, 309-320.
- StataCorp (2013) Stata: Release 13. Statistical Software. Stata Press, Texas, USA.
- Stein, B.A., Kutner, L.S. & Adams, J.S. (2000) Precious heritage: The status of biodiversity in the United States. Oxford University Press, New York.
- Thiéry, D. & Gabel, B. (1993) Inter-specific avoidance of egg-associated semiochemicals in four tortricids. *Experientia*, **49**, 998-1001.

- Thompson, B.M., Bodart, J., McEwen, C. & Gruner, D.S. (2014) Adaptations for symbiont-mediated external digestion in *Sirex noctilio* (Hymenoptera: Siricidae). *Annals of the Entomological Society of America*, **107**, 453e460.
- Tribe, G.D. (1995) The woodwasp *Sirex noctilio* Fabricius (Hymenoptera: Siricidae), a pest of *Pinus* species, now established in South Africa. *African Entomology*, **3**, 215-217.
- Webb, V.D. (1974) Forest and timber entomology in the Republic of South Africa. Entomology Memoir 34, Department of Agricultural Technical Services, Pretoria, South Africa.
- Wermelinger, B. & Thomsen, I.M. (2012) The woodwasp *Sirex noctilio* and its associated fungus *Amylostereum areolatum* in Europe. *The Sirex woodwasp and its fungal symbiont* (ed. by B. Slippers, P. de Groot & M.J. Wingfield). Springer, New York, pp. 65-80.
- Williams, D.J.M. & Langor, D.W. (2002) Morphometric study of the *Pissodes strobi* complex (Coleoptera: Curculionidae). *The Canadian Entomologist*, **134**, 447-466.
- Wingfield, M.J., Garnas, J.R., Hajek, A., Hurley, B.P, de Beer, Z.W. & Taerum, S.J. (2016) Novel and co-evolved associations between insects and microorganisms as drivers of forest pestilence. *Biological Invasions*, **18**, 1045-1056.
- Wondafrash, M., Slippers, B., Garnas, J., Roux, G., Foit, J., Langor, D.W. & Hurley, B.P. (2016) Identification and genetic diversity of two invasive *Pissodes* spp. Germar (Coleoptera: Curculionidae) in their introduced range in the southern hemisphere. *Biological Invasions*, **18**, 2283-2297.
- Yousuf, F., Gurr, G.M., Carnegie, A.J., Bedding, R.A., Bashford, R., Gitau, C.W. & Nicol, H.I. (2014) The bark beetle, *Ips grandicollis*, disrupts biological control of the woodwasp, *Sirex noctilio*, via fungal symbiont interactions. *FEMS Microbiology Ecology*, **88**, 38-47.

Acknowledgments

We thank members of Tree Protection Co-operative Program (TPCP), DST-National Research Foundation (NRF) and the University of Pretoria, South Africa for the financial support. Our special thanks go to Sandile Nkosi, manager at Lothair SAPPI plantation, Eddie Ferrera, manager at Rooihogte SAPPI plantation, and TPCP students, for their assistance in the field.