Supplementary material

Title: Pharmacological evaluation of hydro-ethanol and hot water leaf extracts of *Bauhinia* galpinii (Fabaceae): A South African ethnomedicinal plant

Authors: J.O. Erhabor^{a, b}, A.G. Omokhua^a, M. Ondua^a, M. Ali Abdalla^{a, c} and L.J. McGaw^{a*}

^aPhytomedicine Programme, Department of Paraclinical Sciences, Faculty of Veterinary Science, University of Pretoria, Private Bag X04, Onderstepoort 0110, South Africa.

^bPhytomedicine Unit, Department of Plant Biology and Biotechnology, University of Benin, PMB 154, Benin City, Nigeria.

^cDepartment of Food Science and Technology, Faculty of Agriculture, University of Khartoum, 13314, Khartoum North, Sudan.

Supplementary table 1

Proposed Minimum inhibitory concentrations (MICs) by previous Authors for plant extracts

Authors	Proposed MICs (mg/ml)					
	Excellent	Very	Significant/strong/	moderate	Weak	Inactive
		good	good			
Holetz et			0.1	0.1 - 0.5	0.5 - 1	> 1
al., 2002						
Eloff, 2004				0.1		
Rios and			< 0.1			> 1
Recio,						
2005						
Cos et al.,			< 0.1 (IC ₅₀)			
2006						
van				≤ 0.2		
Vuuren,						
2008						
Kuete,			< 0.1	> 0.1 - ≤ 0.625	> 0.625	
2010						
Omokhua	0.02 - 0.04	0.08	0.16			
et al., 2018						