

Supplemental files

Selective cytotoxic activity of isolated compounds from *Globimetula dinklagei* and *Phragmanthera capitata* (Loranthaceae) against human adenocarcinoma cell lines.

Emmanuel Mfotie Njoya^{1,2*}, Hermine L. D. Maza³, Pierre Mkounga³, Ulrich Koert⁴, Augustin E. Nkengfack³, Lyndy J. McGaw¹




Figure S1: FAB- MS spectra of compound 1




Figure S2: ^{13}C NMR spectra (CDCl_3 , 125 MHz) of compound 1




Figure S3: DEPT NMR spectra (CDCl_3 , 125 MHz) of compound 1




Figure S4: ^1H NMR spectra (CDCl_3 , 500 MHz) of compound 1




Figure S5: HMBC spectra (CDCl_3 , ^1H : 500 MHz ; ^{13}C : 125 MHz) of compound 1




Figure S6: ^{13}C NMR spectra (CDCl_3 , 125 MHz) of compound 2




Figure S7: DEPT NMR spectra (CDCl₃, 125 MHz) of compound 2




Figure S8: ^1H NMR spectra (CDCl_3 , 500 MHz) of compound 2




Figure S9: HMBC spectra 1Spectre (CDCl_3 , ^1H : 500 MHz ; ^{13}C : 125 MHz) of compound 2




Figure S10: ¹³C NMR spectra (CDCl₃, 125 MHz) of compound 3




Figure S11: DEPT NMR spectra (CDCl₃, 125 MHz) of compound 3




Figure S12: ^1H NMR spectra (CDCl_3 , 500 MHz) of compound 3




Figure S13: HMBC spectra (CDCl_3 , ^1H :500 MHz ; ^{13}C : 125 MHz) of compound 3




Figure S14: ¹³C NMR spectra (DMSO, 125 MHz) of compound 4




Figure S15: ¹H NMR spectra (DMSO, 500 MHz) of compound 4




Figure S16: ¹³C NMR spectra (C₅D₅N, 125 MHz) of compound 5




Figure S17: ^1H NMR spectra ($\text{C}_5\text{D}_5\text{N}$, 500 MHz) of compound 5




Figure S18: ${}^1\text{H}$ NMR spectra (DMSO, 300 MHz) of compound 6




Figure S19: HMBC spectra (DMSO, ¹H: 500 MHz ; ¹³C : 75 MHz) of compound 6