

Conservation implications of fine scale population genetic structure of *Ficus* species in South African forests

Jun-Yin Deng, Simon van Noort, Stephen G. Compton, Yan Chen, Jaco M. Greeff

Table S1 The defined distance intervals for every species at each sample site.

Species	Sample site		Distance class															
			1	2	3	4	5	6	7	8	9	10	11	12	13	14		
<i>F. bizanae</i>	MKA	Distance (m)	20	40	60	80	100	150	300									
		No. pairs	245	278	220	217	111	189	171									
	MBO	Distance (m)	40	100	4000	4500	8000	8500	9500									
		No. pairs	82	37	45	32	178	42	80									
<i>F. craterostoma</i>	DWE	Distance (m)	50	100	200	300	400	500	600	700	800	900	1000	1200	1400	1600		
		No. pairs	117	76	72	108	33	33	71	50	109	43	86	154	69	107		
	MAN	Distance (m)	100	200	300	400	500	600	800	1000	1600							
		No. pairs	40	38	40	36	56	31	68	65	61							
	ING (all)	Distance (m)	100	200	300	400	800	900	1000	1500	2000	2500	4000	4600				
		No. pairs	570	374	365	262	199	250	224	1421	793	639	739	159				
	ING (adult)	Distance (m)	100	200	300	400	800	900	1000	1500	2000	2500	4000	4600				
		No. pairs	272	132	184	140	111	140	83	645	368	228	203	50				
	ING (young)	Distance (m)	100	200	400	1500	2000	2500	4600									
		No. pairs	55	49	49	210	70	105	165									
	<i>F. sur</i>	MBO	Distance (m)	100	200	400	1000	2000	4000	8000	10000	13000						
			No. pairs	269	244	106	66	200	367	185	54	49						
MAN		Distance (m)	200	400	600	800	1000	1500	3000	6200								
		No. pairs	122	131	156	206	164	386	269	162								