

Supplementary Material

***Cyclopia intermedia* (honeybush) induces uncoupling protein 1 and peroxisome proliferator-activated receptor alpha expression in obese diabetic female db/db mice**

Babalwa Unice Jack ^{1,*}, Pritika Ramharack ^{1,2}, Christiaan Malherbe ^{3,†}, Kwazi Gabuza ¹, Elizabeth Joubert ^{3,4} and Carmen Pheiffer ^{1,5,6}

¹ Biomedical Research and Innovation Platform, South African Medical Research Council, Tygerberg, Cape Town 7505, South Africa

² Pharmaceutical Sciences, School of Health Sciences, University of KwaZulu-Natal, Westville Campus, Durban 4001, South Africa

³ Plant Bioactives Group, Post-Harvest and Agro-Processing Technologies, Agricultural Research Council (ARC), Infruitec-Nietvoorbij, Stellenbosch 7599, South Africa

⁴ Department of Food Science, University of Stellenbosch, Matieland, Stellenbosch 7602, South Africa

⁵ Centre for Cardio-Metabolic Research in Africa (CARMA), Division of Medical Physiology, Faculty of Medicine and Health Sciences, University of Stellenbosch, Tygerberg, Cape Town 7505, South Africa

⁶ Department of Obstetrics and Gynaecology, Faculty of Health Sciences, University of Pretoria, Pretoria 0001, South Africa

* Correspondence: babalwa.jack@mrc.ac.za; Tel.: +27-219-380336

† Current address: Analytical Services, Afriplex Pty Ltd, Paarl 7620, South Africa

Validation of molecular docking of PPAR α




Figure S1: Validation of molecular docking of PPAR α by superimposing the complexed models with the crystal template structures. All ligands docked at the same active site to the complexed crystal structures (Blue ribbon: 3DZY- ball and stick ligand, Green: 1K7L- ball and stick ligand, Tan: PPAR α model).