

Using FIGO Nutrition Checklist counselling in pregnancy: A review to support healthcare professionals

Sarah Louise Killeen¹ | Niamh Donnellan¹ | Sharleen L. O'Reilly^{1,2} | Mark A. Hanson^{3,4} |
Mary L. Rosser⁵ | Virna P. Medina⁶ | Chandni Maria Jacob^{3,4} | Hema Divakar⁷ |
Moshe Hod^{8,9} | Liona C. Poon¹⁰ | Lina Bergman^{11,12,13} | Patrick O'Brien¹⁴ |
Anil Kapur¹⁵ | Bo Jacobsson^{16,17} | Cynthia V. Maxwell¹⁸ | Harold David McIntyre¹⁹ |
Lesley Regan²⁰ | Esraa Aljurjia^{21,22} | Ronald C. Ma^{23,24} | Sumaiya Adam^{25,26} |
Fionnuala M. McAuliffe¹ | on behalf of the FIGO Committee on Impact of Pregnancy on
Long-term Health and the FIGO Division of Maternal and Newborn Health

¹UCD Perinatal Research Centre, School of Medicine, University College Dublin, National Maternity Hospital, Dublin, Ireland

²School of Agriculture and Food Science, University College Dublin, Dublin, Ireland

³Institute of Developmental Sciences, University Hospital Southampton, Southampton, UK

⁴NIHR Southampton Biomedical Research Centre, University of Southampton, Southampton, UK

⁵Department of Obstetrics and Gynecology, Columbia University Irving Medical Center, New York, New York, USA

⁶Department of Obstetrics and Gynecology, Faculty of Health, Universidad del Valle, Clínica Imbanaco Quirón Salud, Universidad Libre, Cali, Colombia

⁷Divakar's Specialty Hospital, Bengaluru, India

⁸Helen Schneider Hospital for Women, Rabin Medical Center, Petah Tikva, Israel

⁹Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel

¹⁰Department of Obstetrics and Gynecology, Prince of Wales Hospital, The Chinese University of Hong Kong, Hong Kong, Hong Kong SAR, China

¹¹Department of Obstetrics and Gynecology, Institute of Clinical Sciences, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden

¹²Department of Obstetrics and Gynecology, Stellenbosch University, Cape Town, South Africa

¹³Department of Women's and Children's Health, Uppsala University, Uppsala, Sweden

¹⁴Institute for Women's Health, University College London, London, UK

¹⁵World Diabetes Foundation, Bagsvaerd, Denmark

¹⁶Department of Obstetrics and Gynecology, Sahlgrenska University Hospital, Gothenburg, Sweden

¹⁷Department of Genetics and Bioinformatics, Domain of Health Data and Digitalization, Institute of Public Health, Oslo, Norway

¹⁸Maternal Fetal Medicine, Sinai Health and Women's College Hospital, University of Toronto, Toronto, Canada

¹⁹Mater Health, University of Queensland, Mater Health Campus, South Brisbane, Queensland, Australia

²⁰Imperial College London, London, UK

²¹The World Association of Trainees in Obstetrics & Gynecology, Paris, France

²²Elwya Maternity Hospital, Baghdad, Iraq

²³Department of Medicine and Therapeutics, The Chinese University of Hong Kong, Hong Kong, Hong Kong SAR, China

²⁴Hong Kong Institute of Diabetes and Obesity, The Chinese University of Hong Kong, Hong Kong, Hong Kong SAR, China

²⁵Department of Obstetrics and Gynecology, School of Medicine, Faculty of Health Sciences, University of Pretoria, Pretoria, South Africa

²⁶Diabetes Research Centre, Faculty of Health Sciences, University of Pretoria, Pretoria, South Africa

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *International Journal of Gynecology & Obstetrics* published by John Wiley & Sons Ltd on behalf of International Federation of Gynecology and Obstetrics.

Correspondence

Sarah Louise Killeen, UCD Perinatal Research Centre, School of Medicine, University College Dublin, National Maternity Hospital, Dublin, Ireland.
Email: sarah.louise.killeen@ucdconnect.ie

Funding information

Open access funding provided by IReL

Abstract

The period before and during pregnancy is increasingly recognized as an important stage for addressing malnutrition. This can help to reduce the risk of noncommunicable diseases in mothers and passage of risk to their infants. The FIGO Nutrition Checklist is a tool designed to address these issues. The checklist contains questions on specific dietary requirements, body mass index, diet quality, and micronutrients. Through answering these questions, awareness is generated, potential risks are identified, and information is collected that can inform health-promoting conversations between women and their healthcare professionals. The tool can be used across a range of health settings, regions, and life stages. The aim of this review is to summarize nutritional recommendations related to the FIGO Nutrition Checklist to support healthcare providers using it in practice. Included is a selection of global dietary recommendations for each of the components of the checklist and practical insights from countries that have used it. Implementation of the FIGO Nutrition Checklist will help identify potential nutritional deficiencies in women so that they can be addressed by healthcare providers. This has potential longstanding benefits for mothers and their children, across generations.

KEYWORDS

assessment, counselling, diet, FIGO Nutrition Checklist, nutrition, pregnancy

1 | INTRODUCTION

Maternal nutrition is recognized as a high-priority global health issue that requires urgent attention as it is integral to a variety of weight-, nutrition-, and health-related Sustainable Development Goals (SDGs).¹⁻³ Each year, weight-related chronic diseases cause 4 million deaths. Globally, up to 800 million people are undernourished, and at least 1 billion people are deficient in micronutrients.⁴ Investment in maternal and child health, including nutrition, has long-term benefits not only for population health, but also for the educational performance and economic productivity of the next generation.⁵

In many countries, there is a double or triple burden of malnutrition, characterized by concurrent high rates of overnutrition, undernutrition, and micronutrient deficiencies in the population, along with associated cardiometabolic or other health complications.^{6,7} This burden is seen at an individual level where people with obesity or overweight may also have nutritional deficiencies such as iron and iodine.⁸ This challenge is also prevalent in maternal nutrition. Most women do not consume diets in line with the national dietary guidelines before or during pregnancy, and there are high rates of low or late nutritional supplement use.⁹⁻¹¹ The COVID-19 pandemic has further complicated this picture as it has been associated with reduced dietary diversity, increased calorie intake, and changes in food consumption.¹²⁻¹⁵

Suboptimal diets and higher maternal weight during preconception and pregnancy can increase the risk of pregnancy complications and noncommunicable diseases in mothers and their


children in the long term.¹⁶ Improving maternal nutrition is fundamental to improving child health outcomes, protecting women's health in the postpartum period, and potentially interrupting the intergenerational passage of poor health risk.^{17,18} International organizations such as the World Health Organization (WHO) and FIGO recommend that all women receive nutrition and weight counselling during pregnancy.¹⁹ It is known that dietary counselling before and during pregnancy improves maternal nutrition knowledge, dietary intakes, and clinical outcomes like anemia, gestational weight gain, birth weight, with reduced risk of perinatal complications.²⁰⁻²⁴ Dietary counselling during lactation is also of value to support achievement of increased nutrient requirements during this time.²⁵

A woman's engagement with health-related material may differ depending on their pregnancy history or intention to conceive.²⁶⁻²⁸ Healthcare providers are encouraged to discuss pregnancy intention and provide nutrition information to women of childbearing age but engagement and practices around this vary.^{29,30} Although preconception care provides an opportunity to address several risk factors before pregnancy, most women and couples may not access routine preconception care services.³¹⁻³³ In addition, high rates of unplanned pregnancies seen globally act as a barrier to accessing this information.³⁴ Studies have shown that women of childbearing age desire additional nutrition counselling, they consider nutrition during pregnancy important, and see clinicians as the most reliable supplier of this information.³⁵⁻⁴⁰ Despite this, barriers exist such as a lack of nutrition training and supportive resources for healthcare

providers.^{41,42} Healthcare providers can use appropriate protocols and screening tools to identify nutritional risks and implement appropriate interventions.^{43,44}

The FIGO Nutrition Checklist is a validated tool that identifies unbalanced diets during the preconception, pregnancy, and postpartum periods (Figure 1). The checklist aims to facilitate conversations

between healthcare providers and women on optimal dietary intakes.⁴⁵ It can also be completed in advance of antenatal visits, thus saving time in clinical settings.⁴⁵ Beyond this, the checklist can provide useful feedback to women on their dietary issues and fills a gap when nutrition counselling does not or cannot take place.⁴⁵ The FIGO Nutrition Checklist collects information on dietary practices



Good nutrition in the mother, both before and during pregnancy, is important in ensuring healthy outcomes for her and her baby. This checklist is designed for women to complete in conjunction with her health care professional in order to assess whether nutritional intake is sufficient, and provide a basis for the health care professional to advise where changes need to be made (if applicable).

For the woman to complete in conjunction with her healthcare professional:

1). Do you have any special dietary requirements (e.g. vegetarian, vegan, allergies)? If yes, please list below:

.....

2). What is your:

a. Weight kgs

b. Height m

c. (Health care professional to complete): Divide weight in kg by height in metres then divide the answer by your height again to get your BMI.

Your BMI is kg/m²

3). Quality of diet

i) Do you eat meat or chicken 2-3 times per week? **Yes / No**

ii) Do you regularly eat more than 2 – 3 portions of fruit or vegetables per day? **Yes / No**

iii) Do you eat fish at least 1-2 times per week? **Yes / No**

iv) Do you consume dairy products (such as milk, cheese, yogurt) every day? **Yes / No**

v) Do you eat whole grain carbohydrate foods (brown bread, brown pasta, brown rice or other) at least once a day? **Yes / No**

vi) Do you consume packaged snacks, cakes, pastries or sugar-sweetened drinks less than 5 times a week? **Yes / No**

4). What is your:

i) If you are pregnant, did/do you take folate/folic acid supplements in pre-pregnancy and in early pregnancy (first 12 weeks)? **Yes / No**

ii) Do you get regular exposure to the sun (face, arms and hands for at least 10-15 mins per day)? **Yes / No**

iii) Has the doctor/nurse tested your haemoglobin (level of iron in the blood)? **Yes / No**

(Health care professional to complete) If yes, is it more than 110 g/l? **Yes / No** Enter the value:

If you have answered No to any of the questions in section 3 or 4 your nutritional status may need to be assessed in more detail.

FIGURE 1 The FIGO Nutrition Checklist

or 'special diets', body mass index, diet quality (number of servings or frequency of consumption of specific foods) and micronutrients (folic acid, vitamin D, and iron) (Figure 1). The back of the checklist (not shown in the figure, see Figure S1 includes evidence-based information for healthcare providers based on FIGO's recommendations on adolescent, preconception, and maternal nutrition: 'Think Nutrition First'⁴⁶ and the US Institute of Medicine recommendations for gestational weight gain.⁴⁷

Several studies supporting use of the FIGO Nutrition Checklist exist. In Ireland and Hong Kong, the FIGO Nutrition Checklist identified suboptimal diets in over 80% of women.^{45,48} Tsoi et al.⁴⁸ also found that the FIGO Nutrition Checklist was valid when compared with food frequency questionnaire data. Italian evidence suggests that the FIGO Nutrition Checklist was associated with pregnancy outcomes such as pregnancy-associated plasma protein A and placental volume.⁴⁹ Killeen et al.⁴⁵ identified that most women found the checklist easy and quick to complete. Qualitative research on the FIGO Nutrition Checklist suggested a need for improved practices around nutrition counselling in antenatal care and a role for the checklist in meeting this need.³⁵

The aim of this review is to provide an overview of international dietary recommendations for components of the FIGO Nutrition Checklist. A selection of freely available and published nutritional guidelines was reviewed for each component of the checklist.⁵⁰⁻⁵⁴ Countries were selected from Asia, Africa, Europe, Oceania, North America, and South America (Tables 1, 2). This review should be used as a guide only and local dietary and clinical guidelines should be followed where available. Beyond dietary guidelines, the review aims to provide practical recommendations and insights for implementation of the FIGO Nutrition Checklist. This review provides a practical tool for healthcare providers, public health specialists, and decision makers in antenatal care.

2 | SPECIAL DIETS DURING PREGNANCY AND LACTATION

A 'special diet' is a term used in the FIGO Nutrition Checklist to refer to a diet that varies from a traditional diet due to allergy, intolerance, or other medical needs, a religious or cultural diet, or a vegetarian or vegan diet. Women following special diets may be at risk of nutrient deficiencies such as protein and vitamin B12.⁵⁵ Despite this, well-planned special diets can be safe during pregnancy and lactation.⁵⁵ Specific advice can be given to women who follow a special diet to address nutritional concerns highlighted by the FIGO Nutrition Checklist. Women following vegan diets may consume insufficient calcium and/or protein. Healthcare providers can screen for this, counsel on suitable alternative sources, and determine if supplementation is required.⁵⁶ Some countries, such as India, have nutritional supplementation of key nutrients like iron and calcium as standard in antenatal care.⁵¹ In this instance, the FIGO Nutrition Checklist could be used to check compliance with supplementation guidelines.

3 | FRUIT AND VEGETABLES

Fruit and vegetables are considered nutrient-dense foods despite their low energy content.⁵⁷ Polyphenols, oligosaccharides, and fiber found in fruit and vegetables are associated with a decreased risk of chronic diseases.⁵⁸ Consuming fruit and vegetables in pregnancy is associated with a higher fiber diet that may help prevent glucose intolerance, pre-eclampsia, and constipation.^{59,60} Most countries recommend consuming at least five portions of fruit and vegetables per day (Table 1). Women with diabetes or altered glucose tolerance should be aware of high fructose consumption and its prenatal effects.⁶¹ Limited availability, cost of fruit and vegetables, poor tolerance due to nausea, reduced appetite, lack of social support, knowledge deficits, or cultural beliefs are barriers to reaching recommended fruit and vegetable intake.⁶²⁻⁶⁴

4 | DAIRY

Dairy products are high in calcium, which provides numerous health benefits including reduced risk of type 2 diabetes and cardiovascular diseases.⁶⁵ Globally, dairy products, cereals, vegetables, juices, and legumes are the main source of calcium.⁶⁶ Other less bioavailable sources include fish bones, dried fruit, nuts, and seeds.^{66,67} Low calcium intake during pregnancy increases the risk of pre-eclampsia in calcium-depleted women, stunted growth, and reduced peak bone density in teens of deficient mothers.⁶⁸ Most countries recommend consuming 2-3 servings of dairy per day. Butter, ghee, and cream are dairy-derived foods; however, they contain high amounts of saturated fat without calcium and are not recommended as part of this serving guide (Table 1). Pregnant women should avoid eating unpasteurized dairy products and soft cheeses.⁶⁹ In cases where women do not consume dairy, fortified dairy alternatives such as soy, oat, or nut products, or plant foods such as dried fruit, nuts, and beans can be a source of key nutrients.⁶⁵ There is also the option of supplementation for those at risk of calcium deficiency. In a multicenter trial of 500mg calcium or placebo from before pregnancy to 20 weeks, women with over 80% adherence had significantly reduced pre-eclampsia.⁷⁰ In India for example, oral calcium supplementation is routinely advised to women during pregnancy.⁵¹

5 | WHOLEGRAINS

Wholegrain consumption is associated with reduced risk of type 2 diabetes, cardiovascular disease, colorectal cancer, and obesity.⁷¹ The fiber content of wholegrains has the potential to assist in managing constipation, blood pressure, and blood glucose fluctuations.⁷² As illustrated in Table 1, the recommended amount of wholegrain consumption varies between countries. Some countries recommend having up to nine servings of grains, with half of these being wholegrains. These recommendations contrast with those from Ireland, for example, where it is recommended to consume 3-5 portions,

TABLE 1 Global recommendations for food-based components of the FIGO Nutrition Checklist^a

	Component of the FIGO Nutrition Checklist (daily consumption)			
	Fruit and vegetables	Dairy products	Wholegrains	Meat, poultry, or eggs
USA ⁵⁰	1.5–2 servings of fruit and 2.5–3.5 servings of vegetables per day	3 servings per day	3–5 servings per day	60–100g per day (dependent on weight), 20%–25% of total calorie intake during pregnancy Consume with every main meal
India ⁵¹	1–2 servings of fruit and 4–5 servings of vegetables per day	3–5 servings per day	50%–55% of total calorie intake 9 portions of 30g per day	Variable, depending on dietary patterns
Ireland ⁵²	5 servings per day	3 servings per day	3–5 servings per day	2 servings per day in the first and second trimesters 3 servings per day in the third trimester
Canada ⁵³	Daily, consume a variety including dark green vegetables and orange vegetables	Daily. Drink fortified soy beverages if not drinking milk	Daily	Eat lean meats and alternatives daily
Australia ⁵⁴	At least 2 servings of fruit per day and at least 5 servings of vegetables per day	3.5 servings for those aged 18 years or under. 2.5 servings per day for those aged over 18 years	8 servings of carbohydrate foods per day (serving 30–40g) for those aged 18 years or under, to 8.5 servings per day for those aged over 18 years. Choose mostly wholegrains	3.5 servings per day
Kenya ⁵³	5 servings per day	Everyday	With every meal	At least 2 servings per week. Eat red meat and liver when available
Colombia ⁵³	2–3.5 fruit servings 2–3 servings of vegetables	3–4 servings of dairy	3–3.5 servings of grains per day	2.5–3.5 servings of meat or poultry a day One egg per day

^aInformation was gathered from various international sources as referenced. Where relevant, recommendations within guidelines were converted into grams per day to estimate servings.

depending on activity level (Table 1). The number of recommended servings varies, in part because the serving size is variable across different guidelines and reference documents (Table 1).

6 | MEAT, POULTRY, AND EGGS

Adequate maternal protein intake promotes a healthy postnatal outcome and may influence childhood body composition.⁷³ The proteins found in animal products are considered complete sources as they contain all the essential amino acids, whereas plant sources are incomplete but are considered complementary and complete when paired with another source.⁷⁴ Plant proteins are becoming increasingly popular as potentially cost-effective and sustainable protein options.⁷⁵ Adequate protein intake during pregnancy is important as deficiency can cause complications such as miscarriage, fetal growth restriction, and reduced infant growth.⁷⁶ Internationally, variations in guidelines are likely due to variability in serving sizes (Table 1). Protein requirements increase during pregnancy.⁷⁷ In the USA, for

example, the recommended daily intake of protein is 46 g per day (0.8 g/kg body weight/day) in the first trimester and 71 g per day (1.1 g/kg body weight/day) during the second and third trimesters.⁷⁸ American guidelines for pregnancy recommend that women consume a variety of protein sources such as pulses, nuts, and fish.⁵⁰ Recommendations in India state that the diet of pregnant women should contain an additional 0.5 g protein during the first trimester, 6.9 g during the second trimester, and 22.7 g during the third trimester of pregnancy.⁵¹

7 | FISH

The nutritional health benefits of fish consumption primarily come from the long-chain omega-3 fatty acids eicosapentaenoic acid and docosahexaenoic acid. Fish also contain vitamins such as D and B2 (riboflavin), calcium, phosphorus, and minerals, such as iron, zinc, iodine, magnesium, and potassium.⁷⁹ There is significant geographical variation in fish intake.⁸⁰ The nutrients found in fish support

TABLE 2 Recommendations for micronutrients addressed in the FIGO Nutrition Checklist^a

	Folic acid	Vitamin D	Iron
USA ⁵⁰	400–800µg at least 1 month before pregnancy and during the first 12 weeks	No specific recommendation	When recommended by a healthcare provider
India ⁵¹	500µg per day preconceptionally and throughout pregnancy	Only provide vitamin D supplementation if deficient	Supplementation of 100mg elemental iron for 100 days during pregnancy from 16th week onward
Ireland ⁵²	400µg per day throughout pregnancy, 5 mg for those at increased risk of neural tube defects	400IU supplement per day plus dietary sources	Pregnancy 16–20mg per day
Canada ⁵³	400µg per day	600IU per day	Potential benefits of 16–20mg per day however no specific recommendation
Australia ⁵⁴	At least 400µg per day	No specific recommendations	No specific recommendation
Kenya ⁵³	400µg per day for 270 days during pregnancy	No specific recommendations	60mg per day for 270 days during pregnancy
Colombia ⁵³	Use supplement (no dose/duration specification)	No specific recommendations	Use supplement (no dose/duration specification)

^aInformation was gathered from various international sources as referenced. Where relevant, recommendations within guidelines were converted into grams per day to estimate servings.

the prevention of coronary heart disease, metabolic syndrome, and type 2 diabetes.⁸¹ They also provide neurocognitive benefits including IQ-promoting benefits, communication, and other developmental outcomes.⁸² A diet containing adequate amounts of omega-3 fatty acids is essential for fetal neurodevelopment and may protect against other adverse perinatal and longer-term outcomes.⁸³ Many countries recommend that pregnant women avoid consuming predatory fish such as shark and swordfish due to the risk of overexposure to mercury and other heavy metals.⁸⁴ Given the benefits associated with fish consumption, moderate intake of fish, such as tuna, is considered safe when limited to no more than 1–2 times per week. The aim of this is to limit heavy metal exposure.⁸⁵ Overall, most guidelines recommend consuming fish 1–2 times per week.^{50–54}

8 | PACKAGED SNACKS, CAKES, PASTRIES, OR SUGAR-SWEETENED DRINKS

The guidelines for pregnancy and the general population are the same for processed, high sugar, high fat foods, which are that they are not recommended for consumption every day (Table 1). A Global Review of Food-Based Dietary Guidelines found that most countries encourage people to limit salt; 89% to limit fat; and 84% to limit sugar; with 70% encouraging limiting all three.⁸⁶ The aim of the checklist is to identify potential overconsumption of these foods and facilitate discussion around health-promoting alternatives.

9 | FOLIC ACID

Folic acid is the synthetic form of folate, a B vitamin naturally found in leafy green vegetables, citrus fruits, and liver.⁸⁷ Deficiency

increases the risk of neural tube defects in children.⁸⁸ As a result, supplementation of 400µg per day, paired with a healthy balanced diet, is recommended in many countries for women of childbearing age, regardless of their intention to conceive given the high rates of unplanned pregnancy worldwide.^{31,89} The folic acid supplement can continue throughout pregnancy.⁵² Some women are at increased risk of neural tube defects, including those with obesity, and may require a higher dose (up to 5 mg/day) for at least the first 12 weeks of pregnancy^{16,52} (Table 2). More than 40 countries have adopted mandatory folic acid fortification policies to prevent neural tube defects, and this may affect maternal levels.⁹⁰ Folate and vitamin B12 deficiency can also cause anemia.⁹¹

10 | VITAMIN D

Vitamin D is a fat-soluble vitamin that plays an important role in calcium homeostasis and bone metabolism. Vitamin D can be obtained in the diet from a limited number of sources such as UV-grown mushrooms, eggs, and fortified products.⁹² While it is produced by our skin, vitamin D deficiency is common.^{93,94} Adult populations in Middle Eastern countries such as Iran and Syria have a very low average level of circulating vitamin D (14 ng/ml and 10 ng/ml, respectively),⁹⁴ compared with adults in European countries like Denmark and France (26 ng/ml and 24 ng/ml, respectively).⁹⁵ Several reviews have found a high prevalence of vitamin D deficiency even in countries with low latitude, where it was generally assumed that UVB radiation was adequate to prevent vitamin D deficiency, showing the potential benefits of supplementation and fortification.⁹⁶ Foods fortified with vitamin D may contain approximately 100IU per serving.⁹⁷ Severe deficiency can lead to osteomalacia and rickets in both children and adults along with other adverse health outcomes.⁹⁸ Saraf et al.⁹⁹ found vitamin D deficiency, defined as

Use of the FIGO Nutrition Checklist in India

Nutrition assessment is not a part of routine care during antenatal visits in India and there is no standard approach to how nutrition is addressed.

As a result..

1. Work is being done to advocate for use of FIGO Nutrition Checklist in pregnancy and to sensitize the healthcare providers across India during medical education programs through the national professional organization FOGSI.
2. ARTIST, an empaneled training partner for capacity building, has included a module on use of the FIGO Nutrition Checklist on its educational platform.
3. The frontline healthcare providers in urban and rural settings, who are undergoing a skill transfer program, have been sensitized to the FIGO Nutrition Checklist. Work is being done to build capacity on using the resource in clinical practice.
4. Beyond this there is capacity building, to ensure that pregnant women are screened for nutritional status with the appropriate use of the FIGO nutrition checklist.

Impact to date*:

1. The FIGO Nutrition Checklist has been used over 700 times.
2. The providers are capturing data on hemoglobin as per Government of India recommendations.
3. The prevalence of anemia in India is amongst the highest in the world and around half of all pregnant women have anemia. The data collection on micronutrients uptake and Hb levels through the FIGO nutrition checklist helped identify the status of nutrition and key performance indicators for these pregnant women in one snapshot [102].
4. More than one third of study respondents were found to have lower hemoglobin levels (<11 g/dL). This suggests that counselling pregnant women with tailored nutrition messages needs to be focused upon, for improved pregnancy outcomes.

FIGURE 2 Use of the FIGO Nutrition Checklist in India. Abbreviations: FOGSI, Federation of Obstetric and Gynaecological Societies of India; ARTIST, Asian Research and Training Institute for Skill Transfer. *Data unpublished

The FIGO Nutrition Checklist in South Africa

A recent study, the Healthy Life Trajectories Initiative (HeLTI) Bukhali trial in Soweto, incorporated the FIGO nutrition checklist into dietary counselling sessions provided to 387 young women with overweight or obesity.

The study setting was one of prevalent food insecurity and limited health literacy.

Role of the FIGO Nutrition Checklist

1. Most women (97.4%) answered 'no' to at least one diet quality question and this included suboptimal dairy (60.5%) and whole grain carbohydrate (66.3%) consumption.
2. Qualitative research highlighted the FIGO Nutrition Checklist's ease of use and value as a dietary counselling tool. The benefits of use included improved awareness of dietary intakes and a sense of improved support, knowledge, and motivation.

Future directions

Given the need and benefit, future work needs to incorporate additional adaptations, explanations, and language translations

FIGURE 3 The FIGO Nutrition Checklist in South Africa. Full data available from Soepnel et al.¹¹⁵

The FIGO Nutrition Checklist


Key points for healthcare providers

Unmet nutritional needs

Globally, women of reproductive age have unmet nutritional needs which should be addressed regardless of pregnancy intention

Available resource

The FIGO Nutrition Checklist is a free, validated, brief nutrition assessment and counselling tool which can be used to address these gaps. It can be adapted for different dietary patterns and nutritional guidelines



Impact of improved nutritional care

Nutrition should be addressed with women of reproductive age to improve their health across the life course and prevent the risk of noncommunicable diseases

Adaptation

The checklist was designed for use during preconception, pregnancy and postpartum periods. Minor adaptations would widen its application across the lifecourse and range of health settings

FIGURE 4 Key messages for healthcare providers

25-hydroxyvitamin D (25[OH]D) levels below 50nmol/L, in 54% of pregnant women globally. Although poorly defined in many regions, the breakdown by WHO region varied from 87% in Southeast Asia, 83% in the Western Pacific, 64% in the Americas, 57% in Europe, and 46% in the Eastern Mediterranean.⁹⁹ The high prevalence of maternal vitamin D deficiency may be related to differences in ethnicities and/or lifestyles (sun exposure, dietary intake, skin melanin, wearing veiled or covered clothes) rather than increased physiological requirements.

11 | IRON

Iron deficiency anemia is one of the most common health problems in women of reproductive age, affecting over one-third of pregnant women globally.¹⁰⁰ This anemia results in adverse outcomes such as increased maternal and infant mortality, mental and physical development issues, and impaired cognitive function in newborn babies.¹⁰¹ Dietary interventions are more effective in the long term for prevention of iron deficiency anemia than supplementation

and have advantages in relation to compliance, long-term acceptability, and cost-effectiveness.¹⁰² In high-risk populations however, supplementation may be more effective at reaching guideline-specified optimal levels of iron in the diet.¹⁰³ The FIGO Nutrition Checklist addresses iron intake and anemia screening. In relation to diet, it asks about the intake of meat, poultry, fish, vegetables, and fruit. Of these, red meat and other meat or fish are sources of the highly bioavailable heme iron. Other sources of iron include egg yolks, dark green leafy vegetables, beans, peas, dried fruit, and fortified cereals, although these are predominantly non-heme iron, which is a less bioavailable form.¹⁰⁴ Consuming iron-containing foods alongside those with vitamin C, such as citrus juice or fruit or vegetables, may enhance iron absorption.¹⁰³ Similarly, those at risk of deficiency should avoid eating foods that will inhibit iron absorption with the iron-containing food. These include foods that are high in calcium, tannins, or phytates.¹⁰³ The recommendations for iron vary between countries, with some including supplementation (Table 2). A morning dose of supplement may promote optimal response and alternate days could be considered to reduce gastrointestinal symptoms.¹⁰⁵

12 | FUTURE DIRECTIONS FOR THE FIGO NUTRITION CHECKLIST

Disseminating health messages can be an effective way of educating people.¹⁰⁶ Barriers that prevent women from improving their diet include differing priorities, income, and cultural norms.¹⁰⁷ Globally, one-third of adults may have reduced health literacy.¹⁰⁸ Lower levels of health literacy in pregnancy are associated with unhealthy behaviors.¹⁰⁹ Dietary advice in pregnancy should therefore be practical, implementable, and communicated clearly, using plain and simple language.¹¹⁰ The FIGO Nutrition Checklist is a tool that can be used to support this process. It is available online and free to download at: <https://www.figo.org/news/figo-nutrition-checklist>. A digital version of the FIGO Nutrition Checklist is also under development, which will allow for wider access of the resource through mobile or other electronic devices.¹¹¹ Mobile health technologies provide easy access to information and tools, they are highly acceptable to women, and are especially useful for those with lower socioeconomic status, younger age, or raised body mass index.¹¹² The FIGO Nutrition Checklist can also be used to assess response to a dietary intervention in pregnancy and appropriate core outcomes for nutritional studies are being developed.¹¹³ Dietary interventions also show promise in terms of cost-effectiveness.¹¹⁴ For further insights on how the FIGO Nutrition Checklist can be used, see [Figures 2–4](#). Future work will be to review this evidence and adapt the FIGO Nutrition Checklist as required.

13 | CONCLUSION

Using the FIGO Nutrition Checklist supports identification of nutritional deficiencies in women and girls that can be addressed. The resource can be adapted and is suitable for use across a range of health settings, world regions, and throughout the life course.

AUTHOR CONTRIBUTIONS

Sarah Louise Killeen and Niamh Donnellan wrote the manuscript with contributions from all other authors. All authors contributed to and reviewed the final manuscript.

ACKNOWLEDGMENTS

Open access funding provided by IReL.

CONFLICT OF INTEREST

Cynthia Maxwell reports grants from the Canadian Institutes for Health Research and the Crohns and Colitis Foundation of Canada. Harold David McIntyre reports honoraria for lectures from Phillips Health Care, Mead Johnson (China), and Diabetes Ireland. Sharleen O'Reilly reports research grants from the European Commission Horizon 2020, National Health and Medical Research Council of Australia, Health Research Board Ireland, Al Qasimi Foundation, and University of Sharjah. Lina Bergman reports research funds from Thermo Fischer, Roche, and Perkin Elmer and payment from

Homburg and Partner. Ronald Ma reports research support from AstraZeneca, Bayer, Novo Nordisk, Pfizer, and Tricida Inc. Other authors have no conflicts of interests to declare.

REFERENCES

- Lopez de Romaña D, Greig A, Thompson A, Arabi M. Successful delivery of nutrition programs and the sustainable development goals. *Curr Opin Biotechnol*. 2021;70:97-107.
- Heidkamp RA, Piwoz E, Gillespie S, et al. Mobilising evidence, data, and resources to achieve global maternal and child undernutrition targets and the Sustainable Development Goals: an agenda for action. *Lancet*. 2021;303:1400-1418.
- United Nations. Transforming Our World: The 2030 Agenda for Sustainable Development. Accessed July 29 2022. <https://sdgs.un.org/2030agenda>
- FAO, IFAD, UNICEF, WFP, WHO. The State of Food Security and Nutrition in the World 2019. Safeguarding against economic slowdowns and downturns. FAO; 2019. Accessed July 29, 2022. <https://www.fao.org/3/ca5162en/ca5162en.pdf>
- Stenberg K, Sweeny K, Axelson H, Temmerman M, Sheehan P. Returns on investment in the continuum of care for reproductive, maternal, newborn, and child health. In: Black RE, Laxminarayan R, Temmerman M, Walker N, eds. *Reproductive, Maternal, Newborn, and Child Health: Disease Control Priorities*. World Bank; 2016.
- Wells JC, Sawaya AL, Wibaek R, et al. The double burden of malnutrition: aetiological pathways and consequences for health. *Lancet*. 2020;4:75-88.
- Blankenship JL, Rudert C, Aguayo VM. Triple trouble: understanding the burden of child undernutrition, micronutrient deficiencies, and overweight in East Asia and the Pacific. *Matern Child Nutr*. 2020;16 Suppl 2(Suppl 2):e12950.
- de Juras AR, Hsu WC, Hu SC. The double burden of malnutrition at the individual level among adults: a nationwide survey in the Philippines. *Front Nutr*. 2021;8:760437.
- Caut C, Leach M, Steel A. Dietary guideline adherence during pre-conception and pregnancy: a systematic review. *Matern Child Nutr*. 2020;16:e12916.
- Sanghvi T, Nguyen PH, Tharaney M, et al. Gaps in the implementation and uptake of maternal nutrition interventions in antenatal care services in Bangladesh, Burkina Faso, Ethiopia and India. *Matern Child Nutr*. 2022;18:e13293.
- James-McAlpine JM, Vincze LJ, Vanderlelie JJ, Perkins AV. Influence of dietary intake and decision-making during pregnancy on birth outcomes. *Nutr Diet*. 2020;77:323-330.
- O'Connell M, Smith K, Stroud R. The dietary impact of the COVID-19 pandemic. *J Health Econ*. 2022;84:102641.
- Gupta S, Seth P, Abraham M, Pingali P. COVID-19 and women's nutrition security: panel data evidence from rural India. *Econ Polit (Bologna)*. 2022;39:157-184.
- Zhang J, Zhang Y, Huo S, et al. Emotional eating in pregnant women during the COVID-19 pandemic and its association with dietary intake and gestational weight gain. *Nutrients*. 2020;12:2250.
- Acton RB, Vanderlee L, Cameron AJ, et al. Self-reported impacts of the COVID-19 pandemic on diet-related behaviors and food security in 5 countries: results from the international food policy study 2020. *J Nutr*. 2022;152:35s-46s.
- McAuliffe FM, Killeen SL, Jacob CM, et al. Management of pre-pregnancy, pregnancy, and postpartum obesity from the FIGO Pregnancy and Non-Communicable Diseases Committee: a FIGO (International Federation of Gynecology and Obstetrics) guideline. *Int J Gynecol Obstet*. 2020;151:16-36.
- Hanson M, Jacob CM, Hod M, Killeen S, McAuliffe FM. The FIGO Pregnancy Obesity and Nutrition Initiative (PONI). *Int J Gynecol Obstet*. 2019;147:131-133.

18. Hill B, Skouteris H, Teede HJ, et al. Health in preconception, pregnancy and postpartum global alliance: international network preconception research priorities for the prevention of maternal obesity and related pregnancy and long-term complications. *J Clin Med*. 2019;8:2119.
19. World Health Organization. WHO Recommendations on Antenatal Care for a Positive Pregnancy Experience. WHO; 2016. Accessed 29 July, 2022. <https://www.who.int/publications/i/item/9789241549912>
20. Killeen SL, Geraghty AA, O'Brien EC, O'Reilly SL, Yelverton CA, McAuliffe FM. Addressing the gaps in nutritional care before and during pregnancy. *Proc Nutr Soc*. 2022;81:87-98.
21. Teweldemedhin LG, Amanuel HG, Berhe SA, Gebreyohans G, Tsige Z, Habte E. Effect of nutrition education by health professionals on pregnancy-specific nutrition knowledge and healthy dietary practice among pregnant women in Asmara, Eritrea: a quasi-experimental study. *BMJ Nutr Prev Health*. 2021;4:181-194.
22. Girard AW, Olude O. Nutrition education and counselling provided during pregnancy: effects on maternal, neonatal and child health outcomes. *Paediatr Perinat Epidemiol*. 2012;26(Suppl 1):191-204.
23. Cantor AG, Jungbauer RM, McDonagh M, et al. Counseling and behavioral interventions for healthy weight and weight gain in pregnancy: evidence report and systematic review for the US preventive services task force. *JAMA*. 2021;325:2094-2109.
24. Mohsenzadeh-Ledari F, Taghizadeh Z, Motaghi Z, Keramat A, Moosazadeh M, Najafi A. Appropriate interventions for pregnant women with indicators of metabolic syndrome on pregnancy outcomes: a systematic review. *Int J Prev Med*. 2019;15(10):2.
25. Kavle JA, Mehanna S, Khan G, Hassan M, Saleh G, Engmann C. Program considerations for integration of nutrition and family planning: Beliefs around maternal diet and breastfeeding within the context of the nutrition transition in Egypt. *Matern Child Nutr*. 2018;14:e12469.
26. Szwajcer EM, Hiddink GJ, Maas L, Koelen MA, van Woerkum CM. Nutrition-related information-seeking behaviours of women trying to conceive and pregnant women: evidence for the life course perspective. *Fam Pract*. 2008;25(Suppl 1):i99-i104.
27. Phelan S. Pregnancy: a "teachable moment" for weight control and obesity prevention. *Am J Obstet Gynecol*. 2007;202:135.e131-135.e1358.
28. Lindqvist M, Lindqvist M, Eurenus E, Persson M, Mogren I. Change of lifestyle habits - motivation and ability reported by pregnant women in northern Sweden. *Sex Reprod Healthc*. 2017;13:83-90.
29. Gardiner PM, Nelson L, Shellhaas CS, et al. The clinical content of preconception care: nutrition and dietary supplements. *Am J Obstet Gynecol*. 2008;199:345-356.
30. Premji S, McDonald SW, Zaychowsky C, Zwicker JD. Supporting healthy pregnancies: examining variations in nutrition, weight management and substance abuse advice provision by prenatal care providers in Alberta, Canada. A study using the All Our Families cohort. *PLoS One*. 2019;14:e0210290.
31. Jacob CM, Killeen SL, McAuliffe FM, et al. Prevention of noncommunicable diseases by interventions in the preconception period: a FIGO position paper for action by healthcare practitioners. *Int J Gynecol Obstet*. 2020;151(Suppl 1(Suppl 1):6-15.
32. Masinter LM, Dina B, Kjerulff K, Feinglass J. Short interpregnancy intervals: results from the first baby study. *Womens Health Issues*. 2017;27:426-433.
33. Moholdt T, Hawley JA. Maternal lifestyle interventions: targeting preconception health. *Trends Endocrinol Metab*. 2020;31:561-569.
34. Wellings K, Jones KG, Mercer CH, et al. The prevalence of unplanned pregnancy and associated factors in Britain: findings from the third National Survey of Sexual Attitudes and Lifestyles (Natsal-3). *Lancet*. 2013;382:1807-1816.
35. Killeen SL, Callaghan SL, Jacob CM, Hanson MA, McAuliffe FM. "It only takes two minutes to ask"—a qualitative study with women on using the FIGO Nutrition Checklist in pregnancy. *Int J Gynecol Obstet*. 2020;151(Suppl 1):45-50.
36. Bookari K, Yeatman H, Williamson M. Falling short of dietary guidelines - What do Australian pregnant women really know? A cross sectional study. *Women Birth*. 2017;30:9-17.
37. Owusu-Addo SB, Owusu-Addo E, Morhe ES. Health information-seeking behaviours among pregnant teenagers in Ejisu-Juaben Municipality, Ghana. *Midwifery*. 2016;41:110-117.
38. Rahmawati W, Willcox JC, van der Pligt P, Worsley A. Nutrition information-seeking behaviour of Indonesian pregnant women. *Midwifery*. 2021;100:103040.
39. Beckham AJ, Urrutia RP, Sahadeo L, Corbie-Smith G, Nicholson W. "We know but we don't really know": diet, physical activity and cardiovascular disease prevention knowledge and beliefs among underserved pregnant women. *Matern Child Health J*. 2015;19:1791-1801.
40. Porteous HE, Palmer MA, Wilkinson SA. Informing maternity service development by surveying new mothers about preferences for nutrition education during their pregnancy in an area of social disadvantage. *Women Birth*. 2014;27:196-201.
41. Adamski M, Gibson S, Leech M, Truby H. Are doctors nutritionists? What is the role of doctors in providing nutrition advice? *Nutrition Bulletin*. 2018;43:147-152.
42. DiMaria-Ghalili R, Mirtallo J, Tobin B, Hark L, Van Horn L, Palmer C. Challenges and opportunities for nutrition education and training in the health care professions: intraprofessional and interprofessional call to action. *Am J Clin Nutr*. 2014;99:1184S-1193S.
43. Dunneram Y, Jeewon R. Healthy diet and nutrition education program among women of reproductive age: a necessity of multilevel strategies or community responsibility. *Health Promot Perspect*. 2015;7(5):116-127.
44. Atlantis E, John JR, Fahey PP, Hocking S, Peters K. Clinical usefulness of brief screening tool for activating weight management discussions in primary care (AWARE): a nationwide mixed methods pilot study. *PLoS One*. 2021;28(16):e0259220.
45. Killeen SL, Callaghan SL, Jacob CM, Hanson MA, McAuliffe FM. Examining the use of the FIGO Nutrition Checklist in routine antenatal practice: multistakeholder feedback to implementation. *Int J Gynecol Obstet*. 2020;151(Suppl 1):51-56.
46. Hanson MA, Bardsley A, De-Regil LM, et al. The International Federation of Gynecology and Obstetrics (FIGO) recommendations on adolescent, preconception, and maternal nutrition: "Think Nutrition First". *Int J Gynecol Obstet*. 2015;131:13-53.
47. Institute of Medicine and National Research Council Committee to Reexamine IOM Pregnancy Weight Guidelines. In: Rasmussen KM, Yaktine AL, eds. *Weight Gain During Pregnancy: Reexamining the Guidelines*. National Academies Press; 2009.
48. Tsoi KY, Chan RSM, Li LS, et al. Evaluation of dietary pattern in early pregnancy using the FIGO Nutrition Checklist compared to a food frequency questionnaire. *Int J Gynecol Obstet*. 2020;151:37-44.
49. Parisi F, Savasi V, di Bartolo I, Mandia L, Cetin I. Associations between first trimester maternal nutritional score, early markers of placental function, and pregnancy outcome. *Nutrients*. 2020;12:799.
50. US Department of Agriculture, US Department of Health and Human Services. *Dietary Guidelines for Americans, 2020-2025*. 9th ed. USDA; 2020. Accessed July 29, 2022. https://www.dietaryguidelines.gov/sites/default/files/2020-12/Dietary_Guidelines_for_Americans_2020-2025.pdf
51. National Institute of Nutrition. *Dietary guidelines for Indians - a manual*. NIN; 2011. Accessed July 29, 2022. <https://www.nin.res.in/downloads/DietaryGuidelinesforNINwebsite.pdf>
52. Institute of Obstetricians and Gynaecologists, Royal College of Physicians of Ireland, Directorate of Clinical Strategy and

- Programmes, Health Service Executive. Clinical Practice Guideline: Nutrition During Pregnancy, 2019. Accessed July 29 2022. <https://www.hse.ie/eng/about/who/acute-hospitals-division/woman-infants/clinical-guidelines/nutrition-during-pregnancy.pdf>
53. UNICEF. Review of national Food-Based Dietary Guidelines and associated guidance for infants, children, adolescents, and pregnant and lactating women. UNICEF; 2021. Accessed 29 July, 2022. <https://www.unicef.org/media/102761/file/2021-Food-based-Dietary-Guidelines-final.pdf>
 54. National Health and Medical Research Council. Australian Dietary Guidelines. NHMRC; 2013. Accessed July 29, 2022. https://www.eatforhealth.gov.au/sites/default/files/files/the_guidelines/n55_austrian_dietary_guidelines.pdf
 55. Pawlak R. To vegan or not to vegan when pregnant, lactating or feeding young children. *Eur J Clin Nutr*. 2017;71:1259-1262.
 56. Ota E, Hori H, Mori R, Tobe-Gai R, Farrar D. Antenatal dietary education and supplementation to increase energy and protein intake. *Cochrane Database Syst Rev*. 2015;(6):CD000032.
 57. Darmon N, Darmon M, Maillot M, Drewnowski A. A nutrient density standard for vegetables and fruits: nutrients per calorie and nutrients per unit cost. *J Am Diet Assoc*. 2005;105:1881-1887.
 58. van der Merwe M. Gut microbiome changes induced by a diet rich in fruits and vegetables. *Int J Food Sci Nutr*. 2020;72:665-669.
 59. Pretorius R, Palmer D. High-fiber diet during pregnancy characterized by more fruit and vegetable consumption. *Nutrients*. 2020;13:35.
 60. Zerfu TA, Mekuria A. Pregnant women have inadequate fiber intake while consuming fiber-rich diets in low-income rural setting: evidences from analysis of common "ready-to-eat" stable foods. *Food Sci Nutr*. 2019;7:3286-3292.
 61. Zhang H, Li X, Niu Y, et al. Fasting serum fructose is associated with risk of gestational diabetes mellitus. *BMC Pregnancy Childbirth*. 2022;22:446.
 62. Kavle JA, Landry M. Addressing barriers to maternal nutrition in low- and middle-income countries: a review of the evidence and programme implications. *Matern Child Nutr*. 2018;14:12508.
 63. Hromi-Fiedler A, Chapman D, Segura-Pérez S, et al. Barriers and facilitators to improve fruit and vegetable intake among WIC-eligible pregnant Latinas: an application of the health action process approach framework. *J Nutr Educ Behav*. 2016;48:468-477.e1.
 64. Kehoe SH, Dhurde V, Bhaise S, et al. Barriers and facilitators to fruit and vegetable consumption among rural Indian women of reproductive age. *Food Nutr Bull*. 2019;40:87-98.
 65. Thorning TK, Raben A, Tholstrup T, Soedamah-Muthu SS, Givens I, Astrup A. Milk and dairy products: good or bad for human health? An assessment of the totality of scientific evidence. *Food Nutr Res*. 2016;60:32527.
 66. Cormick G, Belizán JM. Calcium intake and health. *Nutrients*. 2019;15(11):1606.
 67. Hansen M, Thilsted SH, Sandström B, et al. Calcium absorption from small soft-boned fish. *J Trace Elem Med Biol*. 1998;12:148-154.
 68. Gomes F, Ashorn P, Askari S, et al. Calcium supplementation for the prevention of hypertensive disorders of pregnancy: current evidence and programmatic considerations. *Ann N Y Acad Sci*. 2022;1510:52-67.
 69. Moran LJ, Verwiel Y, Bahri Khomami M, Roseboom TJ, Painter RC. Nutrition and listeriosis during pregnancy: a systematic review. *J Nutr Sci*. 2018;7:e25.
 70. Hofmeyr GJ, Betrán AP, Singata-Madliki M, et al. Prepregnancy and early pregnancy calcium supplementation among women at high risk of pre-eclampsia: a multicentre, double-blind, randomised, placebo-controlled trial. *Lancet*. 2019;26(393):330-339.
 71. Seal C, Brownlee I. Whole-grain foods and chronic disease: evidence from epidemiological and intervention studies. *Proc Nutr Soc*. 2015;74:313-319.
 72. Nirmala Prasadi VP, Joye I. Dietary fibre from whole grains and their benefits on metabolic health. *Nutrients*. 2020;12:3045.
 73. Elango R, Ball R. Protein and amino acid requirements during pregnancy. *Adv Nutr*. 2016;7:839S-844S.
 74. Young VR, Pellett PL. Plant proteins in relation to human protein and amino acid nutrition. *Am J Clin Nutr*. 1994;59:1203S-1212S.
 75. Aggarwal A, Drewnowski A. Plant- and animal-protein diets in relation to sociodemographic drivers, quality, and cost: findings from the Seattle Obesity Study. *Am J Clin Nutr*. 2019;110:451-460.
 76. Wu G, Imhoff-Kunsch B, Girard A. Biological mechanisms for nutritional regulation of maternal health and fetal development. *Paediatr Perinat Epidemiol*. 2012;26(Suppl):4-26.
 77. Joint WHO/FAO/UNU Expert Consultation. Protein and amino acid requirements in human nutrition. *World Health Organ Tech Rep Ser*. 2007;935:1-265.
 78. Murphy MM, Higgins KA, Bi X, Barraj LM. Adequacy and sources of protein intake among pregnant women in the United States, NHANES 2003-2012. *Nutrients*. 2021;13:795.
 79. Mohanty BP, Mahanty A, Ganguly S, Mitra T, Karunakaran D, Anandan R. Nutritional composition of food fishes and their importance in providing food and nutritional security. *Food Chem*. 2019;293:561-570.
 80. Welch A, Lund E, Amiano P, et al. Variability of fish consumption within the 10 European countries participating in the European Investigation into Cancer and Nutrition EPIC study. *Public Health Nutr*. 2002;5(6B):1273-1285.
 81. Kendall-Tackett K. Long-chain omega-3 fatty acids and women's mental health in the perinatal period and beyond. *J Midwifery Womens Health*. 2010;55:561-567.
 82. Hibbeln J, Davis J, Steer C, et al. Maternal seafood consumption in pregnancy and neurodevelopmental outcomes in childhood (ALSPAC Study): an observational cohort study. *Lancet*. 2007;369:578-585.
 83. Koletzko B, Cetin I, Brenna JT, et al. Dietary fat intakes for pregnant and lactating women. *Br J Nutr*. 2007;98:873-877.
 84. Grandjean P, Lederman SA, Silbergeld EK. Fish consumption during pregnancy. *JAMA Pediatr*. 2019;173:292.
 85. Bramante CT, Spiller P, Landa M. Fish consumption during pregnancy: an opportunity, not a risk. *JAMA Pediatr*. 2018;172:801-802.
 86. Herforth A, Arimond M, Álvarez-Sánchez C, Coates J, Christianson K, Muehlhoff E. A global review of food-based dietary guidelines. *Adv Nutr*. 2019;10:590-605.
 87. Donnelly JG. Folic acid. *Crit Rev Clin Lab Sci*. 2001;38:183-223.
 88. Silva C, Keating E, Pinto E. The impact of folic acid supplementation on gestational and long-term health: critical temporal windows, benefits and risks. *Porto Biomed*. 2017;2:315-332.
 89. Locksmith G. Preventing neural tube defects: the importance of periconceptional folic acid supplements. *Obstet Gynecol*. 1998;91:1027-1034.
 90. Food Safety Authority of Ireland. Report of the national committee on folic acid fortification. Food Safety Authority; 2006. Accessed 29 July, 2022. https://www.fsai.ie/uploadedfiles/folic_acid.pdf
 91. Kangalgil M, Sahinler A, Kırkbir IB, Özcelik AO. Associations of maternal characteristics and dietary factors with anemia and iron-deficiency in pregnancy. *J Gynecol Obstet Hum Reprod*. 2021;50:102137.
 92. Dominguez L, Farruggia M, Verones N, Barbagallo M. Vitamin D sources, metabolism, and deficiency: available compounds and guidelines for its treatment. *Metabolites*. 2021;11:255.
 93. Smith E, Holick M. The skin: the site of vitamin D₃ synthesis and a target tissue for its metabolite 1,25-dihydroxyvitamin D₃. *Steroids*. 1987;49:103-131.
 94. Kaykhaei M, Hashemi M, Narouie B, et al. High prevalence of vitamin D deficiency in Zahedan, Southeast Iran. *Ann Nutr Metab*. 2011;58:37-41.
 95. Souberbielle JC, Massart C, Brailly-Tabard S, Cavalier E, Chanson P. Prevalence and determinants of vitamin D deficiency in healthy French adults: the VARIETE study. *Endocrine*. 2016;53:543-550.

96. Mithal A, Wahl DA, Bonjour JP, et al. Global vitamin D status and determinants of hypovitaminosis D. *Osteoporos Int*. 2009;20:1807-1820.
97. Ritu G, Gupta A. Fortification of Foods with Vitamin D in India. *Nutrients*. 2014;6:3601-3623.
98. Amrein K, Scherkl M, Hoffmann M, et al. Vitamin D deficiency 2.0: an update on the current status worldwide. *Eur J Clin Nutr*. 2020;74:1498-1513.
99. Saraf R, Morton S, Camargo C, Grant C. Global summary of maternal and newborn vitamin D status - a systematic review. *Matern Child Nutr*. 2016;12:647-668.
100. World Health Organization [website]. The Global Health Observatory. *Anaemia in Women and Children*. WHO global anaemia estimates, 2021 edition. Accessed 29 July, 2022. https://www.who.int/data/gho/data/themes/topics/anaemia_in_women_and_children
101. Mettananda S, Suranjan M, Fernando R, et al. Anaemia among females in child-bearing age: relative contributions, effects and interactions of α - and β -thalassaemia. *PLoS One*. 2018;13:e0206928.
102. Patterson A, Brown W, Roberts D, Seldon M. Dietary treatment of iron deficiency in women of childbearing age. *Am J Clin Nutr*. 2021;74:650-656.
103. Prentice A, Mendoza Y, Pereira D, et al. Dietary strategies for improving iron status: balancing safety and efficacy. *Nutr Rev*. 2016;75:49-60.
104. Lv C, Zhao G, Lönnerdal B. Bioavailability of iron from plant and animal ferritins. *J Nutr Biochem*. 2015;26(5):532-540.
105. Stoffel NU, von Siebenthal HK, Moretti D, Zimmermann MB. Oral iron supplementation in iron-deficient women: how much and how often? *Mol Aspects Med*. 2020;75:100865.
106. Hales S, Dunn C, Wilcox S, Turner-McGrievy G. Is a picture worth a thousand words? Few evidence-based features of dietary interventions included in photo diet tracking mobile apps for weight loss. *J Diabetes Sci Technol*. 2016;10:1399-1405.
107. Grenier LN, Atkinson SA, Mottola MF, et al. Be healthy in pregnancy: exploring factors that impact pregnant women's nutrition and exercise behaviours. *Matern Child Nutr*. 2021;17:13068.
108. Moreira L. *Health literacy for people-centred care: Where do OECD countries stand?* OECD Health Working Papers, No. 107, OECD Publishing; 2018.
109. Kilfoyle KA, Vitko M, O'Connor R, Bailey SC. Health literacy and women's reproductive health: a systematic review. *J Womens Health (Larchmt)*. 2016;25:1237-1255.
110. Nawabi F, Krebs F, Vennedey V, Shukri A, Lorenz L, Stock S. Health literacy in pregnant women: a systematic review. *Int J Environ Res Public Health*. 2021;18:3847.
111. FIGO [website]. FIGO's Nutrition Checklist wins semi-final at the HSE-HIHI Spark Ignite awards. 2021. Accessed November 12, 2021. <https://www.figo.org/news/figos-nutrition-checklist-wins-semi-final-hse-hihi-spark-ignite-awards>
112. Greene EM, O'Brien EC, Kennelly MA, et al. Acceptability of the Pregnancy, Exercise, and Nutrition Research Study With Smartphone App Support (PEARS) and the use of mobile health in a mixed lifestyle intervention by pregnant obese and overweight women: secondary analysis of a randomized controlled trial. *JMIR Mhealth Uhealth*. 2021;9:e17189.
113. Killeen SL, O'Brien EC, Jacob CM, O'Reilly SL, Hanson M, McAuliffe FM. PREgnancy Nutrition: a protocol for the development of a Core Outcome Set (PRENCOS). *Int J Gynecol Obstet*. 2019;147:134-139.
114. O'Sullivan EJ, Rokicki S, Kennelly M, Ainscough K, McAuliffe FM. Cost-effectiveness of a mobile health-supported lifestyle intervention for pregnant women with an elevated body mass index. *Int J Obes (Lond)*. 2020;44:999-1010.
115. Soepnel LM, Draper CE, Mabetha K, et al. Evaluating implementation of the FIGO Nutrition Checklist for preconception and pregnancy within the *Bukhali* trial in Soweto, South Africa. *Int J Gynecol Obstet*. 2023;160 (Suppl 1):68-79.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Killeen SL, Donnellan N, O'Reilly SL, et al. Using FIGO Nutrition Checklist counselling in pregnancy: A review to support healthcare professionals. *Int J Gynecol Obstet*. 2023;160(Suppl. 1):10-21. doi:[10.1002/ijgo.14539](https://doi.org/10.1002/ijgo.14539)