












## REVIEW

# Beyond the genomes of *Fulvia fulva* (syn. *Cladosporium fulvum*) and *Dothistroma septosporum*: New insights into how these fungal pathogens interact with their host plants

Carl H. Mesarich<sup>1,2</sup>  | Irene Barnes<sup>3</sup>  | Ellie L. Bradley<sup>1</sup>  | Silvia de la Rosa<sup>1</sup> | Pierre J. G. M. de Wit<sup>4</sup> | Yanan Guo<sup>2,5</sup>  | Scott A. Griffiths<sup>6,7</sup> | Richard C. Hamelin<sup>8,9</sup> | Matthieu H. A. J. Joosten<sup>4</sup>  | Mengmeng Lu<sup>10</sup>  | Hannah M. McCarthy<sup>5</sup>  | Christiaan R. Schol<sup>4,11</sup>  | Ioannis Stergiopoulos<sup>12</sup>  | Mariana Tarallo<sup>5</sup> | Alex Z. Zaccaron<sup>12</sup>  | Rosie E. Bradshaw<sup>2,5</sup> 

<sup>1</sup>Laboratory of Molecular Plant Pathology, School of Agriculture and Environment, Massey University, Palmerston North, New Zealand

<sup>2</sup>Bioprotection Aotearoa, Massey University, Palmerston North, New Zealand

<sup>3</sup>Department of Biochemistry, Genetics and Microbiology, Forestry and Agricultural Biotechnology Institute, University of Pretoria, Pretoria, South Africa

<sup>4</sup>Laboratory of Phytopathology, Wageningen University, Wageningen, Netherlands

<sup>5</sup>Laboratory of Molecular Plant Pathology, School of Natural Sciences, Massey University, Palmerston North, New Zealand

<sup>6</sup>Lazarus Bio Ltd, Llanelli, UK

<sup>7</sup>Hydro Industries Ltd, Llanelli, UK

<sup>8</sup>Department of Forest and Conservation Sciences, University of British Columbia, Vancouver, British Columbia, Canada

<sup>9</sup>Institut de Biologie Intégrative et des Systèmes, Université Laval, Québec City, Québec, Canada

<sup>10</sup>Department of Biological Sciences, University of Calgary, Calgary, Alberta, Canada

<sup>11</sup>Plant Breeding, Wageningen University & Research, Wageningen, Netherlands

<sup>12</sup>Department of Plant Pathology, University of California Davis, Davis, California, USA

## Correspondence

Rosie E. Bradshaw, School of Natural Sciences, Massey University, Palmerston North 4410, New Zealand.

Email: [r.e.bradshaw@massey.ac.nz](mailto:r.e.bradshaw@massey.ac.nz)

## Abstract

*Fulvia fulva* and *Dothistroma septosporum* are closely related apoplastic pathogens with similar lifestyles but different hosts: *F. fulva* is a pathogen of tomato, whilst *D. septosporum* is a pathogen of pine trees. In 2012, the first genome sequences of these pathogens were published, with *F. fulva* and *D. septosporum* having highly fragmented and near-complete assemblies, respectively. Since then, significant advances have been made in unravelling their genome architectures. For instance, the genome of *F. fulva* has now been assembled into 14 chromosomes, 13 of which have synteny with the 14 chromosomes of *D. septosporum*, suggesting these pathogens are even more closely related than originally thought. Considerable advances have also been made in the identification and functional characterization of virulence factors (e.g., effector proteins and secondary metabolites) from these pathogens, thereby providing new insights into how they promote host colonization or activate plant defence responses.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2023 The Authors. *Molecular Plant Pathology* published by British Society for Plant Pathology and John Wiley & Sons Ltd.

For example, it has now been established that effector proteins from both *F. fulva* and *D. septosporum* interact with cell-surface immune receptors and co-receptors to activate the plant immune system. Progress has also been made in understanding how *F. fulva* and *D. septosporum* have evolved with their host plants, whilst intensive research into pandemics of Dothistroma needle blight in the Northern Hemisphere has shed light on the origins, migration, and genetic diversity of the global *D. septosporum* population. In this review, we specifically summarize advances made in our understanding of the *F. fulva*–tomato and *D. septosporum*–pine pathosystems over the last 10 years.

#### KEYWORDS

Dothistroma needle blight (DNB), effector proteins, genome sequences, host susceptibility and resistance, pathogen diversity, secondary metabolites, tomato leaf mould

## 1 | INTRODUCTION

The year 2022 marked 10 years since the first genome sequences of the dothideomycete pathogens *Fulvia fulva* and *Dothistroma septosporum* were published and compared (de Wit et al., 2012). The nomenclature of *F. fulva* has been the subject of debate for decades, with *Cladosporium fulvum*, *F. fulva*, and *Passalora fulva* being proposed (reviewed in Thomma et al., 2005). However, recent multilocus-based DNA barcoding has shown that *F. fulva* is the currently accepted name (Videira et al., 2017). Indeed, this fungus clusters more closely with *Dothistroma* than with members of the *Cladosporium* or *Passalora* genera. *F. fulva* and *D. septosporum* were an unusual choice for a comparative genome analysis as, although they have similar lifestyles, they infect very different hosts. Specifically, *F. fulva* is considered a hemibiotrophic pathogen of tomato, an angiosperm (Hane et al., 2020; Thomma et al., 2005), whilst *D. septosporum*, previously called *Dothistroma pini* (Barnes et al., 2004), is a hemibiotrophic pathogen of gymnosperm conifers, such as pine trees (Kabir et al., 2015b). Here, hemibiotrophs are defined as those pathogens that start infection as a biotroph and then switch to necrotrophy (de Wit et al., 2012; Hane et al., 2020). Both species invade their hosts through stomata and then colonize the apoplastic environment as part of a long latent asymptomatic phase, which is also observed for other closely related hemibiotrophs like *Zymoseptoria tritici* (Brennan et al., 2019). However, whilst *F. fulva* sporulates out of stomata, *D. septosporum* erupts out of needle tissue during sporulation (Kabir et al., 2015b; Thomma et al., 2005). Like *Z. tritici*, but not *F. fulva*, cell death is required for sporulation by *D. septosporum* (Kabir et al., 2015b; Sánchez-Vallet et al., 2015).

In 2012, *F. fulva* was already a well-established model for studying molecular plant–pathogen interactions and was the species from which the first fungal avirulence (*Avr*) effector gene, *CfAvr9*, was cloned (van Kan et al., 1991). In contrast, *D. septosporum* was relatively unknown, except as the causal agent of a forest disease epidemic that was suspected to be associated with climate change (Woods et al., 2005), and for its ability to secrete a secondary

metabolite toxin, dothistromin (Bradshaw et al., 2006). The two species were compared because they are phylogenetically close relatives, as evident in similarities in their genome sequences (de Wit et al., 2012; Ohm et al., 2012). The comparative analysis revealed specific differences in gene and repeat content, but also identified many intriguing similarities such as orthologous effector and dothistromin biosynthetic genes (de Wit et al., 2012).

Since 2012, several additional genomic, transcriptomic, and proteomic resources for *F. fulva* and *D. septosporum* have been made available (Table 1) (Bradshaw et al., 2016; Mesarich et al., 2014, 2018; Zaccaron et al., 2022). In combination with the original genome sequences, these resources have greatly advanced our understanding of how these two pathogens interact with, and have evolved with, their host plants. In this review, we specifically highlight the advances that have taken place since the publication of the original *F. fulva* and *D. septosporum* genome sequences. A glossary of the full gene/protein names mentioned in this review can be found in Table S1.

## 2 | COMPLETION OF THE *F. FULVA* AND *D. SEPTOSPORUM* GENOME SEQUENCES

The first draft genome sequences of *F. fulva* isolate OWU and *D. septosporum* isolate NZE10 were described by de Wit et al. (2012). In the case of *D. septosporum* NZE10, the genome was assembled into 14 chromosomes, while for *F. fulva* OWU a chromosome-level genome assembly could not be obtained. Instead, there was a high degree of fragmentation due mainly to the much higher repetitive DNA content in *F. fulva* OWU (47.2%) compared to *D. septosporum* NZE10 (3.2%) (de Wit et al., 2012).

Recently, by combining long-read sequencing with a chromatin conformation capture technique, a chromosome-level genome assembly for *F. fulva* isolate Race 5 was obtained (Table 2) (Zaccaron et al., 2022). Like *D. septosporum* NZE10, the genome of *F. fulva* Race 5 is predicted to contain 14 chromosomes (Chr), of which nine show a one-to-one match to those from *D. septosporum* NZE10, while a further four match two or

TABLE 1 Genomic, transcriptomic, and proteomic resources available for *Fulvia fulva* and *Dothistroma septosporum*.

Species and isolate	Sequence resource	NCBI BioProject accession or JGI link <sup>a</sup>	Reference
<i>F. fulva</i>			
OWU (CBS 131901)	Genome	PRJNA86753 Clafu1	de Wit et al. (2012)
OWU	Transcriptome (on <i>Solanum lycopersicum</i> ; time course)	PRJNA86753	Mesarich et al. (2014)
OWU	Transcriptome (in culture; nutrient-rich and nutrient-poor media)	PRJNA86753	Mesarich et al. (2014)
Race 5 (5 Kim, IPO 1979)	Genome	PRJNA565804	Zaccaron et al. (2022)
Race 5	Transcriptome (on <i>S. lycopersicum</i> ; single time point)	PRJNA86753	Mesarich et al. (2014)
Race 5	Transcriptome (in culture; nutrient-rich and nutrient-poor media)	PRJNA565804	Zaccaron et al. (2022)
Four global isolates, including OWU and Race 5	Proteome (mixed collection of small, secreted proteins in apoplastic washing fluid from <i>S. lycopersicum</i> at different time points)		Mesarich et al. (2018)
<i>D. septosporum</i>			
NZE10	Genome	PRJNA74753 Dotse1	de Wit et al. (2012)
NZE10	Transcriptome (on <i>Pinus radiata</i> ; time course)	Dotse1	Bradshaw et al. (2016)
NZE10	Transcriptome (in culture; pine extract media <sup>b</sup> )	PRJNA584643	de Wit et al. (2012)
NZE10	Transcriptome (in culture; nutrient-rich media <sup>b</sup> )	PRJNA584708	de Wit et al. (2012)
NZE10 and <i>DsLaeA</i> mutant	Transcriptome (in culture; nutrient-rich and nutrient-poor media)	PRJNA422893	Ozturk et al. (2019)
18 global isolates	Genome	PRJNA381823	Bradshaw, Sim, et al. (2019)
12 New Zealand isolates	Genome	PRJNA426106	Bradshaw, Ormond, et al. (2019)
25 Scottish isolates	Genome	PRJEB14366	Ennos et al. (2020)

<sup>a</sup>NCBI, National Center for Biotechnological Information (<https://www.ncbi.nlm.nih.gov/>); JGI, Joint Genome Institute (<https://mycocosm.jgi.doe.gov/mycocosm/home>).

<sup>b</sup>Unreplicated expressed sequence tags (ESTs) used at the JGI for genome annotation.

more chromosomes (Figure 1a). The remaining chromosome, Chr14, is dispensable, as it is absent from 19 out of 24 *F. fulva* isolates examined to date (but is present in isolate OWU) and has no matches to any of the chromosomes from *D. septosporum* NZE10 (Zaccaron et al., 2022).

Further comparative analysis of the chromosomes from *F. fulva* Race 5 and *D. septosporum* NZE10 revealed a clear pattern of mesosyteny, consisting of a few interchromosomal rearrangements but many intra-chromosomal rearrangements (Figure 1a) (Zaccaron et al., 2022). In addition, large differences in repetitive DNA content meant that while the chromosomes of *D. septosporum* NZE10 are almost entirely composed of gene-dense regions with a high GC content, those of *F. fulva* Race 5 contain long repeat-rich regions with a low GC content that intersperse the gene-dense regions (Figure 1b). Such structural differences account for the large difference in size between matching chromosomes of *F. fulva* Race 5 and *D. septosporum* NZE10 and, consequently, in the genome sizes of these two species (Zaccaron et al., 2022).

Structural differences among isolates of the same species were also noted. In a recent study, the genomes of 18 *D. septosporum* isolates from a worldwide collection were sequenced, resulting in the

identification of large-scale structural variants that included cases of aneuploidy and a reciprocal translocation between Chr5 and Chr13 in isolate NZE10, compared to isolates from other continents (Figure 1a) (Bradshaw, Sim, et al., 2019). Consequently, while Chr3 of *F. fulva* Race 5 matches Chr5 and Chr13 of *D. septosporum* NZE10, this same chromosome matches a single scaffold in the genome assembly of *D. septosporum* isolate SLV1 (Figure 1a).

Comparative analysis of gene content revealed that *F. fulva* Race 5 has a considerably larger number of predicted genes than *D. septosporum* NZE10 (Table 3), which is reflected in the functional gene categories of these two species (de Wit et al., 2012; Zaccaron et al., 2022) and is discussed more thoroughly in the next sections of this review.

### 3 | EFFECTORS AND OTHER SECRETED PROTEINS OF *F. FULVA* AND *D. SEPTOSPORUM*

Like all plant-pathogenic fungi, *F. fulva* and *D. septosporum* secrete effectors and other proteins into their hosts during infection. In

**TABLE 2** Comparison of the genome assemblies of *Fulvia fulva* isolate Race 5 and *Dothistroma septosporum* isolate NZE10.

Assembly statistics	<i>F. fulva</i> Race 5	<i>D. septosporum</i> NZE10
Genome size (bp)	67,169,167	30,209,431
Number of scaffolds <sup>a</sup>	14	14
Longest scaffold (bp)	11,362,290	5,111,597
Scaffold N <sub>50</sub> (Mb)	5,777,465	2,595,548
Scaffold L <sub>50</sub>	5	5
Scaffold N <sub>90</sub> (Mb)	3.3	1.5
Scaffold L <sub>90</sub>	10	11
GC content (%)	48.9	53.1
Complete chromosomes <sup>b</sup>	14	11

<sup>a</sup>Scaffolds longer than 50 kb.

<sup>b</sup>Complete chromosomes refer to the number of assembled scaffolds containing telomeric repeats at both ends.

susceptible plants, these proteins promote everything from host colonization to pathogen lifecycle completion (Bradley et al., 2022; Lo Presti et al., 2015; Rocafort et al., 2020). However, in resistant plants, one or more of these proteins are recognized as avirulence determinants (Avr proteins) by corresponding extracellular or intracellular immune receptors that trigger immune responses (Cook et al., 2015; van der Burgh & Joosten, 2019). These immune responses include the production of antifungal phytoalexins, reactive oxygen species (ROS), and pathogenesis-related (PR) proteins (e.g., chitinases, glucanases, and proteases), and are often accompanied by a hypersensitive cell death response (HR) (de Wit, 2016; de Wit et al., 2009).

### 3.1 | Effector proteins of *F. fulva*

Prior to 2012, a number of effectors from *F. fulva* had already been identified and functionally characterized, including up to eight Avr proteins that are recognized by corresponding Cf (resistance to *C. fulvum*) immune receptors in tomato (de Wit, 2016; de Wit et al., 2009). Since this time, one additional Avr protein, CfAvr5, recognized by the extracellular Cf-5 immune receptor of tomato (Dixon et al., 1998), has been identified (Mesarich et al., 2014). To identify CfAvr5, an expressed transcript sequence comparison between 44 in planta-induced (candidate) effector genes of *F. fulva* isolates OWU and Race 5 (the latter being able to overcome Cf-5-mediated resistance) was carried out. This revealed that one of these genes has a frameshift mutation in Race 5, resulting in a premature stop codon. Through a complementation assay, in which the wild-type (WT) version of this gene from isolate OWU was introduced into isolate Race 5, it was subsequently shown that avirulence could be restored to isolate Race 5 on Cf-5 tomato, confirming identification of CfAvr5. Notably, this complementation also led to an increase in fungal biomass during infection of susceptible tomato, as compared to WT Race 5, thus highlighting


a yet unknown role for CfAvr5 in *F. fulva* virulence (Mesarich et al., 2014).

Shortly after the identification of CfAvr5, significant progress was made towards defining an inventory of all (candidate) effector proteins secreted by *F. fulva* into the leaf apoplast of susceptible tomato plants during infection (Mesarich et al., 2018). Using proteomics, a total of 75 *F. fulva* small, secreted proteins (SSPs) were identified in apoplastic washing fluid (AWF) samples collected from interactions between the fungus (including isolates OWU and Race 5) and susceptible tomato. Crucially, this comprised all *F. fulva* effectors characterized to date, as well as most of the candidate effectors (33 out of 44) encoded by genes examined in the CfAvr5 study. Based on the transcriptome sequencing data of Mesarich et al. (2014), 70 of the 75 SSP-encoding genes were shown to be up-regulated during host colonization (Mesarich et al., 2018).

Following on from this proteomics work, 41 of the 70 in planta-induced SSPs were screened for their ability to trigger a hypersensitive response (HR) across a collection of 25 predominantly wild resistant accessions of tomato, using an approach based on the potato virus X (PVX) expression system (Mesarich et al., 2018). Of these, nine SSPs (CfEcp8-16) triggered an HR in one or more accessions, suggesting that they are Avr proteins recognized by as yet uncharacterized Cf immune receptors. These accessions now represent a valuable resource for the development of new commercial tomato varieties with resistance against leaf mould disease (Mesarich et al., 2018).

One of the HR-eliciting SSPs, CfEcp11-1, has sequence similarity to the AvrLm3 and AvrLm5-9 Avr proteins from *Leptosphaeria maculans* (Ghanbarnia et al., 2018; Mesarich et al., 2018; Plissonneau et al., 2016, 2018; van de Wouw et al., 2014). While the function of CfEcp11-1 is not yet known, sequence and structural analyses determined that all three proteins, as well as two other Avr proteins from *L. maculans*, AvrLm4-7 and AvrLmS-Lep2 (Parlange et al., 2009; Xiang Neik et al., 2022), belong to a conserved group of fungal effectors, termed the “*Leptosphaeria* avirulence and suppressing” (LARS) family (Lazar et al., 2022). This family is characterized by a conserved WR(F/L/V)(R/K) motif that, when mapped onto the crystal structures of CfEcp11-1 (Figure 2a), AvrLm4-7, and AvrLm5-9, is predominantly surface-exposed, suggesting that it may play a role in functional interactions with host proteins. AvrLm4-7 is of particular interest, as it suppresses the recognition of AvrLm3 and AvrLm5-9 by the corresponding immune receptors Rlm3 and Rlm9 of *Brassica* spp., respectively (Ghanbarnia et al., 2018; Plissonneau et al., 2016). Strikingly, transformants of *L. maculans* expressing CfEcp11-1 also trigger Rlm3-mediated immunity in *Brassica* spp. Furthermore, as for AvrLm3 and AvrLm5-9, this recognition is suppressed by AvrLm4-7. These findings suggest that CfEcp11-1 is sufficiently similar to AvrLm3 to be directly or indirectly recognized by Rlm3 (Lazar et al., 2022).

In addition to CfEcp11-1, the crystal structures of the CfAvr4 and CfEcp6 effectors from *F. fulva* have been solved (Hurlburt et al., 2018; Sanchez-Vallet et al., 2013), while for many other effectors and effector candidates of this fungus, tertiary structures have



**FIGURE 1** Whole-genome comparison between *Fulvia fulva* isolate Race 5 and *Dothistroma septosporum* isolate NZE10. (a) Dot plots showing whole genome synteny between *F. fulva* and *D. septosporum*. The plots show a small number of interchromosomal translocations and many intrachromosomal rearrangements. Evidence of a reciprocal translocation between Chr5 and Chr13 in *D. septosporum* is highlighted in grey. (b) Circos plot showing the chromosomes of *D. septosporum* and *F. fulva* with tick marks indicating million base pairs. The other three outermost to innermost tracks represent gene density, repetitive DNA content, and GC content, respectively, within a sliding window of 30kb. Ribbons connect 17 genes encoding candidate effectors and three genes encoding key enzymes for secondary metabolism conserved in both genomes.

been predicted (Karimi-Jashni et al., 2022; Mesarich et al., 2018; Seong & Krasileva, 2022; Tarallo et al., 2022). CfAvr4, which contains a carbohydrate-binding module family 14 (CBM14) domain (van den Burg et al., 2004) and triggers immunity in tomato in the presence of the extracellular Cf-4 immune receptor (Joosten et al., 1997), binds to chitin present in the cell walls of invading *F. fulva* hyphae to protect them against hydrolysis by plant-derived chitinases (Joosten et al., 1994; van den Burg et al., 2003, 2006). The crystal structure of CfAvr4 revealed a dimeric assembly in which two molecules of

CfAvr4 interact to form a three-dimensional molecular sandwich that encapsulates two chitohexaose molecules (Figure 2b) (Hurlburt et al., 2018).

Like CfAvr4, CfEcp6 is a chitin-binding effector; however, CfEcp6 has three lysin motif (LysM) domains that sequester chito-oligosaccharides released from the cell walls of invading *F. fulva* hyphae to prevent their recognition by extracellular chitin-binding tomato immune receptors. This, in turn, prevents activation of chitin-triggered immune responses in tomato (Bolton et al., 2008;

de Jonge et al., 2010). Based on the crystal structure of CfEcp6 in complex with chitotetraose (Figure 2c), it was shown that this sequestration is mediated through the intrachain dimerization of LysM domains 1 and 3. Although not involved in sequestration, LysM domain 2 can also bind chitin, albeit with lower affinity, and has been shown to perturb chitin-triggered immune responses, potentially through interference with the chitin/immune receptor complex (Sanchez-Vallet et al., 2013).

Despite the progress made over the last 10 years in understanding the tertiary structure and function of specific *F. fulva* effectors, very little progress has been made on understanding the regulatory mechanisms underpinning *F. fulva* effector gene expression in planta. One study that attempted to understand this regulation focused on the transcription factor (TF) CfWor1 (Ökmen et al., 2014), as its

homologues in some other plant-pathogenic fungi were shown to regulate effector gene expression (Michielse et al., 2009). However, functional analyses, involving gene deletion and constitutive expression mutants, showed that CfWor1 is a global TF that regulates *F. fulva* development (Ökmen et al., 2014).

### 3.2 | Effector proteins of *D. septosporum*

Although *D. septosporum* and *F. fulva* have fairly dissimilar repertoires of (candidate) effector genes, many are still shared between the two species, consistent with their close phylogenetic relationship (Figure 1b) (de Wit et al., 2012; Mesarich et al., 2018; Stergiopoulos et al., 2012; Tarallo et al., 2022; Zaccaron et al., 2022). Comparison of their chromosomal locations showed that, in many cases, orthologous (candidate) effector gene pairs are present on homologous chromosomes in conserved locations or in locations where interchromosomal rearrangements have occurred (Figure 1b) (Zaccaron et al., 2022). It is likely that many of these orthologous genes encode “core” effectors with conserved biological functions in promoting host colonization (de Wit et al., 2012; Stergiopoulos et al., 2010). In support of this hypothesis, it has been shown that *DsAvr4*, the *D. septosporum* orthologue of *CfAvr4*, specifically binds to chitin and is also recognized by the Cf-4 immune receptor of tomato (de Wit et al., 2012), illustrating that *DsAvr4* and *CfAvr4* share a conserved biological function (Mesarich et al., 2016).

Another likely core effector from *D. septosporum* is *DsEcp2-1*, which is orthologous to *CfEcp2-1* from *F. fulva* (de Wit et al., 2012; van den Ackerveken et al., 1993). Both *CfEcp2-1* and *DsEcp2-1* were previously shown to trigger an HR in the presence of the Cf-Ecp2 immune receptor of tomato, which has not yet been identified (de Wit et al., 2012; Laugé et al., 2000; Stergiopoulos et al., 2010). Likewise, similar to *CfEcp2-1*, which triggers cell death in several *Nicotiana* species (de Kock et al., 2004; Laugé et al., 2000), *DsEcp2-1* was shown to trigger cell death in *Nicotiana tabacum* (Guo et al., 2020; Hunziker et al., 2021), suggesting that, in these nonhost plants, both of these proteins are recognized by an as yet uncharacterized

TABLE 3 Comparison of the genome annotation of *Fulvia fulva* isolate Race 5 and *Dothistroma septosporum* isolate NZE10.

Gene annotation statistics <sup>a</sup>	<i>F. fulva</i> Race 5	<i>D. septosporum</i> NZE10
Number of genes	14,690	12,415
Mean gene length (bp)	1284	1233
Mean exon length (bp)	598	585
Mean intron length (bp)	79	92
Mean exons per gene	2.1	2.1
Mean introns per gene	1.1	1.1
Total coding sequence length (bp)	18,870,333	15,318,053
Total intron length (bp)	1,341,392	1,270,355
BUSCO <sup>b</sup> completeness (%)	98.9	97.9
BUSCO duplicated (%)	0.1	0.1
BUSCO fragmented (%)	0.3	1.4
BUSCO missing (%)	0.8	0.5

<sup>a</sup>Untranslated regions were not considered when calculating the statistics.

<sup>b</sup>BUSCO, Benchmarking Universal Single-Copy Orthologs.




FIGURE 2 Solved tertiary structures of effector proteins from *Fulvia fulva*. Structures are visualized as cartoons using PyMol software (DeLano, 2002). Disulphide bonds are shown in yellow. (a) Crystal structure of the CfEcp11-1 effector protein (RCSB Protein Data Bank (PDB) ID: 6ZUS (Lazar et al., 2022)). The surface-exposed residues of the WRVR motif are shown as red sticks. (b) Crystal structure of the CfAvr4 effector protein (dimer) in complex with chitohexaose (PDB ID: 6BN0 (Hurlburt et al., 2018)). CfAvr4 molecules are coloured pink (left) and light blue (right), while chitohexaose is shown as green sticks. (c) Crystal structure of the CfEcp6 effector protein in complex with chitotetraose (PDB ID: 4B8V; Sanchez-Vallet et al., 2013). Lysin motif (LysM) domains 1, 2 and 3 of CfEcp6 are numbered and are coloured pink, light blue and cyan, respectively, while chitotetraose is shown as green sticks.

immune receptor. In *F. fulva*, *CfEcp2-1* is a virulence factor of unknown function, with deletion or disruption of the encoding gene resulting in reduced virulence (Laugé et al., 1997). Like *CfEcp2-1* (Wubben et al., 1994), *DsEcp2-1* is highly expressed in planta, suggesting that this effector may also function as a virulence factor during infection of pine trees (Guo et al., 2020). But strikingly, deletion of *DsEcp2-1* results in the production of larger necrotic lesions on pine needles by *D. septosporum*, concomitant with increased colonization, suggesting that *DsEcp2-1* is recognized as an Avr protein in pine by an as yet uncharacterized immune receptor, leading to partial resistance (Guo et al., 2020).

Recently, a set of 30 candidate apoplastic effector proteins, encoded by genes that are highly expressed and/or up-regulated during infection of pine seedlings by *D. septosporum* (Bradshaw et al., 2016), was identified (Hunziker et al., 2016, 2021). *Agrobacterium tumefaciens*-mediated transient transformation assays revealed that, in addition to *DsEcp2-1*, five of these proteins (*DsEcp20-3*, *DsEcp32-3*, *Ds71487*, *Ds131885*, and *Ds74283*) trigger cell death in *Nicotiana benthamiana* and/or in *N. tabacum* (Hunziker et al., 2021). As an extension of this work, additional members of the *Ecp20* and *Ecp32* families from *D. septosporum* (and their orthologues from *F. fulva*) were also screened and found to differ in their ability to trigger cell death in these two *Nicotiana* species (Tarallo et al., 2022). In cases where cell death was observed, this response was dependent on the secretion of these proteins to the apoplast, suggesting that they are recognized by extracellular immune receptors in these nonhost plants (Hunziker et al., 2021; Tarallo et al., 2022). In line with this hypothesis, cell death triggered by *DsEcp20-3* (and both *CfEcp20-3* and *CfEcp32-3*) was compromised in *N. benthamiana* plants silenced for *BAK1* or *SOBIR1*, which encode extracellular co-receptors involved in transducing defence response signals following apoplastic effector recognition (Tarallo et al., 2022). Notably, using a protein infiltration method involving clonal microshoots, it was shown that *DsEcp20-3* and *DsEcp32-3* also trigger cell death in three genotypes of the host plant *Pinus radiata*, suggesting that these proteins are recognized by extracellular immune receptors in this species (Hunziker et al., 2021; Tarallo et al., 2022). While not yet performed in *D. septosporum*, deletion of the *CfEcp20-2* gene in *F. fulva* isolate OWU resulted in reduced virulence on tomato, indicating that members of the *Ecp20* family are virulence factors (Karimi-Jashni et al., 2022).

### 3.3 | Glycoside hydrolases of *F. fulva* and *D. septosporum*

Fungal pathogens produce an array of carbohydrate-active enzymes (CAZymes) that are involved in the synthesis, metabolism, and recognition of complex carbohydrates (Bradley et al., 2022). Interestingly, it has recently been shown that fungal pathogens can be classified into different trophic lifestyles based on their complement of CAZyme-encoding genes (Hane et al., 2020). This is of particular relevance to *F. fulva*, which, although previously thought

to be a biotroph (de Wit et al., 2012), is now considered to be a hemibiotroph (Hane et al., 2020; Ökmen et al., 2019). Indeed, the complement of CAZyme-encoding genes in *F. fulva* is more similar to those of hemibiotrophs than biotrophs (de Wit et al., 2012; Hane et al., 2020). Such a classification is further supported by the observation that tomato leaves often become necrotic at the late stages of infection by *F. fulva* (Thomma et al., 2005).

Among the CAZymes, the secreted glycoside hydrolases (GHs) are central to understanding interactions between fungal pathogens and plants, with characterized or anticipated roles in fungal morphogenesis, nutrient acquisition, and host colonization (Bradley et al., 2022). Secreted GHs, however, can also be immunogenic in that they, or the products of their hydrolytic activity, called damage-associated molecular patterns (DAMPs), are recognized by extracellular plant immune receptors to activate the plant immune system (Bradley et al., 2022).

Over the past 10 years, only a handful of secreted GH proteins from *F. fulva* and *D. septosporum* have been studied. The first of these is *CfTom1*, a GH10 enzyme from *F. fulva* with no close homologues in *D. septosporum* (de Wit et al., 2012; Ökmen et al., 2013). *CfTom1* is poorly expressed during the early stages of tomato infection but is significantly up-regulated late in the infection cycle. Functional analysis showed that *CfTom1* is a tomatinase that detoxifies  $\alpha$ -tomatine, an antifungal glycoalkaloid present in the leaf apoplast of tomato, to the nontoxic compound tomatidine by removing the tetrasaccharide lycotetraose. Consistent with this role, *CfTom1* deletion mutants were more sensitive to  $\alpha$ -tomatine and produced less biomass during infection, indicating that *CfTom1* is required for full virulence of *F. fulva* on tomato plants (Ökmen et al., 2013).

Another characterized GH protein from *F. fulva* is *CfGH17-1*, an apoplastic GH17 enzyme with 1,3- $\beta$ -glucanase activity (Ökmen et al., 2019). Expression studies showed that *CfGH17-1* is down-regulated during the early stages of biotrophic infection by *F. fulva*, but is up-regulated late in the infection cycle, when the fungus switches to its necrotrophic state. Notably, *CfGH17-1* elicited cell death in several near-isogenic lines of tomato, but this cell death-eliciting activity was abolished by site-directed mutagenesis of the putative active-site residues present in this enzyme. These results suggest that *CfGH17-1* itself is not immunogenic but, rather, that *CfGH17-1* releases an immunogenic oligosaccharide from a complex polysaccharide or glycoprotein of the plant cell wall that is subsequently recognized as a DAMP by a yet unknown extracellular immune receptor present in tomato plants. In line with this hypothesis, deletion of the *CfGH17-1* gene does not affect symptom development by *F. fulva*, whereas overexpression results in reduced disease symptoms. One possibility is that *CfGH17-1* plays a role in nutrition, releasing sugar molecules from the host cell wall to support the growth and reproduction of *F. fulva* during the late stages of infection when the host tissue is necrotic and no longer capable of recognizing and responding to DAMPs. Interestingly, a paralogue from *F. fulva* (*CfGH17-5*), as well as the orthologue of *CfGH17-1* from *D. septosporum* (*DsGH17-1*), also trigger cell death when heterologously expressed in tomato

plants, suggesting that they might play a similar role in promoting host colonization (Ökmen et al., 2019).

Aside from DsGH17-1, other GH proteins from *D. septosporum* are also candidate virulence factors. One example is DsGH11 (Ds137959), a GH11 enzyme orthologous to an endo-1,4- $\beta$ -xylanase from *Z. tritici* (ZtXyl1) that has been suggested to play a role in lesion development during colonization of wheat (Bradshaw et al., 2016; Siah et al., 2010). In *D. septosporum*, DsGH11 is up-regulated in planta at the stage when disease lesions start to appear on infected pine needles, consistent with a role in lesion development and the switch from biotrophy to necrotrophy (Bradshaw et al., 2016).

### 3.4 | Proteases of *F. fulva* and *D. septosporum*

Proteases of *F. fulva* and *D. septosporum* are also anticipated to play a role in promoting host colonization. The number of protease-encoding genes present in the genome of *F. fulva* is once again more comparable to that of hemibiotrophs than biotrophs (Karimi Jashni et al., 2020). However, protease-encoding genes are more frequently pseudogenized in *F. fulva* than in *D. septosporum* (van der Burgt et al., 2014). Secreted proteases have not yet been studied in detail in *D. septosporum*, but in the *F. fulva* genome 59 of the 147 encoded proteases are predicted to be secreted (Karimi Jashni et al., 2020). Of these, 18 are expressed both in culture and in planta, suggesting that a specific role in virulence for these 18 proteases is unlikely. However, a few genes, such as those encoding the predicted serine proteases CfPro41 and CfPro54, are specifically induced during host infection, while another gene encoding the predicted metalloprotease CfPro19 is more highly expressed in planta than in culture (Karimi Jashni et al., 2020). One possibility is that these proteases promote virulence by removing the chitin-binding domain from tomato chitinases, which decreases their antifungal activity, as was shown for a synergistically acting metallo-/serine-protease pair, FoMep1/FoSep1, from *Fusarium oxysporum* f. sp. *lycopersici* (Jashni et al., 2015).

## 4 | SECONDARY METABOLISM

### 4.1 | Core secondary metabolite genes of *F. fulva* and *D. septosporum*

Fungal secondary metabolites (SMs) play key roles in development, communication, microbial antagonism, and plant disease (Macheleidt et al., 2016). SMs are classified into chemical groups based on their backbone structure, the biosynthesis of which is catalysed by core enzymes, including polyketide synthases (PKSs), nonribosomal peptide synthetases (NRPSs), hybrid PKS-NRPSs, terpene cyclases (TCs), and dimethylallyltryptophan synthases (DMATs) (Macheleidt et al., 2016). Efforts to identify and characterize SM genes in *F. fulva* and *D. septosporum* were greatly

accelerated by analysis of their genomes (de Wit et al., 2012). However, growth of *F. fulva* or *D. septosporum*, either in planta or in culture, yields only one main detectable SM from each species, namely cladofulvin and dothistromin, respectively. The genome of *F. fulva* OWU initially suggested a large chemical diversity with 10 PKS, 10 NRPS, two hybrid PKS-NRPS, and one DMAT core genes, but seven of these genes are truncated or pseudogenized. The *D. septosporum* NZE10 genome contains only five PKS, three NRPS, two hybrid PKS-NRPS, and one DMAT core genes, of which two are truncated or pseudogenized (de Wit et al., 2012). The only three core SM genes shared between *F. fulva* Race 5 and *D. septosporum* NZE10 are located on homologous chromosomes (Figure 1b). These shared genes are Cf/DsPks1 involved in the biosynthesis of DHN melanin, Cf/DsNps2 involved in siderophore production, and Cf/DsPksA involved in dothistromin biosynthesis (Bradshaw et al., 2006; Collemare et al., 2014; de Wit et al., 2012; Ozturk et al., 2017, 2019). Finally, expression analysis showed that, in both species, only a few SM core genes are expressed in planta or under diverse conditions in culture (Bradshaw et al., 2016; Collemare et al., 2014; Griffiths et al., 2015; Ozturk et al., 2019), with CfPks1 and DsPks1 being expressed in both species (Collemare et al., 2014; de Wit et al., 2012; Ozturk et al., 2017). Three additional SM core genes are expressed in *D. septosporum* (DsPksA, DsPks2, and DsNps3) (Bradshaw et al., 2016; Ozturk et al., 2017, 2019) and five in *F. fulva* (CfNps4, CfNps8, CfNps9, CfPks5, and CfPks6) (Table 4) (Collemare et al., 2014; de Wit et al., 2012).

### 4.2 | Functional analysis of SM genes

Functional analysis has been carried out for selected SM genes in *F. fulva* and *D. septosporum* to determine their roles. Deletion studies showed that neither of the core CfPks1 and DsPks1 dihydroxynaphthalene (DHN) melanin biosynthesis genes are pathogenicity or virulence factors, consistent with their low expression in planta (Bradshaw et al., 2016; Griffiths, Cox, et al., 2018; Ozturk et al., 2017). Further studies revealed that *D. septosporum* appears to produce DHN melanin mostly through an alternative PKS-independent L-3,4-dihydroxyphenylalanine (DOPA) pathway (Ozturk et al., 2017), whilst in *F. fulva*, DHN melanin biosynthesis required CfPks1 (Griffiths, Cox, et al., 2018).

In *F. fulva*, deletion of the CfPks6 gene that is required for cladofulvin biosynthesis yielded mutants that were unaltered in pathogenicity or virulence on tomato plants (Griffiths, Mesarich, et al., 2018). CfPks6 is normally only lowly expressed during growth of the fungus inside the apoplast. However, transformants overexpressing the presumed transcriptional activator (CfClaE) from the CfPks6 cluster induced rapid necrosis of tomato leaves and were arrested in growth. Together, these results suggest that *F. fulva* must repress cladofulvin production during early host colonization to sustain biotrophic growth prior to a necrotrophic switch (Griffiths, Mesarich, et al., 2018). The cladofulvin pathway of *F. fulva* was eventually deduced by analysis of metabolites produced by transformants of *Aspergillus oryzae* expressing CfPks6




Core gene	Predicted SM or family	In planta expression <sup>a</sup>	Virulence function	Reference
<i>F. fulva</i>				
<i>CfPks1</i>	Melanin; DHN	Late	No	Griffiths, Cox, et al. (2018)
<i>CfPks5</i>	T-toxin	No	NT	Collemare et al. (2014)
<i>CfPks6</i>	Cladofulvin	Early and late	No <sup>b</sup>	Collemare et al. (2014); Griffiths, Mesarich, et al. (2018)
<i>CfNps4</i>	"EAS (Euascomycete) clade"	No	NT	Collemare et al. (2014)
<i>CfNps8</i>	Siderophore	No	NT	Collemare et al. (2014)
<i>CfNps9</i>	Siderophore	Early	NT	Collemare et al. (2014)
<i>D. septosporum</i>				
<i>DsPksA</i>	Dothistromin	Mid-late	Yes	Kabir et al. (2015a)
<i>DsPks1</i>	Melanin; DHN	Late	No	Ozturk et al. (2017)
<i>DsPks2</i>	Squalestatin-like	Late	No	Ozturk et al. (2017)
<i>DsNps3</i>	Beauvericin-like	Early	Yes	Ozturk et al. (2019)

Abbreviations: DHN, dihydroxynaphthalene; NT, not tested.

<sup>a</sup>In *D. septosporum*, highest gene expression occurred during early (asymptomatic), mid (early lesion) or late (sporulating lesion) stages of disease on *Pinus radiata* needles. In *F. fulva*, highest gene expression was during early (surface growth of runner hyphae 0–4 days after inoculation) or late (conidiation 10–16 days after inoculation) stages on tomato.

<sup>b</sup>Gene deletion mutants were not obviously altered in virulence, but *CfClaE* overexpression transformants were hypervirulent.



**FIGURE 3** Locations of predicted dothistromin biosynthetic pathway genes in *Fulvia fulva* isolate Race 5 and *Dothistroma septosporum* isolate NZE10. Circos plot showing Chr12 of *D. septosporum* and Chr13 of *F. fulva*. From outermost to innermost, tracks represent the chromosomes, gene density, repetitive DNA content, and GC content within a sliding window of 30 kb. Ribbons connect 19 predicted dothistromin genes, which are colour-coded based on their respective subcluster (numbered 1 to 6) in *D. septosporum* (Chettri et al., 2013). All predicted homologues of the dothistromin gene cluster in *F. fulva* are located on Chr13, with all subclusters conserved. The only exception is for *CfDotC*, which is physically distant from *CfVer1* and *CfDotB*. The genes *CfHexB* and *CfAvnA* are fused in the same predicted gene model.

along with the early (*CfClaF* and *CfClaH*) or middle (*CfClaB* and *CfClaC*)-acting genes, as well as by analysis of a *F. fulva* deletion mutant ( $\Delta$ *CfClaM*) that was blocked in the final step of cladofulvin biosynthesis (specifically, the dimerization of the monomeric anthraquinone, nataloe-emodin) (Griffiths et al., 2016).

In *D. septosporum*, deletion of the *DsNps3* (beauvericin-like) gene led to reduced virulence on *P. radiata* but further work is needed to elucidate the role of this potential virulence factor (Ozturk et al., 2019). Deletion of the *DsPks2* (squalestatin-like) gene, however, did not affect virulence (Ozturk et al., 2017).

### 4.3 | Genetics and evolution of dothistromin biosynthesis

The *D. septosporum* *DsPksA* gene is required for the production of dothistromin, an aflatoxin-like anthraquinone (Bradshaw et al., 2006) that was confirmed to be required for full virulence on pine trees (Kabir et al., 2015a). Whilst other genes involved in the biosynthesis of an SM are frequently clustered around the core gene (Macheleidt et al., 2016), the chromosome-level assembly of the *D. septosporum* NZE10 genome sequence revealed that dothistromin pathway genes are in six subclusters spread across chromosome 12 (de Wit et al., 2012). The complete set of dothistromin genes is conserved in *F. fulva* Race 5 (Zaccaron et al., 2022) and is similarly fragmented into subclusters throughout a single chromosome, Chr 13, but with some rearrangements (Figure 3). Dothistromin was not detected in *F. fulva* and genes in the pathway are barely expressed (de Wit et al., 2012). Further analysis revealed that dothistromin biosynthesis in this species is blocked by pseudogenization of early pathway genes, *CfHexA* and *CfNor1* (Chettri et al., 2013). In *D. septosporum*, dothistromin biosynthesis is concomitant with the start of the rapid increase in fungal biomass that occurs during the transition from biotrophic to necrotrophic growth, consistent with its role in expansion of disease lesions (Kabir et al., 2015a, 2015b).

Orthologues of some dothistromin genes are also present in other fungi, including *Rhytidhysterion rufulum*, a saprophytic dothideomycete (Bradshaw et al., 2013; Ohm et al., 2012), and some eurotiomycete species such as *Aspergillus flavus* and *Aspergillus nidulans*, in which the dothistromin orthologues are involved in aflatoxin and sterigmatocystin biosynthesis, respectively (Bradshaw et al., 2013). A comparative genomics and population genetics study of dothistromin and aflatoxin-like genes in these two fungal classes suggested that the dispersed arrangement of dothistromin genes in *D. septosporum* evolved from an ancestral gene cluster, leading to the suggestion that cluster fragmentation might facilitate chemical diversity and provide an adaptive advantage (Bradshaw et al., 2013).

### 4.4 | Global regulation of SM gene expression

Due to the large number of cryptic SM pathways in *F. fulva* and *D. septosporum*, there has been considerable interest in the regulation of SM gene expression. Attempts to activate cryptic SM pathways in *F. fulva* by chemical induction (trichostatin A and 5-azacytidine), or by culturing under different conditions, were unsuccessful (Griffiths et al., 2015). Manipulation of conserved global transcriptional regulators known to affect SM gene expression in other ascomycetes provided another avenue for exploration because these regulators coordinate many gene networks and their manipulation can yield phenotypes that are useful for discovering new SMs or understanding their functions. Both *F. fulva* and *D. septosporum* were previously shown to have genes encoding the global regulators *Cf/DsHdaA* (histone deacetylase), *Cf/DsVeA* (velvet complex scaffold), and *Cf/DsLaeA* (methyltransferase), among others (de Wit et al., 2012).

In *F. fulva*, deletion mutants of the transcriptional regulator genes *CfLaeA* and *CfVeA* were impaired in conidiation but still produced cladofulvin. However, the loss of hyphal pigmentation associated with melanin in  $\Delta$ *CfLaeA* mutants suggested that *CfLaeA* positively regulates melanin production (Griffiths et al., 2015). In contrast, *CfHdaA* mutants do not produce cladofulvin (Griffiths et al., 2015).

In *D. septosporum*, *DsHdaA*, *DsVeA*, and *DsLaeA*, as well as the pathway regulator *DsAflR*, were each shown to play a role in the regulation of dothistromin biosynthesis (Chettri & Bradshaw, 2016; Chettri et al., 2012, 2013, 2018). Gene clustering has been proposed to facilitate coordinated gene expression through control of chromatin structure (Macheleidt et al., 2016). However, co-expression of dothistromin genes in the fragmented dothistromin gene cluster is achieved by both direct chromatin-level H3K27me3 control of the dispersed genes as well as indirect H3K9me3/ac control via *DsAflR* (Chettri et al., 2018).

## 5 | INDUCTION AND CIRCUMVENTION OF DEFENCE RESPONSES IN TOMATO BY *F. FULVA*

### 5.1 | Interactions of *F. fulva* effectors with Cf immune receptors and host virulence targets

Resistance to *F. fulva* in tomato is mediated by cell surface-associated receptor-like proteins (RLPs), referred to as Cf immune receptors. The extracellular domain of Cf immune receptors mostly consists of leucine-rich repeats (LRRs), which are implicated in protein-protein interactions. These RLPs allow tomato plants that are resistant to *F. fulva* to recognize specific secreted effector proteins from the fungus, thereby rendering it an Avr protein. Whether this recognition takes place directly, or indirectly through perception of the manipulated virulence target of the host by the effector, is still an area of intense study.

One of the best-studied Avr proteins from *F. fulva* is *CfAvr2*, a protease inhibitor that interacts with the secreted papain-like cysteine protease of tomato, *Rcr3*. In susceptible host genotypes, *CfAvr2* binds to, and inhibits, *Rcr3* to promote host colonization. However, in resistant tomato genotypes, the *CfAvr2/Rcr3* complex is recognized by the Cf-2 immune receptor to elicit plant immune responses (Rooney et al., 2005). This indirect recognition of *CfAvr2* through the guarded virulence target *Rcr3* is an example of the guard hypothesis, in which the immune receptor (the guard, Cf-2) senses manipulation of a virulence target (the guardee, *Rcr3*) by an effector (Dangl & Jones, 2001; van der Biezen & Jones, 1998). *CfAvr2* behaves as a noncompetitive inhibitor of *Rcr3*. Affinity of *CfAvr2* mutants for *Rcr3* is positively correlated with their ability to trigger a Cf-2-mediated HR (Van't Klooster et al., 2011).

Immune receptor genes are continuously subjected to selective pressure, resulting in high genetic variation and novel effector recognition specificities. Interestingly, the *Rcr3* protease that is guarded by Cf-2 is also subject to selective pressure. Indeed, natural

variants of this protease have been identified, with these variants exhibiting differences in their interactions with CfAvr2 that correlate with variation in immune output (Hörger et al., 2012). In addition, the evolution of Rcr3 is further impacted by its interaction with the Cf-2 immune receptor, and three amino acid polymorphisms in Rcr3 appear to have a major role in co-evolution between this guard and the Cf-2 guard. Based on such observations, it was concluded that, as a result of natural selection, there is a subtle balance between the initiation of a Cf-2-triggered resistance response in the presence of CfAvr2 and the prevention of spontaneous Rcr3/Cf-2-triggered auto-immune responses in the absence of CfAvr2 (Hörger et al., 2012).

In addition to Rcr3, tomato secretes a very closely related protease, Pip1. Notably, CfAvr2 can inhibit both proteases but only Rcr3 is guarded by Cf-2 (Ilyas et al., 2015). Pip1 is up to 10-fold more abundant in tomato than Rcr3 and tomato plants with reduced *Pip1* expression are hypersusceptible to a broad range of pathogens, suggesting that Pip1 is an important protease mediating basal immunity. Strikingly, *Rcr3* knock-out lines do not show significantly higher levels of infection by bacteria and fungi but do show increased susceptibility to the oomycete *Phytophthora infestans*. It has been proposed that gene duplication resulted in Rcr3 and Pip1 being functionally diversified and that co-evolution between Cf-2 and Rcr3 resulted in the indirect detection of Rcr3 inhibitors, such as CfAvr2 (Ilyas et al., 2015). This work suggests that Rcr3 acts as a guarded “decoy” that betrays strains of *F. fulva* secreting Avr2 on their ingress into leaves of Cf-2-containing tomato plants.

A further study revealed that Rcr3 evolved more than 50 million years ago, whereas Cf-2 is evolutionarily much younger (Kourelis et al., 2020). In various solanaceous plants, there are ancient homologues of Rcr3 that can be inhibited by CfAvr2, with four residues in these proteins that vary among the solanaceous species being able to promote this inhibition. Strikingly, the isoform of Rcr3 that is guarded by Cf-2 lacks three of these residues, which suggests that Rcr3 is employed as a type of co-receptor by Cf-2 that is inferior in binding to CfAvr2 (Kourelis et al., 2020).

Interestingly, the P69 protease family of tomato consists of a set of subtilases (SBTs) that proteolytically activate Rcr3 (Paulus et al., 2020; Reichardt et al., 2018). It was observed that Rcr3 is processed by secreted P69B and other SBTs in solanaceous plants. SBTs are aspartic acid-selective proteases, meaning that they cleave amino acid chains in Asp-rich regions. Asp residues are abundantly present in the region where Rcr3-like proteases are cleaved, which suggests that in solanaceous plants the maturation and activation of immune proteases by subtilases generally occurs and supports immunity in the extracellular spaces, contributing to the first line of defence against pathogens such as *F. fulva* (Paulus et al., 2020).

Another well characterized Avr protein of *F. fulva* is CfAvr4, which, as mentioned above, binds to chitin in the cell walls of invading hyphae of the fungus to protect them against hydrolysis by secreted tomato chitinases (van den Burg et al., 2006). As CfAvr4 does not seem to have a virulence target in the plant, a direct interaction between CfAvr4 and the LRRs of Cf-4 is expected. In addition

to DsAvr4 from *D. septosporum*, several orthologues of CfAvr4 have been identified from plant-pathogenic dothideomycete fungi, including CaAvr4 from *Cercospora apii* and CbAvr4 from *Cercospora beticola*. While each of these orthologues binds chitin, only DsAvr4 appears to be recognized by Cf-4. Using protein chimeras and site-directed mutants involving CfAvr4, DsAvr4, CaAvr4, and CbAvr4, a particular conserved proline residue was shown to be required for Avr4 proteins to initiate a Cf-4-dependent HR (Mesarich et al., 2016). The mechanism behind this proline requirement is not known; however, subsequent work suggested that the proline residue itself is not essential, as long as a small aliphatic amino acid residue is present at this position (Hurlburt et al., 2018). In general, mutations in CfAvr4 appear to reduce its stability in the apoplastic environment. It is therefore likely that the small aliphatic residue is required to maintain the structural integrity of the protein, and that this structural integrity is needed for Cf-4 to perceive the globular fold of CfAvr4 and thereby activate host immunity. Crucially, CfAvr4 recognition by Cf-4 is not linked with its chitin-binding properties, meaning that the virulence function of CfAvr4 is not coupled to its ability to be recognized by Cf-4 (Hurlburt et al., 2018).

## 5.2 | Characterization and mapping of novel Cf immune receptor genes

As broad-spectrum resistance against leaf mould disease, to our knowledge, remains elusive in commercial tomato varieties, the last decade has seen significant efforts made in the mapping, identification, and characterization of additional Cf immune receptor genes (Chai et al., 2020; Iakovidis et al., 2020; Jiang et al., 2022; Liu et al., 2018, 2019, 2022; Xu et al., 2018; Xue et al., 2017; Zhang et al., 2020; Zhao et al., 2015, 2016, 2019). For example, the Cf-3 and Cf-11 immune receptor genes, which correspond to the yet-unidentified CfAvr3 and CfAvr11 Avr genes of *F. fulva*, have recently been mapped to a narrow region on chromosome 11 and have been shown to be closely linked (Iakovidis et al., 2020). Interestingly, the target region seems to be devoid of genes encoding RLPs, which suggests that Cf-3 and Cf-11 mediate resistance using a different class of immune receptors.

Little is known about the distribution and presence of Cf immune receptor genes across wild populations of tomato species. Previously, it was shown that homologues of Cf-4 and Cf-9 (*Hcr9s*) are found across the *Solanum* genus and probably pre-date speciation (Kruijt et al., 2005). More recently, *Solanum chilense* accessions originating from the southern part of Peru to the northern part of Chile were screened for their ability to recognize *F. fulva* effectors by infiltrating AWF collected from infected plants (Kahlon et al., 2020). This showed that recognition is mostly lost in southern accessions but retained in northern accessions. Interestingly, whilst Cf-4 and Cf-9 were previously thought to be mutually exclusive alleles (Parniske et al., 1997; Thomas et al., 1997), some *S. chilense* accessions possessed loci associated with both Cf-4 and Cf-9 and consequently recognized both CfAvr4 and CfAvr9 (Kahlon et al., 2020). Such

accessions could prove valuable for introgression into commercial varieties, as the simultaneous presence of both Cf-4 and Cf-9 was previously not thought possible. Finally, in several accessions that do not respond to CfAvr9, there is still expression of Cf-9. This lack of recognition might be explained by inactive polymorphic versions of Cf co-receptors such as BAK1 and SOBIR1. The impact of these inactive versions on CfAvr9 recognition, however, remains to be determined (Kahlon et al., 2020).

More recently, a study on the distribution of the Cf-Ecp5 immune receptor gene (corresponding to the CfEcp5 effector of *F. fulva*) among accessions from various *Solanum* species revealed at least three distinct loci on chromosomes 1, 7, and 12 in four different accessions that harbour functional Cf-Ecp5 alleles (Iakovidis et al., 2020). This includes the previously identified Milky Way cluster on chromosome 1 (Haanstra et al., 2000). Unexpectedly, a seemingly unlinked genetic suppression of CfEcp5 recognition was observed in some biparental crosses with accessions carrying distinct Cf-Ecp5 loci (Iakovidis et al., 2020). This suppression could possibly also be caused by mutations in unlinked genes coding for co-receptors of Cf-Ecp5 or by other pleiotropic effects.

### 5.3 | Immune signalling in response to *F. fulva* effector recognition

In resistant tomato, effector perception triggers an immune response that is typically associated with an HR. In contrast to RLKs, RLPs lack a cytoplasmic kinase domain that can act as a signalling moiety. Therefore, to enable signalling, Cf immune receptors (and many other RLPs) constitutively interact with the regulatory RLK SOBIR1 (Liebrand et al., 2013), which provides the RLP with a cytoplasmic kinase domain. In this way, a bipartite RLK is generated that, on effector perception by the apoplastic LRRs of the RLP, can pass on the immune signal to the cytoplasm (Liebrand et al., 2014).

Notably, as is the case for LRR-RLKs involved in immunity, the bipartite Cf/SOBIR1 complex specifically recruits the LRR-RLK BAK1 on its activation by an Avr effector matching the Cf immune receptor (Postma et al., 2016). When Cf-4 is challenged with CfAvr4, the immune response triggered on formation of the Cf-4/SOBIR1/BAK1 heterotrimer is associated with rapid endocytosis of the complex (Postma et al., 2016) and transphosphorylation events between BAK1 and SOBIR1 (van der Burgh et al., 2019). This observation, along with the requirement of SOBIR1 kinase activity for Cf-4-mediated HR (Liebrand et al., 2013), suggests that, on perception of CfAvr4, SOBIR1 also facilitates the first steps in downstream signalling. The ultimate proof that SOBIR1 is essential for functionality of Cf-4, and probably all LRR-RLPs in general that are involved in plant immunity, was obtained by knocking-out the SOBIR1 gene in stable Cf-4-expressing *N. benthamiana* plants, using CRISPR/Cas technology (Huang et al., 2021). A swift ROS burst is a typical reflection of the Cf-4-triggered immune output and, strikingly, this response

was completely absent in Cf-4-expressing *N. benthamiana* plants lacking a functional SOBIR1 protein. Furthermore, the HR was absent in these *sobir1* knockout plants on challenge with CfAvr4 (Huang et al., 2021).

Information on signalling candidates that function directly downstream of the active Cf-4/SOBIR1/BAK1 complex is scarce. However, Cf-9 requires ACIK1, a class VII tomato receptor-like cytoplasmic kinase (RLCK), for mediating full resistance against CfAvr9-secreting isolates of *F. fulva* (Rowland et al., 2005). Current studies are focusing on whether class VII RLCKs also play a role in propagating the immune response downstream of Cf-4.

Earlier work revealed that the “helper NLR” NRC1 is involved in mounting the Cf-mediated immune response in tomato, as this protein is required for full resistance of Cf-4 tomato to CfAvr4-secreting isolates of *F. fulva* (Gabriëls et al., 2007). The mechanism by which NRC1 contributes to immunity initiated by Cf-4 remained enigmatic until recently when a close homologue of NRC1 in *N. benthamiana*, NRC3, was shown to actually mediate the HR. NRC3 contains an N-terminal coiled-coil (CC) domain, including a so-called “MADA” motif, which is essential for assembly of a resistosome that has the capacity to trigger cell death (Kourelis et al., 2022).

### 5.4 | Circumvention of Cf-mediated resistance in tomato by *F. fulva*

Since 2012, a number of *F. fulva* races have been collected from around the world, including Argentina, China, Croatia, Cuba, Korea, Turkey, and the United States, that have overcome resistance provided by one or more of the Cf-2, Cf-4, Cf-4E, Cf-5, or Cf-9 immune receptor genes in commercial tomato varieties (Altin, 2016; Bernal-Cabrera et al., 2021; Iida et al., 2015; Kubota et al., 2015; Lee et al., 2013; Li et al., 2015; Lucentini et al., 2021; Medina et al., 2015; Mesarich et al., 2014; Novak et al., 2021; Rollan et al., 2013; Sudermann et al., 2022; Yoshida et al., 2021). Analysis of the allelic variation present in the matching Avr genes has revealed many new mutations, in addition to mutations that had previously been characterized (Stergiopoulos, De Kock, et al., 2007), that enable these effectors to escape recognition by their cognate immune receptors (Table 5). Such information will enable the presence and migration of *F. fulva* races to be tracked, which may ultimately play an important role in cultivar selection and deployment.

Recently, the classification system for defining the resistance spectrum of tomato varieties against *F. fulva* has been updated by the International Seed Federation (ISF), and the differential set of tomato lines harbouring the Cf-2, Cf-4, Cf-5, or Cf-9 gene or combinations thereof has been extended with a line harbouring the Cf-6 immune receptor gene, corresponding to the yet unidentified CfAvr6 Avr gene of *F. fulva* (Grushetskaya et al., 2007). This inclusion suggests that breeding companies have (recently) started to deploy Cf-6 in newly developed tomato varieties. More interestingly, a new race of *F. fulva* originating from South Korea has now been included in the differential set of isolates used for classifying the resistance

TABLE 5 Recently identified allelic variation that alters the sequence of avirulence (Avr) proteins from *Fulvia fulva*.

Avr gene (accession number <sup>a</sup> )	Polymorphism at DNA level	Polymorphism at protein level <sup>b</sup>	Loss of Cf-mediated HR	Country	Reference	
CfAvr2 (AJ421628)	Gene deletion	No protein	Yes	Argentina, USA	Medina et al. (2015); Sudermann et al. (2022)	
	c.1A>G	LRG_199p1:p.Met1?	Yes	Japan	lida et al. (2015)	
	<b>c.2 T&gt;C</b>	<b>LRG_199p1:p.Met1?</b>	Yes	USA	Sudermann et al. (2022)	
	c.28T>G	p.Trp10Gly	No	Argentina	Lucentini et al. (2021)	
	c.29G>T	p.Trp10Leu	No	Argentina	Lucentini et al. (2021)	
	c.46del.G	p.Glu16Lysfs*2	Yes	USA	Sudermann et al. (2022)	
	c.50ins.T	p.Ile18Asnfs*25	Yes	Japan	lida et al. (2015)	
	c.52A>C	p.Ile18Leu	No	Japan	lida et al. (2015)	
	c.56del.CAGCAGCCAA	p.Ala19Glu fs*13	Yes	Japan	lida et al. (2015)	
	<b>c.(64–69)del.A</b>	<b>p.Lys23Asnfs*12</b>	Yes	USA	Sudermann et al. (2022)	
	<b>c.(64–69)ins.A</b>	<b>p.Leu24Thrfs*19</b>	Yes	Japan	lida et al. (2015)	
	<b>c.(117_119)ins.A</b>	<b>p.Tyr41Valfs*2</b>	Yes	Croatia	Novak et al. (2021)	
	c.121_122del.TA	p.Tyr41Ilefs*1	Yes	USA	Sudermann et al. (2022)	
	c.209ins.A	p.His70Glnfs*?	Yes	USA	Sudermann et al. (2022)	
	c.188G>T	p.Cys63Phe	Yes	Japan	lida et al. (2015)	
CfAvr4 (Y08356)	Transposon insertion	No protein	Yes	Japan	lida et al. (2015); Yoshida et al. (2021)	
	Large del. after c.37	del. after p.Leu12	Yes	USA	Sudermann et al. (2022)	
	c.118T>C	p.Cys40Arg	Yes	Japan	lida et al. (2015)	
	c.191G>T	p.Cys64Phe	Yes	Japan	lida et al. (2015)	
	c.191G>C	p.Cys64Ser	Yes	Japan	lida et al. (2015); Yoshida et al. (2021)	
	<b>c.191G&gt;A</b>	<b>p.Cys64Tyr</b>	Yes	Japan	lida et al. (2015)	
	c.318del.G	p.Ser107Valfs*5	Yes	Japan	lida et al. (2015)	
	<b>Gene deletion</b>	<b>No protein</b>	Yes	Japan	Yoshida et al. (2021)	
	<b>c.244T&gt;C</b>	<b>p.Phe82Leu</b>	Yes	Japan, Argentina, USA	lida et al. (2015); Lucentini et al. (2021); Sudermann et al. (2022)	
	<b>c.278T&gt;C</b>	<b>p.Met93Thr</b>	Yes	Japan, Argentina, USA	lida et al. (2015); Lucentini et al. (2021); Sudermann et al. (2022)	
	CfAvr5 (KJ452245)	Gene deletion	No protein	Yes	Japan	lida et al. (2015)
		<b>c.11ins.TC</b>	p.Ile5Leufs*3	Yes	France	Mesarich et al. (2014)
		c.268G>C	p.Gly90Arg	Yes	Japan	lida et al. (2015)
	CfAvr9 (X60284)	<b>Gene deletion</b>	<b>No protein</b>	Yes	Japan, Cuba	Bernal-Cabrera et al. (2021); Yoshida et al. (2021)
		<b>c.23T&gt;C</b>	<b>p.Val8Ala</b>	No	Japan, USA	Sudermann et al. (2022)

Abbreviations: c, numbering based on cDNA sequence; del, deletion; fs, frameshift; HR, hypersensitive response; ins, insertion; p, numbering based on protein sequence; \*, termination codon.

<sup>a</sup>National Center for Biotechnology (NCBI) GenBank accession number.

<sup>b</sup>Mutations in bold were previously described by Stergiopoulos, De Kock, et al. (2007). The description of each mutation is based on the nomenclature set forth by the human genome variation society (HGVS) (<http://varnomen.hgvs.org/>).

spectrum of tomato varieties, indicating that Cf-6 has already been overcome in the field (Sangster, 2022).

## 6 | PROGRESS IN UNDERSTANDING *D. SEPTOSPORUM*, DOTHISTROMA NEEDLE BLIGHT, AND PINE DEFENCE

### 6.1 | Defence responses in pine

Although host defences in the *D. septosporum*–pine pathosystem have not been studied to the same level of molecular detail as for those for *F. fulva* and tomato, some recent progress has been made. Such studies were prompted by outbreaks of Dothistroma needle blight (DNB), such as those that caused defoliation and mortality on lodgepole pine trees (*Pinus contorta*) in north-western British Columbia in Canada (Welsh et al., 2009; Woods, 2003). In the first study of its kind to investigate the molecular resistance mechanism(s) of pine to *D. septosporum*, Lu et al. (2021) performed controlled experiments by inoculating tolerant (partially resistant) and susceptible pines with two *D. septosporum* isolates from different genetic lineages (Capron et al., 2021). Subsequent analysis of differentially expressed genes (DEGs) showed a rapid and strong transcriptomic response in the tolerant pines, and a late and weak response in the susceptible pines (Lu et al., 2021). DEGs identified from tolerant pines included those involved in extracellular and intracellular immunity, such as genes encoding calcium-dependent protein kinases, BAK1, WRKY transcription factors, and EDS1. Based on this study, the authors propose that lodgepole pine recognizes *D. septosporum* attack through immune receptors, which then trigger a series of downstream signalling responses, including a Ca<sup>2+</sup> burst and the activation of a mitogen-activated protein kinase (MAPK) cascade, that lead to cell wall reinforcement, stomatal closure, and an HR. Nine candidate immune receptor genes were also identified that contained sites under positive selection. These sites are located close to regions in the genes that encode conserved domains like the NB-ARC domain, LRRs, or the coiled-coil (CC) domain. Comparison of expression profiles between inoculated and control samples showed that these candidate immune receptor genes were differentially regulated, suggesting differential responses to the pathogen (Lu et al., 2021). Further experiments are needed to identify corresponding effectors and to confirm the functions of these candidate immune receptor genes.

### 6.2 | Growth of a *Dothistroma* research community

Research on DNB has not only been accelerated by the availability of the *D. septosporum* genome sequence, but also by the continual improvement of molecular tools for diagnostics and population genetics (Barnes et al., 2008; Groenewald et al., 2007; Iosifidis et al., 2010). In response to the increased spread and severity of DNB in Europe

(Drenkhan et al., 2016), an international consortium of researchers from 41 countries was developed through a European COST (Cooperation in Science and Technology) Action (2011–2015) that included training in the use of molecular tools and culminated in publication of a special issue of *Forest Pathology* on DNB (Bradshaw, 2016). The special issue contained 14 articles, including reviews on the host range and distribution of *D. septosporum* and its lesser known sister species *D. pini*, the role of climatic factors, disease management, and Pinaceae resistance mechanisms (Barnes et al., 2016; Bulman et al., 2016; Drenkhan et al., 2016; Fraser et al., 2016; Woods et al., 2016). Many of these collaborations continue with a particular emphasis on genomic and population genetic studies.

### 6.3 | Genomics of other *D. septosporum* isolates

Next to the genome sequence of *D. septosporum* isolate NZE10, several additional genome sequences have been obtained from various isolates of this pathogen since 2012. Among these are isolates from the New Zealand population of *D. septosporum*, which is almost clonal, with very low diversity (Hirst et al., 1999), and is only of one mating type (Barnes et al., 2014). The availability of isolates collected in the 1960s, soon after the arrival of *D. septosporum* in New Zealand, provided a unique opportunity to determine how the pathogen co-evolved with its host over 50 years in pine plantation forests (Bradshaw, Ormond, et al., 2019). In general, genome sequences of isolates collected over the 50-year period showed low diversity, but, as expected, some diversification over time as well. However, isolates collected in the 1960s were more virulent on *P. radiata* and more potent producers of dothistromin, as compared to those collected from the 1990s onwards. This suggested adaptation towards decreased virulence over time, although no specific changes in the genome sequences could be found that correlated with this phenomenon (Bradshaw, Ormond, et al., 2019).

Genome sequences were also obtained for a global collection of 18 *D. septosporum* isolates from Europe, Africa, Australasia, and the American continent (Bradshaw, Sim, et al., 2019). Several of these isolates appeared to be aneuploid, exhibiting whole or large segmental duplications of particular chromosomes. Included in these isolates was an isolate from the German Alps, which produced significantly higher levels of dothistromin as compared to other isolates, a phenotype that was attributed to the increased copy number of many genes in chromosome duplication regions unique to this isolate (Bradshaw, Sim, et al., 2019).

Analysis of genome sequences from isolates collected from Scotland and western Canada, two regions that experienced severe epidemics of DNB (Mullett et al., 2017; Woods et al., 2005), showed the presence of at least three distinct races of *D. septosporum* in Scotland. One of the races (NPR, native pine race) was mainly associated with native Scots pine (*Pinus sylvestris*) and two others (LPR, lodgepole pine race and SR, southern race) were associated with the introduced pine species lodgepole and Corsican pine, respectively,

reflecting their likely origins (Mullett et al., 2017; Piotrowska et al., 2018). Whole-genome sequencing of 25 isolates revealed low diversity of the indigenous NPR but higher diversity and evidence of sexual reproduction within the introduced SR. The discovery of an NPR-SR hybrid, and the potential for evolution of new variants of the pathogen through intraspecies hybridization, highlighted the increased risk to native pines and the biosecurity implications of introducing more susceptible exotic pine species (Ennos et al., 2020).

In Canada, genotyping-by-sequencing of 119 *D. septosporum* isolates from both plantation and native forests also revealed genetically distinct lineages (Capron et al., 2021). The lineage associated with the Kispiox region of British Columbia, where a severe DNB epidemic occurred in the past (Woods et al., 2005), showed high genetic diversity as well as evidence for sexual reproduction, which had been reported previously in this region (Dale et al., 2011). Two lineages in other regions showed signatures of human-mediated movement and admixture, probably associated with pine breeding (Capron et al., 2021). The authors urged caution in the application of assisted gene flow policies in which pine genotypes used for reforestation are matched with predicted future climate changes. The movement of pines between regions could facilitate hybridization or recombination between genetically distinct pathogen lineages, leading to new variants and increased risk to forest health (Capron et al., 2021).

#### 6.4 | Genetic diversity and origins of *D. septosporum*

The exact origin of *D. septosporum* remains elusive. Population genetic studies using the same set of microsatellite markers across many different studies of individual countries (e.g., Boroń et al., 2021; Jánošíková et al., 2021; Oskay et al., 2020) and intercontinental comparisons have, however, enhanced our understanding of where the pathogen is well established and/or naturalized, and where it has clearly been introduced (Adamson et al., 2018; Barnes et al., 2014; Drenkhan et al., 2013; Mullett et al., 2017). Specific pathways of migration, natural range expansions, and anthropogenic introductions are now becoming more evident.

To consolidate individual population studies, for a more global perspective, Mullett et al. (2021) studied 3800 isolates, including new isolates and those used in nine previous publications, representing the most comprehensive population study on *D. septosporum* (Mullett et al., 2021). Using an evolutionary approach, they suggested a Eurasian origin of the pathogen, with historical migration patterns moving in a westerly direction and emerging from three main clusters. The oldest cluster is represented by the East European cluster that has diverged to produce three smaller subclusters centred in Turkey, central Europe, and north-eastern Europe. The North American cluster, derived from an ancestral population in Eurasia, is exclusively found in the United States and Canada, and is the population causing disease on *P. contorta* in British Columbia (Woods et al., 2005). More

detailed studies using single-nucleotide polymorphism data showed finer level population divergence (Capron et al., 2021), with a single reintroduitory event in Scotland (Ennos et al., 2020). The youngest, western European cluster, which dominates most of the UK and western France, is also the one that has been introduced most dominantly throughout most of the Southern Hemisphere (Mullett et al., 2021).

The populations of *D. septosporum* in the Southern Hemisphere are mostly characterized by clonal lineages with a single mating type, MAT 1-2 (Barnes et al., 2014). This is especially true for populations in South America (Chile and Ecuador) and Australasia (New Zealand and Australia), which have had separate introductory events, most likely with *P. radiata* (Barnes et al., 2014), and coinciding with the expansion of pine plantations. Multiple introductions from other clusters (central and north-eastern Europe) have occurred in Africa (South Africa and Kenya), resulting in populations with both mating types and high diversity, reflecting the long history of pine introduction into these countries (Barnes et al., 2014; Mullett et al., 2021) and highlighting strong anthropogenic influences on global patterns of forest disease (Wingfield et al., 2015).

## 7 | CONCLUSIONS AND FUTURE DIRECTIONS

Although the last 10 years have seen significant advances in our understanding of how *F. fulva* and *D. septosporum* interact (and have evolved) with their host plants, there is still much to learn. For instance, through continued genome sequencing of isolates collected from around the world, and in conjunction with gene editing techniques involving CRISPR-Cas technology, it may be possible to identify a repertoire of core effector genes from *F. fulva* and *D. septosporum* that is essential for host colonization. Crucially, as a starting point for this analysis, CRISPR-Cas technology has already been applied to *D. septosporum* (McCarthy et al., 2022). Such technology may also provide insights into the role(s) of the dispensable chromosome in *F. fulva*.

Further genome sequencing will also assist in the identification of *Avr* genes from *F. fulva*, particularly those deleted or mutated in resistance-breaking strains of the fungus. This is important, as it will enable rapid identification of resistance-breaking strains in commercial greenhouse and high tunnel environments; information that can be used to inform the deployment of resistant tomato varieties. Similarly, the identification of *Avr* genes will increase the speed with which breeding and selection programmes involving cognate Cf immune receptor genes can be carried out, as they can be used as markers. Regarding whether stacked Cf immune receptor genes can provide durable resistance to *F. fulva* in tomato, a key question for the future will be whether this fungus undergoes sexual reproduction, as has been suggested by Stergiopoulos, Groenewald, et al. (2007). If sexual recombination does indeed occur, strains able to overcome stacked resistance could arise more rapidly. Finally, the identification of *Avr* genes from *F. fulva* will also assist in the mapping and cloning of corresponding Cf immune receptor genes, which




will provide new opportunities to identify host virulence targets, and to assess differences in effector recognition, downstream signalling, and immune response outputs. In any case, elucidation of the complete Cf/Avr-triggered signalling pathway leading to leaf mould resistance in tomato will be a major future focus.

Unlike in angiosperms, studies of the molecular basis of disease resistance in gymnosperm trees is in its infancy. Whilst the work described here with the *D. septosporum*-pine pathosystem and some other studies (e.g., Liu et al., 2021) suggest similarities with fungal-angiosperm pathosystems in terms of effector recognition and defence pathways, there is a lot more to discover. However, there is no doubt that the wealth of information learned from the study of how *F. fulva* interacts and evolves with its tomato host will continue to help decipher the molecular mechanisms deployed by *D. septosporum*. In turn, this will help to answer fundamental questions about the extent to which pathogen resistance mechanisms of gymnosperms function and adapt in similar ways to those of angiosperms. Future work will, however, need to rise to the challenges associated with the long lifecycles of most gymnosperm trees, their immense genomes, and the confounding influences of other factors with strong impacts on disease resistance, such as climatic conditions and complex microbiomes, including other potential pathogen species associated with forest trees (Ahmar et al., 2021; Terhonen et al., 2019; Woods et al., 2016; Zimin et al., 2014).

#### DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no new data were created or analysed.

#### ORCID

Carl H. Mesarich  <https://orcid.org/0000-0002-5207-4074>  
 Irene Barnes  <https://orcid.org/0000-0002-4349-3402>  
 Ellie L. Bradley  <https://orcid.org/0000-0002-3699-4305>  
 Yanan Guo  <https://orcid.org/0000-0001-9584-6153>  
 Matthieu H. A. J. Joosten  <https://orcid.org/0000-0002-6243-4547>  
 Mengmeng Lu  <https://orcid.org/0000-0001-5023-3759>  
 Hannah M. McCarthy  <https://orcid.org/0000-0003-1966-1952>  
 Christiaan R. Schol  <https://orcid.org/0000-0001-8807-6699>  
 Ioannis Stergiopoulos  <https://orcid.org/0000-0002-2368-6119>  
 Alex Z. Zaccaron  <https://orcid.org/0000-0001-5554-2059>  
 Rosie E. Bradshaw  <https://orcid.org/0000-0001-5228-2685>

#### REFERENCES

- Adamson, K., Mullett, M.S., Solheim, H., Barnes, I., Müller, M.M., Hantula, J. et al. (2018) Looking for relationships between the populations of *Dothistroma septosporum* in northern Europe and Asia. *Fungal Genetics and Biology*, 110, 15–25.
- Ahmar, S., Ballesta, P., Ali, M. & Mora-Poblete, F. (2021) Achievements and challenges of genomics-assisted breeding in forest trees: from marker-assisted selection to genome editing. *International Journal of Molecular Sciences*, 22, 10583.
- Altin, N. (2016) Identification of race 2.5 of leaf mold (*Passalora fulva*, syn. *Cladosporium fulvum*) on tomato. *Journal of Plant Diseases and Protection*, 123, 279–284.
- Barnes, I., Crous, P.W., Wingfield, B.D. & Wingfield, M.J. (2004) Multigene phylogenies reveal that red band needle blight of *Pinus* is caused by two distinct species of *Dothistroma*, *D. septosporum* and *D. pini*. *Studies in Mycology*, 50, 551–565.
- Barnes, I., Cortinas, M.N., Wingfield, M.J. & Wingfield, B.D. (2008) Microsatellite markers for the red band needle blight pathogen, *Dothistroma septosporum*. *Molecular Ecology Resources*, 8, 1026–1029.
- Barnes, I., Wingfield, M.J., Carbone, I., Kirisits, T. & Wingfield, B.D. (2014) Population structure and diversity of an invasive pine needle pathogen reflects anthropogenic activity. *Ecology and Evolution*, 4, 3642–3661.
- Barnes, I., van der Nest, A., Mullett, M., Crous, P., Drenkhan, R., Musolin, D. et al. (2016) Neotypification of *Dothistroma septosporum* and epitypification of *D. pini*, causal agents of *Dothistroma* needle blight of pine. *Forest Pathology*, 46, 388–407.
- Bernal-Cabrera, A., Martínez-Coca, B., Herrera-Isla, L., Ynfante-Martínez, D., Peteira-Delgado, B., Portal, O. et al. (2021) The first report on the occurrence race 9 of the tomato leaf mold pathogen *Cladosporium fulvum* (syn. *Passalora fulva*) in Cuba. *European Journal of Plant Pathology*, 160, 731–736.
- Bolton, M.D., van Esse, H.P., Vossen, J.H., de Jonge, R., Stergiopoulos, I., Stulemeijer, I.J.E. et al. (2008) The novel *Cladosporium fulvum* lysin motif effector Ecp6 is a virulence factor with orthologues in other fungal species. *Molecular Microbiology*, 69, 119–136.
- Boroń, P., Lenart-Boroń, A., Mullett, M., Grad, B. & Nawrot-Chorabik, K. (2021) Population structure of *Dothistroma septosporum* in Poland: revealing the genetic signature of a recently established pathogen. *Plant Pathology*, 70, 1310–1325.
- Bradley, E.L., Ökmen, B., Doehlemann, G., Henrissat, B., Bradshaw, R.E. & Mesarich, C.H. (2022) Secreted glycoside hydrolase (GH) proteins as effectors and invasion patterns of plant-associated fungi and oomycetes. *Frontiers in Plant Science*, 13, 853106.
- Bradshaw, R.E. (2016) Growth and fruition of a pathology research community: the *Dothistroma* story. *Forest Pathology*, 46, 383–387.
- Bradshaw, R.E., Jin, H.P., Morgan, B.S., Schwelm, A., Teddy, O.R., Young, C.A. et al. (2006) A polyketide synthase gene required for biosynthesis of the aflatoxin-like toxin, dothistromin. *Mycopathologia*, 161, 283–294.
- Bradshaw, R.E., Slot, J.C., Moore, G.G., Chettri, P., de Wit, P.J.G.M., Ehrlich, K.C. et al. (2013) Fragmentation of an aflatoxin-like gene cluster in a forest pathogen. *New Phytologist*, 198, 525–535.
- Bradshaw, R.E., Guo, Y., Sim, A.D., Kabir, M.S., Chettri, P., Ozturk, I.K. et al. (2016) Genome-wide gene expression dynamics of the fungal pathogen *Dothistroma septosporum* throughout its infection cycle of the gymnosperm host *Pinus radiata*. *Molecular Plant Pathology*, 17, 210–224.
- Bradshaw, R.E., Ormond, S., Dupont, P.-Y., Chettri, P., Ozturk, I.K., McDougal, R.L. et al. (2019) Reduced virulence of an introduced forest pathogen over 50 years. *Microorganisms*, 7, 420.
- Bradshaw, R.E., Sim, A.D., Chettri, P., Dupont, P.Y., Guo, Y.N., Hunziker, L. et al. (2019) Global population genomics of the forest pathogen *Dothistroma septosporum* reveal chromosome duplications in high dothistromin-producing strains. *Molecular Plant Pathology*, 20, 784–799.
- Brennan, C.J., Benbow, H.R., Mullins, E. & Doohan, F.M. (2019) A review of the known unknowns in the early stages of septoria tritici blotch disease of wheat. *Plant Pathology*, 68, 1427–1438.
- Bulman, L.S., Bradshaw, R.E., Fraser, S., Martín-García, J., Barnes, I., Musolin, D.L. et al. (2016) A worldwide perspective on the management and control of *Dothistroma* needle blight. *Forest Pathology*, 46, 472–488.
- Capron, A., Feau, N., Heinzelmann, R., Barnes, I., Benowicz, A., Bradshaw, R.E. et al. (2021) Signatures of post-glacial genetic isolation and human-driven migration in the *Dothistroma* needle blight pathogen in western Canada. *Phytopathology*, 111, 116–127.



- Chai, X., Xu, X., Wang, D., Xue, D. & Li, J. (2020) Mapping and candidate gene screening of *Cladosporium fulvum* resistance gene Cf-12 in tomato (*Solanum lycopersicum*) by high-throughput sequencing. *Plant Breeding*, 139, 977–987.
- Chettri, P. & Bradshaw, R.E. (2016) LaeA negatively regulates dothistromin production in the pine needle pathogen *Dothistroma septosporum*. *Fungal Genetics and Biology*, 97, 24–32.
- Chettri, P., Calvo, A.M., Cary, J.W., Dhingra, S., Guo, Y.A., McDougal, R.L. et al. (2012) The *veA* gene of the pine needle pathogen *Dothistroma septosporum* regulates sporulation and secondary metabolism. *Fungal Genetics and Biology*, 49, 141–151.
- Chettri, P., Ehrlich, K.C., Cary, J.W., Collemare, J., Cox, M.P., Griffiths, S.A. et al. (2013) Dothistromin genes at multiple separate loci are regulated by AfIR. *Fungal Genetics and Biology*, 51, 12–20.
- Chettri, P., Dupont, P.-Y. & Bradshaw, R.E. (2018) Chromatin-level regulation of the fragmented dothistromin gene cluster in the forest pathogen *Dothistroma septosporum*. *Molecular Microbiology*, 107, 508–522.
- Collemare, J., Griffiths, S., Iida, Y., Jashni, M.K., Battaglia, E., Cox, R.J. et al. (2014) Secondary metabolism and biotrophic lifestyle in the tomato pathogen *Cladosporium fulvum*. *PLoS One*, 9, e85877.
- Cook, D.E., Mesarich, C.H. & Thomma, B.P.H.J. (2015) Understanding plant immunity as a surveillance system to detect invasion. *Annual Review of Phytopathology*, 53, 541–563.
- Dale, A.L., Lewis, K.J. & Murray, B.W. (2011) Sexual reproduction and gene flow in the pine pathogen *Dothistroma septosporum* in British Columbia. *Phytopathology*, 101, 68–76.
- Dangl, J.L. & Jones, J.D.G. (2001) Plant pathogens and integrated defence responses to infection. *Nature*, 411, 826–833.
- de Jonge, R., van Esse, H.P., Kombrink, A., Shinya, T., Desaki, Y., Bours, R. et al. (2010) Conserved fungal LysM effector Ecp6 prevents chitin-triggered immunity in plants. *Science*, 329, 953–955.
- de Kock, M.J.D., Iskandar, H.M., Brandwagt, B.F., Laugé, R., de Wit, P.J.G.M. & Lindhout, P. (2004) Recognition of *Cladosporium fulvum* Ecp2 elicitor by non-host *Nicotiana* spp. is mediated by a single dominant gene that is not homologous to known Cf-genes. *Molecular Plant Pathology*, 5, 397–408.
- de Wit, P.J.G.M. (2016) *Cladosporium fulvum* effectors: weapons in the arms race with tomato. *Annual Review of Phytopathology*, 54, 1–23.
- de Wit, P.J.G.M., Joosten, M.H.A.J., Thomma, B.H.P.J. & Stergiopoulos, I. (2009) Gene for gene models and beyond: the *Cladosporium fulvum*-tomato pathosystem. In: Deising, H.B. (Ed.) *Plant relationships*. Berlin, Heidelberg: Springer Berlin Heidelberg, pp. 135–156.
- de Wit, P.J.G.M., van der Burgt, A., Okmen, B., Stergiopoulos, I., Abd-El Salam, K.A., Aerts, A.L. et al. (2012) The genomes of the fungal plant pathogens *Cladosporium fulvum* and *Dothistroma septosporum* reveal adaptation to different hosts and lifestyles but also signatures of common ancestry. *PLoS Genetics*, 8, e1003088.
- DeLano, W.L. (2002) Pymol: an open-source molecular graphics tool. *CCP4 Newsletter on Protein Crystallography*, 40, 82–92.
- Dixon, M.S., Hatzixanthis, K., Jones, D.A., Harrison, K. & Jones, J.D.G. (1998) The tomato Cf-5 disease resistance gene and six homologs show pronounced allelic variation in leucine-rich repeat copy number. *The Plant Cell*, 10, 1915–1925.
- Drenkhan, R., Hantula, J., Vuorinen, M., Jankovsky, L. & Muller, M.M. (2013) Genetic diversity of *Dothistroma septosporum* in Estonia, Finland and Czech Republic. *European Journal of Plant Pathology*, 136, 71–85.
- Drenkhan, R., Tomešová-Haataja, V., Fraser, S., Bradshaw, R.E., Vahalík, P., Mullett, M.S. et al. (2016) Global geographic distribution and host range of *Dothistroma* species: a comprehensive review. *Forest Pathology*, 46, 408–442.
- Ennos, R.A., Sjökvist, E.I., Piotrowska, M.J., Riddell, C. & Hoebe, P.N. (2020) Using genome resequencing to investigate racial structure, genetic diversity, sexual reproduction and hybridisation in the pine pathogen *Dothistroma septosporum*. *Fungal Ecology*, 45, 100921.
- Fraser, S., Martín-García, J., Perry, A., Kabir, M.S., Owen, T., Solla, A. et al. (2016) A review of Pinaceae resistance mechanisms against needle and shoot pathogens with a focus on the *Dothistroma*-*Pinus* interaction. *Forest Pathology*, 46, 453–471.
- Gabriëls, S.H., Vossen, J.H., Ekengren, S.K., van Ooijen, G.V., Abd-El-Haliem, A.M., van den Berg, G.C. et al. (2007) An NB-LRR protein required for HR signalling mediated by both extra- and intracellular resistance proteins. *The Plant Journal*, 50, 14–28.
- Ghanbarnia, K., Ma, L., Larkan, N.J., Haddadi, P., Fernando, W.G.D. & Borhan, M.H. (2018) *Leptosphaeria maculans* AvrLm9: a new player in the game of hide and seek with AvrLm4-7. *Molecular Plant Pathology*, 19, 1754–1764.
- Griffiths, S., Saccomanno, B., de Wit, P.J.G.M. & Collemare, J. (2015) Regulation of secondary metabolite production in the fungal tomato pathogen *Cladosporium fulvum*. *Fungal Genetics and Biology*, 84, 52–61.
- Griffiths, S., Mesarich, C.H., Saccomanno, B., Vaisberg, A., de Wit, P.J.G.M., Cox, R. et al. (2016) Elucidation of cladofulvin biosynthesis reveals a cytochrome P450 monooxygenase required for anthraquinone dimerization. *Proceedings of the National Academy of Sciences of the United States of America*, 113, 6851–6856.
- Griffiths, S., Mesarich, C.H., Overdijk, E.J., Saccomanno, B., de Wit, P.J. & Collemare, J. (2018) Down-regulation of cladofulvin biosynthesis is required for biotrophic growth of *Cladosporium fulvum* on tomato. *Molecular Plant Pathology*, 19, 369–380.
- Griffiths, S.A., Cox, R.J., Overdijk, E.J.R., Mesarich, C.H., Saccomanno, B., Lazarus, C.M. et al. (2018) Assignment of a dubious gene cluster to melanin biosynthesis in the tomato fungal pathogen *Cladosporium fulvum*. *PLoS One*, 13, e0209600.
- Groenewald, M., Barnes, I., Bradshaw, R.E., Brown, A.V., Dale, A., Groenewald, J.Z. et al. (2007) Characterization and distribution of mating type genes in the *Dothistroma* needle blight pathogens. *Phytopathology*, 97, 825–834.
- Grushetskaya, Z.E., Lemesh, V.A., Poliksenova, V.D. & Khotyleva, L.V. (2007) Mapping of the Cf-6 tomato leaf mould resistance locus using SSR markers. *Russian Journal of Genetics*, 43, 1266–1270.
- Guo, Y., Hunziker, L., Mesarich, C.H., Chettri, P., Dupont, P.-Y., Ganley, R.J. et al. (2020) DsEcp2-1 is a polymorphic effector that restricts growth of *Dothistroma septosporum* in pine. *Fungal Genetics and Biology*, 135, 103300.
- Haanstra, J., Meijer-Dekens, F., Lauge, R., Seetana, D., Joosten, M.H., de Wit, P.J. et al. (2000) Mapping strategy for resistance genes against *Cladosporium fulvum* on the short arm of chromosome 1 of tomato: Cf-ECP5 near the Hcr9 Milky Way cluster. *Theoretical and Applied Genetics*, 101, 661–668.
- Hane, J.K., Paxman, J., Jones, D.A., Oliver, R.P. & de Wit, P. (2020) “CATAstrophy,” a genome-informed trophic classification of filamentous plant pathogens – how many different types of filamentous plant pathogens are there? *Frontiers in Microbiology*, 10, 3088.
- Hirst, P., Richardson, T.E., Carson, S.D. & Bradshaw, R.E. (1999) *Dothistroma pini* genetic diversity is low in New Zealand. *New Zealand Journal of Forestry Science*, 29, 459–472.
- Hörger, A.C., Ilyas, M., Stephan, W., Tellier, A., van der Hoorn, R.A. & Rose, L.E. (2012) Balancing selection at the tomato RCR3 guard gene family maintains variation in strength of pathogen defense. *PLoS Genetics*, 8, e1002813.
- Huang, W.R., Schol, C., Villanueva, S.L., Heidstra, R. & Joosten, M.H. (2021) Knocking out SOBIR1 in *Nicotiana benthamiana* abolishes functionality of transgenic receptor-like protein Cf-4. *Plant Physiology*, 185, 290–294.
- Hunziker, L., Mesarich, C.H., McDougal, R.L. & Bradshaw, R.E. (2016) Effector identification in the pine pathogen *Dothistroma septosporum*. *New Zealand Plant Protection*, 69, 94–98.

- Hunziker, L., Tarallo, M., Gough, K., Guo, M., Hargreaves, C., Loo, T.S. et al. (2021) Apoplastic effector candidates of a foliar forest pathogen trigger cell death in host and non-host plants. *Scientific Reports*, 11, 19958.
- Hurlburt, N.K., Chen, L.H., Stergiopoulos, I. & Fisher, A.J. (2018) Structure of the *Cladosporium fulvum* Avr4 effector in complex with (GlcNAc)<sub>6</sub> reveals the ligand-binding mechanism and uncouples its intrinsic function from recognition by the Cf-4 resistance protein. *PLoS Pathogens*, 14, e1007263.
- Iakovovidis, M., Soumpourou, E., Anderson, E., Etherington, G., Yourstone, S. & Thomas, C. (2020) Genes encoding recognition of the *Cladosporium fulvum* effector protein Ecp5 are encoded at several loci in the tomato genome. *G3 Genes|Genomes|Genetics*, 10, 1753–1763.
- Iida, Y., van't Hof, P., Beenen, H., Mesarich, C., Kubota, M., Stergiopoulos, I. et al. (2015) Novel mutations detected in avirulence genes overcoming tomato Cf resistance genes in isolates of a Japanese population of *Cladosporium fulvum*. *PLoS One*, 10, e0123271.
- Ilyas, M., Hörger, A.C., Bozkurt, T.O., van den Burg, H.A., Kaschani, F., Kaiser, M. et al. (2015) Functional divergence of two secreted immune proteases of tomato. *Current Biology*, 25, 2300–2306.
- Ioos, R., Fabre, B., Saurat, C., Fourrier, C., Frey, P. & Marçais, B. (2010) Development, comparison, and validation of real-time and conventional PCR tools for the detection of the fungal pathogens causing brown spot and red band needle blights of pine. *Phytopathology*, 100, 105–114.
- Jánošíková, Z., Dutech, C., Ondrušková, E., Adamčíková, K. & Mullett, M. (2021) Population structure and genetic diversity of *Dothistroma septosporum* in Slovakia. *European Journal of Plant Pathology*, 160, 1–17.
- Jashni, M.K., Dols, I.H., Iida, Y., Boeren, S., Beenen, H.G., Mehrabi, R. et al. (2015) Synergistic action of a metalloprotease and a serine protease from *Fusarium oxysporum* f. sp. *lycopersici* cleaves chitin-binding tomato chitinases, reduces their antifungal activity, and enhances fungal virulence. *Molecular Plant-Microbe Interactions*, 28, 996–1008.
- Jiang, X., Li, Y., Li, R., Gao, Y., Liu, Z., Yang, H. et al. (2022) Transcriptome analysis of the Cf-13-mediated hypersensitive response of tomato to *Cladosporium fulvum* infection. *International Journal of Molecular Sciences*, 23, 4844.
- Joosten, M.H.A.J., Cozijnsen, T.J. & de Wit, P.J.G.M. (1994) Host resistance to a fungal tomato pathogen lost by a single base-pair change in an avirulence gene. *Nature*, 367, 384–386.
- Joosten, M.H.A.J., Vogelsang, R., Cozijnsen, T.J., Verberne, M.C. & de Wit, P.J.G.M. (1997) The biotrophic fungus *Cladosporium fulvum* circumvents Cf-4-mediated resistance by producing unstable AVR4 elicitors. *The Plant Cell*, 9, 367–379.
- Kabir, M.S., Ganley, R.J. & Bradshaw, R.E. (2015a) Dothistromin toxin is a virulence factor in dothistroma needle blight of pines. *Plant Pathology*, 64, 225–234.
- Kabir, M.S., Ganley, R.J. & Bradshaw, R.E. (2015b) The hemibiotrophic lifestyle of the fungal pine pathogen *Dothistroma septosporum*. *Forest Pathology*, 45, 190–202.
- Kahlon, P.S., Seta, S.M., Zander, G., Scheikl, D., Hüchelhoven, R., Joosten, M.H. et al. (2020) Population studies of the wild tomato species *Solanum chilense* reveal geographically structured major gene-mediated pathogen resistance. *Proceedings of the Royal Society B*, 287, 20202723.
- Karimi Jashni, M., van der Burgt, A., Battaglia, E., Mehrabi, R., Collemare, J. & de Wit, P.J.G.M. (2020) Transcriptome and proteome analyses of proteases in biotroph fungal pathogen *Cladosporium fulvum*. *Journal of Plant Pathology*, 102, 377–386.
- Karimi-Jashni, M., Maeda, K., Yazdanpanah, F., de Wit, P.J. & Iida, Y. (2022) An integrated omics approach uncovers the novel effector Ecp20-2 required for full virulence of *Cladosporium fulvum* on tomato. *Frontiers in Microbiology*, 13, 919809.
- Kourelis, J., Malik, S., Mattinson, O., Krauter, S., Kahlon, P.S., Paulus, J.K. et al. (2020) Evolution of a guarded decoy protease and its receptor in solanaceous plants. *Nature Communications*, 11, 4393.
- Kourelis, J., Contreras, M.P., Harant, A., Pai, H., Lüdke, D., Adachi, H. et al. (2022) The helper NLR immune protein NRC3 mediates the hypersensitive cell death caused by the cell-surface receptor Cf-4. *PLoS Genetics*, 18, e1010414.
- Kruijt, M., Kip, D.J., Joosten, M.H., Brandwagt, B.F. & de Wit, P.J. (2005) The Cf-4 and Cf-9 resistance genes against *Cladosporium fulvum* are conserved in wild tomato species. *Molecular Plant-Microbe Interactions*, 18, 1011–1021.
- Kubota, M., Morishima, M. & Iida, Y. (2015) First occurrence of tomato leaf mold caused by the novel races 2.5.9 and 4.5.9 of *Passalora fulva* in Japan. *Journal of General Plant Pathology*, 81, 320–323.
- Laugé, R., Joosten, M.H.A.J., van den Ackerveken, G.F.J.M., van den Broek, H.W.J. & de Wit, P.J.G.M. (1997) The in planta-produced extracellular proteins ECP1 and ECP2 of *Cladosporium fulvum* are virulence factors. *Molecular Plant-Microbe Interactions*, 10, 725–734.
- Laugé, R., Goodwin, P.H., de Wit, P.J.G.M. & Joosten, M.H.A.J. (2000) Specific HR-associated recognition of secreted proteins from *Cladosporium fulvum* occurs in both host and non-host plants. *The Plant Journal*, 23, 735–745.
- Lazar, N., Mesarich, C.H., Petit-Houdenot, Y., Talbi, N., Li de la Sierra-Gallay, I., Zélie, E. et al. (2022) A new family of structurally conserved fungal effectors displays epistatic interactions with plant resistance proteins. *PLoS Pathogens*, 18, e1010664.
- Lee, J.H., Park, M.S., Kim, J.-C., Jang, K.S., Choi, Y.H., Kim, H.T. et al. (2013) Occurrence of leaf mold pathogen *Fulvia fulva* isolates infecting tomato Cf-9 cultivars in Korea. *Horticultural Science & Technology*, 31, 740–747.
- Li, S., Zhao, T., Li, H., Xu, X. & Li, J. (2015) First report of races 2.5 and 2.4.5 of *Cladosporium fulvum* (syn. *Passalora fulva*), causal fungus of tomato leaf mold disease in China. *Journal of General Plant Pathology*, 81, 162–165.
- Liebrand, T.W.H., van den Berg, G.C.M., Zhang, Z., Smit, P., Cordewener, J.H.G., America, A.H.P. et al. (2013) Receptor-like kinase SOBIR1/EVR interacts with receptor-like proteins in plant immunity against fungal infection. *Proceedings of the National Academy of Sciences of the United States of America*, 110, 10010–10015.
- Liebrand, T.W., van den Burg, H.A. & Joosten, M.H. (2014) Two for all: receptor-associated kinases SOBIR1 and BAK1. *Trends in Plant Science*, 19, 123–132.
- Liu, G., Liu, J., Zhang, C., You, X., Zhao, T., Jiang, J. et al. (2018) Physiological and RNA-seq analyses provide insights into the response mechanism of the Cf-10-mediated resistance to *Cladosporium fulvum* infection in tomato. *Plant Molecular Biology*, 96, 403–416.
- Liu, G., Zhao, T., You, X., Jiang, J., Li, J. & Xu, X. (2019) Molecular mapping of the Cf-10 gene by combining SNP/InDel-index and linkage analysis in tomato (*Solanum lycopersicum*). *BMC Plant Biology*, 19, 15.
- Liu, J.-J., Schoettle, A.W., Sniezko, R.A., Williams, H., Zamany, A. & Rancourt, B. (2021) Fine dissection of limber pine resistance to *Cronartium ribicola* using targeted sequencing of the NLR family. *BMC Genomics*, 22, 567.
- Liu, G., Zhang, D., Zhao, T., Yang, H., Jiang, J., Li, J. et al. (2022) Integrating omics reveals that miRNA-guided genetic regulation on plant hormone level and defense response pathways shape resistance to *Cladosporium fulvum* in the tomato Cf-10-carrying line. *Research Square*. Available from: <https://doi.org/10.21203/rs.3.rs-2034226/v1>
- Lo Presti, L., Lanver, D., Schweizer, G., Tanaka, S., Liang, L., Tollot, M. et al. (2015) Fungal effectors and plant susceptibility. *Annual Review of Plant Biology*, 66, 513–545.
- Lu, M., Feau, N., Vidakovic, D.O., Ukrainetz, N., Wong, B., Aitken, S.N. et al. (2021) Comparative gene expression analysis reveals mechanism of *Pinus contorta* response to the fungal pathogen *Dothistroma septosporum*. *Molecular Plant-Microbe Interactions*, 34, 397–409.
- Lucentini, C.G., Medina, R., Franco, M.E.E., Saparrat, M.C.N. & Balatti, P.A. (2021) *Fulvia fulva* [syn. *Cladosporium fulvum*, *Passalora fulva*] races in Argentina are evolving through genetic changes and carry

- polymorphic *avr* and *ecp* gene sequences. *European Journal of Plant Pathology*, 159, 525–542.
- Macheleidt, J., Mattern, D.J., Fischer, J., Netzker, T., Weber, J., Schroeckh, V. et al. (2016) Regulation and role of fungal secondary metabolites. *Annual Review of Genetics*, 50, 371–392.
- McCarthy, H.M., Tarallo, M., Mesarich, C.H., McDougal, R.L. & Bradshaw, R.E. (2022) Targeted gene mutations in the forest pathogen *Dothistroma septosporum* using CRISPR/Cas9. *Plants*, 11, 1016.
- Medina, R., López, S.M.Y., Franco, M.E.E., Rollan, C., Ronco, B.L., Saparrat, M.C.N. et al. (2015) A survey on occurrence of *Cladosporium fulvum* identifies race 0 and race 2 in tomato-growing areas of Argentina. *Plant Disease*, 99, 1732–1737.
- Mesarich, C.H., Griffiths, S.A., van der Burgt, A., Okmen, B., Beenen, H.G., Etalo, D.W. et al. (2014) Transcriptome sequencing uncovers the *Avr5* avirulence gene of the tomato leaf mould pathogen *Cladosporium fulvum*. *Molecular Plant-Microbe Interactions*, 27, 846–857.
- Mesarich, C.H., Stergiopoulos, I., Beenen, H.G., Cordovez, V., Guo, Y., Jashni, M.K. et al. (2016) A conserved proline residue in dothideomycete *Avr4* effector proteins is required to trigger a Cf-4-dependent hypersensitive response. *Molecular Plant Pathology*, 17, 84–95.
- Mesarich, C.H., Okmen, B., Rovenich, H., Griffiths, S.A., Wang, C.C., Jashni, M.K. et al. (2018) Specific hypersensitive response-associated recognition of new apoplastic effectors from *Cladosporium fulvum* in wild tomato. *Molecular Plant-Microbe Interactions*, 31, 145–162.
- Michielse, C.B., van Wijk, R., Reijnen, L., Manders, E.M.M., Boas, S., Olivain, C. et al. (2009) The nuclear protein Sge1 of *Fusarium oxysporum* is required for parasitic growth. *PLoS Pathogens*, 5, e1000637.
- Mullett, M.S., Brown, A.V., Fraser, S., Baden, R. & Tubby, K.V. (2017) Insights into the pathways of spread and potential origins of *Dothistroma septosporum* in Britain. *Fungal Ecology*, 26, 85–98.
- Mullett, M.S., Drenkhan, R., Adamson, K., Boroń, P., Lenart-Boroń, A., Barnes, I. et al. (2021) Worldwide genetic structure elucidates the Eurasian origin and invasion pathways of *Dothistroma septosporum*, causal agent of Dothistroma needle blight. *Journal of Fungi*, 7, 111.
- Novak, A., Čosić, J., Vrandečić, K., Jurković, D., Plavec, J., Križanac, I. et al. (2021) Characterization of tomato leaf mould pathogen, *Passalora fulva*, in Croatia. *Journal of Plant Diseases and Protection*, 128, 1041–1049.
- Ohm, R.A., Feau, N., Henrissat, B., Schoch, C.L., Horwitz, B.A., Barry, K.W. et al. (2012) Diverse lifestyles and strategies of plant pathogenesis encoded in the genomes of eighteen Dothideomycetes fungi. *PLoS Pathogens*, 8, e1003037.
- Ökmen, B., Etalo, D.W., Joosten, M.H.A.J., Bouwmeester, H.J., de Vos, R.C.H., Collemare, J. et al. (2013) Detoxification of alpha-tomatine by *Cladosporium fulvum* is required for full virulence on tomato. *New Phytologist*, 198, 1203–1214.
- Ökmen, B., Collemare, J., Griffiths, S., van der Burgt, A., Cox, R. & de Wit, P.J.G.M. (2014) Functional analysis of the conserved transcriptional regulator CfWor1 in *Cladosporium fulvum* reveals diverse roles in the virulence of plant pathogenic fungi. *Molecular Microbiology*, 92, 10–27.
- Ökmen, B., Bachmann, D. & de Wit, P.J. (2019) A conserved GH17 glycosyl hydrolase from plant pathogenic Dothideomycetes releases a DAMP causing cell death in tomato. *Molecular Plant Pathology*, 20, 1710–1721.
- Oskay, F., Tunali, Z., Lehtijärvi, A.T., Doğmuş-Lehtijärvi, H.T., Woodward, S. & Mullett, M. (2020) Distribution and genetic diversity of *Dothistroma septosporum* in *Pinus brutia* forests of south-western Turkey. *Plant Pathology*, 69, 1551–1564.
- Ozturk, I.K., Chettri, P., Dupont, P.-Y., Barnes, I., McDougal, R.L., Moore, G.G. et al. (2017) Evolution of polyketide synthesis in a dothideomycete forest pathogen. *Fungal Genetics and Biology*, 106, 42–50.
- Ozturk, I.K., Dupont, P.-Y., Chettri, P., McDougal, R., Böhl, O.J., Cox, R.J. et al. (2019) Evolutionary relics dominate the small number of secondary metabolism genes in the hemibiotrophic fungus *Dothistroma septosporum*. *Fungal Biology*, 123, 397–407.
- Parlange, F., Daverdin, G., Fudal, I., Kuhn, M.L., Balesdent, M.H., Blaise, F. et al. (2009) *Leptosphaeria maculans* avirulence gene *AvrLm4-7* confers a dual recognition specificity by the *Rlm4* and *Rlm7* resistance genes of oilseed rape, and circumvents Rlm4-mediated recognition through a single amino acid change. *Molecular Microbiology*, 71, 851–863.
- Parniske, M., Hammond-Kosack, K.E., Golstein, C., Thomas, C.M., Jones, D.A., Harrison, K. et al. (1997) Novel disease resistance specificities result from sequence exchange between tandemly repeated genes at the *Cf-4/9* locus of tomato. *Cell*, 91, 821–832.
- Paulus, J.K., Kourelis, J., Ramasubramanian, S., Homma, F., Godson, A., Hörger, A.C. et al. (2020) Extracellular proteolytic cascade in tomato activates immune protease Rcr3. *Proceedings of the National Academy of Sciences of the United States of America*, 117, 17409–17417.
- Piotrowska, M.J., Riddell, C., Hoebe, P.N. & Ennos, R.A. (2018) Planting exotic relatives has increased the threat posed by *Dothistroma septosporum* to the Caledonian pine populations of Scotland. *Evolutionary Applications*, 11, 350–363.
- Plissonneau, C., Daverdin, G., Ollivier, B., Blaise, F., Degrave, A., Fudal, I. et al. (2016) A game of hide and seek between avirulence genes *AvrLm4-7* and *AvrLm3* in *Leptosphaeria maculans*. *New Phytologist*, 209, 1613–1624.
- Plissonneau, C., Rouxel, T., Chèvre, A.M., van de Wouw, A.P. & Balesdent, M.H. (2018) One gene-one name: the *AvrLmJ1* avirulence gene of *Leptosphaeria maculans* is *AvrLm5*. *Molecular Plant Pathology*, 19, 1012–1016.
- Postma, J., Liebrand, T.W., Bi, G., Evrard, A., Bye, R.R., Mbengue, M. et al. (2016) *Avr4* promotes Cf-4 receptor-like protein association with the BAK1/SERK3 receptor-like kinase to initiate receptor endocytosis and plant immunity. *New Phytologist*, 210, 627–642.
- Reichardt, S., Reppe, D., Tuzhikov, A.I., Galiullina, R.A., Planas-Marquès, M., Chichkova, N.V. et al. (2018) The tomato subtilase family includes several cell death-related proteinases with caspase specificity. *Scientific Reports*, 8, 10531–14.
- Rocafort, M., Fudal, I. & Mesarich, C.H. (2020) Apoplastic effector proteins of plant-associated fungi and oomycetes. *Current Opinion in Plant Biology*, 56, 9–19.
- Rollan, C., Protto, V., Medina, R., Lopez, S., Vera Bahima, J., Ronco, L. et al. (2013) Identification of races 0 and 2 of *Cladosporium fulvum* (syn *Passalora fulva*) on tomato in the Cinturón Hortícola de La Plata, Argentina. *Plant Disease*, 97, 992.
- Rooney, H.C.E., van't Klooster, J.W., van der Hoorn, R.A.L., Joosten, M.H.A.J., Jones, J.D.G. & de Wit, P.J.G.M. (2005) *Cladosporium Avr2* inhibits tomato Rcr3 protease required for Cf-2-dependent disease resistance. *Science*, 308, 1783–1786.
- Rowland, O., Ludwig, A.A., Merrick, C.J., Baillieux, F., Tracy, F.E., Durrant, W.E. et al. (2005) Functional analysis of *Avr9/Cf-9* rapidly elicited genes identifies a protein kinase, ACIK1, that is essential for full Cf-9-dependent disease resistance in tomato. *The Plant Cell*, 17, 295–310.
- Sanchez-Vallet, A., Saleem-Batcha, R., Kombrink, A., Hansen, G., Valkenburg, D.-J., Thomma, B.P.H.J. et al. (2013) Fungal effector Ecp6 outcompetes host immune receptor for chitin binding through intrachain LysM dimerization. *eLife*, 2, e00790.
- Sánchez-Vallet, A., McDonald, M.C., Solomon, P.S. & McDonald, B.A. (2015) Is *Zymoseptoria tritici* a hemibiotroph? *Fungal Genetics and Biology*, 79, 29–32.
- Sangster, W. (2022) ISF *Passalora fulva* project tomato. Available from: [https://worldseed.org/dl.php?file\\_url=https://worldseed.org/wp-content/uploads/2022/04/Final-report-ISF-Pf-project-v183.pdf](https://worldseed.org/dl.php?file_url=https://worldseed.org/wp-content/uploads/2022/04/Final-report-ISF-Pf-project-v183.pdf)
- Seong, K. & Krasileva, K. (2022) Comparative computational structural genomics highlights divergent evolution of fungal effectors. *bioRxiv*. Available from: <https://doi.org/10.1101/2022.05.02.490317>

- Siah, A., Deweer, C., Duyme, F., Sanssene, J., Durand, R., Halama, P. et al. (2010) Correlation of *in planta* endo-beta-1,4-xylanase activity with the necrotrophic phase of the hemibiotrophic fungus *Mycosphaerella graminicola*. *Plant Pathology*, 59, 661–670.
- Stergiopoulos, I., De Kock, M.J., Lindhout, P. & De Wit, P.J. (2007) Allelic variation in the effector genes of the tomato pathogen *Cladosporium fulvum* reveals different modes of adaptive evolution. *Molecular Plant-Microbe Interactions*, 20, 1271–1283.
- Stergiopoulos, I., Groenewald, M., Staats, M., Lindhout, P., Crous, P.W. & de Wit, P.J.G.M. (2007) Mating-type genes and the genetic structure of a world-wide collection of the tomato pathogen *Cladosporium fulvum*. *Fungal Genetics and Biology*, 44, 415–429.
- Stergiopoulos, I., van den Burg, H.A., Okmen, B., Beenen, H.G., van Liere, S., Kema, G.H.J. et al. (2010) Tomato Cf resistance proteins mediate recognition of cognate homologous effectors from fungi pathogenic on dicots and monocots. *Proceedings of the National Academy of Sciences of the United States of America*, 107, 7610–7615.
- Stergiopoulos, I., Kourmpetis, Y.A., Slot, J.C., Bakker, F.T., de Wit, P.J. & Rokas, A. (2012) In silico characterization and molecular evolutionary analysis of a novel superfamily of fungal effector proteins. *Molecular Biology and Evolution*, 29, 3371–3384.
- Sudermann, M.A., McGilp, L., Vogel, G., Regnier, M., Jaramillo, A.R. & Smart, C.D. (2022) The diversity of *Passalora fulva* isolates collected from tomato plants in US high tunnels. *Phytopathology*, 112, 1350–1360.
- Tarallo, M., McDougal, R.L., Chen, Z., Wang, Y., Bradshaw, R.E. & Mesarich, C.H. (2022) Characterization of two conserved cell death elicitor families from the Dothideomycete fungal pathogens *Dothistroma septosporum* and *Dulvia fulva* (syn. *Cladosporium fulvum*). *Frontiers in Microbiology*, 13, 964851.
- Terhonen, E., Blumenstein, K., Kovalchuk, A. & Asiegbu, F.O. (2019) Forest tree microbiomes and associated fungal endophytes: functional roles and impact on forest health. *Forests*, 10, 42.
- Thomas, C.M., Jones, D.A., Parniske, M., Harrison, K., Balint-Kurti, P.J., Hatzixanthis, K. et al. (1997) Characterization of the tomato Cf-4 gene for resistance to *Cladosporium fulvum* identifies sequences that determine recognition specificity in Cf-4 and Cf-9. *The Plant Cell*, 9, 2209–2224.
- Thomma, B.P.H.J., van Esse, H.P., Crous, P.W. & de Wit, P.J.G.M. (2005) *Cladosporium fulvum* (syn. *Passalora fulva*), a highly specialized plant pathogen as a model for functional studies on plant pathogenic Mycosphaerellaceae. *Molecular Plant Pathology*, 6, 379–393.
- van de Wouw, A.P., Lowe, R.G.T., Elliott, C.E., Dubois, D.J. & Howlett, B.J. (2014) An avirulence gene, *AvrLmJ1*, from the blackleg fungus, *Leptosphaeria maculans*, confers avirulence to *Brassica juncea* cultivars. *Molecular Plant Pathology*, 15, 523–530.
- van den Ackerveken, G., van Kan, J.A., Joosten, M., Muisers, J.M., Verbakel, H.M. & de Wit, P. (1993) Characterization of two putative pathogenicity genes of the fungal tomato pathogen *Cladosporium fulvum*. *Molecular Plant-Microbe Interactions*, 6, 210–215.
- van den Burg, H.A., Westerink, N., Francoijs, K.-J., Roth, R., Woestenenk, E., Boeren, S. et al. (2003) Natural disulfide bond-disrupted mutants of AVR4 of the tomato pathogen *Cladosporium fulvum* are sensitive to proteolysis, circumvent Cf-4-mediated resistance, but retain their chitin binding ability. *Journal of Biological Chemistry*, 278, 27340–27346.
- van den Burg, H.A., Spronk, C.A.E.M., Boeren, S., Kennedy, M.A., Vissers, J.P.C., Vuister, G.W. et al. (2004) Binding of the AVR4 elicitor of *Cladosporium fulvum* to chitotriose units is facilitated by positive allosteric protein-protein interactions: the chitin-binding site of AVR4 represents a novel binding site on the folding scaffold shared between the invertebrate and the plant chitin-binding domain. *Journal of Biological Chemistry*, 279, 16786–16796.
- van den Burg, H.A., Harrison, S.J., Joosten, M.H., Vervoort, J. & de Wit, P.J. (2006) *Cladosporium fulvum* Avr4 protects fungal cell walls against hydrolysis by plant chitinases accumulating during infection. *Molecular Plant-Microbe Interactions*, 19, 1420–1430.
- van der Biezen, E.A. & Jones, J.D.G. (1998) Plant disease-resistance proteins and the gene-for-gene concept. *Trends in Biochemical Sciences*, 23, 454–456.
- van der Burgh, A.M. & Joosten, M.H. (2019) Plant immunity: thinking outside and inside the box. *Trends in Plant Science*, 24, 587–601.
- van der Burgh, A.M., Postma, J., Robatzek, S. & Joosten, M.H.A.J. (2019) Kinase activity of SOBIR1 and BAK1 is required for immune signaling. *Molecular Plant Pathology*, 20, 410–422.
- van der Burgt, A., Jashni, M.K., Bahkali, A.H. & de Wit, P.J.G.M. (2014) Pseudogenization in pathogenic fungi with different host plants and lifestyles might reflect their evolutionary past. *Molecular Plant Pathology*, 15, 133–144.
- van Kan, J.A., van den Ackerveken, G. & de Wit, P. (1991) Cloning and characterization of cDNA of avirulence gene *avr9* of the fungal pathogen *Cladosporium fulvum*, causal agent of tomato leaf mold. *Molecular Plant-Microbe Interactions*, 4, 52–59.
- Van't Klooster, J.W., van der Kamp, M.W., Vervoort, J., Beekwilder, J., Boeren, S., Joosten, M.A.J. et al. (2011) Affinity of Avr2 for tomato cysteine protease Rcr3 correlates with the Avr2-triggered Cf-2-mediated hypersensitive response. *Molecular Plant Pathology*, 12, 21–30.
- Videira, S., Groenewald, J., Nakashima, C., Braun, U., Barreto, R.W., de Wit, P.J. et al. (2017) Mycosphaerellaceae – chaos or clarity? *Studies in Mycology*, 87, 257–421.
- Welsh, C., Lewis, K. & Woods, A. (2009) The outbreak history of Dothistroma needle blight: an emerging forest disease in northwestern British Columbia, Canada. *Canadian Journal of Forest Research*, 39, 2505–2519.
- Wingfield, M.J., Brockerhoff, E.G., Wingfield, B.D. & Slippers, B. (2015) Planted forest health: the need for a global strategy. *Science*, 349, 832–836.
- Woods, A.J. (2003) Species diversity and forest health in Northwest British Columbia. *The Forestry Chronicle*, 79, 892–897.
- Woods, A.J., Coates, K.D. & Hamann, A. (2005) Is an unprecedented Dothistroma needle blight epidemic related to climate change? *Bioscience*, 55, 761–769.
- Woods, A.J., Martín-García, J., Bulman, L., Vasconcelos, M.W., Boberg, J., La Porta, N. et al. (2016) Dothistroma needle blight, weather and possible climatic triggers for the disease's recent emergence. *Forest Pathology*, 46, 443–452.
- Wubben, J.P., Joosten, M. & de Wit, P. (1994) Expression and localization of two *in planta* induced extracellular proteins of the fungal tomato pathogen *Cladosporium fulvum*. *Molecular Plant-Microbe Interactions*, 7, 516–524.
- Xiang Neik, T., Ghanbarinia, K., Ollivier, B., Scheben, A., Severn-Ellis, A., Larkan, N.J. et al. (2022) Two independent approaches converge to the cloning of a new *Leptosphaeria maculans* avirulence effector gene, *AvrLmS-Lep2*. *Molecular Plant Pathology*, 23, 733–748.
- Xu, W., Chen, L., He, P., Yang, J., Xu, C., Wang, B. et al. (2018) Analysis of Cf-12 tomato transcriptome profile in response to *Cladosporium fulvum* infection with HISAT, StringTie and Ballgown. *International Journal of Agriculture and Biology*, 20, 2590–2598.
- Xue, D.-Q., Chen, X.-L., Zhang, H., Chai, X.-F., Jiang, J.-B., Xu, X.-Y. et al. (2017) Transcriptome analysis of the Cf-12-mediated resistance response to *Cladosporium fulvum* in tomato. *Frontiers in Plant Science*, 7, 2012.
- Yoshida, K., Asano, S., Sushida, H. & Iida, Y. (2021) Occurrence of tomato leaf mold caused by novel race 2.4.9 of *Cladosporium fulvum* in Japan. *Journal of General Plant Pathology*, 87, 35–38.
- Zaccaron, A.Z., Chen, L.-H., Samaras, A. & Stergiopoulos, I. (2022) A chromosome-scale genome assembly of the tomato pathogen *Cladosporium fulvum* reveals a compartmentalized genome architecture and the presence of a dispensable chromosome. *Microbial Genomics*, 8, 000819.
- Zhang, D., Bao, Y., Sun, Y., Yang, H., Zhao, T., Li, H. et al. (2020) Comparative transcriptome analysis reveals the response

- mechanism of *Cf-16*-mediated resistance to *Cladosporium fulvum* infection in tomato. *BMC Plant Biology*, 20, 331–16.
- Zhao, T., Liu, G., Li, S., Li, J., Jiang, J., Zhang, H. et al. (2015) Differentially expressed gene transcripts related to the *Cf-19*-mediated resistance response to *Cladosporium fulvum* infection in tomato. *Physiological and Molecular Plant Pathology*, 89, 8–15.
- Zhao, T., Jiang, J., Liu, G., He, S., Zhang, H., Chen, X. et al. (2016) Mapping and candidate gene screening of tomato *Cladosporium fulvum*-resistant gene *Cf-19*, based on high-throughput sequencing technology. *BMC Plant Biology*, 16, 51–10.
- Zhao, T., Liu, W., Zhao, Z., Yang, H., Bao, Y., Zhang, D. et al. (2019) Transcriptome profiling reveals the response process of tomato carrying *Cf-19* and *Cladosporium fulvum* interaction. *BMC Plant Biology*, 19, 5721–12.
- Zimin, A., Stevens, K.A., Crepeau, M.W., Holtz-Morris, A., Koriabine, M., Marçais, G. et al. (2014) Sequencing and assembly of the 22-Gb loblolly pine genome. *Genetics*, 196, 875–890.

## SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

**How to cite this article:** Mesarich, C.H., Barnes, I., Bradley, E.L., de la Rosa, S., de Wit, P.J.G.M., Guo, Y. et al. (2023) Beyond the genomes of *Fulvia fulva* (syn. *Cladosporium fulvum*) and *Dothistroma septosporum*: New insights into how these fungal pathogens interact with their host plants. *Molecular Plant Pathology*, 24, 474–494. Available from: <https://doi.org/10.1111/mpp.13309>