



Profile of per- and polyfluoroalkyl substances, source appointment, and determinants in Argentinean postpartum women

Solrunn Hansen^{a,*}, Shanshan Xu^b, Sandra Huber^c, Marisa Viviana Alvarez^d, Jon Øyvind Odland^{e,f,g}

^a Department of Health and Care Sciences, UiT The Arctic University of Norway, 9037 Tromsø, Norway

^b Centre for International Health, Department of Global Public Health and Primary Care, University of Bergen, 5009 Bergen, Norway

^c Department of Laboratory Medicine, University Hospital of North Norway, 9038 Tromsø, Norway

^d Hospital Público Materno Infantil de Salta, Sarmiento 1301, 4400 Salta, Argentina

^e Department of Public Health and Nursing, Norwegian University of Science and Technology (NTNU), 7491 Trondheim, Norway


^f Department of General Hygiene I.M. Sechenov First Moscow State Medical University (Sechenov University), 119992 Moscow, Russia

^g School of Health Systems and Public Health, Faculty of Health Sciences, University of Pretoria, Pretoria 0002, South Africa

HIGHLIGHTS

- Maternal serum PFAS from the world's southernmost city and the Andes mountain
- Low detection limits with a high percentage of detected samples for the 24 PFAS
- High detection frequencies of PFBA and PFHxA, which mainly are uncommonly in humans
- Source appointment indicates region-specific PFAS exposure sources despite low concentrations.

GRAPHICAL ABSTRACT



ARTICLE INFO

Editor: Lidia Minguéz Alarcon

Keywords:

Biomonitoring
Source appointment
PFAS
Pregnancy
Predictors
South America

ABSTRACT

Background: Per- and polyfluoroalkyl substances (PFAS) are a group of synthetic chemicals with potential adverse health effects. Information concerning PFAS concentrations in relation to pregnancy is scarce in South America and non-existent in Argentina.

Aim: We aimed to investigate an extended maternal PFAS profile herein serum concentrations in a regional and global view, source appointment, and determinants in Argentinean women.

Methods: A cross-sectional study with a sampling period from 2011 to 2012 included 689 women from Ushuaia and Salta in Argentina. Serum samples collected two days postpartum were analyzed by ultra-high pressure liquid chromatography coupled to electrospray negative ionisation tandem-quadrupole mass-spectrometry. Principal Component Analysis (PCA) following absolute principal component score-multiple linear regression (APCS-MLR) was used for PFAS source appointments. Determinants of PFAS were explored through a MLR approach. A review of previous studies within the same period was conducted to compare with present levels.

* Corresponding author.

E-mail addresses: solrunn.hansen@uit.no (S. Hansen), shanshan.xu@uib.no (S. Xu), sandra.huber@unn.no (S. Huber), jon.o.odland@ntnu.no (J.Ø. Odland).

<https://doi.org/10.1016/j.scitotenv.2024.170096>

Received 25 September 2023; Received in revised form 9 January 2024; Accepted 9 January 2024

Available online 13 January 2024

0048-9697/© 2024 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Results: Argentinean PFAS concentrations were the lowest worldwide, with PFOS (0.74 ng/mL) and PFOA (0.11 ng/mL) as the dominant substances. Detection frequencies largely aligned with the compared studies, indicating the worldwide PFAS distribution considering the restrictions. The PCA revealed region-specific loading patterns of two component groups of PFAS, a mixture of replaced and legacy substances in Ushuaia and long-chain in Salta. This might relate to a mix of non-diet and diet exposure in Ushuaia and diet in Salta. Region, age, lactation, parity, household members, migration, bottled water, and freshwater fish were among the determinants of various PFAS.

Conclusion: This is the first study to monitor human PFAS exposure in Argentina. Maternal PFAS concentrations were the lowest observed worldwide in the same period. Exposure contributions are suggested to be affected by restrictions and substitutions. Given the limited population-based studies and the emergence of PFAS, it is essential to conduct further monitoring of PFAS in Argentina and South America.

1. Introduction

Per- and polyfluoroalkyl substances (PFAS) are synthetic chemicals widely used in industrial and consumer applications. They are environmentally persistent with a potential for global long-range transport and are bio-accumulative and toxic in living organisms (Buck et al., 2011). Thus, PFAS exposure raises health concerns (Fenton et al., 2021), with particular concern apparent to maternal and child issues, both from a short- and long-term perspective (Blake and Fenton, 2020; Rickard et al., 2022). Due to growing concerns, global regulation and phase-outs have been initiated from the 2000s for legacy PFAS, like perfluorooctane sulfonic acid (PFOS) and perfluorooctanoic acid (PFOA) (Brennan et al., 2021), and lately on perfluorohexane sulfonic acid (PFHxS); all listed in the Stockholm Convention in 2009, 2019, and 2023, respectively (Stockholm Convention, 2023).

Abbreviations

AFFFs	aqueous film-forming foams
PFASs	per- and polyfluoroalkyl substances
PFCAs	perfluoroalkyl carboxylic acids
PFSAs	perfluoroalkyl sulfonic acids
FTSAs	fluorotelomer sulfonic acids
PFBS	perfluorobutane sulfonic acid
PFPeS	perfluoropentane sulfonic acid
PFHxS	perfluorohexane sulfonic acid
PFHpS	perfluoroheptane sulfonic acid
PFOS	perfluorooctane sulfonic acid
PFNS	perfluorononane sulfonic acid
PFDS	perfluorodecane sulfonic acid
PFDoDS	perfluorododecane sulfonic acid
PFBA	perfluorobutanoic acid
PFPeA	perfluoropentanoic acid
PFHxA	perfluorohexanoic acid
PFHpA	perfluoroheptanoic acid
PFOA	perfluorooctanoic acid
PFNA	perfluorononanoic acid
PFDA	perfluorodecanoic acid
PFUnDA	perfluoroundecanoic acid
PFDoDA	perfluorododecanoic acid
PFTriDA	perfluorotridecanoic acid
PFTeDA	perfluorotetradecanoic acid
FOSA	perfluorooctane sulfonamide
4:2 FTSA	4:2 fluorotelomer sulfonic acid
6:2 FTSA	6:2 fluorotelomer sulfonic acid
8:2 FTSA	8:2 fluorotelomer sulfonic acid
10:2 FTSA	10:2 fluorotelomer sulfonic acid

The maternal body burden of PFAS has been investigated worldwide, mostly in developed regions, and detected as ubiquitous in varying concentration ranges with PFOS and PFOA as predominant substances (Bjerregaard-Olesen et al., 2017; Liu et al., 2020). Restrictions on production and use have reflected declining human temporal trends of

PFOS and PFOA. From the 1990s until 2018, in the USA, Canada, Germany, Sweden, and Norway, human PFOS has declined from around 60 % to 75 %, also for PFOA, although with more variations across nations (Fan et al., 2022). Similar declines were observed in Australia from 2002 to 2011 (Toms et al., 2014). Also, maternal declining trends were observed in Denmark (2008–2013) (Bjerregaard-Olesen et al., 2016), Japan (2003–2011) (Okada et al., 2013), and in most studies in a global review (1972–2016), except an upward effect for Inuits in Alaska (Liu et al., 2020).

Simultaneous, various trends have been observed in human exposure to replacement PFAS, which include both long-chain substances (with nine or more carbon-fluor chains) and short-chain substances (less than C8) (Fan et al., 2022; Liu et al., 2020; Okada et al., 2013). For example, mostly delayed downward trends were reported for PFHxS, but an upward trend in Sweden for PFHxS and perfluorodecanoic acid (PFDA) (1972–2016) (Liu et al., 2020). Increasing trends were also seen for PFNA perfluorononanoic acid (PFNA), PFDA, and perfluoroundecanoic acid (PFUnDA) in Inuits in Canada (2004–2017) (Caron-Beaudoin et al., 2020) and PFNA and PFDA in Japan, but solely declining trends in Denmark (Bjerregaard-Olesen et al., 2016).

Paralleling the global consensus of regulations, with exemptions, the production of PFOS and PFOA has continued in Asia, particularly in China (Wang et al., 2015). Additionally, PFOS has also been imported into Brazil for Sulfluramid manufacturing (Barbosa Machado Torres et al., 2022). Consequently, human exposure trends in China increased for PFOS until 2009 and PFOA until 2015 (Fan et al., 2022). Trends for South America are less pronounced. To our knowledge, PFAS status is limited to maternal studies from Brazil in 2011 (Souza et al., 2020) and 2017 (Espindola Santos et al., 2021) and global studies with small samples from Latin America (Fiedler and Sadia, 2021) and Brazil (Kannan et al., 2004).

Human PFAS profiles are influenced by several factors, with diet being the primary human exposure. This is followed by drinking water, while dermal absorption, dust, and air inhalation – especially personal care products and indoor environments – also contribute (Jian et al., 2017; Pérez et al., 2014; Wee and Aris, 2023). Pregnancy and lactation are regulators of PFAS through placental and lactation transfer to the baby (Olsen et al., 2009). Race, region, country of origin, and socio-economic status are other relevant determinants (McAdam and Bell, 2023). Age is reported with an inconclusive relationship to PFAS, mainly explained by production, emission, regulation, year of peak, and half-life (McAdam and Bell, 2023; Quinn and Wania, 2012). Exposure contamination grade and the properties of each substance affect the body burden. Humans are exposed to a mixture of PFAS and with variety within and across nations (EFSA, 2020). Profiles of human PFAS exposure might be helpful to understand sources and exposure pathways (Hu et al., 2018), and thus, it is essential to describe the population status beyond legacy substances such as PFOS and PFOA (Sunderland et al., 2019).

This study aimed to enhance understanding of PFAS exposure in South America by providing a comprehensive serum PFAS profile of postpartum women in two Argentinean regions. We include a global

comparison and identify sources and predictors of PFAS levels in our study.

2. Material and methods

2.1. Description of the study area and data collection

The EMASAR study (Estudio del Medio Ambiente y la Salud Reproductiva; Study on the Environment and Reproductive Health) aimed to investigate contaminants related to pregnancy in Argentina. Study areas were designated to Ushuaia, the world's southernmost city, and Salta in the northwestern highland of Argentina. Responsible for the project were UiT, The Arctic University of Norway, Tromsø, and Stavanger University Hospital, both in Norway. Local partners were the private institution Clínica San Jorge in Ushuaia and the Hospital Público Materno Infantil in Salta.

The study period ranged from April 2011 to November 2011 in Ushuaia and June 2011 to March 2012 in Salta and included 698 women (200 from Ushuaia and 498 from Salta). For this present PFAS study, nine participants were excluded due to damaged serum vials. Non-fasting blood samples were obtained at a median of one day with a range of 0–3 days following delivery. Information collected through personal interviews and medical records covered socio-economy, pregnancy, health and lifestyle conditions, diet, and environmental aspects. A detailed description of the study has been given elsewhere (Okland et al., 2017).

Local approvals of the EMASAR study were given by the Ethics Committee of the Salta Medical Association and the Ministries of Health in both the Province of Salta and the Province of Tierra del Fuego (#2010/7317). The Norwegian Regional Committee for Medical and Health Research Ethics (REC North) approved the study (#2011/706). The study was conducted in accordance with the Helsinki Declaration. Informed consent was obtained from the participating woman.

2.2. Chemical analysis of PFAS

Maternal blood was collected in BD Vacutainers® (BD SST II Plus Advance 10/8.5 mL), sampled and treated according to the instruction leaflet, and centrifuged at 2000 relative centrifugal force (RCF) for 10 min. The serum was subsequently apportioned into n-hexane/acetone pre-rinsed glass vials. Samples were shipped in a frozen state to the Biobank at the UiT The Arctic University of Norway for storage at minus 30 degrees Celsius. In 2019, samples were analyzed for 24 PFAS and 6 accompanying linear substances.

The analytical work was conducted at The Environmental Pollution Laboratory, Department of Laboratory Medicine, University Hospital of North Norway, Norway. Aliquots of 50 µL serum samples were extracted by solid-phase microextraction on an Oasis WAX-µElution plate (30 µm, Waters, Milford, USA) and analyzed by ultra-high pressure liquid chromatography coupled to electrospray negative ionisation tandem-quadrupole mass-spectrometry (Waters Acquity UPLC Xevo TQ-S system, Milford, MA, USA) as described previously (Huber and Brox, 2015). The limit of detections (LODs) was set as concentrations calculated by the Targetlynx-software for each sample (LODi) and each PFAS with a signal-to-noise ratio of 3 divided by the related sample amount. For quality assurance, four blank samples, four SRM 1958 (NIST, Gaithersburg, MD, USA), and three bovine serum samples (Sigma Aldrich, Steinheim, Germany) were prepared and analyzed within each batch of 96 samples to control for background and carry-over effects. All the quality controls were within the acceptance limits. Correlation coefficients of variation were < 10 % for all measured PFAS. The limit of detection was in the range of 0.0034 ng/mL to 0.031 ng/mL for analyzed PFAS; five substances were not detected (Table S3). For PFBA and PFOA, a batch-wise blank subtraction was performed. Concentrations detected in the blank samples of each individual sample preparation batch were used and subtracted from the concentration measured in the samples

from the EMASAR study as follows: sample concentration – (average blank concentration + 3 × standard deviation blank concentration). Subtracted blank concentrations were around 0.50 ng/mL for PFBA and 0.11 ng/mL for PFOA, respectively. All PFAS analyses were within the acceptable ranges of the international quality control program: the Arctic Monitoring and Assessment (AMAP) Ring Test for Persistent Organic Pollutants in Human Serum (organized by Centre du Toxicologie du Québec (CTQ), Institut National de Santé Publique du Québec, Canada).

2.3. Statistical analysis

Statistical analyses were done using the IBM SPSS Statistics for Windows statistical package version 28 (SPSS Inc. Chicago, IL, USA). Figures were performed in R. Significant levels were set at $p < 0.05$. PFAS concentrations below LOD were replaced by individual LOD divided by the square root of 2 (Hornung and Reed, 1990). According to the Kolmogorov-Smirnov test, most substances deviated from a normal distribution and were harmonized during \log_{10} transformation. Descriptive analyses described raw data, and the *t*-test, chi-square, or Mann-Whitney test explored variations. The Pearson correlation (*r*) assessed correlations between the PFAS.

Through the dimensional reduction method principal component analysis (PCA), we explored the linear relationship between the log-PFAS by region. Restricted to substances detected above 60 %, only one cluster was revealed for Salta, and rotation was not performed. Thus, the detection frequencies were extended. Initially, ten substances were included, followed by the exclusion of perfluorobutanoic acid (PFBA) for Ushuaia only and perfluoroheptanoic acid (PFHpA) due to a correlation below 0.3 (Table S5). Orthogonal (varimax) rotation was used. Both sampling adequacy measured by Kaiser-Meyer-Olkin (KMO 0.8) and Bartlett's test of sphericity ($p < 0.001$) met the correlation criteria for PCA. Kaiser's criterion ≥ 1 was set in selecting important factors. Factor loading was evaluated as strong (> 0.75), moderate (0.5–0.75), or weak (< 0.05 –0.3) (Cho et al., 2022). Next, the absolute principal component score-multiple linear regression (APCS-MLR) model was performed as described in detail by others (Cho et al., 2022; Wallis et al., 2023). The individual PFAS contribution (dependent variable) in percentage (%) of each factor loading (independent variable) from the PCA was explored in a multiple linear regression (MLR) model. Percentage calculations were based on the equation: $i (\%) = 100 * (Bi / \sum niBi)$, where Bi is the beta coefficient for the individual PFAS (Wallis et al., 2023). To account for the contribution of negative values, negative beta coefficients were treated as positive values (Haji Gholizadeh et al., 2016).

MLR, using a stepwise procedure with backward elimination, described the relationships between seven frequently detected log 10 transformed substances and selected independent factors according to McAdam and Bell (2023). Initial variables were age (year), parity (parous - multiparous), lactation (interval), region (Ushuaia - Salta), people in the household (number), education (primary/secondary - tertiary/university), migration inland/abroad (no-yes), bottled water almost daily (no-yes), dietary factors during pregnancy (never/seldom - weekly/daily): freshwater fish, saltwater fish, seafood, meat, poultry, processed meat, egg, dairy products (milk), butter/cheese, vegetables (bean, tomato, garlic), fruit, bread. Other vegetables and items were not included due to intake below 3 %. Finally, the models were evaluated through diagnostic plots, variance inflation factor (VIF), Mahalanobis and Cook's distance, and casewise diagnostics of standardized residuals with exclusions of the most influential outliers (Field, 2009). The modified Breusch-Pagan test checked for abruption of homogeneity ($p < 0.05$) and with robust standard errors corrected for heteroskedasticity (Mansournia et al., 2020).

For a global comparison of PFAS concentrations, a review of publications published after 2010 was performed on PubMed. The search string was [PFAS per-and polyfluoroalkyl substances pregnancy] in

addition to blood, whole blood, serum, or plasma with 438, 383, 174 or 22 matches, respectively. Additionally, reference lists in review publications were searched. After searching the sampling period, 16 studies collected between 2010 and 2013 were identified. In the aftermath, a recent publication confirmed the included studies (Kuo et al., 2023).

3. Results

3.1. Background characteristics

Compared to those in Salta, women living in Ushuaia were notably older, had fewer children, and had shorter lifetime breastfeeding but the proportion of first-time mothers was similar. Ushuaian women also had higher education and more household members (Table 1). Regarding diet during pregnancy, those from Ushuaia had a significantly higher intake of bottled water, dairy, and marine food, but a lower intake of processed meat, eggs, bread, pasta/cereals, and sugar. The consumption of freshwater fish, meat, vegetables, or fruits did not differ between locations (Supplementary Table S2). Characteristics have been described in detail elsewhere (Okland et al., 2017).

3.2. Serum PFAS detection frequencies and concentrations

The distributions of PFAS in maternal serum samples varied fairly between Ushuaia and Salta (Tables 2, S3). Of the 24 PFAS analyzed, seven substances had a detection frequency above 80 %. Notably, PFOS was detected in all samples of the study group. In Ushuaia, PFOA showed a similar universal presence, followed by the frequency order of PFHxS = PFNA > PFBA > PFDA. In Salta, the order was PFHxS > perfluorohexanoic acid (PFHxA) > PFNA > PFOA > PFDA, with the highest detection frequencies in Ushuaia. Striking were the detection differences between PFBA and PFHxA with alternating detection between the cities. Substances with substantially lower detection are reported in Tables 2, S3, and S4.

Table 1

Characteristic of the study population by region. The EMASAR study in Argentina, 2011–2012.

	Ushuaia n = 193		Salta n = 496		P-value
	Mean, SD	or n (%)	Mean, SD	or n (%)	
Age, year	28.8 ± 6.5		24.7 ± 6.2		<0.001 ^a
Parity, number	1.9 ± 0.97		2.2 ± 1.5		0.005 ^b
para 1	78 (40.4)		221 (44.6)		0.001 ^c
para 2	72 (37.3)		120 (24.2)		
para >2	43 (22.3)		155 (31.3)		
Breastfeeding, months, overall	12.3 ± 18.2		18.6 ± 27.3		<0.001 ^b
Breastfeeding, months, multipara	20.6 ± 19.7		34.0 ± 29.0		<0.001 ^b
BMI, postpartum	28.0 ± 4.1		26.1 ± 4.3		<0.001 ^a
Smoking during last year	54 (28.0)		127 (26.3)		0.525 ^c
Education					
Primary	7 (3.6)		161 (32.5)		0.001 ^c
Secondary	93 (48.2)		283 (57.2)		
Tertiary	55 (28.5)		39 (7.9)		
University	38 (19.7)		12 (2.4)		
Marital status					
Married/cohabitant	172 (89.1)		337 (73.9)		<0.001 ^c
Single/divorced	21 (10.9)		159 (32.1)		
Permanent job, yes	128 (66.3)		86 (31.2)		<0.001 ^c
People living in your home, number	4.2 ± 1.4		6.6 ± 3.5		<0.001 ^b
Country/province born					
Born current province	63 (32.6)		437 (88.1)		
Born other provinces	124 (62.3)		44 (8.9)		
Born other nations	6 (3.1)		15 (3.0)		

SD, standard deviations.

^a T-test.

^b Welch test.

^c Chi-square test.

Table 2 Maternal serum concentrations (ng/mL) of per- and polyfluoroalkyl substances by region. The EMASAR study in Argentina, 2011–2012^a.

Region	GM	AM	Min	25%ile	75%ile	Max	% >	LOD	PFBA	PFHxA	PFHpA	PFOA	PFNA	PFDA	PFUnDA	Linear PFHxS	Sum PFHxS	Linear PFHpS	Sum PFHpS	Linear PFOS	Sum PFOS	Sum PFAS	Sum PFSA	Sum PFCA	Sum PFSA and PFCA
Ushuaia (n = 193)	0.07	0.09	0.01	0.05	0.11	0.29	88.1	19.7	0.07	0.04	0.008	0.24	0.06	0.02	0.02	0.06	0.18	0.007	0.01	0.42	0.84	1.73	1.07	0.60	1.70
	0.09	0.08	0.01	0.05	0.11	0.29	88.1	19.7	0.09	0.06	0.01	0.29	0.07	0.03	0.02	0.09	0.23	0.01	0.02	0.51	1.02	1.97	1.28	0.66	1.94
	0.01	0.01	0.004	0.04	0.04	0.06	0.05	0.007	0.04	0.004	0.004	0.04	0.06	0.005	0.003	0.004	0.006	0.001	0.001	0.14	0.28	0.69	0.38	0.23	0.67
	0.05	0.05	0.007	0.006	0.15	0.40	0.07	0.006	0.04	0.004	0.006	0.15	0.04	0.02	0.008	0.04	0.12	0.003	0.003	0.28	0.58	1.26	0.76	0.42	1.23
	0.11	0.17	0.01	0.38	0.08	0.8	0.29	0.06	0.08	0.08	0.01	0.38	0.08	0.03	0.03	0.09	0.27	0.02	0.034	0.55	1.13	2.23	1.44	0.80	2.20
	0.29	0.48	0.06	1.28	0.42	1.28	0.06	0.06	0.42	0.42	0.06	1.28	0.42	0.20	0.18	3.20	3.30	0.19	0.28	6.02	14.4	15.90	15.2	2.37	15.85
	88.1	88.1	58.5	58.5	85.5	85.5	85.5	85.5	99.5	98.4	55.4	100	99.5	85.5	53.9	98.4	99.5	55.4	100	100	100	100	100	100	100
	0.04	0.06	0.01	0.08	0.03	0.08	0.01	0.01	0.03	0.04	0.007	0.08	0.03	0.02	0.007	0.04	0.21	0.004	0.004	0.40	0.70	1.37	0.97	0.35	1.35
	0.06	0.08	0.01	0.11	0.04	0.11	0.01	0.01	0.04	0.06	0.009	0.11	0.04	0.03	0.009	0.06	0.26	0.007	0.010	0.56	0.97	1.66	1.24	0.39	1.64
	0.05	0.07	0.008	0.09	0.04	0.04	0.008	0.008	0.04	0.04	0.006	0.09	0.04	0.02	0.006	0.04	0.22	0.004	0.004	0.39	0.70	1.38	0.97	0.37	1.36
	0.08	0.08	0.002	0.004	0.004	0.004	0.002	0.002	0.004	0.002	0.001	0.004	0.004	0.004	0.001	0.002	0.002	0.001	0.001	0.06	0.11	0.24	0.12	0.07	0.22
	0.02	0.04	0.006	0.05	0.027	0.16	0.02	0.006	0.026	0.026	0.002	0.05	0.027	0.016	0.006	0.026	0.147	0.002	0.002	0.27	0.50	1.09	0.74	0.28	1.07
	0.09	0.10	0.02	0.14	0.03	0.03	0.09	0.02	0.03	0.08	0.008	0.14	0.48	0.03	0.008	0.67	0.32	0.008	0.013	0.54	0.92	1.74	1.25	0.48	1.72
	0.23	0.27	0.10	2.90	1.14	1.14	0.27	0.10	1.28	0.81	0.45	2.90	1.14	0.55	0.13	0.81	1.28	0.45	0.53	33.1	60.97	60.5	60.5	5.03	60.95
	60.3	96.4	32.1	89.7	85.5	85.5	32.1	32.1	98	97.2	24.8	100	92.7	85.5	19.2	97.2	98	24.8	100	100	100	100	100	100	100
	<0.001	0.693	<0.001	<0.001	0.616	0.616	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	0.616	<0.001	<0.001	<0.001	<0.001	<0.001	0.326	<0.001	<0.001	0.025	<0.001	<0.001
	0.693	0.693	<0.001	<0.001	0.616	0.616	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	0.616	<0.001	<0.001	<0.001	<0.001	<0.001	0.326	<0.001	<0.001	0.025	<0.001	<0.001

^a The limit of detection was in the range of 0.0034 ng/mL to 0.026 ng/mL for all PFAS; Substances with lower detection frequency (<12 %) are reported in the Supplementary file.

^b Group comparison for PFAS detected >50 %, Mann-Whitney test; AM, arithmetic mean; GM, geometric mean; max, maximum; min, minimum; %ile, percentile; abbreviations of the substances, see Table S1.

Median serum PFOS, PFOA, PFBA, and PFNA were significantly highest in Ushuaia women, while PFHxS were highest in Salta women (Table 2). Of the total PFAS, the highest sum concentration was observed in Ushuaia ($p < 0.03$, Table 2). The seven dominating substances accounted for 93 % of the total PFAS in both places. Further, the perfluoroalkyl sulfonic acids (PFSA; PFOS, PFHxS and perfluoroheptane sulfonic acid (PFHpS)) were the leading fraction of total PFAS, comprising 64.9 % in Ushuaia and 75 % in Salta. The fraction of linear PFOS to the total PFOS was 49.8 % and 56.9 %, and linear PFHxS to the total PFHxS was 40.9 % and 21.2 % in Ushuaia and Salta, respectively ($p < 0.001$, data not shown).

3.3. Comparison of Argentinean PFAS profile to global studies

Covering worldwide maternal blood PFAS biomonitoring around a similar period, primarily representing the Northern Hemisphere, present Argentinean concentrations of PFAS were of the lowest reported (Table 3a). In all studies, PFOS was the dominating substance, followed by PFOA. The Argentinean PFOS and PFOA were comparable to Tanzania. Argentinean PFOS were ten times lower than neighboring Brazil, 3–7 times lower than North America (USA > Canada), 4–10 times lower than Europe (Denmark > France), 5 to 15 times lower than both Asia (China > South Korea > Japan) and Inuit women (Greenland > Canada). For PFOA, our levels were half of Brazil, 10–15 times lower than North America and Europe, and >12 times lower than Asia but 350 times lower than China. The remaining PFAS varied across nations. Long-chain PFAS were highest in Inuit women in Greenland and Canada but comparable to China. In general, PFOS and PFOA were detected in all study groups except for PFOA, with detection of 68 % in Brazil and 90 % in Tanzania and Salta. Other PFAS were highly detected, although with more variations across the countries (Table 3b).

3.4. PFAS loading profile and source contribution across regions

In the PCA, two loading components satisfying the Kaisers criterion ≥ 1 and explained around 40 % and 20 % of the variance for their respective component groups. We observed distinct PFAS rotated loading patterns between the two regions (Fig. 1, Table S6). The contribution of each substance's loadings in the APCS-MLR model is presented in brackets (Fig. 2, Table S7). In Ushuaia, component 1 was strongly influenced by PFHxA (85.5 %), PFHpS (78.6 %), and moderately by PFOS (74.0 %), PFHxS (90.5), and PFOA (59.3 %) and PFHpS (86.3 %). Component 2 was clustered strongly by PFDA (88.4 %) and PFUnDA (89.6 %) and moderately by PFNA (53.2 %). In Salta, component 1 was moderately controlled by PFDA (76.4 %), PFUnDA (82.9 %), PFOS (65.8 %), and PFHpS (86.3 %), and component 2 was strongly influenced by PFHxA (82.8 %) and moderately by PFBA (88.8 %), PFOA (57.7 %), and PFHxS (74.9 %). Additionally, The MLR analysis of moderate to strong PFAS included in the factor loadings to each APCS score supported the clustering profile with an explained variance of 92–97 % (Table S7).

3.5. Determinants of maternal PFAS concentrations

Fig. 3 demonstrates the determinants of PFAS discovered through MLR. Compared to Ushuaia, living in Salta was associated with higher concentrations of PFHxA (+74 %) and PFHxS (37 %), but similar lower concentrations of PFOA (–61 %), PFBA (–43 %) and PFNA (–27 %). Age elevated PFOS, PFHxA, PFOA, and PFNA by up to 24 % by five years. Advancing parity from mono- to multiparous reduced all substances up to 45 % except for PFBA and PFDA. Lactation decreased PFOS, PFOA, and PFHxA by around 6 % per half year. Elevating household members lowered PFOA, PFOS, and PFDA by 2 %. Concerning diet, a swift from never/seldom to weekly/daily intake, freshwater fish contributed to PFOS by 12 % and PFDA by 19 %, and fruit and egg elevated PFHxA up to 30 %. Inverse associations were revealed for food

Table 3a
Global comparison of PFAS (ng/mL) in maternal serum, plasma, or whole blood in selected studies during 2010–2013.

Year	Country	n	PFBA	PFHxA	PFHpA	PFOA	PFNA	PFDA	PFUnDA	PFHxS	PFHpS	PFOS	Central tendency	Material	Reference
2013	France (Toulouse)	100				1.05	0.43	ND	ND	0.62	ND	3.07	Median	S	Cariou et al., 2015
2012	China (Tanjing)	141		0.45	0.18	4.0*	1.05	1.33	0.61	0.38	6.7 ^a	6.7 ^a	GM	S	Jiang et al., 2014
2012	China (Shanghai)	981				19.6	1.79	2.05	1.54	2.73	10.8	10.8	GM	P	Tian et al., 2018
2012	Canada (Nunavik)	111	ND	ND	ND	0.67	2.00	0.45	0.44	0.35	3.80	3.80	GM	S	Caron-Beaudoin et al., 2020
2012	Tanzania (Arusha)	48	ND	ND	ND	0.21	0.17	0.14	NC	NC	0.50	0.50	Median	P	Müller et al., 2019
2011–2012	Argentina (Ushuaia)	193	0.07	0.01	0.008	0.24	0.03	0.02	0.02	0.18	0.007	0.84	GM	S	Present study
2011–2012	Argentina (Salta)	496	0.04	0.008	0.01	0.08	0.06	0.02	0.007	0.21	0.004	0.70	GM	S	Present study
2011–2012	USA (Colorado)	NA				1.80	0.50	0.80		1.00		5.30	Median	S	Zell-Baran et al., 2023
2011	Canada (10 cities)	95				1.38				1.04		3.65	GM	P	Fisher et al., 2016
2011	France (national)	277	ND	ND	ND	1.50	0.52	0.26	NC	1.04	NC	3.1	GM	S	Dereumeaux et al., 2016
2011	Japan (Hokkaido)	30		<0.1	<0.1	1.27	1.26	0.69	1.3	0.33		3.52	Median	P	Okada et al., 2013
2011	South Korea (Gyeongbuk)	70				2.62				1.21		9.37	Median	S	Lee et al., 2013
2010–2013	China (Shandong)	369			0.06	39.3	0.78	0.52	0.46	0.32		4.25	GM	S	Wang et al., 2019
2010–2013	Greenland (national)	207		NC	NC	1.19	1.30	0.72	1.60	0.7	0.19	10.2	Median	S	Long et al., 2015
2010–2012	Denmark (Odense)	1436				1.68	0.64	0.29	NC	0.36		7.5	Median	S	Birukov et al., 2021
2010–2011	Brazil (Ribeirão Preto)	243	ND	ND	ND	0.46	NC	NC	NC	NC		7.8	GM	WB ^b	Souza et al., 2020
2010–2011	Canada (Winnipeg)	247	ND	ND	ND	0.89	0.37	0.13	0.069	0.44		2.2	Median	P	Workman et al., 2019
2010	Canada (10 cities)	943				1.62				1.07		4.3	GM	P	Fisher et al., 2016

GM, geometric mean; med., median; n, number of participants; NA, not available; ND; not detectable; NC, quantified, but GM or median concentration not reported/calculated due to low detection.

^a Median.

^b Whole blood concentration multiplied by two for comparison with serum or plasma concentrations as described in Ehresman et al., 2007.

Table 3b

Global comparison of detection frequencies of PFAS (ng/mL) in maternal serum, plasma, or whole blood in selected studies during 2010–2013.

Country	Year	Detection	PFBA	PFHxA	PFHpA	PFOA	PFNA	PFDA	PFUnDA	PFHxS	PFHpS	PFOS	Material	Reference
France (Toulouse)	2013	% > LOD				100	98	93	79	99	50	100	S	Cariou et al., 2015
China (Tanjin)	2012		100	100	100	100	100	100	100	100	100	100	S	Jiang et al., 2014
China (Shanghai)	2012	% > LOD				100	100	100	99.9	100		100	P	Tian et al., 2018
Canada (Nunavik)	2012	% > LOD	ND	ND		100	100	98	92	92		100	S	Caron-Beaudoin et al., 2020
Tanzania (Arusha)	2012	% > LOD		ND	ND	90	81	85	27	38		100	P	Müller et al., 2019
Argentina (Ushuaia)	2011–2012	% > LOD	88.1	58.5	19.7	100	99.5	85.5	53.9	99.5	55.4	100	S	Present study
Argentina (Salta)	2011–2012	% > LOD	60.3	96.4	32.1	89.7	92.7	85.5	19.2	98	24.8	100	S	Present study
USA (Colorado)	2011–2012	% > LOD				99.7	98.2	64.8		98.5		99.3	S	Zell-Baran et al., 2023
Canada (10 cities)	2011	% > LOD				100				97.9		100	P	Fisher et al., 2016
France (national)	2011	% > LOQ	ND	ND	0.4	100	100	67.9	30.3	99.6	7.2	100	S	Dereumeaux et al., 2016
Japan (Hokkaido)	2011	% > LOD		20	50	100	100	100	100	76.7		100	P	Okada et al., 2013
South Korea (Gyeongbuk)	2011					100			100			100	S	Lee et al., 2013
China (Shandong)	2010–2013	% > LOD			85.1	100	100	100	100	99.7		100	S	Wang et al., 2019
Greenland	2010–2013	% > LOQ		ND	16.3	100	100	100	100	100	86.8	100	S	Long et al., 2015
Denmark (Odense)	2010–2012	% > LOQ				100	100	100		96		100	S	Birukov et al., 2021
Brazil (Ribeirão Preto)	2010–2011	% > LOD		ND	ND	67.9	11.1	1.65	0.41	0.41		100	WB ^b	Souza et al., 2020
Canada (Winnipeg)	2010–2011	% > LOQ		ND	ND	100	96	94	85	89		100	P	Workman et al., 2019
Canada (10 cities)	2010	% > LOD				99.7				97.1		99.7	P	Fisher et al., 2016

LOD, limit of detection; LOQ; limit of quantification; ND, not detectable; S, serum; P, plasma; WB, whole blood.

items to mostly all substances, except PFOA and PFHxS, with no dietary relationship. Daily bottled water intake elevated PFHxA by 27 % and PFOA by 16 %. Table S9 shows the detailed MLR results.

4. Discussion

To our knowledge, this is the first study observing PFAS profiles in populations living in Argentina. Analyzing a broad spectrum of PFAS in the context of source contribution brings further novelty to understanding the regional human exposure to PFAS.

4.1. Global comparison of PFAS profile

Present maternal serum concentrations of PFAS were low. PFOS and PFOA were far below the limit with no risk of adverse health effects (5 µg/L and 2 µg/L, respectively) according to Germany Environment Agency (Umweltbundesamt, 2023). Less than a handful exceeded the action limit for women of childbearing age (10 µg/L and 5 µg/L, respectively).

Although the lowest maternal PFAS observed, the Argentinean detection frequencies aligned with the above-referred global studies and reflected the widespread distribution of PFAS. Clearly, the findings in our Argentinean samples demonstrate low maternal exposure. Variations in detection frequencies and concentrations across the countries likely reflect the production, regulations, and phase-out. In contrast to the Western world, but equal to Tanzania (Müller et al., 2019), Argentina has the status of a developing country, and to our knowledge, historically, there has been no PFAS production. North America was the

first to introduce PFOS restrictions and phase-out from 2001, and Europe from around five years later (Brennan et al., 2021). Meanwhile, PFAS production has been shifted from the USA, Europe, and Japan to an extended production of PFOS and PFOA in China (Zhang et al., 2012) and Sulfuramid in Brazil (Guida et al., 2023; Löfstedt Gilljam et al., 2016). This might explain the lower profile in the USA (Zell-Baran et al., 2023) and Canada (Fisher et al., 2016; Workman et al., 2019) to Europe (Birukov et al., 2021; Cariou et al., 2015; Dereumeaux et al., 2016), and Asia (Lee et al., 2013; Wang et al., 2019), and the PFOA concentrations in China likely explained by local contamination (Jiang et al., 2014; Wang et al., 2015; Wang et al., 2019), and the 10-fold higher PFOS in Brazil (Souza et al., 2020) than in Argentina. Further, the distinct PFAS profile in Inuits living in remote Greenland and Canada has been explained by contamination of the marine diet (Caron-Beaudoin et al., 2020; Long et al., 2015).

Considering the short half-life of days to one month for PFBA and PFHxA (EFSA, 2020; Luz et al., 2019), our high PFBA and PFHxA detection frequencies were unexpected. As in the comparison, both substances have, to a small degree, been analyzed in human populations and with infrequent detection, primarily undetected, as reviewed by others (Anderson et al., 2019; EFSA, 2020; Lee and Mabury, 2011). Given the low bioaccumulation, our findings indicate ongoing but low exposure in agreement with the dominance of aquatic biota in Argentina (Llorca et al., 2012).

4.2. Source appointment and contributing factor of PFAS concentrations

Consequences of restriction and phase-out partly align with findings




Fig. 1. Rotated component plot of log-transformed PFAS by region in the EMASAR study, 2011–2012.

Extraction Method: Principal Component Analysis. Rotation Method: Varimax with Kaiser Normalization. Rotation converged in 3 iterations and strong or moderate factor loadings >0.5 are visible in the plot. For details of the analyses, see Table S5. For abbreviations of the PFAS, see Table S1.

from the PCAs with source compounds dominated by short-chain substances used as replacements for legacy PFAS. The region-specific loading patterns of PFAS had distinct contributions (80–90 %) to the specific sources. In Ushuaia, compound 1 was a mixture of replaced and legacy substances across functional groups and chain length (Buck et al., 2011). The dominance of PFHxA, PFHpS, and PFHxS might indicate exposure from consumer products. As indicated by others, the dominance of short-chain substances likely reflects the shift from legacy PFAS to its replacements (Pérez et al., 2014). Short-chain PFAS have been substituted in consumer products, which significantly affects human uptake through ingestion of dust and drinking water (EFSA, 2020; Zheng et al., 2023). Also, due to the legislative control of PFOS and PFOA and

the lower bioaccumulation of PFOA, these two substances may indicate the preference for terrestrial animal food. Thus, dietary contribution seems plausible. The secondary source with solely long-chain PFAS is suggested to fish intake. Bioaccumulation increases by chain length, with the highest bioaccumulative ability in aquatic species (i.e., freshwater $>$ marine origin) compared to terrestrial species (Augustsson et al., 2021). Similarly, PFCA clusters with C9–11 were revealed in a North Atlantic population with high seafood consumption (Hu et al., 2018). In Salta, compound 1, with the dominance of long-chain substances, is likely to reflect diet as the source and preferably diet with freshwater fish due to the long-chain PFAS. For cluster 2, the strong dominance of short-chain PFBA, PFHxA, and PFHxS combined with




Fig. 2. Individual PFAS contributions in % of each factor loading from the PCA analyses through a multiple linear regression model by region. Percentage calculation based on the equation: $i (\%) = 100 * (B_i / \sum B_i)$.

PFOA have a shared affinity to water, and they are linked to consumer products (Domingo and Nadal, 2019; Zheng et al., 2023). Thus, with reservations, this distinct profile of component 2 might suggest low-grade polluted drinking water.

Contributing beyond suggested sources needs to be clarified. Considering the low concentrations, local pollution is of less importance. Known from industrialization and urbanization and pollutions elsewhere are e.g., industry, releases of aqueous film-forming foams (AFFFs) from airport firefighting activities, wastewater discharges, and pesticides from agriculture – i.e., leading to polluted drinking water as well as freshwater fish (Kurwadkar et al., 2022; Wee and Aris, 2023). As previously described, there have been identified general risks to water safety in Salta (Seghezze et al., 2013), and growing urbanization with expanding activities in Ushuaia (Diodato et al., 2020; Ferreira et al., 2021).

4.3. Determinants of PFAS concentrations

After controlling for potential influential factors, regional differences for several PFAS exposures remained. Also, as previously described, there are variations between Ushuaia and Salta concerning latitudes and climate, human activity, economy, and demography (Okland et al., 2017).

Dietary fish intake is considered a major human source to PFAS, with freshwater fish dominating marine origin (Augustsson et al., 2021). In our model, freshwater fish increased PFOS and PFDA. As reviewed by Sunderland et al. (2019), the contribution of fish intake to PFAS is related to the significance of a population's diet. Relative to the high meat consumption, the freshwater fish found as a predominantly PFAS source likely reflects polluted aquatic habitat (van Asselt et al., 2011). The fact that marine intakes were restricted to Ushuaia might have attenuated the results as we observed marine products with borderline association for PFNA and PFHxA. Also, self-reported food frequency questionnaires (FFQ) and their form could be susceptible to bias and




Fig. 3. Results of linear regression model with robust standard errors of maternal log10 transformed-PFAS (ng/mL) serum concentrations with region, socio-demography, obstetric history, and diet as predictors. The EMASAR study in Argentina, 2011–2012. Backward regression with initial variables: age (10 years), parity (monoparous (ref) - multiparous), lifetime lactation (6 months), region (Ushuaia (ref) – Salta), migration (no-yes), people in household (number), education (primary/secondary – tertiary/university), bottled water daily (no-yes), dietary factors during pregnancy (never/seldom ((ref) – weekly/daily): freshwater fish, saltwater fish, seafood, meat, poultry, processed meat, egg, dairy products (milk), butter/cheese, fruit, bread, vegetables.

* The p-value of the selected predictor is < 0.05

limited accuracy (Poothong et al., 2020), as also observed in populations with high fish intake (Caron-Beaudoin et al., 2020). Moreover, the distinct pattern revealed in PCA-MLR analyses with the cluster of PFNA, PFDA, and PFUnDA might underpin the importance of marine fish intake for Ushuaia.

The contribution of specific dietary items to PFAS body burden is reported with dependency on carbon chain length and dietary patterns within geographical variations (Jian et al., 2017; Kärman et al., 2009; van Asselt et al., 2011), which was observed in our models. Also, lower tropic items were inversely related to long-chain PFAS, as animal origin was due to more short-chain substances detected. Besides, our inverse dietary effect on PFAS may reflect a specific pattern with less contamination of the particular substance (Halldorsson et al., 2008; Tian et al., 2018).

Regarding the associations with bottled water, PFAS has infrequently been detected in bottled water, explained by the grade of contamination in water sources, and plastic, (Kaboré et al., 2018; Wee and Aris, 2023), or due to specific production/filling processes (Eschazier et al., 2013). The underlying factor of household members' negative impact on PFOA, PFOS, and PFDA is unclear. However, it might reflect low socio-economic status over the years with less access to consumer products or food items (Buekers et al., 2018), while others have linked to frequent cleaning (DeLuca et al., 2023). Migration is explained by pre-exposure or habits related to origin. Age and pregnancy-related factors align as known predictors (McAdam and Bell, 2023).

4.4. Strengths and limitations

Our study encompassed a broad range of PFAS, providing a comprehensive exposure assessment in a relatively large study group. An expansive suite of PFAS is rarely present in human studies (De Silva et al., 2021; EFSA, 2020). The chemical analyses conducted in the study were highly quality and validated through participation in the Arctic Monitoring and Assessment Programme (AMAP) Ring Test (INSPQ, 2023). Standard sampling time and procedures were followed during this study by trained health-care professionals, ensuring consistency and comparability of results. The present study also has some limitations. Results for PFBA must be interpreted with care due to a limitation in analysis and only one available transition for identification (Huber and Brox, 2015). However, since the detection frequency for PFBA was outstandingly different compared to other previously analyzed and reported cohorts, this substance was worth mentioning in this publication. Next, the study was limited to two specific regions with unique socio-economic characteristics, which may restrict the generalizability of the findings to other populations in Argentina. Moreover, the statistical analyses in the MLR may be subject to biased standard errors due to the violation of homogeneity of variance. However, robust common errors were implemented to mitigate this concern. Regarding the study design and the weakness mentioned, the findings revealed are indicators of associations, not causalities.

5. Conclusion

This study addresses a knowledge gap by investigating maternal PFAS exposure in pregnant women from two distinct geographical areas in Argentina. Low PFAS concentrations were detected, representing the lowest observed in a global comparison within the same period. Regional diversity in PFAS exposure was observed, and potential exposure sources were suggested to reflect the global regulations. In line with previous research, dietary predictors were compound-specific, and age and pregnancy-related factors were regulators. Considering the need for human PFAS information and temporal trends in South America, further maternal monitoring in Argentina and South America is warranted.

Financial support

The Norwegian Department of Foreign Affairs and the Arctic Monitoring and Assessment Programme (AMAP) (2011/706-13) funded the study. The funders had no role in developing the study design or the publication.

CRedit authorship contribution statement

Solrunn Hansen: Writing – original draft, Visualization, Validation, Methodology, Formal analysis, Conceptualization. **Shanshan Xu:** Writing – review & editing, Visualization. **Sandra Huber:** Writing – review & editing, Formal analysis. **Marisa Viviana Alvarez:** Writing – review & editing. **Jon Øyvind Odland:** Writing – review & editing, Methodology, Investigation, Funding acquisition, Conceptualization.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

The authors do not have permission to share data.

Acknowledgments

The authors highly acknowledge the participants, the participating hospitals, and the field workers' efforts. We recognize the Health Ministries of the provinces Salta and Tierra del Fuego, Argentina, for their administrative support and facilitation and Dr. Martin de la Arena for his assistance in obtaining the ethics approval and initiating the study in Argentina. We acknowledge Dra. Silvia dib Ashur, Lic. Maria José Aleman, Maria Florence Bressan, and Silvinia Matiocevic for their critical roles in the local administration. Credit to Stavanger University Hospital for their valuable initiation and effort during funding and implementation and to the administrative staff at UiT. We thank the laboratory engineers Julie S. Kleppe Strømberg, Tone F. Aune, and Christina R. Hansen at the Department of Laboratory Medicine, University Hospital of North Norway, for sample preparation and instrumental analysis and the University Hospital North Norway (UNN) and the North Norway Local Health Authority (Helse Nord) for their financial support through direct strategical funding of the Environmental Pollutant Laboratory at UNN.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.scitotenv.2024.170096>.

References

- Anderson, J.K., Luz, A.L., Goodrum, P., Durda, J., 2019. Perfluorohexanoic acid toxicity, part II: application of human health toxicity value for risk characterization. *Regul. Toxicol. Pharmacol.* 103, 10–20. <https://doi.org/10.1016/j.yrtph.2019.01.020>.
- Augustsson, A., Lennqvist, T., Osbeck, C.M.G., Tibblin, P., Glynn, A., Nguyen, M.A., Westberg, E., Vestergren, R., 2021. Consumption of freshwater fish: A variable but significant risk factor for PFOS exposure. *Environ. Res.* 192, 110284 <https://doi.org/10.1016/j.envres.2020.110284>.
- Barbosa Machado Torres, F., Guida, Y., Weber, R., Machado Torres, J.P., 2022. Brazilian overview of per- and polyfluoroalkyl substances listed as persistent organic pollutants in the Stockholm convention. *Chemosphere* 291 (Pt 3), 132674. <https://doi.org/10.1016/j.chemosphere.2021.132674>.
- Birukov, A., Andersen, L.B., Andersen, M.S., Nielsen, J.H., Nielsen, F., Kyhl, H.B., Jørgensen, J.S., Grandjean, P., Dechend, R., Jensen, T.K., 2021. Exposure to perfluoroalkyl substances and blood pressure in pregnancy among 1436 women from the Odense child cohort. *Environ. Int.* 151, 106442 <https://doi.org/10.1016/j.envint.2021.106442>.

- Bjerregaard-Olesen, C., Bach, C.C., Long, M., Ghisari, M., Bossi, R., Bech, B.H., Nohr, E. A., Henriksen, T.B., Olsen, J., Bonefeld-Jørgensen, E.C., 2016. Time trends of perfluorinated alkyl acids in serum from Danish pregnant women 2008–2013. *Environ. Int.* 91, 14–21. <https://doi.org/10.1016/j.envint.2016.02.010>.
- Bjerregaard-Olesen, C., Bossi, R., Liew, Z., Long, M., Bech, B.H., Olsen, J., Henriksen, T. B., Berg, V., Nøst, T.H., Zhang, J.J., Odland, J.O., Bonefeld-Jørgensen, E.C., 2017. Maternal serum concentrations of perfluoroalkyl acids in five international birth cohorts. *Int. J. Hyg. Environ. Health* 220 (2), 86–93. <https://doi.org/10.1016/j.ijheh.2016.12.005>.
- Blake, B.E., Fenton, S.E., 2020. Early life exposure to per- and polyfluoroalkyl substances (PFAS) and latent health outcomes: A review including the placenta as a target tissue and possible driver of peri- and postnatal effects. *Toxicology* 443, 152565. <https://doi.org/10.1016/j.tox.2020.152565>.
- Brennan, N.M., Evans, A.T., Fritz, M.K., Peak, S.A., von Holst, H.E., 2021. Trends in the regulation of per- and Polyfluoroalkyl substances (PFAS): A scoping review. *Int. J. Environ. Res. Public Health* 18 (20), 10900. <https://doi.org/10.3390/ijerph182010900>.
- Buck, R.C., Franklin, J., Berger, U., Conder, J.M., Cousins, I.T., de Voogt, P., Jensen, A.A., Kannan, K., Mabury, S.A., van Leeuwen, S.P.J., 2011. Perfluoroalkyl and polyfluoroalkyl substances in the environment: terminology, classification, and origins. *Integr. Environ. Assess. Manag.* 7 (4), 513–541. <https://doi.org/10.1002/ieam.258>.
- Buekers, J., Colles, A., Cornelis, C., Morrens, B., Govarts, E., Schoeters, G., 2018. Socio-economic status and health: evaluation of human biomonitoring chemical exposure to per- and polyfluorinated substances across status. *Int. J. Environ. Res. Public Health* 15 (12), 2818. <https://doi.org/10.3390/ijerph15122818>.
- Cariou, R., Veyrand, B., Yamada, A., Berrebi, A., Zalko, D., Durand, S., Pollono, C., Marchand, P., Leblanc, J.-C., Antignac, J.-P., Le Bizec, B., 2015. Perfluoroalkyl acid (PFAA) levels and profiles in breast milk, maternal and cord serum of French women and their newborns. *Environ. Int.* 84, 71–81. <https://doi.org/10.1016/j.envint.2015.07.014>.
- Caron-Beaudoin, É., Ayotte, P., Blanchette, C., Muckle, G., Avard, E., Ricard, S., Lemire, M., 2020. Perfluoroalkyl acids in pregnant women from Nunavik (Quebec, Canada): trends in exposure and associations with country foods consumption. *Environ. Int.* 145, 106169. <https://doi.org/10.1016/j.envint.2020.106169>.
- Cho, Y.-C., Choi, H., Lee, M.-G., Kim, S.-H., Im, J.-K., 2022. Identification and apportionment of potential pollution sources using multivariate statistical techniques and APCS-MLR model to assess surface water quality in Imjin River watershed. *South Korea. Water (Basel)* 14 (5), 793. <https://doi.org/10.3390/w14050793>.
- De Silva, A.O., Armitage, J.M., Bruton, T.A., Dassuncao, C., Heiger-Bernays, W., Hu, X.C., Kärman, A., Kelly, B., Ng, C., Robuck, A., Sun, M., Webster, T.F., Sunderland, E.M., 2021. PFAS exposure pathways for humans and wildlife: A synthesis of current knowledge and key gaps in understanding. *Environ. Toxicol. Chem.* 40 (3), 631–657. <https://doi.org/10.1002/etc.4935>.
- DeLuca, N.M., Thomas, K., Mullikin, A., Slover, R., Stanek, L.W., Pilant, A.N., Cohen Hubal, E.A., 2023. Geographic and demographic variability in serum PFAS concentrations for pregnant women in the United States. *J. Expo. Sci. Environ. Epidemiol.* <https://doi.org/10.1038/s41370-023-00520-6>.
- Dereumeaux, C., Saoudi, A., Pecheux, M., Berat, B., de Crouy-Chanel, P., Zaros, C., Brunel, S., Delamaire, C., le Tertre, A., Lefranc, A., Vandentorren, S., Guldner, L., 2016. Biomarkers of exposure to environmental contaminants in French pregnant women from the Elfe cohort in 2011. *Environ. Int.* 97, 56–67. <https://doi.org/10.1016/j.envint.2016.10.013>.
- Diodato, S., González Garraza, G., Mansilla, R., Moretto, A., Escobar, J., Méndez-López, M., Gómez-Armesto, A., Marcovecchio, J., Nóvoa-Muñoz, J.C., 2020. Quality changes of fluvial sediments impacted by urban effluents in Ushuaia, Tierra del Fuego, southernmost Patagonia. *Environ. Earth Sci.* 79 (20) <https://doi.org/10.1007/s12665-020-09236-4>.
- Domingo, J.L., Nadal, M., 2019. Human exposure to per- and polyfluoroalkyl substances (PFAS) through drinking water: A review of the recent scientific literature. *Environ. Res.* 177, 108648. <https://doi.org/10.1016/j.envres.2019.108648>.
- EFSA, 2020. *Risk to Human Health Related to the Presence of Perfluoroalkyl Substances in Food (1831–4732)*.
- Ehresman, D.J., Froehlich, J.W., Olsen, G.W., Chang, S.-C., Butenhoff, J.L., 2007. Comparison of human whole blood, plasma, and serum matrices for the determination of perfluorooctanesulfonate (PFOS), perfluorooctanoate (PFOA), and other fluorochemicals. *Environ. Res.* 103 (2), 176–184. <https://doi.org/10.1016/j.envres.2006.06.008>.
- Eschauzier, C., Hoppe, M., Schlummer, M., de Voogt, P., 2013. Presence and sources of anthropogenic perfluoroalkyl acids in high-consumption tap-water based beverages. *Chemosphere* 90 (1), 36–41. <https://doi.org/10.1016/j.chemosphere.2012.06.070>.
- Espindola Santos, A.d.S., Meyer, A., Dabkiewicz, V.E., Câmara, V.d.M., Asmus, C.I.R.F., 2021. Serum levels of perfluorooctanoic acid and perfluorooctane sulfonic acid in pregnant women: maternal predictors and associations with birth outcomes in the PIPA project. *J. Obstet. Gynaecol. Res.* 47 (9), 3107–3118. <https://doi.org/10.1111/jog.14883>.
- Fan, X., Tang, S., Wang, Y., Fan, W., Ben, Y., Naidu, R., Dong, Z., 2022. Global exposure to per- and Polyfluoroalkyl substances and associated burden of low birthweight. *Environ. Sci. Technol.* 56 (7), 4282–4294. <https://doi.org/10.1021/acs.est.1c08669>.
- Fenton, S.E., Ducatman, A., Boobis, A., DeWitt, J.C., Lau, C., Ng, C., Smith, J.S., Roberts, S.M., 2021. Per- and Polyfluoroalkyl substance toxicity and human health review: current state of knowledge and strategies for informing future research. *Environ. Toxicol. Chem.* 40 (3), 606–630. <https://doi.org/10.1002/etc.4890>.
- Ferreira, M.F., Lo Nostro, F.L., Fernández, D.A., Genovese, G., 2021. Endocrine disruption in the sub Antarctic fish Patagonotothen tessellata (Perciformes, Nototheniidae) from Beagle Channel associated to anthropogenic impact. *Mar. Environ. Res.* 171, 105478. <https://doi.org/10.1016/j.marenvres.2021.105478>.
- Fiedler, H., Sadiq, M., 2021. Regional occurrence of perfluoroalkane substances in human milk for the global monitoring plan under the Stockholm Convention on persistent organic pollutants during 2016–2019. *Chemosphere* 277, 130287. <https://doi.org/10.1016/j.chemosphere.2021.130287>.
- Field, A., 2009. *Discovering Statistics Using IBM SPSS Statistics: And Sex and Drugs and Rock 'n' Roll*, 3th ed. SAGE.
- Fisher, M., Arbuckle, T.E., Liang, C.L., LeBlanc, A., Gaudreau, E., Foster, W.G., Haines, D., Davis, K., Fraser, W.D., 2016. Concentrations of persistent organic pollutants in maternal and cord blood from the maternal-infant research on environmental chemicals (MIREC) cohort study. *Environ. Health* 15 (1), 59. <https://doi.org/10.1186/s12940-016-0143-y>.
- Guida, Y., Torres, F.B.M., Barizon, R.R.M., Assalin, M.R., Rosa, M.A., 2023. Confirming sulfluramid (EtFOSA) application as a precursor of perfluorooctanesulfonic acid (PFOS) in Brazilian agricultural soils. *Chemosphere* 325, 138370. <https://doi.org/10.1016/j.chemosphere.2023.138370>.
- Haji Gholizadeh, M., Melesse, A.M., Reddi, L., 2016. Water quality assessment and apportionment of pollution sources using APCS-MLR and PMF receptor modeling techniques in three major rivers of South Florida. *Sci. Total Environ.* 566–567, 1552–1567. <https://doi.org/10.1016/j.scitotenv.2016.06.046>.
- Halldorsson, T.I., Fei, C., Olsen, J., Lipworth, L., McLaughlin, J.K., Olsen, S.F., 2008. Dietary predictors of Perfluorinated chemicals: A study from the Danish National Birth Cohort. *Environ. Sci. Technol.* 42 (23), 8971–8977. <https://doi.org/10.1021/es801907r>.
- Hornung, R.W., Reed, L.D., 1990. Estimation of average concentration in the presence of nondetectable values. *Appl. Occup. Environ. Hygiene* 5 (1), 46–51. <https://doi.org/10.1080/1047322X.1990.10389587>.
- Hu, X.C., Dassuncao, C., Zhang, X., Grandjean, P., Weihe, P., Webster, G.M., Nielsen, F., Sunderland, E.M., 2018. Can profiles of poly- and Perfluoroalkyl substances (PFASs) in human serum provide information on major exposure sources? *Environ. Health* 17 (1), 11. <https://doi.org/10.1186/s12940-018-0355-4>.
- Huber, S., Brox, J., 2015. An automated high-throughput SPE micro-elution method for perfluoroalkyl substances in human serum. *Anal. Bioanal. Chem.* 407 (13), 3751–3761. <https://doi.org/10.1007/s00216-015-8601-x>.
- INSPQ, 2023. *AMAP: AMAP Ring Test for Persistent Organic Pollutants in Human Serum*. Institut national de santé publique Québec. Retrieved 19.12.23. <https://www.inspq.qc.ca/en/ctq/eqas/amap/description>.
- Jian, J.-M., Guo, Y., Zeng, L., Liang-Ying, L., Lu, X., Wang, F., Zeng, E.Y., 2017. Global distribution of perfluorochemicals (PFCs) in potential human exposure source—A review. *Environ. Int.* 108, 51–62. <https://doi.org/10.1016/j.envint.2017.07.024>.
- Jiang, W., Zhang, Y., Zhu, L., Deng, J., 2014. Serum levels of perfluoroalkyl acids (PFAAs) with isomer analysis and their associations with medical parameters in Chinese pregnant women. *Environ. Int.* 64, 40–47. <https://doi.org/10.1016/j.envint.2013.12.001>.
- Kaboré, H.A., Vo Duy, S., Munoz, G., Méité, L., Desrosiers, M., Liu, J., Sory, T.K., Sauvè, S., 2018. Worldwide drinking water occurrence and levels of newly-identified perfluoroalkyl and polyfluoroalkyl substances. *Sci. Total Environ.* 616–617, 1089–1100. <https://doi.org/10.1016/j.scitotenv.2017.10.210>.
- Kannan, K., Corsolini, S., Falandysz, J., Fillmann, G., Kumar, K.S., Loganathan, B.G., Mohd, M.A., Olivero, J., Wouwe, N.V., Yang, J.H., Aldous, K.M., 2004. Perfluorooctanesulfonate and related Fluorochemicals in human blood from several countries. *Environ. Sci. Technol.* 38 (17), 4489–4495. <https://doi.org/10.1021/es0493446>.
- Kärman, A., Harada, K.H., Inoue, K., Takasuga, T., Ohi, E., Koizumi, A., 2009. Relationship between dietary exposure and serum perfluorochemical (PFC) levels—A case study. *Environ. Int.* 35 (4), 712–717. <https://doi.org/10.1016/j.envint.2009.01.010>.
- Kuo, K.-Y., Yu, C., Chuang, Y., Lin, P., Lin, Y.-J., 2023. Worldwide serum concentration-based probabilistic mixture risk assessment of perfluoroalkyl substances among pregnant women, infants, and children. *Ecotoxicol. Environ. Saf.* 268 <https://doi.org/10.1016/j.ecoenv.2023.115712>.
- Kurwadkar, S., Dane, J., Kanel, S.R., Nadagouda, M.N., Cawdrey, R.W., Ambade, B., Struckhoff, G.C., Wilkin, R., 2022. Per- and polyfluoroalkyl substances in water and wastewater: A critical review of their global occurrence and distribution. *Sci. Total Environ.* 809, 151003. <https://doi.org/10.1016/j.scitotenv.2021.151003>.
- Lee, H., Mabury, S.A., 2011. A pilot survey of legacy and current commercial fluorinated Chemicals in Human Sera from United States donors in 2009. *Environ. Sci. Technol.* 45 (19), 8067–8074. <https://doi.org/10.1021/es200167q>.
- Lee, Y.J., Kim, M.-K., Bae, J., Yang, J.-H., 2013. Concentrations of perfluoroalkyl compounds in maternal and umbilical cord sera and birth outcomes in Korea. *Chemosphere* 90 (5), 1603–1609. <https://doi.org/10.1016/j.chemosphere.2012.08.035>.
- Liu, Y., Li, A., Buchanan, S., Liu, W., 2020. Exposure characteristics for congeners, isomers, and enantiomers of perfluoroalkyl substances in mothers and infants. *Environ. Int.* 144, 106012. <https://doi.org/10.1016/j.envint.2020.106012>.
- Llorca, M., Farré, M., Tavano, M.S., Alonso, B., Korembli, G., Barceló, D., 2012. Fate of a broad spectrum of perfluorinated compounds in soils and biota from Tierra del Fuego and Antarctica. *Environ. Pollut.* 163, 158–166. <https://doi.org/10.1016/j.envpol.2011.10.027>.
- Löfstedt Gilljam, J., Leonel, J., Cousins, I.T., Benskin, J.P., 2016. Is ongoing Sulfluramid use in South America a significant source of Perfluorooctanesulfonate (PFOS)? Production inventories, environmental fate, and local occurrence. *Environ. Sci. Technol.* 50 (2), 653–659. <https://doi.org/10.1021/acs.est.5b04544>.
- Long, M., Knudsen, A.-K.S., Pedersen, H.S., Bonefeld-Jørgensen, E.C., 2015. Food intake and serum persistent organic pollutants in the Greenlandic pregnant women: the

- ACCEPT sub-study. *Sci. Total Environ.* 529, 198–212. <https://doi.org/10.1016/j.scitotenv.2015.05.022>.
- Luz, A.L., Anderson, J.K., Goodrum, P., Durda, J., 2019. Perfluorohexanoic acid toxicity, part I: development of a chronic human health toxicity value for use in risk assessment. *Regul. Toxicol. Pharmacol.* 103, 41–55. <https://doi.org/10.1016/j.yrtph.2019.01.019>.
- Mansournia, M.A., Nazemipour, M., Naimi, A.I., Collins, G.S., Campbell, M.J., 2020. Reflection on modern methods: demystifying robust standard errors for epidemiologists. *Int. J. Epidemiol.* 50 (1), 346–351. <https://doi.org/10.1093/ije/dyaa260>.
- McAdam, J., Bell, E.M., 2023. Determinants of maternal and neonatal PFAS concentrations: a review. *Environ. Health* 22 (1), 41. <https://doi.org/10.1186/s12940-023-00992-x>.
- Müller, M.H.B., Polder, A., Brynildsrud, O.B., Grønnestad, R., Karimi, M., Lie, E., Manyilizu, W.B., Mdegela, R.H., Mokiti, F., Murtadha, M., Nonga, H.E., Skaare, J.U., Solhaug, A., Lyche, J.L., 2019. Prenatal exposure to persistent organic pollutants in northern Tanzania and their distribution between breast milk, maternal blood, placenta and cord blood. *Environ. Res.* 170, 433–442. <https://doi.org/10.1016/j.envres.2018.12.026>.
- Okada, E., Kashino, I., Matsuura, H., Sasaki, S., Miyashita, C., Yamamoto, J., Ikeno, T., Ito, Y.M., Matsumura, T., Tamakoshi, A., Kishi, R., 2013. Temporal trends of perfluoroalkyl acids in plasma samples of pregnant women in Hokkaido, Japan, 2003–2011. *Environ. Int.* 60, 89–96. <https://doi.org/10.1016/j.envint.2013.07.013>.
- Oklad, I., Odland, J.O., Matiocevic, S., Alvarez, M.V., Aarsland, T., Nieboer, E., Hansen, S., 2017. The Argentinian mother-and-child contaminant study: a cross-sectional study among delivering women in the cities of Ushuaia and Salta. *Int. J. Circumpolar Health* 76 (1), 1364598. <https://doi.org/10.1080/22423982.2017.1364598>.
- Olsen, G.W., Butenhoff, J.L., Zobel, L.R., 2009. Perfluoroalkyl chemicals and human fetal development: an epidemiologic review with clinical and toxicological perspectives. *Reprod. Toxicol.* 27 (3–4), 212–230. <https://doi.org/10.1016/j.reprotox.2009.02.001>.
- Pérez, F., Llorca, M., Köck-Schulmeyer, M., Škrbić, B., Oliveira, L.S., da Boit Martinello, K., Al-Dhabi, N.A., Antić, I., Farré, M., Barceló, D., 2014. Assessment of perfluoroalkyl substances in food items at global scale. *Environ. Res.* 135, 181–189. <https://doi.org/10.1016/j.envres.2014.08.004>.
- Poothong, S., Papadopoulou, E., Padilla-Sánchez, J.A., Thomsen, C., Haug, L.S., 2020. Multiple pathways of human exposure to poly- and perfluoroalkyl substances (PFASs): from external exposure to human blood. *Environ. Int.* 134, 105244. <https://doi.org/10.1016/j.envint.2019.105244>.
- Quinn, C. L., & Wania, F. (2012). Understanding Differences in the Body Burden–Age Relationships of Bioaccumulating Contaminants Based on Population Cross Sections versus Individuals. *Environ. Health Perspect.*, 120(4), 554–559. doi:<https://doi.org/10.1289/ehp.1104236>.
- Rickard, B.P., Rizvi, I., Fenton, S.E., 2022. Per- and poly-fluoroalkyl substances (PFAS) and female reproductive outcomes: PFAS elimination, endocrine-mediated effects, and disease. *Toxicology* 465, 153031. <https://doi.org/10.1016/j.tox.2021.153031>.
- Seghezzo, L., Gatto D'Andrea, M.L., Iribarnegaray, M.A., Liberal, V.I., Fleitas, A., Bonifacio, J.L., 2013. Improved risk assessment and risk reduction strategies in the water safety plan (WSP) of Salta, Argentina. *Water science & technology. Water supply* 13 (4), 1080–1089. <https://doi.org/10.2166/ws.2013.087>.
- Souza, M.C.O., Saraiva, M.C.P., Honda, M., Barbieri, M.A., Bettiol, H., Barbosa, F., Kannan, K., 2020. Exposure to per- and polyfluorinated alkyl substances in pregnant Brazilian women and its association with fetal growth. *Environ. Res.* 187, 109585. <https://doi.org/10.1016/j.envres.2020.109585>.
- Stockholm Convention, S., 2023. The new POPs under the Stockholm Convention. Retrieved 31.11.23 from. <https://chm.pops.int/TheConvention/ThePOPs/TheNewwPOPs/tabid/2511/Default.aspx>.
- Sunderland, E.M., Hu, X.C., Dassuncao, C., Tokranov, A.K., Wagner, C.C., Allen, J.G., 2019. A review of the pathways of human exposure to poly- and perfluoroalkyl substances (PFASs) and present understanding of health effects. *J. Expo. Sci. Environ. Epidemiol.* 29 (2), 131–147. <https://doi.org/10.1038/s41370-018-0094-1>.
- Tian, Y., Zhou, Y., Miao, M., Wang, Z., Yuan, W., Liu, X., Wang, X., Wang, Z., Wen, S., Liang, H., 2018. Determinants of plasma concentrations of perfluoroalkyl and polyfluoroalkyl substances in pregnant women from a birth cohort in Shanghai, China. *Environ. Int.* 119, 165–173. <https://doi.org/10.1016/j.envint.2018.06.015>.
- Toms, L.M.L., Thompson, J., Rotander, A., Hobson, P., Calafat, A.M., Kato, K., Ye, X., Broomhall, S., Harden, F., Mueller, J.F., 2014. Decline in perfluorooctane sulfonate and perfluorooctanoate serum concentrations in an Australian population from 2002 to 2011. *Environ. Int.* 71, 74–80. <https://doi.org/10.1016/j.envint.2014.05.019>.
- Umweltbundesamt. (2023, 15.09.23). Reference and HBM values. German Environment Agency. Retrieved 15.11.23 from <https://www.umweltbundesamt.de/en/topics/health/commissions-working-groups/human-biomonitoring-commission/reference-hbm-values>.
- van Asselt, E.D., Rietra, R.P.J.J., Römkens, P.F.A.M., van der Fels-Klerx, H.J., 2011. Perfluorooctane sulphonate (PFOS) throughout the food production chain. *Food Chem.* 128 (1), 1–6. <https://doi.org/10.1016/j.foodchem.2011.03.032>.
- Wallis, D.J., Barton, K.E., Knappe, D.R.U., Kotlarz, N., McDonough, C.A., Higgins, C.P., Hoppin, J.A., Adgate, J.L., 2023. Source apportionment of serum PFASs in two highly exposed communities. *Sci. Total Environ.* 855, 158842. <https://doi.org/10.1016/j.scitotenv.2022.158842>.
- Wang, T., Wang, P., Meng, J., Liu, S., Lu, Y., Kim, J.S., Giesy, J.P., 2015. A review of sources, multimedia distribution and health risks of perfluoroalkyl acids (PFAAs) in China. *Chemosphere* 129, 87–99. <https://doi.org/10.1016/j.chemosphere.2014.09.021>.
- Wang, Y., Han, W., Wang, C., Zhou, Y., Shi, R., Bonefeld-Jørgensen, E.C., Yao, Q., Yuan, T., Gao, Y., Zhang, J., Tian, Y., 2019. Efficiency of maternal-fetal transfer of perfluoroalkyl and polyfluoroalkyl substances. *Environ. Sci. Pollut. Res. Int.* 26 (3), 2691–2698. <https://doi.org/10.1007/s11356-018-3686-3>.
- Wee, S.Y., Aris, A.Z., 2023. Revisiting the “forever chemicals”, PFOA and PFOS exposure in drinking water. *npj Clean Water* 6 (1). <https://doi.org/10.1038/s41545-023-00274-6> (57–16).
- Workman, C.E., Becker, A.B., Azad, M.B., Moraes, T.J., Mandhane, P.J., Turvey, S.E., Subbarao, P., Brook, J.R., Sears, M.R., Wong, C.S., 2019. Associations between concentrations of perfluoroalkyl substances in human plasma and maternal, infant, and home characteristics in Winnipeg, Canada. *Environ. Pollut.* 249, 758–766. <https://doi.org/10.1016/j.envpol.2019.03.054>.
- Zell-Baran, L.M., Venter, C., Dabelea, D., Norris, J.M., Glueck, D.H., Adgate, J.L., Brown, J.M., Calafat, A.M., Pickett-Nairne, K., Starling, A.P., 2023. Prenatal exposure to poly- and perfluoroalkyl substances and the incidence of asthma in early childhood. *Environ. Res.* 239, 117311. <https://doi.org/10.1016/j.envres.2023.117311>.
- Zhang, L., Liu, J., Hu, J., Liu, C., Guo, W., Wang, Q., Wang, H., 2012. The inventory of sources, environmental releases and risk assessment for perfluorooctane sulfonate in China. *Environ. Pollut.* 165, 193–198. <https://doi.org/10.1016/j.envpol.2011.09.001>.
- Zheng, G., Eick, S.M., Salamova, A., 2023. Elevated levels of ultrashort- and short-chain Perfluoroalkyl acids in US homes and people. *Environ. Sci. Technol.* 57 (42), 15782–15793. <https://doi.org/10.1021/acs.est.2c06715>.