

Supplementary Information

Exploring the modulatory influence on the antimalarial activity of amodiaquine using scaffold hybridisation with ferrocene integration

Mziyanda Mbaba^a, Taryn M. Golding^a, Reinner O. Omondi^a, Roxanne Mohunlal^a, Timothy J. Egan^a, Janette Reader^b, Lyn-Marie Birkholtz^b and Gregory S. Smith^{a,*}

^a*Department of Chemistry, Faculty of Science, University of Cape Town, Rondebosch, 7701, South Africa*


^b*Department of Biochemistry, Genetics and Microbiology, Institute for Sustainable Malaria Control, University of Pretoria, Hatfield 0028, South Africa*

*Corresponding author: gregory.smith@uct.ac.za

Table of contents


Synthesis of aniline 2a	(2)
NMR spectral data of synthesized compounds	(3)
HPLC traces of final compounds	(27)
Representative graphs for β -hematin inhibition	(32)

Synthesis of aniline 2a




Methyl 3-nitrobenzoate (S02a). Two drops of sulfuric acid (100%) were added to a solution of 3-nitrobenzoic acid (2.00 g, 12.0 mmol) in MeOH (25 mL) and the resulting mixture refluxed for 24 h. After 24 h, the reaction mixture was diluted with distilled water (25 mL) and the product extracted with DCM (3 × 25 mL), dried (Na₂SO₄) and the solvent removed under reduced pressure to give an off-white solid. The crude product was purified by silica gel column chromatography (5% MeOH/DCM) to furnish the pure compound as a white solid. Yield: 1.78 g (82%). ¹H NMR (600 MHz, CDCl₃) δ 8.86 (t, *J* = 2.0 Hz, 1H, H₂), 8.41 (ddd, *J* = 8.3, 2.4, 1.2 Hz, 1H, H₄), 8.37 (dt, *J* = 7.8, 1.4 Hz, 1H, H₆), 7.65 (t, *J* = 8.0 Hz, 1H, H₅), 3.99 (s, 3H, OMe). ¹³C NMR (150 MHz, CDCl₃) δ 165.1, 148.5, 135.4, 132.1, 129.8, 127.5, 124.8, 52.9.


Methyl 3-aminobenzoate (2a). Saturated NH₄Cl solution (5 mL) was cautiously added to a suspension of S02a (1.75 g, 9.7 mmol) and Zn dust (3.81 g, 58.2 mmol), and the resulting mixture was refluxed for 1 h. The product was extracted with EtOAc (2 × 25 mL), dried over Na₂SO₄ and the solvent removed *in vacuo* to afford the pure product as a light-yellow oil. Yield: 1.30 g (88 %). ¹H NMR (600 MHz, CDCl₃) δ 7.43 (dt, *J* = 7.7, 1.3 Hz, 1H, H₆), 7.36 (t, *J* = 2.0 Hz, 1H, H₂), 7.21 (t, *J* = 7.8 Hz, 1H, H₅), 6.87 (ddd, *J* = 8.0, 2.5, 1.1 Hz, 1H, H₄), 3.89 (s, 3H, OMe). ¹³C NMR (150 MHz, CDCl₃) δ 167.4, 146.3, 131.3, 129.4, 120.1, 119.7, 116.1, 52.2.




¹³C NMR spectrum of **S02a**



¹H NMR spectrum of **2a**




¹³C NMR spectrum of **2a**




¹H NMR spectrum of **S5b**

WS_FPP_1231.fid
MZM-52b in DMSO
1H Spectrum
Mziyanda




¹H NMR spectrum of **5a**


600.1233.fid
MZM-053a in DMSO
1H Spectrum
Mziyanda



¹H NMR spectrum of **5b**




¹³C NMR spectrum of **5b**




¹H NMR spectrum of 6a

WS_FTP551.fid
 MZM-54a in DMSO
 1H Spectrum
 Mziyanda




¹³C NMR spectrum of 6a


WS_FTP.552.fid
 MZM-54a in DMSO
 13C Spectrum
 Mziyanda




¹H NMR spectrum of **6b**




¹³C NMR spectrum of **6b**




^1H NMR spectrum of **6c**




^{13}C NMR spectrum of **6c**




¹H NMR spectrum of **6d**




¹³C NMR spectrum of **6d**




¹H NMR spectrum of **7a**




¹³C NMR spectrum of **7a**




¹H NMR spectrum of **7b**




¹³C NMR spectrum of **7b**




¹H NMR spectrum of **7c**




¹³C NMR spectrum of **7c**




¹H NMR spectrum of **7d**




¹³C NMR spectrum of **7d**




¹H NMR spectrum of **8**




¹H NMR spectrum of **9a**




¹³C NMR spectrum of **9a**




¹H NMR spectrum of **9b**




¹³C NMR spectrum of **9b**




¹H NMR spectrum of **9c**




¹³C NMR spectrum of **9c**




Representative COSY NMR spectrum of **6b**




Representative HSQC NMR spectrum of **6b**




Representative COSY NMR spectrum of **7a**




Representative HSQC spectrum of **7a**




Representative HMBC spectrum of **7a**




Representative COSY NMR spectrum of **7a**




Representative HSQC spectrum of **7a**




Representative HMBC spectrum of **7a**




Representative HMBC spectrum of **9b**




Representative COSY NMR spectrum of **9c**




Representative HSQC spectrum of **9c**




Representative HMBC spectrum of **9c**



HPLC trace of 6a




HPLC trace of 6b



Injection Results

PK#	Summary	RT (min)	Area	Area%	Height	Concentration	S/N	Tailing
1	VWD1A,Wavelength=300 nm	3.683	10400.824	99.167	1676.551			
2	VWD1A,Wavelength=300 nm	4.074	87.410	0.833	16.964			


HPLC trace of 6c




Injection Results

PK#	Summary	RT (min)	Area	Area%	Height	Concentration	S/N	Tailing
1	VWD1A,Wavelength=300 nm	3.881	18819.561	99.606	2477.335			
2	VWD1A,Wavelength=300 nm	4.076	66.570	0.394	15.964			


HPLC trace of 7a




HPLC trace of 7b




HPLC trace of **7c**




HPLC trace of **7d**




HPLC trace of 9a







HPLC trace of 9b







HPLC trace of 9c



Representative graphs for β -hematin inhibition



Representative graphs for β -hematin inhibition



Representative graphs for β -hematin inhibition

