



Check for updates

AUTHORS:

Teresa A. Coutinho^{1,2,3}
Jacquie E. van der Waals^{2,3,4,5}

AFFILIATIONS:

¹Department of Biochemistry, Genetics and Microbiology, University of Pretoria, Pretoria, South Africa

²Centre for Microbial Ecology and Genomics, University of Pretoria, Pretoria, South Africa

³Forestry and Agricultural Biotechnology Institute, University of Pretoria, Pretoria, South Africa

⁴Department of Plant and Soil Sciences, University of Pretoria, Pretoria, South Africa

⁵Citrus Research International, Mbombela, South Africa

CORRESPONDENCE TO:

Teresa Coutinho

EMAIL:

teresa.coutinho@up.ac.za

HOW TO CITE:

Coutinho TA, van der Waals JE. Historical perspective on 120 years of agriculture: Highlights from research published in the SAJS. S Afr J Sci. 2024;120(Special issue: Celebrating 120 years), Art. #18965. <https://doi.org/10.17159/sajs.2024/18965>

ARTICLE INCLUDES:

- Peer review
- Supplementary material

KEYWORDS:

agriculture, South Africa, climate change, water scarcity, soil quality

PUBLISHED:

07 November 2024



Historical perspective on 120 years of agriculture: Highlights from research published in the SAJS

Significance:

Agriculture, an essential pillar of South Africa’s economy, has evolved through a century of scientific advancements. This Perspective highlights the significant research contributions to the improvement of citrus and maize production, addressing historical challenges and emphasising ongoing issues such as climate change, water scarcity and soil quality. The introduction of modern agricultural practices, including hybrid seeds, genetic modification, and advanced irrigation techniques, has transformed these crops, enabling increased productivity and global competitiveness. However, the sector faces persistent challenges, including the impacts of climate change and food security concerns, particularly in rural communities. We also explore future prospects, underscoring the importance of technology adoption and sustainable practices to enhance food security. South Africa’s agricultural research, although underfunded compared to developed nations, remains crucial for sustaining the agricultural landscape and ensuring the sector’s resilience and growth in an increasingly challenging global environment.

Introduction

The word ‘agriculture’ describes the multitude of ways that crops and livestock provide food and other products for the global human population. It is a practice of agroecosystem manipulation. Agricultural science is a multidisciplinary field that focuses on understanding and enhancing the productivity, sustainability, and profitability of agriculture. In South Africa, agriculture not only includes the commercial sector but also smallholder and subsistence farmers, who have sustained rural communities and shaped the agricultural landscape for centuries. Agriculture has been and still is a pillar of South Africa’s economy, contributing 2.83% to the Gross Domestic Product¹ and providing 19.26% of total employment². Agricultural land covers approximately 80% of the country’s total area of 1.22 million km² of which only 12% is regarded as arable.³ Agriculture has faced many challenges including, for example, the impact of climate change, pests and diseases, and the complexities of land reform and redistribution issues since 1994. Nonetheless, South African agriculture holds promise for future growth and development, both locally and internationally, as there is an increasing global demand for food and its sustainable production.

In this Perspective, we present some highlights of the agricultural research that has been conducted over the past 120 years (1905–2023) – the lifespan of the *South African Journal of Science (SAJS)* from which most citations were taken. However, when subject-specific journals began to appear, for example, the *Agricultural Journal of South Africa* in 1915, *Farming in South Africa* in 1927 and the *South African Journal of Agricultural Science* in 1957, authors preferred to publish in these journals, and fewer articles published on agricultural-related topics were published in the SAJS. For this reason, we included references from other journals. However, agriculture is such a broad and multidisciplinary topic that the focus of this review will be limited to research conducted on the production of citrus and maize and the challenges faced by these two sectors of the agricultural economy.

Citrus

Citrus trees, both oranges and lemons, were brought to the Cape from St Helena Island in 1654 and planted in the Cape Gardens.⁴ By 1661, 1000 trees had been established, and, for the next 200 years, citrus was grown by small-scale farmers in the Western Cape. At the beginning of the 19th century, early settlers in the Eastern Cape and KwaZulu-Natal Provinces planted individual orange and soft citrus trees and, shortly after the Great Trek (1838), citrus trees were planted in the North West Province.

As the era of the SAJS began at the end of the 19th century, the then Prime Minister of the Cape Colony, Cecil John Rhodes, hired horticulturists, HEV Pickstone and RE Davies from California, to develop the citrus industry in the Cape. Thereafter, citrus trees were grown on a commercial scale using the modern scientific methods known at that time. High-quality citrus fruit was exported for the first time to the United Kingdom in 1906 and “made their debut at the Show of the Royal Horticultural Society in London”⁴. The industry flourished thereafter, and between 1906 and 1921, a number of co-operative grower associations formed, of which the most important were the Fruit Growers’ Co-operative Exchange (1922), the South African Co-operative Citrus Exchange (SACCE) (1925) and the Citrus Board (1939).⁵ Prior to 1941, the distribution and sale of citrus fruit were undertaken by individual producers, but in the year that the South African Citrus Scheme was introduced, sales increased as the fruit was sold in a “more profitable manner”⁵. By 1928, more than three million citrus trees were planted in the Cape. Today, the area occupied by citrus trees of all species and varieties is over 100 000ha⁶, equating to approximately 55.5 million trees.

According to Fourie et al.⁷, the first research officer posts were created in 1953 by the SACCE to undertake work on identified problem areas not adequately investigated by government and other research organisations. Between the 1960s and 1970s, three scientists, trained in California, were appointed by the SACCE in the fields of pre- and post-harvest pathology and horticulture. Their research was undertaken at the Citrus and Subtropical Fruit Institute in Nelspruit, which later became part of the Agricultural Research Council. The Citrus Improvement Programme was established in 1973 and later became known as the South African Citrus Improvement Scheme, managed by Citrus Research International. Scientists from this organisation conduct research on a range of topics relevant to their industry, including pathology, entomology and citriculture.

© 2024. The Author(s). Published under a Creative Commons Attribution Licence.

Maize

The introduction of maize into southern Africa is a matter of debate among archaeologists. Burt-Davy⁸ suggested that maize seed was exchanged for water, meat or other commodities by the Portuguese *en route* to the East in the mid-1600s. In the latter part of the 19th century, maize was mainly cultivated for local subsistence needs.⁹ The market changed with the discovery of diamonds (1866) and gold (1886), and black and white farmers alike supplied maize to the rapidly expanding mining population.¹⁰ “Two agricultures” then emerged: the black smallholder farmers and the white commercial farmers, the production potential of the former was suppressed while the opposite was true for the latter farmers who were supported by substantial investment.¹¹ Despite this initial division between the maize producers, which changed in the 1990s, South Africa increased its maize production from 328 000 tons in 1904 to 1.68 million tons in 1935¹⁰ and 15.6 million tons in the 2022/2023 season¹². This significant increase can be attributed to factors such as the use of hybrid and genetically modified seed, and improved fertiliser programmes and farming practices. Our understanding of plant genetics and the use of biotechnological tools such as marker-assisted selection and genetic engineering, for example, have allowed the industry to produce superior quality seed and plants resistant to pest infestations.

Maize is the most important crop in South Africa and is used both as animal feed and as a staple human food crop. Today the area planted to maize ranges from 2.5 to 2.8 million hectares annually.¹³ The cultivation of maize is restricted by climate. It is considered a dry-land crop, and is mainly grown in the summer rainfall areas such as the North West and Limpopo Provinces.⁸ In 1923, the “maize triangle” as it became known with Mafikeng (North West), eNtokozweni (Machadodorp) (Mpumalanga) and Zastron (Free State) forming the apices, produced approximately 60% of the total amount (1.5 million tons) grown in the country. The remaining 40% produced was produced by other provinces (KwaZulu-Natal and the Cape) and subsistence farmers.¹⁴ Today, approximately 14 million tons are produced annually, mainly by commercial farmers, but small-scale farmers also produce a significant proportion of this amount.¹⁵ Approximately 3.2 million tons are exported, mainly to Zimbabwe, South Korea, Japan and Taiwan.¹² The major producers are in the Free State, Mpumalanga and North West Provinces.¹

Yellow and white maize, genetically modified with the Bt toxin, were commercially grown for the first time in the 1998/1999 and 2001/2002 seasons, respectively. South Africa then became the first genetically modified (GM) subsistence crop producer in the world.¹⁶ Food security benefits attributable to GM white maize in South Africa are substantial and “an average of 4.6 million additional white maize rations annually” have been suggested.¹⁷ Thus scientific research has played a pivotal role in the development and widespread adoption of using GM maize in the country.

Challenges

Water availability

Water is scarce in South Africa and severe droughts have impacted agriculture. Water scarcity has been exacerbated in recent years by erratic rainfall patterns¹⁸, the needs of industries and cities, and impending climate change. There have been several drought investigations in the past, with the first published in 1914¹⁹ and the most recent report published in 2020, namely, the National Water Security Framework for South Africa²⁰. Over the years, drought relief plans and management practices were proposed, which included better veld, stock and water management, and improved farm management.

Crop irrigation using flood and furrow methods, as opposed to dry land farming, was first proposed in 1905.^{21,22} Sources of water for irrigation purposes have been and still are from various aquifers²³, catchment areas and rivers²⁴. The benefits of irrigation were highlighted in the SAJS by van Reenen²⁵ and Chunnnett²⁶. Other methods of water delivery and applications such as drip irrigation, developed in the 1900s in Israel, or the use of microsprinklers²⁷, were shown to be more water efficient than flood and furrow methods. These more efficient methods are the generally preferred irrigation systems in citrus orchards, as water is directly placed in the root zone and can also be used for fertigation.²⁸

In maize, supplementing rainfall during the mid-season drought period positively impacted yield²⁹ and this method has been shown to improve and stabilise smallholder farmer maize yields in particular³⁰. Centre pivot-irrigation was first introduced into South Africa in the 1970s to irrigate the farm Soetvelde near Vereeniging.³¹ This method enabled commercial farmers to increase quantities of water applied, and is used in some areas for maize production.³²

Soil quality

As early as 1923, soil erosion, by wind and/or rain, was recognised as a significant problem in South Africa.³³ Policies were developed to prevent and mitigate erosion in the agricultural sector.³⁴ In 1939, the Soil Erosion Advisory Council was formed, and the *Soil Conservation Act* was published in 1946. At that time, all these efforts led to the effective control of erosion in some areas of the country. Today, however, over 70% of South Africa is experiencing some form of soil erosion, with the extent of the problem varying by region.³⁵

Soil quality is multifaceted. Besides soil erosion, South African soils have also been subjected to structural decay, subsoil compaction, acidification, salinisation, pollution, nutrient depletion and surface crusting, resulting in reduced water infiltration, increased run-off and erosion.^{36,37} A number of methods can improve and maintain soil quality, including the use of fertilisers. The low quantities of phosphorus in our soils were first recognised in the early 1900s and found to be the main limiting factor in crop production.³⁸ This led to the development of the first phosphate plant in Durban in 1903.³⁹ Fertilisers were also imported, with 19 000 tons of fertilisers imported in 1906.⁴⁰ By 1962, this amount had grown to 1 385 000 tons⁴¹, and, in 2023, 2 174 911 tons were imported.⁴² South African soils also have low organic matter levels, with about 58% of soils containing less than 0.5% organic carbon and only 4% containing more than 2% organic carbon.⁴³

Pests and pathogens

The fields of entomology and plant pathology have a long history in South Africa that dates to the 17th century.^{44,45} Pests and pathogens were and still are responsible for significant losses in agriculture, not only during production but also during storage. Initially, the most common method of management was pesticide application. Locusts, for example, were controlled with arsenate of soda⁴⁶, while diseases of grapes were controlled using sulfur with an amount of 264 tons used in 1916⁴⁷. After 1919, inorganic pesticides were used, including Bordeaux mixture used to prevent numerous fungal diseases. After 1945, numerous synthetic pesticides were developed, many of which are still used today. However, many were detrimental to the environment and to humans. In the 1980s, sustainable farming was encouraged in an attempt to revise the harmful practices of applying increasing amounts of chemicals.⁴⁸ Today, management has moved from a reliance on pesticides to environmentally friendly options such as the use of biological control agents.⁴⁹

Crop losses due to pests and pathogens are becoming increasingly more common. As pests and pathogens have the capacity to generate new variants containing key pathogenicity, fitness and aggressiveness traits, any new opportunity provided by climate change will be exploited by them.⁵⁰ Fluctuations in rainfall, humidity, temperature and carbon dioxide levels may also result in the emergence of pests in the field, an increase in their reproductive rates, shifts in their life cycle and movement from one area to another.⁵¹

Citrus

Doidge^{52,53} described numerous citrus diseases caused by bacteria and fungi occurring in South Africa between 1919 and 1929. Bacterial, fungal and viral pathogens were and still are problematic pre- and post-harvest. Citrus canker, first detected in the country in 1905/1906, was successfully eradicated by 1927, and was the first plant disease to be successfully eradicated in the world.⁴⁴ Today, one of the most important citrus diseases is black spot (CBD), discovered in 1922 by Doidge.⁷ The disease is not present in the Western Cape, Northern Cape and the Free State⁵⁴ and its presence on the fruit from KwaZulu-Natal, North West, Mpumalanga and Limpopo has impacted on the export market. The EU has strict regulations (zero tolerance) related to the import of

fruit from South Africa, both with regard to CBD and the false codling moth, as neither occurs in Europe. To improve risk assessment and CBD management, spore release prediction models have been developed.⁵⁵ The presence of viruses has also hampered citrus production.^{7,56} The earliest recorded evidence of a virus causing significant damage was the citrus tristeza virus in 1896, which led to the abandonment of sour orange as a rootstock in South Africa.⁷ The identification of the causal agent was, however, only confirmed in 1947.⁴

Entomological problems of citrus described in 1934 included red scale, citrus aphid, fruit fly, mealy bugs and thrips.⁵⁷ Today, many of these insects are still problematic. An additional problem with some pests is their ability to vector plant pathogens, for example, the citrus psyllid, which vectors the citrus greening pathogen.⁵⁸ Although insecticides have been used for many years, an integrated pest management approach is more widely accepted today.

Maize

Maize has been plagued by many pests and pathogens over the past century. In 1911, cutworm, stalk borer, the striped earworm and a leaf disease of unknown cause were problematic.⁸ The number of new incursions of pests and pathogens steadily increased, with the fall armyworm making an appearance in 2017.⁵⁹ Pathogens such as those causing cob and stalk rot and storage rot produce mycotoxins and threaten the health of humans and domestic stock.⁶⁰ Leaf blight diseases include Northern and Southern leaf blights, grey leaf spot, common rust and maize streak virus, and may cause substantial losses each year.^{13,61}

The development of Bt maize, engineered to contain toxins from *Bacillus thuringiensis*, has provided resistance in South African maize to important insect pests such as the European corn borer and other lepidopteran species.⁶² In South Africa, Bt maize makes up more than 90% of the maize planted and has resulted in a significant reduction in pesticides usage, crop damage and attack by fungal pathogens.¹⁶

Climate change

A number of researchers have modelled climate change and the effects thereof on agriculture in southern Africa.^{18,63–66} According to the IPCC 6th Assessment Working Group II report, climate change has increased the mean annual surface temperature of the earth by 1.2 °C, relative to 1850–1900, with the six hottest years ever recorded having occurred in the last decade. According to Blignaut et al.⁶⁷, South Africa became 2% hotter and at least 6% drier in the period 1997 to 2006 compared to the 1970s. During the same period, the use of water also increased significantly, with irrigation agriculture using 60% of total consumption. An increase in temperature and a decrease in rainfall are not the only consequences of climate change; there are increases in extreme weather events, heatwave intensity and frequency, severe droughts and flooding after torrential rains. The average rainfall across South Africa has also decreased in all areas except the northwest and, despite an overall trend in aridity, the intensity and frequency of heavy rainfall events have also increased. In addition to these abiotic factors, biotic stresses, such as an increase in pest and pathogen populations, and increased weed growth have had negative impacts on agriculture.

Climate change will impact both citrus and maize production, leading to possible decreases in yield.^{68,69} In citrus, the effect will be physiological, resulting in a loss of productivity and fruit quality. Adaptation strategies such as canopy management, top netting, soil fertility management and water-saving approaches could be used.⁷⁰ Maize production by both commercial and smallholder farming systems is predicted to decrease by between 10% and 16% because of the projected climatic impacts.⁶⁹ One method of overcoming this challenge is to create new varieties, and in the case of citrus, new rootstocks, that can adapt to the predicted environmental conditions.

Johnston et al.⁷¹ – in a report entitled ‘Climate change impacts in South Africa: what climate change means for a country and its people’ – recorded that commercial, small-scale, and subsistence farmers have implemented various climate change adaptation strategies. Conservation agriculture, which focuses on retaining soil carbon to enhance drought resilience, has become more important in dryland farming regions. Notably, improvements in irrigation efficiency, coupled with practical drought tolerant crop and

cultivar selection, have become standard practices. In addition, there has been a noticeable decrease in reliance on chemical inputs. Access to medium- and short-term weather forecasts has been enhanced, empowering farmers to proactively prepare for extreme weather events. This includes leveraging indigenous and local knowledge of weather and climate patterns, as well as implementing water resource management techniques.⁷² For instance, some farmers now harvest rainwater to sustain agricultural activities during dry periods, thereby prolonging the growing season. Due to these adaptations, the sector’s economic sustainability has largely remained intact. Nevertheless, the escalating temperatures and the intensification of extreme weather phenomena pose difficult challenges to farmers’ adaptive capacity. As such, ongoing efforts to innovate and implement resilient agricultural practices by both smallholder and commercial farmers will be essential in navigating the evolving climate landscape.⁷¹

Food security

The term ‘food security’ was coined in the 1970s.⁷³ Food supplies are deemed secure when every member of the population has access to sufficient food for living an active, healthy life.⁷⁴ There are four elements to the term: food availability, food access, food utilisation and stability.^{51,75}

Despite the constitutional right of every South African citizen to sufficient food and water, food provision is complex and determined by various environmental, health, economic, socio-political, and agro-food-related issues. Over 20% of South Africans are food insecure, that is one in five members of the population, mostly from the lowest socio-economic groups⁷⁶, with about 55% of the population unable to pay for both food and other basic needs.⁷⁷ Widespread poverty, unemployment and inequality have exacerbated food insecurity. This has led to health issues such as obesity and hypertension.⁷⁸ One in four children have stunted growth because of poor nutrition.⁷⁹ There is an urgent need for significant investment in sustainable agriculture, the development of infrastructure, and poverty alleviation strategies to improve food security for all South Africans.

Perspectives

Despite all the challenges, agriculture remains a cornerstone of South Africa’s economy, contributing 2.83% to the GDP. Over the past 120 years – the life of the SAJS – food production has increased due to genetic crop improvements, including the contentious use of GMOs, and cropping practices, notably irrigation and fertilisers.⁸⁰ Key issues that are likely to shape the agricultural future include the adoption of technology; adaption to climate change; sustainable production strategies; production of less studied, locally adapted plant species used in rural communities as food; and education. Technology, such as the use of genetic methodologies like genome editing, artificial intelligence, precision agriculture, drones and satellite imaging for monitoring, and apps for decision-making, have the potential to increase both efficiency and productivity in agriculture. In the case of maize, there is a need to breed new varieties that are drought tolerant, and in citrus, rootstocks and scions that maintain production and fruit quality in the face of climate change. Integrated pest management strategies, including the sensible use of pesticides and the adoption of biological control strategies, will be crucial for managing pests and pathogens while at the same time minimising their environmental impact. Under-utilised agricultural crops, or orphan crops, such as moringa⁸¹ and *Plectranthus aeculentus*⁸², can also contribute to food security. As agriculture will become more technology-intensive, at least in the commercial sector, skilled workers will need to be educated in fields such as data analysis and agricultural engineering. However, low capacity for adaptation, widespread poverty and low technology uptake might hamper agricultural productivity in South Africa.

In conclusion, South Africa’s research focus areas are similar to those in the developed world, especially in terms of technology and infrastructure. Research conducted ranges from cutting-edge molecular studies to system modelling approaches. This is despite the fact that expenditure on research, estimated at 1.04% of the agriculture GDP, is significantly less than that of Europe (2.06%), the USA (3.7%) and Australia (4.02%).⁸³ It is, however, comparable to that of other developing countries. Since 2008⁸⁴, the government has recognised that South Africa needs to progress to a “developed state of agriculture and agroprocessing”, especially in terms of technological innovation, so that



wealth generation and socio-economic development within this sector can be achieved within the next 20 years.⁸³

Acknowledgements

We thank Professor Jane Carruthers for reviewing the first draft of this manuscript.

Declarations

T.A.C. is the SAJS Associate Editor for Agriculture and Forestry. There are no competing interests to declare. There is no AI or LLM use to declare. Both authors read and approved the final manuscript.

References

- Statista. Value added by agriculture to the Gross Domestic Product (GDP) in South Africa from 2016 to 2023 [webpage on the Internet]. c2023 [cited 2024 Jun 05]. Available from: <https://www.statista.com/statistics/1121213/agriculture-value-added-to-gdp-in-south-africa/>
- Trading Economics. South Africa – Employment in agriculture (% of total employment) [website on the Internet]. c2024 [cited 2024 Jun 11]. Available from: <https://tradingeconomics.com/south-africa/employment-in-agriculture-percent-of-total-employment-wb-data.html>
- Food and Agriculture Organization of the United Nations. South Africa: BEFS Country briefs [document on the Internet]. c2013 [cited 2024 Jun 04]. Available from: <https://www.fao.org/4/aq181e/aq181e.pdf>
- Marloth RH. The scientific development of citriculture in southern Africa. *S Afr J Sci.* 1951;47:82–92.
- Moore ID. The development of the South African citrus industry. *Agrekon.* 1962;1:6–16. <https://doi.org/10.1080/03031853.1962.9524618>
- Zondo B. South Africa's citrus industry performance during the 2021/22 season: A focus on oranges and soft citrus. In: South Africa fruit trade flow [document on the Internet]. c2022 [cited Jun 08]. Available from: https://www.namc.co.za/wp-content/uploads/2023/02/SA-Fruit-Flow-Report-Issue-48-December-2022_Final.pdf
- Fourie P, van Niekerk J, Pretorius MC, Schutte T, Pietersen G, Labuschagne N, et al. Citrus pathology. In: Coutinho TA, Wingfield MJ, editors. History of plant pathology in South Africa. Pretoria: Briza; 2022. p. 63–68.
- Burt-Davy J. The maize industry of South Africa. *S Afr Agric J.* 1911;2:588.
- Vorster P. The maize industry. *Farming S Afr.* 1951 March:171–189.
- Greyling JC, Pardey PG. Measuring maize in South Africa: The shifting structure of production during the twentieth century 1904–2015. *Agrekon.* 2019;58:21–41. <https://doi.org/10.1080/03031853.2018.1523017>
- Lipton M. South Africa. Two agricultures? In: Wilson F, Kooy A, Hendrie D, editors. Farm labour in South Africa. Cape Town: David Philip; 1977.
- Statista. Maize production in South Africa 2022/2023, by province [website on the Internet]. c2024 [cited 2024 Jun 05]. Available from: <https://www.statista.com/statistics/1135488/maize-production-in-south-africa-by-province/>
- Flett B. Maize pathology. In: Coutinho TA, Wingfield MJ, editors. History of plant pathology in South Africa. Pretoria: Briza; 2022. p. 89–93.
- Bosman GJ, Osburn IB. The maize industry in South Africa. *Agric J S Afr.* 1924;15:83–86.
- ADAMA. Maize farming in South Africa [website on the Internet]. c2023 [cited 2024 Jun 04]. Available from: <https://www.adama.com/south-africa/en/maize-farming/maize-farming-south-africa>
- Grouse M. GM maize as subsistence crop: The South African smallholder experience. *AgBioForum.* 2012;15:163–174.
- Ala-Kokko K, Nalley LL, Shew AM, Tack JB, Chaminuka P, Matlock MD, et al. Economic and ecosystems impacts of GM maize in South Africa. *Glob Food Secur.* 2021;29, Art. #100544. <https://doi.org/10.1016/j.gfs.2021.100544>
- Mackellar N, New M, Jury C. Observed and modelled trends in rainfall and temperature for South Africa: 1960–2010. *S Afr J Sci.* 2014;110(7/8), Art. #2013-0353. <https://doi.org/10.1590/sajs.2014/20130353>
- Vogel C. (Mis)management of droughts in South Africa: Past, present and future. *S Afr J Sci.* 1994;90:4–6.
- Nepfumbada MP, Seetal A. National water security framework for South Africa: Summary, principles and recommendations [document on the Internet]. c2020 [cited 2024 Jun 08]. Available from: <https://www.nationalplanningcommission.org.za/assets/Documents/National%20Water%20Security%20Framework%20for%20South%20Africa.pdf>
- Diamond C, Braine H. Notes on irrigation in South Africa. *Transvaal Agric J.* 1905;4:8–21.
- Braine CDH. Two methods of farm irrigation. *S Afr J Sci.* 1911;7:242–250.
- Young A. A subterranean tide in the Karroo. *S Afr J Sci.* 1905;2:144–151.
- Stallebras JL. The Orange River development project. *S Afr J Sci.* 1963;59:448–454.
- Van Reenen RJ. Development of irrigation in the Union of South Africa. *S Afr J Sci.* 1925;22:20–41.
- Chunnett EP. Indirect benefits as the major incentive to the establishment of irrigation schemes. *S Afr J Sci.* 1965;61:99–103.
- Smajstrla AG. Microirrigation for citrus production in Florida. *HortSci.* 1993;28:295–298. <https://doi.org/10.21273/HORTSCI.28.4.295>
- Carr MKV. The water relations and irrigation requirements of citrus (*Citrus* spp.): A review. *Exp Agric.* 2012;48:347–377. <https://doi.org/10.1017/S0014479712000038>
- Ndhleve S, Nakin MDV, Longo-Mbenza B. Impacts of supplementary irrigation as a climate change adaptation strategy for maize production: A case of the Eastern Cape Province of South Africa. *Water SA.* 2017;43:222–228. <https://doi.org/10.4314/wsa.v43i2.06>
- Magombeyl MS, Rollin D, Lankford B. The river basin game as a tool for collective water management at community level in South Africa. *Phys Chem Earth.* 2008;33:873–880. <https://doi.org/10.1016/j.pce.2008.06.045>
- South African Irrigation Institute. Africa's first centre pivot: Pioneers. *SABI Magazine* 1(4) [document on the Internet]. c2009 [cited Jun 08]. Available from: <https://hdl.handle.net/10520/EJC92130>
- Van der Walt I, Richter BW. The effectiveness of centre-pivot irrigation from dolomitic ground water in the Northwest Province, South Africa. *Geol J.* 2004;61:197–201. <https://doi.org/10.1007/s10708-004-2888-z>
- Union of South Africa. Final report of the Drought Investigation Commission. Cape Town: Government Printers; 1923. p. 222.
- Du Toit AL. Geological and soil characteristics in relation to deterioration and conservation of natural resources, etc., in the Union. *S Afr J Sci.* 1938;35:470–476.
- Griffin A, Kay A, Sayers P, Bell V, Stewart E, Carr S. Widespread flooding dynamics changing under climate change: Characterizing floods using UKCP18 [preprint]. *Hydrol Earth Syst Sci.* 2022. <https://doi.org/10.5194/hess-2022-243>
- Mills AJ, Fey MV. Declining soil quality in South Africa: Effects of land use on soil organic matter and surface crusting: A review article. *S Afr J Sci.* 2003;99:429–436.
- Du Preez CC, Huyssteen CW. Threats to soil and water resources in South Africa. *Environ Res.* 2019;183, Art. #109015. <https://doi.org/10.1016/j.envres.2019.109015>
- Van Zyl JP. Phosphorus deficiency in South African soils and vegetation. *S Afr J Sci.* 1926;23:244–252.
- Ratlabala ME. An overview of South Africa's mineral based fertilizers. Pretoria: Department of Minerals and Energy; 2003.
- Hall TD, Meredith DBD, Altona RE. The agriculture of the Union of South Africa with reference to major crops and fertilizer usage. *S Afr J Sci.* 1951;41:213–214.
- Altona RE. Agriculture in the Republic of South Africa: Past, present and future. *S Afr J Sci.* 1963;59:517–524.
- Fertasa. Import and export statistics [webpage on the Internet]. c2024 [cited 2024 Jun 04]. Available from: <https://www.fertasa.co.za/>
- Du Preez C, van Huyssteen C, Mnkeni P. Land use and soil organic matter in South Africa 1: Review on spatial variability and the influence of rangeland stock production. *S Afr J Sci.* 2011;107(5/6), Art. #354. <https://doi.org/10.4102/sajs.v107i5/6.354>



44. Coutinho TA, Wingfield MJ. History of plant pathology in South Africa. Pretoria: Briza; 2022. p. 190.
45. Gilliomoe JH. Entomology in South Africa: Where do we come from, where are we now and where are we going? S Afr J Sci. 2013;109(1/2), Art. #a004. <https://doi.org/10.1590/sajs.2013/a004>
46. Price RE. Invasions and local outbreaks of four species of plague locusts in South Africa: Historical review of outbreak dynamics and patterns. Insects. 2023;14(11), Art. #846. <https://doi.org/10.3390/insects14110846>
47. Notes from: Stellenbosch-Elsenburg College of Agriculture University of Stellenbosch. Sulphuring of vines. Farming S Afr. 1929 February:1251.
48. Brundtland GH. Our common future: Report of the World Commission on Environment and Development. UN-Dokument A/42/427. New York: United Nations; 1987.
49. Korsten L. Biological control in Africa: Can it provide a sustainable solution for control of fruit diseases? S Afr J Bot. 2004;70:128–139. [https://doi.org/10.1016/S0254-6299\(15\)30273-8](https://doi.org/10.1016/S0254-6299(15)30273-8)
50. Gregory PJ, Johnson SN, Newton AC, Ingram JSI. Integrating pests and pathogens into the climate change/food security debate. J Environ Bot. 2009;60:2827–2838. <https://doi.org/10.1093/jxb/erp080>
51. Shahzad A, Ullah S, Dar AA, Sardar MF, Mehmood T, Tufail MA, et al. Nexus on climate change: Agriculture and possible solution to cope future climate change stresses. Environ Sci Pollut Res. 2021;28:14211–14232. <https://doi.org/10.1007/s11356-021-12649-8>
52. Doidge EM. Some diseases of citrus prevalent in South Africa. S Afr J Sci. 1929;26:320–325.
53. Doidge EM. The role of bacteria in plant diseases. S Afr J Sci. 1919;16:65–92.
54. Carstens E, le Roux H, Holtzhausen MA, van Rooyen L, Coetzee J, Wentzel R, et al. Citrus black spot is absent in the Western Cape, Northern Cape and Free State Provinces. S Afr J Sci. 2012;108(7/8), Art. #876. <https://doi.org/10.4102/sajs.v108i7.8.876>
55. Moyo P, du Raan S, Fourie PH. Models for predicting pseudothecium maturity and ascospore release of *Phyllosticta* spp. in South African citrus orchards. S Afr J Sci. 2020;116(11/12), Art. #7955. <https://doi.org/10.17159/sajs.2020/7955>
56. McClean APD. Citrus vein-eneation virus. S Afr J Sci. 1954;50:147–151.
57. Smit B. A study of the most important insect pests that confront the citrus growth in the eastern province. S Afr J Sci. 1934;31:439–441.
58. McClean APD, Oberholzer PCJ. Citrus psylla, a vector of the greening disease of sweet orange – disease note. S Afr J Agric Sci. 1965;8:297–298.
59. Mokgoba MC, Tshikhudo PP, Nnzeru LR, Makhado RA. Impact of fall armyworm (*Spodoptera frugiperda*) (J.E. Smith) on small-scale maize farmers and its control strategies in the Limpopo province, South Africa. Jamba – J Disaster Risk Stud. 2021;13, Art. #1016. <https://doi.org/10.4102/jamba.v13i1.1016>
60. Beukes I, Rose LJ, Shephard GS, Flett BC, Viljoen A. Mycotoxigenic *Fusarium* species associated with grain crops in South Africa – a review. S Afr J Sci. 2017;113(3/4), Art. #2016-0121. <https://doi.org/10.17159/sajs.2017/20160121>
61. Berger DK, Mokgobu T, de Ridder K, Christie N, Aveling TAS. Benefits of maize resistance breeding and chemical control against northern leaf blight in smallholder farms in South Africa. S Afr J Sci. 2020;116(11/12), Art. #8286. <https://doi.org/10.17159/sajs.2020/8286>
62. Gerwin V. Genetically modified corn – environmental benefits and risks. PLoS Biol. 2003;1:15–19. <https://doi.org/10.1371/journal.pbio.0000008>
63. Hudson DA. Southern Africa climate change stimulated by GENESIS GCM. S Afr J Sci. 1997;93:389–403.
64. New M. Climate change and water resources in the southwestern Cape, South Africa. S Afr J Sci. 2002;98:369–376.
65. Chami DEI, Moujabber ME. Drought, climate change and sustainability of water in agriculture: A roadmap towards the NWR2. S Afr J Sci. 2016;112(9/10), Art. #2015-0457. <https://doi.org/10.17159/sajs.2016/20150457>
66. Jury MR. Climate trends across South Africa since 1980. Water SA. 2018;44:297–307. <https://doi.org/10.4314/wsa.v44i2.15>
67. Blignaut J, Ueckermann L, Aronson J. Agriculture production's sensitivity to changes in climate in South Africa. S Afr J Sci. 2009;105(1/2):61–67.
68. Balfagón D, Arbona C, Gómez-Cadenas A. The future of citrus fruit: The impact of climate change on citriculture. Méthode Sci Stud J. 2022;12:123–129.
69. Cammarano D, Valdivia RO, Beletse YG, Durand W, Crespo O, Tesfahuney WA, et al. Integrated assessment of climate change impacts on crop productivity and income of commercial farms in northeast South Africa. Food Secur. 2020;12:659–678. <https://doi.org/10.1007/s12571-020-01023-0>
70. Abobatta EF. Citriculture and climate change. Mod Concepts Develop Agron. 2020;6:649–650. <https://doi.org/10.31031/MCDA.2020.06.000639>
71. Johnston P, Egbeyi TS, Zvobgo L, Omar SA, Cartwright A, Hewitson B. Climate change impacts in South Africa: What climate change means for a country and its people [webpage on the Internet]. c2024 [cited 2024 Jun 08]. Available from: <https://www.csag.uct.ac.za/2024/02/28/new-report-what-climate-change-means-forsouth-africa-and-its-people/>
72. Ziervogel G, Lennard C, Midgley G, New M, Simpson NP, Trisos CH, et al. Climate change in South Africa: Risks and opportunities for climate-resilient development in the IPCC Sixth Assessment WGII Report. S Afr J Sci. 2022;118(9/10):14492. <https://doi.org/10.17159/sajs.2022/14492>
73. Hendriks SL. The food security continuum: A novel tool for understanding food insecurity as a range of experiences. Food Secur. 2015;7:609–619. <https://doi.org/10.1007/s12571-015-0457-6>
74. USDA ERS. Overview [webpage on the Internet]. c2024 [cited 2024 Jun 13]. Available from: <https://www.ers.usda.gov/topics/food-nutrition-assistance/food-security-in-the-us/>
75. El Blali H, Bassole IHN, Dambo L, Berjan S. Climate change and food security. Agric For. 2020;3:197–210. <https://doi.org/10.17707/agricultForest.66.3.16>
76. Dlamini SN, Craig A, Mtintsilana A, Mapanga W, du Toit J, Ware LJ, et al. Food insecurity and coping strategies and their association with anxiety and depression: A nationally representative South African survey. Pub Health Nutr. 2023;26:705–715. <https://doi.org/10.1017/S1368980023000186>
77. World Bank. Poverty and equity brief: South Africa [document on the Internet]. c2023 [cited 2024 June 05]. Available from: https://databankfiles.worldbank.org/public/ddpext_download/poverty/987B9C90-CB9F-4D93-AE8C-750588BF00QA/current/Global_POVEQ_ZAF.pdf
78. Mbhatsani HV, Mabapa NS, Ayuk TB, Madiwana TC, Mushaphi LF, Mohlala M, et al. Food security and related health risk among adults in the Limpopo Province of South Africa. S Afr J Sci. 2021;117(11/12), Art. #8848. <https://doi.org/10.17159/sajs.2021/8848>
79. Burger R, du Plessis L, Gangaldzo T, Wills G. Priority areas or reducing stunting in South Africa: Examining the implications of recent international evidence. Ilifa Labantwana & Ewawp ECD Working Paper Series No. ECD WP 004/2022 [document on the Internet]. c2022 [cited 2024 Jun 08]. Available from: <https://ilifalabantwana.co.za/wp-content/uploads/2022/08/Ilifa-Labantwana-Reducing-Stunting-in-SA-report-V05.pdf>
80. Fresco LO. Some thoughts about the future of food and agriculture. S Afr J Sci. 2014;110(5/6), Art. #a0066. <https://doi.org/10.1590/sajs.2014/a0066>
81. Mashamaite CV, Pieterse PJ, Mothapo PN, Phiri M. *Moringa oleifera* in South Africa: A review on its production, growing conditions and consumption as a food source. S Afr J Sci. 2021;117(3/4), Art. #8689. <http://dx.doi.org/10.17159/sajs.2021/8689>
82. Allemann J, Hammes PS. Chemical composition of South African *Plectranthus esculentus* tubers: Research in action. S Afr J Sci. 2003;99:127–129.
83. South African Agriculture and Agroprocessing Sector Working Group. Foresight agricultural report [document on the Internet]. c2024 [cited 2024 Aug 10]. Available from: <https://www.dst.gov.za/images/pdfs/FORESIGHT%20AGRICULTURE%20REPORT.pdf>
84. South African Department of Agriculture. Strategic plan for the Department of Agriculture 2008–2011 [document on the Internet]. c2008 [cited 2024 Aug 10]. Available from: <http://www.daff.gov.za/phocadownloadpdp/Strategy/Strategic%20Plan%20for%20the%20Department%20of%20Agriculture%202008%20092010%2011.pdf>