











Pathogen surveillance and epidemiology in endangered peninsular bighorn sheep (*Ovis canadensis nelsoni*)

Jessica N. Sanchez¹  | Brandon A. Munk²  | Janene Colby³ |
 Steve G. Torres² | Ben J. Gonzales² | James R. DeForge⁴  |
 Aimee J. Byard⁴  | Lora Konde² | Nicholas J. Shirkey²  |
 Pranav S. Pandit¹  | Randy A. Botta³  | Annette Roug⁵  |
 Michael H. Ziccardi⁶  | Christine K. Johnson¹ 

¹EpiCenter for Disease Dynamics, School of Veterinary Medicine, One Health Institute, University of California at Davis, Davis, California, USA

²Wildlife Health Lab, California Department of Fish and Wildlife, Rancho Cordova, California, USA

³California Department of Fish and Wildlife, South Coast Region, San Diego, California, USA

⁴Bighorn Institute, Palm Desert, California, USA

⁵Department of Production Animal Medicine, Faculty of Veterinary Science, Centre for Veterinary Wildlife Research, University of Pretoria, Pretoria, South Africa

⁶School of Veterinary Medicine, One Health Institute, University of California at Davis, Davis, California, USA

Correspondence

Christine K. Johnson, EpiCenter for Disease Dynamics, School of Veterinary Medicine, One Health Institute, University of California at Davis, 1089 Veterinary Medicine Drive, Davis, CA 95616, USA.

Email: ckjohnson@ucdavis.edu

Funding information

National Institutes of Health, Grant/Award Number: T32 OD 011147

Abstract

Peninsular bighorn sheep (*Ovis canadensis nelsoni*) are found exclusively in Southern California and Baja Mexico. They are federally endangered due to multiple threats, including introduced infectious disease. From 1981 to 2017, we conducted surveillance for 16 pathogens and estimated population sizes, adult survival, and lamb survival. We used mixed effects regression models to assess disease patterns at the individual and population levels. Pathogen infection/exposure prevalence varied both spatially and temporally. Our findings indicate that the primary predictor of individual pathogen infection/exposure was the region in which an animal was captured, implying that transmission is driven by local ecological or behavioral factors. Higher *Mycoplasma ovipneumoniae* seropositivity was associated with lower lamb survival, consistent with lambs having high rates of pneumonia-associated mortality, which may be slowing population recovery. There was no association between *M. ovipneumoniae* and adult survival. Adult survival was positively associated with population size and parainfluenza-3 virus seroprevalence in the same year, and orf virus seroprevalence in the previous year. Peninsular bighorn sheep are recovering from small population sizes in a habitat of environmental extremes, compounded by infectious disease. Our research can help inform future pathogen surveillance and population monitoring for the long-term conservation of this population.

KEYWORDS

endangered species, epidemic pneumonia, lamb recruitment, *Mycoplasma ovipneumoniae*, pathogen spillover, Peninsular ranges, survival, wildlife-livestock interface

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *Conservation Science and Practice* published by Wiley Periodicals LLC on behalf of Society for Conservation Biology.

1 | INTRODUCTION

Pneumonia epidemics are a source of mortality and decreased lamb survival in bighorn sheep (*Ovis canadensis nelsoni*) throughout much of their range (Besser et al., 2013; DeForge et al., 1982; Nolen, 2010). Pathogens associated with severe pneumonia are introduced to bighorn sheep herds through contact with domestic sheep (Foreyt & Jessup, 1982) but can be maintained by carrier bighorn sheep for years without continued spillover from domestic animals (Raghavan et al., 2016), causing intermittent epidemics in lambs and suppressing recruitment (Cassirer et al., 2018). Bighorn sheep pneumonia is a disease complex involving coinfection with multiple pathogens, environmental and immune factors, and host behavior (Besser et al., 2013; Wobeser, 1994). Recent research indicates that *Mycoplasma ovipneumoniae* infection can cause pneumonia by decreasing respiratory immune function and allowing colonization by other pathogens (Besser et al., 2012, 2014; Dassanayake et al., 2010). Numerous management tools including vaccination, population reduction, and supplemental feeding have failed to prevent or control pneumonia outbreaks in bighorn sheep (Cassirer et al., 2001, 2018; Ward et al., 1999) but recent efforts to test and remove chronic *M. ovipneumoniae* carriers demonstrate promising results, including improved lamb survival (Garwood et al., 2020).

Peninsular bighorn sheep (*O. c. nelsoni*) reside in the Peninsular Ranges of southern California and Baja Mexico, and are currently considered a genetically distinct metapopulation of desert bighorn sheep (Buchalski et al., 2016). Peninsular bighorn sheep were listed as federally endangered in 1998 due to a multitude of population threats, including habitat loss and fragmentation, infectious disease, predation, and drought (US Fish and Wildlife Service, 2000). The Peninsular metapopulation has been steadily increasing in size from ~300 at the time of listing to ~900 in 2016; however, infectious disease continues to threaten survival and recruitment (Colby & Botta, 2019).

Bighorn sheep behavior and spatial distribution play a role in the transmission and maintenance of disease. The Peninsular bighorn sheep metapopulation consists of at least 19 herds that inhabit the desert slopes, alluvial fans, and washes of the Peninsular Ranges (Colby & Botta, 2019). While most individuals within each herd are philopatric, a subset of ewes and rams will disperse to neighboring herds on a seasonal basis (Bighorn Institute, 2018; Buchalski et al., 2015; Colby & Botta, 2019). The Peninsular mountains are divided into nine “recovery regions” (hereafter, “regions”) defined for bighorn sheep population management (US Fish and Wildlife Service, 2000; Figure 1). Historically, these regions were thought to roughly correspond to different herds (Rubin et al., 1998) but some regions now

contain multiple overlapping herds and inter-regional movements are regularly observed (Bighorn Institute, 2018; Colby & Botta, 2019).

Bighorn sheep movements are driven by food and water availability, which are especially scarce in drought years. California has had chronically low rainfall for several decades, including a severe drought from 2012 to 2016 that significantly reduced the surface water available for wildlife in desert ecosystems where bighorn sheep are found (Lund et al., 2018). Bighorn sheep congregate in high densities at natural and artificial water sources and urban areas where irrigation and landscaping provide resources (Bighorn Institute, 2018; Colby & Botta, 2019; Figure 1). This comingling of animals from different herds, age classes, and disease statuses increases the risk of pathogen transmission, and higher density herds are associated with an increased risk of respiratory disease outbreaks (Monello et al., 2001; Sells et al., 2015).

Peninsular bighorn sheep are recovering in a desert ecosystem that is evolving with climate change. The goal of our research is to identify key epidemiologic factors driving disease prevalence in bighorn sheep, with special attention paid to pathogens associated with epidemic pneumonia. We aim to: (1) Estimate pathogen prevalence by age, sex, and region; (2) Identify demographic and geographic predictors of pathogen infection/exposure in individual bighorn sheep; (3) Identify associations between pathogen infection/exposure prevalence and adult and lamb survival while controlling for important environmental variables associated with climate change; (4) Identify communities of pathogens that co-occur together within individual bighorn sheep and may impact fitness.

2 | METHODS

2.1 | Pathogen infection and exposure prevalence

We sampled wild Peninsular bighorn sheep captured from 1981 to 2017. Animals were captured and sampled by the California Department of Fish and Wildlife (CDFW) or contractors following CDFW guidelines. Protocols were reviewed and approved by CDFW, or other land management agencies when appropriate, and objectives and goals have been directed by the Peninsular Bighorn Sheep Recovery Plan since 2000 (US Fish and Wildlife Service, 2000).

We tested blood samples (735 individuals; 844 sampling events) for infection or exposure to up to 15 pathogens, including *Anaplasma* spp., bluetongue virus (BTV), bovine herpesvirus-1, bovine respiratory syncytial virus (BRSV), bovine viral diarrhea virus Types 1 and 2,




FIGURE 1 Map of the study area within the Peninsular ranges of southern California, USA. Map depicts recovery region boundaries, bighorn sheep herd home ranges, golf course communities bordering, or within bighorn sheep habitat, and primary water sources. Major riparian areas have perennial or intermittent creeks and relatively large amounts of vegetation, including canopy cover and a dense understory. These areas are also utilized by deer and sometimes mountain lions. Artificial ponds and guzzlers provide year-round water through municipal sources or by collecting rainwater then delivering them to a drinking area. Guzzlers tend to be elevated, while ponds are at ground level and therefore vulnerable to contamination by rain runoff and/or animal excrement. Natural seeps and springs are small point sources of water at ground level that contain variable quantities and quality of water throughout the year. Tenajas are small rock depressions that hold water at the bottom of drainages, tend to be poor water quality, and are not dependable during the summer months. Golf course communities shown are those that bighorn sheep utilize on a regular basis; they are in urban areas where human–wildlife conflict is likely, but also have highly nutritious forage and many dependable water sources such as ponds, creeks, canals, reservoirs, and swimming pools

Brucella ovis, *Chlamydia* spp., epizootic hemorrhagic disease virus (EHDV), *Leptospira* spp., *M. ovipneumoniae*, ovine progressive pneumonia virus, orf virus, parainfluenza-3 virus (PI-3), and *Toxoplasma gondii*. Virus isolation (VI) was performed for BT and EHDV, but all other blood tests measured antibodies and more likely indicated previous

exposure. “Prevalence” hereafter refers to the proportion of positive tests, indicating infection or exposure depending on the test used.

In some years, nasal/pharyngeal swabs were also collected (316 individuals; 349 sampling events) and tested for combinations of *M. ovipneumoniae* via polymerase

chain reaction (PCR), *Pasteurellaceae* spp. via culture, and PI-3 via VI to detect active infection (or very recent exposure).

BTV and EHDV are both orbiviruses and cross-react on agar gel precipitin (AGP) and agar gel immunodiffusion (AGID), so we created an “*Orbivirus* spp.” group which included animals positive for BTV and/or EHDV via AGP/AGID. We classified animals as exposed to BTV if they tested positive on the more specific competitive enzyme-linked immunosorbent assay (cELISA).

We did not include *Leptospira* spp. serovars in analyses due to cross-reaction on the modified agglutination test, and an animal was considered positive if any serovar was detected at titers >1:100.

Age, sex, and region were recorded at the time of capture. Age was usually recorded categorically based on dentition and horn growth rings, with lambs and yearlings grouped together and older animals categorized as adults. The dataset was skewed towards adult females (80.3%, $n = 590/735$) since they were the target population for radio-collaring (Colby & Botta, 2019). Most individuals were only captured once ($n = 641$).

We summarized counts of animals that tested positive versus negative for each pathogen, then stratified by age, sex, and region. We tested for differences among 2 groups using Fisher's exact test and among ≥ 3 groups using one-way analysis of variance (significance at $p \leq .05$). These calculations only included samples from first capture events to eliminate retesting errors and biases due to persistent antibodies. We also calculated overall prevalence of each pathogen for each diagnostic test type, summarized for each recovery region and year, including all capture events. All statistics were performed in R version 4.0.4 (R Core Team, 2021).

Annual adult survival rates (June_{t-1} to May_t) for each region were previously calculated by CDFW and Bighorn Institute using Kaplan Meier estimates from radio-collared bighorn sheep, modified to allow for staggered entry (Bighorn Institute, 2018; Colby & Botta, 2019; Ostermann et al., 2001). Lamb survival is considered to be an excellent demographic predictor of health status in bighorn sheep populations (Cassirer et al., 2013). In the Peninsular Ranges, the majority of pneumonia-induced deaths in lambs occur between 8 and 10 weeks (Colby & Botta, 2013). Lamb survival for each region was evaluated based on the ratio of lambs to ewes (lamb:ewe) estimated from observations made during range-wide helicopter surveys, waterhole counts, or ground observations (Bighorn Institute, 2018; Colby & Botta, 2019). This was used as a proxy of lamb survival to ~3–9 months, depending on when surveys were performed. Pregnancy rates in the Peninsular Ranges are consistently high, with 94.3% of radio-collared ewes 2–19 years of age giving

birth from 2005 to 2022, and twins are rare (CDFW, unpublished data). Therefore, lamb:ewe ratios are primarily a reflection of lamb survival rather than birth rates.

2.2 | Population-level risk factors associated with adult and lamb survival

We created “population-level models” using Bayesian, multilevel, ordered beta regression to evaluate associations between annual adult survival or lamb:ewe ratios (outcomes), and pathogen prevalence, population size, and meteorologic covariates. We also evaluated bivariate relationships between model covariates, including year, as part of model building with univariable, ordered beta regression models. The unit of analysis was the year-region unit, and the random intercept was region. We selected weakly informative priors for intercept and beta parameters [Normal(0, 5)], and phi parameter [$\exp(0.1)$] (Kubinec, 2020). Population size, meteorologic covariates, year, and lamb survival were min–max scaled as needed to match outcome variables, so values ranged from 0 to 1 but the relative differences between values were maintained. This was done for each variable by subtracting the minimum value from each x , then dividing by the range of the original variables.

We calculated pathogen prevalence for each year-region unit (i.e., “2010—San Jacinto Mountains”) for which ≥ 5 samples were tested (all capture events included). Models included individual pathogens and combinations of respiratory pathogens (*M. ovipneumoniae*, BRSV, and PI-3) as covariates, and we tested prevalence lag times of -1 to $+1$ years.

We interpolated missing annual population size estimates for each region by averaging values for year_{t-1} and year_{t+1}, but only where estimates were missing for a single year.

Temperature and precipitation for each region were included to control for meteorologic factors influencing survival. We extracted rasters of daily meteorologic data (4×4 km resolution) from “gridMET” (Abatzoglou, 2013) using the “climateR” package (Johnson, 2022), then cropped by the geographic extent of each recovery region and aggregated temporally as described below, resulting in a single summary value for each year-region unit.

We calculated temperature as the average daily maximum temperature (Celsius) from June to September for year_t, which have historically been the hottest months in the Peninsular Ranges (Rubin et al., 2000; Turner et al., 2004).

Precipitation in the Peninsular Ranges is bimodal, with the largest volume and most consistent rains

occurring November to February, and more variable monsoons occurring July to September (Rubin et al., 2000). We calculated annual precipitation as the sum of daily precipitation (centimeters) from November of year_{t-1} through October of year_t. We also aggregated precipitation annually for winter (November_{t-1} to April_t) and summer (May_t to October_t). Winter corresponded to the winter rains and bighorn sheep gestational and peak birthing period, whereas summer corresponded to summer monsoons, postlambing, and the rut (Colby & Botta, 2017; Rubin et al., 2000; Rubin, Boyce et al., 2002).

2.3 | Individual-level risk factors associated with pathogen exposure or infection

We created “individual-level models” using Bayesian, multilevel, logistic regression to evaluate risk factors for an animal being infected or exposed to a pathogen. Predictor variables included age class (lamb/yearling, adult), sex (female, male), and recovery region (categorical, $n = 9$). Reference groups were adults, males, and the San Jacinto Mountains. The unit of analysis was the individual animal, and the random intercept was animal ID (all capture events included). We selected weakly informative priors [Normal(0, 2.5)] for intercept and beta parameters to account for complete or quasiseparation of data (Ghosh et al., 2018).

All regression models were built in package “brms” (Bürkner, 2017, 2018). Models contained 4 chains with 10,000 iterations each. We calculated point estimates as the median value of the posterior and used the 95% highest density interval as the credible interval (CI). We only included models with ≥ 5 observations per covariate in results. We compared models with the same number of observations using leave-one-out cross-validation information criterion (LOO IC) in the “loo” package (Vehtari et al., 2020). Appendix S1 contains heatmaps illustrating model variables by year and region.

2.4 | Pathogen co-occurrence network

We looked for “communities” of pathogens to which individual bighorn sheep were coinfecting/coexposed. Concurrent or subsequent infections can take a toll on host immune function and overall fitness, as has been demonstrated by the polymicrobial nature of pneumonia in bighorn sheep (Asghar et al., 2015; Besser et al., 2008; Jamieson et al., 2013). Clusters of pathogens that regularly co-occur together could be associated with clinical phenotypes and direct future disease surveillance.

Since the diagnostic tests used in this study can indicate infection or previous exposure, we use “co-occurrence” to mean that an animal was infected with a pair of pathogens during its life, but perhaps not concurrently. We generated a weighted, undirected network from the proportion of samples (from first capture events only) that were positive for two pathogens, given that they were tested for both pathogens. Density and betweenness centrality were used to describe how tightly pathogens clustered (package “sna”) (Butts, 2008). We calculated network modularity using the fast greedy modularity optimization algorithm (package “igraph”) (Clauset et al., 2004) and visualized the network in Gephi (Bastian et al., 2009).

3 | RESULTS

3.1 | Pathogen infection and exposure prevalence

A total of 735 first-capture samples were collected from 1981 to 2017. Pathogen and antibody prevalence estimates are in Table 1, including stratifications by age, sex, and region. Not all diagnostic tests were performed in every year or region.

The pathogens with the highest seroprevalence were orf virus (71.8%), *M. ovipneumoniae* (cELISA; 60.2%), *Anaplasma* spp. (49.7%), *Chlamydia* spp. (42.8%), BRSV (39.3%), EHDV (serum virus neutralization [SVN]; 24.4%), *Orbivirus* spp. (21.6%), PI-3 (hemagglutination inhibition [HI]; 21.2%), and *T. gondii* (18.0%).

The most common active infection was *Pasteurella*-*ceae* spp., with all samples tested culturing at least one species. *Mannheimia haemolytica* betahemolytic and *Bibersteinia trehalosi* nonhemolytic were the most common *Pasteurellaceae* spp. (85.0% and 77.9%, respectively). *Pasteurella multocida* was not detected from first capture events, but was cultured in a single sample from a recaptured animal. Prevalence of active infection was much lower for *M. ovipneumoniae* (PCR; 12.0%) and PI-3 (VI; 11.4%). All other pathogens were relatively uncommon or absent (Table 1).

Leptospira spp. was found in 12.1% of animals, across six serovars (although possible cross-reaction makes these diagnoses unreliable): *Leptospira interrogans* serovars bratislava (19.2%, $n = 10/52$), pomona (1.6%, $n = 5/316$), canicola (1.3%, $n = 4/316$), and icterohaemorrhagiae (0.3%, $n = 1/316$); *Leptospira kirschneri* serovar grippotyphosa (5.7%, $n = 18/316$); *Leptospira borgpeterse-nii* serovar hardjo (1.6%, $n = 5/310$).

Females had higher rates of exposure to BRSV ($p = .01$), whereas males had higher exposure to *T. gondii* ($p < .001$) and *Orbivirus* spp. ($p = .03$). Lambs/yearlings

TABLE 1 Summary of pathogen infection/exposure prevalence in Peninsular bighorn sheep across the entire study period (1981–2017), stratified by sex, age, and recovery region (first capture event for each individual).

Pathogen	Test type	Antibody (A) or pathogen (P)	Sex		Recovery region												
			Overall	Female	Male	Age			Northern			Central			Southern		
						Lamb/ yearling	Adult	San Jacinto	Santa Rosa	Santa Rosa	Santa Rosa	Santa Rosa	Santa Rosa	Santa Rosa	Santa Rosa	Santa Rosa	Santa Rosa
<i>Anaplasma</i> spp.	CA	A	49.7% (158/318)	48.1% (117/243)	55.4% (41/74)	34.6% (9/26)	51.0% (148/290)	42.9% (12/28)	29.8% (14/47)	48.3% (14/29)	29.6% (8/27)	57.1% (20/35)	77.1% (27/35)	65.9% (29/44)	48.5% (16/33)	45.0% (18/40)	
Bovine herpesvirus-1	SVN	A	0.6% (3/537)	0.2% (1/407)	1.6% (2/129)	0.0% (0/66)	0.6% (3/469)	0.0% (0/51)	3.0% (3/100)	0.0% (0/57)	0.0% (0/32)	0.0% (0/45)	0.0% (0/51)	0.0% (0/49)	0.0% (0/45)	0.0% (0/107)	
Bovine respiratory syncytial virus	IFA	A	39.3% (259/659)	41.9% (222/530)	28.9% (37/128)	34.8% (23/66)	39.9% (236/592)	12.1% (8/66)	48.7% (38/78)	37.0% (30/81)	72.9% (43/59)	43.8% (21/48)	28.1% (18/64)	44.1% (26/59)	40.0% (30/75)	34.9% (45/129)	
Bovine viral diarrhoea virus type-1	SVN	A	0.7% (4/541)	0.7% (3/411)	0.8% (1/129)	3.0% (2/67)	0.4% (2/472)	0.0% (0/51)	1.0% (1/101)	0.0% (0/57)	0.0% (0/32)	0.0% (0/45)	0.0% (0/54)	0.0% (0/49)	0.0% (0/45)	2.8% (3/107)	
Bovine viral diarrhoea virus type-2	SVN	A	0.0% (0/83)	0.0% (0/65)	0.0% (0/17)	0.0% (0/7)	0.0% (0/76)	0.0% (0/7)	0.0% (0/4)	NT	NT	0.0% (0/7)	0.0% (0/9)	0.0% (0/2)	0.0% (0/42)		
<i>Brucella</i> <i>ovis</i>	ELISA	A	5.0% (23/459)	5.9% (20/340)	2.5% (3/118)	3.4% (2/59)	5.3% (21/399)	0.0% (0/38)	0.0% (0/87)	0.0% (0/50)	0.0% (0/26)	5.3% (2/38)	16.3% (7/43)	10.9% (5/46)	0.0% (0/38)	9.7% (9/93)	
<i>Chlamydia</i> spp.	CF	A	42.8% (199/465)	41.7% (145/348)	45.7% (53/116)	36.5% (19/52)	43.4% (179/412)	48.8% (20/41)	17.0% (9/53)	71.4% (40/56)	12.9% (4/31)	38.1% (16/42)	30.8% (16/52)	42.6% (20/47)	42.2% (19/45)	56.1% (55/98)	
<i>Leptospira</i> spp.	MAT	A	12.1% (38/313)	12.7% (30/237)	10.5% (8/76)	8.1% (3/37)	12.7% (35/275)	3.2% (1/31)	6.6% (6/66)	5.4% (2/37)	30.8% (8/26)	16.0% (4/25)	10.0% (3/30)	33.3% (8/24)	6.5% (2/31)	9.3% (4/43)	
<i>Mycoplasma</i> <i>ovipneumoniae</i>	PCR	P	12.0% (38/316)	11.9% (33/278)	13.5% (5/37)	14.3% (4/28)	11.8% (34/288)	13.3% (4/30)	5.9% (2/34)	3.1% (1/32)	15.2% (5/33)	7.7% (2/26)	0.0% (0/30)	20.0% (6/30)	23.8% (10/42)	13.6% (8/59)	
<i>Chlamydia</i> spp.	cELISA	A	60.2% (336/558)	59.5% (267/449)	63.0% (68/108)	58.3% (28/48)	60.3% (307/509)	55.6% (30/54)	71.2% (42/59)	58.1% (36/62)	55.8% (29/52)	48.9% (22/45)	56.7% (34/60)	50.0% (25/50)	71.0% (49/69)	64.5% (69/107)	
Ovine progressive pneumonia virus	AGID	A	0.0% (0/186)	0.0% (0/138)	0.0% (0/48)	0.0% (0/14)	0.0% (0/170)	0.0% (0/26)	0.0% (0/36)	0.0% (0/9)	0.0% (0/12)	0.0% (0/13)	0.0% (0/17)	0.0% (0/21)	0.0% (0/34)		
Orf virus	CF	A	71.8% (319/444)	70.4% (247/351)	77.2% (71/92)	57.1% (24/42)	73.6% (295/401)	70.3% (26/37)	62.8% (49/78)	50.0% (23/46)	84.2% (32/38)	69.2% (27/39)	83.7% (36/43)	82.2% (37/45)	63.6% (28/44)	82.4% (61/74)	
Parainfluenza-3 virus	VI	P	11.4% (4/35)	13.0% (3/23)	8.3% (1/12)	23.1% (3/13)	4.5% (1/22)	0.0% (0/2)	20.0% (4/20)	NT	NT	NT	NT	NT	NT	0.0% (0/13)	
<i>Toxoplasma</i> <i>gondii</i>	HI	A	21.2% (152/717)	21.4% (123/576)	20.7% (29/140)	12.5% (10/80)	22.2% (141/635)	1.5% (1/66)	32.5% (38/117)	9.9% (8/81)	30.5% (18/59)	8.9% (5/56)	12.3% (8/65)	19.4% (12/62)	35.4% (29/82)	25.6% (33/129)	
<i>Toxoplasma</i> <i>gondii</i>	LA	A	18.0% (16/89)	9.1% (6/66)	43.5% (10/23)	14.3% (1/7)	18.5% (15/81)	20.0% (4/20)	37.0% (10/27)	NT	NT	NT	5.9% (1/17)	7.7% (1/13)	0.0% (0/12)	0.0% (0/12)	
Bluetongue virus	VI	P	3.3% (3/91)	4.5% (3/66)	0.0% (0/25)	11.1% (2/18)	1.4% (1/72)	0.0% (0/16)	5.0% (2/40)	NT	0.0% (0/4)	0.0% (0/5)	NT	0.0% (0/5)	0.0% (1/21)	4.8% (1/21)	
Epizootic haemorrhagic disease virus	cELISA	A	10.4% (60/576)	9.6% (45/467)	13.9% (15/108)	3.8% (2/52)	11.1% (58/523)	8.0% (4/50)	5.2% (3/58)	3.7% (3/81)	8.5% (5/59)	5.4% (3/56)	22.9% (11/48)	12.3% (7/57)	18.3% (15/82)	10.6% (9/85)	
Epizootic haemorrhagic disease virus	VI	P	0.0% (0/22)	0.0% (0/14)	0.0% (0/8)	0.0% (0/10)	0.0% (0/12)	NT	0.0% (0/18)	NT	NT	NT	NT	NT	NT	0.0% (0/4)	
SVN	SVN	A	24.4% (10/41)	20.0% (6/30)	36.4% (4/11)	17.6% (3/17)	29.2% (7/24)	NT	18.9% (7/37)	NT	NT	NT	NT	NT	NT	75.0% (3/4)	

TABLE 1 (Continued)

Pathogen	Test type	Antibody (A) or pathogen (P)	Overall	Sex		Age		Recovery region														
				Female	Male	Lamb/ yearling	Adult	Northern			Central			Southern			Northern			Southern		
								San Jacinto	Santa Rosa	Mtns	Santa Rosa	Santa Rosa	Mtns	Coyote Canyon	Ysidro	Mtns	San Ysidro	San Ysidro	Mtns	Vallecito	Carrizo Canyon	
<i>Orbivirus</i> spp.	AGP/AGID	A	21.6% (145/670)	19.9% (107/538)	29.0% (38/131)	15.1% (11/73)	22.5% (134/595)	30.6% (19/62)	70.0% (7/10)	31.9% (37/116)	7.5% (4/53)	12.1% (7/58)	9.8% (5/51)	21.5% (14/65)	17.7% (11/62)	20.3% (15/74)	25.6% (33/129)	91.7% (11/12)	75.0% (9/12)			
<i>Mannheimia haemolytica</i> beta-hemolytic	Culture	P	85.0% (119/140)	86.5% (90/104)	80.6% (29/36)	100.0% (14/14)	83.3% (105/126)	7.0% (0/10)	85.7% (18/21)	87.0% (20/23)	100.0% (12/12)	90.0% (9/10)	90.0%	83.3% (15/18)	82.4% (13/17)	76.5% (13/17)	91.7% (11/12)	91.7% (11/12)	41.7% (5/12)			
<i>Mannheimia haemolytica</i> nonhemolytic	Culture	P	23.6% (33/140)	24.0% (25/104)	22.2% (8/36)	21.4% (3/14)	23.8% (30/126)	0.0% (0/10)	23.8% (5/21)	34.8% (8/23)	25.0% (3/12)	10.0% (1/10)	10.0%	33.3% (6/18)	17.6% (3/17)	11.8% (2/17)	11.8% (2/17)	11.8% (2/17)	11.8% (2/17)			
<i>Bibersteinia trehalosi</i> beta-hemolytic	Culture	P	12.1% (17/140)	15.4% (16/104)	2.8% (1/36)	7.1% (1/14)	12.7% (16/126)	0.0% (0/10)	0.0% (0/21)	0.0% (0/23)	0.0% (0/12)	0.0% (0/12)	30.0% (3/10)	38.9% (7/18)	0.0% (0/17)	29.4% (5/17)	16.7% (2/12)	16.7% (2/12)	16.7% (2/12)			
<i>Bibersteinia trehalosi</i> nonhemolytic	Culture	P	77.9% (109/140)	77.9% (81/104)	77.8% (28/36)	71.4% (10/14)	78.6% (99/126)	90.0% (9/10)	90.5% (19/21)	82.6% (19/23)	83.3% (10/12)	50.0% (5/10)	50.0%	61.1% (11/18)	94.1% (16/17)	64.7% (11/17)	64.7% (11/17)	64.7% (11/17)	75.0% (9/12)			

Note: Fractions represent the number of positive tests over the number of tests performed. Sample sizes within each stratification level may not sum to the same totals as in the "overall" column because age class and sex were not available for all samples. Bolded text indicates significant differences ($p \leq .05$) among groups within a stratification level (i.e., females vs. males).

Abbreviations: AGP, agar gel precipitin; AGID, agar gel immunodiffusion; CA, card agglutination; cELISA, competitive ELISA; CF, complement fixation; ELISA, enzyme-linked immunosorbent assay; HI, hemagglutination inhibition; IFA, immunofluorescence assay; LA, latex agglutination; MAT, modified agglutination test; NT, not tested; PCR, polymerase chain reaction; SVN, serum virus neutralization; VI, virus isolation.

tested positive for exposure to orf virus more often than adults ($p = .03$). There were differences among regions in the infection/exposure of *Anaplasma* spp. ($p < .001$), BRSV ($p < .001$), *B. ovis* ($p < .001$), *Chlamydia* spp. ($p < .001$), *Leptospira* spp. ($p = .001$), *M. ovipneumoniae* (PCR; $p = .04$), orf virus ($p < .001$), PI-3 (HI; $p < .001$), *T. gondii* ($p = .02$), BTV (cELISA; $p = .01$), EHDV (SVN; $p = .01$), *Orbivirus* spp. ($p = .001$), and *B. trehalosi* beta-hemolytic ($p < .001$).

All pathogens showed temporal changes in prevalence over the 36-year study period (Appendix S1). Some common pathogens (i.e., *M. ovipneumoniae* [cELISA] and *Chlamydia* spp.), were consistently present across the entire study period and all regions. BRSV, PI-3, and orf virus were more variable, sometimes ranging from 0% to 100% in the span of 1 year. *Brucella ovis* was only detected in the 1990s. *M. ovipneumoniae* (PCR) and BRSV showed increasing prevalence over time.

3.2 | Population performance and meteorologic variables

Population size was positively associated ($\beta = 2.9$, CI = 1.7–4.1) with adult survival (Appendix S2). There was no relationship between population size and lamb survival ($\beta = 0.5$, CI = -0.4 – 1.4). Temperature and precipitation were not independently associated with adult or lamb survival (Appendix S2). Despite this lack of association, meteorologic covariates were included in population-level regression model building to test if they improved model fit because they have been previously established as important factors in bighorn sheep survival.

Population size ($\beta = 2.74$, CI = 2.32–3.16) and temperature ($\beta = 0.52$, CI = 0.35–0.69) were positively associated with year (Appendix S2). Annual ($\beta = -0.50$, CI = -0.74 to -0.26), summer ($\beta = -0.67$, CI = -0.98 to -0.37), and winter ($\beta = -0.50$, CI = -0.76 to -0.24) precipitation were negatively associated with year (Appendix S2).

3.3 | Pathogen prevalence and lamb survival

The following pathogens had enough datapoints to be included as regression model covariates to evaluate the impact of pathogens on adult and lamb survival: *Anaplasma* spp., BTV (cELISA), BRSV, *Chlamydia* spp., orf virus, *Leptospira* spp., *M. ovipneumoniae* (PCR, cELISA), *Orbivirus* spp., and PI-3 (HI).

Mycoplasma ovipneumoniae exposure prevalence (cELISA) was negatively associated with current year lamb survival (no lag; $\beta = -1.29$, CI = -2.48 to -0.07;

Appendix S3). This relationship persisted with the addition of population size ($\beta = -1.56$, CI = -2.76 to -0.39 ; Appendix S4), population size and temperature ($\beta = -1.45$, CI = -2.66 to -0.29 ; Appendix S5), population size and annual precipitation ($\beta = -1.62$, CI = -2.84 to -0.38 ; Appendix S6), population size and summer precipitation ($\beta = -1.56$, CI = -2.76 to -0.37 ; Appendix S7), and population size and winter precipitation ($\beta = -1.66$, CI = -2.92 to -0.42 ; Appendix S8). In these models, population size and meteorologic covariates were not associated with lamb survival. *Mycoplasma ovipneumoniae* exposure prevalence was also negatively associated with current year lamb survival ($\beta = -1.47$, CI = -2.96 to -0.04) in the model including BRSV, although BRSV was not a significant predictor (Appendix S3).

Lamb survival was not associated with other pathogens, population size, temperature, or precipitation (Appendices S3–S8). The inclusion of population size improved model fit in just 0.74% ($n = 1/136$) of models. LOO IC standard errors overlapped for all other models.

3.4 | Pathogen prevalence and adult survival

Orf virus prevalence was positively associated with adult survival in the subsequent year (-1 -year lag; $\beta = 1.6$, CI = 0.4 – 2.9 ; Appendix S3). This relationship persisted with the addition of population size ($\beta = 1.2$, CI = 0.0 – 2.5 ; Appendix S4), and population size with annual precipitation ($\beta = 1.2$, CI = 0.0 – 2.4 ; Appendix S6).

PI-3 was associated with increases in current year survival once other covariates were accounted for: population size ($\beta = 1.8$, CI = 0.2 – 4.1 ; Appendix S4), population size and temperature ($\beta = 1.8$, CI = 0.3 – 3.9 ; Appendix S5), population size and annual precipitation ($\beta = 1.8$, CI = 0.2 – 4.0 ; Appendix S6), population size and summer precipitation ($\beta = 1.5$, CI = 0.1 – 3.7 ; Appendix S7), and population size and winter precipitation ($\beta = 1.8$, CI = 1.2 – 4.1 ; Appendix S8). In these models, population size was positively associated with adult survival rates, but meteorologic covariates were not.

No other pathogens were predictors of adult survival, regardless of lag time or covariates (Appendices S3–S8). Population size was associated with higher adult survival rates in 32.4% ($n = 46/142$) of models across all pathogens (Appendices S4–S8). Higher summer temperatures were associated with lower adult survival rates in 7.1% ($n = 2/28$) of models (Appendix S5), and higher summer precipitation was associated with higher adult survival rates in 25.0% ($n = 7/28$) models (Appendix S7). Annual and winter precipitation were not significantly associated with adult survival in any models (Appendices S6 and

S8). Including population size and summer temperature improved model fit in 28.6% ($n = 8/28$) of models. LOO IC standard errors overlapped for all other models. The inconsistency in the significance of population size and meteorologic variables was likely because each model contained a different dataset; samples were not tested for every pathogen and adult survival rates were not available for every year-region unit.

3.5 | Individual-level risk factors associated with pathogen exposure or infection

The following pathogens had enough datapoints to be included as regression model covariates to evaluate infection/exposure risk factors at the individual level: *Anaplasma* spp., BTV (cELISA), BRSV, *B. ovis*, *Chlamydia* spp., *Leptospira* spp., *M. ovipneumoniae* (PCR and cELISA), *Orbivirus* spp., orf virus, PI-3 (HI), and *T. gondii* (Figure 2 and Appendix S9).

Odds of exposure to *M. ovipneumoniae* (cELISA) were higher in the northern Santa Rosa Mountains and Vallecito Mountains (OR = 3.7 and 2.9, respectively), compared with the San Jacinto Mountains. Age, sex, and region were not significant predictors of active infection with *M. ovipneumoniae* (PCR).

BRSV and PI-3 had similar distributions, with most regions having higher odds of exposure compared with the San Jacinto Mountains (OR = 4.4–44.7; Figure 2). Females were more likely to be exposed to BRSV than males (OR = 2.1).

Orbivirus spp. had lower odds of exposure in females (OR = 0.3) and in the northern half of the range (Figure 2). The northern San Ysidro Mountains, in the middle of the range, had higher odds BTV exposure (OR = 8.2). The discrepancies in risk between BTV and *Orbivirus* spp. are likely due to *Orbivirus* spp. models including animals exposed to EHDV and/or BTV, and differences in the spatial/temporal testing for each pathogen (Table 1 and Appendix S1).

Brucella ovis exposure was greater in the southern half of the range, including the northern (OR = 21.0) and southern San Ysidro Mountains (OR = 10.1), and Carrizo Canyon (OR = 9.4). Positive samples were limited to 1990–1997, with 17 of 24 positive results occurring in 1992 in the northern San Ysidro Mountains and Carrizo Canyon. Carrizo Canyon also had higher odds of exposure to orf virus (OR = 3.1).

Anaplasma spp. and *Chlamydia* spp. had patchy geographic distributions. The risk of exposure to *Anaplasma* spp. was lower in the northern Santa Rosa Mountains (OR = 0.1) and higher in the northern San Ysidro Mountains (OR = 8.7). Odds of exposure to *Chlamydia* spp.




FIGURE 2 Legend on next page.




FIGURE 3 A weighted, undirected network of pathogens that co-occurred together within an individual bighorn sheep. Size of nodes (circles) is relative to the number of other pathogens that node is connected too (larger nodes are linked to more pathogens; range 7–15). Width of edges (lines) between nodes is relative to the proportion of bighorn sheep samples which were positive for a pair of pathogens, given that both pathogens were tested for. The color shade of the nodes corresponds to the weighted degree, calculated as the sum of the edges leading into a node, so that darker nodes are linked to a larger number of other nodes and also tested positive in a larger proportion of samples. Abbreviations: *B. ovis*, *Brucella ovis*; BH, beta-hemolytic; cELISA, competitive enzyme-linked immunosorbent assay; *M. ovi*, *Mycoplasma ovipneumoniae*; NH, nonhemolytic; PCR, polymerase chain reaction; *T. gondii*, *Toxoplasma gondii*

were higher in the central Santa Rosa Mountains (OR = 6.3) but lower in the bordering northern and southern Santa Rosa Mountains (OR = 0.1 in both).

Exposure to *Leptospira* spp. was also scattered, with higher exposure odds in the southern Santa Rosa Mountains (OR = 5.6) and southern San Ysidro Mountains (OR = 7.3).

Odds of exposure to *T. gondii* were lower for females (OR = 0.1) and higher for animals in the northern Santa Rosa Mountains (OR = 15.3).

3.6 | Pathogen co-occurrence network

All pathogens were connected to several other pathogens in the network (mean = 13, median = 14, and range = 7–15 out of 15 possible connections) and there was minimal

clustering, as evidenced by low modularity (0.07), low betweenness centrality (0.01), and high density (0.86) of the network (Figure 3 and Appendix S10).

4 | DISCUSSION

4.1 | Pathogen prevalence and bighorn sheep survival

Bighorn sheep infected with *M. ovipneumoniae* can die, clear the infection, or become carriers that persistently or intermittently shed bacteria (Cassirer et al., 2013). After the initial epidemic, adult *M. ovipneumoniae* PCR prevalence tends to be low because most animals stop shedding after <1 year (Plowright et al., 2017). Conversely, *M. ovipneumoniae* seropositivity (cELISA) is a measure of

FIGURE 2 Forest plots demonstrating the relationship between pathogen status (positive, negative) and Peninsular bighorn sheep age class (lamb/yearling, adult), sex (female, male), and recovery region (categorical, $n = 9$). Reference categories were adults (for age), males (for sex), and the San Jacinto Mountains (for recovery region). Numbers and white circle represent the log odds of testing positive for a pathogen, relative to a reference category, using Bayesian, multilevel, logistic regression models. Log odds <0 (blue) indicate a lower risk of testing positive for a given pathogen, and log odds >0 (red) indicate a higher risk of testing positive for a given pathogen, relative to the reference category. Thick bars represent the 80% credible interval (CI), and thin bars represented the 95% CI. A covariate was a significant predictor of pathogen status if the 95% CI did not cross 0. ELISA, enzyme-linked immunosorbent assay; PCR, polymerase chain reaction

past disease exposure and could be indicative of the amount of transmission occurring during the spring lambing season in that year. Lamb survival in this study was lower in years with higher *M. ovipneumoniae* exposure prevalence (cELISA). The highest prevalence and odds of *M. ovipneumoniae* exposure were in the northern Santa Rosa Mountains, which also consistently had the lowest lamb survival across the study period (Table 1 and Appendix S1).

Lambs in the Peninsular Ranges have been observed with clinical signs consistent with pneumonia in all regions, and almost all of the uncollared bighorn mortalities attributed to disease from 2002 to 2019 have been the result of bacterial pneumonia in lambs (Colby & Botta, 2019). Most of these mortalities have been detected in the central Santa Rosa Mountains and northern San Ysidro Mountains, where proximity to urban spaces and areas of heavy recreational use by humans results in decreased predation and increased detection of sick lambs, even though these regions do not consistently have the highest prevalence of respiratory pathogens (Table 1 and Appendix S1).

Increased lamb mortality is associated with the presence of even a few ewes shedding *M. ovipneumoniae* and epidemics of pneumonia affect lamb survival and recruitment to a greater degree than adult survival (Cassirer et al., 2013; Manlove et al., 2014; Manlove et al., 2017; Monello et al., 2001; Plowright et al., 2013). The extinction risk of Peninsular bighorn sheep is inversely related to adult female survival (Rubin, Boyce, & Caswell-Chen, 2002), which was not found to be associated with *M. ovipneumoniae* infection/exposure in this study. However, by reducing lamb recruitment, *M. ovipneumoniae* could both slow population recovery and potentially increase extinction risk by leading to a reduction in adult survival in an aging population.

Poor lamb survival in some regions may be improved by conducting “test and removal” of *M. ovipneumoniae* carrier females, as has been evidenced through other recent studies (Garwood et al., 2020). The relatively linear north–south orientation of these populations may assist in the implementation of this method. However, “test and removal” efforts are costly, requiring the capture and testing of nearly all adult females within a herd, and herds where chronic shedders have been removed are still susceptible to reintroduction of *M. ovipneumoniae*, which could result in epidemics and all age die-offs.

We did not find evidence that orf virus was associated with decreased survival, but it is extremely common and could play a role in individual fitness. Contagious ecthyma (the disease caused by orf virus) is generally self-limiting and resolves within a few months, but can

lead to secondary infections and mortality in young animals or those with comorbidities (Colby & Botta, 2018; Jones et al., 2018; Michelsen & Smith, 2009). Infectious carrier states and reinfections are observed in domestic sheep (Lewis, 1996; Nandi et al., 2011), suggesting the virus may not fade out from herd immunity.

Antibody prevalence of PI-3 was associated with higher adult survival, potentially indicating that age may be associated with cumulative exposure risk. PI-3 generally causes subclinical to mild respiratory signs as a sole agent in domestic sheep, but can predispose animals to fatal secondary bacterial pneumonia, especially from *Pasteurellaceae* spp. (Woolums et al., 2009).

We found a positive relationship between population size and adult survival but could not establish directionality. Larger population sizes may be the result of improving survival rates as the population recovers or there may be a survival benefit to larger groups, such as vigilance against predators.

4.2 | Environmental impacts on survival

We found trends towards higher adult survival rates with lower summer temperatures and higher summer rainfall, once pathogen prevalence and population size were accounted for. Unfortunately, work evaluating a subset of the Peninsular Mountains (overlapping with Regions 3–9) over the same time period as this study (1984–2017) determined that increases in summer temperatures and decreases in precipitation (October_{t-1} to September_t) were associated with widespread declines in perennial vegetation cover, with a stronger magnitude of effect at the lower elevations (<500 m) preferred by P bighorn sheep (Hantson et al., 2021). The increasing temperatures and decreasing precipitation we observed over the course of this study are expected to worsen in the southwestern desert regions as climate change continues (Hess et al., 2008), potentially affecting bighorn sheep through resource limitation and subsequent behavioral adaptations that may alter disease transmission.

While drought and increasing temperatures may drive sheep to aggregated at limited water sources, it may also lead to lower densities and contact rates if sheep are driven to disperse in search of increasingly sparse vegetation (Epps et al., 2004). This has been anecdotally observed recently, with declines in the density of spring vegetation and smaller bighorn sheep nursery groups occurring in the same years as fewer observations of lambs with clinical respiratory disease and lower *M. ovipneumoniae* PCR prevalence (CDFW, unpublished data).

4.3 | Geography is a greater risk factor for pathogen infection/exposure than demographics

The primary risk factor for individual bighorn sheep pathogen infection/exposure was region, which is likely a proxy for local ecological and behavioral factors. The type and number of water sources in a region could influence rates of direct contact between sheep, and contamination of food and/or water with feces and urine could spread pathogens such as *T. gondii* and *Leptospira* spp. (Adler & de la Peña Moctezuma, 2010; Dubey, 2009). Behavioral observations and genetic data have demonstrated a strong matrilineal structure between bighorn sheep herds (Boyce et al., 1999), which could lead to greater disease transmission within herds compared with between herds.

Age was not a significant predictor of an animal's pathogen status, perhaps because lambs and yearlings were categorized together and sampling generally happened in the fall, after the critical window when most pneumonia-associated lamb mortalities occur (Cassirer et al., 2018). Similarly, the relatively low numbers of both lambs/yearlings (11.0%, $n = 81/735$) and males (19.6%, $n = 144/735$) in the dataset may have decreased our power to detect differences among groups.

4.4 | Potential impacts of coinfections and multiple pathogen strains

Bighorn sheep epidemic pneumonia is a complex disease process involving multiple pathogens and non-infectious stressors (Besser et al., 2013). We found that respiratory pathogens were relatively common in Peninsular bighorn sheep across their range, especially *M. ovipneumoniae* and BRSV. Although we found limited evidence for negative population-level effects of pathogens other than *M. ovipneumoniae*, the long-term circulation of multiple pathogens may have a subclinical effect or exacerbate concurrent, nondisease stressors. The co-occurrence network showed that all pathogens co-occurred with numerous other pathogens, and there were no communities that clustered together which could inform future targeted surveillance. This network may have had biases towards detecting highly prevalent pathogens with higher survival rates, long-lasting antibodies, and consistent testing. However, the potential synergism of coinfections implications for individual fitness and long-term population resiliency. For example, lambs that are not feeding well due to painful orf sores around their mouth may be more likely to succumb to pneumonia.

Bighorn sheep do not appear to gain protective cross-immunity against different strains of *M. ovipneumoniae*

after infection (Cassirer et al., 2017), and different *M. ovipneumoniae* strains have been associated with varying levels of morbidity/mortality (Besser et al., 2017). To date, 23 *M. ovipneumoniae* samples from Peninsular bighorn sheep have been genotyped using multilocus sequence typing, and two distinct ovine strains have been identified, possibly representing distinct spillover events. One strain, most closely related to bighorn sheep samples from the nearby Orocopia Mountains, is found throughout all recovery regions (Cassirer et al., 2018). A second strain, most similar to samples from Joshua Tree National Park, was identified in 2020 from sheep in two northern regions (San Jacinto Mountains, central Santa Rosa Mountains; CDFW, unpublished data). Continued monitoring and strain typing will be important to detect the introduction and spread of novel strains which could lead to new outbreaks of pneumonia and all age class mortality.

4.5 | Limitations and opportunities for future surveillance

The lack of associations in this study between most pathogens and bighorn sheep survival may be due to data limitations resulting from the shifting priorities and capabilities of this multi-decade recovery project. Most diagnostic tests measured previous exposure and we do not know the duration of seropositivity for many of these diseases. Our results might differ if we measured clinical disease, active infections, or directly observed lamb survival within the first few months of life. Not measuring lamb survival to a consistent age across years and regions may have masked age-related differences in survival. Using recovery region as the unit of analysis may have masked herd-level differences. More importantly, disease-induced mortality is a multifactorial process that includes variables not included in our models, such as immune function, pathogen virulence, and dynamic behaviors such as contact rates among hosts.

As with many wildlife studies, Peninsular bighorn sheep monitoring suffers from limited funding, staff, and access to rugged and remote habitats. While regular, frequent, and comprehensive molecular surveillance combined with observations of clinical disease is ideal for understanding detailed disease dynamics, it is not always attainable or sustainable. Long-term projects such as this one often need to spread out resources, electing for a lower intensity monitoring plan that can be sustained over many years to detect population-level changes. So, the question becomes, what is the optimal monitoring strategy to detect disease and assess overall population health in a low-density, highly mobile species occurring in remote and difficult to access habitats?

Pairing diagnostic tests that distinguish active from previous infection provides the most information on which diseases could be causing current mortalities versus explain historical population performance. Diseases prioritized for testing should be those that have the greatest potential to negatively impact survival and recruitment, such as the pneumonia-associated pathogens: *M. ovipneumoniae*, BRSV, and PI-3 (WAFWA Wildlife Health Committee, 2016). Although *Pasteurellaceae* spp. have long been implicated as causal organisms in pneumonia, the strength of association is weak and they are common commensal organisms in healthy sheep, so the utility of regular testing is limited (Besser et al., 2013). Highly prevalent pathogens may also be of interest, even if there is no current evidence that they pose a threat to the population. Orf virus is not currently associated with reductions in survival and skin lesions can be detected visually, but it is very common in Peninsular bighorn sheep and active disease may be missed due to the short duration of clinical signs. Continuing to monitor seroprevalence will help us understand how many animals are suffering morbidity or mortality as the population grows.

Throughout most of the Peninsular Ranges, it is not possible to directly observe bighorn sheep on a regular basis due to remote and rugged terrain combined with extreme weather conditions. Continuing to collar a subset of animals for survival monitoring and quick carcass recovery should be a mainstay of bighorn sheep monitoring, especially in areas where directly observing animals is difficult. Collaring young lambs would provide better estimates of lamb survival, an important metric of population health that varies considerably among regions and years, and improve our ability to determine causes of death among juveniles. The large area and hot, dry weather of the Peninsular Ranges make it difficult to retrieve carcasses quickly enough to reliably determine the cause of death, even when animals are radio-collared, limiting our ability to detect disease outbreaks or other population threats. Future monitoring may focus on alternative methods, although they carry detection biases compared with radio-collaring.

As an alternative, remote cameras placed in areas where animals reliably congregate could be used to detect visible signs of morbidity, such as nasal/ocular discharge, postural changes, muscle wasting, lameness, and skin lesions (Brewster et al., 2017; Brown & Elmer, 2019; Carricondo-Sanchez et al., 2017; Muneza et al., 2019). Advancements in machine learning could aid in image processing and automated detection of species, posture, and potentially even disease lesions (Tuia et al., 2022). “Physiologgers,” implantable transmitters that record physiologic variables such as heart rate, body temperature, and respiratory rhythms, are a rapidly developing technology that could one day be utilized to monitor

bighorn sheep for signs of stress and sickness that may not be detected by visual observations alone (Hawkes et al., 2021). If these methods detected signs of morbidity or a decrease in apparent bighorn abundance, a more detailed investigation could be triggered. Combining these tools with survival and movement data from radio-collared animals would provide more comprehensive insights into emerging health threats, bighorn sheep responses to environmental changes, and the effectiveness of management efforts.

5 | CONCLUSIONS

Peninsular bighorn sheep are recovering from critically small population sizes in an ecosystem which includes natural and urban habitats at environmental extremes. This study demonstrates that *M. ovipneumoniae* is associated with lower lamb survival and identified regions with elevated risk of pathogen infection/exposure to guide future surveillance. Changes in bighorn sheep behavior and distribution in response to climate change and anthropogenic development may play a role in the maintenance or amplification of disease, especially where bighorn sheep congregate, such as around limited water sources and on lambing grounds. Long-term, consistent, range-wide pathogen testing and population surveys will be critical to advance our understanding of pathogen transmission and the role of disease in Peninsular bighorn sheep population recovery. Consideration of environmental factors and incorporation of novel technologies will also be important to adjust management strategies in the face of climate change and dynamic disease risks.

AUTHOR CONTRIBUTIONS

Jessica N. Sanchez: conceptualization, methodology, software, validation, formal analysis, investigation, resources, data curation, writing – original draft, writing – review and editing, visualization, supervision, project administration, and funding acquisition. **Brandon A. Munk:** conceptualization, investigation, resources, data curation, writing – review and editing, supervision, project administration, and funding acquisition. **Janene Colby:** conceptualization, investigation, resources, data curation, writing – review and editing, and project administration. **Steve G. Torres:** conceptualization, investigation, resources, data curation, writing – review and editing, supervision, project administration, and funding acquisition. **Ben J. Gonzalez:** conceptualization, investigation, resources, data curation, writing – review and editing, supervision, project administration, and funding acquisition. **James R. DeForge:** investigation, resources, data curation, writing – review and editing,

supervision, project administration, and funding acquisition. **Aimee J. Byard**: investigation, resources, data curation, writing – review and editing, project administration, and funding acquisition. **Lora Konde**: investigation, resources, data curation, writing – review and editing, and project administration. **Nicholas Shirkey**: investigation, resources, data curation, writing – review and editing, and project administration. **Pranav S. Pandit**: conceptualization, methodology, software, and writing – review and editing. **Randy A. Botta**: investigation, resources, writing – review and editing, supervision, project administration, and funding acquisition. **Annette Roug**: conceptualization, data curation, and writing – review and editing. **Michael H. Ziccardi**: methodology, and writing – review and editing. **Christine K. Johnson**: conceptualization, methodology, writing – review and editing, supervision, project administration, and funding acquisition.

ACKNOWLEDGMENTS

Numerous colleagues provided advice, expertise, and support throughout this study: Walter M. Boyce, David A. Jessup, Danielle J. Harvey, Daniel J. Tancredi, Gabriele Maier, E. Robert Atwill, Woutrina A. Smith, Kevin Keel, Beatriz Martínez López, Peregrine L. Wolff, Clinton W. Epps, T. Winston Vickers, Ashley E. Hill, Gabriel A. Reyes, Sarah T. Abusaa, Laura H. Backus, Peter J. Sebastian, Christine T. Chang, Joe Smith, and Shane M. Sanchez.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

DATA AVAILABILITY STATEMENT

All data used in this article are presented in Appendices S11–S13.


ORCID

Jessica N. Sanchez  <https://orcid.org/0000-0002-7689-0324>

Brandon A. Munk  <https://orcid.org/0000-0001-9338-2236>

James R. DeForge  <https://orcid.org/0000-0002-3873-9711>

Aimee J. Byard  <https://orcid.org/0000-0003-0841-8551>

Nicholas J. Shirkey  <https://orcid.org/0000-0002-1919-9596>

Pranav S. Pandit  <https://orcid.org/0000-0001-7649-0649>

Randy A. Botta  <https://orcid.org/0000-0002-7903-7492>

Annette Roug  <https://orcid.org/0000-0002-6844-4076>

Michael H. Ziccardi  <https://orcid.org/0000-0002-5430-2499>

2499

Christine K. Johnson  <https://orcid.org/0000-0001-6673-8743>

REFERENCES

- Abatzoglou, J. T. (2013). Development of gridded surface meteorological data for ecological applications and modelling. *International Journal of Climatology*, 33(1), 121–131. <https://doi.org/10.1002/joc.3413>
- Adler, B., & de la Peña Moctezuma, A. (2010). Leptospira and leptospirosis. *Veterinary Microbiology*, 140(3), 287–296. <https://doi.org/10.1016/j.vetmic.2009.03.012>
- Asghar, M., Hasselquist, D., Hansson, B., Zehindjiev, P., Westerdahl, H., & Bensch, S. (2015). Hidden costs of infection: Chronic malaria accelerates telomere degradation and senescence in wild birds. *Science*, 347(6220), 436–438. <https://doi.org/10.1126/science.1261121>
- Bastian, M., Heymann, S., & Jacomy, M. (2009). Gephi: An open source software for exploring and manipulating networks. Vol. 3, No. 1.
- Besser, T. E., Cassirer, E. F., Highland, M. A., Wolff, P., Justice-Allen, A., Mansfield, K., Davis, M. A., & Foreyt, W. (2013). Bighorn sheep pneumonia: Sorting out the cause of a polymicrobial disease. *Preventive Veterinary Medicine*, 109(3–4), 185. <https://doi.org/10.1016/j.prevetmed.2013.02.020>
- Besser, T. E., Cassirer, E. F., Potter, K. A., & Foreyt, W. J. (2017). Exposure of bighorn sheep to domestic goats colonized with *Mycoplasma ovipneumoniae* induces sub-lethal pneumonia. *PLoS One*, 12(6), e0178707. <https://doi.org/10.1371/journal.pone.0178707>
- Besser, T. E., Cassirer, E. F., Potter, K. A., Lahmers, K., Oaks, J. L., Shanthalingam, S., Srikumaran, S., & Foreyt, W. J. (2014). Epizootic pneumonia of bighorn sheep following experimental exposure to *Mycoplasma ovipneumoniae*. *PLoS One*, 9(10), e110039. <https://doi.org/10.1371/journal.pone.0110039>
- Besser, T. E., Cassirer, E. F., Potter, K. A., VanderSchalie, J., Fischer, A., Knowles, D. P., Herndon, D. R., Rurangirwa, F. R., Weiser, G. C., & Srikumaran, S. (2008). Association of *Mycoplasma ovipneumoniae* infection with population-limiting respiratory disease in free-ranging Rocky Mountain bighorn sheep (*Ovis canadensis canadensis*). *Journal of Clinical Microbiology*, 46(2), 423–430.
- Besser, T. E., Highland, M. A., Baker, K., Cassirer, E. F., Anderson, N. J., Ramsey, J. M., Mansfield, K., Bruning, D. L., Wolff, P., Smith, J. B., & Jenks, J. A. (2012). Causes of pneumonia epizootics among bighorn sheep, western United States, 2008–2010. *Emerging Infectious Diseases*, 18(3), 406–414. <https://doi.org/10.3201/eid1803.111554>
- Bighorn Institute. (2018). *Bighorn Institute 2018 year-end report: Investigations of Peninsular bighorn sheep in the Santa Rosa Mountains and San Jacinto Mountains of California* (p. 15). Bighorn Institute.
- Boyce, W. M., Ramey, R., Rodwell, T., Rubin, E., & Singer, R. (1999). Population subdivision among desert bighorn sheep (*Ovis canadensis*) ewes revealed by mitochondrial DNA analysis. *Molecular Ecology*, 8(1), 99–106. <https://doi.org/10.1046/j.1365-294X.1999.00536.x>
- Brewster, K., Henke, S. E., Hilton, C., & Ortega-S, A., Jr. (2017). Use of remote cameras to monitor the potential prevalence

- of sarcoptic mange in southern Texas, USA. *Journal of Wildlife Diseases*, 53(2), 377–381. <https://doi.org/10.7589/2016-08-180>
- Brown, W. E., & Elmer, J. (2019). Remote detection and monitoring methods for Tasmanian devils. In C. J. Hogg, S. Fox, D. Pemberton, & K. Belov (Eds.), *Saving the Tasmanian devil: Recovery through science-based management* (pp. 139–155). CSIRO Publishing. <https://doi.org/10.1071/9781486307197>
- Buchalski, M. R., Navarro, A. Y., Boyce, W. M., Winston Vickers, T., Tobler, M. W., Nordstrom, L. A., Garcia, J. A., Gille, D. A., Penedo, M. C. T., Ryder, O. A., & Ernest, H. B. (2015). Genetic population structure of Peninsular bighorn sheep (*Ovis canadensis nelsoni*) indicates substantial gene flow across US–Mexico border. *Biological Conservation*, 184, 218–228. <https://doi.org/10.1016/j.biocon.2015.01.006>
- Buchalski, M. R., Sacks, B. N., Gille, D. A., Penedo, M. C. T., Ernest, H. B., Morrison, S. A., & Boyce, W. M. (2016). Phylogeographic and population genetic structure of bighorn sheep (*Ovis canadensis*) in North American deserts. *Journal of Mammalogy*, 97(3), 823–838. <https://doi.org/10.1093/jmammal/gyw011>
- Bürkner, P.-C. (2017). brms: An R package for Bayesian multilevel models using Stan. *Journal of Statistical Software*, 80(1), 1–28. <https://doi.org/10.18637/jss.v080.i01>
- Bürkner, P.-C. (2018). Advanced Bayesian multilevel modeling with the R package brms. *The R Journal*, 10(1), 395–411.
- Butts, C. T. (2008). Social network analysis with sna. *Journal of Statistical Software*, 24(6), 1–51.
- Carricondo-Sanchez, D., Odden, M., Linnell, J. D. C., & Odden, J. (2017). The range of the mange: Spatiotemporal patterns of sarcoptic mange in red foxes (*Vulpes vulpes*) as revealed by camera trapping. *PLoS One*, 12(4), e0176200. <https://doi.org/10.1371/journal.pone.0176200>
- Cassirer, E. F., Manlove, K. R., Almberg, E. S., Kamath, P. L., Cox, M., Wolff, P., Roug, A., Shannon, J., Robinson, R., Harris, R. B., Gonzales, B. J., Plowright, R. K., Hudson, P. J., Cross, P. C., Dobson, A., & Besser, T. E. (2018). Pneumonia in bighorn sheep: Risk and resilience. *Journal of Wildlife Management*, 82(1), 32–45. <https://doi.org/10.1002/jwmg.21309>
- Cassirer, E. F., Manlove, K. R., Plowright, R. K., & Besser, T. E. (2017). Evidence for strain-specific immunity to pneumonia in bighorn sheep. *The Journal of Wildlife Management*, 81(1), 133–143. <https://doi.org/10.1002/jwmg.21172>
- Cassirer, E. F., Plowright, R. K., Manlove, K. R., Cross, P. C., Dobson, A. P., Potter, K. A., & Hudson, P. J. (2013). Spatiotemporal dynamics of pneumonia in bighorn sheep. *Journal of Animal Ecology*, 82(3), 518–528. <https://doi.org/10.1111/1365-2656.12031>
- Cassirer, E. F., Rudolph, K. M., Fowler, P., Coggins, V. L., Hunter, D. L., & Miller, M. W. (2001). Evaluation of ewe vaccination as a tool for increasing bighorn lamb survival following pasteurellosis epizootics. *Journal of Wildlife Diseases*, 37(1), 49–57. <https://doi.org/10.7589/0090-3558-37.1.49>
- Clauset, A., Newman, M. E. J., & Moore, C. (2004). Finding community structure in very large networks. *Physical Review E*, 70(6), 066111. <https://doi.org/10.1103/PhysRevE.70.066111>
- Colby, J., & Botta, R. (2013). California Department of Fish and Wildlife, Peninsular Bighorn sheep 2013 annual report, p. 16. California Department of Fish and Wildlife.
- Colby, J., & Botta, R. (2017). California Department of Fish and Wildlife, Peninsular Bighorn sheep 2016–2017 annual report, p. 29. California Department of Fish and Wildlife.
- Colby, J., & Botta, R. (2018). California Department of Fish and Wildlife Peninsular Bighorn sheep 2017–18 annual report, p. 25. California Department of Fish and Wildlife.
- Colby, J., & Botta, R. (2019). California Department of Fish and Wildlife Peninsular Bighorn Sheep 2018–19 Annual Report and Recovery Program Review 1992–2019, p. 24. California Department of Fish and Wildlife.
- Dassanayake, R. P., Shanthalingam, S., Herndon, C. N., Subramaniam, R., Lawrence, P. K., Bavananthasivam, J., Cassirer, E. F., Haldorson, G. J., Foreyt, W. J., Rurangirwa, F. R., Knowles, D. P., Besser, T. E., & Srikumaran, S. (2010). *Mycoplasma ovipneumoniae* can predispose bighorn sheep to fatal *Mannheimia haemolytica* pneumonia. *Veterinary Microbiology*, 145(3–4), 354–359. <https://doi.org/10.1016/j.vetmic.2010.04.011>
- DeForge, J. R., Jessup, D. A., Jenner, C. W., & Scott, J. E. (1982). Disease investigations into high lamb mortality of desert bighorn in the Santa Rosa Mountains, California. *Desert Bighorn Council Transactions*, 26, 76–81.
- Dubey, J. P. (2009). Toxoplasmosis in sheep—The last 20 years. *Veterinary Parasitology*, 163(1), 1–14. <https://doi.org/10.1016/j.vetpar.2009.02.026>
- Epps, C. W., McCullough, D. R., Wehausen, J. D., Bleich, V. C., & Rechel, J. L. (2004). Effects of climate change on population persistence of desert-dwelling mountain sheep in California. *Conservation Biology*, 18(1), 102–113. <https://doi.org/10.1111/j.1523-1739.2004.00023.x>
- Foreyt, W. J., & Jessup, D. A. (1982). Fatal pneumonia of bighorn sheep following association with domestic sheep. *Journal of Wildlife Diseases*, 18(2), 163–168. <https://doi.org/10.7589/0090-3558-18.2.163>
- Garwood, T. J., Lehman, C. P., Walsh, D. P., Cassirer, E. F., Besser, T. E., & Jenks, J. A. (2020). Removal of chronic *Mycoplasma ovipneumoniae* carrier ewes eliminates pneumonia in a bighorn sheep population. *Ecology and Evolution*, 10(7), 3491–3502. <https://doi.org/10.1002/ece3.6146>
- Ghosh, J., Li, Y., & Mitra, R. (2018). On the use of Cauchy prior distributions for Bayesian logistic regression. *Bayesian Analysis*, 13(2), 359–383. <https://doi.org/10.1214/17-BA1051>
- Hantson, S., Huxman, T. E., Kimball, S., Randerson, J. T., & Goulden, M. L. (2021). Warming as a driver of vegetation loss in the Sonoran Desert of California. *Journal of Geophysical Research: Biogeosciences*, 126(6), e2020JG005942. <https://doi.org/10.1029/2020JG005942>
- Hawkes, L. A., Fahlman, A., & Sato, K. (2021). Introduction to the theme issue: Measuring physiology in free-living animals. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 376(1830), 20200210. <https://doi.org/10.1098/rstb.2020.0210>
- Hess, J. J., Malilay, J. N., & Parkinson, A. J. (2008). Climate change: The importance of place. *American Journal of Preventive Medicine*, 35(5), 468–478. <https://doi.org/10.1016/j.amepre.2008.08.024>

- Jamieson, A. M., Pasman, L., Yu, S., Gamradt, P., Homer, R. J., Decker, T., & Medzhitov, R. (2013). Role of tissue protection in lethal respiratory viral-bacterial coinfection. *Science*, 340(6137), 1230–1234. <https://doi.org/10.1126/science.1233632>
- Johnson, M. (2022). climateR. R package version 0.1.0. <https://github.com/mikejohnson51/climateR>
- Jones, M. E., Gasper, D. J., & Mitchell, E. (2018). Bovidae, Antilocapridae, Giraffidae, Tragulidae, Hippopotamidae. In K. Terio, D. McAloose, & J. St. Leger (Eds.), *Pathology of wildlife and zoo animals* (1st ed., pp. 117–147). Elsevier.
- Kubinec, R. (2020). Ordered beta regression: A parsimonious, well-fitting model for survey sliders and visual analog scales. SocArXiv.
- Lewis, C. (1996). Update on orf. *In Practice*, 18(8), 376–381. <https://doi.org/10.1136/inpract.18.8.376>
- Lund, J., Medellin-Azuara, J., Durand, J., & Stone, K. (2018). Lessons from California's 2012–2016 drought. *Journal of Water Resources Planning and Management*, 144(10), 04018067.
- Manlove, K. R., Cassirer, E. F., Cross, P. C., Plowright, R. K., & Hudson, P. J. (2014). Costs and benefits of group living with disease: A case study of pneumonia in bighorn lambs (*Ovis canadensis*). *Proceedings of the Royal Society B: Biological Sciences*, 281(1797), 20142331. <https://doi.org/10.1098/rspb.2014.2331>
- Manlove, K. R., Cassirer, E. F., Plowright, R. K., Cross, P. C., & Hudson, P. J. (2017). Contact and contagion: Probability of transmission given contact varies with demographic state in bighorn sheep. *Journal of Animal Ecology*, 86(4), 908–920. <https://doi.org/10.1111/1365-2656.12664>
- Michelsen, P. G. P., & Smith, B. P. (2009). Contagious ecthyma (sore mouth, orf, contagious pustular dermatitis, scabby mouth). In B. P. Smith (Ed.), *Large animal internal medicine* (4th ed., pp. 780–790). Elsevier Health Sciences.
- Monello, R. J., Murray, D. L., & Cassirer, E. F. (2001). Ecological correlates of pneumonia epizootics in bighorn sheep herds. *Canadian Journal of Zoology*, 79(8), 1423–1432. <https://doi.org/10.1139/z01-103>
- Muneza, A. B., Ortiz-Calo, W., Packer, C., Cusack, J. J., Jones, T., Palmer, M. S., Swanson, A., Kosmala, M., Dickman, A. J., & Macdonald, D. W. (2019). Quantifying the severity of giraffe skin disease via photogrammetry analysis of camera trap data. *Journal of Wildlife Diseases*, 55(4), 770–781. <https://doi.org/10.7589/2018-06-149>
- Nandi, S., De, U. K., & Chowdhury, S. (2011). Current status of contagious ecthyma or orf disease in goat and sheep—A global perspective. *Small Ruminant Research*, 96(2), 73–82. <https://doi.org/10.1016/j.smallrumres.2010.11.018>
- Nolen, R. S. (2010). Severe pneumonia outbreak kills bighorn sheep. *Journal of the American Veterinary Medical Association*, 236(9), 936.
- Ostermann, S. D., Deforge, J. R., & Edge, W. D. (2001). Captive breeding and reintroduction evaluation criteria: A case study of Peninsular bighorn sheep. *Conservation Biology*, 15(3), 749–760. <https://doi.org/10.1046/j.1523-1739.2001.015003749.x>
- Plowright, R. K., Manlove, K., Cassirer, E. F., Cross, P. C., Besser, T. E., & Hudson, P. J. (2013). Use of exposure history to identify patterns of immunity to pneumonia in bighorn sheep (*Ovis canadensis*). *PLoS One*, 8(4), e61919. <https://doi.org/10.1371/journal.pone.0061919>
- Plowright, R. K., Manlove, K. R., Besser, T. E., Páez, D. J., Andrews, K. R., Matthews, P. E., Waits, L. P., Hudson, P. J., & Cassirer, E. F. (2017). Age-specific infectious period shapes dynamics of pneumonia in bighorn sheep. *Ecology Letters*, 20(10), 1325–1336. <https://doi.org/10.1111/ele.12829>
- R Core Team. (2021). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. <https://www.R-project.org/>
- Raghavan, B., Erickson, K., Kugadas, A., Batra, S. A., Call, D. R., Davis, M. A., Foreyt, W. J., & Srikumaran, S. (2016). Role of carriers in the transmission of pneumonia in bighorn sheep (*Ovis canadensis*). *Biology Open*, 5(6), 745–755. <https://doi.org/10.1242/bio.018234>
- Rubin, E. S., Boyce, W. M., & Bleich, V. C. (2000). Reproductive strategies of desert bighorn sheep. *Journal of Mammalogy*, 81(3), 769–786. [https://doi.org/10.1644/1545-1542\(2000\)081<0769:RSODBS>2.3.CO;2](https://doi.org/10.1644/1545-1542(2000)081<0769:RSODBS>2.3.CO;2)
- Rubin, E. S., Boyce, W. M., & Caswell-Chen, E. P. (2002). Modeling demographic processes in an endangered population of bighorn sheep. *The Journal of Wildlife Management*, 66(3), 796–810. <https://doi.org/10.2307/3803144>
- Rubin, E. S., Boyce, W. M., Jorgensen, M. C., Torres, S. G., Hayes, C. L., O'Brien, C. S., & Jessup, D. A. (1998). Distribution and abundance of bighorn sheep in the Peninsular ranges, California. *Wildlife Society Bulletin*, 26(3), 539–551.
- Rubin, E. S., Boyce, W. M., Stermer, C. J., & Torres, S. G. (2002). Bighorn sheep habitat use and selection near an urban environment. *Biological Conservation*, 104(2), 251–263. [https://doi.org/10.1016/S0006-3207\(01\)00171-9](https://doi.org/10.1016/S0006-3207(01)00171-9)
- Sells, S. N., Mitchell, M. S., Nowak, J. J., Lukacs, P. M., Anderson, N. J., Ramsey, J. M., Gude, J. A., & Krausman, P. R. (2015). Modeling risk of pneumonia epizootics in bighorn sheep. *The Journal of Wildlife Management*, 79(2), 195–210. <https://doi.org/10.1002/jwmg.824>
- Tuia, D., Kellenberger, B., Beery, S., Costelloe, B. R., Zuffi, S., Risse, B., Mathis, A., Mathis, M. W., van Langevelde, F., Burghardt, T., Kays, R., Klinck, H., Wikelski, M., Couzin, I. D., van Horn, G., Crofoot, M. C., Stewart, C. V., & Berger-Wolf, T. (2022). Perspectives in machine learning for wildlife conservation. *Nature Communications*, 13(1), 792. <https://doi.org/10.1038/s41467-022-27980-y>
- Turner, J. C., Douglas, C. L., Hallum, C. R., Krausman, P. R., & Ramey, R. R. (2004). Determination of critical habitat for the endangered Nelson's bighorn sheep in southern California. *Wildlife Society Bulletin*, 32(2), 427–448. [https://doi.org/10.2193/0091-7648\(2004\)32\[427:DOCHFT\]2.0.CO;2](https://doi.org/10.2193/0091-7648(2004)32[427:DOCHFT]2.0.CO;2)
- US Fish and Wildlife Service. (2000). *Recovery plan for bighorn sheep in the Peninsular ranges, California* (p. 251). US Fish and Wildlife Service.
- Vehtari, A., Gabry, J., Magnusson, M., Yao, Y., Bürkner, P., Paananen, T., & Gelman, A. (2020). loo: Efficient leave-one-out cross-validation and WAIC for Bayesian models (2.3.1) [R package].
- WAFWA Wildlife Health Committee. (2016). *Bighorn sheep herd health monitoring recommendations*. Western Association of Fish and Wildlife Agencies.
- Ward, A. C. S., Hunter, D. L., Rudolph, K. M., DeLong, W. J., Bulgin, J. M., Cowan, L. M., McNeil, H. J., & Miller, M. W. (1999). Immunologic responses of domestic and bighorn sheep to a multi-valent *Pasteurella haemolytica* vaccine. *Journal of Wildlife Diseases*, 35(2), 285–296. <https://doi.org/10.7589/0090-3558-35.2.285>

- Wobeser, G. A. (1994). Disease and epizootiology — Basic principles. In: G. A. Wobeser (Ed.), *Investigation and Management of Disease in Wild Animals* (1st ed., pp. 3-12). Springer, Boston, MA
- Woolums, A. R., Ames, T. R., & Baker, J. C. (2009). The bronchopneumonias (respiratory disease complex of cattle, sheep, and goats). In B. P. Smith (Ed.), *Large animal internal medicine* (4th ed., pp. 602–643). Elsevier Health Sciences.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Sanchez, J. N., Munk, B. A., Colby, J., Torres, S. G., Gonzales, B. J., DeForge, J. R., Byard, A. J., Konde, L., Shirkey, N. J., Pandit, P. S., Botta, R. A., Roug, A., Ziccardi, M. H., & Johnson, C. K. (2022). Pathogen surveillance and epidemiology in endangered peninsular bighorn sheep (*Ovis canadensis nelsoni*). *Conservation Science and Practice*, 4(11), e12820. <https://doi.org/10.1111/csp2.12820>