

Blubber thickness in right whales *Eubalaena glacialis* and *Eubalaena australis* related with reproduction, life history status and prey abundance

Running head: Blubber thickness of right whales

Carolyn A. Miller<sup>1,\*</sup>, Desray Reeb<sup>2</sup>, Peter B. Best<sup>2</sup>, Amy R. Knowlton<sup>3</sup>, Moira W. Brown<sup>3</sup>,  
Michael J. Moore<sup>1</sup>

<sup>1</sup>Department of Biology, Woods Hole Oceanographic Institution, Woods Hole, MA 02543  
U.S.A.

<sup>2</sup>Mammal Research Institute, University of Pretoria, C/o Iziko South African Museum, P.O. Box  
61, Cape Town, South Africa

<sup>3</sup>New England Aquarium, Central Wharf, Boston, MA 02110 U.S.A.

\*cmiller@whoi.edu

## ABSTRACT

The high variability in reproductive performance of North Atlantic right whales, *Eubalaena glacialis*, compared to southern right whales, *Eubalaena australis*, may reflect differences in lipid reserves. Amplitude-mode ultrasound was used to measure the thickness of right whale integument (epidermis and blubber; herein referred to as blubber thickness) in *E. glacialis* in the Bay of Fundy, Canada for five summer seasons and in *E. australis* off the South African coast for two winter seasons. *E. glacialis* had significantly thinner blubber layers (mean  $\pm$  1SD,  $n$ :  $12.23 \pm 2.16$  cm, 172) than *E. australis* ( $16.13 \pm 3.88$  cm, 117), suggesting differing planes of nutrition between the two species. Blubber was thickest in females measured three to six months prior to the start of pregnancy (*E. glacialis*); thinner during lactation (*E. glacialis*, *E. australis*) and then thicker with time after weaning (*E. glacialis*). These results suggest that lipids in blubber are used as energetic support for reproduction in female right whales. Blubber thickness increased in calves during suckling (*E. glacialis*, *E. australis*) but subsequently decreased after weaning (*E. glacialis*). Juvenile and adult male *E. glacialis* blubber thickness was compared among years of differing prey (*Calanus finmarchicus*) abundances (abundance data from Pershing et al. [2005]): during a year of low prey abundance they had significantly thinner blubber than during years of greater prey abundance. Taken together, these results suggest that blubber thickness is indicative of right whale energy balance and that the marked fluctuations in North Atlantic right whale reproduction have a nutritional component.

Key Words: right whale, *Eubalaena*, blubber thickness, body condition, reproduction, physiology, energy reserves

## INTRODUCTION

The North Atlantic right whale *Eubalaena glacialis* is listed as an endangered species by the International Union for Conservation of Nature (IUCN) (Reilly et al. 2008a). The population is small, estimated at a minimum number of 325 in 2007, with an annual growth rate of 1.8 % between 1990-2005 (NOAA 2009). This rate is substantially less than that of several populations of southern right whales *Eubalaena australis* that are growing at about 7% a year (South African population 6.8% during 1971-1998 [Best et al. 2001], Argentinian population: 6.9% during 1971-1990 [Cooke et al. 2001]) and are listed as “Least Concern” by the IUCN (Reilly et al. 2008b). Survival rates for *E. glacialis* were lower in the 1990s than in the 1980s (International Whaling Commission 2001) and the continuing mortalities from vessel strikes and fishing gear entanglements (Knowlton & Kraus 2001, Kraus et al. 2005) are a source of concern (International Whaling Commission 2006). Furthermore, a low reproductive rate and high variability in reproduction also contribute to the endangered status of the species (Kraus et al. 2001, Kraus et al. 2005, Kraus et al. 2007). Since 1980, observed annual calf production fluctuated greatly in *E. glacialis*, e.g. one calf was born in 2000 versus 31 calves in 2001 (Kraus et al. 2007), but showed a steady increase in *E. australis* off the coast of South Africa (Best et al. 2001). Mean calving intervals for mothers of two or more calves in the North Atlantic increased from 3.67 years (1987-1992 [Knowlton et al. 1994]) to over five years (1996-2003), and decreased to 3.2 and 3.5 years in 2004 and 2005 respectively (C.I. not stated) (Kraus et al. 2007). In contrast, mean calving intervals for *E. australis* of the South African population remained steady at 3.12 years (95% CI: 3.07, 3.17) from 1979 to 1995 (Best et al. 2001). This striking difference in reproductive performance between the two right whale species suggests that an

investigation into biological and environmental factors contributing to reproductive success is warranted for a better understanding of the critical status of North Atlantic right whales.

Body fat condition is known to affect fertility in a wide variety of mammalian species (see Young 1976, Frisch 1984, Thomas 1990 for reviews). For example, a minimum level of body fat reserves is necessary for ovulation in humans *Homo sapiens* (Frisch 1984). Likewise, in domestic ewes, *Ovis aries*, a certain body weight and condition score are necessary for optimum ovulation and ovum quality (see McEvoy & Robinson 2003 for review) and body fat condition was strongly correlated with ovulation (Doney et al. 1982). The primary link between body fat condition and fertility appears to be regulated by leptin, a hormone produced by adipose tissue which acts at the level of the hypothalamus and pituitary to stimulate the release of gonadotropin releasing hormone (GnRH; a hormone which regulates the synthesis and release of the gonadotropins: luteinizing hormone [LH] and follicle stimulating hormone [FSH]) and LH (a hormone which regulates many aspects of gonadal function, including ovulation), respectively (see Zieba et al. 2005 for review).

Relationships between body fat condition and fertility have also been observed in a species of large whale. In fin whales *Balaenoptera physalus*, strong correlations between body fat condition, food resource and potential fecundity (determined by proportion of mature non-lactating females with a corpus luteum – a measure of ovulation rate) were observed between 1977 and 1984 (Lockyer 1987). This led us to examine body fat condition of right whales as a potential contributor to the reduced and variable reproductive output of the North Atlantic population.

Blubber is a part of cetacean integument which underlies and interdigitates with the epidermis. It is a highly specialized organ composed of adipose tissue supported by a matrix of

fibrous connective tissue, peripheral nerves and vasculature (Simpson & Gardiner 1972, Ling 1974, Haldiman & Tarpley 1993, Reeb et al. 2007). Cetacean blubber serves a number of functions, including maintaining the hydrodynamic shape of the body (e.g. Koopman 1998, Hamilton et al. 2004), insulating the body core (e.g. Doidge 1990, Dunkin et al. 2005), adjusting buoyancy (e.g. Kipps et al. 2002, Dunkin et al. 2010), and storing energy reserves (e.g. Lockyer 1987, Castellini 2000, Koopman et al. 2002). Blubber can account for 15 – 43% of total body mass in large whales (Lockyer 1991) and accounts for 35 – 45% of total body mass in congeneric right whales in the North Pacific *Eubalaena japonica* (Omura et al. 1969). In fin whales, 96% of the total adipose tissue of the body was located in the blubber (Pond & Mattacks 1988). The adipose tissue in blubber had a similar capacity for glucose utilization (an estimation of relative metabolic activity) as internal adipose depots (Pond & Mattacks 1988), suggesting that blubber is an important indicator of large whale energy reserves. Examining patterns in lipid composition (i.e. total lipids, fatty acid composition and protein content) of the blubber also can lead to inferences about certain functions of the blubber, such as metabolic activity and energy storage (e.g. Ackman et al. 1975, Lockyer et al. 1984, Aguilar & Borrell 1990, Koopman 2007). The lipid content and composition of balaenopterid blubber varied: across the body; with depth from the epidermis; with season and annually (paralleling variation in food availability); and with reproductive status in females (Lockyer 1986, 1987, Aguilar & Borrell 1990). This suggests that changes to the physiological state of the whale increase mobilization and deposition of lipids to the adipose tissue of the blubber. Furthermore, seasonal trends in fattening and thinning in balaenopterid whales were observable in measurements of blubber thickness (see Lockyer 1981 for review).

Information about right whale blubber thickness and body fat condition is comparatively limited. Post mortem measurements combined from a six year collection period suggested that *Eubalaena australis* blubber thickness does not increase during the summer feeding season (Tormosov et al. 1998). According to photogrammetric measurements of body width in *Eubalaena glacialis* and *E. australis*, dorsal body shape was significantly thinner in lactating females than non-lactating, non-pregnant females (Miller et al. In Review). Subjective assessments of dorsal body contours in *E. glacialis* indicated that the external appearance of pregnant females was better than that of lactating and resting females (Pettis et al. 2004). However, just how these assessments relate to blubber thickness (Miller Angell 2006) or more generally, to total adiposity, remains undetermined.

Measurements of blubber thickness are commonly used to assess marine mammal body fat condition. Such measurements of blubber thickness are typically collected during necropsy with a ruler (e.g. Lockyer et al. 1985, Koopman 1998, Tormosov et al. 1998) or during live-captures with ultrasound (Curran & Asher 1974, Gales & Burton 1987). A technique to measure blubber thickness in free-ranging right whales using amplitude-mode ultrasound was developed by Moore et al. (2001). Using this method, acoustic blubber thickness measurements were collected from free-ranging *Eubalaena glacialis* in the North Atlantic and *Eubalaena australis* off the coast of South Africa to investigate the relationships between blubber thickness and reproduction. Blubber thickness measurements collected from *E. glacialis* were also compared with life history parameters (data from North Atlantic Right Whale Consortium Photo-Identification Database [New England Aquarium, Boston, MA, U.S.A.]) and prey abundance in the Gulf of Maine (data from Pershing et al. 2005).

## METHODS

### Data collection

Right whale blubber thickness was measured using amplitude-mode ultrasound (Panametrics *Epoch* III and 9100, Olympus NDT, 48 Ward St, Waltham, MA, USA). A 0.5 MHz Panametrics A301S transducer mounted on the end of a cantilevered 12m carbon fiber pole (Composite Engineering, 277 Baker Ave, Concord, MA, 01742, USA) was deployed to make brief and perpendicular contact with the dorsal aspect of surfacing right whales from small (6 and 7.6 m) boats (Moore et al. 2001). The transducer was cleaned with alcohol between animals and a medical grade gel was used for acoustic coupling. Data were recorded in real time on a laptop computer using custom software (Upper Cape Systems, 29 Prince Henry Drive, East Falmouth, MA 02536 USA).

Acoustic blubber thickness measurements were collected from *Eubalaena glacialis* in Cape Cod Bay, MA, USA, (a winter feeding area) during February-March of 1998 and in the Bay of Fundy, Canada, (a summer feeding area) during August and sometimes early September, 1998 to 2002. *Eubalaena australis* were measured in St. Sebastian Bay, De Hoop Marine Reserve and Walker Bay (Gansbaai), off the southern coast of South Africa (a winter calving area) during August-September, 1999 and October-November, 2000. Parallel data from southern right whales in summer feeding areas were not pursued due to logistical constraints of working in remote, open oceans.

A zero-phase forward and reverse digital filter was applied to each ultrasound record to smooth the acoustic signal in a custom graphical user interface (GUI) (Miller Angell 2006) created in Matlab Version 6.1.0.450, Release 12.1 (The MathWorks Inc., Natick, MA, U.S.A.). The strongest, most consistent acoustic reflections with the greatest transmission times, which

Moore et al. (2001) demonstrated are characteristic of the deepest subdermal connective tissue sheath ([SDS] as defined by Pabst 1996) at the blubber-muscle interface (see discussion), were chosen for measurement. The resultant acoustic transmission times of flight from the chosen reflections were converted to a measurement of distance (cm) using the sound transmission velocity for right whale blubber,  $1483 \text{ m s}^{-1}$  (Miller Angell 2006). The half-range mode (HRM) (Bickel 2002) of these extracted data was calculated and used as the measurement of blubber thickness for each contact with a whale. Note: the measurements of ‘blubber’ thickness presented here are measurements of the entire integument because they include all tissue from the surface of the epidermis to the SDS at the blubber-muscle interface.

Blubber thickness varies across the body in cetaceans and the pattern of variability appears to be species specific (Lockyer et al. 1985, Doidge 1990, Koopman 1998). Although the pattern of variability for right whale blubber has not been defined, blubber thickness data collected at necropsy suggested that the dorsal blubber exhibits low variability in the area approximately between 30 and 50% of the body length from the snout (Figure 1). Blubber thickness measurements from free-swimming right whales inadvertently measured more than once in the same field season also indicated low variability in the same area (C. A. Miller unpubl. data). Therefore, to minimize the effect of variability from measurement location along the body, the following analyses were restricted to dorsal measurements collected between 30 and 50% of the body length from the snout. Measurement locations were determined from the oblique-aerial, digital video footage of each measurement event. With the exception of calves, if an individual whale was measured more than once during a field season within the restricted area on the body, the mean of the measurements from that season was used as the blubber thickness for that individual whale. The 30 to 50% region also was the most consistently accessible.



Total length was measured using aerial photogrammetry (W. L. Perryman unpubl. data – Southwest Fisheries Science Center, NOAA, La Jolla, CA, U.S.A.). Perpendicular aerial photographs of *Eubalaena glacialis* were collected with a 126mm format KA76 military reconnaissance camera (image size 114mm X 114mm) from a Twin Otter airplane during August 2000, 2001 and 2002 in the Bay of Fundy, Canada. The camera was mounted vertically over a camera port in the deck of the aircraft and was operated by an observer stationed at the right forward bubble window. As each photograph was taken, a computer based data acquisition system recorded time, GPS position and altitude from a Honeywell AA300 series radar altimeter. Altitudes for photo passes ranged between 150 m and 250 m. Camera cycle rate was adjusted to provide 60% to 80% overlap between successive images. The photographs used in the study were taken with Kodak Aerochrome HS film (SO-359).

### **Ultrasound comparability, accuracy and precision**

The *Epoch* III ultrasound was used to measure blubber thickness only in 1998 (Moore et al. 2001) because, with such brief contact durations (often less than one second) with the whales, its 2Hz sampling rate limited the quantity and quality of data collected. The Panametrics 9100 ultrasound, deployed during the 1999-2002 field seasons, provided a faster sampling rate of 20Hz, which supplied ample data for measuring blubber thickness. Laboratory testing of ultrasound comparability on *Eubalaena glacialis* blubber samples collected at necropsy indicated that the two ultrasound units provide comparable data (paired t-test:  $t = 1.845$ ,  $df = 4$ ,  $p > 0.05$ ) (Miller Angell 2006). Therefore, the datasets collected by both systems were combined and used in the following analyses.

Laboratory testing of ultrasound accuracy on *Eubalaena glacialis* blubber samples collected at necropsy indicated that the mean difference between blubber thicknesses measured

with a ruler and those measured by ultrasound ( $3.4 \pm 2.8$  mm,  $n = 5$ ) was not significantly different from zero (paired t-test:  $t = 2.679$ ,  $df = 4$ ,  $p > 0.05$ ) (Miller Angell 2006). As such, the ultrasound systems can be regarded as being accurate to 3 mm.

To assess measurement precision, coefficients of variation (CV) were calculated from the extracted data from 30 measurement events chosen at random (mean = 1.7%, range = 0.2% to 4.2%). The low CVs indicated low variability in the blubber thickness data from within a measurement event on free-ranging whales. This variability from within a measurement event was likely a result of three factors: (1) the quality of the contact with the whale, (2) movement of the whale at the time of contact, and (3) choice of acoustic peak during data extraction. To minimize the effects of these sources of variability, the half-range mode (Bickel 2002) was applied to the extracted data and used as the blubber measurement for each contact with a whale. CVs also were calculated from blubber thickness measurements collected from whales measured three or more times during one field season (mean = 6.5%, range = 1.1% to 17.8%,  $n = 16$ ). These CVs were higher, most likely because blubber thickness varies along the body of an individual whale (see above). Although measurement location was limited to an area of low variability along the body, this area is not uniform (Figure 1). It is important to remember that the at-sea blubber thickness measurements are of fibro-elastic material in an animal that is swimming by flexing and extending its body; thus, some degree of variability is inherent in the live system.

### **Right whale identification and demographic classification**

Right whales were individually identified by the patterns of callosities on the top and sides of their heads, patterns of skin pigmentation, and any scarring (Best 1990, Hamilton et al. 2007) from 35 mm photographs taken at the time of measurement. Life history data on individual

*Eubalaena glacialis* were obtained from the North Atlantic Right Whale Consortium Photo-Identification Database, version dated June 29, 2004 (New England Aquarium, Boston, MA, U.S.A.). Life history data on individual adult female *Eubalaena australis* of the South African population were obtained from a photo-identification database created from aerial photographs collected annually on helicopter surveys between 1979 and 2002 (P. B. Best unpubl. data).

For *Eubalaena glacialis*, the sex of individual whales was established from photographic observation of the genital area, molecular markers (Brown et al. 1994) or, for females, close and consistent association with a new born calf (Knowlton et al. 1994). Actual age was determined when year of birth was known, while age class (calf, juvenile, or adult) was based on actual age or length of sighting history. Whales were considered calves for all sightings within the first year of life. Since data from Hamilton & Cooper (2010) suggest that peak calving occurs in December or January and the blubber thickness measurements were collected from late July to early September, calves were approximately eight to nine months old when measured. Sexual maturity occurs at approximately nine years of age (Kraus et al. 2001); therefore, whales were classified as juveniles if they were between the ages of one and eight years and as adults if they were  $\geq$  nine years old, had a sighting history of at least nine years (Hamilton et al. 1998) or earlier for females if they were consistently sighted in close association with a calf (Knowlton et al. 1994). Note: it was assumed that males reach sexual maturity at the same age as females.

Adult female *Eubalaena glacialis* were classified according to their reproductive status (lactating, pre-pregnant, pregnant, resting, nulliparous) at the time of measurement. Lactating females were those that had been sighted consistently with a calf since the winter calving season. At the time of measurement, females in this category had been lactating for approximately eight months and because weaning typically occurs approximately 10 – 12 months after calving

(Hamilton et al. 1995, Hamilton & Cooper 2010), were considered to be in 'late lactation.' Pre-pregnant females were those measured three to six months prior to the estimated start of pregnancy. The estimated start of pregnancy was back-calculated by subtracting gestation duration (estimated as 357 to 396 days [Best 1994]) from the estimated month of calving (month of the first sighting with a calf during the calving season). Only one female was measured while pregnant; she was sighted in consistent, close association with a calf during the subsequent winter calving season. Resting females were those that had been sighted consistently with a calf at least once in their sighting history and were not classified as pre-pregnant, pregnant, or lactating. Resting females could not be subdivided by the number of years resting because of limited sample sizes, so the category includes females that were resting between eight months and six years. Nulliparous females were those that had never been observed consistently in close association with a calf in their sighting histories. Females were also classified as nulliparous from age nine until their first pregnancy (back-calculated from the first calving event).

*Eubalaena australis* of the South African population were measured during the winter calving season, which extends from late June to late October with peak calving in August (Best 1994). Whales were classified as mothers if observed in close, consistent association with a calf. Calves were classified as such based on small size and close, consistent association with a large whale, the presumed mother. Juveniles were classified subjectively in the field as any whale obviously smaller than an adult female, larger than a calf and lacking the dip in the posterior rostrum characteristic of neonate and younger calf right whales. The consequence of subjectively classifying juveniles in the field was that only the smaller, younger juveniles could be consistently determined. Whales were classified as adult females if they had been sighted with a calf in previous years or were known to be a mother in the given measurement season. Mothers

and calves were considered to be lactating and suckling, respectively, given that they were measured within the first few months of birth and weaning is thought to occur 10 to 12 months after birth (Hamilton et al. 1995). Based on the characterization of the *E. australis* calving season (see above) by Best (1994), mothers and calves measured in September 1999 were considered to have been in the early stage of lactation - suckling. The mothers and calves measured in November 2000 may have been lactating and suckling a few days but more likely two to four months and so the November measurements were classified as 'mid-lactation - suckling' measurements.

### **Prey Abundance**

To examine blubber thickness in the context of prey abundance, a *Calanus finmarchicus* abundance index created by Pershing et al. (2005) was chosen. This annual index used raw *C. finmarchicus* abundance measured by the Gulf of Maine Continuous Plankton Recorder (CPR) Program during 1961-1974 and 1978-2003 in the Gulf of Maine. The Gulf of Maine encompasses and is juxtaposed to many of the *Eubalaena glacialis* feeding habitats. Furthermore, interannual variability in the Gulf of Maine appears to capture variability over much of the NW Atlantic shelf (Pershing et al. 2010).

### **Statistical analyses**

Statistical analyses were conducted using Systat, version 10.0 (SPSS, Inc., Chicago, IL) and Matlab, version 6.1.0.450, release 12.1 (The MathWorks Inc., Natick, MA, U.S.A.). Since a relationship between blubber thickness and total length was found only in juveniles and unknown age whales (see results) and since total length measurements were not available for all of the whales in the blubber thickness dataset, the raw, uncorrected blubber thickness data were

used for all analyses, except one. The ratio of blubber thickness to total length was used in a comparison between calves and yearlings.

## RESULTS

*Eubalaena glacialis*. Between 1998 and 2002, 172 blubber thickness measurements were collected. Mean blubber thickness was 12.23 cm (95% CI = 11.90, 12.55; range = 7.93 to 22.33).

*Eubalaena australis*. During 1999 and 2000, 117 blubber thickness measurements were collected. Mean blubber thickness was 16.13 cm (95% CI = 15.41, 16.84; range = 4.89 to 25.82).

### Blubber thickness and total length

*Eubalaena glacialis*. No apparent relationship was found between blubber thickness and length in whales for which total length and blubber thickness measurements were available ( $r = 0.21$ ,  $p = 0.22$ ,  $n = 36$ , Figure 2). However, when these whales were divided according to age class, blubber thickness of juvenile and unknown age whales increased with increasing total length ( $y = 0.01988x - 9.89610$ ,  $R^2 = 0.56$ ,  $p = 0.0004$ ,  $n = 18$ , Figure 2).

### Blubber thickness and age

*Eubalaena glacialis*. Blubber thickness measurements were tested among calves and juveniles of various ages by a one-way ANOVA and a Tukey pairwise comparisons test. Uncorrected blubber thickness data were used in this ANOVA because total length measurements were lacking for most individuals. The results of the post-hoc tests indicate that eight-month old calves had significantly thicker blubber (mean  $\pm$  1SD,  $n$ :  $12.58 \pm 1.16$  cm, 8) than two year olds ( $10.01 \pm 1.42$  cm, 12;  $p = 0.006$ , Figure 3a) and eight year olds had significantly thicker blubber than one, two, three and four year olds ( $p < 0.01$ , Figure 3a).

Three individual right whales measured as calves (eight months old) and again as yearlings (20 months old) showed a 1.7 cm decrease in blubber thickness from calf to yearling (Figure 3b). Also, among the calves and yearlings measured in 2001 and 2002, blubber thickness corrected for total length was significantly greater for calves (mean  $\pm$  1SD,  $n$ :  $1.38 \pm 0.12$ , 7) than for yearlings ( $1.07 \pm 0.12$ , 6) (two-sample t-test:  $t = 4.771$ ,  $df = 11$ ,  $p = 0.0003$ ; Figure 3c).

*Eubalaena australis*. Blubber thickness of calves measured during early suckling in September 1999 (mean  $\pm$  1SD,  $n$ :  $9.22 \pm 2.28$  cm, 8) was significantly thinner than that of calves measured during mid-suckling in November 2000 ( $11.28 \pm 1.8$  cm, 10) (two-sample t-test:  $t = -2.133$ ,  $df = 16$ ,  $p = 0.024$ ; Figure 4a).

### **Blubber thickness and prey abundance**

*Calanus finmarchicus*. *C. finmarchicus* abundance in the Gulf of Maine declined dramatically in 1998 to the third lowest value in the entire Pershing et al. (2005) time series behind those of 1961 and 1970. Abundance recovered in 1999 and 2000 relative to 1998, although the values for these two years still remain below the mean of the values of the entire time series (as indicated by the dotted line in Figure 5). *C. finmarchicus* abundance increased above the time series average in 2001-2003.

*Eubalaena glacialis*. Yearlings measured in 1998 had significantly thinner blubber (mean  $\pm$  1SD,  $n$ :  $9.85 \pm 0.77$  cm, 7) than those measured in 2002 ( $11.42 \pm 1.46$  cm, 6) (two-sample t-test:  $t = -2.462$ ,  $df = 11$ ,  $p = 0.016$ ; Figure 5).

Adult males measured in 1998 had significantly thinner blubber (mean  $\pm$  1SD,  $n$ :  $11.89 \pm 1.31$  cm, 21) than those measured in 1999 ( $13.25 \pm 1.77$  cm, 10; Smith-Satterthwaite (unpooled) t-test (Devore 1995):  $t = 2.162$ ,  $df = 13.9$ ,  $p = 0.024$ ) and in 2000 ( $13.98 \pm 1.21$  cm, 3; Smith-Satterthwaite t-test (Devore 1995):  $t = 2.759$ ,  $df = 2.7$ ,  $p = 0.039$ ; Figure 5). The blubber

thickness of adult males measured in 1999 did not differ statistically from that of the males measured in 2000, according to a Smith-Satterthwaite t-test (Devore 1995) ( $t = 0.811$ ,  $df = 4.9$ ,  $p = 0.455$ ; Figure 5).

### **Blubber thickness and reproduction**

*Eubalaena glacialis*. Pre-pregnant females had significantly thicker blubber (mean  $\pm$  1SD,  $n$ :  $17.23 \pm 0.67$  cm, 4) than lactating females ( $12.73 \pm 1.31$  cm, 11) (Smith-Satterthwaite t-test (Devore 1995):  $t = 8.712$ ,  $df = 10.8$ ,  $p = 1.60e-06$ ; Figure 6) and had significantly thicker blubber than nulliparous females ( $13.64 \pm 1.44$  cm, 8) (Smith-Satterthwaite t-test (Devore 1995):  $t = 5.898$ ,  $df = 10.0$ ,  $p = 7.75e-05$ ; Figure 6). The blubber of nulliparous females did not differ from lactating females (two-sample t-test:  $t = 1.427$ ,  $df = 17$ ,  $p = 0.17$ ; Figure 6). Due to a sample size of one, the measurement from the pregnant female was not included in the analysis.

Four individual females were measured in multiple years at different stages of the reproductive cycle (Figure 7) and are referred to by their identification number from the North Atlantic Right Whale Catalog (Hamilton et al. 2007). The blubber of #1407, measured after resting for three years in 1998, was 4.6 cm thicker than the measurement collected after eight months of lactation in 2001, while #1701's blubber was 1.8 cm thicker when measured within a few months of pregnancy in 1999, than after eight months of lactation in 2001. When #1911 was measured in 1998 as an immature, nulliparous female, her blubber was 1.1 cm thinner than when she was measured after eight months of lactation in 2001. The whale #2050 was measured three times. Between 1998 and 1999, her blubber thickness increased by 4.2 cm as she transitioned from nulliparous to pre-pregnant. Her blubber then decreased by 3.4 cm after eight months of lactation in 2001.



In reproductively active females, a positive relationship was found between blubber thickness and number of years since last calving (Pearson's correlation;  $r = 0.75$ ,  $p = 0.02$ ,  $n = 9$ ; Figure 8).

*Eubalaena australis*. Mean blubber thickness from mothers measured in September 1999, early lactation, (mean  $\pm$  1SD,  $n$ :  $19.82 \pm 2.89$  cm, 12) was significantly thicker than the mean of those measured in November 2000, mid-lactation, ( $17.24 \pm 1.84$  cm, 19) (two-sample t-test:  $t = 2.761$ ,  $df = 16.7$ ,  $p = 0.007$ ; Figure 4b).

*Eubalaena glacialis* and *Eubalaena australis*. No difference was found between the blubber thickness of primiparous females (females that have had only one calf) and that of multiparous females (females that have had two or more calves). *E. australis* lactating females mean  $\pm$  1SD,  $n$ : primiparous =  $18.78 \pm 2.69$  cm, 9, vs. multiparous =  $18.15 \pm 3.06$  cm, 17 (Smith-Satterthwaite t-test (Devore 1995):  $t = -0.538$ ,  $df = 18.4$ ,  $p = 0.597$ ). *E. glacialis* lactating females mean  $\pm$  1SD,  $n$ : primiparous =  $12.83 \pm 1.36$  cm, 4, vs. multiparous =  $12.68 \pm 1.38$  cm, 7 (two-sample t-test:  $t = -0.174$ ,  $df = 9$ ,  $p = 0.866$ ).

### **North Atlantic vs. South African right whale blubber thickness**

Blubber was significantly thinner in *Eubalaena glacialis* (mean  $\pm$  1SD,  $n$ :  $12.23 \pm 2.16$  cm, 172) than in *Eubalaena australis* ( $16.13 \pm 3.88$  cm, 117) (Smith-Satterthwaite t-test (Devore 1995):  $t = -9.869$ ,  $df = 165.1$ ,  $p = 0.000$ ; Figure 9a). When the species were compared by age classes, *E. glacialis* blubber was significantly thinner than *E. australis* blubber for all age classes except calves (Adult females: *E. glacialis* ( $13.93 \pm 2.49$  cm, 32) vs. *E. australis* ( $18.13 \pm 2.69$  cm, 35) ( $t = -6.167$ ,  $df = 65$ ,  $p = 0.000$ ; Figure 9b); Juveniles: *E. glacialis* ( $11.25 \pm 1.76$  cm, 75) vs. *E. australis* ( $15.73 \pm 2.11$  cm, 19) ( $t = -8.549$ ,  $df = 24.8$ ,  $p = 0.000$ ; Figure 9c). *E. glacialis* calves ( $12.58 \pm 1.16$  cm, 8) were significantly fatter than *E. australis* calves ( $10.36 \pm 2.24$  cm,

18) ( $t = 3.324$ ,  $df = 23.2$ ,  $p = 0.003$ ; Figure 9d), presumably because *E. glacialis* calves were approximately five to seven months older.

## DISCUSSION

A combination of at-sea acoustic blubber thickness measurements and more than two decades of life history and reproductive data were used to assay body fat condition of two species of free-ranging right whales, *Eubalaena glacialis* in the North Atlantic and *Eubalaena australis* off the southern coast of South Africa. Adult females had the thickest blubber prior to pregnancy, calves showed a loss of blubber thickness after weaning, juveniles gained blubber thickness with length, males and yearlings had blubber thickness that varied with prey abundance, and *E. australis* had thicker blubber than *E. glacialis*. *E. australis* mothers were also shown to lose blubber thickness as their calves gained it during suckling.

Acoustic blubber thickness data from *Eubalaena glacialis* presented here (range = 8 to 22 cm) were comparable to ruler measurements of blubber thickness from along the dorsal midline at the level of the axilla collected during necropsy of *E. glacialis* (range = 8 to 17 cm [Moore et al. 2004]). *Eubalaena australis* acoustic blubber thickness data (range = 5 to 26 cm) were comparable to those reported during Soviet whaling (range = 5 to > 30 cm [Tormosov et al. 1998]), where the measurement was taken mid-laterally at the level of the anus. It should be noted that the difficulties inherent in measuring blubber thickness on carcasses often result in inconsistencies within and between datasets; therefore, comparisons among and between such datasets should be made with caution.

### **Integument morphology and the acoustic assay**

Acoustic reflections from boundaries between materials of different acoustic impedance allow measurement of blubber thickness. Upon gross visual examination of the innermost aspect of *Eubalaena glacialis* integument (including underlying skeletal muscle) collected at necropsy, Moore et al. (2001) described two tendinous connective tissue (CT) sheaths separated by loose CT. The distal sheath (described as the ‘upper’ sheath by Moore et al. [2001] ) occurred at the innermost aspect of the blubber while the proximal sheath (described as the ‘lower’ sheath) overlaid the skeletal muscle. The ultrasound examination of this same sample of integument from *E. glacialis* clearly showed acoustic reflections from both CT sheaths (Moore et al. 2001). The two sheaths are interconnected periodically and the distal sheath may not always be present. The proximal sheath is homologous to the subdermal connective tissue sheath (SDS) described for dolphins by Pabst (1996). In the sei whale *Balaenoptera borealis*, two ‘layers’ (a 3-5 mm thick sheath comprised of elastic and collagen fibers and a region of dense collagenous bundles overlaying muscle) separated by well-vascularized adipose tissue interspersed with collagen fibers were observed in the innermost aspect of the blubber (Ackman et al 1975). Also, Ackman et al (1975) pointed out that according to Heyerdahl (1932), the ‘isterlag’ (a layer with elevated lipid content) was situated between two layers of connective tissue at the innermost aspect of fin *Balaenoptera physalus* and blue whale *Balaenoptera musculus* blubber. In bowhead whale *Balaena mysticetus* integument, Haldiman and Tarpley (1993) described a highly tendinous CT sheath (1-2 mm thick) bordering the innermost (deepest) part of the ‘true’ blubber (reticular dermis) which overlays a layer of loose CT containing varying amounts of adipose tissue (hypodermis), followed by the skeletal muscle. Haldiman and Tarpley (1993) did not describe a second tendinous CT sheath (a proximal or ‘lower’ sheath) overlaying the skeletal muscle. It is likely that the loose CT between the two CT sheaths described by Moore et al. (2001) is

homologous to the hypodermis (loose CT layer) between the CT sheath and skeletal muscle described for bowhead whales, the region of well-vascularized adipose tissue layer observed in sei whales and the 'isterlag' observed in blue and fin whales. Our technology and analysis failed to consistently define a distal ('upper') sheath but was successful and consistent in acquiring reflections from the proximal sheath. As a result, the measurements of 'blubber' thickness presented here include all tissue from the surface of the epidermis to the proximal CT sheath at the blubber-muscle interface.

The thick epidermis of cetaceans (Ling 1974) can vary by species (Simpson & Gardiner 1972). Because the measurements presented here include epidermis, it is important to understand the contribution of epidermis to the thickness of the entire integument for both *Eubalaena glacialis* and *Eubalaena australis*. Yet, the thickness of the epidermis could not be measured because the ultrasound was optimized for detecting structures (the tendinous CT sheath) deeper in the integument. Epidermis was measured on full-core biopsies of *E. australis* adults (mean  $\pm$  1SE,  $n$ :  $1.42 \pm 0.09$  cm, 13) and calves ( $1.39 \pm 0.07$  cm, 19) (Reeb et al. 2007) and on the body of *E. glacialis* at necropsy (1.1, 1.1, 1.2 cm for three whales; M. J. Moore unpubl. data). It is difficult to determine whether or not there is a difference in epidermal thickness between the two species based on these data. The measurements of epidermis from *E. glacialis* may appear thinner than measurements from biopsied *E. australis* because measurements from *E. australis* were not corrected for the angle of the biopsy (Reeb et al. 2007) and/or because of possible post-mortem compaction of the epidermis due to post-mortem rendering of the epidermal lipids, post-mortem sloughing, the stresses and tension from the fibro-elastic integument being measured while attached to the body (whereas a measurement of a biopsy is free of such stresses and tensions), and/or the distribution of the whale's weight with respect to the location of

measurement. However, based on the aforementioned epidermal thicknesses and the mean 'blubber thickness' (integument thickness) for each right whale species, epidermis only accounts for a small proportion of the entire thickness of the integument and any difference in epidermal thicknesses between the two species is likely to be on the scale of a few millimeters or less. Therefore, the thickness of the epidermis most likely does not make a sizeable contribution to any differences in thickness of the integument presented here. As such, the observed differences in ultrasound 'blubber thickness' measurements are driven primarily by changes in dermal and/or hypodermal thickness.

The blubber of large whales is not metabolized as a whole (Castellini 2000), but rather lipids are deposited to or mobilized from adipocytes, which are supported by a matrix of collagen. In several other cetaceans, trends in blubber lipid content (either measured directly or inferred from adipocyte morphology) parallel trends in blubber thickness with respect to reproductive (adult females) and nutritive status (Lockyer 1981, 1986, 1987, Aguilar & Borrell 1990, Koopman et al. 2002). In *Eubalaena australis* mothers, changes in blubber lipid composition (Reeb 2001) paralleled the observed thinning of blubber (Figure 4) during the initial months of lactation. Such evidence suggests that variability in lipid content and composition contributed to the changes/differences in blubber thickness presented here. Moreover, lipid content and composition of blubber can vary both across the body and with depth from the epidermis in cetaceans, which has led to the hypothesis that certain areas and blubber depths are more metabolically active than others (e.g. Ackman et al. 1975, Lockyer et al. 1984, Aguilar & Borrell 1990, Koopman 2007). Although such variability in blubber lipids across the body and with depth from the epidermis needs to be determined in right whales, the measurements of 'blubber' thickness presented here include all tissue from the surface of the epidermis to the

proximal CT sheath at the blubber-muscle interface, thus integrating the entire integumentary energy store.

### **Blubber thickness and body length**

Positive linear relationships between blubber thickness and body length were found in fin and blue whales (Nishiwaki & Hayashi 1950, Nishiwaki & Oye 1951), while a negative relationship was observed in harbor porpoise *Phocoena phocoena* (Koopman 1998). Blubber thickness and body length were not related in gray whales *Eschrichtius robustus* (Rice & Wolman 1971). Tormosov *et al.* (1998) plotted a linear relationship between blubber thickness and body length in *Eubalaena australis* taken by Soviet whaling operations. Blubber thickness and body length were not related in *Eubalaena glacialis* when all data were analyzed together (Figure 2). However, when the life history stages were analyzed separately, blubber thickness significantly increased with increasing body length in juveniles. Interestingly, by visualizing this strong relationship in juveniles with the data from other life history stages (Figure 2), the relationship for all whales combined may have been skewed by calves having thicker blubber and lactating females having thinner blubber than expected for their respective body lengths.

### **Blubber thickness and nutritional regime**

In *Eubalaena australis* calves of the South African population, blubber thickness increased significantly during the initial months of suckling (Figure 4a), a time during which rapid growth in length occurs,  $2.8 \pm 0.7 \text{ cm d}^{-1}$  (Best & Rüther 1992). The blubber of *Eubalaena glacialis* calves measured in the late months of suckling was as thick as older juvenile whales (seven and eight year old whales), much thicker than the younger juveniles (Figure 3a). The substantially thicker blubber of calves suggests that the mother's milk allows right whale calves

to be in a positive energy balance, storing lipids in the blubber while simultaneously growing in length at a rapid rate.

Significant thickening of blubber during suckling reversed to a significant thinning once the calves were weaned (Figure 3). Weaning appears to occur typically at approximately 10 – 12 months of age (Hamilton et al. 1995, Hamilton & Cooper 2010). Similarly in harbor porpoise, blubber thickness decreased from calves to adults, with the greatest relative decline occurring immediately after weaning (Koopman 1998). Effects of weaning, such as interruptions in the growth curve have been observed in other mammalian species (Pond 1977). A post-weaning hiatus in growth of body length appeared to occur in juvenile bowhead whales (Lubetkin et al. 2008) and in juvenile *Eubalaena australis* (Best & Schell 1996). The post-weaning thinning of blubber is likely an indication that young right whale juveniles rely on the lipids in their blubber to help support energetic challenges as they transition to nutritional independence.

The principal prey species of *Eubalaena glacialis* is the later stages of the copepod *Calanus finmarchicus* (Kenney et al. 1986, Baumgartner & Mate 2003, Michaud & Taggart 2007), the abundance of which can be highly variable, both seasonally and annually, and appears to be influenced by climate (Conversi et al. 2001, Greene et al. 2003, Pershing et al. 2005). Right whales must find prey patches dense enough for feeding to be energetically efficient (Kenney et al. 1986, Baumgartner & Mate 2003, Michaud & Taggart 2007) because they are essentially ram filter feeders without the ability to concentrate prey (Baumgartner et al. 2007). Differences in blubber thickness of adult males and yearlings were detected among years during which *C. finmarchicus* abundance in the Gulf of Maine (GoM) fluctuated greatly (Pershing et al. 2005) (Figure 5). In 1998, *C. finmarchicus* abundance in the GoM declined to the third lowest value of the entire Pershing et al. (2005) time series (1961-1974, 1978-2003). In 1999 and 2000 *C.*

*finmarchicus* abundance in the GoM recovered relative to 1998 but was still below the Pershing et al. (2005) time series average. Notably, blubber thickness of adult male *E. glacialis* mirrored *C. finmarchicus* abundance in these years: significantly thinner in 1998 when compared with 1999 and 2000 (Figure 5). *C. finmarchicus* abundance in the GoM (Pershing et al. 2005) then increased to above average values in 2001 and 2002, but only one adult male was measured in each of these years.

Likewise, yearlings measured in 1998 had significantly thinner blubber than those measured in 2002, the year of the second highest *Calanus finmarchicus* abundance in the GoM (Pershing et al. 2005) of our study period (Figure 5). As discussed above, the significant thinning of blubber thickness between calf and yearling *Eubalaena glacialis* suggests that the transition to nutritional independence is energetically challenging. For the 1998 yearlings, this already challenging transition to nutritional independence likely was intensified by the dramatically low abundance of *C. finmarchicus*. Furthermore, the 1998 yearlings may have been weaned earlier than the 2002 yearlings: many of the 2002 yearlings were sighted in close association with their mothers well into their second year of life, a phenomenon which rarely has been observed in *E. glacialis* and may indicate that suckling was prolonged for the 2002 yearlings (Hamilton & Cooper 2010). Nutrition in early postnatal life, particularly prior to weaning, influences the number of adipocytes and total adiposity, as observed in woodchucks (Young 1976) and rats (Knittle & Hirsch 1968). As such, the thicker blubber of the 2002 yearlings may have been indicative of better nutrition, whether the result of greater availability of *C. finmarchicus* during the transition to nutritional independence and/or prolonged suckling.

### **Blubber thickness and reproduction**



Female mammals, particularly from those species that fast during portions of the reproductive cycle, can increase their chances of reproductive success by acquiring adequate fat reserves to ensure survival of themselves and healthy offspring. The mechanism for this appears to occur prior to the start of the reproductive cycle: ample reserves of body fat are necessary for ovulation to occur (e.g. Doney et al. 1982, Frisch 1984, Albon et al. 1986) because sufficient levels of leptin, the hormone produced by adipose tissue which is correlated with adipose tissue mass (see Houseknecht et al. 1998, Chilliard et al. 2005 for reviews), are necessary for successful reproduction (see Zieba et al. 2005, Barb et al. 2008, Hill et al. 2008 for reviews). Leptin appears to serve as the primary metabolic signal linking energy status with the neuroendocrine system and subsequent reproduction (see Houseknecht et al. 1998, Zieba et al. 2005, Barb et al. 2008, Hill et al. 2008 for reviews). Through a complex process involving an array of neuroendocrine and endocrine signals, leptin acts on the hypothalamus and pituitary to stimulate the release of GnRH and LH, respectively (see Zieba et al. 2005, Barb et al. 2008, Hill et al. 2008 for reviews) thereby, influencing ovulation and subsequent reproduction.

Given the importance of body fat and nutritional status to mammalian reproduction, particularly with regards to ovulation, it is noteworthy that adult females measured prior to the start of pregnancy had the thickest blubber of all life history classes in *Eubalaena glacialis* (Figure 6). Also, the blubber of an individual female increased by 4.2 cm during the year leading up to the start of pregnancy (Figure 7). In balaenopterids, pregnant females had the thickest blubber (Lockyer 1981, 1987) and the highest blubber lipid content (Lockyer 1987, Aguilar & Borrell 1990) of all age classes but peri-ovulatory females were not specified. Body fat condition data from prior to the potential start of pregnancy appear to exist for only one other species of large whale. The blubber of female gray whales which had recently ovulated was thicker than

that of immature males and females, adult males and adult anoestrous and postpartum females but similar in thickness to that of females taken during the early and late stages of pregnancy (Rice & Wolman 1971). Based on these results, in which improved body condition was apparent prior to the start of pregnancy (right whales) and peri-ovulatory and during gestation (gray whales), and the preponderance of evidence in this respect from other mammals, body fat appears to play a critical, and possibly permissive, role in the initiation of the large whale reproductive cycle. The significant thickening of blubber prior to the start of pregnancy suggests that, like terrestrial mammals, a critical level of body fat appears to be important for successful ovulation and subsequent reproduction in right whales.

Hlista et al. (2009) observed a significant positive relationship between the number of right whale calves born each year between 1998 and 2007 (an indicator of reproductive success) and a weighted annual chlorophyll concentration (a realization of a right whale nutritional index based on concentrations coinciding with peak right whale sightings in high-use habitats) averaged over the prior two years, the significance of which appeared to be driven primarily by the strong relationship between annual right whale calf counts and weighted annual chlorophyll concentrations from two years prior. Such relationships concur with the hypothesis that feeding conditions in the year leading up to conception likely lead to substantial accumulation of blubber lipid reserves (as indicated by the thickest blubber observed in pre-pregnant females) subsequently affecting calf production through the short and long term metabolic cues necessary for successful reproductive function (e.g. Prunier & Quesnel 2000, Barb et al. 2008, Hill et al. 2008).

As mentioned above, pregnant balaenopterids and peri-ovulatory and pregnant gray whales had the thickest blubber of all life history classes in these species (Rice & Wolman 1971,

Lockyer 1981, 1987). Moreover, the similarity in blubber thickness among peri-ovulatory, early pregnant and late pregnant gray whales (Rice & Wolman 1971) suggests that the blubber thickness of gray whale females does not change between ovulation and late gestation. The lack of relationship between annual calf count and weighted annual chlorophyll concentration from one year prior (Hlista et al. 2009) may indicate that feeding conditions during gestation are not as critical to calf production as those leading up to conception. Having measured only one pregnant right whale, further study is necessary to examine such trends during gestation in right whales.

In other large whales, lactation was calculated as the costliest part of the reproductive cycle (Lockyer 1981) and in other mammals, lactation was estimated to require three to five times more energy than gestation (Young 1976). Much of these energetic costs of lactation likely occur during the initial months: right whale mothers do not appear to feed while on the calving grounds, and consequently are believed to support their calves and themselves during these months by relying on endogenous nutrient reserves – a concept supported by these results. Blubber thickness decreased significantly between the second month of lactation (September 1999) and the fourth month (November 2000) among lactating *Eubalaena australis* females (Figure 4b). Likewise, when individual *E. australis* mothers were measured on multiple occasions within the same calving season, dorsal body shape (determined by a series of aerial photogrammetric measurements of body width) thinned significantly during the initial months of lactation (Miller et al. In Review), suggesting that the difference in blubber thickness was due to lactation rather than an inter-annual difference in blubber thickness. Corresponding to this significant thinning of blubber during the initial months of lactation, Reeb (2001) detected a significant change in blubber fatty acid composition in lactating *E. australis* mothers between these same months (September 1999 and November 2000), not only suggesting that lipids in

right whale blubber are affected by lactation, but also supporting the assertion that the thinning of blubber during lactation likely is a result of catabolism of blubber lipids.

Among the resting *Eubalaena glacialis* females, blubber thickness was positively related to the number of years since the last calving event, suggesting recovery and replenishment of body fat reserves during the time between calves (Figure 8). Right whale females are capable of a three-year calving interval (Knowlton et al. 1994, Best et al. 2001, Kraus et al. 2007): one year of gestation, one year of lactation, and one year of recovery. Considering that the greatest difference in blubber thickness was observed between females that had been lactating for eight months and those measured a few months prior to the start of gestation, right whale females would need to encounter enough food not only to cover daily metabolic needs, but also to replenish lipid stores during the year of recovery. If right whale females do not encounter adequate food supply, the replenishment of body fat reserves may slow and increases in calving interval may occur. For example, in primiparous pigs, a longer weaning to first oestrus interval was observed in sows with lower backfat as compared to sows with higher backfat (Tummaruk et al. 2001). Moreover, nutritional status not only is important to the replenishment of fat reserves, but also can affect reproductive performance. For instance, chronic feed restriction interfered with normal cyclicity in the pig where ovarian activity was altered through the alteration of hypothalamic release of GnRH (Armstrong & Britt 1987). In this regard, it is interesting that a significant increase in calving interval among *E. glacialis* from 3.67 years (1987-1992) to greater than five years (1993-1998) (Kraus et al. 2001) paralleled changes in *Calanus finmarchicus* abundance, which was low throughout the 1990s relative to the late 1980s, when abundance was at its highest in the Gulf of Maine (Pershing et al. 2005).

In *Eubalaena glacialis*, the blubber thickness of nulliparous females was significantly thinner than that of pre-pregnant females but not statistically different from the energetically drained lactating females (Figure 6). A few of the nulliparous females that were measured during this study subsequently produced calves, including *E. glacialis* #2050, whose blubber increased by more than 4 cm in the year leading up to the start of pregnancy (see below). As of 2005, 12% of adult *E. glacialis* females were reproductively inactive (Kraus et al. 2007) which in part, as our results suggest, may be a consequence of females not attaining adequate body fat reserves necessary for successful reproduction.

Four individual *Eubalaena glacialis* females were measured in multiple years at different stages in their reproductive cycles (Figure 7). These measurements offer evidence that annual changes in blubber thickness of individual whales are detectible and serve to corroborate our population-based results. In particular, the series of measurements from *E. glacialis* #2050 highlight that blubber thickness changes among reproductive stages can be marked. *E. glacialis* #2050 was thinnest as a nulliparous female when measured in 1998, when *Calanus finmarchicus* abundance in the GoM (Pershing et al. 2005) was dramatically low. One year later, as pre-pregnant, her blubber was 4.2 cm thicker, the thickest measurement collected during the series. She was not measured the next year during gestation but was measured again the following year after 8 months of lactation. Blubber thickness decreased by 3.4 cm between the pre-pregnant measurement and the measurement collected after 8 months of lactation.

The mean blubber thickness of adult male *Eubalaena glacialis* was similar to that of older juveniles, and lactating, resting, and nulliparous adult females (Figure 6). Mean adult male blubber thickness was affected by the thin blubber measurements collected from males in 1998, but the mean blubber thicknesses of males in 1999 and 2000 were essentially the same as that of

nulliparous females (see results). In human males, weight loss and low body fat may result in decreased sperm mobility and longevity, decreased hypothalamic GnRH, and decreased levels of testosterone and prolactin in blood serum levels (see Frisch 1997 for review) and such interactions between body fat and fertility may be mediated by the hormone leptin. In bulls, rams, boars and stallions, malnutrition was associated with decreases in spermatogenesis (Foote 1978). Given that male right whales have the largest testes among the baleen whale species, both in absolute weight and relative to body weight (Brownell & Ralls 1986), and are believed to use sperm competition as a mating strategy (Brownell & Ralls 1986, Kraus & Hatch 2001), sexual activity may be energetically costly for male right whales. Male blubber thickness and body fat condition should therefore, in the future, be analyzed in the context of paternity.

### **North Atlantic vs. South African right whale blubber thickness**

When reproductive parameters of the *Eubalaena glacialis* are compared with those of the *Eubalaena australis* populations, *E. glacialis* reproduction appears to be much more variable and at times compromised (see introduction). In this regard, it is interesting that blubber thickness, measured during a depressed time in *E. glacialis* reproduction (Kraus et al. 2007), was significantly thinner than in *E. australis* of the South African population (Figure 9). The difference in blubber thickness remained significant when the species were compared within age classes, except for the comparison between calves (Figure 9d), where *E. glacialis* calves had thicker blubber because they were approximately three to nine months older than the *E. australis* calves.

A number of variables could influence the observed difference in blubber thickness between *Eubalaena glacialis* and *Eubalaena australis*: differences in body size, differences in measurement season, differences in habitat temperature regime, genetic differences, or

differences in planes of nutrition. Whales belonging to these two species appear to be comparable in size (Best & Rüther 1992, W.L. Perryman, pers. comm.) but without measurements of total length from the individuals for which measurements of blubber thickness were collected, it was not possible to test whether or not differences in total length influenced the observed difference in blubber thickness.

*Eubalaena glacialis* blubber was measured during the summer feeding season while *Eubalaena australis* was measured during the winter breeding season. In balaenopterid whales, blubber thickens during the summer feeding season and thins during the winter migration (see Lockyer 1981 for review). Summer seasonal fattening was not detected in measurements of blubber thickness in southern right whales taken by whaling (Tormosov et al. 1998). Such fattening may not have been detected due to the pooling of the data (data for all individuals were pooled over a six year period [Tormosov et al. 1998]), differences in individual whale arrival times to the feeding grounds (e.g. Lockyer 1981, Lockyer et al. 1985), inter-annual differences in prey abundance or in seasonality of prey abundance, or because of limited seasonal coverage (no winter measurements were available). However, our results indicate that the thickness of right whale blubber can change seasonally in lactating females and annually depending on nutritional and reproductive status. If seasonal fattening and thinning does occur in right whales, everything else being equal, measurements of blubber thickness from *E. glacialis* collected during summer feeding should be greater than those from *E. australis* collected during winter fasting, minimizing what would have been an even larger difference had the seasons been comparable.

Right whales experience a wide range of water temperatures on a daily (swimming between surface and depth) and seasonal basis (both species are seasonally migratory in subtropical and temperate waters), so it is difficult to imagine that the whales would respond to

changing temperatures by changing the thickness of the blubber on a short term time scale. However, if the overall temperature range experienced by *Eubalaena australis* was consistently lower than that experienced by *Eubalaena glacialis*, it is possible that *E. australis* had thicker blubber necessary for insulation against the colder temperatures. But equally, it is possible that right and other large whales are constrained by the need to lose heat (Alexander 1998).

Recent analysis of genetic markers from both mitochondrial (Rosenbaum et al. 2000) and nuclear DNA (Gaines et al. 2005), indicate that right whales of the North Atlantic and of the Southern Hemisphere are two distinct species, *Eubalaena glacialis* and *Eubalaena australis* respectively. It is possible that genetic factors controlling fat storage and/or utilization may affect the observed difference in blubber thickness between the two species.

Finally, the difference in blubber thickness between the two species could be influenced by differing planes of nutrition. Both species feed on lipid-rich, late stage calanoid copepods (Kenney et al. 1986, Tormosov et al. 1998, Baumgartner & Mate 2003, Michaud & Taggart 2007); southern right whales also feed on euphausiids (Hamner et al. 1988, Tormosov et al. 1998, Reeb 2001). Prey species, life stage, available biomass and patch density will impact relative foraging success; therefore, in aggregate, southern right whales appear to be more successful at accumulating energy reserves.

Although further investigation is needed to fully understand the observed difference in blubber thickness between *Eubalaena glacialis* of the North Atlantic and *Eubalaena australis* off of the southern coast of South Africa, the difference is suggestive in the context of the differences in reproductive performance of the two species. According to our results, adequate food supply appears to be necessary for adequate body fat reserves which, in turn, appear to be necessary for successful reproduction in right whales. Therefore, the thicker blubber of *E.*



*australis*, which may indicate a better nutritional regime, likely contributes to the more successful and stable reproductive performance of this population, where the population growth rate is close to the maximum biologically possible.

## ACKNOWLEDGEMENTS

We thank all those who assisted with field work: M.S. Morss, J.N. Ciano, M.K. Marx, P.K. Hamilton, L.A. Conger, C.K. Slay, S.D. Kraus, B.M. McKay, J.N. Wocial, T.R. Frasier, C.J. Harper, M. Levine, L. Morse, O. Nichols, M. Piche, B. Bjermi, H.T.C. Moore and the Moore family, M. Best, M. Thornton, A. Du Randt, D. Kemp, M. Addison, and D.P. Nowacek, A. Shorter and the WHOI D-tag project. We thank the New England Aquarium Right Whale Research Project for field support, matching of the individual whales and the creation and ongoing maintenance of the right whale database. The Right Whale Consortium generously granted access to the right whale database; v. 29 June 2004 was used. S. Berube and D. Kass of Panametrics, Inc. assisted with the ultrasound equipment. K. Prada, Upper Cape Systems, Inc. created the software to acquire and record the ultrasound data. B. Lange and D. Wright made possible the video documentation of the blubber thickness measurements. Mazda Wildlife Fund provided invaluable vehicle support in South Africa. W.L. Perryman generously shared total length data. G. Terray, A. Solow and A. Beet assisted with data extraction. We thank M.F. Baumgartner and A.J. Pershing for discussions about right whale prey, body condition and reproduction. R. Asmutis-Silvia worked on the development of the project in the early years. We also acknowledge four anonymous reviewers whose thorough efforts significantly improved the manuscript. Acoustic blubber measurements of *Eubalaena glacialis* were collected under license numbers 1999-026, 2000-014, 2001-020, and 2002-029 from the Canadian Department of Fisheries and Ocean and National Marine Fisheries Service permit number 1014 to S.D. Kraus. Measurements of *Eubalaena australis* were collected under permits to P.B.B. from the South African Department of Environment Affairs and Tourism. This project was made possible with funds provided by: Massachusetts Environmental Trust, Office of Naval Research, National

Marine Fisheries Service – National Oceanic and Atmospheric Administration, Northeast Consortium, Hussey Foundation, and National Research Foundation in South Africa.

## LITERATURE CITED

- Ackman RG, Hingley JH, Eaton CA, Sipos JC, Mitchell ED (1975) Blubber fat deposition in mysticeti whales. *Can J Zool* 53:1332-1339
- Aguilar A, Borrell A (1990) Patterns of lipid content and stratification in the blubber of fin whales (*Balaenoptera physalus*). *J Mammal* 71:544-554
- Albon SD, Mitchell B, Huby BJ, Brown D (1986) Fertility in female red deer (*Cervus elaphus*): The effects of body composition, age and reproductive status. *J Zool Lond* 209:447-460
- Alexander RM (1998) All-time giants: the largest animals and their problems. *Palaeontology* 41:1231-1245
- Armstrong JD, Britt JH (1987) Nutritionally-induced anestrus in gilts: metabolic and endocrine changes associated with cessation and resumption of estrus cycles. *J Anim Sci* 65:508-523
- Barb CR, Hausman GJ, Lents CA (2008) Energy metabolism and leptin: effects on neuroendocrine regulation of reproduction in the gilt and sow. *Reprod Domest Anim* 2:324-330
- Baumgartner MF, Mate BR (2003) Summertime foraging ecology of North Atlantic right whales. *Mar Ecol Prog Ser* 264:123-135
- Baumgartner MF, Mayo CA, Kenney RD (2007) Enormous carnivores, microscopic food, and a restaurant that's hard to find. In: Kraus SD, Rolland RM (eds) *The urban whale: North Atlantic right whales at the crossroads*. Harvard University Press, Cambridge, Massachusetts, p 138-171
- Best PB (1990) Natural markings and their use in determining calving intervals in right whales off South Africa. *S Afr J Zool* 25:114-123
- Best PB (1994) Seasonality of reproduction and the length of gestation in southern right whales *Eubalaena australis*. *J Zool Lond* 232:175-189
- Best PB, Brandão A, Butterworth DS (2001) Demographic parameters of southern right whales off South Africa. *J Cetacean Res Manag (Special Issue)* 2:161-169
- Best PB, R  ther H (1992) Aerial photogrammetry of southern right whales, *Eubalaena australis*. *J Zool Lond* 228:595-614
- Best PB, Schell DM (1996) Stable isotopes in southern right whale (*Eubalaena australis*) baleen as indicators of seasonal movements, feeding and growth. *Mar Biol* 124:483-494
- Bickel DR (2002) Robust estimators of the mode and skewness of continuous data. *Comput Stat Data Anal* 39:153-163
- Brown MW, Kraus SD, Gaskin DE, White BN (1994) Sexual composition and analysis of reproductive females in the North Atlantic right whale, *Eubalaena glacialis*, population. *Mar Mamm Sci* 10:253-265
- Brownell RL, Ralls K (1986) Potential for sperm competition in baleen whales. *Rep Int Whaling Comm (Special Issue)* 8:97-112
- Castellini M (2000) History of polar whaling: insights into the physiology of the great whales. *Comp Biochem Physiol A* 126:153-159
- Chilliard Y, Delavaud C, Bonnet M (2005) Leptin expression in ruminants: nutritional and physiological regulations in relation with energy metabolism. *Domest Anim Endocrinol* 29:3-22

- Conversi A, Piontkovski S, Hameed S (2001) Seasonal and interannual dynamics of *Calanus finmarchicus* in the Gulf of Maine (Northeastern US shelf) with reference to the North Atlantic Oscillation. *Deep-Sea Res Part II* 48:1-3
- Cooke JG, Rowntree VJ, Payne R (2001) Estimates of demographic parameters for southern right whales (*Eubalaena australis*) observed off Península Valdés, Argentina. *J Cetacean Res Manag (Special Issue)* 2:125-132
- Curran MP, Asher WM (1974) Investigation of blubber thickness in a gray whale using ultrasonography. *Mar Fish Rev* 36:15-20
- Devore JL (1995) Probability and Statistics for Engineering and the Sciences. Duxbury Press, NY 4th Edition.
- Doidge DW (1990) Integumentary heat loss and blubber distribution in the beluga, *Delphinapterus leucas*, with comparisons to the narwhal, *Monodon monoceros*. *Can J Fish Aquat Sci* 224:129-140
- Doney JM, Gunn RG, Horak F (1982) Reproduction. In: Coop IE (ed) Sheep and Goat Production. Elsevier Scientific, Amsterdam, p 57-80
- Dunkin RC, McLellan WA, Blum JE, Pabst D (2010) The buoyancy of the integument of Atlantic bottlenose dolphins (*Tursiops truncatus*): Effects of growth, reproduction, and nutritional state. *Mar Mamm Sci* 26:573-587
- Dunkin RC, McLellan WA, Blum JE, Pabst DA (2005) The ontogenetic changes in the thermal properties of blubber from Atlantic bottlenose dolphin *Tursiops truncatus*. *J Exp Biol* 208:1469-1480
- Foote RH (1978) Factors influencing the quantity and quality of semen harvested from bulls, rams, boars and stallions. *J Anim Sci* 47:1-11
- Frisch R (1997) Body weight, body fat and ovulation: relation to the natural fertility of populations. In: Dorfman R, Rogers P (eds) Science with a Human Face: In Honor of Roger Randall Revelle. Harvard University Press, Cambridge, p 139-166
- Frisch RE (1984) Body fat, puberty and fertility. *Biol Rev* 59:161-188
- Gaines CA, Hare MP, Beck SE, Rosenbaum HC (2005) Nuclear markers confirm taxonomic status and relationships among highly endangered and closely related right whale species. *Proc R Soc Lond B* 272:533-542
- Gales NJ, Burton HR (1987) Ultrasonic measurement of blubber thickness of the southern elephant seal, *Mirounga leonina* (Linn.). *Aust J Zool* 35:207-217
- Greene CH, Pershing AJ, Conversi A, Planque B and others (2003) Trans-Atlantic responses of *Calanus finmarchicus* populations to basin- scale forcing associated with the North Atlantic Oscillation. *Prog Oceanogr* 58:301-312
- Haldiman JT, Tarpley RJ (1993) Anatomy and Physiology. In: Burns JJ, Montague JJ, Cowles CJ (eds) The bowhead whale. Special publication number 2. The Society for Marine Mammalogy. Allen Press, Inc., New York, p 71-156
- Hamilton JL, Dillaman RM, McLellan WA, Pabst DA (2004) Structural fiber reinforcement of keel blubber in harbor porpoise (*Phocoena phocoena*). *J Morphol* (1931) 261:105-117
- Hamilton PK, Cooper LA (2010) Changes in North Atlantic Right Whale (*Eubalaena glacialis*) Cow-Calf Association Times and Use of the Calving Ground: 1993-2005. *Mar Mamm Sci* 26:896-916
- Hamilton PK, Knowlton AR, Marx MK (2007) Right whales tell their own stories: The Photo-Identification *Catalog*. In: Kraus SD, Rolland RM (eds) The urban whale: North Atlantic

right whales at the crossroads. Harvard University Press, Cambridge, Massachusetts, p 75-104

- Hamilton PK, Knowlton AR, Marx MK, Kraus SD (1998) Age structure and longevity in North Atlantic right whales *Eubalaena glacialis* and their relation to reproduction. *Mar Ecol Prog Ser* 171:285-292
- Hamilton PK, Marx MK, Kraus SD (1995) Weaning in North Atlantic right whales. *Mar Mamm Sci* 11:386-390
- Hamner WM, Stone GS, Obst BS (1988) Behavior of southern right whales, *Eubalaena australis*, feeding on the Antarctic krill, *Euphausia superba*. *Fish Bull* 86:143-150
- Heyerdahl E (1932) Hvalindustrien en tekniskjemisk undersøkelse. I. Råmaterialet. Kommandør Chr-Christensens Hvalfangstmuseum, Sandefjord Publ Kasjon Nr 7
- Hill JW, Elmquist JK, Elias CF (2008) Hypothalamic pathways linking energy balance and reproduction. *Am J Physiol Endocrin Metab* 294:E827-E832
- Hlista BL, Sosik HM, Traykovski LVM, Kenney RD, Moore MJ (2009) Seasonal and interannual correlations between right-whale distribution and calving success and chlorophyll concentrations in the Gulf of Maine, USA. *Mar Ecol Prog Ser* 394:289-302
- Houseknecht KL, Baile CA, Matteri RL, Spurlock ME (1998) The biology of leptin: a review. *J Anim Sci* 76:1405-1420
- International Whaling Commission (2001) Report of the workshop on the status and trends in western North Atlantic right whales. *J Cetacean Res Manag (Special Issue)* 2:61-87
- International Whaling Commission (2006) Report of the Scientific Committee. *J Cetacean Res Manag* 8:1-65
- Kenney RD, Hyman MAM, Owen RE, Scott GP, Winn HE (1986) Estimation of prey densities required by western North Atlantic right whales. *Mar Mamm Sci* 2:1-13
- Kipps EK, McLellan WA, Rommel SA, Pabst DA (2002) Skin density and its influence on buoyancy in the manatee (*Trichechus manatus latirostris*), harbor porpoise (*Phocoena phocoena*), and bottlenose dolphin (*Tursiops truncatus*). *Mar Mamm Sci* 18:765-778
- Knittle JL, Hirsch J (1968) Effect of early nutrition on the development of rat epididymal fat pads: cellularity and metabolism. *J Clin Investig* 47:2091-2098
- Knowlton AR, Kraus SD (2001) Mortality and serious injury of northern right whales (*Eubalaena glacialis*) in the western North Atlantic Ocean. *J Cetacean Res Manag (Special Issue)* 2:193-208
- Knowlton AR, Kraus SD, Kenney RD (1994) Reproduction in North Atlantic right whales (*Eubalaena glacialis*). *Can J Zool* 72:1297-1305
- Koopman HN (1998) Topographical distribution of the blubber of harbor porpoises (*Phocoena phocoena*). *J Mammal* 79:260-270
- Koopman HN (2007) Phylogenetic, ecological, and ontogenetic factors influencing the biochemical structure of the blubber of odontocetes. *Mar Biol* 151:277-291
- Koopman HN, Pabst DA, McLellan WA, Dillaman RM, Read AJ (2002) Changes in blubber distribution and morphology associated with starvation in the harbor porpoise (*Phocoena phocoena*): evidence for regional differences in blubber structure and function. *Physiol Biochem Zool* 75:498-512
- Kraus SD, Brown MW, Caswell H, Clark CW and others (2005) North Atlantic right whales in crisis. *Science* 309:561-562
- Kraus SD, Hamilton PK, Kenney RD, Knowlton A, Slay C (2001) Reproductive parameters of the North Atlantic right whale. *J Cetacean Res Manag (Special Issue)* 2:231-236

- Kraus SD, Hatch JJ (2001) Mating strategies in the North Atlantic right whale (*Eubalaena glacialis*). J Cetacean Res Manag (Special Issue) 2:237-244
- Kraus SD, Pace RM, Frasier TR (2007) High investment, low return: the strange case of reproduction in *Eubalaena glacialis*. In: Kraus SD, Rolland RM (eds) The urban whale: North Atlantic right whales at the crossroads. Harvard University Press, Cambridge, Massachusetts, p 172-199
- Ling JK (1974) The Integument of Marine Mammals. In: Harrison RJ (ed) Functional Anatomy of Marine Mammals, Vol 2. Academic Press, Inc., New York, p 1-44
- Lockyer C (1981) Growth and energy budgets of large baleen whales from the southern hemisphere. In: Mammals in the Seas; general papers and large cetaceans, Vol 3. Food and Agriculture Organization of the United Nations, Rome, p 379-488
- Lockyer C (1986) Body fat condition in Northeast Atlantic fin whales, *Balaenoptera physalus*, and its relationship with reproduction and food resource. Can J Fish Aquat Sci 43:142-147
- Lockyer C (1987) The relationship between body fat, food resource and reproductive energy costs in north Atlantic fin whales (*Balaenoptera physalus*). Symp Zool Soc Lond 57:343-361
- Lockyer C (1991) Body composition of the sperm whale, *Physeter catodon*, with special reference to the possible functions of fat depots. Rit Fiskideildar 12:1-24
- Lockyer CH, McConnell LC, Waters TD (1984) The biochemical composition of fin whale blubber. Can J Zool 62:2553-2562
- Lockyer CH, McConnell LC, Waters TD (1985) Body condition in terms of anatomical and biochemical assessment of body fat in North Atlantic fin and sei whales. Can J Zool 63:2328-2338
- Lubetkin SC, Zeh JE, Rosa C, George JC (2008) Age estimation for young bowhead whales (*Balaena mysticetus*) using annual baleen growth increments. Can J Zool 86:525-538
- McEvoy TG, Robinson JJ (2003) Nutrition and its interaction with reproductive processes. In: Holt WV, Pickard AR, Rodger JC, Wildt DE (eds) Reproductive Science and Integrated Conservation. Cambridge University Press, Cambridge, p 42-56
- Michaud J, Taggart CT (2007) Lipid and gross energy content of North Atlantic right whale food, *Calanus finmarchicus*, in the Bay of Fundy. Endang Species Res 3:77-94
- Miller Angell C (2006) Body fat condition of free-ranging right whales, *Eubalaena glacialis* and *Eubalaena australis*. PhD dissertation, Boston University
- Miller CA, Perryman WL, Best PB, Baumgartner MF, Moore MJ (In Review) Body shape changes associated with reproductive status, body fat condition and growth in free-ranging right whales *Eubalaena glacialis* and *Eubalaena australis*. Mar Ecol Prog Ser
- Moore MJ, Knowlton AR, Kraus SD, McLellan WA, Bonde RK (2004) Morphometry, gross morphology and available histopathology in North Atlantic right whale (*Eubalaena glacialis*) mortalities (1970 to 2002). J Cetacean Res Manag 6:199-214
- Moore MJ, Miller CA, Morss MS, Arthur R and others (2001) Ultrasonic measurement of blubber thickness in right whales. J Cetacean Res Manag (Special Issue) 2:301-309
- Nishiwaki M, Hayashi K (1950) Biological survey of fin and blue whales taken in the Antarctic season 1947-48 by the Japanese fleet. Sci Rep Whales Res Inst Tokyo 3:132-190
- Nishiwaki M, Oye T (1951) Biological investigation on blue whales (*Balaenoptera sic musculus*) and fin whales (*Balaenoptera physalus*) caught by the Japanese Antarctic whaling fleets, July 1948. Sci Rep Whales Res Inst Tokyo 5:91-167

- NOAA (2009) Stock Assessment Report: North Atlantic right whale (*Eubalaena glacialis*): Western Atlantic stock. <http://www.nmfs.noaa.gov/pr/pdfs/sars/ao2009whnr-w.pdf>
- Omura H, Oshumi S, Nemoto T, Nasu K, Kasuya T (1969) Black right whales in the North Pacific. *Sci Rep Whales Res Inst Tokyo* 21:1-78
- Pabst DA (1996) Morphology of the subdermal connective tissue sheath of dolphins: A new fibre-wound, thin-walled, pressurized cylinder model for swimming vertebrates. *J Zool Lond* 238:35-52
- Pershing AJ, Greene CH, Jossi JW, O'Brien L, Brodziak JKT, Bailey BA (2005) Interdecadal variability in the Gulf of Maine zooplankton community, with potential impacts on fish recruitment. *ICES J Mar Sci* 62:1511-1523
- Pershing AJ, Head EHJ, Greene CH, Jossi JW (2010) Pattern and scale of variability among Northwest Atlantic Shelf plankton communities. *J Plankton Res* 32:1661-1674
- Pettis HM, Rolland RM, Hamilton PK, Brault S, Knowlton AR, Kraus SD (2004) Visual health assessment of North Atlantic right whales (*Eubalaena glacialis*) using photographs. *Can J Zool* 82:8-19
- Pond CM (1977) The significance of lactation in the evolution of mammals. *Evolution* 31:177-199
- Pond CM, Mattacks CA (1988) The distribution, cellular structure, and metabolism of adipose tissue in the fin whale, *Balaenoptera physalus*. *Can J Zool* 66:534-537
- Prunier A, Quesnel H (2000) Nutritional influences on the hormonal control of reproduction in female pigs. *Livestock Production Science* 63:1-16
- Reeb D (2001) Structure, development and composition of the integument of the southern right whale, *Eubalaena australis*. *Philosophiae Doctor, University of Pretoria*
- Reeb D, Best PB, Kidson SH (2007) Structure of the integument of southern right whales, *Eubalaena australis*. *Anat Rec* 290:596-613
- Reilly SB, Bannister JL, Best PB, Brown M and others (2008a) *Eubalaena glacialis* 2008 IUCN Red List of Threatened Species
- Reilly SB, Bannister JL, Best PB, Brown M and others (2008b) *Eubalaena australis* 2008 IUCN Red List of Threatened Species
- Rice DW, Wolman AA (1971) The life history and ecology of the gray whale (*Eschrichtius robustus*). Special Publication No. 3. American Society of Mammalogists, Stillwater, Oklahoma
- Rosenbaum HC, Brownell LR, Brown WM, Schaeff C and others (2000) World-wide genetic differentiation of *Eubalaena*: questioning the number of right whale species. *Mol Ecol* 9:1793-1802
- Simpson JG, Gardiner MB (1972) Comparative microscopic anatomy of selected marine mammals. In: Ridgway SH (ed) *Mammals of the Sea - Biology and Medicine*. Charles C. Thomas, Springfield, IL, p 363-378
- Thomas VG (1990) Control of reproduction in animal species with high and low body fat reserves. In: Frisch R (ed) *Adipose Tissue and Reproduction*, Vol 14. Karger, Basel, p 27-41
- Tormosov DD, Mikhaliev YA, Best PB, Zemsky VA, Sekiguchi K, Brownell RL, Jr. (1998) Soviet catches of southern right whales *Eubalaena australis*, 1951-1971. Biological data and conservation implications. *Biol Conserv* 86:185-197



- Tummaruk P, Lundeheim N, Einarsson S, Dalin AM (2001) Effect of birth litter size, birth parity number, growth rate, backfat thickness and age at first mating of gilts on their reproductive performance as sows. *Anim Reprod Sci* 66:225-237
- Young RA (1976) Fat, energy and mammalian survival. *Am Zool* 16:699-710
- Zieba DA, Amstalden M, Williams GL (2005) Regulatory roles of leptin in reproduction and metabolism: a comparative review. *Domest Anim Endocrinol* 29:166-185

## FIGURE LEGENDS:

Figure 1. *Eubalaena glacialis*. Measurements of blubber thickness (epidermis and blubber) collected with a ruler during necropsies from positions along the dorsal midline of four adult females and one adult male. *E.g.* #s refer to individual whale identification numbers in the North Atlantic Right Whale Catalog (Hamilton et al. 2007).

Figure 2. *Eubalaena glacialis*. Acoustic measurements of blubber thickness (epidermis and blubber) compared with photogrammetric measurements of total body length (snout to fluke notch) Bay of Fundy, Canada, August 2000-2002. The solid line indicating the relationship between total length and blubber thickness found in juveniles is extrapolated on both ends with a lighter line.

Figure 3. *Eubalaena glacialis*. (a) Acoustic blubber thickness measurements from calves and juveniles (randomly jittered with respect to age with mean  $\pm 1$  standard deviation) between ages 0 and 8 years old, Bay of Fundy, Canada, 1998-2002. Significance, indicated by letters, was determined at  $p < 0.05$ ; (b) Blubber thickness of three individuals sampled as 8 month old calves and again, 12 months later, as yearlings, Bay of Fundy, Canada, 2001-2002. *E.g.* #s refer to individual whale identification numbers in the North Atlantic Right Whale Catalog (Hamilton et al. 2007); (c) Acoustic blubber thickness measurements as a percentage of total body length for calves and yearlings (randomly jittered with respect to age with mean  $\pm 1$  standard deviation), Bay of Fundy, Canada, 2001-2002.

Figure 4. *Eubalaena australis*. Blubber thickness measurements (randomly jittered) collected by ultrasound from (a) calves and (b) mothers during early (September 1999) and mid (November 2000) lactation with mean  $\pm 1$ SD, St. Sebastian Bay, De Hoop Marine Reserve and Walker Bay (Gansbaai), South Africa 1999 – 2000. Significance was determined at  $p < 0.05$  and indicated by asterisks.

Figure 5. *Eubalaena glacialis*. Mean blubber thickness  $\pm 1$  SD ( $N$ ) measured with ultrasound from adult male and yearling right whales, Bay of Fundy, Canada, 1998-2002. Significance, determined at  $p < 0.05$ , is denoted by letters as marked.

*Calanus finmarchicus*. Yearly means of the differences between observed log-transformed abundance data (measured by Continuous Plankton Recorder, Gulf of Maine) and expected abundance (climatological seasonal cycle) (Pershing et al. 2005). Mean of the yearly abundance deviations from 1961-1974 and 1978-2003 plotted as dotted line for reference to the entire Pershing et al. (2005) time-series.

Figure 6. *Eubalaena glacialis*. Mean blubber thickness  $+ 1$ SD ( $N$ ) measured with ultrasound from adult females of different reproductive status, adult males, juveniles and calves, Bay of Fundy and Cape Cod Bay 1998-2002. Significance was determined at  $p < 0.05$ . Mean values marked with different letters were statistically different.

Figure 7. *Eubalaena glacialis*. Blubber thickness collected by ultrasound from four individual females measured in multiple years at different stages of the reproductive cycle, Bay of Fundy,

Canada 1998 – 2002. *E.g.* #s refer to individual whale identification numbers in the North Atlantic Right Whale Catalog (Hamilton et al. 2007).

Figure 8. *Eubalaena glacialis*. Blubber thickness and number of years since last calf in reproductively active adult females, Bay of Fundy, Canada 1998 – 2002. Data points labeled with year measured, number of years to subsequent calving and reproductive status if pregnant (P) or pre-pregnant (PP).

Figure 9. *Eubalaena glacialis* (+) and *Eubalaena australis* (□). Blubber thickness measurements (randomly jittered with mean  $\pm$  1 SD) collected by ultrasound from (a) all whales; (b) adult females; (c) juveniles; (d) calves; Bay of Fundy, Canada 1998 – 2002 and St. Sebastian Bay, De Hoop Marine Reserve and Walker Bay (Gansbaai), South Africa 1999 – 2000. Asterisks indicate a significant difference between means within each age class at  $\alpha = 0.05$ . Note: the *E. glacialis* calves were eight months old while *E. australis* calves were neonate to four months old.

Figure 1





Figure 2.



- ◇ Calf
- Juvenile
- ⊕ Male - Unknown Age
- × Female - Unknown Age
- Adult Female (lactating)
- \* Adult Female (never been observed with a calf in her 26 year sighting history)
- Juvenile & Unknown Age Regression

Figure 3.




Figure 4.





Figure 5.



□ Adult Males

■ Yearlings

—×— Calanus Abundance Index (Pershing et al. 2005)

..... Mean of Calanus Abundance Index for 1961-1974 and 1978-2003 (Pershing et al. 2005)



Figure 6.





Figure 7.



- R = Resting; N = Nulliparous; I = Immature
- Pre-Pregnant
- Lactating

Figure 8.




Figure 9.

