



## ONDERSTEPSOORT JOURNAL OF VETERINARY RESEARCH

Volume 69 • Numbers 1-4 • 2002

### Subject index

#### Acarl: Ixodidae

- Hosts, seasonal occurrence and life cycle of *Rhipicephalus nuttalli* (Acarl: Ixodidae)—L.J. FOURIE, I.G. HORAK, D.J. KOK and W. VAN ZYL ..... 177-187

#### Acaricide

- Acaricide resistance profiles of single- and multi-host ticks from communal and commercial farming areas in the Eastern Cape and North-West Provinces of South Africa—S. MEKONNEN, N.R. BRYSON, L.J. FOURIE, R.J. PETER, A.M. SPICKETT, R.J. TAYLOR, T. STRYDOM and I.G. HORAK ..... 99-105

#### African buffaloes

- The gamma-interferon test, its usefulness in a bovine tuberculosis survey in African buffaloes (*Syncerus caffer*) in the Kruger National Park—D.G. GROBLER, ANITA L. MICHEL, LIN-MARI DE KLERK and R.G. BENGIS ..... 221-227

#### Albendazole

- A survey on the occurrence of resistance to anthelmintics of gastrointestinal nematodes of goats in Mozambique—A. ATANÁSIO, J. BOOMKER and C. SITOE ..... 215-220

- Strategic control of gastrointestinal nematodes of sheep in the highlands of central Kenya—N. MAINGI, R.O. OTIENO, V.M. MUMBI and E.H. WEDA ..... 229-235

#### Amphistome infection

- Amphistome infection of goats farmed under resource-poor conditions in South Africa—A.F. VATTA and R.C. KRECEK ..... 327-329

#### Anaplasmosis

- Occurrence of concurrent trypanosomosis, theileriosis, anaplasmosis and helminthosis in Friesian, Zebu and Sahi-wal cattle in Uganda—J.W. MAGONA and J.S.P. MAYENDE ..... 133-140

#### Animal health

- Review of current problems and shortcomings in the Tanzanian animal health information system with suggestions on improvement—F.M. KIVARIA and A.M. KAPAGA ..... 305-314

#### Anthelmintic resistance

- A survey on the occurrence of resistance to anthelmintics of gastrointestinal nematodes of goats in Mozambique—A. ATANÁSIO, J. BOOMKER and C. SITOE ..... 215-220

#### Antigen preparation

- Development of solid phase antigen for indirect ELISA for the detection of specific antibody responses to infection with Newcastle disease virus—M. SAHLE and G.W. BURGESS ..... 237-242

#### Arbovirus

- Seasonal abundance of four *Culicoides* spp. (Diptera: Ceratopogonidae) at Al-Ahsa oasis, Eastern Province, Saudi

## Subject index

|  |         |
|--|---------|
| Arabia—E.M.E. ABU ELZEIN, M.A. HILALI, A.I. AL-AFALEQ, P.S. MELLOR, J. BOORMAN, S. AL-ATIYA and A AL NAIEM .....   | 115–122 |
| <b>Atelerix frontalis</b>  |         |
| Hosts, seasonal occurrence and life cycle of <i>Rhipicentor nuttalli</i> (Acari, Ixodidae)—L.J. FOURIE, I.G. HORAK, D.J. KOK and W. VAN ZYL .....  | 177–187 |
| <b>Babesia caballi</b>   |         |
| <i>In vitro</i> isolation of equine piroplasms derived from Cape mountain zebra ( <i>Equus zebra zebra</i> ) in South Africa—E. ZWEYGARTH, LAURA M. LOPEZ-REBOLLAR and P. MEYER .....  | 197–200 |
| <b>Boophilus decoloratus</b>   |         |
| Seasonal abundance of adult ixodid ticks infesting cattle belonging to resource-limited farmers in the north-eastern Free State Province of South Africa—M. HLATSHWAYO, P.A. MBATI and O.O. DIPEOLU .....  | 1–6     |
| <b>Bovine fasciolosis</b>  |         |
| The prevalence and economic importance of bovine fasciolosis in Kenya—an analysis of abattoir data—J.M. KI-THUKA, N. MAINGI, F.M. NJERUH and J.N. OMBUI .....  | 255–262 |
| <b>Bovine tuberculosis</b>   |         |
| The gamma-interferon test: its usefulness in a bovine tuberculosis survey in African buffaloes ( <i>Syncerus caffer</i> ) in the Kruger National Park—D.G. GROBLER, ANITA L. MICHEL, LIN-MARI DE KLERK and R.G. BENGIS .....   | 221–227 |
| <b>Brucella abortus</b>  |         |
| Validation of an indirect enzyme-linked immunosorbent assay for the detection of antibody against <i>Brucella abortus</i> in cattle sera using an automated ELISA workstation—J.T. PAWESKA, A.D. POTTS, H.J. HARRIS, S.J. SMITH, G.J. VILJOEN, B. DUNGU, O.L. BRETT, M. BUBB and L. PROZESKY ..... | 61–77   |
| <b>Brucella melitensis</b>   |         |
| An overview of the eradication of <i>Brucella melitensis</i> from KwaZulu-Natal—F.R. EMSLIE and J.R. NEL .....   | 123–127 |
| <b>Bufadienolide</b>   |         |
| The effect of three bufadienolide cardiac glycosides on contraction of isolated rat jejunum—C.J. BOTHA, R. GEHRING, J.M. VAN ROOYEN and D. VENTER .....  | 243–246 |
| <b>Camels</b>  |         |
| Animal-level risk factors for <i>Trypanosoma evansi</i> infection in camels in eastern and central parts of Kenya—J.M. NGAIRA, B. BETT and S.M. KARANJA .....  | 263–271 |
| <b>Cape mountain zebra</b>   |         |
| <i>In vitro</i> isolation of equine piroplasms derived from Cape mountain zebra ( <i>Equus zebra zebra</i> ) in South Africa—E. ZWEYGARTH, LAURA M. LOPEZ-REBOLLAR and P. MEYER .....  | 197–200 |
| <b>Cardiac glycosides</b>  |         |
| The effect of three bufadienolide cardiac glycosides on contraction of isolated rat jejunum—C.J. BOTHA, R. GEHRING, J.M. VAN ROOYEN and D. VENTER .....  | 243–246 |
| <b>Cattle</b>  |         |
| Seasonal abundance of adult ixodid ticks infesting cattle belonging to resource-limited farmers in the north-eastern Free State Province of South Africa—M. HLATSHWAYO, P.A. MBATI and O.O. DIPEOLU .....  | 1–6     |
| Validation of an indirect enzyme-linked immunosorbent assay for the detection of antibody against <i>Brucella abortus</i> in cattle sera using an automated ELISA workstation—J.T. PAWESKA, A.D. POTTS, H.J. HARRIS, S.J. SMITH, G.J. VILJOEN, B. DUNGU, O.L. BRETT, M. BUBB and L. PROZESKY ..... | 61–77   |
| Parasitological prevalence of bovine trypanosomosis in Kindo Koisha district, Wolloita zone, south Ethiopia—A. KIDANE MARIAM, K. HADGU and M. SAHLE .....  | 107–113 |
| Occurrence of concurrent trypanosomosis, theileriosis, anaplasmosis and helminthosis in Friesian, Zebu and Sahi-wal cattle in Uganda—J.W. MAGONA and J.S.P. MAYENDE .....  | 133–140 |
| Seasonal availability of gastrointestinal nematode larvae to cattle on pasture in the central highlands of Kenya—R.M. WARUIRU, E.H. WEDA, R.O. OTIENO and J.W. NGOCHO .....  | 141–146 |
| Non-immune control of trypanosomosis: <i>In vitro</i> oxidative burst of PMA- and trypanosome-stimulated neutrophils of Boran and N'Dama cattle—V.O. TAIWO, J.O. ADEJINMI and J.O. OL UWANIYI .....  | 155–161 |

|   |         |
|---|---------|
| The prevalence and economic importance of bovine fasciolosis in Kenya—an analysis of abattoir data—J.M. KITHUKA, N. MAINGI, F.M. NJERUH and J.N. OMBUI . . . . .  | 255-262 |
| <b>Chelonia</b>   |         |
| Description of <i>Pelonia alcana</i> n. g., n. sp. (Pentastomida: Sebekidae) from the lungs of <i>Pelomedusa subrufa</i> and <i>Pelusios sinuatus</i> (Chelonia) in South Africa—K. JUNKER and J. BOOMKER . . . . .                             | 53-59   |
| <b>Closantel</b>  |         |
| Strategic control of gastrointestinal nematodes of sheep in the highlands of central Kenya—N. MAINGI, R.O. OTIENO, V.M. MUMBI and E.H. WEDA . . . . .   | 229-235 |
| <b>Colon</b>  |         |
| Water absorption in relation to fermentation in the colon of the ostrich ( <i>Struthio camelus</i> )—C. MUSARA, J.P. CHAMUNORWA, K. HOLTUG and E. SKADHAUGE . . . . .   | 315-320 |
| <b>Complement fixation test</b>   |         |
| Apparent prevalence of dourine in the Khomas region of Namibia—F.F. KUMBA, B. CLAASEN and P. PETRUS . . . . .   | 295-298 |
| <b>Coryza</b>   |         |
| Isolation of serovar C-3 <i>Haemophilus paragallinarum</i> from Zimbabwe: A further indication of the need for the production of vaccines against infectious coryza containing local isolates of <i>H. paragallinarum</i> —R.R. BRAGG . . . . . | 129-132 |
| Virulence of South African isolates of <i>Haemophilus paragallinarum</i> . Part 3: experimentally produced NAD-independent isolate—M. TAOLE, J. ALBERTYN, E. VAN HEERDEN and R.R. BRAGG . . . . .   | 189-196 |
| <b>Cotyledoside</b>   |         |
| The effect of three bufadienolide cardiac glycosides on contraction of isolated rat jejunum—C.J. BOTHA, R. GEHRING, J.M. VAN ROOYEN and D. VENTER . . . . .   | 243-246 |
| <b>Cryopreservation</b>   |         |
| Freezing of sheep faeces invalidates <i>Haemonchus contortus</i> faecal egg counts by the McMaster technique—J.A. VAN WYK and LAETITIA VAN WYK . . . . .  | 299-304 |
| <b>Culicoides</b>   |         |
| Seasonal abundance of four <i>Culicoides</i> spp. (Diptera: Ceratopogonidae) at Al-Ahsa oasis, Eastern Province, Saudi Arabia—E.M.E. ABU ELZEIN, M.A. HILALI, A.I. AL-AFAEQ, P.S. MELLOR, J. BOORMAN, S. AL-ATIYA and A. AL NAIEM . . . . .     | 115-122 |
| <b>Cycloheximide</b>  |         |
| The Kümm isolate of <i>Ehrlichia ruminantium</i> . In vitro isolation, propagation and characterization—E. ZWEYGARTH, ANTOINETTE I. JOSEMANS, M. FRANSI VAN STRIJP, HENRIETTE VAN HEERDEN, MARIA T.E.P. ALLSOPP and B.A. ALLSOPP . . . . .      | 147-153 |
| <b>Diptera: Ceratopogonidae</b>   |         |
| Seasonal abundance of four <i>Culicoides</i> spp. (Diptera: Ceratopogonidae) at Al-Ahsa oasis, Eastern Province, Saudi Arabia—E.M.E. ABU ELZEIN, M.A. HILALI, A.I. AL-AFAEQ, P.S. MELLOR, J. BOORMAN, S. AL-ATIYA and A. AL NAIEM . . . . .     | 115-122 |
| <b>Dourine</b>  |         |
| Apparent prevalence of dourine in the Khomas region of Namibia—F.F. KUMBA, B. CLAASEN and P. PETRUS . . . . .   | 295-298 |
| <b><i>Ehrlichia (Cowdria) ruminantium</i></b>   |         |
| The Kümm isolate of <i>Ehrlichia ruminantium</i> : In vitro isolation, propagation and characterization—E. ZWEYGARTH, ANTOINETTE I. JOSEMANS, M. FRANSI VAN STRIJP, HENRIETTE VAN HEERDEN, MARIA T.E.P. ALLSOPP and B.A. ALLSOPP . . . . .      | 147-153 |
| <b><i>Elephantulus myurus</i></b>   |         |
| Hosts, seasonal occurrence and life cycle of <i>Rhipicephalus nuttalli</i> (Acar: Ixodidae)—L.J. FOURIE, I.G. HORAK, D.J. KOK and W. VAN ZYL . . . . .  | 177-187 |
| <b><i>Epoxyssiliroldin</i></b>  |         |
| The effect of three bufadienolide cardiac glycosides on contraction of isolated rat jejunum—C.J. BOTHA, R. GEHRING, J.M. VAN ROOYEN and D. VENTER . . . . .   | 243-246 |

## Subject index

### Equine encephalosis virus

- The classification of seven serotypes of equine encephalosis virus and the prevalence of homologous antibody in horses in South Africa—P.G. HOWELL, DALEEN GROENEWALD, CARINA W. VISAGE, ANNA-MARI BOSMAN, J.A.W. COETZER and A.J. GUTHRIE ..... 79-93

### Equine piroplasmosis

- In vitro* isolation of equine piroplasms derived from Cape mountain zebra (*Equus zebra zebra*) in South Africa—E. ZWEYGARTH, LAURA M. LOPEZ-REBOLLAR and P. MEYER ..... 197-200

### *Equus zebra zebra*

- In vitro* isolation of equine piroplasms derived from Cape mountain zebra (*Equus zebra zebra*) in South Africa—E. ZWEYGARTH, LAURA M. LOPEZ-REBOLLAR and P. MEYER ..... 197-200

### Ethiopia

- Parasitological prevalence of bovine trypanosomosis in Kindo Koisha district, Wollaia zone, south Ethiopia—A. KIDANEMARIAM, K. HADGU and M. SAHLE ..... 107-113

### Faecal egg counts

- Freezing of sheep faeces invalidates *Haemonchus contortus* faecal egg counts by the McMaster technique—J.A. VAN WYK and LAETITIA VAN WYK ..... 299-304

- Amphistome infection of goats farmed under resource-poor conditions in South Africa—A.F. VATTA and R.C. KRECEK ..... 327-329

### *Fasciola*

- The prevalence and economic importance of bovine fasciolosis in Kenya—an analysis of abattoir data—J.M. KITHUKA, N. MAINGI, F.M. NJERUH and J.N. OMBUI ..... 255-262

- Amphistome infection of goats farmed under resource-poor conditions in South Africa—A.F. VATTA and R.C. KRECEK ..... 327-329

### Feed aversion

- Continuous exposure to an aversive mixture as a means of maintaining aversion to vermeerbos (*Geigeria ornativa* O. Hoffm.) in the presence of non-averted sheep—L.D. SNYMAN, R.A. SCHULTZ, T.S. KELLERMAN and L. LABU-SCHAGNE ..... 321-325

### Fenbendazole

- A survey on the occurrence of resistance to anthelmintics of gastrointestinal nematodes of goats in Mozambique—A. ATANÁSIO, J. BOOMKER and C. SITOE ..... 215-220

### Gamma-Interferon test

- The gamma-interferon test: its usefulness in a bovine tuberculosis survey in African buffaloes (*Synacerus caffer*) in the Kruger National Park—D.G. GROBLER, ANITA L. MICHEL, LIN-MARI DE KLERK and R.G. BENGIS ..... 221-227

### *Geigeria ornativa* (vermeerbos)

- Continuous exposure to an aversive mixture as a means of maintaining aversion to vermeerbos (*Geigeria ornativa* O. Hoffm.) in the presence of non-averted sheep—L.D. SNYMAN, R.A. SCHULTZ, T.S. KELLERMAN and L. LABU-SCHAGNE ..... 321-325

### Gekkonidae

- Redescription of some *Spauligodon* spp. and *Parapharyngodon* spp., and of *Skriabinodon mabuyaee* Sandground, 1936 (Inglis, 1968) (Pharyngodonidae: Oxyuroidea) from insectivorous South African lizards—S.F.B.N. HERING-HAGENBECK, A.J. PETTER and J. BOOMKER ..... 7-29

### Gerrhosauridae

- Redescription of some *Spauligodon* spp. and *Parapharyngodon* spp., and of *Skriabinodon mabuyaee* Sandground, 1936 (Inglis, 1968) (Pharyngodonidae: Oxyuroidea) from insectivorous South African lizards—S.F.B.N. HERING-HAGENBECK, A.J. PETTER and J. BOOMKER ..... 7-29

- Redescription of some *Thelandsos* and *Tachygoneira* spp. (Pharyngodonidae: Oxyuroidea) from the omnivorous plated lizard, *Gerrhosaurus validus validus* A. Smith, 1849 in South Africa—S.F.B.N. HERING-HAGENBECK, A.J. PETTER and J. BOOMKER ..... 31-51

### Goats

- An overview of the eradication of *Brucella melitensis* from KwaZulu-Natal—F.R. EMSLIE and J.R. NEL ..... 123-127

|  |         |
|--|---------|
| A survey on the occurrence of resistance to anthelmintics of gastrointestinal nematodes of goats in Mozambique—A. ATANÁSIO, J. BOOMKER and C. SITOÉ .....  | 215–220 |
| Amphistome infection of goats farmed under resource-poor conditions in South Africa—A.F. VATTA and R.C. KRECEK .....   | 327–329 |
| <b>Haematology</b>   |         |
| Effects of the scab mite <i>Psoroptes ovis</i> on the haematology and live mass of Merino and Dorper sheep—T. MEINTJES, L.J. FOURIE and I.G. HORAK .....   | 285–293 |
| <b><i>Haemonchus contortus</i></b>   |         |
| Freezing of sheep faeces invalidates <i>Haemonchus contortus</i> faecal egg counts by the McMaster technique—J.A. VAN WYK and LAETITIA VAN WYK .....   | 299–304 |
| <b><i>Haemophilus paragallinarum</i></b>   |         |
| Isolation of serovar C-3 <i>Haemophilus paragallinarum</i> from Zimbabwe: A further indication of the need for the production of vaccines against infectious coryza containing local isolates of <i>H. paragallinarum</i> —R.R. BRAGG .....  | 129–132 |
| Virulence of South African isolates of <i>Haemophilus paragallinarum</i> . Part 1: NAD-dependent field isolates—R.R. BRAGG .....   | 163–169 |
| Virulence of South African isolates of <i>Haemophilus paragallinarum</i> . Part 2: Naturally occurring NAD-independent field isolates—R.R. BRAGG .....   | 171–175 |
| Virulence of South African isolates of <i>Haemophilus paragallinarum</i> . Part 3: experimentally produced NAD-independent isolate—M. TAOLE, J. ALBERTYN, E. VAN HEERDEN and R.R. BRAGG .....  | 189–196 |
| <b>Heartwater</b>  |         |
| The Kümm isolate of <i>Ehrlichia ruminantium</i> : <i>In vitro</i> isolation, propagation and characterization—E. ZWEYGARTH, ANTOINETTE I. JOSEMANS, M. FRANS VAN STRIJP, HENRIETTE VAN HEERDEN, MARIA T.E.P. ALLSOPP and B.A. ALLSOPP .....   | 147–153 |
| <b>Helminthosis</b>  |         |
| Occurrence of concurrent trypanosomosis, theileriosis, anaplasmosis and helminthosis in Friesian, Zebu and Sahi-wal cattle in Uganda—J.W. MAGONA and J.S.P. MAYENDE .....  | 133–140 |
| Seasonal availability of gastrointestinal nematode larvae to cattle on pasture in the central highlands of Kenya—R.M. WARUIRU, E.H. WEDA, R.O. OTIENO and J.W. NGOCHO .....  | 141–146 |
| <b>Hippopotami</b>   |         |
| Hepatic and renal concentrations of copper and other trace elements in hippopotami ( <i>Hippopotamus amphibius</i> L.) living in and adjacent to the Kafue and Luangwa Rivers in Zambia—M. MWASE, B. ALMLI, T. SVERTSEN, M.M. MUSONDA and A. FLÅØYEN .....   | 207–214 |
| <b>Horses</b>  |         |
| The classification of seven serotypes of equine encephalitis virus and the prevalence of homologous antibody in horses in South Africa—P.G. HOWELL, DALEEN GROENEWALD, CARINA W. VISAGE, ANNA-MARI BOSMAN, J.A.W. COETZER and A.J. GUTHRIE .....   | 79–93   |
| Apparent prevalence of dourine in the Khomas region of Namibia—F.F. KUMBA, B. CLAASEN and P. PETRUS .....  | 295–298 |
| <b>Indirect ELISA</b>  |         |
| Validation of an indirect enzyme-linked immunosorbent assay for the detection of antibody against <i>Brucella abortus</i> in cattle sera using an automated ELISA workstation—J.T. PAWESKA, A.D. POTTS, H.J. HARRIS, S.J. SMITH, G.J. VILJOEN, B. DUNGU, O.L. BRETT, M. BUBB and L. PROZESKY ..... | 61–77   |
| Development of solid phase antigen for indirect ELISA for the detection of specific antibody responses to infection with Newcastle disease virus—M. SAHLE and G.W. BURGESS .....   | 237–242 |
| <b>Information system</b>  |         |
| Review of current problems and shortcomings in the Tanzanian animal health information system with suggestions on improvement—F.M. KIVARIA and A.M. KAPAGA .....   | 305–314 |
| <b>Kenya</b>   |         |
| Seasonal availability of gastrointestinal nematode larvae to cattle on pasture in the central highlands of Kenya—R.M. WARUIRU, E.H. WEDA, R.O. OTIENO and J.W. NGOCHO .....  | 141–146 |
| Strategic control of gastrointestinal nematodes of sheep in the highlands of central Kenya—N. MAINGI, R.O. OTIENO, V.M. MUMBII and E.H. WEDA .....   | 229–235 |

## Subject index

|  |         |
|--|---------|
| The prevalence and economic importance of bovine fasciolosis in Kenya—an analysis of abattoir data—J.M. KITHUKA, N. MAINGI, F.M. NJERUH and J.N. OMBUI . . . . .   | 255–262 |
| Animal-level risk factors for <i>Trypanosoma evansi</i> infection in camels in eastern and central parts of Kenya—J.M. NGAIRA, B. BETT and S.M. KARANJA . . . . .  | 263–271 |
| <b>Kruger National Park</b>  |         |
| The gamma-interferon test: its usefulness in a bovine tuberculosis survey in African buffaloes ( <i>Syncerus caffer</i> ) in the Kruger National Park—D.G. GROBLER, ANITA L. MICHEL, LIN-MARI DE KLERK and R.G. BENGIS . . . . .   | 221–227 |
| <b>Kümm Isolate</b>  |         |
| The Kümm isolate of <i>Ehrlichia ruminantium</i> : <i>In vitro</i> isolation, propagation and characterization—E. ZWEYGARTH, ANTOINETTE I. JOSEMANS, M. FRANSI VAN STRIJP, HENRIETTE VAN HEERDEN, MARIA T.E.P. ALLSOPP and B.A. ALLSOPP . . . . .                          | 147–153 |
| <b>KwaZulu-Natal</b>   |         |
| An overview of the eradication of <i>Brucella melitensis</i> from KwaZulu-Natal—F.R. EMSLIE and J.R. NEL . . . . .   | 123–127 |
| <b>Levamisole</b>  |         |
| A survey on the occurrence of resistance to anthelmintics of gastrointestinal nematodes of goats in Mozambique—A. ATANÁSIO, J. BOOMKER and C. SITOE . . . . .  | 215–220 |
| <b>Lizards</b>   |         |
| Redescription of some <i>Spauligodon</i> spp. and <i>Parapharyngodon</i> spp., and of <i>Skjababinodon mabuyae</i> Sandground, 1936 (Pharyngodonidae: Oxyuroidea) from insectivorous South African lizards—S.F.B.N. HERING-HAGENBECK, A.J. PETTER and J. BOOMKER . . . . . | 7–29    |
| Redescription of some <i>Thelandon</i> and <i>Tachygonaetria</i> spp. (Pharyngodonidae: Oxyuroidea) from the omnivorous plated lizard, <i>Gerrhosaurus validus validus</i> A. Smith, 1849 in South Africa—S.F.B.N. HERING-HAGENBECK, A.J. PETTER and J. BOOMKER . . . . .  | 31–51   |
| <b>Madin Derby bovine kidney cells (MDBK)</b>  |         |
| Multiplication of the V4 strain of Newcastle disease virus in Madin Derby bovine kidney cells—M. SAHLE, G.W. BURGESS and A. KIDANEMARIAM . . . . .   | 201–206 |
| <b>McMaster technique</b>  |         |
| Freezing of sheep faeces invalidates <i>Haemonchus contortus</i> faecal egg counts by the McMaster technique—J.A. VAN WYK and LAETITIA VAN WYK . . . . .   | 299–304 |
| <b>Methacholine</b>  |         |
| The effect of three bufadienolide cardiac glycosides on contraction of isolated rat jejunum—C.J. BOTHA, R. GEHRING, J.M. VAN ROOYEN and D. VENTER . . . . .  | 243–246 |
| <b>Mites</b>   |         |
| On-host ecology and off-host survival of the sheep scab mite <i>Psoroptes ovis</i> —T. MEINTJES, L.J. FOURIE and I.G. HORAK . . . . .  | 273–283 |
| Effects of the scab mite <i>Psoroptes ovis</i> on the haematology and live mass of Merino and Dorper sheep—T. MEINTJES, L.J. FOURIE and I.G. HORAK . . . . .   | 285–293 |
| <b>Mozambique</b>  |         |
| A survey on the occurrence of resistance to anthelmintics of gastrointestinal nematodes of goats in Mozambique—A. ATANÁSIO, J. BOOMKER and C. SITOE . . . . .  | 215–220 |
| <b>Mycobacterium bovis</b>   |         |
| The gamma-interferon test: its usefulness in a bovine tuberculosis survey in African buffaloes ( <i>Syncerus caffer</i> ) in the Kruger National Park—D.G. GROBLER, ANITA L. MICHEL, LIN-MARI DE KLERK and R.G. BENGIS . . . . .   | 221–227 |
| <b>NAD-dependence/NAD-independence</b>   |         |
| Virulence of South African isolates of <i>Haemophilus paragallinarum</i> . Part 1: NAD-dependent field isolates—R.R. BRAGG . . . . .   | 163–169 |
| Virulence of South African isolates of <i>Haemophilus paragallinarum</i> . Part 2: Naturally occurring NAD-independent field isolates—R.R. BRAGG . . . . .   | 171–175 |

|   |         |
|---|---------|
| Virulence of South African isolates of <i>Haemophilus paragallinarum</i> . Part 3: experimentally produced NAD-independent isolate—M. TAOLE, J. ALBERTYN, E. VAN HEERDEN and R.R. BRAGG   | 189–196 |
| <b>Namibia</b>  |         |
| Apparent prevalence of dourine in the Khomas region of Namibia—F.F. KUMBA, B. CLAASSEN and P. PETRUS  | 295–298 |
| <b>Nematodes</b>  |         |
| Redescription of some <i>Spauligodon</i> spp. and <i>Parapharyngodon</i> spp., and of <i>Skrjabinodon mabuyaе</i> Sandground, 1936 (Pharyngodonidae: Oxyuroidea) from insectivorous South African lizards—S.F.B.N. HERING-HAGENBECK, A.J. PETTER and J. BOOMKER | 7–29    |
| Redescription of some <i>Thelandros</i> and <i>Tachygonetria</i> spp. (Pharyngodonidae: Oxyuroidea) from the omnivorous plated lizard, <i>Gerrhosaurus validus validus</i> A. Smith, 1849 in South Africa—S.F.B.N. HERING-HAGENBECK, A.J. PETTER and J. BOOMKER | 31–51   |
| Seasonal availability of gastrointestinal nematode larvae to cattle on pasture in the central highlands of Kenya—R.M. WARUIRU, E.H. WEDA, R.O. OTIENO and J.W. NGOTHO   | 141–146 |
| A survey on the occurrence of resistance to anthelmintics of gastrointestinal nematodes of goats in Mozambique—A. ATANASIO, J. BOOMKER and C. SITOE   | 215–220 |
| Strategic control of gastrointestinal nematodes of sheep in the highlands of central Kenya—N. MAINGI, R.O. OTIENO, V.M. MUMBI and E.H. WEDA   | 229–235 |
| <b>Neutralizing antibody</b>  |         |
| The classification of seven serotypes of equine encephalitis virus and the prevalence of homologous antibody in horses in South Africa—P.G. HOWELL, DALEEN GROENEWALD, CARINA W. VISAGE, ANNA-MARI BOSMAN, J.A.W. COETZER and A.J. GUTHRIE                      | 79–93   |
| <b>Neutrophils</b>  |         |
| Non-immune control of trypanosomiasis. <i>In vitro</i> oxidative burst of PMA- and trypanosome-stimulated neutrophils of Boran and N'Dama cattle—V.O. TAIWO, J.O. ADEJINMI and J.O. OLUWANIYI   | 155–161 |
| <b>Newcastle disease virus</b>  |         |
| Multiplication of the V4 strain of Newcastle disease virus in Madin Derby bovine kidney cells—M. SAHLE, G.W. BURGESS and A. KIDANEMARIAM  | 201–206 |
| Development of solid phase antigen for indirect ELISA for the detection of specific antibody responses to infection with Newcastle disease virus—M. SAHLE and G.W. BURGESS  | 237–242 |
| <b>Nigeria</b>  |         |
| Non-immune control of trypanosomiasis: <i>In vitro</i> oxidative burst of PMA- and trypanosome-stimulated neutrophils of Boran and N'Dama cattle—V.O. TAIWO, J.O. ADEJINMI and J.O. OLUWANIYI   | 155–161 |
| <b>Orbivirus</b>  |         |
| The classification of seven serotypes of equine encephalitis virus and the prevalence of homologous antibody in horses in South Africa—P.G. HOWELL, DALEEN GROENEWALD, CARINA W. VISAGE, ANNA-MARI BOSMAN, J.A.W. COETZER and A.J. GUTHRIE                      | 79–93   |
| <b>Osmosis</b>  |         |
| Water absorption in relation to fermentation in the colon of the ostrich ( <i>Struthio camelus</i> )—C. MUSARA, J.P. CHAMUNORWA, K. HOLTUG and E. SKADHAUGE   | 315–320 |
| <b>Ostrich</b>  |         |
| Water absorption in relation to fermentation in the colon of the ostrich ( <i>Struthio camelus</i> )—C. MUSARA, J.P. CHAMUNORWA, K. HOLTUG and E. SKADHAUGE   | 315–320 |
| <b>Oxyuroidea</b>   |         |
| Redescription of some <i>Spauligodon</i> spp. and <i>Parapharyngodon</i> spp., and of <i>Skrjabinodon mabuyaе</i> Sandground, 1936 (Pharyngodonidae: Oxyuroidea) from insectivorous South African lizards—S.F.B.N. HERING-HAGENBECK, A.J. PETTER and J. BOOMKER | 7–29    |
| Redescription of some <i>Thelandros</i> and <i>Tachygonetria</i> spp. (Pharyngodonidae: Oxyuroidea) from the omnivorous plated lizard, <i>Gerrhosaurus validus validus</i> A. Smith, 1849 in South Africa—S.F.B.N. HERING-HAGENBECK, A.J. PETTER and J. BOOMKER | 31–51   |
| <b>Pachydactylus</b>  |         |
| Redescription of some <i>Spauligodon</i> spp. and <i>Parapharyngodon</i> spp., and of <i>Skrjabinodon mabuyaе</i> Sandground,   |         |

## Subject Index

|  |         |
|--|---------|
| 1936) Inglis, 1968 (Pharyngodonidae: Oxyuroidea) from insectivorous South African lizards—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER .....   | 7-29    |
| <b>Parapharyngodon</b>   |         |
| Redescription of some <i>Spauligodon</i> spp. and <i>Parapharyngodon</i> spp., and of <i>Skrjabinodon mabuyae</i> Sandground, 1936) Inglis, 1968 (Pharyngodonidae: Oxyuroidea) from insectivorous South African lizards—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER ..... | 7-29    |
| <b>Pelomedusa</b>  |         |
| Description of <i>Pelonia africana</i> n. g., n. sp. (Pentastomida: Sebekidae) from the lungs of <i>Pelomedusa subrufa</i> and <i>Pelusios sinuatus</i> (Chelonia) in South Africa—K. JUNKER and J. BOOMKER .....  | 53-59   |
| <b>Pelonia africana</b>  |         |
| Description of <i>Pelonia africana</i> n. g., n. sp. (Pentastomida: Sebekidae) from the lungs of <i>Pelomedusa subrufa</i> and <i>Pelusios sinuatus</i> (Chelonia) in South Africa—K. JUNKER and J. BOOMKER .....  | 53-59   |
| <b>Pelusios</b>  |         |
| Description of <i>Pelonia africana</i> n. g., n. sp. (Pentastomida: Sebekidae) from the lungs of <i>Pelomedusa subrufa</i> and <i>Pelusios sinuatus</i> (Chelonia) in South Africa—K. JUNKER and J. BOOMKER .....  | 53-59   |
| <b>Pentastomida</b>  |         |
| Description of <i>Pelonia africana</i> n. g., n. sp. (Pentastomida: Sebekidae) from the lungs of <i>Pelomedusa subrufa</i> and <i>Pelusios sinuatus</i> (Chelonia) in South Africa—K. JUNKER and J. BOOMKER .....  | 53-59   |
| <b>Pharyngodonidae</b>   |         |
| Redescription of some <i>Thelandros</i> and <i>Tachygonetria</i> spp. (Pharyngodonidae: Oxyuroidea) from the omnivorous plated lizard, <i>Gerrhosaurus validus validus</i> A. Smith, 1849 in South Africa—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER .....               | 31-51   |
| <b>Plant poisoning</b>   |         |
| Continuous exposure to an aversive mixture as a means of maintaining aversion to vermeerbos ( <i>Geigeria ornativa</i> O. Hoffm.) in the presence of non-averted sheep—LD. SNYMAN, R.A. SCHULTZ, T.S. KELLERMAN and L. LABUSCHAGNE .....   | 321-325 |
| <b>Psoroptes ovis</b>  |         |
| On-host ecology and off-host survival of the sheep scab mite <i>Psoroptes ovis</i> —T. MEINTJES, L.J. FOURIE and I.G. HORAK .....  | 273-283 |
| Effects of the scab mite <i>Psoroptes ovis</i> on the haematology and live mass of Merino and Dorper sheep—T. MEINTJES, L.J. FOURIE and I.G. HORAK .....   | 285-293 |
| <b>Rat jejunum</b>   |         |
| The effect of three butadienolide cardiac glycosides on contraction of isolated rat jejunum—C.J. BOTHA, R. GEHRING, J.M. VAN ROOYEN and D. VENTER .....  | 243-246 |
| <b>Rhipicentor nuttalli</b>  |         |
| Hosts, seasonal occurrence and life cycle of <i>Rhipicentor nuttalli</i> (Acar: Ixodidae)—L.J. FOURIE, I.G. HORAK, D.J. KOK and W. VAN ZYL .....   | 177-187 |
| <b>Rhipicephalus spp.</b>  |         |
| Seasonal abundance of adult ixodid ticks infesting cattle belonging to resource-limited farmers in the north-eastern Free State Province of South Africa—M. HLATSHWAYO, P.A. MBATI and O.O. DIPEOLU .....  | 1-6     |
| <b>Rift Valley fever virus</b>   |         |
| Teratogenicity of a mutagenised Rift Valley fever virus (MVP 12) in sheep—P. HUNTER, B.J. ERASMUS and J.H. VORSTER .....   | 95-98   |
| Development of a diagnostic one-tube RT-PCR for the detection of Rift Valley fever virus—A. ESPACH, M. ROMITO, L.H. NEL and G.J. VILJOEN .....   | 247-252 |
| <b>RT-PCR</b>  |         |
| Development of a diagnostic one-tube RT-PCR for the detection of Rift Valley fever virus—A. ESPACH, M. ROMITO, L.H. NEL and G.J. VILJOEN .....   | 247-252 |

**Saudi Arabia**

Seasonal abundance of four *Culicoides* spp. (Diptera: Ceratopogonidae) at Al-Ahsa oasis, Eastern Province, Saudi Arabia—E.M.E. ABU ELZEIN, M.A. HILALI, A.I. AL-AFALEQ, P.S. MELLOR, J. BOORMAN, S. AL-ATIYA and A. AL NAIEM ..... 115-122

**Scincidae**

Redescription of some *Spauligodon* spp. and *Parapharyngodon* spp., and of *Skrjabinodon mabuyaе* Sandground, 1936 (Pharyngodonidae: Oxyuroidea) from insectivorous South African lizards—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER ..... 7-29

**Sera**

Validation of an indirect enzyme-linked immunosorbent assay for the detection of antibody against *Brucella abortus* in cattle sera using an automated ELISA workstation—J.T. PAWESKA, A.D. POTTS, H.J. HARRIS, S.J. SMITH, G.J. VILJOEN, B. DUNGU, O.L. BRETT, M. BUBB and L. PROZESKY ..... 61-77

**Serotypes**

The classification of seven serotypes of equine encephalitis virus and the prevalence of homologous antibody in horses in South Africa—P.G. HOWELL, DALEEN GROENEWALD, CARINA W. VISAGE, ANNA-MARI BOSMAN, J.A.W. COETZER and A.J. GUTHRIE ..... 79-93

**Sheep**

Teratogenicity of a mutagenised Rift Valley fever virus (MVP 12) in sheep—P. HUNTER, B.J. ERASMUS and J.H. VORSTER ..... 95-98

Strategic control of gastrointestinal nematodes of sheep in the highlands of central Kenya—N. MAINGI, R.O. OTIENO, V.M. MUMBI and E.H. WEDA ..... 229-235

On-host ecology and off-host survival of the sheep scab mite *Psoroptes ovis*—T. MEINTJES, L.J. FOURIE and I.G. HORAK ..... 273-283

Effects of the scab mite *Psoroptes ovis* on the haematology and live mass of Merino and Dorper sheep—T. MEINTJES, L.J. FOURIE and I.G. HORAK ..... 285-293

Continuous exposure to an aversive mixture as a means of maintaining aversion to vermeerbos (*Geigeria ornativa* O. Hoffm.) in the presence of non-averted sheep—L.D. SNYMAN, R.A. SCHULTZ, T.S. KELLERMAN and L. LABU-SCHAGNE ..... 321-325

***Skrjabinodon***

Redescription of some *Spauligodon* spp. and *Parapharyngodon* spp., and of *Skrjabinodon mabuyaе* Sandground, 1936 (Pharyngodonidae: Oxyuroidea) from insectivorous South African lizards—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER ..... 7-29

**South Africa**

Seasonal abundance of adult ixodid ticks infesting cattle belonging to resource-limited farmers in the north-eastern Free State Province of South Africa—M. HLATSHWAYO, P.A. MBATI and O.O. DIPEOLU ..... 1-6

Redescription of some *Spauligodon* spp. and *Parapharyngodon* spp., and of *Skrjabinodon mabuyaе* Sandground, 1936 (Pharyngodonidae: Oxyuroidea) from insectivorous South African lizards—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER ..... 7-29

Redescription of some *Thelanos* and *Tachygonetria* spp. (Pharyngodonidae: Oxyuroidea) from the omnivorous plated lizard, *Gerrhosaurus validus validus* A. Smith, 1849 in South Africa—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER ..... 31-51

Description of *Pelonia africana* n. g., n. sp. (Pentastomida: Sebekidae) from the lungs of *Pelomedusa subrufa* and *Pelusios sinuatus* (Chelonidae) in South Africa—K. JUNKER and J. BOOMKER ..... 53-59

The classification of seven serotypes of equine encephalitis virus and the prevalence of homologous antibody in horses in South Africa—P.G. HOWELL, DALEEN GROENEWALD, CARINA W. VISAGE, ANNA-MARI BOSMAN, J.A.W. COETZER and A.J. GUTHRIE ..... 79-93

Acaricide resistance profiles of single- and multi-host ticks from communal and commercial farming areas in the Eastern Cape and North-West Provinces of South Africa—S. MEKONNEN, N.R. BRYSON, L.J. FOURIE, R.J. PETER, A.M. SPICKETT, R.J. TAYLOR, T. STRYDOM and I.G. HORAK ..... 99-105

Virulence of South African isolates of *Haemophilus paragallinarum*. Part 1: NAD-dependent field isolates—R.R. BRAGG ..... 163-169

Virulence of South African isolates of *Haemophilus paragallinarum*. Part 2: Naturally occurring NAD-independent field isolates—R.R. BRAGG ..... 171-175

|   |         |
|---|---------|
| <i>In vitro</i> isolation of equine piroplasms derived from Cape mountain zebra ( <i>Equus zebra zebra</i> ) in South Africa—E. ZWEYGARTH, LAURA M. LOPEZ-REBOLLAR and P. MEYER .....   | 197-200 |
| Amphistome infection of goats farmed under resource-poor conditions in South Africa—A.F. VATTA and R.C. KRECEK .....  | 327-329 |
| <b><i>Spauligodon</i></b>   |         |
| Redescription of some <i>Spauligodon</i> spp. and <i>Parapharyngodon</i> spp., and of <i>Skrjabinodon mabuyaæ</i> Sandground, 1936 (Pharyngodontidae: Oxyuroidea) from insectivorous South African lizards—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER .....                 | 7-29    |
| <b><i>Struthio camelus</i></b>  |         |
| Water absorption in relation to fermentation in the colon of the ostrich ( <i>Struthio camelus</i> )—C. MUSARA, J.P. CHAMUNORWA, K. HOLTUG and E. SKADHAUGE .....   | 315-320 |
| <b><i>Synicerus caffer</i></b>  |         |
| The gamma-interferon test: its usefulness in a bovine tuberculosis survey in African buffaloes ( <i>Synicerus caffer</i> ) in the Kruger National Park—D.G. GROBLER, ANITA L. MICHEL, LIN-MARI DE KLERK and R.G. BENGIS .....   | 221-227 |
| <b><i>Tachygometria</i></b>   |         |
| Redescription of some <i>Thelandros</i> and <i>Tachygometria</i> spp. (Pharyngodontidae: Oxyuroidea) from the omnivorous plated lizard, <i>Gerrhosaurus validus validus</i> A. Smith, 1849 in South Africa—S F B N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER .....                 | 31-51   |
| <b>Tanzania</b>   |         |
| Review of current problems and shortcomings in the Tanzanian animal health information system with suggestions on improvement—F.M. KIVARIA and A.M. KAPAGA .....  | 305-314 |
| <b>Teratogenicity</b>   |         |
| Teratogenicity of a mutagenised Rift Valley fever virus (MVP 12) in sheep—P. HUNTER, B.J. ERASMUS and J.H. VORSTER .....  | 95-98   |
| <b><i>Theileria equi</i></b>  |         |
| <i>In vitro</i> isolation of equine piroplasms derived from Cape mountain zebra ( <i>Equus zebra zebra</i> ) in South Africa—E. ZWEYGARTH, LAURA M. LOPEZ-REBOLLAR and P. MEYER .....   | 197-200 |
| <b>Theileriosis</b>   |         |
| Occurrence of concurrent trypanosomosis, theileriosis, anaplasmosis and helminthosis in Friesian, Zebu and Sahi-wal cattle in Uganda—J.W. MAGONA and J.S.P. MAYENDE .....   | 133-140 |
| <b><i>Thelandros</i></b>  |         |
| Redescription of some <i>Thelandros</i> and <i>Tachygometria</i> spp. (Pharyngodontidae: Oxyuroidea) from the omnivorous plated lizard, <i>Gerrhosaurus validus validus</i> A. Smith, 1849 in South Africa—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER .....                 | 31-51   |
| <b>Ticks</b>  |         |
| Seasonal abundance of adult ixodid ticks infesting cattle belonging to resource-limited farmers in the north-eastern Free State Province of South Africa—M. HLATSHWAYO, P.A. MBATI and O.O. DIPEOLU .....   | 1-6     |
| Acaricide resistance profiles of single- and multi-host ticks from communal and commercial farming areas in the Eastern Cape and North-West Provinces of South Africa—S. MEKONNEN, N.R. BRYSON, L.J. FOURIE, A.J. PETTER, A.M. SPICKETT, R.J. TAYLOR, T. STRYDOM and I.G. HORAK ..... | 99-105  |
| <b>Trematode</b>  |         |
| The prevalence and economic importance of bovine fasciolosis in Kenya—an analysis of abattoir data—J.M. KITHUKA, N. MAINGI, F.M. NJERUH and J.N. OMBUI .....  | 255-262 |
| Amphistome infection of goats farmed under resource-poor conditions in South Africa—A.F. VATTA and R.C. KRECEK .....  | 327-329 |
| <b><i>Trypanosoma/trypanosomosis</i></b>  |         |
| Parasitological prevalence of bovine trypanosomosis in Kindo Koisha district, Wolaita zone, south Ethiopia—A. KIDANEMARIAM, K. HADGU and M. SAHLE .....   | 107-113 |
| Occurrence of concurrent trypanosomosis, theileriosis, anaplasmosis and helminthosis in Friesian, Zebu and Sahi-wal cattle in Uganda—J.W. MAGONA and J.S.P. MAYENDE .....   | 133-140 |

|   |         |
|---|---------|
| Non-immune control of trypanosomosis: <i>In vitro</i> oxidative burst of PMA- and trypanosome-stimulated neutrophils of Boran and N'Dama cattle—V.O. TAIWO, J.O. ADEJINMI and J.O. OLUWANIYI . . . . .  | 155-161 |
| Animal-level risk factors for <i>Trypanosoma evansi</i> infection in camels in eastern and central parts of Kenya—J.M. NGAIRA, B. BETT and S.M. KARANJA . . . . .   | 263-271 |
| Apparent prevalence of dourine in the Khomas region of Namibia—F.F. KUMBA, B. CLAASEN and P. PETRUS . . . . .   | 295-298 |
| <b>Tyledoside D</b>   |         |
| The effect of three bufadienolide cardiac glycosides on contraction of isolated rat jejunum—C.J. BOTHA, R. GEHRING, J.M. VAN ROOYEN and D. VENTER . . . . .   | 243-246 |
| <b>Uganda</b>   |         |
| Occurrence of concurrent trypanosomosis, theileriosis, anaplasmosis and helminthosis in Friesian, Zebu and Sahi-wal cattle in Uganda—J.W. MAGONA and J.S.P. MAYENDE . . . . .   | 133-140 |
| <b>V4 strain replication</b>  |         |
| Multiplication of the V4 strain of Newcastle disease virus in Madin Derby bovine kidney cells—M. SAHLE, G.W. BURGESS and A. KIDANEMARIAM . . . . .  | 201-206 |
| <b>Vaccines</b>   |         |
| Isolation of serovar C-3 <i>Haemophilus paragallinarum</i> from Zimbabwe: A further indication of the need for the production of vaccines against infectious coryza containing local isolates of <i>H. paragallinarum</i> —R.R. BRAGG . . . . .                 | 129-132 |
| <b>Vermeersiekte</b>  |         |
| Continuous exposure to an aversive mixture as a means of maintaining aversion to vermeerbos ( <i>Geigeria ornativa</i> O. Hoffm.) in the presence of non-averted sheep—L.D. SNYMAN, R.A. SCHULTZ, T.S. KELLERMAN and L. LABU-SCHAGNE . . . . .                  | 321-325 |
| <b>Zambia</b>   |         |
| Hepatic and renal concentrations of copper and other trace elements in hippopotami ( <i>Hippopotamus amphibius</i> L.) living in and adjacent to the Kafue and Luangwa Rivers in Zambia—M. MWASE, B. ALMLI, T. SIVERTSEN, M.M. MUSONDA and A. FLÅØYEN . . . . . | 207-214 |
| <b>Zimbabwe</b>   |         |
| Isolation of serovar C-3 <i>Haemophilus paragallinarum</i> from Zimbabwe: A further indication of the need for the production of vaccines against infectious coryza containing local isolates of <i>H. paragallinarum</i> —R.R. BRAGG . . . . .                 | 129-132 |