



Author index

ABAIGAR, T. Oral administration of mebendazole failed to reduce nematode egg shedding in captive African gazelles—J. ORTIZ, M.R. RUIZ DE YBAÑEZ, T. ABAIGAR, M. GARIJO, G. ESPESO and M. CANO	79–82
AHARISOLO, C. Validity of intradermal tuberculin testing for the screening of bovine tuberculosis in Madagascar—R. QUIRIN, V. RASOLOFO, R. ANDRIAMBOLOLONA, A. RAMBOASOLO, T. RASOLONAVALONA, C. AHARISOLO, H. RAKOTARITAHINA, S. CHANTEAU and P. BOISIER	231–238
ALCIGIR, G. Immunohistochemical and histopathological studies of fixed rabies virus in goats—S. ATALAY VURAL, G. ALCIGIR and S. BERKIN	83–89
ANDONOVA, M. Changes in some factors of the innate immunity and serum zinc and iron concentrations in pigs following intravenous administration of <i>Escherichia coli</i> lipopolysaccharide—M. ANDONOVA, I. BORISSOV and L. SOTIROV	91–99
ANDRIAMBOLOLONA, R. Validity of intradermal tuberculin testing for the screening of bovine tuberculosis in Madagascar—R. QUIRIN, V. RASOLOFO, R. ANDRIAMBOLOLONA, A. RAMBOASOLO, T. RASOLONAVALONA, C. AHARISOLO, H. RAKOTARITAHINA, S. CHANTEAU and P. BOISIER	231–238
ATALAY VURAL, S. Immunohistochemical and histopathological studies of fixed rabies virus in goats—S. ATALAY VURAL, G. ALCIGIR and S. BERKIN	83–89
BASSON, K.M. Effect of pavetamine on protein synthesis in rat tissue—R.A. SCHULTZ, N. FOURIE, K.M. BASSON, L. LABUSCHAGNE and L. PROZESKY	325–331
BEDANE, B. The distribution of ixodid ticks (Acari: Ixodidae) in central Ethiopia—S. MEKONNEN, I. HUSSEIN and B. BEDANE ..	243–251
BENGIS, R.G. The epidemiology of tuberculosis in free ranging African buffalo (<i>Syncerus caffer</i>) in the Kruger National Park, South Africa—V. DE VOS, R.G. BENGIS, N.P.J. KRIEK, A. MICHEL, D.F. KEET, J.P. RAATH and H.F.K.A. HUCHZERMAYER	119–130
Tuberculosis in kudu (<i>Tragelaphus strepsiceros</i>) in the Kruger National Park—D.F. KEET, N.P.J. KRIEK, R.G. BENGIS and A.L. MICHEL	225–230
Tuberculosis, caused by <i>Mycobacterium bovis</i> , in a kudu (<i>Tragelaphus strepsiceros</i>) from a commercial game farm in the Malelane area of the Mpumalanga Province, South Africa—R.G. BENGIS, D.F. KEET, A.L. MICHEL and N.P.J. KRIEK	239–241

BERKIN, S.	
Immunohistochemical and histopathological studies of fixed rabies virus in goats—S. ATALAY VURAL, G. ALCIGIR and S. BERKIN	83–89
BIGALKE, R.D.	
A review of the infectious disease of African wild ruminants—R.W. WORTHINGTON and R.D. BIGALKE	291–323
BINEPAL, Y.S.	
Alternative cell lines for the propagation of lumpy skin disease virus—Y.S. BINEPAL, F.A. ONGADI and J.C. CHEPKWONY	151–153
BINTA, M.G.	
Antibodies to Newcastle disease virus in the sera of indigenous chickens in Oodi, Kgatleng, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, J.M.K. HYERA, K.M. THIBANYANE and J. MKARIA	69–70
Seroprevalence of <i>Toxoplasma gondii</i> and <i>Chlamydia psittaci</i> in domestic pigeons (<i>Columba livia domestica</i>) at Sebele, Gaborone, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, R. NDEBELE and R. PANZIRAH	159–161
BLAYLOCK, R.S.M.	
Normal oral bacterial flora from some southern African snakes—R.S.M. BLAYLOCK	175–182
BOISIER, P.	
Validity of intradermal tuberculin testing for the screening of bovine tuberculosis in Madagascar—R. QUIRIN, V. RASOLOFO, R. ANDRIAMBOLOLONA, A. RAMBOASOLO, T. RASOLONAVALONA, C. AHARISOLO, H. RAKOTARITAHINA, S. CHANTEAU and P. BOISIER	231–238
BORISSOV, I.	
Changes in some factors of the innate immunity and serum zinc and iron concentrations in pigs following intravenous administration of <i>Escherichia coli</i> lipopolysaccharide—M. ANDONOVA, I. BORISSOV and L. SOTIROV	91–99
BOTHA, C.J.	
Seasonal variation in cotyledoside concentration of <i>Tylecodon wallichii</i> (Harv.) Tölken subsp. <i>wallichii</i> sampled in a krimpsiekte-prevalent region—C.J. BOTHA, T. RUNDBERGET, A.L. WILKINS, M.S.G. MÜLDERS, A. FLÅØYEN and M.P. VAN AARDT	1–9
BURSEY, C.R.	
Intestinal helminths of four species of skinks (<i>Mabuya</i>) (Sauria: Scincidae) from southern Africa—S.R. GOLDBERG and C.R. BURSEY	143–147
CANO, M.	
Oral administration of mebendazole failed to reduce nematode egg shedding in captive African gazelles—J. ORTIZ, M.R. RUIZ DE YBÁÑEZ, T. ABAIGAR, M. GARIJO, G. ESPESO and M. CANO	79–82
CHABO, R.G.	
Antibodies to Newcastle disease virus in the sera of indigenous chickens in Oodi, Kgatleng, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, J.M.K. HYERA, K.M. THIBANYANE and J. MKARIA	69–70
Seroprevalence of <i>Toxoplasma gondii</i> and <i>Chlamydia psittaci</i> in domestic pigeons (<i>Columba livia domestica</i>) at Sebele, Gaborone, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, R. NDEBELE and R. PANZIRAH	159–161
The impact of traditional management on seasonal internal parasite burdens and productivity of indigenous Tswana goats in southern Botswana—S.J. NSOSO, J.B. MACHETE, M. MOLATOLE, R.T. NDEBELE, N.N. LEBANI, R.G. CHABO, A.M. KALAKE, L. JACYNA, B.W. SEGADIMO and O.M. MINE	101–104
CHANTEAU, S.	
Validity of intradermal tuberculin testing for the screening of bovine tuberculosis in Madagascar—R. QUIRIN, V. RASOLOFO, R. ANDRIAMBOLOLONA, A. RAMBOASOLO, T. RASOLONAVALONA, C. AHARISOLO, H. RAKOTARITAHINA, S. CHANTEAU and P. BOISIER	231–238
CHEPKWONY, J.C.	
Alternative cell lines for the propagation of lumpy skin disease virus—Y.S. BINEPAL, F.A. ONGADI and J.C. CHEPKWONY	151–153
COHEN, M.	
Hosts of the immature stages of the rhinoceros tick, <i>Dermacentor rhinocerinus</i> (Acari, Ixodidae)—I.G. HORAK and M. COHEN	75–77

- DE VOS, V.**
The epidemiology of tuberculosis in free ranging African buffalo (*Syncerus caffer*) in the Kruger National Park, South Africa—V. DE VOS, R.G. BENGIS, N.P.J. KRIEK, A. MICHEL, D.F. KEET, J.P. RAATH and H.F.K.A. HUCHZERMEYER 119–130
- DIPPENAAR, S.M.**
Lamproglena hoi n. sp. (Copepoda: Lernaedidae) from two yellowfish hosts, *Barbus marequensis* and *Barbus polylepis*, caught in a river in Mpumalanga, South Africa—S.M. DIPPENAAR, W.J. LUUS-POWELL and F. ROUX 209–215
- DORNY, P.**
A survey on anthelmintic resistance in nematode parasites of sheep in Lusaka, Zambia—S. GABRIEL, I.K. PHIRI, P. DORNY and J. VERCRUYSE 271–274
- DZIVA, F.**
Vaccination to control an outbreak of *Mycoplasma crocodyli* infection—K. MOHAN, C.M. FOGGIN, F. DZIVA and P. MUVAVARIRWA 149–150
- ELKAN, P.W.**
Multifactorial mortality in bongos and other wild ungulates in the north of the Congo Republic—F.W. HUCHZERMEYER, M.-L. PENRITH and P.W. ELKAN 263–269
- EMSLIE, F.R.**
Parasites of domestic and wild animals in South Africa. XL. Ticks on dogs belonging to people in rural communities and carnivore ticks on the vegetation—I.G. HORAK, F.R. EMSLIE and A.M. SPICKETT 135–141
- ESMANN, J.B.**
A survey of parasitic nematode infections of chickens in rural Zimbabwe—S. MUKARATIRWA, T. HOVE, J.B. ESMANN, C.J. HOJ, A. PERMIN and P. NANSEN 183–186
- ESPESO, G.**
Oral administration of mebendazole failed to reduce nematode egg shedding in captive African gazelles—J. ORTIZ, M.R. RUIZ DE YBÁÑEZ, T. ABAIGAR, M. GARIJO, G. ESPESO and M. CANO 79–82
- FLÅØYEN, A.**
Seasonal variation in cotyledoside concentration of *Tylecodon wallichii* (Harv.) Tölken subsp. *wallichii* sampled in a krimpsiekte-prevalent region—C.J. BOTHA, T. RUNDBERGET, A.L. WILKINS, M.S.G. MÜLDERS, A. FLÅØYEN and M.P. VAN AARDT 1–9
- FOGGIN, C.M.**
Transmission studies on *Trichinella* species isolated from *Crocodylus niloticus* and efficacy of fenbendazole and levamisole against muscle L1 stages in Balb C mice—S. MUKARATIRWA, K. MAGWEDERE, E. MATENGA and C.M. FOGGIN 21–25
Vaccination to control an outbreak of *Mycoplasma crocodyli* infection—K. MOHAN, C.M. FOGGIN, F. DZIVA and P. MUVAVARIRWA 149–150
- FOURIE, L.J.**
Diversity, seasonality and sites of attachment of adult ixodid ticks on dogs in the central region of the Free State Province, South Africa—P.A.H. JACOBS, L.J. FOURIE, D.J. KOK and I.G. HORAK 281–290
- FOURIE, N.**
Effect of pavetamine on protein synthesis in rat tissue—R.A. SCHULTZ, N. FOURIE, K.M. BASSON, L. LABUSCHAGNE and L. PROZESKY 325–331
- FUKUSHI, H.**
Prevalence of virulent *Rhodococcus equi* in isolates from soil collected from two horse farms in South Africa and restriction fragment length polymorphisms of virulence plasmids in the isolates from infected foals, a dog and a monkey—S. TAKAI, M.M. HENTON, J.A. PICARD, A.J. GUTHRIE, H. FUKUSHI and C. SUGIMOTO 105–110
- GABRIEL, S.**
A survey on anthelmintic resistance in nematode parasites of sheep in Lusaka, Zambia—S. GABRIEL, I.K. PHIRI, P. DORNY and J. VERCRUYSE 271–274
- GARIJO, M.**
Oral administration of mebendazole failed to reduce nematode egg shedding in captive African gazelles—J. ORTIZ, M.R. RUIZ DE YBÁÑEZ, T. ABAIGAR, M. GARIJO, G. ESPESO and M. CANO 79–82

GOLDBERG, S.R.

Intestinal helminths of four species of skinks (*Mabuya*) (Sauria: Scincidae) from southern Africa—S.R. GOLDBERG and C.R. BURSEY 143–147

GOTHE, R.

Studies on the critical water mass, rehydration capability and potential, acute chill tolerance and supercooling point of *Argas (Pescicargas) walkerae* (Acari: Argasidae)—U. STARK and R. GOTHE 11–20

GREEFF, C.J.

Parasites of domestic and wild animals in South Africa. XXXIX. Helminth and arthropod parasites of Angora goats in the southern Karoo—I.G. HORAK, K.M. DE F. MACIVOR and C.J. GREEFF 27–35

GUTHRIE, A.J.

Prevalence of virulent *Rhodococcus equi* in isolates from soil collected from two horse farms in South Africa and restriction fragment length polymorphisms of virulence plasmids in the isolates from infected foals, a dog and a monkey—S. TAKAI, M.M. HENTON, J.A. PICARD, A.J. GUTHRIE, H. FUKUSHI and C. SUGIMOTO 105–110

HEATH, L.E.

The 3A non-structural-protein coding region of the southern African SAT type isolates differs from that of other foot-and-mouth disease viruses—L.E. HEATH, H.G. VAN RENSBURG, W. VOSLOO and L.H. NEL 253–262

HENTON, M.M.

Prevalence of virulent *Rhodococcus equi* in isolates from soil collected from two horse farms in South Africa and restriction fragment length polymorphisms of virulence plasmids in the isolates from infected foals, a dog and a monkey—S. TAKAI, M.M. HENTON, J.A. PICARD, A.J. GUTHRIE, H. FUKUSHI and C. SUGIMOTO 105–110

HOJ, C.J.

A survey of parasitic nematode infections of chickens in rural Zimbabwe—S. MUKARATIRWA, T. HOVE, J.B. ESMANN, C.J. HOJ, A. PERMIN and P. NANSEN 183–186

HORAK, I.G.

Parasites of domestic and wild animals in South Africa. XXXIX. Helminth and arthropod parasites of Angora goats in the southern Karoo—I.G. HORAK, K.M. DE F. MACIVOR and C.J. GREEFF 27–35

Hosts of the immature stages of the rhinoceros tick, *Dermacentor rhinoceros* (Acari, Ixodidae)—I.G. HORAK and M. COHEN 75–77

Parasites of domestic and wild animals in South Africa. XL. Ticks on dogs belonging to people in rural communities and carnivore ticks on the vegetation—I.G. HORAK, F.R. EMSLIE and A.M. SPICKETT 135–141

Diversity, seasonality and sites of attachment of adult ixodid ticks on dogs in the central region of the Free State Province, South Africa—P.A.H. JACOBS, L.J. FOURIE, D.J. KOK and I.G. HORAK 281–290

HOVE, T.

A survey of parasitic nematode infections of chickens in rural Zimbabwe—S. MUKARATIRWA, T. HOVE, J.B. ESMANN, C.J. HOJ, A. PERMIN and P. NANSEN 183–186

Epidemiological observations of Zimbabwean theileriosis: Disease incidence and pathogenicity in susceptible cattle during *Rhipicephalus appendiculatus* nymphal and adult seasonal activities—A.A. LATIF, T. HOVE, G.K. KANHAI, S. MASAKA and R.G. PEGRAM 187–195

Exposure of cattle immunized with different stocks of *Theileria parva* to buffalo-associated *Theileria* challenge on two game parks in Zimbabwe—A.A. LATIF, T. HOVE, G.K. KANHAI and S. MASAKA 197–201

Laboratory and field investigations into the *Theileria parva* carrier-state in cattle in Zimbabwe—A.A. LATIF, T. HOVE, G.K. KANHAI and S. MASAKA 203–208

HUCHZERMEYER, F.W.

Multifactorial mortality in bongos and other wild ungulates in the north of the Congo Republic—F.W. HUCHZERMEYER, M.-L. PENRITH and P.W. ELKAN 263–269

HUCHZERMEYER, H.F.K.A.

The epidemiology of tuberculosis in free ranging African buffalo (*Syncerus caffer*) in the Kruger National Park, South Africa—V. DE VOS, R.G. BENGIS, N.P.J. KRIEK, A. MICHEL, D.F. KEET, J.P. RAATH and H.F.K.A. HUCHZERMEYER 119–130

HUNTER, P.

A monovalent attenuated serotype 2 bluetongue virus vaccine confers homologous protection in sheep—P. HUNTER and J. MODUMO 331–333

- HUSSEIN, I.**
The distribution of ixodid ticks (Acari: Ixodidae) in central Ethiopia—S. MEKONNEN, I. HUSSEIN and B. BEDANE . 243–251
- HYERA, J.M.K.**
Antibodies to Newcastle disease virus in the sera of indigenous chickens in Oodi, Kgatleng, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, J.M.K. HYERA, K.M. THIBANYANE and J. MKARIA 69–70
- JACOBS, P.A.H.**
Diversity, seasonality and sites of attachment of adult ixodid ticks on dogs in the central region of the Free State Province, South Africa—P.A.H. JACOBS, L.J. FOURIE, D.J. KOK and I.G. HORAK 281–290
- JACYNA, L.**
The impact of traditional management on seasonal internal parasite burdens and productivity of indigenous Tswana goats in southern Botswana—S.J. NSOSO, J.B. MACHETE, M. MOLATOLE, R.T. NDEBELE, N.N. LEBANI, R.G. CHABO, A.M. KALAKE, L. JACYNA, B.W. SEGADIMO and O.M. MINE 101–104
- JOSEMANS, A.I.**
A chemically defined medium for the growth of *Cowdria ruminantium*—E. ZWEYGARTH and A.I. JOSEMANS 37–40
Continuous *in vitro* propagation of *Cowdria ruminantium* (Welgevonden stock) in a canine macrophage-monocyte cell line—E. ZWEYGARTH and ANTOINETTE I. JOSEMANS 155–157
- KALAKE, A.M.**
The impact of traditional management on seasonal internal parasite burdens and productivity of indigenous Tswana goats in southern Botswana—S.J. NSOSO, J.B. MACHETE, M. MOLATOLE, R.T. NDEBELE, N.N. LEBANI, R.G. CHABO, A.M. KALAKE, L. JACYNA, B.W. SEGADIMO and O.M. MINE 101–104
- KANHAI, G.K.**
Epidemiological observations of Zimbabwean theileriosis: Disease incidence and pathogenicity in susceptible cattle during *Rhipicephalus appendiculatus* nymphal and adult seasonal activities—A.A. LATIF, T. HOVE, G.K. KANHAI, S. MASAKA and R.G. PEGRAM 187–195
Exposure of cattle immunized with different stocks of *Theileria parva* to buffalo-associated *Theileria* challenge on two game parks in Zimbabwe—A.A. LATIF, T. HOVE, G.K. KANHAI and S. MASAKA 197–201
Laboratory and field investigations into the *Theileria parva* carrier-state in cattle in Zimbabwe— A.A. LATIF, T. HOVE, G.K. KANHAI and S. MASAKA 203–208
- KEET, D.F.**
The epidemiology of tuberculosis in free ranging African buffalo (*Syncerus caffer*) in the Kruger National Park, South Africa—V. DE VOS , R.G. BENIGIS, N.P.J. KRIEK, A. MICHEL, D.F. KEET, J.P. RAATH and H.F.K.A. HUCHZERMEYER 119–130
Tuberculosis in kudu (*Tragelaphus strepsiceros*) in the Kruger National Park—D.F. KEET, N.P.J. KRIEK, R.G. BENIGIS and A.L. MICHEL 225–230
Tuberculosis, caused by *Mycobacterium bovis*, in a kudu (*Tragelaphus strepsiceros*) from a commercial game farm in the Malelane area of the Mpumalanga Province, South Africa—R.G. BENIGIS, D.F. KEET, A.L. MICHEL and N.P.J. KRIEK 239–241
- KOK, D.J.**
Diversity, seasonality and sites of attachment of adult ixodid ticks on dogs in the central region of the Free State Province, South Africa—P.A.H. JACOBS, L.J. FOURIE, D.J. KOK and I.G. HORAK 281–290
- KRECEK, R.C.**
Helminths in dogs belonging to people in a resource-limited urban community in Gauteng, South Africa—W.N. MINNAAR and R.C. KRECEK 111–117
- KRIEK, N.P.J.**
The epidemiology of tuberculosis in free ranging African buffalo (*Syncerus caffer*) in the Kruger National Park, South Africa—V. DE VOS , R.G. BENIGIS, N.P.J. KRIEK, A. MICHEL, D.F. KEET, J.P. RAATH and H.F.K.A. HUCHZERMEYER 119–130
Tuberculosis in kudu (*Tragelaphus strepsiceros*) in the Kruger National Park—D.F. KEET, N.P.J. KRIEK, R.G. BENIGIS and A.L. MICHEL 225–230
Tuberculosis, caused by *Mycobacterium bovis*, in a kudu (*Tragelaphus strepsiceros*) from a commercial game farm in the Malelane area of the Mpumalanga Province, South Africa—R.G. BENIGIS, D.F. KEET, A.L. MICHEL and N.P.J. KRIEK 239–241

LABUSCHAGNE, L.

- Effect of pavetamine on protein synthesis in rat tissue—R.A. SCHULTZ, N. FOURIE, K.M. BASSON, L. LABUSCHAGNE and L. PROZESKY 325–331

LANGONI, H.

- Seroprevalence of bovine leptospirosis in Garanhuns municipal district, Pernambuco State, Brazil—A.A.F. OLIVEIRA, R.A. MOTA, G.C. PEREIRA, H. LANGONI, M.I. SOUZA, W.A. NAVEGANTES and M.E.P. SÁ 275–279

LATIF, A.A.

- Epidemiological observations of Zimbabwean theileriosis: Disease incidence and pathogenicity in susceptible cattle during *Rhipicephalus appendiculatus* nymphal and adult seasonal activities—A.A. LATIF, T. HOVE, G.K. KANHAI, S. MASAKA and R.G. PEGRAM 187–195

- Exposure of cattle immunized with different stocks of *Theileria parva* to buffalo-associated *Theileria* challenge on two game parks in Zimbabwe—A.A. LATIF, T. HOVE, G.K. KANHAI and S. MASAKA 197–201

- Laboratory and field investigations into the *Theileria parva* carrier-state in cattle in Zimbabwe—A.A. LATIF, T. HOVE, G.K. KANHAI and S. MASAKA 203–208

LEBANI, N.N.

- The impact of traditional management on seasonal internal parasite burdens and productivity of indigenous Tswana goats in southern Botswana—S.J. NSOSO, J.B. MACHETE, M. MOLATOLE, R.T. NDEBELE, N.N. LEBANI, R.G. CHABO, A.M. KALAKE, L. JACYNA, B.W. SEGADIMO and O.M. MINE 101–104

LUUS-POWELL, W.J.

- Lamproglena hoi* n. sp. (Copepoda: Lernaecidae) from two yellowfish hosts, *Barbus marequensis* and *Barbus polylepis*, caught in a river in Mpumalanga, South Africa—S.M. DIPPENAAR, W.J. LUUS-POWELL and F. ROUX 209–215

MACHETE, J.B.

- The impact of traditional management on seasonal internal parasite burdens and productivity of indigenous Tswana goats in southern Botswana—S.J. NSOSO, J.B. MACHETE, M. MOLATOLE, R.T. NDEBELE, N.N. LEBANI, R.G. CHABO, A.M. KALAKE, L. JACYNA, B.W. SEGADIMO and O.M. MINE 101–104

MACIVOR, K.M. DE F.

- Parasites of domestic and wild animals in South Africa. XXXIX. Helminth and arthropod parasites of Angora goats in the southern Karoo—I.G. HORAK, K.M. DE F. MACIVOR and C.J. GREEFF 27–35

MAGWEDERE, K.

- Transmission studies on *Trichinella* species isolated from *Crocodylus niloticus* and efficacy of fenbendazole and levamisole against muscle L1 stages in Balb C mice—S. MUKARATIRWA, K. MAGWEDERE, E. MATENGA and C.M. FOGGIN 21–25

MALAN, F.S.

- Clinical evaluation of anaemia in sheep: early trials—F.S. MALAN, J.A. VAN WYK and C.D. WESSELS 165–174

MASAKA, S.

- Epidemiological observations of Zimbabwean theileriosis: Disease incidence and pathogenicity in susceptible cattle during *Rhipicephalus appendiculatus* nymphal and adult seasonal activities—A.A. LATIF, T. HOVE, G.K. KANHAI, S. MASAKA and R.G. PEGRAM 187–195

- Exposure of cattle immunized with different stocks of *Theileria parva* to buffalo-associated *Theileria* challenge on two game parks in Zimbabwe—A.A. LATIF, T. HOVE, G.K. KANHAI and S. MASAKA 197–201

- Laboratory and field investigations into the *Theileria parva* carrier-state in cattle in Zimbabwe—A.A. LATIF, T. HOVE, G.K. KANHAI and S. MASAKA 203–208

MATENGA, E.

- Transmission studies on *Trichinella* species isolated from *Crocodylus niloticus* and efficacy of fenbendazole and levamisole against muscle L1 stages in Balb C mice—S. MUKARATIRWA, K. MAGWEDERE, E. MATENGA and C.M. FOGGIN 21–25

MBOGO, S.K.

- Immunization of cattle against East Coast fever using *Theileria parva* (Marikebuni) and relaxation of tick control in North Rift, Kenya—J.M. WANJOHI, J.N. NGERANWA, R.M. RUMBERIA, G.R. MURAGURI and S.K. MBOGO 217–223

MEKONNEN, S.

- The distribution of ixodid ticks (Acari: Ixodidae) in central Ethiopia—S. MEKONNEN, I. HUSSEIN and B. BEDANE 243–251

- MICHEL, A.**
The epidemiology of tuberculosis in free ranging African buffalo (*Syncerus caffer*) in the Kruger National Park, South Africa—V. DE VOS, R.G. BENGIS, N.P.J. KRIEK, A. MICHEL, D.F. KEET, J.P. RAATH and H.F.K.A. HUCHZERMEYER 119–130
- MICHEL, A.L.**
Tuberculosis in kudu (*Tragelaphus strepsiceros*) in the Kruger National Park—D.F. KEET, N.P.J. KRIEK, R.G. BENGIS and A.L. MICHEL 225–230
Tuberculosis, caused by *Mycobacterium bovis*, in a kudu (*Tragelaphus strepsiceros*) from a commercial game farm in the Malelane area of the Mpumalanga Province, South Africa—R.G. BENGIS, D.F. KEET, A.L. MICHEL and N.P.J. KRIEK 239–241
- MINE, O.M.**
The establishment, composition and severity of infection of gastro-intestinal parasites and their impact on productivity of Tswana kids in southern Botswana—S.J. NSOSO, M.M. SENKU and O.M. MINE 71–74
The impact of traditional management on seasonal internal parasite burdens and productivity of indigenous Tswana goats in southern Botswana—S.J. NSOSO, J.B. MACHETE, M. MOLATOLE, R.T. NDEBELE, N.N. LEBANI, R.G. CHABO, A.M. KALAKE, L. JACYNA, B.W. SEGADIMO and O.M. MINE 101–104
- MINNAAR, W.N.**
Helminths in dogs belonging to people in a resource-limited urban community in Gauteng, South Africa—W.N. MINNAAR and R.C. KRECEK 111–117
- MITCHELL, J.O.**
Rapid detection and differentiation of Newcastle disease virus isolates by a triple one-step RT-PCR—Z. WANG, F.T. VREEDE, J.O. MITCHELL and G.J. VILJOEN 131–134
- MKARIA, J.**
Antibodies to Newcastle disease virus in the sera of indigenous chickens in Oodi, Kgatleng, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, J.M.K. HYERA, K.M. THIBANYANE and J. MKARIA 69–70
- MODUMO, J.**
A monovalent attenuated serotype 2 bluetongue virus vaccine confers homologous protection in sheep—P. HUNTER and J. MODUMO 331–333
- MOHAN, K.**
Vaccination to control an outbreak of *Mycoplasma crocodyli* infection—K. MOHAN, C.M. FOGGIN, F. DZIVA and P. MUVAVARIRWA 149–150
- MOLATOLE, M.**
The impact of traditional management on seasonal internal parasite burdens and productivity of indigenous Tswana goats in southern Botswana—S.J. NSOSO, J.B. MACHETE, M. MOLATOLE, R.T. NDEBELE, N.N. LEBANI, R.G. CHABO, A.M. KALAKE, L. JACYNA, B.W. SEGADIMO and O.M. MINE 101–104
- MOTA, R.A.**
Seroprevalence of bovine leptospirosis in Garanhuns municipal district, Pernambuco State, Brazil—A.A.F. OLIVEIRA, R.A. MOTA, G.C. PEREIRA, H. LANGONI, M.I. SOUZA, W.A. NAVAGANTES and M.E.P. SÁ 275–279
- MUKARATIRWA, S.**
Transmission studies on *Trichinella* species isolated from *Crocodylus niloticus* and efficacy of fenbendazole and levamisole against muscle L1 stages in Balb C mice—S. MUKARATIRWA, K. MAGWEDERE, E. MATENGA and C.M. FOGGIN 21–25
A survey of parasitic nematode infections of chickens in rural Zimbabwe—S. MUKARATIRWA, T. HOVE, J.B. ESMANN, C.J. HOJ, A. PERMIN and P. NANSEN 183–186
- MÜLDERS, M.S.G.**
Seasonal variation in cotyledoside concentration of *Tylecodon wallichii* (Harv.) Tölken subsp. *wallichii* sampled in a krimpsiekte-prevalent region—C.J. BOTHA, T. RUNDBERGET, A.L. WILKINS, M.S.G. MÜLDERS, A. FLÅØYEN and M.P. VAN AARDT 1–9
- MURAGURI, G.R.**
Immunization of cattle against East Coast fever using *Theileria parva* (Marikebuni) and relaxation of tick control in North Rift, Kenya—J.M. WANJOHI, J.N. NGERANWA, R.M. RUMBERIA, G.R. MURAGURI and S.K. MBOGO 217–223

MUSHI, E.Z.	
Antibodies to Newcastle disease virus in the sera of indigenous chickens in Oodi, Kgatleng, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, J.M.K. HYERA, K.M. THIBANYANE and J. MKARIA	69–70
Seroprevalence of <i>Toxoplasma gondii</i> and <i>Chlamydia psittaci</i> in domestic pigeons (<i>Columba livia domestica</i>) at Sebele, Gaborone, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, R. NDEBELE and R. PANZIRAH	159–161
MUVAVARIRWA, P.	
Vaccination to control an outbreak of <i>Mycoplasma crocodyli</i> infection—K. MOHAN, C.M. FOGGIN, F. DZIVA and P. MUVAVARIRWA	149–150
NANSEN, P.	
A survey of parasitic nematode infections of chickens in rural Zimbabwe—S. MUKARATIRWA, T. HOVE, J.B. ESMANN, C.J. HOJ, A. PERMIN and P. NANSEN	183–186
NAVEGANTES, W.A.	
Seroprevalence of bovine leptospirosis in Garanhuns municipal district, Pernambuco State, Brazil—A.A.F. OLIVEIRA, R.A. MOTA, G.C. PEREIRA, H. LANGONI, M.I. SOUZA, W.A. NAVEGANTES and M.E.P. SÁ	275–279
NDEBELE, R.	
Seroprevalence of <i>Toxoplasma gondii</i> and <i>Chlamydia psittaci</i> in domestic pigeons (<i>Columba livia domestica</i>) at Sebele, Gaborone, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, R. NDEBELE and R. PANZIRAH	159–161
NDEBELE, R.T.	
The impact of traditional management on seasonal internal parasite burdens and productivity of indigenous Tswana goats in southern Botswana—S.J. NSOSO, J.B. MACHETE, M. MOLATOLE, R.T. NDEBELE, N.N. LEBANI, R.G. CHABO, A.M. KALAKE, L. JACYNA, B.W. SEGADIMO and O.M. MINE	101–104
NEL, L.H.	
The 3A non-structural-protein coding region of the southern African SAT type isolates differs from that of other foot-and-mouth disease viruses—L.E. HEATH, H.G. VAN RENSBURG, W. VOSLOO and L.H. NEL	253–262
NGERANWA, J.N.	
Immunization of cattle against East Coast fever using <i>Theileria parva</i> (Marikebuni) and relaxation of tick control in North Rift, Kenya—J.M. WANJOHI, J.N. NGERANWA, R.M. RUMBERIA, G.R. MURAGURI and S.K. MBOGO	217–223
NSOSO, S.J.	
The establishment, composition and severity of infection of gastro-intestinal parasites and their impact on productivity of Tswana kids in southern Botswana—S.J. NSOSO, M.M. SENKU and O.M. MINE	71–74
The impact of traditional management on seasonal internal parasite burdens and productivity of indigenous Tswana goats in southern Botswana—S.J. NSOSO, J.B. MACHETE, M. MOLATOLE, R.T. NDEBELE, N.N. LEBANI, R.G. CHABO, A.M. KALAKE, L. JACYNA, B.W. SEGADIMO and O.M. MINE	101–104
OFFICE INTERNATIONAL DES EPIZOOTIES	
Comprehensive reports on technical items presented to the International Committee or to Regional Commissions 2000—OFFICE INTERNATIONAL DES EPIZOOTIES	163–164
OHALE, L.O.C.	
Closure of the ductus arteriosus of indigenous South African goats at high altitude—L.O.C. OHALE	47–53
OLIVEIRA, A.A.F.	
Seroprevalence of bovine leptospirosis in Garanhuns municipal district, Pernambuco State, Brazil—A.A.F. OLIVEIRA, R.A. MOTA, G.C. PEREIRA, H. LANGONI, M.I. SOUZA, W.A. NAVEGANTES and M.E.P. SÁ	275–279
ONGADI, F.A.	
Alternative cell lines for the propagation of lumpy skin disease virus—Y.S. BINEPAL, F.A. ONGADI and J.C. CHEP-KWONY	151–153
ORTIZ, J.	
Oral administration of mebendazole failed to reduce nematode egg shedding in captive African gazelles—J. ORTIZ, M.R. RUIZ DE YBÁÑEZ, T. ABAIGAR, M. GARIJO, G. ESPESO and M. CANO	79–82
PANZIRAH, R.	
Seroprevalence of <i>Toxoplasma gondii</i> and <i>Chlamydia psittaci</i> in domestic pigeons (<i>Columba livia domestica</i>) at Sebele, Gaborone, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, R. NDEBELE and R. PANZIRAH	159–161

- PEGRAM, R.G.**
Epidemiological observations of Zimbabwean theileriosis: Disease incidence and pathogenicity in susceptible cattle during *Rhipicephalus appendiculatus* nymphal and adult seasonal activities—A.A. LATIF, T. HOVE, G.K. KANHAI, S. MASAKA and R.G. PEGRAM 187–195
- PENRITH, M.-L.**
Multifactorial mortality in bongos and other wild ungulates in the north of the Congo Republic—F.W. HUCHZERMEYER, M.-L. PENRITH and P.W. ELKAN 263–269
- PEREIRA, G.C.**
Seroprevalence of bovine leptospirosis in Garanhuns municipal district, Pernambuco State, Brazil—A.A.F. OLIVEIRA, R.A. MOTA, G.C. PEREIRA, H. LANGONI, M.I. SOUZA, W.A. NAVAGANTES and M.E.P. SÁ 275–279
- PERMIN, A.**
A survey of parasitic nematode infections of chickens in rural Zimbabwe—S. MUKARATIRWA, T. HOVE, J.B. ESMANN, C.J. HOJ, A. PERMIN and P. NANSEN 183–186
- PHIRI, I.K.**
A survey on anthelmintic resistance in nematode parasites of sheep in Lusaka, Zambia—S. GABRIEL, I.K. PHIRI, P. DORNY and J. VERCRUYSSÉ 271–274
- PICARD, J.A.**
Prevalence of virulent *Rhodococcus equi* in isolates from soil collected from two horse farms in South Africa and restriction fragment length polymorphisms of virulence plasmids in the isolates from infected foals, a dog and a monkey—S. TAKAI, M.M. HENTON, J.A. PICARD, A.J. GUTHRIE, H. FUKUSHI and C. SUGIMOTO 105–110
- PROZESKY, L.**
Effect of pavetamine on protein synthesis in rat tissue—R.A. SCHULTZ, N. FOURIE, K.M. BASSON, L. LABUSCHAGNE and L. PROZESKY 325–331
- QUIRIN, R.**
Validity of intradermal tuberculin testing for the screening of bovine tuberculosis in Madagascar—R. QUIRIN, V. RASOLOFO, R. ANDRIAMBOLOLONA, A. RAMBOASOLO, T. RASOLONAVALONA, C. AHARISOLO, H. RAKOTOARITAHINA, S. CHANTEAU and P. BOISIER 231–238
- RAATH, J.P.**
The epidemiology of tuberculosis in free ranging African buffalo (*Syncerus caffer*) in the Kruger National Park, South Africa—V. DE VOS, R.G. BENGIS, N.P.J. KRIEK, A. MICHEL, D.F. KEET, J.P. RAATH and H.F.K.A. HUCHZERMEYER 119–130
- RAKOTOARITAHINA, H.**
Validity of intradermal tuberculin testing for the screening of bovine tuberculosis in Madagascar—R. QUIRIN, V. RASOLOFO, R. ANDRIAMBOLOLONA, A. RAMBOASOLO, T. RASOLONAVALONA, C. AHARISOLO, H. RAKOTOARITAHINA, S. CHANTEAU and P. BOISIER 231–238
- RAMBOASOLO, A.**
Validity of intradermal tuberculin testing for the screening of bovine tuberculosis in Madagascar—R. QUIRIN, V. RASOLOFO, R. ANDRIAMBOLOLONA, A. RAMBOASOLO, T. RASOLONAVALONA, C. AHARISOLO, H. RAKOTOARITAHINA, S. CHANTEAU and P. BOISIER 231–238
- RASOLOFO, V.**
Validity of intradermal tuberculin testing for the screening of bovine tuberculosis in Madagascar—R. QUIRIN, V. RASOLOFO, R. ANDRIAMBOLOLONA, A. RAMBOASOLO, T. RASOLONAVALONA, C. AHARISOLO, H. RAKOTOARITAHINA, S. CHANTEAU and P. BOISIER 231–238
- RASOLONAVALONA, T.**
Validity of intradermal tuberculin testing for the screening of bovine tuberculosis in Madagascar—R. QUIRIN, V. RASOLOFO, R. ANDRIAMBOLOLONA, A. RAMBOASOLO, T. RASOLONAVALONA, C. AHARISOLO, H. RAKOTOARITAHINA, S. CHANTEAU and P. BOISIER 231–238
- REGASSA, A.**
Tick infestation of Borana cattle in the Borana Province of Ethiopia—A. REGASSA 41–45
- ROUX, F.**
Lamproglana hoi n. sp. (Copepoda: Lernaieidae) from two yellowfish hosts, *Barbus marequensis* and *Barbus polylepis*, caught in a river in Mpumalanga, South Africa—S.M. DIPPENAAR, W.J. LUUS-POWELL and F. ROUX 209–215

RUIZ DE YBÁÑEZ, M.R.

Oral administration of mebendazole failed to reduce nematode egg shedding in captive African gazelles—J. ORTIZ, M.R. RUIZ DE YBÁÑEZ, T. ABAIGAR, M. GARIJO, G. ESPESO and M. CANO 79–82

RUMBERIA, R.M.

Immunization of cattle against East Coast fever using *Theileria parva* (Marikebuni) and relaxation of tick control in North Rift, Kenya—J.M. WANJOHI, J.N. NGERANWA, R.M. RUMBERIA, G.R. MURAGURI and S.K. MBOGO 217–223

RUNDBERGET, T.

Seasonal variation in cotyledoside concentration of *Tylecodon wallichii* (Harv.) Tölken subsp. *wallichii* sampled in a krimpsiekte-prevalent region—C.J. BOTHA, T. RUNDBERGET, A.L. WILKINS, M.S.G. MÜLDERS, A. FLÅØYEN and M.P. VAN AARDT 1–9

SÁ, M.E.P.

Seroprevalence of bovine leptospirosis in Garanhuns municipal district, Pernambuco State, Brazil—A.A.F. OLIVEIRA, R.A. MOTA, G.C. FERREIRA, H. LANGONI, M.I. SOUZA, W.A. NAVAGANTES and M.E.P. SÁ 275–279

SCHULTZ, R.A.

Effect of pavetamine on protein synthesis in rat tissue—R.A. SCHULTZ, N. FOURIE, K.M. BASSON, L. LABUSCHAGNE and L. PROZESKY 325–331

SEGADIMO, B.W.

The impact of traditional management on seasonal internal parasite burdens and productivity of indigenous Tswana goats in southern Botswana—S.J. NSOSO, J.B. MACHETE, M. MOLATOLE, R.T. NDEBELE, N.N. LEBANI, R.G. CHABO, A.M. KALAKE, L. JACYNA, B.W. SEGADIMO and O.M. MINE 101–104

SENKU, M.M.

The establishment, composition and severity of infection of gastro-intestinal parasites and their impact on productivity of Tswana kids in southern Botswana—S.J. NSOSO, M.M. SENKU and O.M. MINE 71–74

SOTIROV, L.

Changes in some factors of the innate immunity and serum zinc and iron concentrations in pigs following intravenous administration of *Escherichia coli* lipopolysaccharide—M. ANDONOVA, I. BORISSOV and L. SOTIROV 91–99

SOUZA, M.I.

Seroprevalence of bovine leptospirosis in Garanhuns municipal district, Pernambuco State, Brazil—A.A.F. OLIVEIRA, R.A. MOTA, G.C. PEREIRA, H. LANGONI, M.I. SOUZA, W.A. NAVAGANTES and M.E.P. SÁ 275–279

SPICKETT, A.M.

Parasites of domestic and wild animals in South Africa. XL. Ticks on dogs belonging to people in rural communities and carnivore ticks on the vegetation—I.G. HORAK, F.R. EMSLIE and A.M. SPICKETT 135–141

STARK, U.

Studies on the critical water mass, rehydration capability and potential, acute chill tolerance and supercooling point of *Argas (Persicargas) walkerae* (Acari: Argasidae)—U. STARK and R. GOTHE 11–20

SUGIMOTO, C.

Prevalence of virulent *Rhodococcus equi* in isolates from soil collected from two horse farms in South Africa and restriction fragment length polymorphisms of virulence plasmids in the isolates from infected foals, a dog and a monkey—S. TAKAI, M.M. HENTON, J.A. PICARD, A.J. GUTHRIE, H. FUKUSHI and C. SUGIMOTO 105–110

TAKAI, S.

Prevalence of virulent *Rhodococcus equi* in isolates from soil collected from two horse farms in South Africa and restriction fragment length polymorphisms of virulence plasmids in the isolates from infected foals, a dog and a monkey—S. TAKAI, M.M. HENTON, J.A. PICARD, A.J. GUTHRIE, H. FUKUSHI and C. SUGIMOTO 105–110

THIBANYANE, K.M.

Antibodies to Newcastle disease virus in the sera of indigenous chickens in Oodi, Kgatleng, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, J.M.K. HYERA, K.M. THIBANYANE and J. MKARIA 69–70

VAN AARDT, M.P.

Seasonal variation in cotyledoside concentration of *Tylecodon wallichii* (Harv.) Tölken subsp. *wallichii* sampled in a krimpsiekte-prevalent region—C.J. BOTHA, T. RUNDBERGET, A.L. WILKINS, M.S.G. MÜLDERS, A. FLÅØYEN and M.P. VAN AARDT 1–9

VAN RENSBURG, H.G.

The 3A non-structural-protein coding region of the southern African SAT type isolates differs from that of other foot-and-mouth disease viruses—L.E. HEATH, H.G. VAN RENSBURG, W. VOSLOO and L.H. NEL 253–262

VAN WYK, J.A.

Refugia—overlooked as perhaps the most potent factor concerning the development of anthelmintic resistance—J.A. VAN WYK 55–67

Clinical evaluation of anaemia in sheep: early trials—F.S. MALAN, J.A. VAN WYK and C.D. WESSELS 165–174

VERCRUYSSSE, J.

A survey on anthelmintic resistance in nematode parasites of sheep in Lusaka, Zambia—S. GABRIEL, I.K. PHIRI, P. DORNY and J. VERCRUYSSSE 271–274

VILJOEN, G.J.

Rapid detection and differentiation of Newcastle disease virus isolates by a triple one-step RT-PCR—Z. WANG, F.T. VREEDE, J.O. MITCHELL and G.J. VILJOEN 131–134

VOSLOO, W.

The 3A non-structural-protein coding region of the southern African SAT type isolates differs from that of other foot-and-mouth disease viruses—L.E. HEATH, H.G. VAN RENSBURG, W. VOSLOO and L.H. NEL 253–262

VREEDE, F.T.

Rapid detection and differentiation of Newcastle disease virus isolates by a triple one-step RT-PCR—Z. WANG, F.T. VREEDE, J.O. MITCHELL and G.J. VILJOEN 131–134

WANG, Z.

Rapid detection and differentiation of Newcastle disease virus isolates by a triple one-step RT-PCR—Z. WANG, F.T. VREEDE, J.O. MITCHELL and G.J. VILJOEN 131–134

WANJOHI, J.M.

Immunization of cattle against East Coast fever using *Theileria parva* (Marikebuni) and relaxation of tick control in North Rift, Kenya—J.M. WANJOHI, J.N. NGERANWA, R.M. RUMBERIA, G.R. MURAGURI and S.K. MBOGO 217–223

WESSELS, C.D.

Clinical evaluation of anaemia in sheep: early trials—F.S. MALAN, J.A. VAN WYK and C.D. WESSELS 165–174

WILKINS, A.L.

Seasonal variation in cotyledoside concentration of *Tylecodon wallichii* (Harv.) Tölken subsp. *wallichii* sampled in a krimpsiekte-prevalent region—C.J. BOTHA, T. RUNDBERGET, A.L. WILKINS, M.S.G. MÜLDERS, A. FLÅØYEN and M.P. VAN AARDT 1–9

WORTHINGTON, R.W.

A review of the infectious disease of African wild ruminants—R.W. WORTHINGTON and R.D. BIGALKE 291–323

ZWEYGARTH, E.

A chemically defined medium for the growth of *Cowdria ruminantium*—E. ZWEYGARTH and A.I. JOSEMANS 37–40

Continuous *in vitro* propagation of *Cowdria ruminantium* (Welgevonden stock) in a canine macrophage-monocyte cell line—E. ZWEYGARTH and ANTOINETTE I. JOSEMANS 155–157