

**PECTINOLYTIC BACTERIA ASSOCIATED WITH POTATO SOFT ROT AND
BLACKLEG IN SOUTH AFRICA AND ZIMBABWE**

**Elizabeth Ngadze¹, Carrie L. Brady², Teresa A. Coutinho³ and Jacquie E. van der
Waal⁴**

¹Department of Microbiology and Plant Pathology, University of Pretoria, Pretoria, 0002,
South Africa; and Department of Crop Science, University of Zimbabwe, P.O. Box MP167,
Mount Pleasant Harare, Zimbabwe

Fax: +27 12 420 4588

Email: engadze@agric.uz.ac.zw

²Department of Microbiology and Plant Pathology, Forestry and Agricultural Biotechnology
Institute (FABI), University of Pretoria, Pretoria, 0002, South Africa

Fax: +27 12 420 3960

Email: Carrie.Brady@fabi.up.ac.za

³Department of Microbiology and Plant Pathology, Forestry and Agricultural Biotechnology
Institute (FABI), University of Pretoria, Pretoria, 0002, South Africa

Fax: +27 12 420 3960

Email: Teresa.Coutinho@up.ac.za

⁴Department of Microbiology and Plant Pathology, University of Pretoria, Pretoria, 0002,
South Africa

Fax: +27 12 420 4588

Email: Jacquie.vdwaals@up.ac.za (Corresponding author)

ABSTRACT

Soft rot and blackleg can cause severe economic losses in potato production in South Africa and Zimbabwe depending on climatic conditions. The aim of the study was to identify the predominant bacteria causing potato soft rot and blackleg in these countries. Samples, comprising of stems and tubers from potato plants with blackleg and soft rot symptoms were collected throughout the 2008/9 growing season from potato production areas where disease outbreaks occurred. The isolates from these plants and tubers yielded Gram negative, pectinolytic bacteria on crystal violet pectate and inoculated tubers. Identification was based on biochemical and phenotypic characteristics, rep-PCR, Amplified Fragment Length Polymorphisms and sequences of *gyrB* and *recA* genes. Isolates from Zimbabwe were identified as *P. c.* subsp. *brasiliensis* (21 isolates), *Dickeya dadantii* subsp. *dadantii* (20 isolates), *P. c.* subsp. *carotovorum* (16 isolates) and *P. atrosepticum* (4 isolates). *Pcb*, *Pcc* and *Dd* subsp. *dadantii* were isolated from samples collected from all the regions, while *Pa* was isolated from Nyanga the coolest region in Zimbabwe. In South Africa, however, *Pectobacterium carotovorum* subsp. *brasiliensis* (*Pcb*) was the most common causal agent of soft rot and blackleg. *P. atrosepticum* was the only pathogen isolated from samples collected in Nyanga, Zimbabwe and was not isolated from any South African samples. AFLP analysis separated the *Pcb* strains into 12 clusters, reflecting subdivision in terms of geographic origin, and *Pectobacterium carotovorum* subsp. *carotovorum* (*Pcc*) isolates were clearly differentiated from *Pcb* isolates. A large degree of DNA polymorphism was evident between these 12 clusters. The study identified all the pathogens associated with the blackleg / soft rot disease complex.

Keywords: *Erwinia*, genetic diversity, pathogenicity, soft rot, *Solanum tuberosum*

INTRODUCTION

The major source of tuber and stem rot in potatoes in tropical and subtropical climates is contamination by one or a combination of soft rotting pectinolytic enterobacteria. These bacteria were until recently classified within the genus *Erwinia*, which was divided into various species and subspecies on the basis of molecular, biochemical and host range differences (Lelliot and Dickey 1984, Gallois et al. 1992, Hélias et al. 1998). Subsequent revisions have led to the taxonomic reclassification of pectinolytic *Erwinia* into several genera. Strains formerly described as *Erwinia carotovora* have been incorporated into the genus *Pectobacterium*, and strains classified as *Erwinia chrysanthemi* are now assigned to the genus *Dickeya* (Hauben et al. 1998, Gardan et al. 2003, Samson et al. 2005).

Pectobacterium atrosepticum almost exclusively infects potato, causing blackleg of the stem and tuber soft rot. Blackleg is an economically important disease and is characterised by the blackening of the stem base of potato plants, which originates from the rotting mother tuber (Pèrombelon 2002, Czajkowski et al. 2011). Blackleg is unpredictable in the field due to the ability of the pathogen to remain latent in the seed tuber (Laurila et al. 2008). *Pa* is the causal agent of blackleg disease in many geographical areas (Duarte et al. 2004).

By contrast *Pectobacterium carotovorum* subsp. *carotovorum* (*Pcc*) has a broad host range, causing soft rot disease in various crops including potato, carrot, capsicum and calla lily (Wright 1998, Toth et al. 2003). *Pcc* can also be virulent in temperate climates causing blackleg symptoms (De Haan et al. 2008). Another atypical, highly virulent enterobacterial subspecies, *Pectobacterium carotovorum* subsp. *brasiliensis* (*Pcb*) was recently described in Brazil and South Africa. The pathogen causes severe soft rotting of tubers and blackleg of the stems (Duarte et al. 2004, van der Merwe et al. 2010).

The genus *Dickeya* is extremely diverse and comprised of species that are pathogenic to many important crops and ornamental plants, mostly in tropical and subtropical climates. Species belonging to this genus cause soft rot and blackleg on potato in temperate regions (Pèrombelon 2002, Toth et al. 2011, Tsrer et al 2012). The species structure of *Dickeya* is well defined, but not all species have been described and there is considerable overlap in host range. Isolates infecting potato are now distributed among the six described species of *Dickeya* (Samson et al. 2005) and it remains unclear whether these strains produce the same symptoms during disease development, or whether only some are responsible for blackleg-like symptoms. The discovery of distinct species of *Pectobacterium* and *Dickeya* associated with certain hosts supports the idea that identifiable groups of pectinolytic enterobacteria are found in specific ecological niches and may be responsible for variable disease symptoms on potato depending on the prevailing environmental conditions (De Boer 2003, Czajkowski et al. 2011).

Pectobacterium atrosepticum and *Pectobacterium carotovorum* subsp. *carotovorum* have been identified as the causal agents of potato blackleg and soft rot diseases in Zimbabwe (Masuka et al. 1998, Manzira 2010). Recently *D. dadantii* subsp. *dadantii* was isolated from plants with typical wilting symptoms and rotting tubers (Ngadze et al. 2010). *Pcb* is the main causal agent of potato blackleg and soft rot diseases in South Africa (van der Merwe, 2010).

Detection and differentiation of pathogen strains are important aspects of disease management strategies and several molecular and biochemical tools have been developed for the identification of pectinolytic enterobacteria (De Boer and McNaughton 1987, De Boer and Ward 1995, Kang et al. 2003). PCR restriction fragment length polymorphism (RFLP) and amplified fragment length polymorphism (AFLP) techniques have also been used in the study of genetic diversity (Toth et al. 2001, Avrova et al. 2002). AFLP analysis has demonstrated that a greater diversity of pectinolytic enterobacteria infect potato than

previously thought (Avrova et al. 2002, Pitman et al. 2010). Repetitive extragenic palindromic-PCR (rep-PCR), which can differentiate between closely related strains of bacteria, is considered one of the most effective methods to differentiate between bacterial species (Rademaker et al. 2004). In rep-PCR DNA fingerprinting, PCR amplification of the DNA between adjacent repetitive extragenic elements is used to obtain strain-specific DNA fingerprints, which can be easily analysed with pattern recognition computer software (Rademaker et al. 2004).

Pcb has recently been identified in South Africa and Zimbabwe as one of the primary causal agents of the blackleg / soft rot complex. This was the first report of the pathogen affecting potato in both of these countries (van der Merwe et al. 2010). As Zimbabwe has been importing potato seed from South Africa for almost a decade, it is of great interest to examine the genetic diversity of *Pcb* isolates from these two countries.

The objectives of the study were (i) to isolate and identify the causal agents of potato soft rot and blackleg in South Africa and Zimbabwe by rep-PCR and sequences of two genes involved in bacterial metabolic processes, *recA* (encoding recombinase A) and *gyrB* (encoding DNA gyrase), and (ii) determine and compare the genetic diversity of *Pcb* isolates from South Africa to those occurring in Zimbabwe using AFLP analysis. .

MATERIALS AND METHODS

Isolation of bacterial strains

The stems and tubers of potato plants with soft rot and blackleg symptoms (soft rot, wilting, internal and external darkening on stems) were collected throughout the 2008/9 growing season from six production areas where soft rot outbreaks occurred in Zimbabwe. Severe

outbreaks occurred in Darwendale, Gwebi, Harare, Marondera, Mazowe and Nyanga. South African isolates came from the Eastern Cape, Eastern Freestate, KwaZulu-Natal, Limpopo, Mpumalanga, Northern Cape, North West, Sandveld and Western Freestate. Isolates from South Africa and mini-tubers imported from China and India (various varieties; sources not identified) were obtained from the Potato Pathology Program at the University of Pretoria culture collection. The isolates in the culture collection were collected from all growing regions in South Africa, during most growing seasons, from 2005 – 2009. Pieces of infected stems and tubers were macerated in 0.01M magnesium sulphate (MgSO₂). Isolations were made on the selective medium, Crystal Violet Pectate (CVP) (Hyman et al. 2001) and plates were incubated at 25°C for 48h. Isolates that tested positive for pectinolytic cavity formation (formed pits on CVP medium) were purified on the same medium and then transferred to nutrient agar (NA) (Merck, Darmstadt, Germany), on which they were maintained. All isolates were stored in sterile water at room temperature and in 15% (v/v) Nutrient Agar glycerol medium at -80°C and maintained in the Potato Pathology Programme culture collection at the University of Pretoria, Pretoria, South Africa. A list of isolates and type strains of the various species and subspecies of *Pectobacterium* and *Dickeya* used in this study is shown in Table 1.

Biochemical and physiological identification of isolates

Single colonies of each isolate were used for biochemical tests. The isolates were identified by standard bacterial methods based on Cother and Sivasithamparam (1983). The tests performed were Gram reaction, oxidase activity, glucose metabolism, pectate degradation in Sutton's medium, production of phosphatase, indole production from tryptophan, gelatine hydrolysis, production of reducing substances from sucrose, production of acid from α -methylglucoside and trehalose, malonate utilisation, sensitivity to erythromycin (15 μ g),

growth at 37°C which was determined after 24h in nutrient broth (NB, Difco), salt tolerance which was checked after 48h growth in NB with 5g.l⁻¹ NaCl.

Rep-PCR analyses

Whole genome rep-PCR fingerprints were generated from all strains (including type strains and reference strains) according to Rademaker et al. (2004). Cluster analysis of rep-PCR profiles was performed in BioNumerics 4.0 (Applied Maths) using Pearson's correlation indices of similarity with the unweighted pair-group method using arithmetic averages (UPGMA). The patterns of each of the independently processed strains analyzed on independent gels always shared > 70% similarity. Thus, the cut-off used to define rep types was 70%.

Sequencing and phylogenetic analysis

Genomic DNA was extracted from overnight broth cultures using a DNeasyTM Blood and Tissue Kit (Qiagen, Southern Cross Biotechnology). For selected bacterial strains, the *recA* and *gyrB* genes were amplified using *recAF*, *recAR* and *gyrB-01F*, *gyrB-02F* respectively (Table 2). PCR amplification was performed with a DNA Thermal Cycler 480 (Perkin-Elmer Corp). PCR products were purified using a QIAquickTM Purification Kit (Qiagen, Southern Cross Biotechnology). The *gyrB* and *recA* genes were sequenced in both directions using primers and conditions previously described (Young and Park 2007, Brady et al. 2008). Sequencing products were cleaned using a sodium acetate precipitation step and sequenced on an ABI Prism DNA Automated Sequencer (Perkin Elmer). Incorrect basecalls were corrected using Chromas Lite v 2.01. The corrected sequences were edited and aligned with BioEdit Sequence Alignment v 7.0.0 and both ends were trimmed to the following final sizes *recA* 697 positions and *gyrB*, 740 positions. Searches were performed on each consensus sequence generated in BioEdit using the BLAST algorithm from GenBank.

All the selected sequences were aligned using MAFFT. Phylogenetic trees for the individual gene sets were inferred by using the neighbour-joining programme of MEGA v 4 (Tamura et al. 2007). The stability of the phylogenetic trees was assessed by performing bootstrap analysis of the neighbour-joining data based on 1 000 bootstrap replicates (Kwon et al. 1997).

Pectinolytic activity

Preliminary test on potato tuber slices Fourteen representative samples from Zimbabwe were selected from the five growing regions and used in this assay. The isolates used were: EN4, EN6, EN16, EN31, EN38, EN44, EN47, EN54, EN58, EN59, EN60, EN63 and EN66. Type strains of the various species and subspecies of *Pectobacterium* and *Dickeya* were also included in this assay (Table 1). All the bacterial isolates were tested for pectinolytic activity on potato tubers (*Solanum tuberosum* cv Amethyst). The tubers were washed under running tap water, air dried and then dipped in 96% ethanol, flamed and cut into 10 mm thick slices. The slices were placed on wet sterile filter papers in Petri dishes. Two loops of each bacterial isolate grown on Nutrient Agar were uniformly spread on the upper surface of 20 slices. Development of rot on the slices was examined 24 – 48h after incubation at 25°C. The number of rotten slices was expressed as a percent of the total number of inoculated slices. Control tubers were treated in the same manner but sterile distilled water was used in place of the bacterial isolate.

Quantitative test on potato tuber slices using filter paper discs This test was performed as described by Wegener (2002). A filter paper disc 10mm in diameter, previously dipped in a water suspension of each bacterial isolate containing 10^8 cfu.ml⁻¹, was placed at the centre of the upper surface of each potato slice. Each isolate was tested on 20 potato slices. After incubation at 25°C for 2 days, the rotting zone diameter on each tissue slice was measured. For the control, tubers were inoculated with filter paper discs soaked in sterile distilled water.

Quantitative test on whole tubers using filter paper discs Pectinolytic assays were performed on whole potato tubers. For tuber maceration assays, potatoes (cv. Amethyst) were surface sterilised by submersion in 10% sodium hypochlorite for 10min and each potato inoculated with 10µl of a 48h culture of each strain grown in Luria-Bertani broth. Holes of a fixed depth of 20mm were stabbed into the potatoes using a sterile pipette tip, filled with inoculum and then sealed with Vaseline. The inoculated potato was then wrapped in a plastic bag and incubated at 25°C for 72h. Rotting tissue was scraped from the potato and weighed to establish the degree of potato maceration.

Fluorescent amplified fragment length polymorphism analysis

The diversity of *Pcb* causing soft rot of potatoes in Zimbabwe and South Africa was evaluated using AFLP fingerprinting. Sixty-four *Pcb*, 4 *Dd* and 4 *Pcc* isolates from 16 different South African and Zimbabwean potato-growing regions were used in the analysis. Fluorescent AFLP analysis was performed according to the method described by Pitman et al. (2010). AFLP products were visualised by a PRISM 310 Genetic Analyser (Applied Biosystems) using the GeneScan 500 size standard (Applied Biosystems). All primers used in AFLP reactions are listed in Table 2. Band patterns were analysed with BioNumerics 4.0 (Applied Maths) and compared with a database containing profiles of reference strains of all *Pectobacterium* and *Dickeya* species with validly published names. A UPGMA dendrogram was constructed using Pearson's correlation coefficient.

RESULTS

Biochemical and physiological characterisation

The analysis of the biochemical properties of the enterobacterial isolates from potatoes in Zimbabwe indicated that the majority (35% of isolates) were *D. dadantii* subsp. *dadantii* (Table 3). In addition four isolates were identified as *Pectobacterium atrosepticum* and 21 as *Pectobacterium carotovorum* subsp. *carotovorum*. *Pcb* could not be differentiated from *Pcc* and *D. dadantii* subsp. *dadantii* based on the biochemical properties examined. Most of the isolates, except the putative *Pectobacterium atrosepticum* and the reference sample of the same species, grew at 37°C supporting the provisional classification of the majority as *Dickeya dadantii* subsp. *dadantii* and *Pcc*.

Rep-PCR analyses

The rep-PCR fingerprint patterns consisted of 10 to 45 PCR products, ranging in size from about 200 to 10 000bp. When analyzed using the Pearson's correlation coefficient, the rep-PCR fingerprints had an average similarity of 86%. Cluster analysis of the profiles revealed four groups, which separated the isolates into *Pa*, *Pcc*, *Pcb* and *Dd* (Fig. 1) The reference isolates for each of these species (LMG 2404^T, LMG 2386^T, Ech 3937 and ATCC BAA 417) fell into their respective clusters and grouped with the appropriate isolates. Polymorphic patterns sharing multiple bands of equal mobility could be distinguished for some of the isolates.

Sequencing and phylogenetic analysis

Partial *recA* and *gyrB* gene sequences of the potato isolates confirmed their identity as belonging to species of *Pectobacterium* and *Dickeya*. Phylogenetic analyses grouped the strains and reference samples into four distinct clusters in both trees (Fig.2, Suppl. Fig. 1). In

the *gyrB* phylogenetic tree (Fig. 2), all strains identified biochemically as *Dickeya dadantii* subsp. *dadantii* clustered in a single clade, containing the type strain and reference strains, which was supported by a bootstrap value of 86%. Fifteen strains formed a stable cluster with reference strains of *Pectobacterium carotovorum* subsp. *brasiliensis* (ATCC BAA-417, ATCC BAA-419, LMG 21370 and LMG 21371) with a bootstrap value of 82%. *RecA* and *gyrB* sequencing also confirmed the identity of both *P. atrosepticum* and *P. c.* subsp. *carotovorum* strains, as the isolates fell into separate distinct clusters with their respective type and reference strains supported by high bootstrap values in both trees.

Pectinolytic activity

All isolates evaluated showed pectinolytic activity on potato tuber slices. *Pa* (LMG 2386^T, EN54), *Pcb* (ATCC BAA-417, EN16, EN66), *Pcc* (EN31) and *Dd* (EN47, EN58) caused rots which penetrated the entire thickness of potato slices within 24 h whereas isolates LMG 2404^T (*Pcc*), *E. c* 3937^T, EN38, EN44, EN59 (*Dd*), EN4, EN6 (*Pcb*) and EN60 (*Pa*) caused the same effect in 48 h (Table 4). Evaluation of rotting zone diameters after 4 days showed variable responses. The largest rotting zone diameter of 36 mm was observed in potato slices inoculated with LMG 2386^T, EN54 (*Pa*), ATCC BAA-417, EN66 (*Pcb*), EN 12 (*Dd*) and EN63 (*Pcc*). The reference sample LMG 2404^T (*Pcc*) and EN6 (*Pcb*) recorded the smallest rotting zone diameters (Table 4).

The relative pectinolytic activity of strains was quantified using whole tuber assays. The amount of tissue macerated was notably higher for EN12 (*Dd*) and EN54 (*Pa*). These strains also showed greater levels of maceration than the reference samples used in the study (Fig. 3). Pectinolytic activity was not consistent within groups of genetically and biochemically similar strains, for example, EN54 and EN60 which are *Pa* isolates, exhibited different levels of maceration on the potato tubers (Fig. 3). The tuber maceration results were supported by

the results of the quantitative pectinolytic activity in which the rotting zone diameters were significantly ($P < 0.05$) larger on potato slices inoculated with EN12 and EN54 (Table 4). The reference samples were not as virulent as strains EN12 (*Dd*) and EN54 (*Pa*). EN44 was the least pectinolytic strain, as it showed very little maceration of tubers in the virulence assays and a relatively small rotting zone diameter for the quantification of pectinolytic activity.

Fluorescent amplified fragment length polymorphism

The restriction enzymes *EcoRI* and *MseI* together with selective primers *E00* and *MSeI-C* generated between 40 and 60 distinguishable bands within a range of 40 – 550bp for each isolate tested (Fig. 4). Following numerical analysis of the AFLP banding profiles by UPGMA, the isolates clustered according to identity as determined by rep-PCR and *gyrB*- and *recA*-gene sequencing. *D. dadantii* subsp. *dadantii*, *Pcb* and *Pcc* isolates clustered separately in different clades. Within the *Pcb* clade, isolates grouped into 12 distinct clusters, several according to geographical origin, at a similarity level of 70%. The reference strain ATCC BAA419 grouped with the isolates in cluster 1. In clusters 1, 2, 3 and 7 isolates from South Africa clustered with those from Zimbabwe. Clusters 4 and 10 contained isolates from Limpopo and Mpumalanga provinces respectively (South Africa), while clusters 11 and 12 were exclusively made up of isolates from the Sandveld region, also in South Africa. Isolates from Zimbabwe (Marondera and Harare) grouped together in cluster 5, while clusters 6 and 9 contained isolates from Zimbabwe, China and India. Cluster 8 included isolates from South Africa, Zimbabwe and China. *Pcc* isolates clustered at a similarity level of 75%, while *Dd* isolates clustered at a relatively low similarity level of 30%.

DISCUSSION

The results of biochemical, physiological and genetic analyses revealed the identity of pectinolytic enterobacteria infecting potatoes in Zimbabwe and South Africa. The majority of isolates in the collection were shown to be *D. dadantii* subsp. *dadantii* (35% of isolates), followed by *Pectobacterium carotovorum* subsp. *brasiliensis* (25%). *Pectobacterium carotovorum* subsp. *carotovorum* and *Pectobacterium atrosepticum* were also identified, but in lower quantities.

Dickeya chrysanthemi (formerly *Erwinia chrysanthemi*) is the causal agent of tuber soft rot and blackleg-like symptoms in many potato growing areas of the world (Young et al. 1992, Toth et al. 2011, Tsrer et al. 2012). However, our study found *D. dadantii* subsp. *dadantii* to be the most predominant species in Zimbabwe. This finding was not surprising because *D. dadantii* subsp. *dadantii* is a warm climate pathogen (Pèrombelon 2002) and confirms earlier findings of Ngadze et al. (2010). The strains isolated in this study shared biochemical and genetic features with the species *D. dadantii*. Phylogenetic analysis of the *recA* and *gyrB* gene sequences demonstrated high genetic homogeneity between the potato isolates from Zimbabwe and *D. dadantii* subsp. *dadantii* reference strains, with sequence similarity ranging from 86 to 97%.

Pectobacterium carotovorum subsp. *carotovorum* is prevalent on potatoes worldwide (Laurila et al. 2008). This pathogen has been identified and reported as a causal agent of potato soft rot in Zimbabwe (Masuka et al. 1998), and the present study confirms this, although *Pcc* was isolated less frequently than *Pcb*. The ubiquitous nature of *Pcc* may be due to an earlier divergence, wider geographical distribution or broader host range and it is thought to have resulted in the extensive genetic variability of this species (Avrova et al. 2002). These observations are supported by genetic analysis of *Pcc* isolates in the present

study. Phylogenetic analysis of partial *gyrB* sequences demonstrated a high genetic divergence between potato isolates from Zimbabwe and reference strains of *Pcc*. The type strain of *Pcc* (LMG 2404T) clustered with a reference strain (LMG 2407) on the border of *Pcb*, while the Zimbabwe potato isolates formed a strongly supported cluster with a second reference strain on a separate branch at a lower level. The polyphyletic arrangement of *Pcc* suggests that another subspecies may exist within *Pectobacterium carotovorum*, although a greater number of strains would have to be examined to provide further proof.

Pectobacterium carotovorum subsp. *brasiliensis* is the most important causal agent of blackleg in South Africa and it is widely distributed in the country (van der Merwe et al. 2010). The bacterium could have entered Zimbabwe through import of contaminated seed tubers since Zimbabwe has been importing seed and table potatoes from South Africa. Other researchers have also highlighted the importance of seed tubers in the transmission of blackleg disease (Czajkowski et al. 2009, 2011).

Pectobacterium atrosepticum has been identified as the causal agent of potato blackleg in Zimbabwe (Masuka et al. 1998). The pathogen almost exclusively infects potato causing blackleg of the stem and tuber soft rot (Pèrombelon 2002). *Pa* was isolated from samples which came from Nyanga and Marondera, which are the cooler potato growing regions in Zimbabwe. The low abundance of *P. atrosepticum* compared to *Pcb* and *Pcc* may be due to limited host range, temperature requirements and origin (Avrova 2002).

Several studies have shown the usefulness of the rep-PCR technique to fingerprint a large variety of bacteria as well as its ability to differentiate at the species, pathovar or strain level (Versalovic et al. 1991, Rademaker et al. 2004). Therefore selected isolates from potatoes in Zimbabwe were characterized by rep-PCR fingerprinting of genomic DNA. The technique clustered the isolates in this study into four groups, each containing a reference strain.

Isolates identified as either *Dd* or *Pa* produced fingerprints which were nearly identical to those of their respective reference strains, suggesting a high level of homogeneity. In contrast, several visually different fingerprints were observed in both the *Pcc* and *Pcb* clusters providing further evidence of the genetic divergence of this species.

All potato isolates tested were pectinolytic on potato tuber slices and CVP medium, showing that they produce cell wall degrading enzymes. These enzymes break down pectin and release nutrients which are essential for microbial growth. The effect of these enzymes results in tissue disintegration, cell damage and seepage of cell contents (Pèrombelon 2002), which produces the disease symptoms seen on plants and tubers. The isolates exhibited differences in pectinolytic activity which can be attributed to their origin. The isolates are from different geographical locations and were isolated from different potato cultivars. Other studies have also reported a link between differences in virulence and taxonomic diversity based on location and host (Pitman et al. 2010, Tsrör et al. 2012).

AFLP analysis separated the *Pcb* strains into 12 clusters, reflecting subdivision in terms of geographic origin, and *Pcc* isolates were clearly differentiated from *Pcb* isolates, which displayed clearer and more intense banding profiles. Such intraspecies diversity has been noted in *Pcc* isolates analyzed with PCR-RFLP (Darrasse et al. 1994, Hèlias et al. 1998) and this may reflect molecular diversity within subspecies. Four clusters contained isolates from both South Africa and Zimbabwe, four clusters contained isolates only from South Africa and one cluster contained isolates only from Zimbabwe. A large degree of DNA polymorphism was evident between these 12 clusters, suggesting that several different *Pcb* populations, derived from a common ancestor, are causing the blackleg symptoms seen in these two countries. It also appears that there are two different lineages of *Pcb* in the Sandveld region (South Africa) as clusters 11 and 12 contain isolates from this region that are genotypically different. As Zimbabwe has been importing potato seed from South Africa for many years, it

is possible that *Pcb* was introduced to Zimbabwe on infected seed. This would explain the similar banding patterns of South African and Zimbabwean isolates within clusters 1, 2, 3 and 7.

AFLP analysis provides a basis for an evolutionary analysis of *Pcb* pathotypes. Most strains collected from the same geographical area clustered together in the UPGMA tree, showing that *Pcb* populations from each area represent a distinct phenetic group. However in some cases, where numerous phenetic groups exist, it is possible that different founder effects occurred, probably because different potato cultivars are grown in diverse geographical areas in South Africa and Zimbabwe. The inconsistencies within some clusters (isolates from different geographical areas grouping together) may also reflect genetic exchanges that have occurred between groups.

Despite the genetic diversity evident in *Pcb*, a characteristic pattern of 2 to 3 bands was clearly visible in all isolates, providing potential molecular markers for identification and diagnosis. In most cases the subspecies-specific bands appeared more intense than others. This could be attributed to their higher genomic copy numbers (Avrova et al. 2002) and it has been suggested that rDNA genes could be the source of these bands (Keim et al. 1997).

In conclusion, *D. dadantii* subsp. *dadantii* was the most frequently isolated pectinolytic pathogen from potatoes in Zimbabwe, followed by *P. carotovorum* subsp. *brasiliensis*. *P. carotovorum* subsp. *carotovorum* and *P. atrosepticum* were also isolated but in lower frequencies. The majority of isolates collected in South Africa in the 2006/7 growing seasons were *P. carotovorum* subsp. *brasiliensis*, which is widely spread throughout the potato production regions. Both *gyrB* and *recA* sequencing could reliably identify isolates to the subspecies level, although *gyrB* demonstrated a greater intraspecies diversity. Selected isolates belonging to the four taxa could also be differentiated by rep-PCR and a high degree

of genetic diversity was evident for *Pcc* and *Pcb*. The genetic diversity in *Pcb* was also observed by AFLP analysis which showed great discriminatory power by separating isolates into 12 different clusters. The rep-PCR and AFLP results did not correlate in the present study and more strains must be analyzed by rep-PCR to determine the existence or lack of correlation between these techniques.

ACKNOWLEDGEMENTS

This work was funded by Potatoes South Africa and the National Research Foundation of South Africa. The authors would like to acknowledge the help of Mr Charles Wairuri in sequence alignment and drawing of phylogenetic trees.

REFERENCES

- Avrova, A. O., Hyman, L. J., Toth, R. L. and Toth, I. K. (2002). Application of amplified fragment length polymorphism fingerprinting for taxonomy and identification of the soft rot bacteria *Erwinia carotovora* and *Erwinia chrysanthemi*. *Applied Environmental Microbiology*, 68, 1499-1508
- Brady, C. L., Cleenwerck, I., Venter, S. N., Vancanneyt, M., Swings, J. and Coutinho, T. A. (2008). Phylogeny and identification of *Pantoea* species associated with the environment, humans and plants based on multilocus sequence analysis (MLSA). *Systematic Applied Microbiology*, 31, 447-460

- Cother, E. J. and Sivasithamparam, K. (1983). *Erwinia*: the “carotovora” group. (In P. C. Fahy and G. J. Persley (Eds) *Plant Bacterial Diseases. A Diagnostic Guide*, (pp. 87-101). Sydney: Academic Press)
- Czajkowski, R., Grabe, G. J. and Van der Wolf, J. M. (2009). Distribution of *Dickeya* spp. and *Pectobacterium carotovorum* subsp. *carotovorum* in naturally infected seed potatoes. *European Journal of Plant Pathology*, 125, 263-75
- Czajkowski, R., Pérombelon, M. C. M., van Veen, J. A. and van der Wolf, J. M. (2011). Control of blackleg and tuber soft rot of potato caused by *Pectobacterium* and *Dickeya* species: a review. *Plant Pathology*, 60, 999-1013
- Darrasse, A., Priou, S., Kotoujansky, A. and Bertheau, Y. (1994). PCR and restriction fragment length polymorphism of a *pel* gene as a tool to identify *Erwinia carotovora* in relation to potato diseases. *Applied Environmental Microbiology*, 60, 1437-1447
- De Boer, S. H. and McNaughton, M. E. (1987). Monoclonal antibodies to the lipopolysaccharide of *Erwinia carotovora* subsp. *atroseptica* serogroup-1. *Phytopathology*, 77, 828-832
- De Boer, S. H. and Ward, L. J. (1995). PCR detection of *Erwinia carotovora* subsp. *atroseptica* associated with potato tissue. *Phytopathology*, 85, 854-858
- De Boer, S. H. (2003). Characterisation of pectolytic erwinias as highly sophisticated pathogens of plants. *European Journal of Plant Pathology*, 109, 893-899
- De Haan, E. G., Dekker-Nooren, T. C. E. M., Van den Bovenkamp, G. W., Speksnijder, A. G. C. L., van der Zouwen, P. S. and Van der Wolf, J. M. (2008). *Pectobacterium*

carotovorum subsp. *carotovorum* can cause potato blackleg in temperate climates. *European Journal of Plant Pathology*, 122, 561-568

Duarte, V., De Boer, S. H., Ward, L. J. and De Oliveira, A. M. R. (2004). Characterization of atypical *Erwinia carotovora* strains causing blackleg of potato in Brazil. *Journal of Applied Microbiology*, 96, 535-5

Gallois, A., Samson, R. Ageron, E. and Grimont, P. A. D. (1992). *Erwinia carotovora* subsp. *odorifera* subsp. nov associated with odorous soft rot of chicory (*Gichorium intybus* L). *International Journal of Systematic Bacteriology*, 4, 582-588

Gardan, L., Gouy, C., Christen, R. and Samson R. (2003). Elevation of three subspecies of *Pectobacterium carotovora* to species level: *Pectobacterium atrosepticum* sp. nov., *Pectobacterium betavascularum* sp. nov. and *Pectobacterium wasabiae* sp. nov.. *International Journal of Systematic Evolutionary Microbiology*, 53, 381-391

Hauben, L., Moore, E. R., Vauterin, L., Steenackers, M., Mergaert, J., Verdonck, L. and Swings, J. (1998). Phylogenetic position phytopathogens within *Enterobacteriaceae*. *Systematic and Applied Microbiology*, 21, 384 – 397.

Hèlias, V., Le Roux, A. C., Bertheau, Y., Andrivon, D., Gauthier, J. P. and Jouan, B. (1998). Characterization of *Erwinia carotovora* subspecies and detection of *Erwinia carotovora* subspecies *atroseptica* in potato plants, soil and water extracts with PCR-based methods. *European Journal of Plant Pathology*, 104, 685-699

Hyman, L. J., Sullivan, L., Toth, I. K., and Pérombelon, M. C. M. (2001). Modified crystal violet pectate medium (CVP) based on a new polypectate source (Slendid) for the detection and isolation of soft rot erwinias. *Potato Research*, 44, 265-270

- Kang, H. W., Kwon, S. W. and Go, S. J. (2003). PCR-based specific and sensitive detection of *Pectobacterium carotovorum* ssp. *carotovorum* by primers generated from a URP-PCR fingerprinting-derived polymorphic band. *Plant Pathology*, 52, 127-133
- Keim, P., Kalif, A., Schupp, J., Hill, K., Travis, S. E., Richmond, K., Adair, D. M., Hugh-Jones, M., Kuske, C. R., and Jackson, P. (1997). Molecular evolution and diversity in *Bacillus anthracis* as detected by amplified fragment length polymorphism markers. *Journal of Bacteriology*, 179, 818-824
- Kwon, S., Go, S., Kang, H. Ryu, J., and Jo, J. (1997). Phylogenetic analysis of *Erwinia* species based on 16S rRNA gene sequences. *International Journal of Systematic Bacteriology*, 47, 1061-1067
- Laurila, J., Ahola, V., Lehtinen, A., Joutsjoki, T., Hannukkala, A., Rahkonen, A. and Pirhonen, M.. (2008). Characterization of *Dickeya* strains isolated from potato and river water samples in Finland. *European Journal of Plant Pathology*, 122, 213-225
- Lelliot, R. A. and Dickey, R. S. (1984). Genus VII. *Erwinia* Winslow, Broadhurst, Buchanan, Krumwiede, Rogers and Smith 1920. (In Krieg, N.R. and Holt, J.G. Baltimore (Eds) *Bergey's Manual of Systematic Bacteriology*, 1, 469-476
- Manzira, C. (2010). Potato production handbook. Potato Seed Association Zimbabwe. (Harare: Jongwe Printers)
- Masuka, A. J., Cole D. L., and Mguni, C. (Eds.) (1998). *List of plant diseases in Zimbabwe*. (Harare: Jongwe Printers)
- Ngadze, E., Coutinho, T .A. and van der Waals, J. E. (2010). First report of soft rot of potatoes caused by *Dickeya dadantii* in Zimbabwe. *Plant Disease*, 94, 1263

- Pérombelon, M. C. M. (2002). Potato diseases caused by soft rot erwinias: an overview of pathogenesis. *Plant Pathology*, 51, 1-12
- Pitman, A. R., Harrow, S. A. and Visnovsky, S. B. (2010). Genetic characterisation of *Pectobacterium wasabiae* causing soft rot disease of potato in New Zealand. *European Journal of Plant Pathology*, 126, 423-435
- Rademaker, J. L. W., Louws, F. J. and de Bruijn F. J. (2004). Characterization of the diversity of ecological important microbes by rep-PCR genomic fingerprinting. In: de Bruijn, F.J., Head, I.M., Akkermans, A.D. and van Elsas, J.D. (Eds.) *Molecular Manual*, (pp. 1 – 33) (Dordrecht: Kluwer)
- Samson, R., Legendre, J. B., Christen, R., Fischer-Le Saux, M., Achouak, W. and Gardan, L. (2005). Transfer of *Pectobacterium chrysanthemi* (Burkholder *et al.*, 1953) Brenner *et al.*, 1973 and *Brenneria paradisiaca* to the genus *Dickeya* gen. nov. known as *Dickeya chrysanthemi* comb. nov and *Dickeya paradisiaca* combi. nov. and delineation of four novel species, *Dickeya dadantii* sp nov., *Dickeya dianthicola* sp. nov., *Dickeya diefferenbachiae* sp. nov. and *Dickeya zae* sp. nov. *International Journal of Systematic and Evolutionary Microbiology*, 55, 1415-1427
- Tamura, K., Dudley, J., Nei, M., and Kumar, S. (2007). MEGA4: molecular evolutionary genetics analysis (MEGA) software version 4.0. *Molecular Biology and Evolution*, 24, 1596-1599
- Toth, I. K., Avrova, A. O. and Hyman, L. J. (2001). Rapid identification and differentiation of the soft rot Erwinias by 16S-23S intergenic transcribed spacer and restriction

fragment length polymorphism analysis. *Applied and Environmental Microbiology*, 67, 4070-4076

Toth, I. K., Bell, K. S., Holeva M. C. and Birch, P. R. J. (2003). Soft rot Erwinia: from genes to genomes. *Molecular Plant Pathology*, 4, 17-30

Toth, I. K., van der Wolf, J. M., Saddler, G., Lojkowska, E., Hèlias, V., Pirhonen, M., Tsrör, L. and Elphinstone, J. G. (2011). *Dickeya* species: an emerging problem for potato production in Europe. *Plant Pathology*, 60, 385-399

Tsrör, L., Erlich, O., Hazanovsky, M., Ben Daniel, B., Zig, U. and Lebiush, S. (2012). Detection of *Dickeya* spp. latent infection in potato seed tubers using PCR or ELISA and correlation with disease incidence in commercial field crops under hot-climate conditions. *Plant Pathology* 61, 161-168

Van der Merwe, J. J., Coutinho, T. A., Korsten, L. and van der Waals, J. E. (2010). *Pectobacterium carotovorum* subsp. *brasiliensis* causing blackleg on potatoes in South Africa. *European Journal Plant Pathology*, 126, 175-185

Versalovic, J., Koeuth, T., and Lupski, R. (1991). Distribution of repetitive DNA sequences in eubacteria and application to fingerprinting of bacterial genomes. *Nucleic Acids Research*, 19, 6823-6831

Wegener, C. B. (2002). Induction of defense responses against Erwinia soft rot by an endogenous pectate lyase in potatoes. *Physiological and Molecular Plant Pathology*, 60, 91-100

- Wright, P. J. (1998). A soft rot of Calla (*Zantedeschia* spp.) caused by *Erwinia carotovora* subspecies *carotovora*. *New Zealand Journal of Crop Horticultural Science*, 26, 331-334
- Young, J. M. and Park, D. -C. (2007) Relationship of plant pathogenic enterobacteria based on *atpD*, *carA* and *recA* and individual and as concatenated nucleotide and peptide sequences. *Systematic and Applied Microbiology*, 30, 343-354
- Young, J. M., Takikawa, Y., Gardan, L. and Stead, D. E. (1992). Changing concepts in the taxonomy of plant pathogenic bacteria. *Annual Review of Phytopathology*, 30, 67-105




Figure 1 Cluster analysis of Rep-PCR fingerprint pattern generated from purified DNA of 23 *Pectobacterium* and *Dickeya dadantii* strains. The dendrogram was generated using Bionumerics with an UPGMA algorithm applied to the similarity matrix generated by Pearson's correlation coefficient from whole patterns of individual gel tracks.




Figure 2. Phylogenetic tree based on *gyrB* gene sequence showing the phylogenetic relationship among Zimbabwean strains, *D. dadantii* subsp. *dadantii* and different *Pectobacterium* spp. The phylogram was produced by the neighbour-joining programme (Tamura *et al.* 2007). The numbers on the branches indicate bootstrap value support based on neighbour-joining analyses of 1000 bootstrap replication. Accession numbers of references strains in GenBank are in parenthesis. T, Type strain




Figure 3 Relative pectinolytic activity of atypical strains of pectinolytic enterobacteria isolated from potatoes grown in Zimbabwe and the type strains of *Pectobacterium atrosepticum*, *Pectobacterium carotovorum* subsp. *carotovorum* and *Dickeya dadantii* subsp. *dadantii*.





Figure 4 Dendrogram derived from the UPGMA linkage of correlation coefficient between AFLP profiles from strains of *P. carotovorum* subsp. *brasilensis*, *P. carotovorum* subsp. *carotovorum* and *D. dadantii* subsp. *dadantii*. The levels of linkage representing the Pearson product moment correlations are expressed as percentages for convenience. The banding profiles against each branch each branch represent normalized and background-subtracted digitized gel strips processed in Bionumerics.



Supplementary Figure 1 Phylogenetic tree based on *recA* gene sequence showing the phylogenetic relationship among Zimbabwean strains, *D. dadantii* subsp. *dadantii* and different *Pectobacterium* spp. The phylogram was produced by the neighbour-joining programme (Tamura *et al.* 2007). The numbers on the branches indicate bootstrap value support based on neighbour-joining analyses of 1000 bootstrap replication. Accession numbers of reference strains in GenBank are in parenthesis. T, type strain

Table 1 *Dickeya* spp., *Pectobacterium atrosepticum*, *Pectobacterium carotovorum* subsp. *brasiliensis* and *Pectobacterium carotovorum* subsp. *carotovorum* used in this study

Strain	Host	Isolated from	Origin
<i>Pectobacterium atrosepticum</i> LMG 2386 ^T	<i>Solanum tuberosum</i>		Belgium Coordinated Collections of Microorganisms (BCCM TM)
EN53	<i>Solanum tuberosum</i>	Seed tuber	Nyanga (Zimbabwe)
EN54	<i>Solanum tuberosum</i>	Seed tuber	Nyanga (Zimbabwe)
EN55	<i>Solanum tuberosum</i>	Seed tuber	Nyanga (Zimbabwe)
EN60	<i>Solanum tuberosum</i>	Seed tuber	Nyanga (Zimbabwe)
<i>Pectobacterium carotovorum</i> subsp. <i>brasiliensis</i> strain 8 ATCC BAA-417	<i>Solanum tuberosum</i>		American Type Culture Collection (ATCC) – originated from Brazil
<i>Pectobacterium carotovorum</i> subsp. <i>brasiliensis</i> strain 371 ATCC BAA-419	<i>Solanum tuberosum</i>		American Type Culture Collection (ATCC) – originated from Brazil
JJ1	<i>Solanum tuberosum</i>	Harvested tuber	Mpumalanga (South Africa)
JJ13	<i>Solanum tuberosum</i>	Harvested tuber	Eastern Freestate (South Africa)
JJ18	<i>Solanum tuberosum</i>	Harvested tuber	KwaZulu-Natal (South Africa)
JJ19	<i>Solanum tuberosum</i>	Harvested tuber	KwaZulu-Natal (South Africa)
JJ24	<i>Solanum tuberosum</i>	Harvested tuber	KwaZulu-Natal (South Africa)
JJ27	<i>Solanum tuberosum</i>	Harvested tuber	KwaZulu-Natal (South Africa)
JJ30	<i>Solanum tuberosum</i>	Harvested tuber	KwaZulu-Natal (South Africa)
JJ38	<i>Solanum tuberosum</i>	Harvested tuber	Northern Cape (South Africa)
JJ46	<i>Solanum tuberosum</i>	Harvested tuber	Limpopo (South Africa)
JJ47	<i>Solanum tuberosum</i>	Harvested tuber	Limpopo (South Africa)
JJ49	<i>Solanum tuberosum</i>	Harvested tuber	Limpopo (South Africa)
JJ52	<i>Solanum tuberosum</i>	Seed tuber	Sandveld (South Africa)

Table 1 (continued)

Strain	Host	Isolated from	Origin
JJ54	<i>Solanum tuberosum</i>	Seed tuber	Sandveld (South Africa)
JJ55	<i>Solanum tuberosum</i>	Seed tuber	Sandveld (South Africa)
JJ57	<i>Solanum tuberosum</i>	Harvested tuber	Limpopo (South Africa)
JJ59	<i>Solanum tuberosum</i>	Seed tuber	Sandveld (South Africa)
JJ60	<i>Solanum tuberosum</i>	Seed tuber	Sandveld (South Africa)
JJ64	<i>Solanum tuberosum</i>	Seed tuber	Western Freestate (South Africa)
JJ68	<i>Solanum tuberosum</i>	Stems	Mpumalanga (South Africa)
JJ70	<i>Solanum tuberosum</i>	Stems	Mpumalanga (South Africa)
JJ71	<i>Solanum tuberosum</i>	Stems	Limpopo (South Africa)
JJ72	<i>Solanum tuberosum</i>	Harvested tuber	Limpopo (South Africa)
JJ74	<i>Solanum tuberosum</i>	Seed tuber	Sandveld (South Africa)
JJ75	<i>Solanum tuberosum</i>	Seed tuber	Sandveld (South Africa)
JJ76	<i>Solanum tuberosum</i>	Seed tuber	Sandveld (South Africa)
JJ77	<i>Solanum tuberosum</i>	Seed tuber	Western Freestate (South Africa)
JJ81	<i>Solanum tuberosum</i>	CBS	Mpumalanga (South Africa)
JJ83	<i>Solanum tuberosum</i>	Seed tuber	Eastern Cape (South Africa)
JJ85	<i>Solanum tuberosum</i>	Seed tuber	Eastern Cape (South Africa)
JJ86	<i>Solanum tuberosum</i>	Techni-tubers	CH1A (China)
JJ88	<i>Solanum tuberosum</i>	Techni-tubers	CH3A (China)
JJ90	<i>Solanum tuberosum</i>	Techni-tubers	CH3Bo (China)
JJ91	<i>Solanum tuberosum</i>	Techni-tubers	CH5A (China)
JJ92	<i>Solanum tuberosum</i>	Techni-tubers	CH5B (China)

Table 1 (continued)

Strain	Host	Isolated from	Origin
JJ95	<i>Solanum tuberosum</i>	Techni-tubers	IN3C (china)
JJ96	<i>Solanum tuberosum</i>	Techni-tubers	IN3D (China)
JJ97	<i>Solanum tuberosum</i>	Techni-tubers	IN3E (China)
JJ99	<i>Solanum tuberosum</i>	Techni-tubers	CH1C (China)
JJ107	<i>Solanum tuberosum</i>	Harvested tuber	North West (South Africa)
JJ114	<i>Solanum tuberosum</i>	Seed tuber	Sandveld (South Africa)
JJ128	<i>Solanum tuberosum</i>	Seed tuber	Eastern Cape (South Africa)
JJ130	<i>Solanum tuberosum</i>	Harvested tuber	Eastern Cape (South Africa)
JJ135	<i>Solanum tuberosum</i>	Stems	Eastern Cape (South Africa)
JJ136	<i>Solanum tuberosum</i>	Stems	Eastern Cape (South Africa)
JJ145	<i>Solanum tuberosum</i>	Stems	Limpopo South Africa
JJ147	<i>Solanum tuberosum</i>	Stems	Mpumalanga (South Africa)
JJ149	<i>Solanum tuberosum</i>	Stems	Mpumalanga (South Africa)
JJ150	<i>Solanum tuberosum</i>	Stems	Mpumalanga (South Africa)
JJ153	<i>Solanum tuberosum</i>	Stems	Mpumalanga (South Africa)
EN1	<i>Solanum tuberosum</i>	Harvested tuber	Gwebi (Zimbabwe)
EN2	<i>Solanum tuberosum</i>	Harvested tuber	Gwebi (Zimbabwe)
EN4	<i>Solanum tuberosum</i>	Harvested tuber	Darwendale (Zimbabwe)
EN5	<i>Solanum tuberosum</i>	Harvested tuber	Harare (Zimbabwe)
EN6	<i>Solanum tuberosum</i>	Seed tuber	Harare (Zimbabwe)
EN16	<i>Solanum tuberosum</i>	Seed tuber	Mazowe (Zimbabwe)
EN18	<i>Solanum tuberosum</i>	Seed tuber	Mazowe (Zimbabwe)

Table 1 (continued)

Strain	Host	Isolated from	Origin
EN19	<i>Solanum tuberosum</i>	Seed tuber	Mazowe (Zimbabwe)
EN23	<i>Solanum tuberosum</i>	Seed tuber	Nyanga (Zimbabwe)
EN24	<i>Solanum tuberosum</i>	Seed tuber	Nyanga (Zimbabwe)
EN25	<i>Solanum tuberosum</i>	Seed tuber	Harare (Zimbabwe)
EN26	<i>Solanum tuberosum</i>	Harvested tuber	Harare (Zimbabwe)
EN36	<i>Solanum tuberosum</i>	Harvested tuber	Mazowe (Zimbabwe)
EN37	<i>Solanum tuberosum</i>	Harvested tuber	Mazowe (Zimbabwe)
EN51	<i>Solanum tuberosum</i>	Seed tuber	Shurugwi (Zimbabwe)
EN52	<i>Solanum tuberosum</i>	Seed tuber	Nyanga (Zimbabwe)
EN66	<i>Solanum tuberosum</i>	Seed tuber	Marondera (Zimbabwe)
EN67	<i>Solanum tuberosum</i>	Seed tuber	Nyanga (Zimbabwe)
EN68	<i>Solanum tuberosum</i>	Seed tuber	Nyanga (Zimbabwe)
EN78	<i>Solanum tuberosum</i>	Stem	Gweru (Zimbabwe)
EN79	<i>Solanum tuberosum</i>	Stem	Gweru (Zimbabwe)
<i>Pectobacterium carotovorum</i> subsp. <i>carotovorum</i> LMG 2404 ^T	<i>Solanum tuberosum</i>		Belgium Coordinated Collections of Microorganisms (BCCM TM)
EN7	<i>Solanum tuberosum</i>	Seed tuber	Harare (Zimbabwe)
EN20	<i>Solanum tuberosum</i>	Seed tuber	Mazowe (Zimbabwe)
EN21	<i>Solanum tuberosum</i>	Harvested tuber	Nyanga (Zimbabwe)
EN28	<i>Solanum tuberosum</i>	Harvested tuber	Harare (Zimbabwe)
EN30	<i>Solanum tuberosum</i>	Stem	Nyanga (Zimbabwe)
EN31	<i>Solanum tuberosum</i>	Stem	Darwendale (Zimbabwe)

Table 1 (continued)

Strain	Host	Isolated from	Origin
EN32	<i>Solanum tuberosum</i>	Stem	Darwendale (Zimbabwe)
EN35	<i>Solanum tuberosum</i>	Stem	Gweru (Zimbabwe)
EN39	<i>Solanum tuberosum</i>	Stem	Mazowe (Zimbabwe)
EN58	<i>Solanum tuberosum</i>	Stem	Shamva (Zimbabwe)
EN61	<i>Solanum tuberosum</i>	Stem	Gweru (Zimbabwe)
EN63	<i>Solanum tuberosum</i>	Harvested tuber	Harare (Zimbabwe)
EN69	<i>Solanum tuberosum</i>	Harvested tuber	Nyanga (Zimbabwe)
EN70	<i>Solanum tuberosum</i>	Harvested tuber	Darwendale (Zimbabwe)
EN71	<i>Solanum tuberosum</i>	Stem	Harare (Zimbabwe)
EN75	<i>Solanum tuberosum</i>	Stem	Mazowe (Zimbabwe)
<i>Dickeya dadantii</i> subsp. <i>dadantii</i> (Erwinia chrysanthemi 3937)	<i>Saintpaulia ionantha</i>		Scottish Crop Research Institute (SCRI)
EN3	<i>Solanum tuberosum</i>	Harvested tuber	Gwebi (Zimbabwe)
EN10	<i>Solanum tuberosum</i>	Harvested tuber	Marondera (Zimbabwe)
EN11	<i>Solanum tuberosum</i>	Seed tuber	Marondera (Zimbabwe)
EN12	<i>Solanum tuberosum</i>	Seed tuber	Marondera (Zimbabwe)
EN13	<i>Solanum tuberosum</i>	Seed tuber	Mazowe (Zimbabwe)
EN14	<i>Solanum tuberosum</i>	Stem	Mazowe (Zimbabwe)
EN17	<i>Solanum tuberosum</i>	Stem	Mazowe (Zimbabwe)
EN29	<i>Solanum tuberosum</i>	Stem	Nyanga (Zimbabwe)
EN38	<i>Solanum tuberosum</i>	Stem	Mazowe (Zimbabwe)
EN40	<i>Solanum tuberosum</i>	Stem	Gweru (Zimbabwe)
EN44	<i>Solanum tuberosum</i>	Stem	Harare (Zimbabwe)

Table 1 (continued)

Strain	Host	Isolated from	Origin
EN45	<i>Solanum tuberosum</i>	Stem	Harare (Zimbabwe)
EN47	<i>Solanum tuberosum</i>	Stem	Darwendale (Zimbabwe)
EN48	<i>Solanum tuberosum</i>	Harvested tuber	Gwebi (Zimbabwe)
EN50	<i>Solanum tuberosum</i>	Harvested tuber	Gwebi (Zimbabwe)
EN59	<i>Solanum tuberosum</i>	Harvested tuber	Nyanga (Zimbabwe)
EN72	<i>Solanum tuberosum</i>	Stem	Harare(Zimbabwe)
EN74	<i>Solanum tuberosum</i>	Stem	Gweru (Zimbabwe)
EN76	<i>Solanum tuberosum</i>	Stem	Harare (Zimbabwe)
EN77	<i>Solanum tuberosum</i>	Seed tuber	Harare (Zimbabwe)

Table 2 List of primers used in study

Primer Code	Sequence	Source
<i>gyrB</i> 01-F	TAA RTT YGA YGA YAA CTC YTA YAA AGT	Brady et al., 2008
<i>gyrB</i> 02-R	CMC CYT CCA CCA RGT AMA GTT	Brady et al., 2008
<i>gyrB</i> 07-F	GTV CGT TTC TGG CCV AG	Brady et al., 2008
<i>gyrB</i> 08-R	CTT TAC GRC GKG TCA TWT CAC	Brady et al., 2008
<i>recA</i> F	GAR KCB TCN GGT AAA ACV AC	Young & Park 2007
<i>recA</i> R	TTC GCY TTR CCC TGR CCR ATC	Young & Park 2007
<i>recA</i> R2	RTT GAT RCC TTC GCC GTA SA	Young & Park 2007
REP IR	III ICG ICG ICA TCI GGC	Versalovic et al., 1991
REP 2IR	ICG ICT TAT CIG GCC TAC	Versalovic et al., 1991
E00	GAC TGC GTA CCA ATT C	Pitman et al., 2010
M00	GAT GAG TCC TGA GTA A	Pitman et al., 2010
MseI-C	GAT GAG TCC TGA GTA AC	Pitman et al., 2010

Table 3 Biochemical and physiological properties of selected *Dickeya dadantii* subsp. *dadantii*, *Pectobacterium atrosepticum*, *P. carotovorum* subsp. *carotovorum* and *P. carotovorum* subsp. *brasiliensis* strains used in this study

Strain ^a	Phosphate activity	Indole from tryptophan	Gelatin hydrolysis	Reduced substances from sucrose ^c	Acid from α -methylglucoside	Acid from threulose	Malonate utilization	Erythromycin sensitivity	Growth 37°C	Growth 5 g ⁻¹ NaCl
<i>P. atrosepticum</i> LMG 2386 ^T (BCCM ^T _M)	-	-	+	+	+	+	-	-	-	+
EN54 ^b	+	-	+	+	+	+	-	-	-	+
EN60 ^b	-	-	+	+	+	+	-	-	-	+
<i>P. c.</i> subsp <i>brasiliensis</i> strain8 ATCC BAA-417	-	-	+	-	-	+	-	-	+	+
<i>P. c.</i> subsp <i>brasiliensis</i> strain371 ATCC BAA-419	-	-	+	-	-	+	-	-	+	+
JJ38	+	-	+	-	-	-	+	+	+	-
JJ54	+	-	+	-	-	-	+	+	+	-
JJ68	-	-	+	-	-	+	-	-	+	+
JJ74	+	-	+	-	-	-	+	+	+	-
JJ145	+	-	+	-	-	-	+	+	+	-
JJ147	+	-	+	-	-	-	+	+	+	-
EN4 ^b	-	-	+	-	-	+	-	-	+	+
EN6 ^b	-	-	+	-	-	+	-	-	+	+
EN16 ^b	-	-	+	-	-	+	-	-	+	+

Table 3 Biochemical and physiological properties of selected *Dickeya dadantii* subsp. *dadantii*, *Pectobacterium atrosepticum*, *P. carotovorum* subsp. *carotovorum* and *P. carotovorum* subsp. *brasiliensis* strains used in this study

Strain ^a	Phosphate activity	Indole from tryptophan	Gelatin hydrolysis	Reduced substances from sucrose ^c	Acid from α -methylglucoside	Acid from threose	Malonate utilization	Erythromycin sensitivity	Growth 37°C	Growth 5 g ⁻¹ NaCl
EN66 ^b	-	-	+	-	-	+	-	-	+	+
<i>P. carotovorum</i> subsp. <i>carotovorum</i> LMG 2404 ^T (BCM TM)	-	-	+	-	-	+	-	-	+	+
EN31 ^b	-	-	+	-	-	+	-	-	+	+
EN39 ^b	-	-	+	-	-	+	-	-	+	-
EN63 ^b	-	-	+	-	-	+	-	-	+	+
<i>D. dadantii</i> subsp. <i>dadantii</i> (<i>E. chrysanthemi</i> 3937) (SCRI)	+	+	+	-	-	-	+	+	+	-
EN12 ^b	+	-	+	-	-	-	+	+	+	-
EN38 ^b	+		+	-	-	-	+	+	+	-
EN44 ^b	+	+	+	-	-	-	+	+	+	W+
EN47 ^b	+		+	-	-	-	+	+	+	W+
EN59 ^b	+		+	-	-	-	+	+	+	-

ATCC American Type Culture Collection, BCCM Belgian Coordinated Collections of Microorganisms, SCRI Scottish Crop Research Institute, Dundee, Scotland, UK

EN, Isolated from potatoes grown in Zimbabwe

JJ, Isolated from potatoes grown in South Africa

W, weak reaction

^aGram, Oxidase, glucose metabolism and pectate degradation in Sutton's medium test results of all the strains agreed with those expected for *Dickeya dadantii* subsp. *dadantii*, *Pectobacterium atrosepticum*, *Pectobacterium carotovorum* subsp. *carotovorum* and *Pectobacterium carotovorum* subsp. *brasiliensis*, according to Cother and Sivasithaparam (1983)

^bStrains considered as representative of bacteria obtained from different potato growing regions in Zimbabwe and selected for further analysis

^cOrange zones developing in the culture medium after 5 min boiling when treated with Benedict's reagent

Table 4 Soft rot thickness and diameter on potato slices after inoculation with bacterial isolates from potato tubers

Symbol of isolate/ strain	Penetration of potato slice tissue by soft rot (% of slice thickness affected) ^a		Diameter of soft rot (mm) After 48 h of incubation ^b
	24 h	48 h	
Control – water	0	0	0a
LMG 2404 ^T (Pcc)	30	100	24b
EN31	100	100	30cde
EN63	100	100	36e
LMG 2386 ^T (Pa)	100	100	36e
EN54	100	100	36e
EN60	50	100	27bcd
ATCC BAA – 417 (Pcb)	100	100	36e
ATCC BAA-419 (Pcb)	100	100	31cde
EN4	50	100	30de
EN6	50	100	25b
EN16	100	100	31cde
EN66	100	100	36e
Ech 3937 (Dd)	50	100	28bcd
EN12	100	100	36e
EN38	50	100	28bcd
EN44	50	100	27bcd
EN47	100	100	27bcd
EN58	100	100	32cde
EN59	50	100	25b

^a Preliminary test (average value of 20 slices per isolate)

^b Quantitative test; values with the same letter in a column are not significantly different at P< 0.05