

Genotype composition of populations of *Citrus tristeza virus* grapefruit cross-protecting GFMS12 in different host plants and aphid-transmitted sub-isolates

Katherine Anne Scott^a, Quinsile Hlela^a, Olivier Zablocki^a, David Read^a, Stephanus van Vuuren^c and Gerhard Pietersen^{a, b}

^aDepartment of Microbiology and Plant Pathology and FABI, New Agricultural building, University of Pretoria/Citrus Research International, Lunnon Road, Hatfield, South Africa, 0002.

Email address: ScottKA@arc.agric.za

Tel: +27124203279

Fax: +27124203266

^bAgricultural Research Council - Plant Protection Research Institute, Private Bag X134, Queenswood, 0121, South Africa

Email address: gerhard.pietersen@up.ac.za

^cCitrus Research International, P.O.Box 28, Nelspruit, 1200, South Africa

Note: Nucleotide sequence data reported are available in the DDBJ/EMBL/GenBank databases under the accession numbers JQ819902 to JG820013.

Abstract

Citrus tristeza virus (CTV) causes severe losses in grapefruit production in South Africa and requires mild strain cross-protection to maintain production. Unfortunately cross-protection breakdown of the pre-immunizing CTV grapefruit mild source GFMS12 is prevalent in grapefruit in South Africa. The CTV genotype composition of the GFMS12 population inoculated onto different hosts was determined by sequencing part of ORF1a and the p23 gene of multiple clones from each plant. Analysis of the GFMS12 population in Mexican lime, Marsh and Star Ruby grapefruit varieties, revealed that at least four genotypes occur in the GFMS12 population, and that genotype compositions differed amongst the populations in different host plants. Single aphid transmitted sub-isolates derived from the GFMS12 mother population on Mexican lime appeared to contain three populations of a mixture of VT-like and recombinant B165/VT-like genotypes; a mixture of recombinant RB/VT- and B165/VT-like genotypes; and a single recombinant B165/VT-like genotype. This study underlines the importance of determining the genotype composition of a potential CTV pre-immunizing source on a range of inoculated host species before utilization.

Introduction

Citrus tristeza virus (CTV) is an aphid-borne closterovirus, endemic in South Africa. Stem pitting and decline symptoms of CTV have been major factors limiting productivity of grapefruit locally. Without CTV cross-protection, it has been suggested that grapefruit production would be uneconomical in South Africa [19]. Marsh grapefruit trees planted in the 1920's and infected with CTV but still producing excellent fruit in the 1970's served as sources of mild sources to potentially protect grapefruit trees. These sources, Grapefruit mild strain GFMS 12, GFMS 35 and others were evaluated for mildness, initially in the greenhouse [21], and then in field trials [19]. In 1973, the Southern African Citrus Improvement Scheme (SACIS) was initiated and until 2007 all grapefruit selections were pre-immunized with the GFMS12 source. Star Ruby grapefruit, pre-immunized with the GFMS12 source however later showed varying degrees of stem-pitting and variable fruit size in field trials and on trees for budwood production [20, 22]. The heterogeneity of the mother GFMS12 source was demonstrated and, based on biological indexing, a severe CTV strain with severe stem-pitting found in it [23]. CTV strains within GFMS12 were separated by single *Toxoptera citricida* transmissions and sub-isolates GFMS12-1 to GFMS12-9 were established [23]. Two sub-isolates were shown to be less virulent while one sub-isolate was more virulent than the original GFMS12 source [23]. Additionally, the presence of seedling yellows (CTV-SY) symptoms, not found in GFMS12 plants maintained in greenhouses, but in some plants in

the field, suggested that at least some GFMS12 pre-immunized plants had become super-infected with additional CTV strains probably introduced by aphids [18]. Altered single strand conformational polymorphic (SSCP) patterns, not corresponding to the SSCP profile of the original GFMS12 source supported this [18]. Recent studies [5] have shown that hyper infection exclusion in CTV is genotype specific; suggesting that cross protecting sources should contain mild variants of all CTV genotypes circulating within a given area in order to remain effective.

The purpose of the study was to characterise the genotype composition of the greenhouse maintained GFMS12 mother source population on various citrus hosts as well as that of single aphid transmitted sub-isolates derived from it [23], in order to better investigate the role of various CTV genotypes in cross protection breakdown of this mild source. The genotype composition in each population was determined by sequencing multiple clones (between 22 and 64) of two genomic regions.

Materials & Methods

Virus isolates and sub-isolates

The original field-collected GFMS12 isolate is maintained on Marsh grapefruit on rough lemon rootstock in a glasshouse. This source was used to pre-immunize grapefruit plants within the Citrus Improvement Scheme (S.P van Vuuren, personal communication). The Star Ruby and Marsh grapefruit varieties (*C. paradisi* Macf.) on rough lemon rootstocks and Mexican lime (*Citrus aurantifolia* (L.) Swingle) plants were inoculated from the original GFMS12 isolate (S.P van Vuuren, personal communication) and maintained in an insect-free greenhouse in Nelspruit for eight years prior to this study. GFMS12 sub-isolates (12-1; 12-4; 12-5; 12-6; 12-7; 12-8; and 12-9) on Mexican lime were obtained by single aphid transmissions (*Toxoptera citricida* (Kirkaldy)) from the GFMS12 inoculated Mexican lime plant obtained in a previous study [23]. Sub-isolates were sampled and their A-region and p23 gene DNA products sequenced at two different time points; initially when plants were introduced into the glasshouse in Pretoria and then repeated between 2 and 3 years later.

RNA extraction, cDNA synthesis, and PCR amplification

Bark from different parts of the CTV-infected plant was pulverized with liquid nitrogen and total RNA extracted with the SV Total RNA Isolation System (Promega) according to the manufacturer's protocol. A two-step RT-PCR protocol was utilized to detect and amplify CTV RNA. Two pairs of primers targeting the A

region (ORF1a) [14] and the p23 gene [15] (Table 1) were used. Reverse transcription was performed according to Herron [9] using either the PM51 or A-R reverse primers (Table 1). PCR reactions of 50 µl contained 10 µl of the primed cDNA and a mixture containing: 100 pmoles of each primer (PM50/PM51 or A-F/A-R); 2.5 mM MgCl₂; 1x Bioline Buffer; 2.5U BioTaq (Bioline); 0.14 mM each of dATP, dCTP, dGTP and dTTP; and 0.2 µg/µl BSA. The thermal cycle conditions were: 92 °C for 2 min, 40 cycles of 92 °C for 30 sec, 55 °C for 45 sec, and 72 °C for 1 min, with extension for 10 min at 72 °C. The resulting PCR products were separated by 1 % agarose gel electrophoresis. The PCR products were purified using a Wizard® SV gel and PCR clean-up kit (Promega) according to the manufacturer's instructions.

Cloning and sequence analysis of GFMS12 sources

Characterization of the genotype composition of GFMS12 populations on Mexican lime, Star Ruby and Marsh grapefruit plants was performed by cloning the PCR product of the A region and the p23 gene from each source and sequencing multiple clones. Cloning was with the pGEM-T Easy vector cloning system (Promega) according to the manufacturer's instructions, followed by transformation of *Escherichia coli* JM109 competent cells (Promega). Recombinant colonies were selected by blue/white selection and grown overnight in Luria broth with 100 µg/ml ampicillin (USB). Plasmids were isolated using the alkaline lysis method [16], and screened by PCR using pGEM-T vector specific primers T7 and Sp6 (Integrated DNA Technologies; IDT) (Table 1) and PCR products were purified [25]. Sequences of cloned plasmids were determined using primers SP6 or T7.

Sequences obtained were trimmed using the BioEdit sequence alignment editor (v7.0, Tom Hall, Isis Pharmaceuticals, Inc 1997-2004), and alignments carried out by the ClustalW alignment software (EBI) incorporated into the BioEdit software [17]. Sequences were aligned with the cognate regions of 18 full genome reference sequences retrieved from GenBank. Cognate regions from the full genome sequences of CTV isolates T30 (AF260651) and T36 (U16304) from Florida; T318A (DQ151548) and T385 (Y18420) from Spain; VT from Israel (U56902); SY568 (AF001623) from California; Mexican CTV isolate (DQ272579); NUagA (AB046398) from Japan; B165 (EU076703) and Kpg3 (HM573451) from India; Qaha (AY340974) from Egypt; HA16-5 (GQ454870) and HA18-9 (GQ454869) from Hawaii; and New Zealand resistance breaking isolates NZRB-M12 NZRB-G90, NZRB-TH28, NZRB-TH30 and NZRB-M17 (FJ525431-35) were retrieved from GenBank and included in the analysis to provide phylogenetic references. The Mega 3.1

package was used to perform the phylogenetic and distance analyses [11]. Unrooted dendrograms were constructed after bootstrapping to 1000 replicates by the Neighbour-joining (NJ) method using the Jukes-Cantor base substitution model. Dendrograms with bootstrap values of more than 70% were generally regarded as providing evidence for a phylogenetic grouping. The graphical output for the 50% majority rule consensus trees was obtained by using the MEGA 3 tree explorer version 3.1 [11]. Recombination was analyzed using the GENCOV function of RDP-V2 Beta 08.

Direct sequencing without DNA cloning was done on GFMS12 single aphid transmission sub-isolates: 12-1; 12-4; 12-5; 12-6; 12-7; 12-8 and 12-9 [23] to obtain the consensus sequence of the population in order to select a representative of each potentially isolated genotype. Sequencing was performed: a) initially when plants were obtained from Citrus Research International, Nelspruit, and; b) repeated after a 2-3 year period of maintenance under glasshouse conditions in Pretoria. Direct nucleotide sequence determination was done in both directions by means of an ABI Prism DNA sequencer 377 (Perkin-Elmer) using A-F, A-R, PM50 and PM51 primers. The DNA was purified by the EDTA/NaOAc/EtOH method [16]. The genotype composition of the population for representative sub-isolates was determined by sequencing multiple clones of the A-region and p23 gene amplicons as described above for the GFMS12 mother sources.

Biological indexing

The pathogenicity of GFMS12 sub-isolates was determined by inoculating the virus source on greenhouse maintained container-grown plants of five recognized [7] citrus indicators: Mexican lime (ML); sour orange (SO) (*C. aurantium*); Duncan grapefruit (DG) (*C. paradisi*); Madam Vinous (MV) sweet orange (*C. sinensis*); and sweet orange grafted on sour orange rootstock (SW/SO). Three plants of each indicator/virus source were graft inoculated with bark-pieces from CTV-infected donors (GFMS12 source on Mexican lime; sub-isolates 12-7; 12-8; and 12-9) with un-inoculated plants kept as controls. Symptoms were monitored periodically over a six month period for SO, SW/SO, ML plants while DG and MV plants were monitored for a further four months [7]. Intensity of stem-pitting was scored as follows: 0-20 pits/cm² was considered mild; 21-50 pits/cm², moderate; and >51 pits/cm² was considered severe [23].

Results

The CTV genotypes present within the maintained GFMS12 sources on grapefruit varieties Marsh and Star Ruby, and Mexican lime were identified from sequences of multiple clones of amplicons of the A-region (ORF1a) and the p23 gene and used to prepare dendrograms vs. cognate regions of known CTV genotypes (Figs. 1 and 2). The number of clones of each identified genotype in the GFMS12 sources gives an indication of the genotype composition present (Table 2). From this it is evident that GFMS12 consists of multiple genotypes and that the population on different plants appear to contain differing genotype compositions. Incongruences in genotype composition using the A-region and the p23 gene within the same population are also observed. The genotypes present in the GFMS12 population on the two grapefruit cultivars show similar incongruences. Both populations are composed of CTV genotypes with only VT-like p23 genes, and both contain CTV genotypes with B165-like and T30-like A-regions (Table 2). In the GFMS12 population on Star Ruby an additional genotype with a VT-like A-region is observed (Table 2). No clones containing RB-, HA16-5-, Kpg-3- or T36-like sequences in the A-region or p23 gene were obtained in the grapefruit cultivars. The relative proportion of cloned plasmids with specific genotype sequences in the two grapefruit populations differed however. Most cloned A-region amplicons have sequences that are B165-like in Marsh, while being T30-like in Star Ruby. The Mexican lime population differs considerably from the two grapefruit populations with most sequenced clones of both the A-region and the p23 gene being RB-like, a genotype sequence not observed in clones of either sequenced region from either of the grapefruit populations. The GFMS12 population on Mexican lime also contained B165- and VT-like genotypes, as was obtained in the two grapefruit populations.

Identical nucleotide sequences within cloned amplicons from different GFMS12 populations were often obtained. These were collapsed into single branches on the dendrogram (branches with multiple coloured circles in Figs. 1 and 2). Commonly however, some variation in sequence within any given genotype, even within a single GFMS12 population, was observed. The following example illustrates these two occurrences; eight clones of A-region amplicons from GFMS12 on Star Ruby and one from GFMS12 on Marsh yielded identical sequences clustering within the T30-like group (Fig. 1), however clones of this sequence, along with five other clones all from the GFMS12 population in Star Ruby, within the T30 clade had 1% nucleotide diversity (Table 3). The average intra-genotype nucleotide diversity for the A-region and p23 gene of cloned amplicons from GFMS12 Mexican lime, Marsh and Star Ruby grapefruit populations are shown in Table 3. Higher nucleotide diversity exists amongst strains in the variable A-region than in the more conserved p23

gene. Recombination analysis was performed using the GENCOV function of RDP-V2 Beta 08, with no genotype recombination events being found within any of the A-region sequences of any of the GFMS12 populations. Recombination with an unknown parental sequence was detected in the p23 gene of the RB-like genotype identified within the GFMS12 Mexican lime population (data not presented).

Direct sequencing of the A-region and the p23 gene of single aphid-transmission sub-isolates of GFMS12 on Mexican lime yielded the population consensus sequence, effectively that of the dominant genotype (Fig. 3). Based on direct sequences of the A region when sources were initially moved from greenhouses in Nelspruit to those in Pretoria, a population consensus of a VT-like dominant genotype was observed in GFMS12 sub-isolates 12-1, 12-4, 12-6, and 12-9; a B165-like genotype in sub-isolate 12-8; and an RB-like genotype in sub-isolates 12-5 and 12-7 while after a 2 to 3 year period in the Pretoria greenhouse the population consensus sequences shifted to that of a dominant B165-like genotype for all sub-isolates (Fig. 3A). Analysis of the p23 gene of the sub-isolates over the three year time period suggested that they all contain VT-like dominant genotypes, with the consensus sequence obtained from sub-isolates 12-5 and 12-7 grouping more closely to the T318A and NUagA variants of the VT-genotype, and sub-isolates, 12-1, 12-4, 12-6, 12-8 and 12-9 being more related to the type VT isolate (Fig. 3B).

The combined A-region and p23 gene analyses of the GFMS12 sub-isolates suggest that three populations with apparently different CTV genotypes are present; sub-isolates 12-1, 12-4, 12-6 and 12-9 consisted of a B165-like and VT-like sequence type in the A-region but VT-like sequence type in the p23 gene and represents a mixture of a possible recombinant B165/VT-like strain and a VT-like strain; sub-isolates 12-5 and 12-7 had a RB-like and B165-like sequence type in the A-region and a VT-like sequence in the p23 gene and represents a mixture of possible recombinant RB/VT-like and B165/VT-like strains. Sub-isolate 12-8 had B165-like sequence type in the A-region but VT-like sequence type in the p23 gene and represents a single possible recombinant B165/VT-like strain that was maintained over time. Based on the direct sequencing information sub-isolates 12-7, 12-8 and 12-9 were chosen as representatives of the three populations found. Dendrograms derived from sequences of 20-30 clones of the A-region and p23 gene of the representative sub-isolates compared to the cognate region of known CTV genotypes are not presented as clones were all of the same sequences identified with direct sequencing. Based only on the second time point after direct sequencing sub-isolates 12-7 and 12-8 were B165-like, sub-isolate 12-9 was a mixture of VT-like and B165-like in the A-region

while all of them still had VT-like p23 gene sequences. These three sub-isolates were indexed in triplicate in each of five citrus indicator hosts and their symptoms compared to the GFMS12 pre-immunizing source (Table 4). Sub-isolate 12-7 yielded mild symptoms, whereas symptoms of sub-isolates 12-8, 12-9 and the GFMS12 source were moderate under the experimental conditions used. The GFMS12 source had moderate vein clearing and SP in Mexican lime; mild stunting and moderate SP in Duncan grapefruit. Sub-isolate 12-7 was milder than the GFMS12 source for Mexican lime and Duncan grapefruit, and in addition displayed decline in SW/SO. Sub-isolate 12-8 (B165-like) did not display any stem-pitting symptoms in grapefruit. Sub-isolate 12-9 had symptoms on all five indicators with mild SP and stunting occurring. The symptom profile induced by sub-isolates 12-9 and 12-8 differed from that of the GFMS12 source.

Discussion

In a recent study by Folimonova *et al.* [5] it was shown that super-infection exclusion by CTV sources is genotype specific, with a given CTV genotype only providing super-infection exclusion against other sources of the same T36 genotype. These results have direct bearing on the strategy of mild strain cross protection of citrus, where it is now evident that mild forms of all CTV genotypes circulating in a given citrus production area are required as pre-immunizing sources to achieve sustainable mild strain cross protection. It is therefore critical to fully characterise, at least to genotype level, the CTV populations used for pre-immunization, as well as to identify and characterise those able to overcome the cross protecting infections. In this study we therefore determined the genotype composition of the GFMS12 pre-immunizing source which, while initially showing great promise as a cross protection source, fell into disuse due to cross protection breakdown in the field [22]. We also determined the genotype composition of populations obtained from the mother source by single aphid transmission to determine their usefulness as pre-immunizing sources.

We have shown that GFMS12 population contains at least RB-; T30-; VT-; and B165-like genotype sequences observed amongst the Mexican lime, Marsh and Star Ruby GFMS12 populations tested. We suspect that all four genotypes are in the GFMS12 population even though the full complement was not found in any of the individual populations, and that the populations in the different plants represent various combinations and differing ratios of the components present in the original GFMS12 population. We conclude this as the possibility of introduction of additional CTV genotypes by aphid transmission was negligible since populations were maintained in insect-free greenhouses throughout their lifetime. The study therefore also confirms the

mixed “strain” nature of GFMS12 suggested previously by differing dsRNA and SSCP patterns [2, 23]. In all three GFMS12 populations a predominant genotype with additional minor genotypes was observed similar to that previously reported for CTV populations [10].

The GFMS12 population from Mexican lime was composed of: 66% RB-; 30% B165-; and 4% VT-like genotypes when based on the A-region, but composed of 95% RB- and 5% VT-like genotypes when based on the p23 gene. Such incongruent relationships have been found in the past for CTV [14] and may represent recombination events between diverged sequence variants. This may suggest that about 30% of the genotypes present in the GFMS12 population on Mexican lime are probably a B165 (A-region)/RB (p23 gene) recombinant. To determine the actual recombination sites further sequence determination, of the region between the A-region and the p23 gene, but preferably the whole genome, is envisaged, a goal made possible with next generation sequencing technology. It will be important to determine the extent of genotype specific hyper-exclusion [5] that occurs amongst a range of variants and recombinants within a specific genotype. Full genome sequencing will also help to distinguish if any sequence variants are dRNAs or recombinant full-length genomes.

Interestingly, the GFMS12 populations differed between varieties of the same host, with the genotype composition on grapefruit cv. Marsh being 96% B165- and 4% T30-like; and in contrast, on grapefruit cv. Star Ruby, it was 81% T30-, 13% B165- and 6% VT-like based on the A-region. Only VT-like sequences were obtained amongst multiple clones of the p23 gene on both grapefruit cultivars. This suggests that along with some VT-like genotypes within the Star Ruby GFMS12 plant, some are likely to be recombinants between the B165- or T30-like genotypes in the 5' end and VT-like genotypes in the 3' end. A previous study found that a large number of genetic variants were generated by recombination between major genotypes [24]. Potentially cv. Marsh may be selecting for the B165-VT recombinant and cv. Star Ruby selecting for the T30-VT recombinants. Analyses of GFMS12 populations in replicated individuals of these two hosts are however required to confirm this.

Three of the four genotypes observed in the parental GFMS12 populations were detected amongst seven Mexican lime single aphid transmission sub-isolates [23] tested by direct sequencing and hence could represent the most dominant genotypes or the most efficiently transmitted. These included VT-like, possible RB (A-

region)-/VT (p23-gene)-like and B165/VT-like recombinant genotypes. None of the sub-isolates contained the RB-like or recombinant B165/RB sequences, found in the GFMS12 parental population. Direct sequencing of the A-region of GFMS12 sub-isolates showed the detection of different genotypes over time. Initially four sub-isolates had VT-like genotype and two sub-isolates had a RB-like sequence type but two to three years later consisted of the B165-like sequence type. The change in the population of sub-isolates 12-7, 12-8 and 12-9 was confirmed with cloning the A-region at the second the time period. The change in genotypes present in sub-isolates over time suggests that the B165/VT genotype was repressed initially or segregated with graft transmission but with time was able to replicate and dominate the other sequence types (VT-like or RB/B165-like recombinant) present under the experimental conditions in the Pretoria greenhouse. A mixture of RB/VT-like and B165/VT-like recombinant genotypes for sub-isolate 12-7; a B165/VT-like recombinant genotype for sub-isolate 12-8; and a mixture of VT-like and B165/VT recombinant genotypes for sub-isolate 12-9 were selected as representatives of the three genotype populations found within the sub-isolates. To accurately determine the pathogenicity of genotypes present in pre-immunizing sources, representatives of the genotype populations were evaluated. The symptoms observed for sub-isolate 12-7 are possibly due to the putative recombinant RB/VT genotype. RB isolates are generally mild in most indicators, with no SP or SY reactions (S. Harper, personal communication). Sub-isolate 12-8 (B165-like) had no stem-pitting in grapefruit and was very similar to the B165 isolate from India [13]. It cannot be predicted whether symptoms are associated with particular regions of the genome or if they are induced by the predominant variant, minor variants or by a combination of variants of the population. Sub-isolate 12-9 had symptoms on all five indicators with mild SP and stunting possibly attributed to the VT-like and putative recombinant B165/VT genotype. In previous studies, Mexican lime and grapefruit varieties Star Ruby and Marsh infected with GFMS12 displayed moderate, mild, and mild to moderate stem-pitting symptoms respectively [19, 23]. These symptoms are not generally caused by RB-like or T30-like genotypes but could potentially be caused by the B165- and VT-like genotypes present as minor or major components of the population, as shown by Černi *et al* [3]. Since a SY pathogenicity determinant was mapped to the p23 gene [1], it is possible that the symptoms observed for GFMS12 are attributed to the RB- (Mexican lime) and VT-like strains (Marsh and Star Ruby) found in the p23 gene.

In this study, the RB-like strain of CTV is reported for the first time in South Africa previously only being reported from New Zealand [8]. RB strains are able to infect the economically important rootstock species *Poncirus trifoliata* [4] of which hybrid rootstocks are commonly used in South Africa, which is resistant to

most isolates of CTV [6]. We did not confirm the resistance breaking nature of GFMS12, or sub-isolate 12-7 by inoculating it onto *P. trifoliata* but would be essential to test local RB-like isolates in the future.

This study underlines the importance of determining the genotype composition of a potential CTV pre-immunizing source on a range of inoculated host species before utilization. Even with the current study the possibility of additional genotypes being present in sub-detectable levels in the three GFMS12 populations tested cannot be excluded, and these may only present themselves at detectable levels in additional host species and with larger numbers of cloned sequences analysed. Additionally, a clearly unbiased method of determining the genotype composition of greater numbers of CTV populations must be developed. While we have tried to minimize genotype bias by PCR by using primers to conserved sequences amongst genotypes which do not preferentially amplify specific genotypes and have been used in similar studies [14, 15], the occurrence of bias cannot be discounted completely. The absence of some CTV genotypes in the GFMS12 populations (e.g. HA16-5, T30 and T36) could leave trees pre-immunized with this source vulnerable to these genotypes should the genotype specific hyper-exclusion phenomenon [5] be a general principle amongst CTV genotypes. It is possible that at least some of the instances of cross protection breakdown observed in South Africa grapefruit orchards may be due to the super-infection of these additional genotypes in GFMS12 pre-immunized plants. Studies are in progress to determine whether such additional genotypes are found in GFMS12 pre-immunized grapefruit trees in the field displaying severe CTV symptoms (Read, unpublished results).

Acknowledgements:

Financial support was provided by Citrus Research International (CRI) (South Africa). We wish to acknowledge Glynnis Cook for her critical reading of the manuscript.

References

1. Albiach-Martí MR, Robertson C, Gowda S, Tatineni S, Belliure B, Garnsey SM, Folimonova SY, Moreno P, Dawson WO (2010) The pathogenicity determinant of Citrus tristeza virus causing the seedling yellows syndrome maps at the 3'-terminal region of viral genome. *Mol Plant Pathol* 11: 55-67
2. Albiach MR, da Graca JV, van Vuuren SP, Guerri J, Cambra M, Laigret F, Moreno P (1996) The effects of different hosts and natural disease pressure on molecular profiles of mild isolates of citrus tristeza virus (CTV).

In: da Graca JV, Moreno P, Yokomi RK (eds) Proceedings of the 13th Conf. of the IOCV. IOCV, Riverside, pp 147-153

3. Černi S, Ruscic J, Nolasco G, Gatin Z, Krajacic M, Skoric D (2008) Stem-pitting and seedling yellows symptoms of *Citrus tristeza virus* infection may be determined by minor sequence variants. *Virus Genes* 36: 241-249

4. Dawson TE, Mooney PA (2000) Evidence for trifoliolate resistance breaking isolates of *Citrus tristeza virus* in New Zealand. In: Yokomi RF, Lee RK, Da Graca JV (eds) Proceedings of the 14th Conf. of the IOCV, IOCV, Riverside, pp 69-76

5. Folimonova SY, Robertson CJ, Shilts T, Folimonov AS, Hilf ME, Garnsey SM, Dawson WO (2010) Infection with strains of *Citrus Tristeza virus* does not exclude superinfection by other strains of the virus. *Virology* 84: 1314-1325

6. Garnsey S M, Barret HC, Hutchison DJ (1987) Identification of *Citrus tristeza virus* resistance in citrus relatives and its potential applications. *Phytophylactica* 19: 187-192

7. Garnsey SM, Gumpf DJ, Roistacher CN, Civerolo EL, Lee RF, Yokomi RK, Bar-Joseph M (1987) Towards a standardized evaluation of the biological properties of *Citrus tristeza virus*. *Phytophylactica* 19: 151-157

8. Harper SJ, Dawson TE, Pearson MN (2010) Isolates of *Citrus tristeza virus* that overcome *Poncirus trifoliata* resistance comprise a novel strain. *Arch. of Virol* 155: 471-480

9. Herron CM (2003) *Citrus tristeza virus*: Characterization of Texas isolates, studies on aphid transmission and pathogen-derived control strategies. Dissertation, Texas A&M University

10. Kong P, Rubio L, Polek M, Falk BW (2000) Population structure and genetic diversity within California *Citrus tristeza virus* (CTV) isolates. *Virus Genes* 21: 139-145

11. Kumar S, Tamura K, Nei M (2004) MEGA3: integrated software for molecular evolutionary genetics analysis and sequence alignment. *Brief Bioinformatics* 5: 150-163

12. Roy A, Brlansky RH (2009) Population dynamics of a Florida *Citrus tristeza virus* isolate and aphid-transmitted sub-isolates: Identification of three genotypic groups and recombinants after aphid transmission. *Phytopathology* 11: 1297-1306
13. Roy A, Brlansky RH (2010) Genome Analysis of an Orange Stem Pitting *Citrus Tristeza Virus* Isolate Reveals a Novel Recombinant Genotype. *Virus Res* 151: 118-130
14. Rubio L, Ayllón MA, Kong P, Fernandez A, Polek M, Guerri J, Moreno P, Falk B (2001) Genetic variation of *Citrus tristeza virus* isolates from California and Spain: Evidence for mixed infections and recombination. *J Virol* 75: 8054-8062
15. Sambade A, López C, Rubio L, Flores R, Guerri J, Moreno P (2003) Polymorphism of a specific region in gene p23 of *Citrus tristeza virus* allows discrimination between mild and severe isolates. *Arch Virol* 148: 2325-2340
16. Sambrook J, Russell DW (2001) *Molecular cloning: a laboratory manual* 3rd ed, Cold Spring Harbour Laboratory Press, New York
17. Thompson JD, Higgins DJ, Gibson TJ (1994) CLUSTAL W: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucl Acids Res* 22: 4673-4680
18. Van der Vyver J, van Vuuren SP, Luttig M, da Graca JV (2002) Changes in the *Citrus tristeza virus* status of the pre-immunized grapefruit field trees. In: Duran-Vila N, Milne RG, da Graca JV (eds) *Proceedings of the 15th Conf. IOCV, IOCV, Riverside*, pp
19. Van Vuuren SP, Collins RP, da Graça JV (1993) Evaluation of *Citrus tristeza virus* isolates for cross protection of grapefruit in South Africa. *Plant Dis* 77: 24-28

20. Van Vuuren SP, Manicom BQ (2005) The response of Star Ruby Grapefruit to different *Citrus tristeza virus* isolates. In: Proceedings of the 16th Conf. of the IOCV, IOCV, Riverside pp 112-116
21. Van Vuuren SP, Moll JN (1987) Glasshouse evaluation of citrus *tristeza virus* isolates. *Phytophylactica* 19: 219-221
22. Van Vuuren SP, van der Vyver J (2000) Comparison of South African ore-immunizing citrus *tristeza* isolates with foreign isolates in three grapefruit selections. In: da Graca JV, Lee RF, Yokomi RK (eds) Proceedings of the 14th Conf. of the IOCV, IOCV, Riverside, pp 50-56
23. Van Vuuren SP, van der Vyver JB, Luttig M (2000) Diversity among sub-isolates of cross-protecting *Citrus tristeza virus* isolates in South Africa. In: da Graca JV, Lee RF, Yokomi RK (eds) Proceedings of the 14th Conf. of the IOCV, IOCV, Riverside, pp 103-109
24. Weng Z, Barthelson R, Gowda S, Hilf ME, Dawson WO (2007) Persistent infection and promiscuous recombination of multiple genotypes of an RNA virus within a single host generate extensive diversity. *PLoS ONE* 2(9): e917. Doi: 10.1371/journal.pone.0000917.
25. Werle E, Schneider C, Renner M, Völker M, Fiehn W (1994) Convenient single-step, one tube purification of PCR products for direct sequencing. *Nucl Acids Res* 22: 4354-4355

Figure 1: Neighbour-joining dendrogram depicting phylogenetic relationships between clones of A-region nucleotide sequences of CTV GFMS12 on Marsh (M) (circles); Star Ruby (SR) (squares); Mexican lime (ML) (triangles) and cognate reference CTV sequences. Bootstrap values (1000 replicates) supporting a particular phylogenetic grouping is indicated on the branch points.

Figure 2: Neighbour-joining dendrogram depicting phylogenetic relationships between clones of p23 gene nucleotide sequences of CTV GFMS12 on Marsh (M) (circles); Star Ruby (SR) (squares); Mexican lime (ML) (triangles) and cognate reference CTV sequences. Bootstrap values (1000 replicates) supporting a particular phylogenetic grouping is indicated on the branch points.

Figure 3: Neighbour-joining dendrogram of the (A): A-region (2021-2548 nt) and (B) p23 gene, depicting the phylogenetic relationships of nucleotide sequences of GFMS12 sub-isolates sequenced (a) initially upon introduction in the greenhouse (black dots) and then (b) after 2-3 years of maintenance under greenhouse conditions (black triangles) and references using the Jukes-Cantor parameter. Bootstrap values (1000 replicates) supporting a particular phylogenetic grouping is indicated on the branch points.

Figure 1




Figure 2




Figure 3





Table 1. Primers used for the RT-PCR amplification of GFMS 12 sources and sub-isolates.

Primer	Sequence (5'-3')	Region	Position¹	Product size	Author
A-F forward A-R Reverse	ACGTGTTTCGTGAAACGCGG GTCGATAACTCGACAAACGAGC	ORF1a	2021-39 2527-48	528 bp	Rubio <i>et al.</i> , 2001
PM 50 Forward PM51 Reverse	ACTAACTTTAATTCGAACA AACTTATCCGTCCACTTC	p23	18347-65 19026-44	697 bp	Sambade <i>et al.</i> , 2002
T7 SP6	TAATACGACTCACTATAGGG ATTTAGGTGACACTATA GAA	pGEM-T vector	N/A	N/A	Promega pGEM-T Easy vector system (Promega, USA)

¹Nucleotide positions of primers relative to the T36 isolate (GenBank accession U16304)

Table 2. The population diversity in the A-region (ORF1a) and p23 gene of GFMS 12 sources on Mexican lime, Marsh and Star Ruby grapefruits. The percentage of cloned sequences from each source that grouped with reference genotypes (B165, HA16-5, Kpg3, RB, T30, T36 and VT) are shown.

Source	Sequence	No. of clones	Percentage CTV genotype							
			B165	HA16-5	Kpg3	RB	T30	T36	VT	
GFMS12 Mexican Lime	A-region	27	30*	0	n/a	66.0	0	0	4.0	
	p23 gene	22	n/a	0	0	95.0	0	0	5.0	
GFMS12 Marsh	A-region	24	96.0	0	n/a	0	4.0	0	0	
	p23 gene	40	n/a	0	0	0	0	0	100.0	
GFMS12 Star Ruby	A-region	16	13.0	0	n/a	0	81.0	0	6.0	
	p23 gene	42	n/a	0	0	0	0	0	100.0	

* = Percentage of clones sequenced

n/a = not applicable, no differentiation of such a genotype based on this region

Table 3. Intra-group nucleotide diversity values from sequences derived from the A-region (ORF1a) and p23 gene that were estimated for the B165-, HA16-5-, Kpg3-, RB-, T30-, T36- and VT-like genotypes of CTV from each GFMS12 source (Marsh and Star Ruby grapefruits and Mexican lime). The number of sequences (n) used to estimate nucleotide diversity is shown.

Source	Sequence	CTV genotype						
		B165	HA16-5	Kpg3	RB	T30	T36	VT
GFMS12 Marsh	A-region	0.008 (n=23)	-	n/a	-	n/c	-	-
GFMS12 Star Ruby	A-region	0.065 (n=2)	-	n/a	-	0.01 (n=13)	-	n/c
GFMS12 Mexican Lime	A-region	0.001 (n=8)	-	n/a	0.06(n=17)	-	-	n/c
GFMS12 Marsh	p23 gene	n/a	-	-	-	-	-	0.009 (n=39)
GFMS12 Star Ruby	p23 gene	n/a	-	-	-	-	-	0.009 (n=42)
GFMS12 Mexican Lime	p23 gene	n/a	-	-	0.018 (n=20)	-	-	n/c

n/a = not applicable, differentiation of genotype based on this region not generally accepted

n/c = no comparison possible, single sequence observed

- = none such sequences obtained

Table 4. Biological indexing of GFMS12 source and sub-isolates on different host responses.

Isolate	Host response ^x							
	Mexican lime (ML)		Madam Vinous (SW)		Sour Orange (SO)	SW/SO	Duncan Grapefruit (DG)	
	Vein clearing	Stem-pitting	Stunting	Stem-pitting	Stunting/ Seedling Yellows	Stunting	Stunting	Stem-pitting
Sub-isolate 12-7	+	+	-	-	-/-	+	+	+
Sub-isolate 12-8	++	++	+	+	+/+	+	+	-
Sub-isolate 12-9	++	++	+	-	+/+	+	+	+
GFMS 12	++	++	-	-	-/-	-	+	++

^x symptom consistently observed and mild (+), moderate (++), not observed (-).