

## Chapter eight

## REFERENCES

### 8. Bibliography

- AAA. (2003). *Position Statement: Preventing Noise-Induced Occupational Hearing Loss*. Reston, VA: American Academy of Audiology.
- Abbot, E. J. (1935). Scales for Sound Measurements Used in Machinery Noise Reduction. *JASA*, *VI*, January, 137-150.
- Agrawal, Y., Niparko, J. K., & Dobie, R. A. (2010). Estimating the Effect of Occupational Noise exposure on hearing thresholds: The importance of adjusting for confounding variables. *Ear and Hearing*, *31*, 234–237.
- Agrawal, Y., Platz, E. A., & Niparko, J. K. (2009). Risk Factors for Hearing Loss in US Adults: Data From the National Health and Nutrition Examination Survey, 1999 to 2002. *Otol Neurotol*, *30*, 139-145.
- Agricola, G. (1950). *De Re Metallica* (1950 ed.). (H. Hoover, & L. Hoover, Trans.) New York: Dover.
- Alberti, P. W. (2001). The anatomy and physiology of the ear and hearing. *Occupational exposure to noise: Evaluation, prevention and control*, pp. 51-63.
- AMA. (1955). American Medical Association-Council on Physical medicine and Rehabilitation. Principles for evaluating hearing loss. *JAMA*, *157*(16), 1408-1409.
- Amedofu, G. K. (2002). Hearing-impairment among workers in a surface gold mining company in Ghana. *African Journal of Health Sciences*, *9*(1-2), 91-97.
- American College of Occupational and Environmental Medicine (ACOEM). (2003). ACOEM evidence-based statement: noise-induced hearingloss. *Journal of Occupational and Environmental Medicine*, *45*, 579–581.
- American Medical Association (AMA). (2001). *Guides for the evaluation of permanent impairment* (5th ed.). Chicago: AMA.
- American Speech-Language-Hearing Association . (2007). *Noise and hearing loss*. . Retrieved June 12, 2008, from www.asha.org: <http://www.asha.org/public/hearing/disorders/noise/htm>
- American Speech-Language-Hearing Association. (2004). The audiologist's role in occupational hearing conservation and hearing loss prevention programs. *ASHA Supplement*, *24*, 58-59.
- American Standards Association. (1936). The calibration of microphones . *J. Acoust. Soc. Am.*, *7*(4), 300-305.
- Anglogold Ashanti. (2011). *Sustainability report*. Retrieved May 12, 2012, from www.anglogoldashanti.com: <http://www.anglogoldashanti.com/Sustainability>

AngloGoldAshanti. (2007). *Country report South Africa Vaal River operations*. Vaal River: AngloGold Ashanti.

AngloGoldAshanti. (2007). *Country report South Africa West Wits operations*. West Wits: AngloGold Ashanti.

AngloGoldAshanti. (2010, 11 17). *Employment equity report*. Retrieved 05 17, 2011, from [www.anglogoldashanti.com/NR/rdonlyres/D2FAF96B-EF67-4438-B454-33ED60F9FB99/0/EEReport2010.pdf](http://www.anglogoldashanti.com/NR/rdonlyres/D2FAF96B-EF67-4438-B454-33ED60F9FB99/0/EEReport2010.pdf)

ANSI. (1996). *Determination of Occupational Noise Exposure and Estimation of Noise-Induced Hearing Impairment. ANSI S344*. New York: American National Standards Institute; ANSI S3.44, Acoustical Society of America.

Armitage, P., & Berry, G. (1994). *Statistical methods in medical research* (3 ed.). London: Blackwell.

Attias, J., Bresloff, I., Reshef, I., Horowitz, G., & Furman, V. (1998). Evaluating noise induced hearing loss with distortion product otoacoustic emissions. *British Journal of Audiology*, 32(1), 39-46.

Axelsson, A. (1979). Diagnosis and treatment of occupational noise induced hearing loss. *Acta Otolaryngol, Suppl* 1979(360), 86–87.

Bankaitis, A. E., & Keith, R. W. (1995). Audiological changes associated with HIV infection. *Ear, Nose & Throat Journal*, 74(5), 353-359.

Barnes, D. (2008, November 11). Occupational Health department, AngloGoldAshanti. (S. Strauss, Interviewer)

Baughn, W. L. (1966). Noise control- Precent of population potential. *Int. Aud*, 5, 331-338.

Baughn, W. L. (1971). *Relation Between Daily Noise Exposure and Hearing Loss Based on the Evaluation of 6835 Industrial Noise Exposure Cases*. Pensacola: Aerospace Medical Research Laboratory (AMRL-TR-73-53).

Becker, P. (2008, 05 28). Personal communication with Prof Becker, chief investigator/ statistician at the Medical Research Council. (S. Strauss, Interviewer)

Berger, E. H., Royster, L. H., & Thomas, W. G. (1978). Presumed noise-induced permanent threshold shift resulting from exposure to an A-weighted leq of 89dB. *J Acoust Soc Am*, 64, 192-197.

Berger, K. W. (1976). Genealogy of the words "audiology" and "audiologist". *J Am Audiol Soc*, 2(2), 38-44.

Bohne, B. A., Harding, G. W., & Lee, S. C. (2006). Death pathways in noise-damaged outer hair cells. *Hearing Research*, 223, 61–70.

Bohne, B. A., Zahn, S. J., & Bozzay, D. G. (1985). Damage of the cochlea following interupted exposure to low frequency noise. *Ann. Otol. Rhinol. Laringol*, 94(2 pt. 1), 122-128.

- Botsford, J. H. (1967). Simple Method for Identifying Acceptable Noise Exposures. *J. Acous. Soc. Am.*, 4, 810.
- Bovo, R., Ciorba, A., & Martini, A. (2007). Genetic factors in noise induced hearing loss. *Audiological Medicine*, 5.
- Brewer, E. W., & Stockton, S. (2010). Directional Hypothesis. *Encyclopedia of Research Design*. (N. J. Salkind, Ed.) Thousand Oaks, CA, USA. Retrieved March 4, 2012, from SAGE Reference Online. Web.
- Brits, J., Strauss, S., Eloff, Z., Becker, P. J., & Swanepoel, D. W. (2011). Hearing profile of gold miners with and without tuberculosis . *Occup Environ Med*, 1-7.
- Bunch, C. C., & Raiford, T. S. (1931). Race and sex variations in auditory acuity. *Arch Otolaryngol*, 31, 423-434.
- Carlin, M., & McCroskey, R. (1980). Is eye color a predictor of noise induced hearing loss? *Ear Hear.*, 1, 191-196.
- Carter, H. A. (1942). Review of methods used for estimating percentage loss of hearing. *The Laryngoscope*, 52(11), 879–890.
- Carter, N. L. (1980). Eye colour and susceptibility to noise-induced permanent threshold shift. *Audiology*, 19, 86-93.
- Chamber of Mines. (2007). *Facts and figures 2006/2007*. Retrieved September 6, 2008, from [www.bullion.org.za: http://www.bullion.org.za/content](http://www.bullion.org.za/content)
- Chamber of Mines. (2009). *Annual report 2008-2009*. Retrieved September 5, 2010, from [www.bullion.org.za:  
http://www.bullion.org.za/content/?pid=70&pagename=Annual+Reports](http://www.bullion.org.za/content/?pid=70&pagename=Annual+Reports)
- Chamber of Mines. (2009). *Annual report 2008-2009*. Retrieved September 5, 2010, from [www.bullion.org.za:  
http://www.bullion.org.za/content/?pid=70&pagename=Annual+Reports](http://www.bullion.org.za/content/?pid=70&pagename=Annual+Reports)
- Chamber of Mines. (2012). *Annual report 2010/2011*. Retrieved 05 20, 2012, from [www.bullion.org.za:  
http://www.bullion.org.za/content/?pid=70&pagename=Annual+Reports](http://www.bullion.org.za/content/?pid=70&pagename=Annual+Reports)
- Chamber of Mines. (2012). *Annual report 2010/2011*. Retrieved 05 20, 2012, from [www.bullion.org.za:  
http://www.bullion.org.za/content/?pid=70&pagename=Annual+Reports](http://www.bullion.org.za/content/?pid=70&pagename=Annual+Reports)
- Chamber Of Mines Research Organization (COMRO). (1988). *User guide no11: Guidelines for the implementation and control of a hearing conservation programme in the South African mining industry*. Johannesburg: Chamber of Mines Research Organization.

- Chandrasekhar, S. S., Connelly, P. E., Brahrnbhatt, S. S., Shah, C. S., Kloser, P. C., & Baredes, S. (2000). Otologic and audiologic evaluation of human immunodeficiency virus-infected patients. *American Journal of Otolaryngology*, 121, 1-9.
- Chang, S. J., & Chang, C. K. (2009). Prevalence and risk factors of noise-induced hearing loss among liquefied petroleum gas (LPG) cylinder infusion workers in Taiwan. *Ind Health*, 603-610.
- Chesky, K. (2011). Schools of music and conservatories and hearing loss prevention . *International Journal of Audiology* 2011; 50: S32–S37, 50, S32–S37.
- Cheyip, M. Y., Nelson, G., Ross, M. H., & Murray, J. (2007). South African platinum mine employees reduce smoking in 5 years. *Tobacco Control*, 16, 197–201.
- Ciletti, L., & Flamme, G. A. (2008). Prevalence of hearing impairment by gender and audiometric configuration: results from the national health and nutrition examination survey (1999–2004) and the Keokuk County rural health study (1994–1998). *J Am Acad Audiol*, 19, 672–685.
- Clark, S. (2000). *Five myths in assessing the effects of noise on hearing*. Retrieved August 20, 2008, from www.healthyhearing.com:  
[http://www.healthyhearing.com/hearing\\_library/article\\_content.asp?article\\_id=42](http://www.healthyhearing.com/hearing_library/article_content.asp?article_id=42)
- COIDA. (1994). *Compensation for Occupational Diseases and Injuries Act, 130*. Pretoria: Republic of South Africa.
- COIDA. (2001). *Compensation for Occupational Injuries and Diseases Act, No. 130 of 1993. Circular Instruction No. 171*. South Africa: Government Gazette, no. 22296, A61311, A10/4/3/4.
- Coles, R. R., Lutman, M. E., & Buffin, J. T. (2000). Guidelines on the diagnosis of noise-induced hearing loss for medicolegal purposes. *Clinical Otolaryngology and Allied Sciences*, 25, 264-273.
- Conradie, A. S. (2007). Part two: Review of selected commodities:Gold. In DME, I. Robinson, S. Mabuza, S. Sikhosana, T. Masetlana, J. A. Duval, P. Mwape, . . . D. Dlambulo (Eds.), *South African mining industry*. Pretoria: Department of Minerals and Energy.
- Cooper, C. J. (1994). Ear and race effects in hearing. Health and Nutrition Examination Survey of 1971-75: Part I. *J Am Acad Audiol*, 5, 30–36.
- Cooper, C. J. (1994). Ear and race effects in hearing. Health and Nutrition Examination Survey of 1971-75: Part I. *J Am Acad Audiol*, 5, 30–36.
- Crichton, N. (2000). Information point: prevalence and incidence. *Journal of clinical nursng*, 178-188.
- Cunningham, D. R., & Norris, M. L. (1982). Eye color and noise induced hearing loss: a population study. *Ear and Hearing*, 3, 211-214.
- Dane, F. C. (1990). *Research Methods*. Pacific Grove, California: Brooks/ Cole.

- Davis, A. C. (1989). The prevalence of hearing impairment and reported hearing disability among adults in Great Britain. *Int J Epidemiol*, 18(4), 911–917.
- Davis, H. (1971). A historical introduction. In B. A. Series, & D. Robinson (Ed.), *Volume I- Occupational hearing Loss* (pp. 7-12). New York: Academic Press.
- Davis, H. (1971). A historical introduction. *Brittish Acoustical Society Special Series: Volume I- Occupational Hearing Loss*, 7-12.
- De Koker, E. (2003). *SIM070201. Part one: Evaluate the viability of auditory steady state reponse testing for pseudohypacusis workers in the South African mining industry*. Johannesburg: Safety in Mines Advisory Committee (SIMRAC).
- De Toro, M. A., Ordoñez, R., Reuter, K., & Hammershøi, D. (2011). Is it necessary to penalize impulsive noise +5 dB due to higher risk of hearing damage? *Acoust. Soc. Am.*, 129(6), 3808-3817.
- Deanne, P. (2000). *The First Industrial Revolution. Second Edition*. Cambridge, United Kingdom: Cambridge University Press.
- Dear, T. A. (2006). Noise criteria regarding risk and prevention of hearing injury in industry. In R. T. Sataloff, & J. Sataloff, *Occupational Hearing Loss* (3 ed., pp. 665-686). New York: Taylor & Francis Group.
- DeForge, B. R. (2010). Research Design Principles . *Encyclopedia of research design*. (N. J. Salkind, Ed.) Thousand Oaks, CA, USA. Retrieved February 29, 2012, from Encyclopedia of Research Design: SAGE Reference Online. Web.
- Dekker, J. J., Edwards, A. L., Franz, R. M., van Dyk, R. M., & Banyini, A. (2011). Meeting the milestones: are South African small-to-medium-scale mines up to the task? *Journal of the Southern African Institute of Mining and Metallurgy*, 111, 309-313.
- Department of Minerals and Energy. (1996). *Mine Health and Safety Act (MHSA, Act 29 of 1996)*. Pretoria, South Africa: DME.
- DME. (2003). *DME 16/3/2/4-A3 Guideline for the compilation of a mandatory code of practice for an occupational health program for noise*. Pretoria, South Africa: Department of Minerals and Energy. Mine Health and Safety Inspectorate.
- Dobie, R. A. (1992). The relative contributions of occupational noise and aging in individual cases of hearing loss. *Ear and Hearing*, 13(1), 19-27.
- Dobie, R. A. (2001). *Medical-legal evaluation of hearing loss* (2nd Edition ed.). San Diego, CA: Singular. Thomson Learning.
- Dobie, R. A. (2001). *Medical-legal evaluation of hearing loss* (2nd Edition ed.). San Diego, CA: Singular. Thomson Learning.
- Dobie, R. A. (2005). Noise-induced permanent threshold shift from audiometric shape: The ISO-1999 model. *Ear & Hearing*, 26(6), 630-636.

- Dobie, R. A. (2006). Methodological issues when comparing hearing thresholds of a group with population standards: The case of the ferry engineers. *Ear & Hearing*, 27(5), 526-538.
- Dobie, R. A. (2007). Noise-induced permanent threshold shifts in the occupational noise and hearing survey: An explanation of elevated risk estimates. *Ear and Hearing*, 28(4), 580-591.
- Dobie, R. A. (2008). The burdens of age-related and occupational noise-induced hearing loss in the United States. *Ear & Hearing*, 29(4), 565–577.
- Dollinger, A. (2000). *Diodorus Siculus on mining in Egypt*. Retrieved May 17, 2010, from <http://www.reshafim.org.il/ad/egypt/timelines/topics/diodorus.htm>.
- Dreisbach, L. E., Kramer, S. J., Cobos, S., & Cowart, K. (2007). Racial and gender effects on pure-tone thresholds and distortion-product otoacoustic emissions (DPOAEs) in normal-hearing young adults. *International Journal of Audiology*, 46, 419-426.
- Edelson, J., Neitzel, R., Meischke, F., Daniell, W., Sheppard, L., Stover, B., & Seixas, N. S. (2009). Predictors of hearing protection use in construction. *Ann. Occup. Hyg.*, 53(6), 605-615.
- Edwards, A. (2010). *Unpublished doctoral thesis: Measurement of distortion product otoacoustic emissions in South African gold miners at risk for noise-induced hearing loss*. Johannesburg: Wits.
- Edwards, A., Dekker, J. J., & Franz, R. M. (2011). Profiles of noise exposure levels in South African mining . *Journal of the Southern African Institute of Mining and Metallurgy*, 11, 315-322.
- Ehrlich, J. R., Johnston, M., & Ross, H. (2008). *Handbook of occupational health practice in the South African mining industry*. Braamfontein: The Safety in Mines Research Advisory Committee (SIMRAC).
- Eldred, K. M., Gannon, W. J., & von Gierke, H. E. (1955). *Criteria for short time exposure of personnel to high-intensity jet aircraft noise*. OH: Wright- Patterson AFB.
- Eloff, Z. (2009, November 11). Dr. (S. Strauss, Interviewer)
- EPA. (1973). *Public Health and Welfare Criteria for Noise*. EPA Report No. 550/9-73-002. Washington D.C.: Environmental Protection Agency.
- EU. (2003). Directive 2003/10/EC of the European Parliament and the Council of the European Union on the minimum health and safety requirements regarding the exposure of workers to the risks arising from physical agents (noise). *Official Journal of the European Union*, 38-44.
- Ferrite, S., & Santana, V. (2005). Joint effects of smoking, noise exposure and age on hearing loss. *Occupational Medicine*, 55, 48–53.
- Flamme, G. A., Deiters, K., & Needham, T. (2011). Distributions of pure-tone hearing threshold levels among adolescents and adults in the United States by gender, ethnicity, and age: Results from the US National Health and Nutrition Examination Survey. *International Journal of Audiology*, 50, S11-S20.

- Fletcher, H. (1938). Loudness, masking and their relation to the hearing process and the problem of noise measurement. *JASA*, 9(4), 275-294.
- Fletcher, H., & Munsen, W. A. (1933). Loudness, its definition, measurement and calculation. *JASA*, V, October, 82-108.
- Flood, L. M. (1987). Occupational hearing loss in the United Kingdom. In R. T. Sataloff, & J. Sataloff, *Occupational hearing loss* (pp. 605-612). New York: Marcell Dekker.
- Franks, R. (1996). *Analysis of audiograms for a large cohort of noise-exposed miners*. Cincinnati , OH: National Institute for Occupational Safety and Health.
- Fransen, E., Topsakal, V., Hendrickx, J. J., Van Laer, L., Huyghe, J. R., Van Eyken, E., . . . Manninen. (2008). Occupational noise, smoking, and a high body mass index are risk factors for age-related hearing impairment and moderate alcohol consumption is protective: A European population-based multicenter study. *JARO*, 9, 264–276.
- Franz, R. M., & Phillips, J. I. (2001). Noise and Vibration. In R. Guild, R. I. Ehrlich, J. R. Johnston, & M. H. Ross, *Handbook of Occupational Health Practice in the South African Mining Industry* (pp. 193-230). Braamfontein: The Safety in Mines Research Advisory Committee (SIMRAC).
- Free, E. E. (1930). Practical methods of noise measurement. *Journal of the Acoustical Society of America*, July, 18-29.
- Fuente, A., & Hickson, L. (2011). Noise-induced hearing loss in Asia. *International Journal of Audiology*, 50, S3-S10.
- Galt, R. H. (1929). Methods and Apparatus for Measuring the Noise Audiogram. *Journal of the Acoustical Society of America*, October, 147-157.
- Galt, R. H. (1930). Results of Noise Surveys Part I. Noise Out-of-Doors. *Journal of the Acoustical Society of America*, July, 30-60.
- Gao, W. W., King, D. L., Zheng, X. Y., Ruan, F. M., & Liu, Y. J. (1992). Comparison in the changes in the stereocilia between temporary and permanent threshold shift. *Hearing Research*, 62, 27-41.
- Garth, R. (1994). Blast injury of the auditory system: A review of the mechanisms and pathology. *J. Laryngol. Otol.*, 108, 925-929.
- Gates, G. A., Schmid, P., Sharon, G., Kujawa, A., Nam, B., & D'Agostino, R. (2000). Longitudinal threshold changes in older men with audiometric notches. *Hearing Research*, 141, 220-228.
- Girard, S. A., Picard, M., Davis, A. C., Simard, M., Larocque, R., Leroux, T., & Turcotte, F. (2009). Multiple work-related accidents: tracing the role of hearing status and noise exposure. *Occup. Environ. Med.*, 66, 319-324.
- Given, L. M. (2008). *The Sage encyclopedia of qualitative research methods*. Los Angeles, California: Sage .
- Glorig, A. (1980). Noise: Past, Present and Future. *Ear and Hearing*, 1(1), 4-19.

- Greville, A. (2010, February). *The NAL percentage loss of hearing*. Retrieved 05 06, 2012, from [www.acc.co.nz:](http://www.acc.co.nz/PRD_EXT_CSMP/groups/external_communications/documents/reference_tools/wpc091009.pdf)  
[http://www.acc.co.nz/PRD\\_EXT\\_CSMP/groups/external\\_communications/documents/reference\\_tools/wpc091009.pdf](http://www.acc.co.nz/PRD_EXT_CSMP/groups/external_communications/documents/reference_tools/wpc091009.pdf)
- Guida, H. L., De Sousa, A. L., & Cardoso, A. C. (2012). Relationship between the findings of pure-tone audiometry andtoacoustic emission tests on military police personnel. *International Archives of Otorhinolaryngology*, 16(1), 67-73.
- Guyton, G. P. (1999). A brief history of workers' compensation. *The Iowa Orthopaedic Journal*, 19, 106-110.
- Harding, G. W., & Bohne, B. A. (2007). Distribution of focal lesions in the chinchilla organ of Corti following exposure to a 4-kHz or a 0.5-kHz octave band of noise. *Hearing Research*, 225 , 50–59.
- Harding, G. W., & Bohne, B. A. (2009). Relation of focal hair-cell lesions to noise-exposure parameters from a 4- or a 0.5-kHz octave band of noise. *Hearing Research*, 254 , 54–63.
- Harding, G. W., Bohne, B. A., & Vos, J. D. (2005). The effect of an age-related hearing loss gene (Ahl) on noise-induced hearing loss and cochlear damage from low-frequency noise. *Hearing Research*, 204, 90–100.
- Helleman, H. W., & Dreschler, W. A. (2012). Overall versus individual changes for otoacoustic emissions and audiometry in a noise-exposed cohort. *International Journal of Audiology*, 51(5), 362-372.
- Helleman, H. W., Jansen, E. J., & Dreschler, W. A. (2010). Otoacoustic emissions in a hearing conservation program: General applicability in longitudinal monitoring and the relation to changes in pure-tone thresholds. *International Journal of Audiology*, 49(6), 410-419.
- Henderson, D., & Hamernik, R. P. (1995). *The biological basis of noise-induced hearing loss*. Buffalo: Hearing Research Laboratory, Dept of Communicative Disorders and Science, State University of New York .
- Henderson, D., Bielefeld, E. C., Harris, K. C., & Hu, B. H. (2006). The role of oxidative stress in noise-induced hearing loss. *Ear and Hearing*, 27(1), 1-19.
- Henselman, L. W., Henderson, D., Shadoan, J., Subramaniam, M., Saunders, S., & Ohlin, D. (1995). Effects of noise exposure, race, and years of service on hearing in U.S. Army soldiers. *Ear & Hearing* , 16, 382-391.
- Hermanus, M. A. (2006). The status of occupational health in the South African mining industry and key challenges for practitioners. Paper presented at the 6th International Scientific Conference of the participating International Occupational Hygiene Association. *Journal of the Mine Ventilation Society of South Africa*, 59(1), 8-13.

- Hermanus, M. A. (2007). Occupational health and safety in mining—status, new developments, and concern. *The Journal of The Southern African Institute of Mining and Metallurgy*, 107, 531–538.
- Hessel, P. A., & Sluis-Cremer, G. K. (1987). Hearing loss in white South African goldminers. *South African Medical Journal*, 364-367.
- Ho, P. M., Peterson, P. N., & Masoudi, F. A. (2012). *Key issues in outcomes research: evaluating the evidence. Is there a rigid hierarchy?* Retrieved 06 16, 2012, from American Heart Association: <http://circ.ahajournals.org/content/118/16/1675.full>
- Hoffman, H. J., Dobie, R. A., Ko, C., Themann, C. L., & Murphy, W. J. (2010). Americans hear as well or better today compared with 40 years ago: Hearing threshold levels in the unscreened adult population of the United States, 1959–1962 and 1999–2004. *Ear & Hearing*, 13, 725-734.
- Hyde, M. (2005). Evidence-based practice, ethics and EHDI program quality. In R. C. Seewald, & J. C. Bamford (Ed.), *A sound foundation through early amplification 2004: Proceedings of the third international conference* (pp. 281-303). Stäfa, Switzerland: Phonak AG.
- Ishii, E. K., & Talbott, E. O. (1998). Race/ethnicity differences in the prevalence of noise-induced hearing loss in a group of metal fabricating workers. *J Occup Environ Med*, 40(8), 661–666.
- ISO. (1990). *Acoustics: Determination of Occupational Noise Exposure and Estimation of Noise-induced Hearing Impairment*. Geneva, Switzerland: International Organization for Standardization: ISO-1999.
- ISO. (2010). <http://www.iso.org/iso/about.htm>. Retrieved July 26, 2010, from <http://www.iso.org>.
- Jerger, J. (2009). *Audiology in the USA*. San Diego: Plural Publishing, Inc.
- Johansson, M. S., Arlinger, & D, S. (2002). Hearing threshold levels for an otologically unscreened, non-occupationally noise exposed population in Sweden. *Int J Aud*, 41, 180– 194.
- Johnson, D. L. (1978). *Derivation of presbycusis and noise induced permanent threshold shift (NIPTS) to be used for the basis of a standard on the effects of noise on hearing. (AMRL-TR-78-128)*. Ohio: Aerospace Medical Research Laboratory, Wright-Patterson Air Force Base.
- Johnson, D. L. (1999). The major historical events in the relating of noise-induced hearing loss to noise exposure (A). *J. Acoust. Soc. Am.*, 105(2), 1183-1183.
- Johnson, D. L., Papadopoulos, P., Watka, N., & Takala, J. (2006). *Exposure criteria: Occupational exposure levels*. Retrieved 03 01, 2011, from [www.who.int/occupational\\_health/publications/noise4.pdf](http://www.who.int/occupational_health/publications/noise4.pdf).
- Killion, M. C., Monroe, T., & Drambarean, V. (2011). Better protection from blasts without sacrificing situational awareness. *International Journal of Audiology*, 50, S38–S45.
- Konings, A., Van Laer, L., Wiktorek-Smagur, A., Rajkowska, E., Pawelczyk, M., Carlsson, P. I., . . . Van Camp, G. (2009). Candidate gene association study for noise-induced hearing loss in two independent noise-exposed populations. *Annals of Human Genetics*, 73, 215–224.

- Kopke, R., Bielefeld, E., Liu, J., Zheng, J., Jackson, R., Henderson, D., & Coleman, J. K. (2005). Prevention of impulse noise-induced hearing loss with antioxidants. *Acta Oto-Laryngologica*, 125, 235-243.
- Kothari, C. R. (1990). *Research Methods* (2nd ed.). New Delhi: New Age International Publishers.
- Kryter, K. D. (1950). The effects of noise on man. *Journal of Speech and Hearing, Monograph supplement 1*.
- Kryter, K. D. (1965). Damage risk criterion and contours based on permanent and temporary hearing loss data. *J Am Ind Hyg Assoc*, 26, 34–44.
- Kryter, K. D. (1985). *The effects of noise on man, 2nd edition*. Orlando, FL: Academic Press.
- Kryter, K. D. (2009, June). Acoustical model and theory for predicting effects of environmental noise on people. *J. Acoust. Soc. Am.*, 125(6), 3707-3721.
- Kryter, K. D., & Williams, C. (1960). The loudness and noisiness of sounds of different bandwidth. *Journal of the Acoustical Society of America*, 32(November), 1523.
- Kujawa, S. G., & Liberman, M. C. (2006). Acceleration of age-related hearing loss by early noise exposure: Evidence of a misspent youth. *The Journal of Neuroscience*, 26(7), 2115–2123.
- Kurmis, A. P., & Apps, S. A. (2007). Occupationally-acquired noise-induced hearing loss: A senseless workplace hazard. *International Journal of Occupational Medicine and Environmental Health*, 20(2), 127-136.
- Lawton, B. W. (2005). Variation of young normal-hearing thresholds measured using different audiometric earphones: Implications for the acoustic coupler and the ear simulator. *Int J Audiol*, 44, 444-451.
- Le Prell, C. G., Hensley, B. N., Campbell, K. C., Hall III, J. W., & Guire, K. (2011). Evidence of hearing loss in a ‘normally-hearing’ college-student population. *International Journal of Audiology*, 50, S21–S31.
- Le Prell, C. G., Yamashita, D., Minami, S. B., Yamasoba, T., & Miller, J. M. (2007). Mechanisms of noise-induced hearing loss indicate multiple methods of prevention. *Hearing Research*, 226(1-2), 22-43.
- Leedy, P. D., & Ormrod, J. E. (2001). *Practical research: Planning and design*. New Jersey: Prentice-Hall, Inc.
- Lutman, M. E., & Davis, A. C. (1994). The distribution of hearing threshold levels in the general population aged 18 – 30 years. *International Journal of Audiology*, 33, 327–350.
- Macrae, J. H. (1988). *Improved procedure for determinng percentage loss of hearing*. Report no 118: NAL.
- Majno, G., & Joris, I. (1995). Apoptosis, oncosis, and necrosis. An overview of cell death. *Am. J. Path.*, 146, 3-15.

- Margolis, R. H., & Saly, G. L. (2007). Toward a standard description of hearing loss. *International Journal of Audiology, 46*, 746-758.
- Martini, A., Stephens, D., & Read, A. P. (2007). *Genes, hearing and deafness. From molecular biology to clinical practice*. London: Informa UK Ltd.
- Marvin, H. B. (1932). On the loudness of noise. *Journal of the Acoustical Society of America, January*, 388-392.
- Maxwell, D. L., & Satake, E. (2006). *Research and statistical methods in communication sciences and disorders*. Clifton Park, NY: Thomson Delmar.
- May, J. J. (2000). Occupational Hearing Loss. *American Journal Of Industrial Medicine, 37*, 112-120.
- McBride, D. I. (2004). Noise-induced hearing loss and hearing conservation in mining. *Occupational Medicine, 54*, 290-296.
- McBride, D. I., & Williams, S. (2001). Audiometric notch as a sign of noise induced hearing loss. *Occup. Environ. Med., 58*, 46-51.
- McCulloch, J. (2005). Asbestos, lies and the state: Occupational disease and the South African science. *African Studies, 64*(2), 201-217.
- McIlwain, D. S., Gates, K., & Ciliax, D. (2008). Heritage of army audiology and the road ahead: The army hearing program. *American Journal of Public Health, 98*(128), 2167-2172.
- McReynolds, M. C. (2005). Noise-induced hearing loss. *Air Medical Journal, 73*-79.
- MRC. (2002). *Ethics in research*. Retrieved May 15, 2008, from Medical Research Council: <http://www.mrc.ac.za/ethics/doc>
- MRC. (2002). *Ethics in research*. Retrieved May 15, 2008, from www.mrc.ac.za: <http://www.mrc.ac.za/ethics/doc>
- Mwape, R., Roberts, M. J., & Mokwena, E. (2007). South African mining industry: Part one: South Africa's mineral industry; general review. In I. Robinson, S. Mabuza, S. Sikhosana, T. R. Masetlana, J. A. Duval, R. Mwape, . . . N. Dlambulo (Eds.), *South African Mining Industry* (24 ed.). Pretoria: Department: Mineral and Energy (DME), Republic of South Africa.
- Myers, J. E., & Macun, I. (1989). The sociologic context of occupational health in South Africa. *American Journal of Public Health, 79*(2), 216-224.
- National Institute for Occupational Safety and Health. (1976). Survey of hearing loss in the coal mining industry. *Publication 76-172, June*, 70pp.
- National Institute of Hearing (NIH). (1990). *Noise and hearing loss. NIH Consensus Statement*. Bethesda, MD, USA: NIH.
- Neitzel, R., Seixas, N., Goldman, B., & Daniell, W. (2004). Contributions of non-occupational activities to total noise exposure of construction workers. *Ann. occup. Hyg., 48*(5), 463-473.

- Nelson, D. I., & Schwela, D. (2001). *Occupational exposure to noise: Evaluation, prevention and control*. Retrieved March 03, 2011, from [www.who.int/occupational\\_health/publications/noise.pdf](http://www.who.int/occupational_health/publications/noise.pdf).
- Nelson, D. I., Concha-Barrientos, M., Driscoll, T., Steenland, K., Fingerhut, M., Punnett, L., . . . Corvalan, C. (2005). The global burden of selected occupational diseases and injury risks: Methodology and summary. *American Journal Of Industrial Medicine*, 48, 400-418.
- Nelson, D. I., Nelson, R. Y., Concha-Barrientos, C., & Fingerhut, M. (2005). The global burden of occupational noise-induced hearing loss. *American Journal Of Industrial Medicine*(48), 446-458.
- Newcombe, R. G. (1998). Two-sided confidence intervals for the single proportion. *Statistics in medicine*, 17(8), 857-872.
- NIOSH. (1972). *Criteria for a recommended standard: Occupational exposure to noise*. Washington: US Department of Health, Education and Welfare. Health Services and Mental Health Administration. National Institute for Occupational Safety and Health.
- NIOSH. (1998). *Criteria for a recommended standard; Occupational noise exposure. Revised criteria*. Cincinnati: OH: U.S. Department of Health and Human service, Centre for Disease Control and Prevention, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication No. 98-126.
- Niskar, A. S., Kieszak, S. M., Holmes, A. E., Esteban, E., Rubin, C., & Brody, D. J. (2001). Estimated prevalence of noise induced hearing threshold shifts among children 6 to 19 years of age: the Third National Health and Nutrition Examination Survey, 1988-1994, United States. *Pediatrics*, 108, 40-43.
- NOHSC. (2000). *National Occupational Health and Safety Commission: National standard for occupational noise*, NOHSC: 1007 (2000). Canberra: Australian Government; .
- Nondahl, D. M., Cruickshanks, K. J., Wiley, T. L., Klein, R., & Klein, B. E. (2000). Recreational firearms use and hearing loss. *Arch Fam Med*, 9, 352-357.
- OHSA. (1993). *Occupational Health and Safety Act as amended by Occupational Health and Safety Act no. 181 of 1993*. The South African Department of Labour: Republic of South Africa.
- OHSA. (2003). *Occupational Health and Safety Act, 1993: Noise induced-hearing loss regulations*. South Africa: Government Gazette.
- Ologe, F. E., Akande, T. M., & Olajide, T. G. (2005). Noise exposure, awareness, attitudes and use of hearing protection in a steel rolling mill in Nigeria. *Occup Med*, 55(6), 487-489.
- Osbon, W. O., & Oplinger, K. A. (1933). A new portable meter for noise measurement and analysis. *JASA*, July, 39-47.
- Osei-Lah, V. H., & Yeoh, L. (2010). High frequency audiometric notch: An outpatient clinic survey. *International Journal of Audiology*, 49, 95-98.

- Passchier-Vermeer, W. (1974). Hearing loss due to continuous exposure to steady-state broad-band noise. *Journal of Acoustical Society of America*, 56(5), 1585-1593.
- Passchier-Vermeer, W., & Passchier, W. F. (2000). Noise exposure and public health. *Environmental health perspective*, 108(Suppl 1), 123-131.
- Patuzzi, R. (2002). Non-linear aspects of outer hair cell transduction and the temporary threshold shifts after acoustic trauma. *Audiology & Neuro-otology*, 7, 17-20.
- Perez, R., Freeman, S., & Sohmer, H. (2004). Effect of an initial noise induced hearing loss on subsequent hearing loss. *Hearing Research*, 192, 101–106.
- Phillips, J. I., Heyns, P. S., & Nelson, G. (2007). Comparative study of noise and vibration levels. *Annals of Occupational Hygiene*, 51(3), 305–310.
- Picard, M. (2012, 02 10). Correspondence regarding the use of Yantis (1994) categories for hearing loss. (S. Strauss, Interviewer)
- Picard, M. (2012, 02 10). PhD. (S. Strauss, Interviewer)
- Picard, M., Girard, S. A., Simard, M., Larocque, R., Leroux, T., & Turcotte, F. (2008). Association of work-related accidents with noise exposure in the workplace and noise-induced hearing loss based on the experience of some 240,000 person-years of observation. *Accident Analysis and Prevention*, 40, 1644–1652.
- Plontke, S., Zenner, J., & Tübingen, H. (2004). Current aspects of hearing loss from occupational and leisure noise. In H. J. Schultz-Coulon, *Environmental and occupational health disorders* (pp. 233-325). Videel OHG, Germany.
- Pouryaghoub, G., Mehrdad, R., & Mohammadi, S. (2007). Interaction of smoking and occupational noise exposure on hearing loss: a cross-sectional study. *BMC Public Health*, 7, 137-142.
- Prince, M. M., Stayner, L. T., Smith, R. J., & Gilbert, S. J. (1998). A re-examination of the risk estimates from the NIOSH occupational noise and hearing survey ONHS. *The Journal of the Acoustical Society of America*, 101(2), 950-963.
- Pyykkö, I., Toppila, E., Howard, J. Z., Jacobs, T., & Kentala, E. (2007). Noise related hearing impairment. In A. Martini, D. Stephans, & A. P. Read, *Genes, hearing and deafness. From molecular biology to clinical practice* (pp. 91-109). London: Informa UK. Ltd.
- Pyykkö, I., Toppila, E., Zou, J., & Kentala, E. (2007). Individual susceptibility to noise-induced hearing loss. *Audiological Medicine*, 5, 41-53.
- Rabinowitz, P. M., Pierce Wise, J., Hur Mobo, B., Antonucci, P. G., Powell, C., & Slade, M. (2002). Antioxidant status and hearing function in noise-exposed workers. *Hear. Res.* 173, 164–171. *Hear. Res.*, 173, 164-171.
- Rabinowitz, P. M., Sircar, K. D., Tarabar, S., Galusha, D., & Slade, M. D. (2005). Hearing loss in migrant agricultural workers. *J Agromedicine*, 10(4), 9–17.

- Rabinowitz, R. M., Galusha, D., Slade, M. D., Dixon-Ernst, M., Sircar, K. D., & Dobie, R. A. (2006). Audiogram notches in noise-exposed workers. *Ear & Hearing, December*, 742-750.
- Randera, F. (2007, October 17). Presentation: Chamber of mines. *Held at the Sustainable Development Conference*. Pretoria, South Africa.
- Rashaad-Hansia, M., & Dickinson, D. (2010). Hearing protection device usage at a South African gold mine. *Occupational Medicine (Oxford)*, 60(1), 72-74.
- Raynal, M., Kossowski, M., & Job, A. (2006). Hearing in military pilots: One-time audiometry in pilots of fighters, transports, and helicopters. *Aviat Space Environ Med* 2006, 77(1), 57-61.
- RMA. (2002). *Rand Mutual Assurance Company Limited: Guide to Noise Induced Hearing Loss (NIHL)*.
- Robinson, D. W. (1988). Threshold of hearing as a function of age and sex for the typical unscreened population. *British Journal of Audiology*, 22, 5-20.
- Rosenhall, U. (2003). The influence of ageing on noise-induced hearing loss. *Noise Health*, 5, 47-53.
- Rowe, D. (2003). Occupational hygiene practices in the South African mining industry – inspectorate perspective. *30th International Conference of Safety in Mines Research Institutes* (pp. 337-350). Pretoria: South African Institute of Mining and Metallurgy.
- RSA Compensation Fund. (2006). *Annual report of the compensation fund of the Department of Labour, RSA*. Pretoria: Compensation Fund.
- RSA Department of Labour. (1993). *No.85 of 1993. Occupational Health and Safety Act as amended by the Occupational Health and Safety Amendment Act, No.181 of 1993*. Pretoria: Department of Labour.
- Salmivalli, A. (1979). Military audiological aspects in noise-induced hearing losses. *Acta Otolaryngol, Suppl* 1979, 96-97.
- SANS10083:2004. (2004). *The measurement and assessment of occupational noise for hearing conservation purposes*. Pretoria: South African National Standards.
- SANS10083:2007. (2007). *The measurement and assessment of occupational noise for hearing conservation purposes*. Pretoria: South African National Standards.
- SANS10154-1:2004. (2004). *Calibration of pure-tone audiometers Part 1: Air conduction*. Pretoria: South African National Standards.
- SANS10154-2:2004. (2004). *Calibration of pure-tone audiometers Part2: Bone conduction*. Pretoria: South African National Standards.
- SANS10182:2004. (2004). *The measurement and assessment of acoustic environments for audiometric tests*. Pretoria: South African National Standards.
- Santam. (2005). *Compensation for Occupational Injuries and Diseases Act (COIDA)*. Retrieved April 09, 2010, from

[www.santam.co.za/FAIR/RPL/Compensation%20for%20Occupational%20Injuries%20and%20Diseases%20Act.pdf](http://www.santam.co.za/FAIR/RPL/Compensation%20for%20Occupational%20Injuries%20and%20Diseases%20Act.pdf).

- Sataloff, R. T. (2006). *The Council for Accreditation in Occupational Hearing Conservation: a good model for training allied health personnel*. Retrieved April 25, 2010, from <http://www.thefreelibrary.com/The+Council+for+Accreditation+in+Occupational+Hearing+Conservation:+a...-a0146629316>.
- Sataloff, R. T., & Sataloff, J. (1987). *Occupational hearing loss*. New York: Marcel Dekker, Inc.
- Sataloff, R. T., & Sataloff, J. (2006). *Occupational hearing loss* (3 ed.). New York: CRC publishers.
- Saunders, J. C., Cohen, Y. E., & Szymko, Y. M. (1991). The structural and functional consequences of acoustic injury in the cochlea and peripheral auditory system: A five year update. *Journal of the Acoustical Society*, 1(July), 136-147.
- Saunders, J. C., Dear, S. P., & Schneider, M. E. (1985). The anatomical consequences of acoustic injury: A review and tutorial. *Journal of the Acoustical Society*, 78(3), 833-860.
- Scott, D. F., Grayson, R. L., & Metz, E. A. (2004). Disease and illness in U.S. mining, 1983–2001. *Journal of Occupational and Environmental Medicine*, 46, 1272-1277.
- Seidman, M. (1999). Noise-Induced Hearing Loss (NIHL). *Volta Review*, 101(1), 29.
- Seixas, N. S., Neitzel, R., Stover, B., Sheppard, L., Daniell, B., Edelson, J., & Meischke, H. (2011). Multi-component intervention to promote hearing protector use among construction workers. *International Journal of Audiology*, 50, S46–S56A.
- Shupak, A., Tal, D., Sharoni, Z., Oren, M., Ravid, A., & Pratt, H. (2007). Otoacoustic emissions in early noise-induced hearing loss. *Otology & Neurotology*, 28, 745-752.
- Silverstein, M. (2008). Meeting the challenges of an aging workforce. *American Journal of Industrial Medicine*, 51, 269–280.
- Singh, A., Georgalas, C., Patel, N., & Papesch, M. (2003). ENT presentations in children with HIV. *Clin. Otolaryngol.*, 28, 240-243.
- Sivian, L. J., & White, S. D. (1933). On minimum audible fields. *Journal of the Acoustical Society of America*, 288.
- Śliwińska-Kowlaska, M., Dudarewicz, D., Kotyło, P., Zamysłowska-Szmytko, E., Wlaczek-Łuszczynska, M., & Gajda-Szadkowska, A. (2006). Individual susceptibility to noise- induced hearing loss: Choosing an optimum method of retrospective classification of workers into noise-susceptible and noise resistant groups. *International Journal of Occupational Medicine and Environmental Health*, 19(4), 235 – 245.
- Smith, P. A., Davis, A., Ferguson, M., & Lutman, M. E. (2000). The prevalence and type of social noise exposure in young adults in England. *Noise Health*, 2, 41-56.

- Sobieraj, J. A., Gibbs, S. G., Harris, M. D., & Tarwater, P. M. (2009). Hearing threshold comparisons between 2001-02 NHANES and 2003-05 Fort Bliss U.S. Army service components. *Military Medicine*, 174(1), 42-49.
- Soer, M., Pottas, L., & Edwards, A. L. (2002). Characteristics of noise induced hearing loss in gold miners. *Health SA Gesondheid*, 7(2), 78-85.
- Sonjica, B., & Nogxina, S. (2008). *Presidential mine and health safety audit*. Pretoria: Department of Minerals and Energy.
- Sonnenberg, P., Glynn, J. R., Fielding, K., Murray, J., Godfrey-Faussett, P., & Shearer, S. (2005). How soon after infection with HIV does the risk of tuberculosis start to increase? A retrospective cohort study in South African gold miners. *J Infect Dis*, 191:150e8.
- Stander, I., & Sataloff, R. T. (2006). Formulae differences in state and federal hearing loss compensation. In R. T. Sataloff, & J. Sataloff, *Occupational hearing loss* (3 ed., pp. 798-806). New-York: Taylor & Francis.
- Steenkamp, R. J. (2007). A six sigma-based management model to eliminate the noise-induced hearing loss pandemic in South African mines. *Southern African Business Review*, 11(1), 104-124.
- Stephenson, M. R., Nixon, C. W., & Johnson, D. L. (1980). Identification of the minimum noise level capable of producing a asymptotic temporary threshold shift. *Aviat Space Environ Med*, 51(4), 391-396.
- Sulkowski, W. J. (1980). *Industrial noise pollution and hearing impairment, problems of prevention diagnosis and certification criteria*. Bethesda, MD: NTIS USA.
- Swanepoel, D. W., & Hall, J. W. (2010). Football match spectator sound exposure and effect on hearing: A pretest–post-test study . *SAMJ*, 239-242.
- Szanto, C., & Ionesco, M. (1983). Influence of age and sex on hearing thresholds levels in workers exposed to different intensity levels of occupational noise. *Audiology*, 22, 339-356.
- Tak, S. W., Davis, R. R., & Calvert, G. M. (2009). Exposure to hazardous workplace noise and use of hearing protection devices among US workers—NHANES, 1999–2004. *American Journal of Industrial Medicine*, 52, 358–371.
- teWaterNaude, J. M., Ehrlich, M. I., Churchyard, G. J., Pemba, L., Dekker, K., Vermeis, M., . . . Myers, J. E. (2006). Tuberculosis and silica exposure in South African gold miners. *Occupational and Environmental Medicine*, 63(3), 187-192.
- Toynbee, J. (1860). *Diseases of the ear: Their nature, diagnosis, and treatment*. London: Churchill.
- Uddin, J. F., Dingle, A. F., Sharp, J. F., & Flood, L. M. (2006). Occupational hearing loss in the United Kingdom. In R. T. Sataloff, & J. Sataloff, *Occupational hearing loss* (pp. 859-870). New York: CRC publishers.

- US Department of Labor. (1983). Occupational Health and Safety Administration (OSHA): Occupational noise exposure: Hearing Conservation Amendment, final rule. *Federal Regulations*, 48:9738-85.
- Vermaas, R. L., Edwards, A. L., & Soer, M. (2007, September). Noise exposure in gold miners: utilising audiogram configuration to determine hearing handicap. *Occupational Health Southern Africa*, 16-19.
- Viljoen, D. A., Nie, V., & Guest, M. (2006). Is there a risk to safety when working in the New South Wales underground coal-mining industry while having binaural noise-induced hearing loss? *Internal Medicine Journal*, 36, 180–184.
- Vipperman, J. S., Bauer, E. R., & Babich, D. R. (2007). Survey of noise in coal preparation plants. *J. Acoust. Soc. Am.*, 197-205.
- Welleschik, B. (1979). The effect of the noise level on the occupational hearing loss. Observations carried out in 25,544 industrial workers (author's transl). *Laryngol Rhinol Otol (Stuttg)*, 58(11), 832-841.
- WHO. (2009). *Global Burden of Disease*. Retrieved June 30, 2010, from www.who.int: [www.who.int/topics/global\\_burden\\_of\\_disease/en/](http://www.who.int/topics/global_burden_of_disease/en/)
- WHO. (2009a). *Global Health Risk: Mortality and burden of disease attributable to selected major risks*. Geneva, Switzerland: World Health Organization.
- WHO. (2011). *Burden of disease from environmental noise-Quantification of healthy life years lost in Europe*. Bonn,: The WHO European Centre for Environment and Health.
- Wikipedia. (2010). *The Industrial Revolution*. Retrieved April 9, 2010, from www.wikipedia.org.
- Wild, D. C., Brewster, M. J., & Banerjee, A. R. (2005). Noise-induced hearing loss is exacerbated by long-term smoking. *Clin. Otolaryngol.*, 30, 517–520.
- Wilson, R. H. (2011). Some observations on the nature of the audiometric 4000 Hz notch: data from 3430 veterans. *Journal of the American Academy of Audiology*, 22, 23-33 .
- World Health Organisation. (2012). *Country statistics: South Africa*. Retrieved 06 02, 2012, from www.who.int: <http://www.who.int/countries/zaf/en/>
- World Health Organization (WHO). (1997). *Prevention of noise-induced hearing loss. Report of an informal consultation*. Geneva : WHO.
- World Health Organization. (2009). *TB/HIV Factsheet*. Retrieved March 2012, from www.who.int: [http://www.who.int/tb/challenges/hiv/factsheet\\_hivtb\\_2009update.pdf](http://www.who.int/tb/challenges/hiv/factsheet_hivtb_2009update.pdf)
- Wu, T. N., Liou, S. H., Shen, C. Y., Hsu, C. C., & Chao, S. L. (1998). Surveillance of noise-induced hearing loss in Taiwan, ROC: A report of the PRESSNIHL results. *Prev Med*, 27, 65 – 69.
- Yantis, P. A. (1994). Pure-tone air-conduction threshold testing. In J. Katz (Ed.), *Handbook of clinical audiology* (4 ed., pp. 97-108). Baltimore: Williams and Wilkens.

Zinsser, M E. (2006). *Annual Report 2005/2006*. Pretoria: Mine Health and Safety Council (MHSC).



## Appendices

---

### Appendix A

### Ethical clearance and collaboration agreements



## **Appendix B**

# **Structure and mandate of the Mine Health and Safety Council (MHSC)**



## **Appendix C**

**COIDA, Compensation for Occupational  
Injuries and Diseases Act, No. 130 of 1993.  
Circular Instruction No. 171, 2001**



UNIVERSITEIT VAN PRETORIA  
UNIVERSITY OF PRETORIA  
YUNIBESITHI YA PRETORIA

## Appendix D

# Equipment calibration certificate