

Phytosociology of the north-eastern Transvaal high mountain grasslands

by

Priscilla Merle Burgoyne

Submitted in partial fulfilment of the requirements for the degree of

MAGISTER SCIENTIAE

In the Department of Botany
Faculty of Biological and Agricultural Sciences

University of Pretoria

PRETORIA

Supervisor: Prof. Dr. G.J. BREDENKAMP
Co-supervisor: Prof. Dr. N. VAN ROOYEN

PREFACE

"THE HIGHEST FUNCTION OF ECOLOGY
IS UNDERSTANDING CONSEQUENCES"

Dune, pp. 314

The novel by Frank Herbert.

TABLE OF CONTENTS

CHAPTER 1: INTRODUCTION	1
CHAPTER 2: STUDY AREA	3
2.1 LOCALITY	3
CHAPTER 3: PHYSICAL FACTORS AFFECTING THE VEGETATION	5
3.1 GEOLOGY	5
3.1.1 Pretoria Group of the Transvaal Sequence.	5
3.1.2 Bushveld Igneous Complex	12
3.1.3 Ecca Series of the Karoo Supergroup	14
3.1.4 Quarternary Deposits	14
3.2 MINING IN THE AREA	15
3.3 LAND TYPES	18
3.4 PHYSIOGRAPHY	20
3.4.1 Topography	20
3.4.2 Drainage	21
3.4.3 Altitude	23
3.5 CLIMATE	26
3.5.1 Insolation	26
3.5.2 Temperature	28
3.5.3 Precipitation	30
Rainfall	30
Fog or mist	32
Hail	34
Frost	34
Snow	37
3.5.4 Wind	38
CHAPTER 4: BIOTIC FACTORS AFFECTING THE VEGETATION	43
4.1 HISTORY	43
4.2 PREVIOUS BOTANICAL AND VEGETATION SURVEYS IN THE AREA	47
4.3 EXPLOITATION AND LAND-USSES	49

4.4 FIRE AND GRAZING	52
4.5 EROSION	55
CHAPTER 5 PHYTOSOCIOLOGICAL METHODS	56
5.1. ANALYTICAL PHASE	56
5.1.1 Reconnaissance	56
5.1.2 Sampling strategy	57
5.1.3 Distribution of sample sites	57
5.1.4 Phytosociological sampling	59
5.1.5 Floristic composition	62
5.2 SYNTHETIC PHASE	63
5.2.1 Classification	63
5.2.2 The Braun-Blanquet method	65
5.2.3 Ordination	65
CHAPTER 6: RESULTS	68
6.1 WETLANDS.	68
6.1.1 The <u>Phragmites australis</u> deep wetlands.	70
6.1.1.1 <u>Phragmites australis</u> - <u>Ficinia acuminata</u> wetlands.	71
6.1.1.2 <u>Phragmites australis</u> - <u>Senecio microglossus</u> deep wetlands. .	73
6.1.2 <u>Miscanthus junceus</u> wetlands.	74
6.1.2.1 <u>Alepidea amatymbica</u> - <u>Miscanthus junceus</u> moist river-banks.	74
6.1.2.2 <u>Agrostis gigantea</u> - <u>Miscanthus junceus</u> moist grassland. .	75
6.1.2.3 <u>Panicum schinzii</u> - <u>Miscanthus junceus</u> shallow wetlands. .	76
6.1.2.4 <u>Carex cognata</u> - <u>Miscanthus junceus</u> wetlands.	77
6.1.2.5 <u>Ischaemum fasciculatum</u> - <u>Miscanthus junceus</u> wetlands. .	80
6.1.3 <u>Eragrostis biflora</u> - <u>Stiburus alopecuroides</u> moist grassland. .	81
6.1.3.1 <u>Helichrysum aureonitens</u> - <u>Eragrostis biflora</u> - <u>Stiburus</u> .	82
6.1.3.2 <u>Disa patula</u> - <u>Eragrostis biflora</u> - <u>Stiburus alopecuroides</u> .	84
6.1.4 <u>Arundinella nepalensis</u> moist turf grassland.	86
6.1.4.1 <u>Hypericum hirtellum</u> - <u>Arundinella nepalensis</u> moist turf grassland.	86
6.1.4.2 <u>Imperata cylindrica</u> - <u>Arundinella nepalensis</u> moist turf grassland.	87
6.2 BOULDERIES.	90
6.2.1 <u>Koeleria capensis</u> drainage lines	91

6.2.1.1 <u><i>Diospyros whyteana</i></u> - <u><i>Koeleria capensis</i></u> steep wet woodlands.	91
6.2.1.2 <u><i>Pentaschistis natalensis</i></u> - <u><i>Koeleria capensis</i></u> moist rocky grassland.	92
6.2.2 <u><i>Protea caffra</i></u> moist sparse woodland	93
6.2.2.1 <u><i>Protea caffra</i></u> - <u><i>Senecio erubescens</i></u> sparse open woodland.	94
6.2.2.2 <u><i>Protea caffra</i></u> - <u><i>Themeda triandra</i></u> sparse open woodland.	96
6.2.2.3 <u><i>Aristida junciformis</i></u> moist rocky grassland.	97
6.2.3 <u><i>Coleochloa setifera</i></u> - <u><i>Aristida junciformis</i></u> moist rocky grassland.	97
6.2.4 <u><i>Dicoma anomala</i></u> - <u><i>Aristida junciformis</i></u> moist rocky grassland	98
6.2.5 <u><i>Vernonia natalensis</i></u> - <u><i>Protea roupelliae</i></u> moist rocky woodland.	100
6.3 GRASSLANDS.	103
6.3.1 <u><i>Loudetia simplex</i></u> grassland.	106
6.3.1.1 <u><i>Rabdosiella calycina</i></u> - <u><i>Loudetia simplex</i></u> moist rocky grassland.	106
6.3.1.2 <u><i>Koeleria capensis</i></u> - <u><i>Loudetia simplex</i></u> moist grassland.	110
6.3.1.3 <u><i>Pteridium aquilinum</i></u> - <u><i>Loudetia simplex</i></u> bracken patches.	112
6.3.1.4 <u><i>Harpochloa falx</i></u> - <u><i>Loudetia simplex</i></u> broken rocky grassland.	113
6.3.1.5 <u><i>Andropogon schirensis</i></u> - <u><i>Loudetia simplex</i></u> grassland.	114
6.3.2 <u><i>Eragrostis chloromelas</i></u> grassland.	121
6.3.2.1 <u><i>Aristida aequiglumis</i></u> - <u><i>Eragrostis chloromelas</i></u> moist grassland.	123
6.3.2.2 <u><i>Agrostis eriantha</i></u> - <u><i>Eragrostis chloromelas</i></u> moist grassland.	124
6.3.2.3 <u><i>Phymaspernum acerosum</i></u> - <u><i>Eragrostis chloromelas</i></u> grassland.	124
6.3.2.4 <u><i>Hyparrhenia anamesa</i></u> - <u><i>Eragrostis chloromelas</i></u> tall grassland.	127
6.3.2.5 <u><i>Helichrysum rugulosum</i></u> - <u><i>Eragrostis chloromelas</i></u> grassland.	128
6.3.3 <u><i>Gladiolus ecklonii</i></u> - <u><i>Themeda triandra</i></u> grassland.	128
6.3.3.1 <u><i>Crocosmia paniculata</i></u> - <u><i>Gladiolus ecklonii</i></u> - <u><i>Themeda</i></u>	129
6.3.3.2 <u><i>Cymbopogon validus</i></u> - <u><i>Gladiolus ecklonii</i></u> - <u><i>Themeda</i></u> <u><i>triandra</i></u> grassland.	131

6.3.3.3 <u>Acacia karroo</u> - <u>Gladiolus ecklonii</u> - <u>Themeda triandra</u> open woodland.	131
6.3.3.4 <u>Halleria lucida</u> - <u>Gladiolus ecklonii</u> - <u>Themeda triandra</u> rocky grassland.	133
6.3.4 <u>Bromus firmior</u> grassland.	134
 CHAPTER 7 ENDEMISM AND BIODIVERSITY	137
7.1 THREATENED SPECIES	137
7.2 ENDEMISM	138
7.3 PLANT SPECIES RICHNESS	139
 CHAPTER 8: DISCUSSION	145
8.1 SITE SELECTION.	145
8.3 DATA ANALYSIS.	146
8.3 TIME AT WHICH SAMPLING TOOK PLACE.	147
8.4 SIMILARITIES BETWEEN OTHER AREAS	147
8.5 STATUS OF THE COMMUNITIES	148
8.5 COMPARISON WITH ACOCK'S SURVEY.	148
8.6. SPECIFICS OF SPECIES	149
 CHECKLIST OF SPECIES	151
 REFERENCES.	202

LIST OF ILLUSTRATIONS

Figure 1. Location of the study area.	4
Figure 2. The deposition of the Pretoria Group in the eastern Transvaal.	6
Figure 3. Geological Groups occurring in the study area.	9
Figure 4. Example of a diabase intrusion near Dullstroom.	13
Figure 5. Mines and mining activities found in the study area.	16
Figure 6. Land types occurring in the study area.	19
Figure 7. Map of the study area showing altitude and rivers.	22
Figure 8. Transect of the study area including geological strata (this transect was taken from Figure 3).	24
Figure 9. Different physiographical units occurring in the study area (Scheepers 1988).	25
Figure 10. Line chart comparing the cloud cover of three weather stations in the area.	27
Figure 11. Rainfall map of the study area (Weather Bureau 1968a) underlain by topography.	31
Figure 12. Line chart showing the monthly occurrence of mist at three weather stations in the area (Weather Bureau 1968b).	33
Figure 13. Line chart of the monthly occurrence of hail at four stations in the study area (Weather Bureau 1968b).	35
Figure 14. Duration of frost for two stations in the study area at 10% and 80% probabilities (Weather Bureau 1968b).	36
Figure 15. Wind roses for Lydenburg during four months of the year (Weather Bureau 1968b).	39
Figure 16. Walter Climate Diagrams for four weather stations in the study area.	41
Figure 17. Ruins of an ancient settlement ± 2km east of Machadodorp. .	44
Figure 18. Graves from the Anglo-Boer war ± 16 km north of Dullstroom on the road to Lydenburg.	46
Figure 19. Acocks' classification of the eastern Transvaal (Taken from Acocks 1988).	48
Figure 20. Sheet and donga erosion in the south-eastern part of the study area.	54

Figure 21. The <u><i>Phragmites australis</i></u> - <u><i>Ficinia acuminata</i></u> wetlands showing the dominant reed <u><i>Phragmites australis</i></u>	72
Figure 22. <u><i>Carex cognata</i></u> dominates the <u><i>Carex cognata</i></u> - <u><i>Miscanthus junceus</i></u> wetland (<u><i>Nerine angustifolia</i></u> in foreground).	79
Figure 23. The <u><i>Helichrysum aureonitens</i></u> - <u><i>Eragrostis biflora</i></u> - <u><i>Stiburus alopecuroides</i></u> moist grassland.	83
Figure 24. Model of wetland degradation.	89
Figure 25. <u><i>Protea caffra</i></u> - <u><i>Themeda triandra</i></u> sparse open woodland showing <u><i>Protea caffra</i></u>	95
Figure 26. <u><i>Dicoma anomala</i></u> - <u><i>Aristida junciformis</i></u> moist rocky grassland showing the dominant grass <u><i>Aristida junciformis</i></u>	99
Figure 27. <u><i>Vernonia natalensis</i></u> - <u><i>Protea roupelliae</i></u> moist rocky woodland.	101
Figure 28. <u><i>Rabdosiella calycina</i></u> - <u><i>Loudetia simplex</i></u> - <u><i>Rhus montana</i></u> moist rocky grassland	107
Figure 29. <u><i>Pteridium aquilinum</i></u> - <u><i>Loudetia simplex</i></u> bracken patches showing almost pure stands of <u><i>Pteridium</i></u>	111
Figure 30. <u><i>Andropogon shirensis</i></u> - <u><i>Loudetia simplex</i></u> - <u><i>Clerodendrum triphyllum</i></u> grassland showing <u><i>Clerodendrum triphyllum</i></u>	119
Figure 31. <u><i>Andropogon schirensis</i></u> - <u><i>Loudetia simplex</i></u> - <u><i>Indigofera sanguinea</i></u> grasslands showing <u><i>Indigofera sanguinea</i></u> when flowering.	120
Figure 32. The blue-green leaves of <u><i>Tristachya leucothrix</i></u> are prominent in the <u><i>Eragrostis chloromelas</i></u> grasslands.	122
Figure 33. <u><i>Phymaspermum acerosum</i></u> distinguishes the <u><i>Phymaspermum acerosum</i></u> - <u><i>Eragrostis chloromelas</i></u> grassland.	126
Figure 34. <u><i>Crocosmia paniculata</i></u> - <u><i>Gladiolus ecklonii</i></u> - <u><i>Themeda triandra</i></u> grassland.	130
Figure 35. The <u><i>Acacia karroo</i></u> - <u><i>Gladiolus ecklonii</i></u> - <u><i>Themeda triandra</i></u> community.	132
Figure 36. The <u><i>Halleria lucida</i></u> - <u><i>Gladiolus ecklonii</i></u> - <u><i>Themeda triandra</i></u> rocky grassland (<u><i>Leucosidea sericea</i></u> dominates the vegetation).	134
Figure 37. The <u><i>Bromus firmior</i></u> grassland showing the prominent grass <u><i>Bromus firmior</i></u>	136
Figure 38. Plot of number of species against lithological groups.	141
Figure 39. Line chart showing the species per 200 m ² per community	

for the wetlands.	142
Figure 40. Line chart showing the species per 200 m ² per community for the boulderries.	143
Figure 41. Line chart showing the species per 200 m ² per community for the grasslands.	144

ABSTRACT

Phytosociology of the north-eastern Transvaal high mountain grassland

by

PRISCILLA MERLE BURGOYNE

Supervisor: Prof. Dr. G.J.BREDENKAMP

Co-supervisor: Prof. Dr. N. VAN ROOYEN

DEPARTMENT OF BOTANY

UNIVERSITY OF PRETORIA

MAGISTER SCIENTIAE

The vegetation of the high mountain grasslands of the north-eastern Transvaal was sampled by using stratification based on geology and land types. Data was classified by TWINSPAN procedures and refined by using the Braun-Blanquet method. This resulted in the recognition of three vegetation units namely wetlands, boulderies and grasslands which represent a moisture gradient. Wetlands comprised four major communities and ten minor plant communities while boulderies resulted in four major communities and seven minor communities. The grasslands comprised four major plant communities which were subdivided into twenty eight minor communities.

All identified communities were described and ecologically interpreted and a species list was compiled combining the data gathered in the area by other authors.