

Biological activities of extracts and isolated compounds from *Bauhinia galpinii* (Fabaceae) and *Combretum vendae* (Combretaceae) as potential antidiarrhoeal agents

Ahmed Aroke Shahid

BSc (Chemistry) (ABU), MSc (Chemistry) (UNILAG)

A dissertation submitted in fulfilment of the requirements for the degree of Doctor of

Philosophy (PhD)

in the

Phytomedicine Programme, Department of Paraclinical Sciences,

Faculty of Veterinary Science



UNIVERSITEIT VAN PRETORIA
UNIVERSITY OF PRETORIA
YUNIBESITHI YA PRETORIA

Promoter: Prof. Jacobus N. Eloff

Co-promoters: Dr. Nivan Moodley (CSIR)

Prof. Vinny Naidoo

Dr. Lyndy McGaw

January 2012

Declaration

The research work described in the thesis was conducted in the Phytomedicine Programme in the Department of Paraclinical Sciences, Faculty of Veterinary Science, University of Pretoria under the supervision of Professor JN. Eloff, Dr. N. Moodley, Prof. V. Naidoo and Dr. LJ. McGaw

The results presented herewith were generated from my own experiments, except where the work of others are quoted and referenced. There is no part of this work that has been submitted to any other University.

Aroke Shahid, Ahmed

Dedication

This work is dedicated to the memory of the following: My Father (Late Mr. Ahmed Aninya Aroke), my brothers (Late Salihu Aroke and Late Ibrahim Onimisi Ahmed), Late Olukemi Ore Udom (A friend and colleague who started her PhD, but could not finish the programme before death) and my dear sister (Late Mrs. Husseinatu Ohunene Abubakar).

Acknowledgements

This Ph.D. study would not have been possible without the support and encouragement of many people. I will thank my mum Mrs Omeneke Aminatu Ahmed who has been the pillar of my life. The love of my wife Mrs Rabiatu Isoyiza Ahmed and my children Enehu Mazidah, Onize Nusrah, Ometere Shamsiyah, Eneze Azeemah and Ava'ami AbdulAleem have been essential to my success. I thank you all for your understanding and endurance during my absence. I also wish to thank my siblings Mr. Aroke Haruna, Mr. Umar Omeiza Aroke and his wife Rukkayat, Mrs. Zulaiha Mubarak, Dr. Halidu Aroke Ahmed, Mrs. Khaltum Khamilu and Aisha Ahmed for their enormous support. I am also deeply indebted to my friends who stuck by me through the years: Dr. Muhammed Awwalu Usman, Engr. Yakubu Adajah, Mr. Yahaya Ohida Yusuf, Mr. Tijani Muhammed Isah, Abdulmumuni Enesi Umar, Dr. Muhammed Onujagbe Onoda, Mrs. Ukachi Ezenwa Igbo, Dr. Caroline Anyakorah, Dr. Oluwatoyin Taiwo and Dr. Chima Cartney Igwe among others. Financial support was a crucial element of my ability to devote so much time to this research. Major support that keeps me and my family afloat during the study was provided by a study leave with pay provide by Federal Institute of Industrial Research Oshodi (FIIRO), Lagos, Nigeria. The University of Pretoria has provided support through University bursary, National Research Foundation (NRF) of South Africa provide fund for the research, Faculty of Veterinary Science also provide research fund through the research committee (RESCOM), and Department of Paraclinical Science also provide research fund.

I am extremely grateful to my supervisor, Prof. J. N. Eloff (overall supervisor) and my co-supervisors, Dr. Nivan Moodley (characterization of isolated compounds), Prof. Vinny Naidoo (isolated organ studies), and Dr. Lyndy J. McGaw (cellular toxicology) for allowing me to tap from the wealth of their knowledge and also giving me much of their time and energy through the duration of this study. I have a great respect and appreciated the instruction and assistance I received from each and every one of you. I sincere thank the Secretary to Phytochemical Programme, Tharien de Winnaar for all her assistance in coordinating the purchase of materials and other important aspects of this project.

I feel honoured to have opportunity to do some of the work at Bioscience, CSIR South Africa and lucky to meet with Dr. Vinesh Maharaj, Dr. Jacqueline Ndlebe, and Ms. Teresa Faleschini who are friends as well as crucial resources of information, techniques and instructions. I express my gratitude to CSIR for allowing me to use her NMR spectroscopic facilities.

At the UPBRC where I got my isolated organ part, the assistance of Dr. Tamsyn Pulleer, Mrs. Stephanie Keulder and Mrs. Ilse Janse van Rensburg were highly appreciated. The ability to collect plant material was essential to this project, and for that I want to recognize the help and outstanding collaboration of Ms Magda Nel of the Manie van der Schyff Botanical Garden and Ms Elsa van Wyk (Curator of the HGW Schweickert Herbarium of the University of Pretoria) for assistance in collection, identification and authentication of the plant samples. My appreciation also goes to Ms Lita Pauw of the Phytomedicine Programme for allowing me access her stored

plant samples. I also appreciated working in harmony with all other students of the Phytomedicine Programme, most especially Mr. Thanyani Ramandwa, Ms Bellonah Sakong, Ms Salaelo Raphatlelo, Ms Imelda Ledwaba and Mrs Edwina Muleya. I thank you all for your co-operation and immeasurable assistance.

Table of Contents

	Page number
Abstract	xiv
List of abbreviations	xvi
List of figures	xix
List of tables	xxi
List of appendix	xxii
 CHAPTER ONE	
Gastrointestinal disorders in diarrhoea diseases mechanisms and medicinal plants potentiality as therapeutic agents	
1.0. Introduction	1
1.1. Plant metabolites as potential therapeutic agent	2
1.2. Aims	3
1.3. Specific objectives	4
1.4. Hypothesis	4
 CHAPTER TWO	
2.0 Literature review	
2.1. Diarrhoea as a disease	5
2.2. Pathophysiology of Diarrhoea	6
2.3. Detailed pathophysiology of diarrhoea	8
2.3.1. Inflammation in diarrhoea	8
2.3.2. Oxidative damage in diarrhoea	11
2.3.3. Enteric nervous system in diarrhoea	15
2.3.4. Cystic fibrosis transmembrane conductance regulator (CFTR) regulation	16
2.4. Specific Agents of Diarrhoea	16
2.4.1. Bacterial causes of diarrhoea	16
2.4.1.1. <i>Escherichia coli</i>	16
2.4.1.2. <i>Staphylococcus aureus</i>	17
2.4.1.3. <i>Campylobacter jejuni</i>	18
2.4.1.4. <i>Shigella</i> spp	18
2.4.1.5. <i>Vibrio cholerae</i>	18
2.4.1.6. <i>Bacillus cereus</i>	19
2.4.1.7. <i>Yersinia enterocolitica</i>	19
2.4.1.8. <i>Listeria monocytogenes</i>	20

2.4.1.9. <i>Clostridium</i> spp	20
2.4.1.10. <i>Salmonella typhimurium</i>	20
2.4.1.11. <i>Enterococcus faecalis</i>	21
2.5. Fungal induced diarrhoea symptoms	21
2.5.1. <i>Candida albicans</i>	21
2.6. Viral induced diarrhoea	21
2.6.1. Rotavirus	21
2.6.2. Norovirus	22
2.6.3. Hepatitis A virus	22
2.6.4. Human immunodeficiency virus (HIV)	22
2.7. Protozoa induced diarrhoea	22
2.7.1. <i>Giardia intestinalis</i>	22
2.7.2. <i>Entamoeba histolytica</i>	23
2.7.3. <i>Cryptosporidium parvum</i>	23
2.7.4. <i>Cyclospora cayetanensis</i>	23
2.8. Parasitic induced diarrhoea	23
2.8.1. <i>Trichinella spiralis</i>	23
2.9. Immune disordered induced diarrhoea	24
2.9.1. Compromised immune system	24
2.9.2. Hyperactive immune system	24
2.10. Antibiotic therapy induced diarrhoea	24
2.10.1. Antibiotic toxicity	24
2.10.2. Alteration of digestive functionality	25
2.10.3. Overgrowth of pathogenic microorganisms	25
2.11. Diabetic complications induced diarrhoea	25
2.12. Food allergy induced diarrhoea	26
2.13. Potential mechanisms in the control of diarrhoea	26
2.13.1. Oxidative damage and antioxidants in diarrhoeal management	26
2.13.2. Inflammation and anti-inflammatory agents in diarrhoea management	26
2.13.3. Enteric nervous system in diarrhoea symptoms and treatment	26
2.14. Plants as potential source of therapeutic agents in alleviating diarrhoeal symptoms	28
2.14.1. Anti-infectious mechanisms of plant secondary metabolites against diarrhoeal pathogens	28
2.14.2. Antioxidative mechanisms of plant phytochemical as potential antidiarrhoeal agents	29
2.14.3. Anti-inflammatory mechanisms of plant phytochemical in diarrhoea management	29
2.14.4. Antidiarrhoeal mechanisms of plant phytochemical	30
2.15. Classification of phytochemicals with antidiarrhoea potential	30
2.15.1. Terpenoids	30

2.15.2. Alkaloids	33
2.15.3. Phenolic	35
2.16. Ethnobotany and scientific investigation of plant species used traditionally in treating diarrhoea in South Africa	38
2.17. Conclusion	38
CHAPTER THREE	
Plant selection, collection, extraction and analysis of selected species	
3.1. Introduction	39
3.2. Solid-liquid extraction	40
3.3. Liquid-liquid fractionation	41
3.4. Thin layer chromatography (TLC)	41
3.4.1. Phytochemical fingerprints	41
3.5. Materials and Methods	42
3.5.1. Selection of South Africa medicinal plants for antidiarrhoeal screening	42
3.5.2. Collection of plant materials	42
3.5.3. Preparation of plant material and optimization of phenolic-enriched extraction process	42
3.5.4. Phytochemical profiling	44
3.6. Quantification of the phenolic constituents of the extracts	45
3.6.1. Determination of total phenolic constituents	45
3.6.2. Determination of total tannin	45
3.6.3. Determination of proanthocyanidin	45
3.6.4. Determination of condensed tannin	46
3.6.5. Determination of hydrolysable tannin (gallotannin)	46
3.6.6. Determination of total flavonoids and flavonol	46
3.6.7. Determination of anthocyanin	47
3.7. Results	47
3.7.1. Yield of extractions and fractionations processes	47
3.7.2. Phytochemical screening (fingerprints)	48
3.7.3. Phenolic composition of the crude extracts	52
3.8. Discussion	57
3.8.1. Yield	57
3.8.2. Thin layer chromatogram	57
3.8.3. Phenolic constituents of the crude extract	58
3.9. Conclusion	60
CHAPTER FOUR	
Antimicrobial activities of the plant extracts against potential diarrhoeal pathogens	
4.0. Introduction	61

4.1. Qualitative antimicrobial (Bioautography) assay	62
4.2. Quantitative antimicrobial activity (Minimum inhibitory concentration (MIC)) assay	63
4.3. Selection of microorganisms used in the study	63
4.4. Material and Methods	64
4.4.1. Microorganism strains	64
4.4.2. Culturing of the Bacteria	64
4.4.3. Bioautography against some pathogenic microorganisms	64
4.4.4. Determination of Minimum Inhibitory Concentration (MIC) against the bacteria pathogens	64
4.4.5. Determination of Minimum Inhibitory Concentration (MIC) against the fungal pathogens	65
4.5. Results	65
4.5.1. Microbial bioautography	65
4.5.2. Minimum inhibitory concentration against bacteria	70
4.5.3. Minimum inhibitory concentration (MIC)	73
4.6. Discussion	75
4.6.1. Antimicrobial bioautography	75
4.6.2. Minimum inhibitory concentration (MIC)	75
4.7. Conclusion	77
CHAPTER FIVE	
Free radical scavenging and antioxidant activities of the extracts and fractions as antidiarrhoeal mechanism	
5.1. Introduction	79
5.1.1. Superoxide ion	81
5.1.2. Hydrogen peroxide	81
5.1.3. Hydroxyl radical	81
5.1.4. Peroxyl radical	82
5.1.5. Hypochlorous acid	82
5.1.6. Nitric oxide	82
5.2. Antioxidant assays	83
5.2.1. Antioxidant bioautography	83
5.2.2. The chemistry of some common antioxidant assays	83
5.2.2.1. Hydroxyl radical	83
5.2.2.2. Hydrogen peroxide scavenging	84
5.2.2.3. Superoxide scavenging capacity	84
5.2.2.4. DPPH	84
5.2.2.5. ABTS	85
5.2.2.6. Ferric reducing antioxidant power (FRAP)	85
5.3. Materials and Methods	85

5.3.1. Antioxidative profile of the crude extracts and fractions using DPPH radical solution	86
5.3.2. Antioxidative assays	86
5.3.2.1. DPPH free radical-scavenging method	86
5.3.2.2. ABTS free radical-scavenging method	86
5.3.2.3. Ferric reducing antioxidant power (FRAP)	87
5.3.2.4. Hydroxyl radical scavenging assay	87
5.3.2.5. Lipid peroxidation inhibition assay	87
5.4. Result	87
5.4.1. TLC-DPPH analyses	87
5.4.2. DPPH effective concentration (EC ₅₀)	90
5.4.3. ABTS effective concentration (EC ₅₀)	92
5.4.4. FRAP gradient	93
5.4.5. Hydroxyl radical effective concentration (EC ₅₀)	94
5.4.6. Lipid peroxidation inhibition effective concentration (EC ₅₀)	95
5.5. Discussion	96
5.5.1. Qualitative antioxidant analyses (DPPH-TLC bioautography)	96
5.6. Conclusion	98
CHAPTER SIX	
Anti-inflammatory activities of the crude extracts as antidiarrhoeal mechanisms	
6.0. Introduction	100
6.1. Effect of cyclooxygenases (COX) on GIT	101
6.2. Effects of lipoyxygenase (LOX) on GIT	101
6.3. Effects of cytokines on GIT	102
6.4. Oxidative species as inflammatory mediator	102
6.5. Allopathic anti-inflammatory therapies and adverse effects on GIT	103
6.6. Plant phytochemicals as anti-inflammatory agents	105
6.7. Mechanisms of anti-inflammatory assay models	105
6.8. Materials and Methods	106
6.8.1. COX assay	106
6.8.2. LOX assay	106
6.9. Results	107
6.9.1. COX	107
6.9.2. LOX	108
6.10. Discussion	109
6.10.1. COX	109
6.10.2. LOX	109
6.11. Conclusion	110

CHAPTER SEVEN

Cytotoxicity evaluation of the crude extracts against Vero African green monkey kidney cell lines

7.0. Introduction	111
7.1. Materials and Methods	112
7.1.1. Preparation of plant extracts	112
7.1.2. Cytotoxicity assay against Vero cell	112
7.2. Results	113
7.3. Discussion	114
7.4. Conclusion	115

CHAPTER EIGHT

Motility modulation potential of *Bauhinia galpinii* and *Combretum vendae* phenolic-enriched leaf extracts on isolated rat ileum

8.0. Introduction	116
8.1. Drugs and reagents	117
8.2. Animal care	117
8.2.1. Isolated ileum preparation	118
8.3. Contractility test	118
8.3.1. Spasmogen assay	118
8.3.2. Spasmolytic assays	118
8.3.2.1. Effects on acetylcholine-induced contractility	118
8.3.2.2. Effects on serotonin-induced contractility	118
8.3.2.3. Effects on KCl-induced contractility	119
8.4. Data analysis	119
8.5. Results	119
8.5.1. Effect of <i>B. galpinii</i> crude extract on isolated rat ileum	119
8.5.2. Effect of <i>C. vendae</i> crude extract on isolated rat ileum	122
8.6. Discussion	123
8.7. Conclusion	126

CHAPTER NINE

Isolation and characterization of antimicrobial and antioxidant compounds from *Bauhinia galpinii* and *Combretum vendae*

9.0. Introduction	127
9.1.1. Column chromatography	128
9.1.2. Mass spectrometry	128
9.2. Materials and Methods	128
9.2.1. Preparation of plant extracts	128
9.2.2. Bioautography	128

9.2.3. Isolation of bioactive triterpenoids from <i>C. vendae</i>	129
9.2.4. Isolation of phenolic compounds from <i>C. vendae</i>	130
9.3. Isolation of compounds from <i>B. galpinii</i>	130
9.3.1. Isolation of bioactive triterpenoids from <i>B. galpinii</i>	130
9.3.2. Isolation of phenolic compounds from <i>B. galpinii</i>	130
9.4. Characterization of the isolated compounds	131
9.4.1. NMR spectroscopy	132
9.4.2. Mass spectrometry	132
9.4.3. Ultra-violet spectroscopy	132
9.5. Results	132
9.5.1. Identification of the chemical structures of isolated compounds from <i>C. vendae</i>	132
9.5.2. Antimicrobial activity of isolated compounds from <i>C. vendae</i>	135
9.5.3. Identification of the chemical structures of isolated compounds from <i>B. galpinii</i>	136
9.5.4. Antimicrobial activity of isolated compounds from <i>B. galpinii</i>	140
9.6. Discussion	140
9.6.1. Bioactive compounds from <i>C. vendae</i>	140
9.6.2. Bioactive compounds from <i>B. galpinii</i>	141
9.7. Conclusion	143
CHAPTER 10	
General conclusion and future prospects	
10. Introduction	144
10.1. Identification of diarrhoeal pathogenesis and medicinal plants used as therapeutic Agents	145
10.2. Antimicrobial evaluation of the extracts against infectious pathogens	145
10.3. Antioxidant evaluation of the extracts	145
10.4. Anti-inflammatory potential of the extracts	146
10.5. Toxicity risk of the extracts	146
10.6. Motility modulatory effects of <i>Bauhinia galpinii</i> and <i>Combretum vendae</i>	147
10.7. Isolation and characterisation of bioactive compounds	147
CHAPTER 11	
References	149

Abstract

Diarrhoea is one of the killer diseases resulting from the dehydration and loss of electrolytes through profuse and excessive excretion of loose stool. The pathoaetiologies include infections, intestinal inflammation, imbalanced intestinal oxidative homeostasis and altered motility. Treatment with oral rehydration therapy (ORT) is a key intervention especially in secretory diarrhoea as supportive therapy. Symptomatic and non-symptomatic therapies directed at treating the intestinal tissues are available. However, these conventional treatments are still not sufficient in curing diarrhoea due to their associated hazards such as the development and spread of drug-resistant pathogens, changes in normal intestinal bacteria flora and potential chronic toxicity. Therapies targeted at intestinal tissue include antimotility and antisecretory agents have adverse effects such as addictiveness, constipation and fatal ischaemic colitis. Many ethnopharmacological and ethnobotanical therapies for treating diarrhoea exist among different cultures. The aims of this study were to evaluate the biological activities of plant extracts against some diarrhoeal pathophysiologicals.

A literature search in English of published articles and books that discussed ethnobotanical uses of medicinal plants in southern Africa was conducted. A list of 230 medicinal plants used in South African traditional medicines for treating diarrhoea and associated complications was created. The list included family, genus, species, biological activities and bioactive isolates as well as the remedies for diarrhoea. Twenty seven species were selected to evaluate for antimicrobial, antioxidant and anti-inflammatory activities. Safety of the plants was determined by determining the cytotoxicity of the crude extracts against Vero African green monkey kidney cell lines using a standard method. Motility effects of *Bauhinia galpinii* (BGE) and *Combretum vendae* (CVE) were determined by modulation of the contractility process of the isolated rat ileum induced by spasmogens.

Phenolic compositions of the crude extract were determined using various standard methods and finally bioactivity guided isolation of antimicrobial and antioxidant compounds from BGE and CVE were carried out using open column chromatography. Identification and characterization of the isolated compounds was achieved by NMR, EI-MS and UV spectroscopy.

The non-polar fractions had good antimicrobial activities with MIC ranged between 19 – 1250 µg/ml while the polar fraction had moderate antimicrobial activities with MIC ranged between 39 - >2500 µg/ml. In general the non-polar fractions had a higher antimicrobial activity.

The crude extracts contained wide range phenolic compounds with a total phenolic (74.91±1.26 to 467.04±15.82 mg GAE/g plant material), and total flavonoids (11.27±3.37 to 176±5.96 mg EQ/g plant material). The antioxidant activities were concentrated and potentiated in the polar fractions. The non-polar fractions had poor antioxidant activities with EC₅₀ values ranging from 0.21±0.03 to 303.65±3.84 µg/ml for DPPH radical scavenging and 0.43±0.03 to 1709±91.44 µg/ml for ABTS radical scavenging.

The crude extracts had selective COX-1 inhibitory activities ranging between 41.70 to 84.61% and had no COX-2 inhibitory activity. All the extracts tested had 15-LOX inhibitory capacity with LC₅₀ values ranging between 0.86±0.27 and 111.44±37.28 µg/ml. The cytotoxicity results indicated a wide variation in toxic potential of the crude extracts with LC₅₀ values ranging from 3.51 to 741.90µg/ml.

The BGE extracts had dual activities as spasmolytic by stimulating the spontaneous contractility and also agonised contractions induced by spasmogens but it inhibited K^+ induced contraction. CVE had spasmodic activities through a multiple mechanisms inhibiting contractions induced by spasmogens and K^+ in a dose-dependent manner.

Several bioactive compounds were isolated from the *Combretum vendae* leaves, There were triterpenoids (ursol-12-en-28-oic acid, mixtures of corosolic acid and maslinic acid, and asiatic acid and arjunolic acid) as well as bibenzyls combretastatin B5-O-2'- β -D-glucopyranoside, combretastatin B1-O-2'- β -D-glucopyranoside and a flavonoid (apigenin)..

From *Bauhinia galpinii* the following bioactive compounds were isolated and characterized: β -3 ethoxy sitosterol, one new flavone (5, 7, 4', 5' tetrahydroxy-2'-methoxyflavone (isoetin 2'-methyl ether) or 5, 7, 2', 5' tetrahydroxy-4'-methoxyflavone (isoetin 4'-methyl ether)), 3, 5, 7, 3', 4'-pentahydroxyflavone and 3, 5, 7, 3', 4', 5'-hexahydroxyflavone, quercetin-3-O- β -galactopyranoside and myricetin-3-O- β -galactopyranoside

The extraction protocol used in this work potentiated the antimicrobial activities in the non-polar fractions while antioxidant activities were potentiated in the polar fractions. This indicated that using polar solvents as extractant for treating infectious diarrhoea may not be quite effective unless some other antidiarrhoeal mechanisms are involved. Therefore, mixture of organic solvent (ethanol) and water can be recommended for broad-based activity.

Bauhinia galpinii extracts had a dual- mechanism of action (prokinetic and relaxant) on gastro-intestinal motility, depending on the prevalent patho-physiological condition and *Combretum vendae* mediated spasmolytic effects on isolated rat ileum through multiple inhibitions of a wide range of contractile stimuli. Hence, the presence of multiple acting spasmolytic activities in the plant extract might be contributing towards its effectiveness in treating diarrhoea and abdominal spasm. The uses of these plants in traditional medicine need to be monitored closely because of the selective inhibition of COX-1 and its associated GIT injury, and the high toxicity potential of some of the extracts.

Further work evaluating the antidiarrhoea mechanisms, identification and isolation of bioactive compounds, sub-acute and acute toxicity of the plant extracts is recommended.

Key words: Antimicrobial, antioxidant, anti-inflammatory, diarrhoeal, antispasmodic, enteric nervous system, cytotoxicity.

List of Abbreviations

A

ABTS=2,2'-azino-bis-(3-ethylbenzothiazoline-6-sulfonic acid)

AMP=Antimicrobial peptides

B

BAB= *Bauhinia bowkeri*

BAG= *Bauhinia galpinii*

BAP= *Bauhinia petersiana*

BAV= *Bauhinia variegata*

BGE= *Bauhinia galpinii* extract

C

Ca²⁺= Calcium ion

Cl⁻= chloride ions

CNF-1= Cytotoxic necrotising factor 1

CNS= Central nervous system

COB= *Combretum bracteosum*

COP= *Combretum padoides*

COV= *Combretum vendae*

COX= Cyclooxygenase

COW= *Combretum woodii*

CVE= *Combretum vendae* extract

D

DAEC= diffusively adherent *Escherichia coli*

DNA

DPPH=2, 2-diphenyl-1-picrylhydrazyl

E

EAEC= Enteroaggregative *Escherichia coli*

EHEC= Enterohaemorrhagic *Escherichia coli*

EIEC= Enteroinvasive *Escherichia coli*

ENS= Enteric nervous system

EPEC= Enteropathogenic *Escherichia coli*

ETEC= Enterotoxigenic *Escherichia coli*

EUC=*Euclea crispa*

EUN= *Euclea natalensis*



F

FIC= *Ficus cratostoma*

FIG=*Ficus glumosa*

FRAP= Ferric reducing antioxidant capacity

G

GIT= Gastrointestinal tract

H

HIV/AIDS= Human immune deficiency virus/Acquired immune deficiency syndrome

HOCl= hypochlorite

HUB= Haemolytic uremic syndrome

I

IBS= Irritable bowel syndrome

IL= Interleukin

INC= *Indigofera cylindrica*

iNOS= inducible nitric oxide synthase

INT= p-iodonitrotetrazolium

L

LT= Heat labile enterotoxin

LTB= Leukotriene B

M

MDA= Malondialdehyde

MCP-1= Monocyte chemoattractant protein

MIC= Minimum inhibitory concentration

MPD= *Maytenus peduncularis*

MPR= *Maytenus procumbens*

MSE= *Maytenus senegalensis*

MUN= *Maytenus undata*

N

Na⁺= sodium ions

NAME= nitro

NH₂Cl= Ammonium chloride

NO= Nitric oxide

O

OH⁻ = Hydroxyl radical

ORT=Oral rehydration therapy

OZM= *Ozoroa mucronata*

OZP= *Ozoroa paniculosa*

P

PG= Prostaglandin

R

ROS= Reactive oxygen species

RNS= Reactive nitrogen species

S

SCB=*Schotia brachypetala*

SLE= *Searsia leptodictya*

SPD= *Searsia pendulina*

SPT= *Searsia pentheri*

ST= Heat stable enterotoxins

SYP= *Syzygium paniculatum*

T

TLC=Thin layer chromatography

TNF- α = Tumour necrosis factor- α

Trolox= 6-hydroxy-2, 5, 7, 8-tetrahydroxyl-chroman-2-carboxylic acid

U

UNICEF=United Nation Children Fund

W

WHO= World Health Organization

List of Figures

Chapter 2

Fig. 2.1. Classification of the diarrhoea and the stimulants	6
Fig. 2.2. Cytokines production network in the tissues	8
Fig. 2.3. Biosynthetic pathways for the eicosanoids	9
Fig. 2.4. Intestinal epithelial TJs as a physical barrier	10
Fig.2.5. The integrative pathophysiology and mechanism of diarrhoeal disease	13
Fig. 2.6. Lipid peroxidation chain reactions	14
Fig. 2.7. Chemical structures of the lipid peroxidation intermediates	14
Fig. 2.8. Mechanisms of antibiotic-induced diarrhoea	25
Fig. 2.9. Chemical structures of bioactive terpenoids against diarrhoeal mechanisms	31
Fig. 2.10. Chemical structures of bioactive alkaloids against diarrhoeal mechanisms	33
Fig. 2.11. Sub-classes of biologically important phenolic compounds	34
Fig. 2.12. Chemical structures of bioactive phenolics against diarrhoeal mechanisms	36

Chapter 3

Fig.3.1. Flow chart for the extraction, phytochemical analysis and fractionation of the crude extracts	43
Fig.3.2. TLC phytochemical profile of the crude extracts	48
Fig.3.3. TLC phytochemical profile of the hexane fractions	49
Fig.3.4. TLC phytochemical profile of the dichloromethane fraction	50
Fig.3.5. TLC phytochemical profile of the ethyl acetate fraction	51
Fig.3.6. Total phenolic and non-tannin constituents of the crude extract	52
Fig.3.7. Total tannin and condensed tannin constituents of the crude extracts	53
Fig.3.8. Proanthocyanidin and gallotannin constituents of the crude extract	54
Fig.3.9. Total flavonoid and flavonol constituents of the crude extract	56

Chapter 4

Fig.4.1. The classification of microbiological methods for biological detection	63
Fig.4.2. Bioautography of the hexane fractions against <i>S. aureus</i>	65
Fig.4.3. Bioautography of the dichloromethane fractions against <i>S. aureus</i>	66
Fig.4.4. Bioautography of hexane fractions of different plant species against <i>E. faecalis</i>	66
Fig.4.5. Bioautography of dichloromethane fractions of different plant species against <i>E. coli</i>	67
Fig.4.6. Bioautography of dichloromethane fractions of different plant species against <i>E. faecalis</i>	67
Fig.4.7. Bioautography of hexane of different plant species against <i>C. neoformans</i>	68
Fig.4.8. Bioautography of dichloromethane fractions against <i>C. neoformans</i>	68
Fig.4.9. Bioautography of hexane fractions against <i>A. fumigatus</i>	69
Fig.4.10. Bioautography of dichloromethane fractions against <i>A. fumigatus</i>	69
Fig.4.11. Bioautography of hexane fractions against <i>C. albicans</i>	70

Fig.4.12. Bioautography of dichloromethane fractions against <i>C. albicans</i>	70
Chapter 5	
Fig.5.1. Some deleterious reactions from the production of reactive free radicals in biological systems	80
Fig.5.2. TLC-DPPH profiles of the crude extracts of extracts of different plants	88
Fig.5.3. TLC-DPPH profile of the hexane fractions of different plants	89
Fig.5.4. TLC-DPPH profiles of the dichloromethane fractions of different plants	89
Fig.5.5. TLC-DPPH profiles of the ethyl acetate fractions of different plants	90
Chapter 6	
Fig.6.1: Roles of COX in the pathogenesis mechanism of NSAID-induced intestinal damage	103
Fig.6.2: Factors involved in the pathogenesis of indomethacin-induced small intestinal lesions	104
Fig.6.3: COX-1 inhibitory activity of some selected phenolic-enriched crude extracts	106
Chapter 8	
Fig.8.1.Schematic presentation of the contractility assay using isolated rat ileum	118
Fig.8.2.Stimulatory effects <i>B. galpinii</i> on spontaneous contraction of isolated rat ileum	119
Fig.8.3.Effects of 70% acetone crude leaf extract of <i>B. galpinii</i> on the acetylcholine cumulative concentration dependent-induced contraction in the absence and presence of atropine	119
Fig.8.4.Agonized effects of <i>B. galpinii</i> on serotonin-induced contraction of the isolated rat ileum	120
Fig.8.5.Relaxant effects of <i>B. galpinii</i> on KCl-induced contraction of the isolated rat ileum	120
Fig.8.6.Spasmolytic effects of 70% acetone crude leaf extract of <i>C. vendae</i> on acetylcholine-induced contraction of the isolated rat ileum	121
Fig.8.7.Spasmolytic effects of 70% acetone crude leaf extract of <i>C. vendae</i> on serotonin-induced contraction of the isolated rat ileum	121
Fig.8.8.Spasmolytic effect of the <i>C. vendae</i> on the depolarised KCl-induced isolated rat ileum contractions	122
Chapter 9	
Fig.9.1. Extraction, fractionation and isolation of bioactive compounds from the leaf extract of <i>Combretum vendae</i>	128
Fig.9.2. Extraction, fractionation and isolation of bioactive compounds from the leaf extract of <i>Bauhinia galpinii</i>	130
Fig. 9.3 Chemical structures isolated bioactive compounds from the leaf extract of <i>C. vendae</i>	134
Fig. 9.4 Chemical structures isolated bioactive compounds from the leaf extract of <i>B. galpinii</i>	138

List of Tables

Chapter 2

Table 2.1. The mechanism of action and symptoms of enteric pathogenic <i>E.coli</i>	17
Table 2.2. Neurotransmitters of ENS causing intestinal secretion in diarrhoea	27

Chapter 3

Table 3.1. Medicinal plants selected for the antidiarrhoeal investigations in this study	42
Table 3.2. The percentage yield of the crude extract and fractions (g/g dried plant material)	47

Chapter 4

Table 4.1. The minimum inhibitory concentration (MIC) of the crude extracts and fractions against bacterial strains tested	72
Table 4.2. The minimum inhibitory concentration (MIC) of the crude extracts and fractions against fungal strains tested	74

Chapter 5

Table 5.1. DPPH radical scavenging potential of the crude extracts and fractions expressed as EC ₅₀ (µg/ml)	91
Table 5.2. ABTS radical scavenging potential of the crude extracts and fractions expressed as EC ₅₀ (µg/ml)	92
Table 5.3. FRAP	94
Table 5.4. Hydroxyl radical scavenging potential of the crude extracts and fractions expressed as EC ₅₀ (µg/ml)	94
Table 5.5. Linoleic acid peroxidation inhibition expressed as LC ₅₀ (µg/ml)	95

Chapter 6

Table 6.1. Lipoxygenase inhibitory activity of the crude extracts	107
---	-----

Chapter 7

Table 7.1. The LD ₅₀ of the cytotoxicity assay of some medicinal plants used in South African traditional medicine to treat diarrhoea and related ailments	112
---	-----

Chapter 9

Table 9.1: NMR experiments commonly applied for natural product structural elucidation	126
Table 9.2: Minimum inhibitory concentration (µg/ml) of the isolated compounds from the leaf extract of <i>C. vendae</i>	135
Table 9.2: Minimum inhibitory concentration (µg/ml) of the isolated compounds from the leaf extract of <i>B. galpinii</i>	139

List of Appendix

	Page number
Appendix 2.1. Ethnobotanical and literature information of medicinal plant species used traditionally for treating diarrhoea in South Africa	172
Appendix 9.1. 1D and 2D NMR spectra data of Ursolic acid	197
Appendix 9.2. 1D and 2D NMR spectra data of mixture of corosolic acid and maslinic acid	198
Appendix 9.3. 1D and 2D NMR spectra data of mixture of asiatic acid and arjunolic acid	199
Appendix 9.4. 1D and 2D NMR spectra data of combretastatin B5-2'-O- glucopyranoside	200
Appendix 9.5. 1D and 2D NMR spectra data of combretastatin B1-2'-O- glucopyranoside	200
Appendix 9.6. 1D and 2D NMR spectra data of 3 β -ethoxy sitosterol	201
Appendix 9.7. 1D and 2D NMR spectra data of quercetin	201
Appendix 9.8. 1D and 2D NMR spectra data of myricetin	202
Appendix 9.9. 1D and 2D NMR spectra data of isoetin 2' methyl ether/ isoetin 4' methyl ether	202
Appendix 9.10. 1D and 2D NMR spectra data of quercetin-3-O- β -galactopyranoside	203
Appendix 9.11. 1D and 2D NMR spectra data of myricetin-3-O- β -galactopyranoside	203