

CHAPTER 3

3. HYPOTHESES AND OBJECTIVES

3.1. Hypotheses

3.1.1. Acidified water extracts prepared from marama bean seed coats and condensed tannin sorghum bran will have higher total phenolic content, flavonoid content, condensed tannin content, antioxidant activity, and a higher protective effect against erythrocyte haemolysis, LDL oxidation and oxidative DNA damage initiated by radicals compared to water extracts. This is because acidic condition during preparation of acidified water extracts will enhance extraction of free and esterified phenolic compounds and hydrolysis and release of bound phenolic compounds (Liyana-Pathirana & Shahidi, 2005; Li *et al.*, 2007). Phenolic compounds will exert antioxidant activity by scavenging free radicals and chelation of metal cations (Aherne & O'Brien, 2000).

3.1.2. Acidified water extracts prepared from marama bean seed coats and condensed tannin sorghum bran will have higher levels of free phenolic acids such as *p*-coumaric, ferulic and caffeic acids, and flavonoid aglycones compared to water extracts on analysis by HPLC-MS. The acids exist mostly in bound forms esterified to cell wall components (Manach *et al.*, 2004) and the flavonoids exist mostly as glycosides and upon pH treatment bound phenolic acids will be released as free phenolic acids (Liyana-Pathirana & Shahidi, 2005) and flavonoid glycosides hydrolysed to their aglycone forms (Careri *et al.*, 1998).

3.2. Objectives

3.2.1. To determine the effect of extraction under acidic condition on total phenolic, total flavonoid, condensed tannin contents and protein precipitation capacity of aqueous extracts prepared from marama bean seed coat and condensed tannin sorghum bran and to quantify and characterize their phenolic compounds using HPLC-MS.

3.2.2. To determine the effect of extraction under acidic condition on antioxidant activity of aqueous extracts prepared from marama bean seed coat and condensed tannin sorghum bran using the 2,2'-azinobis(3-ethylbenzothiazoline-6-sulphonic acid (ABTS⁺), 2,2-diphenyl-1-

picrylhydrazyl (DPPH[•]) scavenging assays and oxygen radical absorbancy capacity (ORAC) assay.

3.2.3. To determine the effect of aqueous extracts prepared from marama bean seed coats and condensed tannin sorghum bran on AAPH-induced oxidative DNA damage and AAPH-induced human erythrocyte haemolysis and copper-catalysed human LDL oxidation.

CHAPTER 4

4. RESEARCH

4.1. Characterization of phenolic acid and flavonoid compounds in aqueous extracts prepared from marama bean seed coats and condensed tannin sorghum bran

Part of this chapter has been published in the International Journal of Food Science and Technology¹ and accepted for publication in the Journal of Food Biochemistry².

¹Shelembe, J.S., Cromarty, D., Bester, M.J., Minnaar, A. and Duodu, K.G. (2012) Characterisation of phenolic acids, flavonoids, proanthocyanidins and antioxidant activity of water extracts from seed coats of marama bean (*Tylosema esculentum*) – an underutilised food legume. *International Journal of Food Science and Technology* 47, 648-655.

²Shelembe, J.S., Cromarty, D., Bester, M.J., Minnaar, A. and Duodu, K.G. (2012) Effect of acidic condition on phenolic composition and antioxidant potential of aqueous extracts from sorghum (*Sorghum bicolor*) bran. *Journal of Food Biochemistry* (in press)

4.1.1. Abstract

Marama bean is a legume plant growing wild in arid and semi arid regions of Southern Africa and sorghum is an important cereal crop for populations in arid and semi-arid regions of the world. Freeze-dried aqueous extracts with possible application as antioxidant functional food ingredients were prepared from marama bean seed coats and condensed tannin sorghum bran by extracting with water (water extract) or acidified water pH 2 (acidified water extract). The extracts were analyzed for total phenolic and flavonoid content using spectrophotometric methods and for phenolic acid and flavonoid compounds using HPLC-MS. Water and acidified water extracts from marama bean seed coats had 10 and 5 fold higher total phenolic and total flavonoid content, respectively than equivalent extracts from sorghum bran. Acidified water extract from marama bean seed coats had 36, 33% and 38% lower total phenolic content, flavonoid content and overall phenolic compound concentration, respectively, compared to water extracts. The reduction in phenolic content was probably as a result of co-precipitation of phenolic compounds with other polymeric species. In contrast, acidified water extracts from sorghum bran had 15% and 19% higher total phenolic content and overall phenolic compounds concentration, respectively, compared to water extracts, but no significant difference in total flavonoid content was found. Acidic condition enhanced extraction of free, esterified phenolic compounds and released bound phenolic compounds resulting in increased phenolic compound content. Phenolic acids in extracts from marama bean seed coats were benzoic acid derivatives and flavonoids were flavanols esterified to gallic acid. In extracts from sorghum, phenolic compounds were benzoic and cinnamic acid derivatives, phenolic aldehydes and flavonoids from the flavanone subclass.

Kew words: Marama bean seed coat; Sorghum bran; Phenolic compounds; Total phenolic content; Total flavonoid content; HPLC-MS.

4.1.2. Introduction

Nowadays there is considerable interest in dietary phenolics because of their presumed role in the prevention of various degenerative diseases associated with oxidative stress (Scalbert, Morand, Manach & Rémésy, 2002). Oxidative stress is implicated in the development of various chronic diseases including atherosclerotic cardiovascular disease, several cancers (Hodgson, Croft, Puddy, Mori & Beilin, 1996; Xu & Chang, 2007), neurodegenerative conditions (Chen & Blumberg, 2008) and inflammatory conditions (Erlejman, Jagggers, Fraga & Oteiza, 2008). It is initiated by reactive oxygen and nitrogen (ROS/RNS) species. These free radicals are continuously produced in the body and are essential in energy metabolism, detoxification, chemical signalling and immune function and are controlled by endogenous enzymes such as superoxide dismutase, glutathione peroxidase and catalase (Jacob & Burri, 1996; Dimitrios, 2006). However, failure of the antioxidant defence systems to scavenge and minimize the formation of free radicals as well as the overproduction of free radicals leads to oxidative damage of biomolecules such as DNA, lipids and proteins in the human body (Jacob & Burri, 1996; Baublis, Decker & Clydesdale, 2000; Kamath, Chandrashekar & Rajin, 2004). Phenolic compounds are non-nutritive components found in many food and medicinal plants and have various biological functions such as antioxidative, anti-inflammatory and antimutagenic activities (Suda, Ishikawa, Hatakeyama, Miyawaki, Kudo, Hirano, Ito, Yamakawa & Horiuchi, 2008). The antioxidative functions are mediated by their ability to scavenge free radicals and to chelate of transition metals such as copper and iron (Aherne & O'Brien, 2000; Scalbert *et al.*, 2002) which catalyze oxidation reactions.

The marama bean (*Tylosema* species) plant is drought-tolerant and has potential for cultivation as a food crop (Bower, Hertel, Oh & Storey, 1988). The bean is a potential source of value-added processed products similar to those from soy bean (Maruatona, Duodu & Minnaar, 2010). The seed coats of marama beans are not edible and are discarded as waste material during consumption/processing. A study has shown that organic extracts prepared from the seed coats of marama beans contain phenolic compounds with antioxidant activity. Total phenolic content of acidified methanol (1% HCl in methanol) extract prepared from marama bean seed coats was found to be 24.6 mg catechin equivalents/100 mg on a dry weight basis and phenolic compounds in the extract were phenolic acids and flavonoids (Chingwaru, Duodu, van Zyl, Schoeman, Majinda, Yeboah, Jackson, Kapewangolo, Kandawa-Shultz, Minnaar & Cencic, 2011).

Sorghum (*Sorghum bicolor* (L.) Moench) bran, a by-product of sorghum milling, is a potential source of natural antioxidant compounds with potential health benefits (Awika & Rooney, 2004). During milling, the grain is de-hulled to remove phenolic compounds which are concentrated in the pericarp or outer layers of the grain (bran) and this improves colour, reduces astringency and improves protein digestibility of products produced from the sorghum meal (Dlamini, Taylor & Rooney, 2007). Therefore the bran fraction is an ideal material source for the extraction of phenolic compounds. All sorghums contain phenolic acids and flavonoid compounds but only type II and III sorghums contain high molecular weight condensed tannins (Dykes & Rooney, 2006).

Phenolic compounds are commonly extracted with organic solvents. However, organic solvents may not be compatible with food because they have some degree of toxicity towards human health (Tsuda, Mizuno, Kawakishi & Osawa, 1995). Extraction with water might be a preferred option for pharmaceutical and food grade commercial processes (Oomah, Corbé & Balasubramanian, 2010). Water does not pollute the environment and does not require special management and it is cheaper compared to organic solvents (Llorach, Tomás-Barberán & Ferreres, 2004). Extraction under aqueous acidic condition (using acidified water) may help release bound phenolic compounds from complex carbohydrates and thus increasing yield of antioxidant phenolics (Liyana-Pathirana & Shahidi, 2005). Furthermore, acidification also simulates the pH condition of the stomach and can give an indication of the availability of phenolic compounds during the early stages of digestion in the stomach.

The objective of this study was to determine the effect of extraction under acidic condition on yield, total phenolic content, total flavonoid content and phenolic compound profile (phenolic acid and flavonoid compounds) of aqueous extracts from marama bean seed coats and condensed tannin sorghum bran.

4.1.3. Materials and methods

4.1.3.1. Materials

Marama beans from the 2008 season were obtained from Ghanzi province in Botswana. Condensed tannin sorghum (PAN 3860, a type III sorghum) was from the 2004 season and was cultivated in South Africa. Folin Ciocalteu reagent, sodium carbonate, 1 M hydrochloric acid solution, 1 M sodium hydroxide solution, 32% concentrated hydrochloric acid, HPLC

grade methanol, HPLC grade acetonitrile, ethyl acetate, aluminium chloride, sodium nitrate and sodium hydroxide were purchased from Merck Chemicals (Johannesburg, South Africa). Gallic acid, protocatechuic acid, (+)-catechin hydrate, *p*-hydroxybenzoic acid, vanillic acid, caffeic acid, *p*-coumaric acid, (-)-epicatechin-3-O-gallate, ferulic acid, taxifolin, naringen, hesperetin were purchased from Sigma-Aldrich (Johannesburg, South Africa).

4.1.3.2. Methods

4.1.3.2.1. Sample preparation

Marama beans were cracked using a DF sample cracker (WMC Metal Sheet Works, Tzaneen, South Africa) and the seed coats were separated from the cotyledons manually. The seed coats were then ground into a powder using a Waring blender, sieved through a 500 μm aperture test sieve and packed under vacuum. The powdered samples were stored at 4 °C until extraction.

Five kilograms of condensed tannin sorghum grain was hand cleaned and sorted to remove diseased, broken grains and foreign materials. The grain was decorticated using a Prairie Research Laboratory (PRL) type sorghum dehuller (Rural Industries Innovation Center, Kanye, Botswana) for 5 min to obtain a bran yield of approximately 7%. Bran was separated from de-hulled grain by sieving through a 1000 μm aperture test sieve. The bran was ground into a powder using a Waring blender to pass through a 500 μm test sieve. Powdered bran sample was packed under vacuum and stored at 4 °C until extraction.

4.1.3.2.2. Preparation of extracts

Aqueous extracts from marama bean seed coats and condensed tannin sorghum bran were prepared with water adjusted to pH 2 (acidified water extracts) or water only (water extracts) according to the methods of Baublis *et al.* (2000) and Liyana-Pathirana and Shahidi (2005) with some modifications. Acidified water extracts were prepared by mixing 10 g sample with 150 ml distilled water and stirring for 30 min using a magnetic stirrer. The pH was adjusted to 2 using 1 M HCl and then incubated in a shaking water bath at 37 °C for 30 min. The pH was then adjusted to 6 using 1 M NaOH and incubation continued in a shaking water bath at 37 °C for 30 min. Water extracts were prepared by mixing 10 g sample with 150 ml distilled water containing 0.3 g NaCl (equivalent to the amount produced by the reaction

between 1 M HCl and 1 M NaOH added in acidified water extract) and stirring for 30 min using a magnetic stirrer. Extraction was further carried out in a shaking water bath at ambient temperature for 60 min. Both water and acidified water extracts were centrifuged at 7500 g, 25 °C for 15 min using a model Rotanta 460R centrifuge (Andreas Hettich GmbH & Co.KG, Tuttlingen, Germany). The supernatant was decanted and freeze-dried. Extracts were weighed and stored at -20 °C in sealed zip-lock bags until analyzed.

4.1.3.2.3. Total phenolic content

Total phenolic contents of the extracts were determined using the Folin-Ciocalteu method (Singleton & Rossi, 1965). Freeze-dried marama bean seed coat (25 mg) and sorghum bran (75 mg) extracts were weighed into 25 ml volumetric flasks and made up to volume with distilled water. Catechin standard serial dilutions ranging from 0 to 1000 mg/L were prepared from a stock solution of catechin (1250 mg/L). Extract solution or catechin standard solution (0.5 ml) was added into a 50 ml volumetric flask containing 10 ml distilled water. Folin-Ciocalteu's reagent (2.5 ml) was added and the contents were mixed before 7.5 ml sodium carbonate solution (20 g/100 ml) was added within 1-8 minutes after the addition of Folin-Ciocalteu's reagent. The contents were mixed again and made-up to the mark with distilled water. Two hours after the addition of sodium carbonate solution the absorbance was measured at 760 nm using a T80+ UV-VIS spectrophotometer (PG Instruments, Wibtoft, UK) and data acquired with UVWin software version 5.0.5 (PG Instruments, Wibtoft, UK). Total phenolic content were reported as mg CE/g (mg catechin equivalents per gram) on dry sample weight basis.

4.1.3.2.4. Total flavonoid content

Total flavonoid content was determined as described by Ali-Farsi and Lee (2008) with some modification. Five milligrams of freeze-dried marama bean seed coat and condensed tannin sorghum bran extracts were dissolved in 2 ml and 1 ml distilled water, respectively, to a final concentration of 2.5 mg/ml and 5 mg/ml, respectively. (+)-Catechin standard serial dilutions ranging from 0–0.8 mg/ml were prepared from a 1.2 mg/ml stock solution. Extract or catechin standard solution (30 µl) was added into the wells of a 96 well plate. Then 20 µl of 36 mM sodium nitrate, 20 µl 94 mM aluminium chloride and 100 µl of 0.5 M sodium hydroxide solutions were added to each well. Absorbance was read at 450 nm on a 96 well

plate reader (Bio-Tek EL X800, Bio-tek Instruments Inc, Winooski, USA). A calibration curve was prepared from catechin standards and flavonoid content reported as mg catechin equivalents per gram of sample (mg CE/g) on dry weight basis.

4.1.3.2.5. Characterization of phenolic acids and flavonoids in aqueous extracts by HPLC-MS

Extracts were subjected to acid catalyzed hydrolysis and extracted with ethyl acetate according to the methods described by Hahn, Faubion and Rooney (1983) and Svensson, Sekwati-Manang, Lutz, Schieber and Ganzle (2010) with some modification. Freeze-dried marama bean seed coat (6 mg) and condensed tannin sorghum bran (20 mg) extracts were dissolved in 5 ml of 2 M HCl. The mixture was heated in a shaking water bath at 98 °C for 1 hour. The solution was allowed to cool and phenolic acid and flavonoid compounds were extracted twice with 5 ml ethyl acetate using a separation funnel. After partitioning, the ethyl acetate fractions were combined and then evaporated to dryness in a rotary vacuum evaporator model Rotavapor RE 120 (Büchi Labortechnik AG, Flawil, Switzerland) at 30 °C. The residues (after evaporation of ethyl acetate) from marama bean seed coat and condensed tannin sorghum bran were re-dissolved in 2 ml and 1 ml methanol, respectively, filtered through a 0.2 µm PTFE filter and stored at 4°C until analysis.

HPLC conditions: The HPLC system consisted of a Waters 1525 binary pump, Waters 2487 dual wavelength absorbance detector (Waters Associates, Milford, USA) and a YMC-Pack ODS AM-303 (250 x 4.6 mm i.d., 5 µm particle size) reverse phase column (Waters Associates, Milford, USA). The HPLC conditions were according to Kim, Hyun, Kim, Park, Kim, Kim, Lee, Chun & Chung (2007). Mobile phase A was 0.1% acetic acid in water and mobile phase B was 0.1% acetic acid in acetonitrile. Solvents were delivered in a linear gradient as follows: 8-10% B (2 min), 10-30% B (25 min), 30-90% B (23 min), 90-100% B (2 min), 100-8% B (2 min) and 8% B (7 min). Sample injection volume was 20 µl. Flow rate was maintained at 0.8 ml/min and total run time was 61 min. Phenolic compounds were detected at 280 nm and data acquired by Breeze system software (Waters Associates, Milford, USA). Calibration curves were prepared from phenolic standards serial dilutions and concentration of each compound in samples calculated and data was reported as µg/g of sample on dry weight basis.

HPLC/ESI-MS conditions: The system consisted of an Agilent 1100 series HPLC system with model G1312A binary pump, model G1322A degasser, model G1367A auto sampler, model G1316A column heater and model G1315B photo diode array detector (Agilent Technologies, Santa Clara, CA, USA). A YMC-Pack ODS AM-303 (250 x 4.6 mm i.d., 5 µm particle size) reverse phase column (Waters Associates, Milford, MA) was used. The HPLC system was coupled to a triple quadrupole / linear ion trap mass spectrometer, model 4000 Q TRAP LC/MS/MS system (AB SCIEX, Concord, Canada). Injection sample volume was 20 µl and solvent gradient was the same as previously outlined under HPLC conditions except that column temperature was set at 30 °C and detector wavelength at 190-600 nm. The operating conditions for the mass spectrometer were set as follows: The source temperature was maintained at 400 °C, ion spray voltage at -4000 V (negative mode). Nitrogen at a pressure of 35 psi was used as a nebulizer gas (GS1) and at 40 psi as a heating gas (GS2) and at 23 psi as a curtain gas (CUR). Declustering (DP) and entrance (EP) potential were set at -70 and -10 V, respectively. The mass of compounds was scanned from m/z 70 Da to m/z 1200 Da in 0.75 s and the data acquired and monitored by Analyst software, version 1.5 (Applied Biosystems / MDS Analytical Technologies Instruments, Concord, Canada).

4.1.3.3. Statistical analysis

Results from two independent experiments performed in duplicate were expressed as means ± standard deviations. Data were analyzed using one-way analysis of variance (ANOVA) and post hoc testing using Fishers LSD test for multiple comparisons of means. Significant differences between means were accepted at $p \leq 0.05$. Statistica 8.0 program (StatSoft, Inc, Tulsa, OK) was used for statistical data analysis.

4.1.4. Results and discussion

4.1.4.1. Yield, total phenolic and total flavonoid content of aqueous extracts

Table 4.1.1 shows that water extracts from marama bean seed coats had the highest yield amongst all the extracts. Extraction of marama bean seed coats under acidic condition (acidified water extract) resulted in a 20% significant reduction in yield compared to extraction with water (water extract). In contrast, extraction of sorghum bran under acidic condition resulted in a 7% significant increase in yield compared to extraction with water.

The total phenolic and flavonoid content of water extract from marama bean seed coats were the highest amongst all the extracts following the same trend as the yield. Extraction of marama bean seed coats under acidic condition resulted in 36% and 33% reduction in total phenolic content and flavonoid content, respectively compared to water extracts.

Table 4.1.1. Effect of extraction under acidic condition on yield, total phenolic and total flavonoid content of aqueous extracts from marama bean seed coats and condensed tannin sorghum bran

Sample	% Yield ¹	Total phenolic content (mg CE ² / g)	Total flavonoid content (mg CE/g)
<u>Marama bean seed coat</u>			
Water extract	13.4 ³ d ⁴ ± 0.16	62.1 d ± 1.3	22.2 c ± 0.9
Acidified water extract	10.7 c ± 0.44	39.5 c ± 2.3	14.9 b ± 1.2
<u>Sorghum bran</u>			
Water extract	9.7 a ± 0.05	5.8 a ± 0.1	4.3 a ± 0.2
Acidified water extract	10.4 b ± 0.05	6.7 b ± 0.1	4.7 a ± 0.2

¹ Grams of freeze dried aqueous extract/100 g of dry sample weight

²CE, Catechin equivalent

³Data are means ± standard deviations of four determinations, on dry weight basis

⁴Means followed by different letters in the same column are significantly different ($p \leq 0.05$)

In contrast, extraction of condensed tannin sorghum bran under acidic condition resulted in a 15% and 9% increase in total phenolic and flavonoid content, respectively. However the increase in flavonoid content was not significant ($p > 0.05$) when compared to water extracts.

Water extract from marama bean seed coats appeared opaque without coagulation before removal of insoluble solids by centrifugation (Fig. 4.1.1A). However, in acidified water

extracts a precipitate was observed after acid treatment to pH 2 before removal of insoluble solids by centrifugation (Fig. 4.1.1B). The precipitate appeared to be as a result of a coagulation/flocculation process, leaving a transparent brownish supernatant. The precipitate was likely to be an interpolymer complex formed between non-ionic polymers such as cellulose or other hydroxyl containing polymers and ionic polymers (Khutoryanskiy, Dubolazov & Mun, 2009) such as tannins due to their ionic behaviour in solution (Roussy, Chastellan, van Vooren & Guibal, 2005).




Figure 4.1.1. Appearance of aqueous extracts from marama bean seed coats and condensed tannin sorghum bran after treatment; A) Marama water extract; B) Marama acidified water extract; C) Sorghum water extract; D) Sorghum acidified water extract.

These complexes are formed under acidic conditions (below pH 3) due to the formation of hydrogen bonds between ionic polymers and proton-accepting non-ionic polymers (Khutoryanskiy *et al.*, 2009). Low molecular weight compounds (such as phenolic acids and flavonoids) may be incorporated into the interpolymer complex through reversible non-

covalent binding (Molyneux & Vekavakayanondha, 1986). Therefore the decrease in yield, total phenolic and flavonoid content of acidified water extracts from marama bean seed coat could be as a result of co-precipitation of phenolic compounds with interpolymer complexes initiated by change in pH.

Acidified water extracts from condensed tannin sorghum bran (Fig. 4.1.1D) did not show as much coagulation and precipitation, and the medium was slightly cloudier and more opaque compared to the water extract (Fig. 4.1.1C). Liyana-Pathirana & Shahidi (2005) reported a two fold increase in yield and 2–5 fold increase in total phenolic content of extracts from soft and hard wheat bran, as a result of extraction under acidic condition. According to the authors the increase was attributed to the release of phenolic compounds esterified to non-starch polysaccharides such as arabinoxylans under simulated acidic conditions. Extraction of phenolic compounds following acid or base hydrolysis has been shown to results in extracts with higher levels of phenolic compounds (Hahn *et al.*, 1983; Madhujith & Shahidi, 2009), which is attributed to release of bound phenolic compounds. Therefore the increase in yield, total phenolic and flavonoid content in acidified water extracts from sorghum bran could be as a result of a similar process. However, in acidified water extracts from marama bean seed coats bound phenolic compounds released by acid hydrolysis were possibly co-precipitated with interpolymer complexes at low pH resulting in lower yield, total phenolic content and total flavonoid content.

The total phenolic content of water and acidified water extracts from marama bean seed coat were 75% and 84%, respectively lower than that of acidified methanolic extracts prepared from marama bean seed coats (246 mg catechin equivalent/g, dry weight basis) (van Zyl, 2007). This is because water is a poor extraction solvent for phenolic compounds and it can also extract soluble polysaccharides and proteins compared to methanol. The solubility of phenolic compounds in water is very low compared to organic solvents such as methanol and acetone (Ali-Farsi & Lee, 2008).

Kobue-Lekalake, Taylor and de Kock (2007) reported that the total phenolic content of infusion from bran of the same sorghum cultivar used in this study (PAN 3860) prepared by boiling in water for 20 min was 48.6 g tannic acid equivalents per kg (4.9% (w/w)). These results were comparatively higher than that of water extracts and acidified water extracts as shown in Table 4.1.1 (0.58% and 0.67% (w/w), respectively). This could be due to enhanced

extraction of phenolic compounds by boiling which breaks down cell walls and thus releasing compounds into extraction solution. Awika, Rooney, Wu, Prior and Cisneros-Zevallos (2003) reported that the total phenolic content of aqueous acetone extracts from bran of Hi tannin and Sumac (SU99) sorghum were 55 and 66 mg gallic acid equivalent/g on dry weight basis (5.5 % (w/w) and 6.6 % (w/w) respectively). Sikwese and Duodu (2007) reported that the total phenolic content of 100% methanol extracts from bran of a condensed tannin sorghum variety (*Phatafuli*) was 101 mg tannic acid equivalent/g on dry weight basis (10.1% (w/w)). The comparatively lower total phenolic content in the water and acidified water extracts from the condensed tannin sorghum bran may be mainly due to water being a poor solvent for the extraction of phenolic compounds compared to organic solvents (Ali-Farsi & Lee, 2008). Nevertheless, the amount of phenolic compounds in the aqueous extracts is an indication of the bio-available fraction i.e. phenolic compounds that may have health promoting effects when added to the diet. It must be mentioned that it is generally difficult to compare total phenolic contents in literature directly due to the fact that there is no uniformity in the standards used.

Total phenolic contents of water and acidified water extracts from marama bean seed coat were 10 and 5 fold respectively, higher than equivalent extracts from condensed tannin sorghum bran. Also, the flavonoid contents were 6 and 3 fold respectively, higher than equivalent extracts from condensed tannin sorghum bran.

4.1.4.2. Characterization of phenolic acid and flavonoid compounds by HPLC-MS

The retention times, mass spectra and λ_{\max} of selected phenolic compound standards were determined under the same separation conditions as samples and the results are shown in Table 4.1.2.

4.1.4.2.1. Marama bean seed coats

The phenolic compound profiles of hydrolysed water and acidified water extracts from marama bean seed coats were similar. The HPLC and total ion current chromatograms are presented in Fig. 4.1.2 and the mass spectra data is presented in Fig. 4.1.3.

Table 4.1.2. Mass spectra and UV data for selected phenolic acid and flavonoid standard compounds

Peak No.	Compound	Molecular weight (g/mol)	t_R^* (min)	m/z [M-H] ⁻ (% intensity)	Fragment ions m/z MS ⁿ (% intensity)	HPLC-DAD (λ_{max} nm)
1	Gallic acid	170.12	7.1	169 (100)	125 (51)	216, 272
2	Protocatechuic acid	154.10	10.8	153 (100)	109 (55)	218, 260, 294
3	Catechin	290.29	14.1	289 (100)	-	208, 228, 278
4	<i>p</i> -Hydroxybenzoic acid	138.12	14.7	137 (100)	93 (53)	198, 208, 256
5	Vanillic acid	168.10	16.2	167 (100)	123 (27), 109 (13)	216, 260, 292
6	Caffeic acid	180.20	16.6	179 (100)	135 (61)	216, 324
7	<i>p</i> -Coumaric acid	164.20	21.4	163 (100)	119 (91)	212, 222, 310
8	(-)-Epicatechin 3-O-gallate	442.40	22.6	441 (100)	289 (15), 208 (36), 179 (25)	208, 324
9	Ferulic acid	194.19	23.0	193 (100)	178 (39), 149 (19), 134 (26)	216, 322
10	Taxifolin	304.25	24.1	303 (100)	285 (16)	202, 290
11	Naringenin	272.26	35.7	271 (100)	151 (7)	194, 288
12	Hesperetin	302.30	36.4	301 (100)	242 (8)	204, 288

t_R^* - Retention time

- No fragment ion for parent compound


The identity of the major phenolic acids and flavonoids are summarized in Table 4.1.3. Peak 1 with the most abundant $[M-H]^-$ molecular ion at m/z 169.8 (100) (Fig. 4.1.3A) corresponded to gallic acid (MW=170.12). The major fragment ion at m/z 125.6 (63) corresponded to decarboxylated gallic acid ion. The retention time, fragmentation pattern and UV λ_{max} were similar to that of gallic acid standard (Table 4.1.2). Gallic acid has been reported in alkaline hydrolysates of seed coats from mung beans, lentils, faba beans, pigeon and cowpeas (Sosulski & Dabrowski, 1984). It has also been reported to occur as a component of condensed tannins in legumes in the form of (-)-epicatechin 3-O-gallate as extender and terminal units (Dueñas, Hernández & Estrella, 2006) and procyanidin gallate in red lentils (Amarowicz, Estrella, Hernández, Dueñas, Agnieszka, Kosińska & Pegg, 2009).

However, gallic acid was not found in aqueous acetone extracts from the seed coats of black soy beans (Xu & Chang, 2008) and aqueous methanol extracts from the seed coats of fifteen Brazilian and Peruvian bean cultivars (Rannilla, Genovese & Lajolo, 2007). From these reports it would appear that gallic acid does not occur in free form in the seed coats but as part of proanthocyanidin structure, which would explain its absence in aqueous methanol and acetone extracts and its presence in the alkaline hydrolysates.


The compound eluting in peak 2 did not match any of the standards. The most abundant $[M-H]^-$ molecular ion at m/z 167.6 (100) (Fig. 4.1.3B) suggested that this compound could be homogentisic acid (MW = 168.14). The fragment ion at m/z 123.6 (100) corresponded to decarboxylated homogentisic acid $[M-H-44]^-$ molecular ion. Homogentisic acid has been reported in lentils (Zadernowski, Pierzynowska-Korniak, Ciepiewska & Forniak, 1992). The compound eluting in peak 3 with abundant $[M-H]^-$ molecular ion at m/z 137.6 (100) (Fig. 4.1.3C) corresponded to *p*-hydroxybenzoic acid (MW = 138.12). The retention time spectra matched that of *p*-hydroxybenzoic acid standard (Table 4.1.2). *p*-Hydroxybenzoic acid has been reported in the seed coats of black soybeans (Xu & Chang, 2008), lentils (Zadernowski *et al.*, 1992), mung beans, fababeans, pigeon pea, lima beans, chickpea and cowpeas (Sosulski & Dabrowski, 1984).

The compound eluting in peak 4 did not match any of the standard phenolic compounds. It had an abundant $[M-H]^-$ molecular ion at m/z 455.9 (100) (Fig. 4.1.3D) and fragment ion at

m/z 169.3 (10) and it was tentatively identified as methyl (epi)catechin-3-O-gallate (MW = 456).



(A)



(B)

Figure 4.1.2. HPLC chromatogram recorded at 280 nm (top) and total ion current (TIC) chromatogram (bottom) of phenolic compounds in A) water and B) acidified water extracts from marama bean seed coats. Peak identities are shown in Table 4.1.3.




Figure 4.1.3. Mass spectra of phenolic acid and flavonoid compounds in aqueous extracts from marama bean seed coats.

This compound has been previously identified in green tea (*Camelia sinensis*) by the same molecular ion m/z 455 (Bastos, Saldanha, Catharino, Sawaya, Cunha & Eberlin, 2007). However, the position of the methyl group could not be established by mass spectrometry so this compound could be 4'-O-methyl (epi)catechin-3-O-gallate or 3'-O-methyl (epi)catechin-3-O-gallate. Also the ion at m/z 239.7 (58) which appeared to be a major fragment ion could not be identified. It is likely that it was a molecular ion of another compound co-eluting with methyl (epi)catechin-3-O-gallate.

Peak 5 did not match any of the standard phenolic compounds. The mass spectra (Fig. 4.1.3E) showed an abundant $[M-H]^-$ molecular ion at m/z 439.9 (100) and fragment ion at m/z 169.3 (91). The molecular weight of this compounds was 16 Da (one oxygen) lower than that of methyl (epi)catechin-3-O-gallate and the compound was tentatively identified as methyl (epi)afzelechin-3-O-gallate (MW = 440).

In their work on identifying phenolic antioxidants Bastos *et al.* (2007) reported a similar molecular ion at m/z 439 for one of the phenolic compounds in the mass spectrum of ethanolic extracts from green tea. However, it could not be identified. Peak 6 which was the third largest peak had the same abundant $[M-H]^-$ molecular ion at m/z 439.9 (100) and fragment ion at m/z 169.2 (13) (Fig. 4.1.3F) as peak 5. This compound was tentatively identified as an isomer of methyl (epi)afzelechin-3-O-gallate.

The compound eluting in Peak 7 with abundant $[M-H]^-$ molecular ion at m/z 441.9 (100) (Fig. 4.1.3G) corresponded to (-)-epicatechin-3-O-gallate standard (MW=442) (Table 4.1.2). Fragmentation gave rise to m/z 289.6 (7) ion corresponding to epicatechin ion and m/z 169.6 (60) ion corresponding to gallic acid ion. Similar molecular and fragment ions have been observed for epicatechin-3-O-gallate from green tea (Miketova, Schram, Whitney, Kerns, Valcic, Timmermann & Volk, 1998) and oolong tea (Dou, Lee, Tzen & Lee, 2007). The compound eluting in peak 8 with abundant $[M-H]^-$ molecular ion at m/z 441.9 (100) (Fig.

4.1.3H) had the same molecular weight as (-)-epicatechin-3-O-gallate in peak 7, suggesting that it could be its isomer (+)-catechin-3-O-gallate. Fragmentation gave rise to m/z 289.6 (6) and m/z 169.6 (59) ions which corresponded to catechin and gallic acid ions, respectively.

Table 4.1.3. Phenolic acid and flavonoid compounds identified in aqueous extracts prepared from marama bean seed coats

Peak No	t_R^* min	m/z [M-H] ⁻ (% Intensity)	Fragment ion m/z (% Intensity)	HPLC-DAD λ_{max} (nm)	Compound identity
1	7.2	169.8 (100)	125.6 (63)	214, 270	Gallic acid
2	11.8	167.6 (100)	123.6 (88)	208	Homogentisic acid
3	13.9	137.6 (100)	-	208, 278	<i>p</i> -Hydroxybenzoic acid
4	14.6	455.9 (100),	239.7 (58), 169.4 (10)	214, 280	Methyl (epi)catechin-3-O-gallate
5	16.7	439.9 (100)	169.3 (91)	210, 280	Methyl (epi)afzelechin-3-O-gallate
6	20.5	439.9 (100)	169.2 (13)	208, 280	Methyl (epi)afzelechin-3-O-gallate
7	22.6	441.9 (100)	289.6 (7), 169.6 (59)	208, 278	(-)-Epicatechin-3-O-gallate
8	22.9	441.9 (100)	169.6 (48)	208, 278	(+)-Catechin-3-O-gallate

* t_R -retention time

- No fragment ion for the parent compound was observed

The concentration of phenolic acid and flavonoid compounds in the aqueous extracts from the seed coats of marama bean is shown in Table 4.1.4. Methyl (epi)afzelechin-3-O-gallate isomers were the major compounds in the extracts accounting for at least 40% of total phenolic acid and flavanoid compound content. Methyl (epi)catechin-3-O-gallate was the second major compound accounting for at least 28% and gallic acid was third accounting for 10%. The concentration of all the phenolic compounds in acidified water extracts was significantly higher than that of water extracts. Extraction under acidic condition resulted in a significant reduction in concentration of all the compounds. The percentage reduction ranged from 24% for gallic acid to 54% for *p*-hydroxybenzoic acid. The overall reduction in phenolic acid and flavonoid compounds as a result of extraction under acidic condition was 38% and this was similar to the 36% reduction in total phenolic content as determined with the Folin Ciocalteu method. As explained earlier, the reduction could be as a result of phenolic compounds co-precipitating with interpolymer complex triggered by low pH.

Table 4.1.4. Effect of extraction under acidic condition (pH 2) on phenolic acid and flavonoid compounds in aqueous extracts from marama bean seed coat¹

Peak No	Compound	Water extract μg/g	Acidified water extract μg/g
1	Gallic acid	1981.0 b ² ± 178.0	1510.2 a ± 115.2
2	Homogentisic acid ³	489.9 b ± 57.6	277.6 a ± 13.6
3	<i>p</i> -Hydroxybenzoic acid	1760.2 b ± 156.6	802.4 a ± 34.2
4	Methyl (epi)catechin-3-O-gallate ⁴	5839.9 b ± 424.4	3265.5 a ± 250.6
5, 6	Methyl (epi)afzelechin-3-O-gallate ⁴	8339.2 b ± 334.0	5464.7 a ± 502.6
7, 8	(Epi)catechin 3-O-gallate ⁴	1744.5 b ± 159.7	1133.5 a ± 80.9
	Total	20154.7 b ± 1067.5	12453.9 a ± 673.9

¹ Results are means ± standard deviation of four determinations from two independent experiments on dry weight basis

² Means with different letters in the same row are significantly different ($p \leq 0.05$)

³ Values expressed as protocatechuic acid equivalents

⁴ Values expressed as (-)-epicatechin-3-O-gallate equivalents



4.1.4.2.2. Condensed tannin sorghum bran

HPLC and total ion current chromatograms of phenolic compounds in water extracts and acidified water extracts from condensed tannin sorghums bran are shown in Fig. 4.1.4 and mass spectra data in Fig 4.1.5. The peak identities are shown in Tables 4.1.5. The profiles of the HPLC chromatograms and total ion current chromatograms for water and acidified water extracts were similar and eleven phenolic compounds were identified in the extracts.



Peak 1 with an abundant $[M-H]^-$ molecular ion at m/z 153.7 (100) (Fig. 4.1.5A, which corresponded to authentic protocatechuic acid standard (MW=154.10). On fragmentation it gave rise to $[M-H-44]^-$ ion at m/z 109.6 (31) as a result of decarboxylation. The retention time, mass spectra and λ_{max} were similar to that of protocatechuic acid authentic standard (Table 4.1.2).

Peak 2 did not match any of the authentic phenolic standards. The most abundant $[M-H]^-$ molecular ion at m/z 137.6 (100) (Fig. 4.1.5B) suggested that it could be protocatechualdehyde (MW = 138.12). Protocatechualdehyde has been reported in aqueous methanol extracts from sorghum grain (Svensson *et al.*, 2010). The compound eluting in peak 3 with abundant $[M-H]^-$ molecular ion at m/z 137.6 (100) (Fig. 4.1.5C) corresponded to *p*-hydroxybenzoic acid (MW = 138.12) and on decarboxylation resulted in $[M-H-44]^-$ fragment ion at m/z 93.4 (61). The retention time, mass spectrum, and λ_{max} matched that of *p*-hydroxybenzoic acid authentic standard (Table 4.1.2).

Compound in peak 4 did not match any of the phenolic standards. The abundant $[M-H]^-$ molecular ion at m/z 253.7 (100) (Fig. 4.1.5D) suggested that it could be caffeoylglycerol (MW = 254.08) which has been reported in aqueous methanol extract from sorghum grain (Svensson *et al.*, 2010). Traces of vanillic acid (MW=168.15) identified by $[M-H]^-$ molecular ion at m/z 167.6 appeared to co-elute with caffeoylglycerol at a retention time of approximately 15.9 min. Peak 5 with abundant $[M-H]^-$ molecular ion at m/z 179.6 (Fig. 4.1.5E) corresponded to caffeic acid standard (MW=180.2) (Table 4.1.2) and on decarboxylation gave rise to $[M-H-44]^-$ fragment ion at m/z 135.3 (60).



(A)



(B)

Figure 4.1.4. HPLC chromatogram recorded at 280 nm (top) and total ion current (TIC) chromatogram (bottom) of phenolic compound profile of A) water and B) acidified water extracts from condensed tannin sorghum bran.




Figure 4.1.5. Mass spectra of phenolic acid and aldehyde compounds in water extracts prepared from condensed tannin sorghum bran.

Peak 6 did not match any of the phenolic standards. The abundant $[M-H]^-$ molecular ion at m/z 121.6 (100) (Fig. 4.1.5F) suggested that it could be *p*-hydroxybenzaldehyde (MW= 122.12) which has been reported in methanolic extracts from sorghum grain (Hahn *et al.*, 1983) and ethanol extracts from aerial parts of sorghum (Sène, Gallet & Doré, 2001). Peak 7 with abundant $[M-H]^-$ molecular ion at m/z 153.5 (100) (Fig. 4.1.5G) did not match any of the authentic phenolic standards and it was tentatively identified as gentisic acid (MW = 154.14). Gentisic acid has been reported in acidified acetonitrile extracts from Korean sorghum varieties (Chung, Kim, Yeo, Kim, Seo & Moon, 2011), aqueous methanol extracts from Sudanese sorghum cultivars (Awadelkareem, Muralikrishna, EL Tinay & Mustafa, 2009) and in extracts from sorghum aerial parts (Sène *et al.*, 2001). Co-eluting with this compound was the $[M-H]^-$ molecular ion at m/z 163.6 (49) which corresponded to *p*-coumaric acid standard (MW = 164.2) (Table 4.1.2) and $[M-H-44]^-$ fragment ion at m/z 119 (31) was as result of decarboxylation of the parent compound.




Figure 4.1.6. Mass spectra of flavonoid compounds in hydrolysed water extracts from condensed tannin sorghum bran.

Table 4.1.5. Mass spectra and UV data of phenolic compounds identified in aqueous extracts from condensed tannin sorghum bran

Peak No.	t_R^* (min)	Compound Identity	m/z [M-H] ⁻ (% intensity)	Fragment ion m/z (% intensity)	HPLC-DAD λ_{max} (nm)
1	10.7	Protocatechuic acid	153.7 (100)	109 (31)	208, 260, 292
2	13.8	Protocatechualdehyde	137.6 (100)	-	208, 282
3	14.9	<i>p</i> -Hydroxybenzoic acid	137.5 (100)	93 (61)	208, 256
4	16.2	Caffeyolglycerol	253.7 (100)		208, 266, 288
5	16.5	Caffeic acid	179 (100)	135 (60), 121 (15)	210, 288, 322
6	18.7	<i>p</i> -Hydroxybenzaldehyde	121.6 (100)	-	210, 280
7	21.4	Gentisic acid	153.6 (100)	-	210, 292
8	23.0	Ferulic acid	193.6 (100)	-	212, 322
9	24.0	Taxifolin	303.8 (100)	285 (15)	204, 288
10	32.6	Eriodictyol	287.7 (100)	151 (14), 135 (7)	202, 288
11	35.4	Naringenin	271.6 (100)	151(11)	121, 288

* t_R , Retention time; -, No corresponding fragment ion identified

The compound eluting in Peak 8 with abundant [M-H]⁻ molecular ion at m/z 193.6 (100) (Fig. 4.1.5H) corresponded to ferulic acid standard (MW=194.18) (Table 4.1.2). The phenolic acids, protocatechuic acid, *p*-hydroxybenzoic acid, vanillic acid, caffeic acid, *p*-coumaric acid and ferulic acid have been previously identified in aqueous methanolic (Hahn *et al.*, 1983; Svensson *et al.*, 2010) and acidified acetonitrile (Chung *et al.*, 2011) extracts from sorghum grain as well as aerial parts of sorghum plant (Sène, Gallet & Doré, 2001).

The mass spectra of flavonoid compounds identified in sorghum bran aqueous extracts are shown in Fig. 4.1.6 and their identities and concentrations are shown in Tables 4.1.5 and

4.1.6, respectively. Peak 9 with abundant $[M-H]^-$ molecular ion at m/z 303.8 (100) and fragment ion at m/z 285 (15) (Fig. 4.1.6A) corresponded to taxifolin (MW = 304.25) standard (Table 4.1.2). The compound eluting in peak 10 did not match any of the authentic phenolic standards. Its abundant $[M-H]^-$ molecular ion at m/z 287.7 (100) (Fig. 4.1.6B) suggested that it could be eriodictyol (MW= 288.25). The compound eluting in peak 11 with abundant molecular ions at m/z 271.6 (100) and fragment ion at m/z 151.0 (11) (Fig. 4.1.6C) corresponded to naringenin authentic standard (MW=272.26) (Table 4.1.2). The three flavonoid compounds, taxifolin, eriodictyol and naringenin have been previously identified in organic extracts from sorghum grain (Gujer, Magnolato & Self, 1986; Svensson *et al.*, 2010).

Extraction of phenolic compounds from condensed tannin sorghum bran under acidic condition resulted in a significant increase in concentration of seven of the phenolic compounds (Table 4.1.6). The increase in concentration of phenolic acids ranged from 13% for caffeic acid to 36% for ferulic acid. Flavonoid content increased by 34, 41 and 48% for naringenin, eriodictyol and taxifolin, respectively.

The sum of all phenolic compounds identified in the acidified water extract was 19% higher than that of the water extract. This increase was similar to the 15% increase in total phenolic content as determined by the Folin Ciocalteu method. The increase could be as a result of enhanced extraction of free and esterified phenolic compounds and hydrolysis and release of bound phenolic compounds under acidic condition (Liyana-Pathirana & Shahidi, 2005).

The phenolic compound profiles of marama bean and condensed tannin sorghum bran were different. Three phenolic acids were identified in aqueous extracts from marama bean seed coats compared to six in sorghum bran aqueous extracts. The common phenolic acid in both marama bean seed coat and condensed tannin sorghum bran aqueous extracts was *p*-hydroxybenzoic acid. Sorghum bran aqueous extracts had two phenolic aldehydes and none were present in marama bean seed coat aqueous extracts.

Table 4.1.6. Effect of extraction under acidic condition on phenolic compounds in aqueous extracts from condensed tannin sorghum bran

Peak No.	t_R^* (min)	Compound Identity	Compound concentration ¹	
			Water extract ($\mu\text{g/g}$)	Acidified water extract ($\mu\text{g/g}$)
1	10.7	Protocatechuic acid	281.5 a ² \pm 12.8	280.2 a \pm 30.3
2	13.8	Protocatechualdehyde ³	178.9 a \pm 32.0	239.9 b \pm 30.0
3	14.9	<i>p</i> -Hydroxybenzoic acid	368.7 a \pm 40.9	384.2 a \pm 51.1
4	16.2	Caffeoylglycerol ⁴ / vanillic acid	78.1 a \pm 4.2	113.0 b \pm 14.7
5	16.5	Caffeic acid	147.9 a \pm 16.7	172.1 b \pm 5.5
6	18.7	<i>p</i> -Hydroxybenzaldehyde ⁵	49.2 a \pm 7.5	95.4 b \pm 6.5
7	21.4	Gentisic acid ⁴ / <i>p</i> -coumaric acid	6.4 a \pm 1.9	5.6 a \pm 0.6
8	23.0	Ferulic acid	10.9 a \pm 1.6	17.2 b \pm 1.8
9	24.0	Taxifolin	39.2 a \pm 2.4	68.5 b \pm 2.7
10	32.6	Eriodictyol ⁶	78.0 a \pm 9.5	134.2 b \pm 11.6
11	35.4	Naringenin	99.7 a \pm 9.4	152.5 b \pm 14.9
		Total	1338.6 a \pm 97.8	1663.4 b \pm 97.2

t_R^* - Retention time

¹Results are means \pm standard deviation of four determinations from two independent experiments on dry weight basis

²Means with different letters in the same row are significantly different ($p \leq 0.05$)

³Values expressed as protocatechuic acid equivalent

⁴Values expressed as caffeic acid equivalent

⁵Values expressed as *p*-hydroxybenzoic acid equivalent

⁶Values expressed as taxifolin equivalent

Flavonoid compounds were also different in marama bean seed coat and condensed tannin sorghum bran aqueous extracts. In marama bean seed coat extracts flavonoid compounds

were flavanols esterified with gallic acid while in sorghum these were flavanones. Taxifolin sometimes may be classified as a flavanonol (also called dihydroflavonol or 3-hydroxyflavanone) because of the presence of a 3-hydroxyl group in its chemical structure (Harbone, 1989; Naczki and Shahidi, 2004). The concentration of phenolic compounds in water and acidified water extracts from marama bean seed coats were 15 and 7 fold higher, respectively than equivalent extracts from condensed tannin sorghum bran. This trend was similar to the total phenolic content as determined by the Folin Ciocalteu method where total phenolic content of water and acidified water extracts from marama bean seed coat were 10 and 5 fold higher than equivalent extracts from condensed tannin sorghum bran.

The decrease in levels of phenolic compounds in acidified water extracts from marama bean seed coats but corresponding increase for extracts from condensed tannin sorghum bran suggests that there may be differences in the polymer compound matrix in marama bean seed coats and condensed tannin sorghum bran. Marama bean seed coats possibly contain both ionic and non-ionic polymers at relatively high concentration which interact at low pH through hydrogen bonds resulting in interpolymer complex precipitate (Khutoryanskiy *et al.*, 2009). Phenolic compounds of which some may have been released from bound forms by acidic condition (Liyana-Pathirana & Shahidi, 2005) co-precipitated with the interpolymer complex (Molyneux & Vekavakayanondha, 1986).

4.1.5. Conclusion

Aqueous extracts from marama bean (*Tylosema esculentum*) seed coats have significantly higher levels of total phenolic content, total flavonoid content and phenolic compound concentration than aqueous extracts from the bran of condensed tannin sorghum, variety PAN 3860. Extraction of marama bean seed coats under acidic condition results in reduction in phenolic content possibly due to co-precipitation of phenolic compounds with polymeric species. However, extraction of condensed tannin sorghum bran under acidic conditions results in an increase in phenolic content. Acidic conditions effect release of bound phenolic compounds and depending on the type of high molecular weight polymeric species present in the sample matrix it may result in co-precipitation of phenolic compounds due to interpolymer complex precipitation triggered by low pH. Sorghum bran aqueous extracts contain phenolic acids, phenolic aldehydes and flavonoid compounds while extracts from marama bean seed coats contain phenolic acid and flavonoid compounds at relatively higher

concentrations. The flavonoid compounds in aqueous extracts from sorghum bran are flavanones while in aqueous extracts from marama bean seed coats, they are flavanols esterified to gallic acid.

Marama bean (*Tylosema esculentum*) seed coats may be a better material source for aqueous extraction of phenolic compounds than bran of condensed tannin sorghum, variety PAN 3860 because of the higher phenolic content of the resultant extracts. These extracts from marama bean seed coats are likely to have higher antioxidant activity and potential health benefits than extracts from the condensed tannin sorghum bran in view of the higher phenolic content. Extraction of sorghum bran under acid conditions may be the preferred method because it increases the recovery of phenolic compounds. In contrast it is not recommended for marama bean seed coats because it causes co-precipitation of phenolic compounds with other polymeric species.

4.2. Characterization of proanthocyanidins in aqueous extracts from marama bean seed coats and condensed tannin sorghum bran

This chapter has been published in part in the International Journal of Food Science and Technology

Shelembe, J.S., Cromarty, D., Bester, M.J., Minnaar, A. and Duodu, K.G. (2011) Characterisation of phenolic acids, flavonoids, proanthocyanidins and antioxidant activity of water extracts from seed coats of marama bean (*Tylosema esculentum*) – an underutilised food legume. *International Journal of Food Science and Technology* 47, 648-655.

4.2.1. Abstract

Legumes with dark seed coats have been shown to contain condensed tannins which are important dietary antioxidants. However, there are no reports on condensed tannins in marama bean seed coats therefore the need for this investigation. Condensed tannin content of seed coats from marama bean and bran from condensed tannin sorghum was determined in methanol and aqueous extracts using the vanillin-HCl and butanol-HCl assays. The capacity of the extracts to precipitate proteins was determined using the protein precipitation capacity assay. The constitutive unit composition of proanthocyanidins in aqueous acetone and aqueous extracts was determined using the thiolysis degradation method and HPLC-MS to identify the compounds. Methanol extracts from marama bean seed coats had 3.7 fold higher condensed tannin content and 7.8 fold higher protein precipitation capacities than methanol extracts from condensed tannin sorghum bran. Aqueous extracts from marama also had higher condensed tannin content and higher protein precipitation capacity than equivalent extracts from condensed tannin sorghum bran. Extraction under acidic conditions resulted in a significant reduction in condensed tannin content of extracts from marama bean seed coats while it had no significant effect on extracts from condensed tannin sorghum bran. The reduction may be as a result of precipitation of interpolymer complexes formed through hydrogen bonding between tannins and cell wall polysaccharide triggered by low pH. The constitutive units of proanthocyanidins in marama seed coats were epigallocatechin-3-O-gallate and epicatechin-3-O-gallate occurring as major extension and terminal units and epigallocatechin and epicatechin as minor extension unit components. Proanthocyanidins in aqueous extracts from condensed tannin sorghum bran were composed of catechin and epicatechin. In conclusion, marama bean proanthocyanidins are predominantly prodelphinidins due to the presence of (epi)gallocatechin units while sorghum bran proanthocyanidins, due to the presence of (epi)catechin units are procyanidins.

Key words: Marama bean seed coat; Sorghum bran; Proanthocyanidin; Prodelphinidin; Procyanidin; Thiolysis

4.2.2. Introduction

Proanthocyanidins (condensed tannins) are widely distributed in the plant kingdom (Strumeyer & Malin, 1975) and they are widespread in many foods derived from plants (Hagerman, Riedl, Jones, Sovik, Ritchard, Hartfeld & Riechel, 1998; Serrano, Puupponen-Pimiä, Dauer, Aura & Saura-Calixto, 2009). Tannins are normally regarded as antinutrients because of their ability to precipitate proteins (Strumeyer & Malin, 1975), inhibit digestive enzymes and decrease absorption of minerals and vitamins (Kunyanga, Imungi, Okoth, Momanyi, Biesalski & Vadivel, 2011). However, proanthocyanidins have attracted increasing attention due to the rapidly increasing body of evidence associating this class of phenolic compounds with a wide range of health benefits (Lazarus, Adamson, Hammerstone & Schmitz, 1999). Proanthocyanidins are thought to be important dietary antioxidants (Hagerman *et al.*, 1998) and are the major phenolic constituents presumed to have a health-protective action (Guyot, Marnet & Drilleau, 2001).

Tannins are biologically active compounds that may have beneficial effects (Xu, Yuan & Chang, 2007). A diet supplemented with 1% (w/w) tannins from young persimmon fruit promoted faecal bile acid excretion in rats suggesting that tannins can be used clinically to reduce hypercholesterolemia (Matsumoto, Kadowaki, Ozaki, Takenaka, Ono, Yokoyama & Gato, 2011) often associated with cardiovascular disease. Condensed tannins were reported to inhibit α -amylase and α -glucosidase enzymatic activity thereby having the ability to delay intestinal glucose absorption and the onset of insulin-dependent diabetes mellitus (Kunyanga *et al.*, 2011). Procyranidins (dimer–hexamer) were shown to inhibit liposome oxidation in a polymer length dependent manner (Verstraeten, Keen, Schmitz, Fraga & Oteiza, 2003). Condensed tannins and flavonoid compounds from legumes (lentils, black beans, red kidney beans and pinto beans) were reported inhibit *in vitro* copper-catalysed LDL oxidation (Xu *et al.*, 2007) where LDL oxidation is a contributing factor in the development of cardiovascular disease.

Purification of condensed tannins is an important step in their characterization. Solid phase extraction is the most commonly used method for the purification of condensed tannins in plant extracts (Strumeyer & Malin, 1975; Hagerman & Butler, 1980; Prieur, Rigaud, Cheynier & Moutounet, 1994; Sun, Leandro, Ricardo da Silva & Spranger, 1998a; Prior, Lazarus, Cao, Muccitelli & Hammerstone, 2001; Dueñas, Sun, Hernández, Estrella & Spranger, 2003).

In the purified fractions constitutive unit composition and mean degree of polymerization of proanthocyanidin polymers have been determined by reverse-phase HPLC analysis of acid catalyzed degradation products. Acid catalyzed degradation of proanthocyanidins is carried out under mild conditions in the presence of a nucleophilic reagent such as toluene- α -thiol (benzyl mercaptan) (Prieur *et al.*, 1994; Guyot, Marnet, Laraba, Sanoner & Drilleau, 1998; Fu, Loo, Chia & Huang, 2007). This is an important analysis because it distinguishes between extension and terminal units in the polymer (Prieur *et al.*, 1994; Guyot *et al.*, 1998). Extension units are captured by benzyl mercaptan to form benzylthioether derivatives and terminal units are released as free flavan-3-ols (Prieur *et al.*, 1994) or flavan-3-O-gallates units (Yousef, Grace, Cheng, Belolipov, Raskin & Lila, 2006). HPLC coupled to mass spectrometry, with electrospray interface (ESI) has proven to be extremely useful for peak assignment and characterization of individual phenolic compounds (Weisz, Kammerer & Carle, 2009).

Marama beans are gathered by hand from the wild in arid regions of Southern Africa and the cotyledons are utilized as food (Amarteifio & Moholo, 1998) and the seed coats discarded. The seed coats contain phenolic compounds (Chingwaru, Duodu, van Zyl, Schoeman, Majinda, Yeboah, Jackson, Kapewangolo, Kandawa-Shultz, Minnaar & Cencic, 2011). However nothing is known about proanthocyanidins and their structural features in marama bean seed coats. Condensed tannins in other legumes are predominantly procyanidins consisting of catechin and epicatechin subunits, located in the seed coat (Dueñas, Hernández & Estrella, 2006). However, small amounts of propelargonidin in common bean varieties (Beninger, Gu, Prior, Junk, Vanderberg & Bett, 2005) and prodelphinidins in lentils (Dueñas *et al.*, 2003) have also been reported.

The aim of this study was to characterize and compare proanthocyanidins or condensed tannins in aqueous extracts prepared from seed coats of marama beans and bran of condensed tannin sorghum using high performance liquid chromatography and mass spectrometry.

4.2.3. Materials and methods

4.2.3.1. Materials

Chemicals. (+)-catechin, (-)-epicatechin, (-)-gallocatechin, (-)-epigallocatechin, (-)-epicatechin gallate, (-)-epigallocatechin gallate, bovine serum albumin, were purchased from

Sigma-Aldrich (Johannesburg, South Africa). Ethyl acetate, diethyl ether, benzyl mercaptan (toluene- α -thiol), HPLC grade methanol, HPLC grade acetonitrile, 32% hydrochloric acid, glacial acetic acid (analytical grade reagent), 1-butanol, ferric ammonium sulphate, vanillin, sodium chloride, sodium dodecyl sulphate, triethanolamine, ferric chloride, acetone, sodium hydroxide were purchased from Merck Chemicals (Johannesburg, South Africa).

4.2.3.2. Methods

4.2.3.2.1. Sample preparation

Marama bean seed coat and condensed tannin sorghum bran powders were prepared as described in section 4.1.3.2.1. Freeze-dried aqueous extracts (water extracts and acidified water extracts) from marama bean seed coats and condensed tannin sorghum bran powders were prepared as described in section 4.1.3.2.2. Aqueous acetone extracts were also prepared to compare with aqueous extracts because condensed tannins are usually extracted with aqueous acetone for characterization (Lazarus, Adamson, Hammerstone & Schmitz, 1999; Prior *et al.*, 2001; Gu, Kelm, Hammerstone, Beecher, Cunningham, Vannozzi, & Prior, 2002; Gu, Kelm, Hammerstone, Beecher, Holden, Haytowitz & Prior, 2003b).

4.2.3.2.2. Determination of condensed tannin content by vanillin-HCl method

The vanillin-HCl assay (Price, Van Scoyoc & Butler, 1978) was used for the determination of condensed tannin content of methanol extracts and freeze-dried aqueous extracts prepared from marama bean seed coat and condensed tannin sorghum bran. Methanol is the commonly used solvent in this assay (Price, Van Scoyoc & Butler, 1978). The assay is based on the condensation of the aromatic aldehyde vanillin with monomeric flavanols and their oligomers to form a red adduct with absorbance maxima at 500 nm (Sarkar and Howarth, 1976; Dykes & Rooney, 2006). Methanol extracts were prepared by extracting 0.2 g marama bean seed coat or condensed tannin sorghum bran powder twice with 10 ml of 100% methanol. After the addition of each 10 ml portion of methanol, the mixture was shaken for 20 min using an orbital shaker model POS 300 (Grant Instruments Ltd, Shepreth, UK) and then centrifuged at 4500 rpm (5200 x g) for 10 min at 25 °C using a model Rotanta 460R centrifuge (Andreas Hettich GmbH & Co. KG, Tuttlingen, Germany) and the supernatants were decanted and combined. One millilitre of the methanol extract from marama bean seed coat and condensed tannin sorghum bran was diluted with 24 ml and 5 ml of 100% methanol,

respectively for analysis. Serial standard solutions of (+)-catechin were prepared in 100% methanol in a concentration range of 0 – 1000 mg/L. Vanillin reagent was prepared on the day required by mixing equal volumes of 1% vanillin in methanol and 8% concentrated HCl in methanol. Freeze-dried aqueous extracts from marama bean seed coat (35 mg) and condensed tannin sorghum bran (50 mg) were weighed into 100 ml and 25 ml volumetric flasks, respectively and made up to volume with 100% methanol to give a concentration of 0.35 and 2 mg/ml, respectively. Five millilitres of the vanillin-HCl reagent was added to 1 ml aliquot of sample or catechin standard solution. Sample blanks (without vanillin) were prepared by adding 5 ml of 4% HCl in methanol to 1 ml extract solutions. Absorbance was measured at 500 nm after 20 min from the time of addition of vanillin-HCl reagent using a T80+ UV-VIS spectrophotometer and UVWin software version 5.0.5 (PG Instruments, Wibtsoft, UK). Blank readings were subtracted from sample absorbance readings. Results were reported as mg catechin equivalent/g (mg CE/g) of dry sample weight.

4.2.3.2.3. Determination of condensed tannin content by Butanol-HCl assay

Condensed tannin content of methanol extracts and freeze-dried aqueous extracts was also determined with the butanol-HCl assay. The assay was performed as described by Porter, Liana, Hrstich and Chan (1986) with some modifications. In this assay proanthocyanidins are converted to anthocyanidins through an autoxidation reaction catalysed by Fe^{3+} following acid-catalysed cleavage of the interflavanoid bonds (Porter *et al.*, 1986). The anthocyanidins formed have a red colour and a maximum absorbance wavelength at around 550 nm (Sun *et al.*, 1998b). Methanol extracts were prepared by extracting 0.2 g marama bean seed coat or condensed tannin sorghum bran powder with 10 ml of 100% methanol. Mixture was shaken for 20 min on an orbital shaker model POS 300 and then centrifuged at 4500 rpm (5200 x g) for 10 min at 25 °C using a model Rotanta 460R centrifuge (Andreas Hettich GmbH & Co. KG, Tuttlingen, Germany) and supernatant decanted. Five millilitre of supernatant was diluted with 20 ml methanol for analysis. Freeze-dried extracts (25 mg) were weighed into centrifuge tubes and 25 ml methanol was added to each tube. The tubes were shaken at 300 rpm for 20 min on an orbital shaker model POS 300. Samples were then centrifuged at 4500 rpm (5200 x g) for 10 min at 25 °C using model Rotanta 460 R centrifuge (Andreas Hettich GmbH & Co., Tuttlingen, Germany). One millilitre of sample supernatant, 0.2 ml ferric reagent (2% (w/v) $NH_4Fe(SO_4)_2 \cdot 12H_2O$ in 2 M HCl) and 6 ml of 5% (v/v) 32% concentrated HCl in 1-butanol were added into glass test tube. The test tube was covered

with a glass marble, heated in a boiling water bath for 60 min, cooled and absorbance measured at 550 nm using a T80+ UV-VIS spectrophotometer and UVWin software version 5.0.5 (PG Instruments, Wibtoft, UK). Results were reported as absorbance units per gram (Abs_{550}/g) of sample on dry weight basis.

4.2.3.2.4. Determination of protein precipitation capacity

The protein precipitation capacity assay (Hagerman & Butler, 1978) was used to evaluate the ability of tannins in methanol extracts and aqueous extracts from marama bean seed coats and condensed tannin sorghum bran to precipitate proteins. This assay is based on the capacity of tannins to interact with proteins (added in excess) to form a tannin-protein complex precipitate (Hagerman & Butler, 1978). The tannin-protein precipitate is then dissolved in sodium dodecyl sulphate-triethanolamine solution and ferric ions (added as ferric chloride) react with tannins in the complex in the alkaline solution to form a violet complex with maximum absorbance at 510 nm. Methanol extracts were prepared by extracting 0.2 g marama bean seed coat or condensed tannin sorghum bran powder twice with 10 ml methanol. After the addition of each 10 ml portion of methanol, the mixture was shaken for 20 min on an orbital shaker model POS 300 and then centrifuged at 4500 rpm (5200 x g) for 10 min at 25 °C using a model Rotanta 460R centrifuge and the supernatant was then decanted. Supernatants were combined. One millilitre of the methanol extract from marama bean seed coat was diluted with 4 ml of methanol and extract from sorghum was analyzed without dilution. Freeze-dried aqueous extract (25 mg) from marama bean seed coat and condensed tannin sorghum bran were dissolved in 20 ml and 2 ml distilled water to give a concentration of 0.25 and 2.5 mg/ml, respectively. Two millilitres of bovine serum albumin (1 mg/ml) in 0.2 M acetate buffer solution, pH 5 containing 0.17 M sodium chloride and 1 ml of methanol or aqueous extract solution were added into the centrifuge tube, mixed and allowed to stand for 15 min. The tubes were centrifuged at 5000 g for 15 min at 25 °C and the pellet carefully washed with 2 ml of 0.2 M acetate buffer solution pH 5. The pellet was then dissolved in 4 ml of 1% (w/v) sodium dodecyl sulphate-5% (w/v) triethanolamine solution. One millilitre of 0.01 M ferric chloride reagent in 0.01 M HCl solution was added and absorbance measured at 510 nm after 20 min using T80+ UV-VIS spectrophotometer and UVWin software version 5.0.5. Results were reported as absorbance units at 510 nm per gram (Abs_{510}/g) on sample dry weight basis.

4.2.3.2.5. Identification of terminal and extension units in proanthocyanidins by HPLC-MS

Proanthocyanidins in aqueous acetone extracts and aqueous extracts were subjected to acid catalysed thiolysis degradation in the presence of benzyl mercaptan nucleophile under mild conditions to preserve flavan-3-O-gallate units and the products analysed by HPLC-MS. (Gu, Hammerstone, Zhang, Beecher, Holden, Howtowitz & Prior, 2003a).

4.2.3.2.5.1. Fractionation of proanthocyanidins using Sep-Pak C₁₈ cartridges

Proanthocyanidins in aqueous acetone extracts and aqueous extracts from marama bean seed coat and condensed tannin sorghum bran were fractionated into monomer (FI), oligomer (FII) and polymer (FIII) fractions using Sep Pak C₁₈ cartridges according to the method of Sun *et al.* (1998a). Aqueous acetone extracts were prepared as described by Gu *et al.* (2002) with some modification. One gram of marama bean seed coat or condensed tannin sorghum bran powder was extracted with 10 ml acetone/water/acetic acid (70:29.5:0.5, v/v/v) solution by vortexing for 30 sec and shaking for 50 min on an orbital shaker model POS 300. The mixture was centrifuged at 3500 rpm (3150 x g) for 15 min at 25 °C using a model Rotanta 460R centrifuge (Andreas Hettich GmbH & Co. KG, Tuttlingen, Germany). Acetone in six millilitres of supernatant was evaporated off at 25 °C under vacuum (200 tor) using a vacuum rotary evaporator model Rotavapor RE 120. Residue was made up to 6 ml with distilled water adjusted to pH 7 with 1 M NaOH. Fifty milligrams of freeze-dried extracts was dissolved in 6 ml distilled water adjusted to pH 7.0 and then sonicated for 5 sec. Samples were subjected to solid phase extraction by passing through two preconditioned neutral Sep Pak C₁₈ (plus short, 0.7 ml, 360 mg) and SepPak tC₁₈ (plus short, 0.8 ml, 400 mg) cartridges (Waters Associates, Milford, USA) connected in series under vacuum. The inferior C₁₈ was at the top and the superior tC₁₈ at the bottom. Cartridges were activated with 6 ml methanol (Tian, Nakamura, Cui & Kayahara, 2005; Walsh, Haak, Bohn, Tian, Schwartz & Failla, 2007) and reconditioned with 12 ml distilled water adjusted to pH 7 with 0.1 M NaOH solution. Phenolic acids and sugars (Fraction 0) were eluted with 10 ml distilled water adjusted to pH 7. Mixture of monomer and oligomer proanthocyanidins (Fraction I + II) was eluted with 15 ml ethyl acetate. Polymeric proanthocyanidins (Fraction III) were eluted with 5 ml methanol. The ethyl acetate solution was evaporated to dryness under vacuum (200 tor) at 25 °C using a rotary vacuum evaporator model Rotavapor RE 120 (Büchi Labortechnik AG, Flawil, Switzerland) and residue re-dissolved in 2 ml distilled water and re-deposited on

to the same cartridges after preconditioning with 12 ml distilled water adjusted to pH 7. Monomers (Fraction I) were eluted with 15 ml diethyl ether. Oligomeric proanthocyanidins (Fraction FII) were eluted with 5 ml methanol.

4.2.3.2.5.2. Thiolysis degradation of proanthocyanidins

Thiolysis degradation of oligomeric (FII) and polymeric (FIII) fractions was carried out according to the method of Prieur *et al.* (1994) modified by Gu *et al.* (2003a). A 50 μ l aliquot of oligomeric or polymeric fraction, 50 μ l of methanol acidified with concentrated HCl (3.3% v/v) and 100 μ l of toluene- α -thiol (5% v/v in methanol) were added into safety lock eppendorf tube. The tube was sealed and heated at 40 °C for 30 min in a water bath and then allowed to stand at room temperature for 10 h to complete the reaction. Reaction mixtures were kept at -20 °C until analysis.

4.2.3.2.5.3. Determination of mean degree of polymerization

Separation of oligomeric (FII) and polymeric (FIII) thiolysis degradation products in the reaction mixtures was carried out according to Gu *et al.* (2002) with some modification. Briefly, 20 μ l reaction mixture was injected into an HPLC system (Water Associates, Milford, USA) consisting of a model 1525 binary pump, model 2487 dual wavelength absorbance detector and a YMC-Pack ODS AM-303 (250 x 4.6 mm i.d., 5 μ m particle size) reverse phase column (Water Associates, Milford, USA). Solvent system consisted of mobile phase A (2% acetic acid in water, v/v) and B (methanol) delivered in a linear gradient of B from 15–80% in 45 min, 80–15% in 10 min and 15% for 10 min. Flow rate was 0.5 ml/min. The detector was set at 280 nm and the chromatograms acquired and monitored by Waters Breeze system software, version 3.30 SPA (Water Associates, Milford, USA). The mean degree of polymerization was calculated from peak area using the following equation adapted from Gu *et al.*, (2002).

$$\text{mDP} = \left\{ \frac{\text{total area of benzylthioether derivatives}}{\text{total area of flavan-3-ol units}} \right\} + 1$$

4.2.3.2.5.4. Identification of thiolysis degradation products by HPLC-MS

Thiolysis degradation products were analyzed by HPLC-MS to confirm their identity according to the method described by Gu *et al.* (2002) with some modification. The chromatographic system used consisted of an Agilent 1100 series HPLC system (Agilent Technologies, Santa Clara, CA) with model G1312A binary pump, model G1322A degasser, model G1367A auto sampler, model G1316A column heater and model G1315B photo diode array detector, reverse phase column (YMC-Pack ODS AM-303, 250 x 4.6 mm i.d., 5 μ m particle size), (Water Associates, Milford, USA) and a triple quadruple / linear ion trap mass spectrometer, model 4000 Q TRAP LC/MS/MS (AB SCIEX, Concord, Canada). Solvent system consisted of mobile phase A (2% acetic acid in water) and mobile phase B (methanol) delivered at 0.5 ml/min in a linear gradient of B from 15 – 80% in 45 min, 80% – 15% in 10 min and 15% for 10 min. Injection volume was 20 μ l, column temperature was 40 °C and detector wavelength was set at 275-285 nm. The operating conditions for the mass spectrometer were as follows: Source temperature was maintained at 400 °C, spray voltage at -4000 V (negative mode). Nitrogen at a pressure of 35 psi was used as a nebulizer gas (GS1) and at 40 psi as heating gas (GS2) and at 23 psi as curtain gas (CUR). Declustering (DP) and entrance potential (EP) were set at -60 V and -10 V, respectively. The mass of compounds was scanned from m/z 70 to m/z 1200 and chromatograms acquired and monitored by Analyst software, version 1.5 (Applied Biosystems and MDS Analytical Technologies, Concord, Canada)

4.2.3.3. Statistical analysis

Results from two independent experiments performed in duplicate were expressed as means \pm standard deviations. Data were analyzed using t-test to compare two means and one-way analysis of variance (ANOVA) and post hoc testing using Fishers LSD test for multiple comparisons of means. Statistica 8.0 program (StatSoft Inc, Tulsa, USA) was used for statistical data analysis and significance was accepted at $p \leq 0.05$.

4.2.4. Results and discussion

4.2.4.1. Condensed tannin content and protein precipitation capacity of methanolic extracts

The condensed tannin content and protein precipitation capacity of methanolic extract from marama bean seed coat were higher than that of methanolic extract from condensed tannin sorghum bran (Table 4.2.1). Condensed tannin content of marama bean seed coat determined with the vanillin-HCl method was 3.7 fold higher than that of condensed tannin sorghum bran. However the protein precipitation capacity of methanolic extract from marama bean seed coats extract was 7.8 fold higher than that of methanolic extract from condensed tannin sorghum bran which may be an indication that condensed tannins in marama bean seed coats have a higher affinity for proteins compared to condensed tannins from sorghum bran.

Table 4.2.1. Condensed tannin content and protein precipitation capacity of methanol extracts from marama bean seed coat and condensed tannin sorghum bran

Extract	Vanillin-HCl assay (mg CE ¹ /g)	Butanol-HCl assay (Abs ₅₅₀ units/g)	Protein precipitation capacity (Abs ₅₁₀ units/g)
Marama seed coat	1226.4 ² b ³ ± 16.3	198.6b ± 15.8	226.6 b ± 13.9
Sorghum bran	331.1a ± 9.6	119.3a ± 5.0	29.1a ± 1.1

¹CE, Catechin equivalents

²Data are means ± standard deviations of six determinations, on dry weight basis

³Means followed by different letters in the same column are significantly different ($p \leq 0.05$)

Compared to other legumes the condensed tannin content of the methanol extracts from marama bean seed coats were higher than levels reported in methanol extracts from seed coats of dark coloured common beans which ranged from 216 to 449 mg CE/g on wet weight basis depending on cultivar (Ranilla, Genovesse & Lajolo, 2007) and seed coats of black soybean which was approximately 45 mg CE/g of seed coat (Xu & Chang, 2008).

Condensed tannin content of sorghum bran was comparable to that reported in methanol extracts from *Phatafuli* (a condensed tannin sorghum cultivar from Malawi) sorghum bran which was 451 mg CE/g, dry weight (Sikwese & Duodu, 2007). However, it was higher than

that reported for 70% acetone extract from sorghum variety Sumac bran which was approximately 175 mg CE/g (Awika, McDonough & Rooney, 2005). The differences could be due to genetics and environmental factors which have been suggested to influence condensed tannin content in sorghum (Awika, Rooney, Wu, Prior & Cisneros-Zevallos, 2003). The protein precipitation capacity of sorghum bran was comparable to that reported for whole grain methanol extracts from NK300, a type III condensed tannin sorghum, which was 2.84 absorbance units per gram of whole grain (Hagerman & Butler, 1978) which translated to 28.4 absorbance units per gram of bran, assuming that bran account for 10% of total grain weight (Sikwese & Duodu, 2007).

4.2.4.2. Condensed tannin content and protein precipitation capacity of aqueous extracts

Water extracts from marama bean seed coats had the highest condensed tannin content measured with both the vanillin-HCl and the Butanol-HCl methods and protein precipitation capacity amongst all the extracts (Table 4.2.2). The condensed tannin contents of acidified water extracts from marama bean seed coats were 2 and 2.3 fold lower than water extracts as measured with the vanillin-HCl and butanol-HCl assays, respectively. The reduction in condensed tannins could be as a result of their interaction with cell wall polysaccharides (Hanlin, Hrmova, Harbertson & Downey, 2010) through the formation of strong hydrogen bonds at low pH resulting in the formation of an insoluble interpolymer complex precipitate (Khutoryanskiy, Dubolazov & Mun, 2009) as mentioned earlier (section 4.1.4.1). However, there were no significant differences in condensed tannin content between water extracts and acidified water extracts from condensed tannin sorghum bran as condensed tannins from sorghum bran probably interacted with cell wall polysaccharides to a lesser extent compared to condensed tannins from marama bean seed coats under acidic condition.

Condensed tannin content of water extract and acidified water extract from marama bean seed coats were 19 and 9 fold respectively higher, as measured with the vanillin-HCl method and 7 and 4 fold, respectively higher as measured with the butanol-HCl assay than equivalent extracts from condensed tannin sorghum bran. The vanillin-HCl assay gave a measure of the amount of monomeric, oligomeric and polymeric species in the sample, while the butanol-HCl assay gave a measure of the amount of anthocynins formed from the hydrolysis and autoxidation of tannins.

The protein precipitation capacity of acidified water extracts from marama bean seed coats were 2 fold lower than that of water extracts. The significant reduction in protein precipitation capacity was proportional to the reduction in condensed tannin content, which was also by 2 fold as a result of extraction under acidic conditions. However, there were no significant differences in protein precipitation capacities between water extracts and acidified water extracts from condensed tannin sorghum bran and this may be due to the lack of significant differences in tannin contents between the extracts. Water extracts and acidified water extracts from marama bean seed coats showed a 47 and 37 fold higher protein precipitation capacities respectively compared to equivalent extracts from condensed tannin sorghum bran.

Table 4.2.2. Effect of extraction under acidic condition on condensed tannin content and protein precipitation capacity of aqueous extracts from marama bean seed coats and condensed tannin sorghum bran

Extract	Vanillin-HCl assay (mg CE ¹ /g)	Butanol-HCl assay (Abs ₅₅₀ units/g)	Protein precipitation capacity (Abs ₅₁₀ units/g)
<u>Marama bean seed coat</u>			
Water	199.9 ² c ³ ± 1.1	122.0 c ± 3.0	83.9 c ± 2.7
Acidified water	99.2 b ± 3.7	52.7 b ± 4.7	48.8 b ± 1.0
<u>Sorghum bran</u>			
Water	10.5 a ± 1.9	13.3 a ± 1.1	1.8 a ± 0.1
Acidified water	13.9 a ± 1.9	14.6 a ± 2.0	1.3 a ± 0.2

¹CE, Catechin equivalents

²Data are means ± standard deviations of six determinations, on dry weight basis

³Means followed by different letters in the same column are significantly different (p ≤ 0.05)

The higher protein precipitation capacity may not be accounted for by the condensed tannin content only. It appears that marama bean seed coat proanthocyanidins may have a higher affinity for proteins to form complexes compared to condensed tannin sorghum bran proanthocyanidins.

4.2.4.3. Characterization of condensed tannins in extracts

4.2.4.3.1. Marama bean seed coat proanthocyanidins

Total ion current chromatograms and HPLC chromatograms of thiolysis degradation products of proanthocyanidins in the polymer fraction (FIII) from aqueous acetone, water and acidified water extracts from marama bean seed coats had similar phenolic compound profiles (Fig. 4.2.1) and the identities of the component peaks are shown in Table 4.2.3.

The chromatograms were similar to those of thiolysis degradation products of proanthocyanidins from the oligomeric fraction (FII) (data not shown). The chromatograms showed six distinct peaks labelled 1 to 6 and the mass spectrum of each compound is shown in Fig. 4.2.2.

The compound eluting in Peak 1 with an abundant $[M-H]^-$ molecular ion at m/z 457.7 (100) (Fig. 4.2.2A) was identified as (epi)catechin-3-O-gallate (MW = 458.37). The peak also showed a fragment ion at m/z 305.9 (14) which was identified as the deprotonated epigallocatechin ion and at m/z 169.5 (21) corresponding to the gallic acid ion. This compound has been identified in aqueous methanol extracts from adzuki beans also by the molecular ion $[M-H]^-$ ion at m/z 457 (Amarowicz, Estrella, Hernández & Troszyńska, 2008). Similar fragmentation pattern for epigallocatechin-3-O-gallate was reported in green tea (Miketova, Schram, Whitney, Kerns, Valcic, Timmermann, Volk, 1998) and oolong tea (Dou, Lee, Tzen & Lee, 2007) extracts.

Compounds eluting in Peak 2a and 2b with abundant molecular ion at m/z 441.8 (100) (Fig. 4.2.2B and 4.2.2C) were identified as isomers of (epi)catechin-3-O-gallate (MW=442.37). Fragmentation resulted in ions at m/z 289.5 (10) and at m/z 169.3 (40) which were identified as (epi)catechin and gallic acid ions. The retention time and mass spectrum of (-)-epicatechin-3-O-gallate standard was the same as that of peak 2b (Data not shown) therefore 2a could be (+)-catechin-3-O-gallate. Miketova *et al.* (1998) reported similar mass spectra data for (epi)catechin-3-O-gallate with m/z values at 441(100), 289 (16) and 169 (18).







Figure 4.2.1. Total ion current chromatogram (top) and HPLC chromatogram (bottom) recorded at 280 nm of thiolysis degradation products of proanthocyanidins in polymer fraction (FIII) from extracts prepared from marama beans seed coats.




A) (Epi)gallocatechin-3-O-gallate




B) (Epi)catechin-3-O-gallate




C) (Epi)catechin-3-O-gallate




D) (Epi)gallocatechin benzylthioether



E) (Epi)gallocatechin-3-O-gallate benzylthioether



F) (Epi)catechin benzylthioether



G) (Epi)catechin-3-O-gallate benzylthioether

Figure 4.2.2. Mass spectra of thiolysis degradation products of proanthocyanidins in acetone extracts from marama bean seed coats.

Table 4.2.3. Mass spectra data of thiolysis degradation products of proanthocyanidins in extracts from marama bean seed coats

Peak No.	t_R^* (min)	Identification	Molecular Weight	m/z [M-H] ⁻ (% intensity)	m/z MS ⁿ (% intensity)
Terminal units release as free flavon-3-ol					
1	18.1	(Epi)gallo catechin-3-O-gallate	458	457.7 (100)	305.9 (14), 169.6 (32)
2	23.0	(Epi)catechin-3-O-gallate	442	441.8 (100)	289.5 (10), 169.3 (40)
Extension units released as benzylthioether derivatives					
3	33.9	(Epi)gallo catechin benzylthioether	428	427.7 (100)	303.5 (80)
4	35.3	(Epi)gallo catechin-3-O-gallate benzylthioether	580	579.8 (100)	457.0 (14), 303.6 (20), 169.6 (11)
5	38.4	(Epi)catechin benzylthioether	412	411.7 (90)	287.7 (100)
6	39.3	(Epi)catechin-3-O-gallate benzylthioether	564	563.5 (100)	439.8 (17), 287.7 (53), 169.1 (38)

* t_R – retention time

Extension units released as benzylthioether derivatives from marama bean seed coat proanthocyanidins are shown as peaks 3 to 6 (Fig. 4.2.1). Peak 3 with abundant $[M-H]^-$ molecular ion at m/z 427.3 (100) (Fig. 4.2.2D) and fragment ion at m/z 303.5 (80) was identified as (epi)gallocatechin benzylthioether (MW=428).

Compound in Peak 4 with abundant $[M-H]^-$ molecular ion at m/z 579.8 (100) (Fig. 4.2.2E) was identified as (epi)gallocatechin-3-O-gallate benzylthioether (MW=580). Fragmentation gave rise to m/z 303.6 (20) and m/z 169.3 (11) which corresponded to (epi)gallocatechin ion and gallic acid fragment ions, respectively. Similar fragmentation patterns for (epi)gallocatechin benzylthioether and (epi)gallocatechin-3-O-gallate benzylthioether have been reported in thiolysis degradation products of a highly galloylated proanthocyanidin polymer from persimmon fruit (Li, Laveverence, Trombley, Xu, Yang, Tian, Reed & Hagerman, 2010).

The compound eluting in peak 5 with $[M-H]^-$ molecular ion at m/z 411.7 (90) and major fragment ion at m/z 287.7 (100) (Fig. 4.2.2F) was identified as (epi)catechin benzylthioether (MW=412). Peak 6 with abundant $[M-H]^-$ molecular ion at m/z 563.8 (100) was identified as (epi)catechin-3-O-gallate benzylthioether (MW=564) (Figure 4.2.2.G). Fragment ion at m/z 439.8 (17) corresponded to (epi)catechin-3-O-gallate ion after cleavage of thiol group, at m/z 287.7 (44) corresponded to (epi)catechin ion and at m/z 169.1 (35) corresponded to galloyl group. Gu *et al.* (2003b) also identified epicatechin benzylthioether by molecular ion at m/z 411 and fragment ion at m/z 287 and epicatechin-3-O-gallate benzylthioether by the molecular ion at m/z 563 and fragment ion at m/z 439 in grape proanthocyanidin thiolysis degradation products.

4.2.4.3.2. Condensed tannin sorghum bran proanthocyanidins

Total ion current (TIC) and HPLC chromatograms of thiolytic degradation products of proanthocyanidins in polymer fraction (FIII) of aqueous acetone, water and acidified water extracts from condensed tannin sorghum bran are shown in Fig. 4.2.3 and the identities of the peaks are shown in Table 4.2.4. The compound profiles of the three extracts were similar. The total ion current and HPLC chromatograms of thiolytic degradation products of proanthocyanidins from the oligomer fraction (FII) (data not shown) were similar to those of the polymer fraction (FIII).




Figure 4.2.3. Total current ion (top) and HPLC (bottom) chromatograms of thiolytic degradation products of proanthocyanidins in polymer fraction from extracts prepared from condensed tannin sorghum bran.




Figure 4.2.4. Mass spectra of thiolysis degradation products of proanthocyanidins in acetone extracts from condensed tannin sorghum bran.

Table 4.2.4. Mass spectra data of thiolysis degradation products of proanthocyanidins from condensed tannin sorghum bran

Peak No.	t_R^* (min)	m/z [M-H] ⁺ (intensity, %)	m/z MS ⁿ (intensity, %)	Identification
1	15.6	289.6 (100)	-	(+)-catechin
2	38.4	441.8 (100)	287.2 (100)	(Epi)catechin benzylthioether

* t_R – retention time

- No fragment ion of parent compound was observed

Peak 1 with abundant molecular ion at m/z 289.6 (100) (Fig. 4.2.4A) corresponded to (+)-catechin (MW=290.27) occurring as a terminal units. (+)-Catechin standard showed abundant molecular ion at m/z 290.1 and its retention time was the same as that for peak 1 therefore it was concluded that peak 1 was (+)-catechin. Duo *et al.* (2007) also identified (+)-catechin by the molecular ion at m/z 289 in oolong tea extracts. Terminal units in proanthocyanidins of condensed tannin sorghum bran were reported to be composed of (+)-catechin accounting for 89% and (-)-epicatechin accounting for 11% (Gu *et al.*, 2002). In this work (-)-epicatechin was not detected, probably because it occurred in low amounts.

The extension unit benzylthioether derivative from condensed tannin sorghum bran proanthocyanidins is shown as peak 2 (Fig. 4.2.4B). The compound in peak 2 with molecular

ion at m/z 411.8 (90) and major fragment ion at m/z 287.4 (100) was identified as (epi)catechin benzylthioether (MW=412). (-)-Epicatechin extension units released as (epi)catechin benzylthioether were identified by the parent ion at m/z 411 and fragment ion at m/z 287, in cocoa, sorghum bran, lowbush blueberry and cranberry proanthocyanidin thiolytic degradation products (Gu *et al.*, 2002). The peak marked (i) (Fig. 4.2.3A) had an abundant molecular ion at m/z 577(100) and was identified as A-type procyanidin dimer (MW=578) which has been previously reported in sorghum (Awika & Rooney, 2004).

4.2.4.3.3. Flavan-3-ol composition and mean degree of polymerization of proanthocyanidins in extracts

Flavan-3-ol composition and mean degree of polymerization (mDP) of proanthocyanidins in marama bean seed coat and condensed tannin sorghum bran extracts were calculated from HPLC chromatogram peak areas and are presented in Table 4.2.5. The structural composition of proanthocyanidins from marama bean seed coat was different from that of condensed tannin sorghum bran. The constitutive units in marama bean seed coat proanthocyanidins were (epi)gallo catechin-3-O-gallate, (epi)catechin-3-O-gallate, (epi)gallo catechin and epicatechin while in sorghum these were (+)-catechin and (-)-epicatechin. Marama bean seed coat proanthocyanidins differed from those of other legumes such as the common bean and lentils because of the higher percentage of galloylated subunits. Terminal units of marama bean seed coat proanthocyanidins were epigallo catechin-3-O-gallate and (epi)catechin-3-O-gallate, while in pinto beans, small red beans and red kidney beans were reported to be catechin (7.3–14.5%) and epicatechin (0.4–1.1%) (Gu *et al.*, 2003b) and in lentils were reported to be catechin (8.6–11.9%), epicatechin (1.3–1.7%) and traces of epicatechin-3-O-gallate (Dueñas *et al.*, 2003).

Extension units in marama bean seed coats proanthocyanidins were predominantly epigallo catechin-3-O-gallate and (Epi)catechin-3-O-gallate, while (Epi)gallo catechin and (epi)catechin (EC) were present as minor extension unit components, suggesting that the proanthocyanidins in the seed coats are predominantly prodelphinidins.

Table 4.2.5. Constitutive unit composition (relative percentage by peak area) of proanthocyanidins in polymer (FIII) and oligomer (FII) fractions from aqueous extracts from marama bean seed coat and condensed tannin sorghum bran¹

Extract	% Terminal units released as free flavan-3-ol			% Extension units released as benzylthioether derivatives					
	C ²	EGCG	ECG	EGC	EGCG	EC	ECG	mDP	
<u>Marama bean seed coat</u>									
FIII	Water	-	2.7 a ³ ± 0.2	8.2 a ± 0.7	6.6 a ± 1.2	43.3 d ± 0.7	5.7 a ± 0.4	33.5 b ± 1.5	9.2 e ± 0.6
	Acidified water	-	3.9 a ± 0.5	10.3 b ± 0.8	7.0 a ± 1.4	40.0 c ± 2.2	6.2 a ± 0.1	32.6 b ± 0.2	7.1 d ± 0.5
FII	Water	-	5.9 b ± 0.8	13.6 c ± 0.8	8.2 a ± 2.1	35.1 b ± 0.8	7.0 a ± 0.5	30.1 a ± 2.1	5.1 b ± 0.3
	Acidified water	-	7.5 b ± 1.5	18.7 d ± 2.2	7.1 a ± 1.2	29.3 a ± 2.7	8.0 a ± 1.5	29.4 a ± 1.9	3.8 a ± 0.2
<u>Sorghum bran</u>									
FIII	Water	18.4 a ± 2.5	-	-	-	-	81.6 d ± 2.5	-	5.5 bc ± 0.7
	Acidified water	15.8 a ± 1.8	-	-	-	-	84.2 d ± 1.8	-	6.4 cd ± 0.7
FII	Water	26.8 b ± 2.1	-	-	-	-	73.2 c ± 2.1	-	3.7 a ± 0.3
	Acidified water	33.4 c ± 1.6	-	-	-	-	66.6 b ± 1.6	-	3.0 a ± 0.1

¹ Data are means and standard deviations of four determinations from two independent experiments.

² Abbreviations: C, catechin; EGCG, (epi)gallocatechin-3-O-gallate; ECG, (epi)catechin-3-O-gallate; EGC, (epi)gallocatechin; EC, (epi)catechin; mDP, mean degree of polymerization

³ Means with different letters in the same column are significantly different ($p \leq 0.05$).

- Not detected

In comparison, extension units of proanthocyanidin from pinto beans, small red beans and red kidney beans were found to be composed of epicatechin (43.1–73.9%), catechin (5.4–33.2%) and (epi)afzelechin (8.7–14.6%) (Gu *et al.*, 2003b) and those in polymer fraction from the seed coats of lentils were composed of catechin (49.5–61.9%), epicatechin (20.8–33.8), epigallocatechin (3.8–6.7%) and traces of epigallocatechin-3-O-gallate (Dueñas *et al.*, 2003). The flavan-3-ol constitutive unit composition of sorghum bran proanthocyanidins were similar to those reported by Gu *et al.* (2003b) except that in this study (-)-epicatechin occurring as terminal unit was not detected. These authors reported that terminal units in sorghum bran proanthocyanidins were predominantly (+)-catechin (89%) and (-)-epicatechin (11%) and extension units were exclusively (-)-epicatechin.

Aqueous extracts from marama bean seed coats had a higher mDP compared to equivalent extracts from condensed tannin sorghum bran. Extraction under acidic conditions of marama bean aqueous extracts resulted in a significant reduction in the mDP of proanthocyanidins in both oligomer (FII) and polymer (FIII) fractions. The reduction in mDP may be as a result of interaction of high molecular weight proanthocyanidins with cell wall polysaccharides (Hanlin *et al.*, 2010) at low pH resulting in the formation of an insoluble interpolymer complex precipitate (Khutoryanskiy *et al.*, 2009) as mentioned earlier (section 4.1.4.1) resulting in extraction of lower molecular weight polymers. The higher molecular weight proanthocyanidin polymers may be more involved in the interaction because of their higher number of hydroxyl groups that may form hydrogen bonds with other polymers causing precipitation. The mDP of marama bean seed coat proanthocyanidins was similar to that of other legumes such as the common bean and lentils. The mDP of proanthocyanidins in aqueous acetone extracts from pinto beans, small red beans and red kidney beans ranged from 6.3–12.0, depending on variety (Gu *et al.*, 2003b) and in polymer and oligomer fractions from lentils it ranged from 6.9–9.4 and 3.8–4.7, respectively (Dueñas *et al.*, 2003). Extraction under acidic conditions did not have a significant effect on the mDP of proanthocyanidins in acidified water extracts from condensed tannin sorghum bran. The change in pH had little or no effect on the formation of hydrogen bonds and eventual precipitate formation when compared to proanthocyanidins from marama bean seed coats. This may be due to the lower number of hydroxyl groups on the procyanidin polymer as well as the lower condensed tannin content.

The mDP of proanthocyanidins from condensed tannin sorghum bran were lower than that of aqueous acetone extracts from Sumac sorghum, a condensed tannin sorghum cultivar, which was reported to be 8.4 (Gu *et al.*, 2002). This indicated that lower molecular weight polymers were extracted under aqueous condition compared to extraction with organic solvent. The mDP obtained for oligomeric (FII) and polymeric (FIII) fractions from both marama bean seed coats and sorghum bran were lower than 10 which suggested that the proanthocyanidins in both fractions obtained from the Sep Pak cartridges were all oligomers.

4.2.5. Conclusions

Aqueous extracts from marama bean (*Tylosema esculentum*) seed coats have higher condensed tannin content and protein precipitation capacity than equivalent extracts from bran of condensed tannin sorghum variety PAN 3860. Proanthocyanidins in marama bean seed coats are predominantly prodelphinidins and those from condensed tannin sorghum bran are procyanidins. Prodelphinidins have more hydroxyl groups within their polymer structure than procyanidins and this may render the polymer more water soluble than procyanidins resulting an extract with higher condensed tannin content. The higher number of hydroxyl groups in the polymer structure of prodelphinidins also results in higher interaction with proteins probably through hydrogen bonding resulting in higher protein precipitation capacity compared to procyanidins from condensed tannin sorghum bran.

Extraction under acidic condition results in a decrease in condensed tannin content of aqueous extracts from seed coats of marama beans. However, it does not have any significant effect on condensed tannin content of aqueous extracts from condensed tannin sorghum bran. The decrease in condensed tannin content of acidified water extracts from marama bean seed coats may be due to the large number of hydroxyl groups on the prodelphinidin structure which allows the polymer to interact with other high molecular weight cell wall polysaccharides through hydrogen bonding resulting in interpolymer complexation and precipitation at low pH.

4.3. *In vitro* antioxidant activity and protective effect of aqueous extracts prepared from marama bean seed coat and condensed tannin sorghum bran against oxidative damage of biological molecules

Part of this chapter has been published in the International Journal of Food Science and Technology¹ and accepted for publication in the Journal of Food Biochemistry².

¹Shelembe, J.S., Cromarty, D., Bester, M.J., Minnaar, A. and Duodu, K.G. (2011) Characterisation of phenolic acids, flavonoids, proanthocyanidins and antioxidant activity of water extracts from seed coats of marama bean (*Tylosema esculentum*) – an underutilised food legume. *International Journal of Food Science and Technology* 47, 648-655.

²Shelembe, J.S., Cromarty, D., Bester, M.J., Minnaar, A. and Duodu, K.G. (2011) Effect of acidic condition on phenolic composition and antioxidant potential of aqueous extracts from sorghum (*Sorghum bicolor*) bran. *Journal of Food Biochemistry* (in press)

4.3.1. Abstract

Water and acidified water extracts prepared from marama bean seed coats and condensed tannin sorghum bran for possible use as natural dietary antioxidant were evaluated for free radical scavenging activities using the 2,2'-azinobis- (3-ethyl-benzothiazoline-6-sulphonic acid (ABTS), 2,2-diphenyl-1-picrylhydrazyl (DPPH) and oxygen radical absorbance capacity (ORAC) assays. The extracts were also evaluated for their protective effect against free radical oxidative damage of biomolecules using physiological relevant assays. These were AAPH-induced human red blood cell haemolysis, AAPH-induced supercoiled plasmid pBR 322 DNA oxidative damage, and copper-catalyzed LDL oxidation assays. Aqueous extracts from marama bean seed coats had higher free radical scavenging activities than equivalent extracts from condensed tannin sorghum bran. Extraction of marama bean seed coats under acidic conditions resulted in an extract with lower free radical scavenging activity, while it had no significant effect on extracts from condensed tannin sorghum bran. The free radical scavenging activity of the extracts correlated with total phenolic, flavonoid and condensed tannin contents. Extracts from marama bean seed coats protected against AAPH-induced red blood cell haemolysis and copper-catalyzed LDL oxidation at lower concentration than equivalent extracts from condensed tannin sorghum bran. Extracts from condensed tannin sorghum bran showed a clear protective effect against AAPH-induced oxidative DNA damage. However, extracts from marama bean seed coats gave inconclusive results probably due to prodelphinidins binding to DNA causing poor mobility of the DNA within the agarose gel. In conclusion, the extracts from marama bean seed coats and condensed tannin sorghum bran have a potential to reduce oxidative stress implicated in the development of chronic diseases.

Keyword: Marama bean seed coat; Sorghum bran; Antioxidant activity; Erythrocyte haemolysis, Oxidative DNA damage; LDL oxidation

4.3.2. Introduction

Numerous studies have shown that a number of non-nutrient dietary compounds, known as antioxidants might have a beneficial role in protecting against the development of chronic degenerative diseases (Paiva-Martins, Fernandes, Rocha, Nascimento, Vitrino, Amado, Borges, Belo & Santos-Silva, 2009) associated with oxidative stress (Gheldof & Engeseth, 2002; Somparn, Phisalaphong, Nakornchai, Unchern & Morales, 2007) including cardiovascular disease (CVD) (Hodgson, Croft, Puddy, Mori & Beilin, 1996), neurodegenerative diseases and cancer (Somparn *et al.*, 2007). Oxidative stress is caused by reactive oxygen species (ROS) which are harmful by-products produced during normal cell metabolism (Liyana-Pathirana & Shahidi, 2005; Dimitrios, 2006). Excessive generation of ROS can modify or cause oxidative damage to DNA, proteins and polyunsaturated fatty acids (Heo & Lee, 2005) in the human body and these changes have been implicated in the pathogenesis of human chronic diseases (Jacob & Burri, 1996).

Lipid peroxidation is a major cause of many pathological effects such as CVD, cancer, brain dysfunction and the ageing process (Lim, Cheung, Ooi & Ang, 2002). The major sites of lipid peroxidation damage are biomembranes due to the presence of polyunsaturated fatty acids in the membrane (Mak, Misra & Weglicki, 1983). Destruction of membrane lipids compromises the function of membrane localized receptors and channels and it also affects the transmembrane passage of solutes and fluidity of the membrane (Reiter, 1998) and thereby affecting the function of the cell and subcellular organelles such as mitochondria, microsomes and lysosomes (Cejas, Casado, Belda-Iniesta, Catro, Espinosa, Redondo, Sereno, García-Cabezas, Vara, Domínguez-Cáceresi, Perona & González-Barón, 2004). Lipid peroxides decomposes to relatively stable toxic aldehyde products which can diffuse and attack biomolecules such as proteins, DNA and phospholipids forming intra and inter molecular covalent adducts (Cejas *et al.*, 2004) leading to decline in physiological functions of the cell and ultimately cell death.

In European countries CVD is ranked as the number one cause of death, accounting for 48% of all deaths in 2008 (European Heart Network, 2008) and worldwide it is also a rapidly growing problem (de Lima Portella, Barcelos, de Bern, Carratu, Bresolin, da Rocha & Soares, 2008). It is reported that in South Africa, between 1997 and 2004, 195 people died per day because of some form of heart and blood vessel disease (CVD) (Steyn, 2007). LDL oxidation is implicated in the development of coronary heart disease (Regnström, Ström,

Moldeus & Nilsson, 1993). Oxidized LDL is recognizable by the non down-regulating macrophage scavenger receptor (Mosinger, 1995) resulting in the uptake of the oxidized LDL by the macrophage (foam cells) (Jacob & Burri, 1996) and smooth muscle cells leading to the formation of fatty streaks (Baba, Osakabe, Kato, Natsume, Yasuda, Kido, Kukuda, Muto & Kondo, 2007). In recent years there has been great interest in antioxidants that retard LDL oxidation (Paiva-Martins *et al.*, 2009). In human subjects, phenolic compounds in cocoa powder increased resistance of LDL to oxidation (Baba *et al.*, 2007) and those found in virgin olive oil resulted in a decrease in LDL oxidation markers (de la Torre-Carbor, Chávez-Servin, Jaúregui, Castellote, Mamuela-Raventós, Nurmi, Paulsen, Gaddi, Kaikkonen, Zunt, Fitó, Covas & López-Sabater, 2010) demonstrating their potential to prevent atherosclerotic processes which lead to CVD.

Worldwide cancer is the leading cause of deaths and in 2008 it accounted for 13% of all deaths (WHO, 2011). Most cancers begin with somatic mutation of normal cells (Jacob & Burri, 1996) which may be initiated by oxidative DNA damage (Laparra, Vélez, Barberá, Farré & Montoro, 2008) caused by ROS (Ames, 1983). Phenolic compounds of plant origin have been shown to have anticancer and apoptosis inducing properties (Zhang, Spitz, Tomlinson, Schabath, Minna & Wu, 2002). Green tea extracts were found to have chemoprotective effect against DNA damage (Bhat, Azmi, Hanif & Hadi, 2006) and epigallocatechin-3-O-gallate from tea was found to inhibit the growth of nasopharyngeal carcinoma cells in a dose-dependent manner (Yan, Yong-Guang, Fei-Jun, Fa-Qing, Min & Ya, 2004).

The aims of this study were (1) to determine the effect of extraction under acidic condition on free radical scavenging activity of aqueous extracts from marama bean seed coats and condensed tannin sorghum bran, (2) to determine the protective effect of the extracts against erythrocyte membrane, LDL and plasmid DNA oxidative damage.

4.3.3. Materials and methods

4.3.3.1. Materials

Marama beans were obtained from Botswana, Ghanzi province from the 2008 season and condensed tannin sorghum was from the 2004 season grown in South Africa (Section 4.1.3.1). Supercoiled plasmid vector pBR322 DNA, agarose (D1 LE) and bromophenol blue

were purchased from Whitehead Scientific (Pty) Ltd (Johannesburg, South Africa). Potassium persulphate, ethylenediaminetetraacetate (EDTA), Tris(hydroxymethyl) aminomethane, glacial acetic acid, saccharose, sodium chloride, potassium chloride, sodium hydrogen phosphate 1-hydrate ($\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$), di-sodium hydrogen phosphate (Na_2HPO_4), methanol, copper sulphate, thiobarbituric acid, sodium hydroxide, trochloroacetic acid were purchased from Merck (Pty) Ltd South Africa (Johannesburg, South Africa). 2,2'-azinobis-(3-ethyl-benzothiazoline-6-sulphonic acid (ABTS), ethidium bromide, 2,2'-Azobis (2-methyl-propionamide) dihydrochloride (AAPH), 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid (Trolox), fluorescein (FL), 2,2-diphenyl-1-picrylhydrazyl (DPPH), human low density lipoprotein (LDL) were supplied by Sigma-Aldrich (Pty) Ltd (Johannesburg, South Africa).

4.3.3.2. Methods

4.3.3.2.1. Sample preparation and extraction

Marama bean seed coats and condensed tannin sorghum bran powder were prepared as described in section 4.1.3.2.1 and their freeze-dried aqueous extracts were prepared as described in section 4.1.3.2.2.

4.3.3.2.2. ABTS free radical scavenging assay

Total antioxidant activities of the extracts were determined using the $\text{ABTS}^{\bullet+}$ radical scavenging assay (Awika, Rooney, Wu, Prior & Cisneros-Zevallos, 2003; Siddhuraju, 2006; Thaipong, Boonprakob, Crosby, Cisneros-Zevallos & Byrne, 2006; Xu, Ye, Chen & Liu, 2007). In this assay, phenolic antioxidants react directly with preformed stable ABTS radical cations causing decrease in absorbance or colour of the radical (Hagerman, Riedl, Jones, Sovik, Ritchard, Hartfeld & Richel, 1998). The stable radical is generated by potassium persulphate oxidation of ABTS^{2-} to form a blue-green $\text{ABTS}^{\bullet+}$ chromophore (Macdonald, Wood & Garg, 2006). The $\text{ABTS}^{\bullet+}$ radical cation stock solution was produced by mixing 5 ml of 7.4 mM ABTS solution and 5 ml of 2.45 mM potassium persulphate solution and allowed to react in the dark at room temperature for 16 hours. $\text{ABTS}^{\bullet+}$ radical cation working solution was prepared by diluting 5 ml of the $\text{ABTS}^{\bullet+}$ radical cation stock solution with 145 ml Phosphate buffer saline (PBS) pH 7.4 (0.2 M Na_2HPO_4 , 0.2 M NaH_2PO_4 and 150 mM NaCl).

Freeze-dried aqueous extracts from marama bean seed coat (25 mg) and condensed tannin sorghum bran (50 mg) were weighed into a 25 ml volumetric flask and made up to volume with PBS to give a concentration of 1 mg/ml and 2 mg/ml, respectively and then further diluted to 0.1 mg/ml and 0.4 mg/ml respectively. Trolox standard serial dilutions (0–1000 μ M Trolox) were prepared in PBS. To 100 μ l of diluted sample extract or Trolox standard solution 2850 μ l of ABTS^{•+} radical working solution was added and allowed to react for 30 min and the absorbance measured at 734 nm using a T80+ UV-VIS spectrophotometer (PG Instruments, Wibtsoft, UK). Results were reported as micromoles of Trolox equivalents/g (μ mol TE/g), on dry sample weight basis.

4.3.3.2.3. DPPH free radical scavenging assay

The antioxidant activity of the extracts was also determined with the 2,2-diphenyl-1-picrylhydrazyl (DPPH) free radical scavenging assay (Brand-Williams, Cuvelier & Berset, 1995) which measures the ability of an antioxidants to quench the DPPH[•] radical (Cardador-Martinez, Loarca-Pina & Oomah, 2002; Nuengchamnonng, Krittasilp & Ingkaninan, 2009). DPPH is a free radical that is stable at room temperature which upon reduction by an antioxidant results in loss of absorbance at 515–517 nm (Brand-Williams *et al.*, 1995). The degree of discolouration indicates the scavenging efficiency of the added antioxidant (Xu & Chang, 2007). The DPPH[•] stock solution was prepared by weighing 12 mg of DPPH into a 50 ml volumetric flask and made to volume with methanol. Ten millilitres of this stock solution was further diluted with 50 ml methanol to obtain an absorbance value of approximately 1.1 at 515 nm. Trolox standard serial dilutions ranging from 0 to 1000 μ M were prepared in methanol. A 1 mg/ml solution in methanol of each extract was prepared. These were further diluted with methanol to final working concentrations of 0.5 and 0.1 mg/ml for extracts from marama bean seed coats and condensed tannin sorghum bran, respectively. To 150 μ l of the diluted sample extract solution or Trolox solution, 2850 μ l of DPPH working solution was added and the tubes were sealed and covered with foil. Samples were shaken for 60 min and Trolox standards for 15 min and absorbance was measured at 515 nm using a T80+ UV-VIS spectrophotometer. Results were reported as micromoles of Trolox equivalents/g (μ mol TE/g), on dry sample weight basis.

4.3.3.2.4. Oxygen radical absorbance capacity assay (ORAC)

The method of Ou, Hampsch-Woodill and Prior (2001) was used to measure the oxygen radical absorbance capacity of the extracts with some modifications. In this method fluorescein (FL) is used as a probe which is attacked by peroxy radicals generated by 2,2'-Azobis (2-methyl-propionamidine) dihydrochloride (AAPH) resulting in loss of fluorescence which is recorded at specific time intervals as the reaction goes to completion. The presence of an antioxidant compound inhibits free radical damage of fluorescein (Xu & Chang, 2007) and its protective effect is measured by assessing the area under the fluorescence decay curve (AUC) (Ou *et al.*, 2001).

Freeze-dried marama bean seed coat and condensed tannin sorghum bran aqueous extracts were dissolved in distilled water at a concentration of 0.12 mg/ml and 0.15 mg/ml, respectively. Trolox standards ranging from 0 to 1 mM were prepared in distilled water. FL stock solution was prepared by dissolving 3.76 mg FL in 50 ml PBS pH 7.4 (prepared as described in section 4.3.3.2.2). FL working solution was made by mixing 120 µl FL stock solution, 5 ml PBS and 45 ml distilled water. AAPH solution (74 mM) was prepared by dissolving 80 mg AAPH in 4 ml distilled water. Trolox standard or sample extract (10 µl) was added into 96 well plate in triplicates. A FL-AAPH solution was prepared by mixing the 4 ml of AAPH solution with 16 ml of FL working solution and immediately after mixing, 200 µl was added to each well containing Trolox, or extract. For blank and negative control 200 µl FL working solution and 200 µl FL-AAPH solution respectively was added to wells containing 10 µl distilled water. The well plates were placed in a FLUOstar OPTIMA multifunctional plate reader (BMG LABTECH GmbH, Ortenberg, Germany) set at 37 °C. Fluorescence at 485 nm excitation and 520 nm emission (Macdonald *et al.*, 2006; Thaipong *et al.*, 2006) was recorded at every 5 min interval for 245 min. Area under curve (AUC) was calculated for samples and standards using Microcal Origin software, version 6 (Microcal Software Inc, Northampton, USA). AUC for samples and standards were subtracted from AUC for blank. Antioxidant activity for samples was calculated from Trolox calibration curves and results reported as micromoles Trolox equivalents/g (µmol TE/g) on dry weight basis.

4.3.3.2.5. AAPH-mediated red blood cell haemolysis assay

Aqueous extracts were evaluated for their capacity to protect biomembrane against oxidative damage by free radicals using human red blood cell as a model system. In this assay haemolysis of red blood cell membrane is induced with AAPH (Paiva-Martins *et al.*, 2009). AAPH decomposes at physiological temperature and generates peroxy radicals (Somparn *et al.*, 2007) which cause peroxidation of lipids and oxidation of proteins in the membrane leading to membrane rupture or formation of haemolytic holes (Sato, Sato & Suzuki, 1999) and resultant leakage of cell contents. Antioxidants that scavenge peroxy radicals would reduce the rate of red blood cell haemolysis.

The red blood cell haemolysis assay was performed according to Tang and Liu (2008). Human red blood cells were obtained from the Department of Pharmacology of the University of Pretoria and were suspended in isotonic phosphate buffer saline PBS, pH 7.4 (8.1 mM Na₂HPO₄, 1.9 mM NaH₂PO₄, 3 mM KCl and 0.137 M NaCl) in a ratio of 1:1, stored at 4 °C and used within 3 days. Prior to use, the red blood cell suspension was centrifuged at 2750 rpm (1184 x g) for 3 min using model Z300 centrifuge (Hermle Labortechnik GmbH, Wehingen, Germany) and the supernatant removed and discarded. Red blood cells were washed 3 times by adding isotonic PBS in a ratio of 1:1 and centrifuging at 2750 rpm (1184 x g) for 3 min and with each step the supernatant was discarded. Freeze-dried aqueous extracts from marama bean seed coat and condensed tannin sorghum bran were dissolved in distilled water at a concentration of 1.2 and 12 mg/ml, respectively. Extract solutions from marama bean seed coats were further diluted to three different concentrations 10, 40 and 80 µg/ml concentrations and extracts solution from sorghum bran were further diluted to 100, 200 and 400 µg/ml concentrations. For treatments 100 µl isotonic PBS, 10 µl red blood cells, 40 µl extract solution and 40 µl of 110 mM AAPH solution (prepared in isotonic PBS) were added into 1.5 ml safe lock eppendorf tubes. For sample blanks 140 µl isotonic PBS, 10 µl red blood cells and 40 µl extract were added into each tube. Positive controls were prepared by adding 140 µl isotonic PBS, 10 µl red blood cells and 40 µl of 110 mM AAPH solution into tubes and negative controls were prepared by adding 180 µl isotonic PBS and 10 µl red blood cells. Tubes were sealed, mixed by vortexing and incubated at 37 °C for 16 h. After incubation the tubes were mixed by vortexing and centrifuged at 2750 rpm (1184 x g) for 3 min. Supernatant (50 µl) was transferred into a 96 well plate and absorbance measured at 405 nm on a Bio Tek ELx 800 plate reader (Biotek Instruments Inc, Winooski, USA).

4.3.3.2.6. AAPH-mediated oxidative DNA damage

The protective effects of marama bean seed coat and condensed tannin sorghum bran aqueous extracts on oxidative DNA damage induced by AAPH were assessed by the method as described by Wei, Zhou, Cai, Yang & Liu (2006) with some modification. Under oxidative stress supercoiled plasmid pBR322 DNA is converted into relaxed circular form due to single strand breaks and into linear form due to double strand breaks and these three forms are separated by agarose gel electrophoresis because of their differences in electrophoretic mobility (Aronovitch, Godinger, Israeli, Krishna, Samuni & Goldstein, 2007).

DNA solution was prepared by diluting 10 µg supercoiled plasmid pBR322 DNA with 100 µl Tris-acetate buffer pH 8 working solution and vortexed. Treatment samples were prepared by adding 2.5 µl DNA solution, 5 µl sample extract or Trolox solution and 5 µl of 11 µM AAPH solution into eppendorf tubes. Positive controls were prepared by adding 2.5 µl DNA solution, 5 µl Tris-acetate buffer, pH 8 and 5 µl of 11 µM AAPH into eppendorf tubes and negative controls by adding 2.5 µl DNA solution and 10 µl Tris-acetate buffer solution. Tubes were vortexed and then incubated for 1.5 h at 37 °C. Twelve microliters of loading buffer (40% sucrose and 0.025% bromophenol) solution was added into each tube and vortexed. Twenty microlitres of sample was loaded into the well in the gel in electrophoresis apparatus (Owl Scientific Inc, Woburn, USA) containing 80 mM Tris-acetate buffer pH 8. The gel was run at 60 V, 30 mA for 2 h using electrophoresis power supply unit model EPS 301 (Amersham Pharmacia Biotech, Uppsala, Sweden). DNA bands were visualized and photographed on a UV transilluminator (Vilber Lourmat, Marne la Vallee, France). Band intensities were measured using Image Tool for Windows software version 3.00 (The University of Texas Health Center, San Antonio, USA) and % intensity was calculated relative to the negative control (without AAPH).

4.3.3.2.7. Copper-catalyzed human low density lipoprotein oxidation

The effect of extracts against copper-catalyzed human low density lipoprotein (LDL) oxidation was evaluated using the thiobarbituric assay (TBA assay) (Rüfer & Kulling, 2006; Xu, Yuan, & Chang, 2007a). Freeze-dried marama bean seed coat and condensed tannin sorghum bran aqueous extracts were dissolved in distilled water at a concentration of 1.2 mg/ml and 6 mg/ml, respectively. Extract solutions were further diluted into three different concentrations, 10, 40 and 80 µg/ml for extract solution from marama bean seed coats and

100, 250 and 500 µg/ml for extract solution from condensed tannin sorghum bran. For treatments, 2 µl of LDL solution (5.5 mg/ml), 168 µl 0.02 M PBS solution and 10 µl sample extract or Trolox solution were added into eppendorf safety lock tubes and oxidation was initiated by adding 20 µl of 100 mM copper sulphate solution. Positive controls were prepared by adding 2 µl LDL, 178 µl PBS and 20 µl copper sulphate solutions. Negative controls were prepared by adding 2 µl LDL and 198 µl PBS solutions. Tubes were sealed and incubated in a water bath at 37 °C for 3 h. After incubation, 200 µl of 10 mM EDTA, 200 µl of 20 % (w/v) trichloroacetic acid solution and 200 µl of 0.67 % (w/v) thiobarbituric acid solution in 0.2 M NaOH were added and sealed. Tubes were then heated at 80 °C (Lim *et al.*, 2002) for 30 min in a water bath and after cooling were centrifuged at 1500 g for 15 min and the supernatant was then transferred into a 1 ml cuvette. Absorbance was measured at 532 nm using a T80+ UV-VIS spectrophotometer.

4.3.3.3. Statistical analysis

Results from two independent experiments performed in duplicate were expressed as means ± standard deviations. Data were analyzed using one-way analysis of variance (ANOVA) and post hoc testing using Fishers LSD test for multiple comparisons of means was used to determine significant differences between means. Statistica 8.0 program (StatSoft Inc, Tulsa, USA) was used for statistical data analysis and significance was accepted at $p \leq 0.05$.

4.3.4. Results and discussion

4.3.4.1. Antioxidant capacity

Water extracts from marama bean seed coats exhibited the highest antioxidant activity amongst all the extracts as determined with the ABTS^{•+} and DPPH[•] radical scavenging and ORAC assays (Table 4.3.1).

Extracts from marama bean seed coats had higher antioxidant activity than equivalent extracts from condensed tannin sorghum bran. Water extracts from marama bean seed coats had 35, 9 and 4 fold higher antioxidant activities than water extracts from condensed tannin sorghum bran as measured with the DPPH, ABTS and ORAC assays, respectively. Acidified water extract from marama bean seed coats had 19, 5, and 2 fold higher antioxidant activities than acidified water extract from condensed tannin sorghum bran as measured by DPPH, ABTS and ORAC assays, respectively. The DPPH antioxidant activities of marama bean

aqueous extracts were lower but comparable to those reported for methanol extracts from the seed coats of common beans with dark seed coats which had the highest antioxidant activity averaging 518 $\mu\text{mol TE/g}$, on wet weight basis amongst 28 cultivars (Rannilla, Genovese & Lajolo, 2007).

Table 4.3.1. Effect of extraction under acidic condition on antioxidant activity of aqueous extracts from marama bean seed coats and condensed tannin sorghum bran

Extract	ABTS ($\mu\text{mol TE/g}$)	DPPH ($\mu\text{mol TE/g}$)	ORAC ($\mu\text{mol TE/g}$)
<u>Marama bean seed coat</u>			
Water extract	707.1 ¹ c ² \pm 10.6	492.4 c \pm 17.9	439.3 c \pm 34.9
Acidified water extract	343.2 b \pm 21.3	325.4 b \pm 13.8	273.1 b \pm 42.3
<u>Condensed tannin sorghum bran</u>			
Water extract	75.2 a \pm 7.6	14.0 a \pm 0.8	111.1 a \pm 20.1
Acidified water extract	70.7 a \pm 2.3	17.0 a \pm 1.0	134.9 a \pm 13.5

¹Data are means \pm standard deviations on dry weight basis of four determinations from two independent experiments.

²Means with different letters in the same column are significantly different ($p \leq 0.05$).

However, they were higher than that of acidified acetone extracts from seed coats of black soybean which was approximately 165 $\mu\text{mol TE/g}$ on dry weight basis (Xu & Chang, 2008). The ORAC values for both marama bean aqueous extracts were lower than that reported for acidified acetone extract from seed coats of black soybeans, which was approximately 450 $\mu\text{mol TE/g}$ on dry weight basis (Xu & Chang, 2008).

The DPPH[•], ABTS^{•+} scavenging activities and ORAC values of aqueous extracts from sorghum bran were lower than values reported for aqueous acetone extracts from sorghum variety Sumac (SU99) bran which were 716 $\mu\text{mol TE/g}$, 768 $\mu\text{mol TE/g}$ and 3124 $\mu\text{mol TE/g}$ (Awika *et al.*, 2003) on dry weight basis, respectively. The differences could be due to differences in cultivars, extraction methods and solvents. Water is a poor solvent for extracting phenolic compounds compared to organic solvents such as acetone and methanol and their aqueous mixtures (Yu, Ahmedna & Goktepe, 2005).

Acidified water extract from marama bean seed coats had 2, 1.6 and 1.5 fold lower antioxidant activities as determined with the ABTS^{•+}, DPPH[•] scavenging assays and ORAC assay, respectively compared to water extracts. The lower antioxidant activity was as a result of lower phenolic content which may have been caused by interpolymer precipitation and coprecipitation of phenolic compounds triggered by low pH as discussed in section 4.1.4.1. Extraction of condensed tannin sorghum bran under acidic condition caused a slight increase in DPPH[•] scavenging activity and ORAC value, however, this increase was not statistically significant and there were also no significant differences in ABTS^{•+} scavenging activities.

There was a strong correlation between phenolic content as reported in sections 4.1.4.1 and 4.1.4.2 and antioxidant activity of aqueous extracts from marama beans seed coats and condensed tannin sorghum bran (Table 4.3.2). Similar Pearson's correlation coefficients between phenolic content and ABTS ($r = 0.99$) and DPPH ($r = 0.98$) have been reported in acidified methanol and aqueous 70% acetone extracts, respectively from sorghum (Dykes, Seitz, Rooney & Rooney, 2009). The results show that ORAC values were more correlated to condensed tannin content. A high correlation ($r^2 = 0.994$) between ORAC values and procyanidins levels were reported in cocoa extracts (Adamson, Lazarus, Mitchell, Prior, Cao, Jacobs, Kremers, Hammerstone, Rucker, Ritter & Schmitz, 1999).

DPPH free radical scavenging activity was strongly correlated to total phenolic content, total flavonoid content and the sum of phenolic compound concentration. Xu and Chang (2008) also found similar results where DPPH was strongly correlated to total phenolic content ($r = 0.99$) and total flavonoid content ($r = 0.98$) and ORAC values showed lower correlations to total phenolic content ($r = 0.81$) and total flavonoid ($r = 0.77$), and poor correlation to phenolic acids ($r = 0.49$) and the flavanol catechin ($r = 0.06$). The difference were explained by the authors to be due to differences in reaction mechanisms, ORAC is based on the hydrogen transfer mechanism whereas DPPH is based on the electron transfer mechanism.

ABTS^{•+} scavenging activity was strongly correlated to condensed tannin content as measured with the vanillin-HCL method. Hagerman *et al.* (1998) reported that procyanidins (tannins) were more effective quenchers of the ABTS^{•+} because of the high molecular weight and proximity of many aromatic rings and hydroxyl groups.

Table 4.3.2. Pearson's correlation coefficients between antioxidant activities and phenolic contents determined with different methods

	ORAC	DPPH	ABTS
Total phenolic content (Folin Ciocalteu method)	0.965(**)	0.995(**)	0.984(**)
Condensed tannin content (Vanillin-HCl method)	0.980(**)	0.981(**)	0.997(**)
Total flavonoid content (Colorimetric method)	0.957(**)	0.991(**)	0.985(**)
Sum of phenolic compounds (HPLC method)	0.967(**)	0.994(**)	0.984(**)

** Correlation is significant at the 0.01 level (2-tailed), N=16

4.3.4.2. Effect of extracts on AAPH-mediated red blood cell haemolysis

Aqueous extracts from marama bean seed coats inhibited AAPH-induced red blood cell haemolysis in a dose dependent manner as shown in Fig. 4.3.1. Water and acidified water extracts from marama bean seed coats at 10 µg/ml concentrations resulted in a 16 and 20% reduction in haemolysis, respectively and at 80 µg/ml concentrations reduction in haemolysis was as much as 64 and 57%, respectively compared to positive control. Acidified water extracts from marama bean seed coats had significantly lower protective effect at 40 and 80 µg/ml concentrations than water extracts which is similar to the relative phenolic contents and antioxidant activities of these samples.

The results also show significant differences between the negative control and sample controls especially the water extracts indicating that the extracts on their own were also causing red blood cell haemolysis. This may be ascribed to the presence of homogentisic acid in the extracts (section 4.1.4.2.1) which has been reported to cause haemolysis when incubated with blood (Heoedus & Nayak, 1994). Homogentisic acid is metabolized by homogentisic acid oxidase during normal catabolism of phenylalanine and tyrosine in the liver (Gutzmer, Herbst, Kiehl, Kapp & Weiss, 1997) therefore it may not be expected to cause any harm when ingested. Homogentisic acid is also present in honey used for food depending on the source of nectar (Cabras, Angioni, Tuberoso, Floris, Reniero, Guilou & Ghelli, 1999).




Figure 4.3.1. Effect of water and acidified water extracts prepared from marama bean seed coats on AAPH-mediated human red blood cell haemolysis. Each bar on the graph is an average of four determinations from two independent experiments. Error bars represent standard deviation of the mean. Bars with different letters are significantly different ($p \leq 0.05$).

Aqueous extracts from condensed tannin sorghum bran also inhibited AAPH-induced red blood cell haemolysis in a dose dependent manner as shown in Fig. 4.3.2. Treatment with water and acidified water extracts at 100 µg/ml concentrations resulted in 42 and 52% reduction in haemolysis, respectively and at 400 µg/ml concentrations by as much as 72 and 77%, respectively compared to positive control. Acidified water extracts showed a significantly higher protective effect at all concentration levels compared to water extracts

(Fig. 4.3.2). This may be due to the significantly higher total phenolic content and phenolic compound concentrations as measured with the Folin Ciocalteu method (section 4.1.4.1) and HPLC methods (section 4.1.4.2), respectively compared to water extracts. There was no significant difference between the negative control and sample controls indicating that the extracts from condensed tannin sorghum bran did not cause any haemolysis.




Figure 4.3.2. Effect of water and acidified water extracts from condensed tannin sorghum bran on AAPH-mediated human red blood cell haemolysis. Each bar on the graph is an average of four determinations from two independent experiments. Error bars represent standard deviation of the mean. Graph bars with different letters are significantly different ($p \leq 0.05$).

Extracts from marama bean seed coats were effective at lower concentrations 10–80 µg/ml (Figure 4.3.1) compared to extracts from condensed tannin sorghum bran which were effective at 100–400 µg/ml (Fig. 4.3.2). This is because extracts from marama bean seed

coats had higher phenolic content and antioxidant activity compared to equivalent extracts from condensed tannin sorghum bran.

Phenolic compounds from olive oil (Paiva-Martins, Fernandes, Santos, Silval, Borges, Rocha, Belo, & Bogdanov, 2010) and green tea (Lanping, Zaiqun, Bo, Li & Zhongli, 2000) showed a similar dose response protective effect against H₂O₂ and AAPH-induced red blood cell haemolysis, respectively. The protective effect of phenolic compounds in the extracts against red blood cell biomembrane oxidation is through antioxidant activity (Paiva-Martins *et al.*, 2009) by reducing lipid peroxides (Lin, Wu, Wang, Yang & Chang, 2001). Condensed tannins and flavonoid compounds in the extracts may also provide protection by accumulating at the membrane surface through hydrogen bonding to polar groups of membrane phospholipids thus preventing access of the deleterious peroxy radicals to the hydrophobic region of the bilayer (Verstraeten, Keen, Schmitz, Fraga & Oteiza, 2003). The results suggest that phenolic compounds in the extracts have a potential to reduce oxidative stress on cell biomembranes which are the main sites of lipid peroxidation (Mak *et al.*, 1983). Low molecular weight proanthocyanidins in the extracts, such as dimers, may be absorbed in the upper gastrointestinal tract in their intact form (Holt, Lazarus, Sullards, Zhu, Schramm, Hammerstone, Fraga, Schmitz & Keen, 2002). However, high molecular weight condensed tannins (oligomers and polymers) may not be absorbed into circulation in their intact form (Rios, Gonthier, Rémésy, Mila, Lapierra, Lazarus, Williamson & Scalbert, 2003). They may therefore only directly provide protection to cells lining the gastrointestinal tract against oxidative damage and may also play a unique role in sparing other antioxidants and thus indirectly increasing antioxidant levels in other tissues (Hagerman *et al.*, 1998). However, the high molecular weight proanthocyanidins may be metabolised into phenolic acid metabolites by colonic bacteria and these metabolites may be absorbed into circulation and therefore may confer health beneficial effects (Rios *et al.*, 2003; Saura-Calixto, Pérez-Jiménez, Touriño, Serrano, Fuguet, Torres & Goñi, 2010).

4.3.4.3. Effect of aqueous extracts on AAPH-mediated oxidative DNA damage

The effects of water and acidified water extracts from marama bean seed coats and condensed tannin sorghum bran on AAPH-induced oxidative supercoiled plasmid pBR 322 DNA damage are shown in Fig. 4.3.3.




Figure 4.3.3. Effect of water and acidified water extracts from marama bean seed coats and condensed tannin sorghum bran on AAPH-induced oxidative supercoiled plasmid pBR 322 DNA damage. The S band represents supercoiled DNA and R band represents relaxed circular DNA.

Supercoiled DNA shown as S band had higher mobility than relaxed circular DNA shown as R band. In the negative control sample (without AAPH) the intensity of the super-coiled DNA band (S) was set to 100% and the intensities of the other bands were calculated relative to this band. Addition of 11 µmol AAPH solution to super-coiled plasmid DNA (positive control) caused single strand breaks resulting in decrease in intensity of the supercoiled DNA

band (S) to a mean value of 86 % of negative control and there was a slight increase in intensity of the relaxed circular DNA band (R) (Fig. 4.3.3).




Figure 4.3.4. Effect of water and acidified water extracts from marama bean seed coats on AAPH-induced oxidative supercoiled plasmid pBR 322 DNA damage. Each bar on the graph is an average of four determinations from two independent experiments. Error bars represent standard deviation of the mean. Graph bars with different letters are significantly different ($p \leq 0.05$).

Addition of 5 µmol Trolox solution (5 µm Trolox + AAPH) resulted in protection against APPH-mediated single strand breaks and the intensity of the supercoiled DNA band was not significantly different from the negative control. Treatment with both water and acidified extracts from marama bean seed coats affected the mobility of DNA. All the lanes treated with extracts from marama bean seed coats (8 µg/ml) left behind a trail of DNA from the

wells resulting in lower intensities of the R and S bands especially the water extracts with higher levels of condensed tannin content (Fig. 4.3.3). However, all lanes treated with aqueous extracts from condensed tannin sorghum (200 µg/ml) did not show this trail of DNA.

Water and acidified water extracts from marama bean seed coats did not appear to provide protection against AAPH-mediated oxidative supercoiled DNA damage because there was no significant difference between the positive control and samples treated with the aqueous extracts (Fig. 4.3.4). However, it would appear that these results were inconclusive because there was poor migration of the DNA bands through the gel leaving behind a trail. The interference in the lanes treated with extracts from marama bean seed coats could be as a result of an interaction between DNA and the galloylated condensed tannins (section 4.2.5.3.1) in the extracts possibly through hydrogen bonding to form a complex with a higher molecular mass and low mobility. Condensed tannins in plant tissue have been reported to bind to DNA upon cell lyses during isolation of DNA for research work (Maliyakal, 1992). Also hydrolysable tannins have been reported to bind to DNA from calf thymus and the binding was attributed to the large number of hydroxyl groups and the high molecular weight (Labieniec & Gabryelak, 2006).

Treatment with water and acidified water extract solutions (200 µg/ml) from condensed tannin sorghum bran resulted in super-coiled DNA band (S) intensities of 96 and 97 %, respectively (Fig. 4.3.5). There was no significant difference between these bands and the negative control and the 5 µM Trolox treated sample indicating that both extracts from condensed tannin sorghum bran provided some protection against AAPH-mediated single strand breaks.




Figure 4.3.5. Effect of water and acidified water extracts (200 µg/ml) from condensed tannin sorghum bran on AAPH-induced oxidative supercoiled plasmid pBR 322 DNA damage. Each bar on the graph is an average of four determinations from two independent experiments. Error bars represent standard deviation of the mean. Graph bars with different letters are significantly different ($p \leq 0.05$).

The phenolic compounds epigallocatechin-3-O-gallate, epigallocatechin and epicatechin isolated from green tea were reported to have a protective effect against AAPH-induced supercoiled plasmid vector DNA (pBR 322) oxidative damage (Wei *et al.*, 2006). In another study the phenolic compounds caffeic acid, hydroxytyrosol and tyrosol from olive oil were also reported to have a protective effect against H₂O₂ induced oxidative DNA damage in cell lines from human prostate epithelium in a dose dependent manner (Quiles, Farquharson, Simpson, Grant & Wahle, 2002). The protective effect of phenolic compounds against oxidative DNA damage may be through free radical scavenging activity and transition metal ion chelation (Aherne & O'Brien, 2000). The results of this study suggest that the extracts by

virtue of their phenolic compound content may play a role in the prevention of cancer which may be initiated by free radical oxidative DNA damage.

4.3.4.4. Effect of extracts on copper-catalyzed human LDL oxidation

In this assay LDL oxidation was initiated with Cu (II) and the breakdown products of lipid peroxides, thiobarbituric reactive substances (TBARS) which are the late stage products of LDL oxidation (Xu *et al.*, 2007) were measured at the end of the incubation period using the TBA assay. Incubation of human LDL with copper sulphate solution (positive control) caused oxidation of LDL and produced maximum TBARS (Fig. 4.3.6 and Fig. 4.3.8).




Figure 4.3.6. Effect of water and acidified water extracts prepared from marama bean seed coats on copper-catalyzed LDL oxidation determined with the thiobarbituric (TBA) assay. Bars are means of four determinations from two independent experiments. Error bars represent standard deviation of the mean. Bars with different letters are significantly different ($p \leq 0.05$).

Addition of 100 μM Trolox reduced LDL oxidation by 54% of positive control. Treatment with water and acidified water extracts from marama bean seed coats at 10 and 40 $\mu\text{g}/\text{ml}$ concentrations resulted in 11–17% reduction in TBARS levels, however the TBARS levels were not significantly different from the positive control (Fig. 4.3.6). Treatment with water and acidified water extracts at 80 $\mu\text{g}/\text{ml}$ concentrations resulted in 73 and 75% significant reduction in TBARS levels, respectively. The reduction was higher than that of 100 μM Trolox and the levels of TBARS were not significantly different from that of the negative control (without AAPH) which suggests that there was almost complete inhibition of LDL oxidation. There were no significant differences in the level of inhibition between water and acidified water extracts at all concentrations even though there were significant differences in phenolic compound content and antioxidant activity between the extracts. In Fig. 4.3.7 a possible explanation is provided as to why extracts with different levels of phenolic contents could exhibit similar TBARS levels. LDL oxidation is a lipid peroxidation reaction which is characterized by three phases; initiation (lag phase), propagation and decomposition. Antioxidants such as phenolic compounds lower the rate of initiation by scavenging lipid peroxy radicals (Abuja, Murkovic & Pfannhauser, 1998) thereby prolonging the lag phase during incubation (Esterbauer, Gebicki, Puhl, & Jürgens, 1992). The extracts at 10 and 40 $\mu\text{g}/\text{ml}$ concentrations had short lag phases and allowed the reaction to go through all the three lipid peroxidation phases resulting in similar TBARS levels. Both extracts at 80 $\mu\text{g}/\text{ml}$ concentrations suppressed the initiation phase through-out the incubation period resulting in similar low TBARS levels.

Similar results were reported in an experiment investigating the protective effect of an antioxidant compound, oxime against copper-catalyzed LDL oxidation (de Lima Portella *et al.*, 2008). The authors found that at low concentrations levels (0.5, 1, 2 and 3 μM) the TBARS levels were not significantly different amongst treatments and the positive control even though the lag phases as measured by conjugated diene formation were significantly different amongst treatments and the positive control at all concentrations except at 0.5 μM . However at 5 μM there was a significant reduction in TBARS compared to the other treatments and the positive control.




Figure 4.3.7. LDL oxidation kinetic curves showing how two extracts with significantly different antioxidant concentration could exhibit similar TBARS levels (Adapted from de Lima Portella *et al.* (2008)).

Addition of water and acidified water extracts from condensed tannin sorghum bran also showed a similar trend as extracts from marama bean seed coats (Fig. 4.3.8). Treatment with 100 and 250 µg/ml extract concentrations resulted in a 12–20% reduction in LDL oxidation and the TBARS levels at these concentrations were significantly different from that of the positive control. Treatment with water and acidified water extracts at 500 µg/ml concentration resulted in 72 and 71% significant reduction in TBARS levels, respectively and the levels were not significantly difference from that of the negative control (without AAPH) indicating an almost complete inhibition. The inhibition at this level was higher than that of 100 µm Trolox treated samples.




Figure 4.3.8. Effect of aqueous extracts prepared from condensed tannin sorghum bran on copper-catalyzed LDL oxidation using thiobarbituric assay (TBA assay). Bars are means of four determinations from two independent experiments. Error bars represent standard deviations. Bars with the same letter are not significantly different ($p \leq 0.05$).

Phenolic compounds protect against LDL oxidation by lowering the rate of initiation by scavenging lipid peroxy radicals (Abuja *et al.*, 1998) interrupting propagation of free radicals and thus acting as chain breaking antioxidants (Reiter, 1998). These compounds may also provide protection through chelation of copper ions and by stabilizing LDL structure through interaction with apolipoprotein B thereby preventing binding of copper ions to the particles (Rüfer & Kulling, 2006).

The results show that water and acidified water extracts from marama bean seed coats were more effective than equivalent extracts from condensed tannin sorghum bran. This effect was observed at a concentration of 10–80 µg/ml for extracts from marama bean seed coats compared to 100–500 µg/ml for extracts from condensed tannin sorghum bran. This is due to

the higher phenolic content and antioxidant activity of extracts from marama bean seed coats compared to equivalent extracts from condensed tannin sorghum bran.

The results did not show a dose-response effect. Also there were no significant differences between water and acidified water extract treated samples, especially samples treated with extracts from marama bean seed coat which had significantly different phenolic content and antioxidant activity. The reason is that antioxidants only prolong the lag phase of lipid peroxidation during the incubation period (Abuja *et al.*, 1998; Esterbauer, Gebicki, Puhl & Jürgens, 1992) and if the reaction goes through all the phases of lipid peroxidation the amount of TBARS produced becomes dependent on the initial amount of LDL present and not on the extract concentration.

Other plant extracts from elderberry (Abuja *et al.*, 1998), grape juice (Frankel, Bosanek, Meyer, Silliman & Kirk, 1998) have been shown to inhibit LDL oxidation *in vitro*. Purified phenolic compounds extracted from plant material such as hydroxycinnamic acids from grape (Meyer, Donovan, Pearson, Waterhouse & Frankel, 1998), isoflavonoids and their metabolites from soybean (Hodgson *et al.*, 1996; Rüfer & Kulling, 2006) have also been shown to have a protective effect against LDL oxidation. Overall, the results provide some evidence that extracts from marama bean seed coats and condensed tannin sorghum bran, by virtue of their content of phenolic compounds could provide protection against LDL oxidation, which is a risk factor in cardiovascular disease.

4.3.5. Conclusions

Aqueous extracts from marama bean seed coats have higher free radical scavenging activity than equivalent extracts from condensed tannin sorghum bran because of their higher phenolic compound content. Extraction of marama bean seed coats under acidic condition results in an extract with lower free radical scavenging activity because of lower phenolic content as a result of co-precipitation of phenolic compounds with interpolymer complex precipitate which may be as a result of interaction between prodelphinidins (condensed tannins) and cell wall polysaccharides. In contrast, extraction of condensed tannin sorghum bran under acidic condition does not have a significant effect on free radical scavenging activity, probably because of the similar levels of condensed tannins in water and acidified water extracts.

Extracts from marama bean seed coats are effective at lower concentration against biomembrane oxidative damage than equivalent extracts from condensed tannin sorghum bran due to their higher phenolic compound content and antioxidant activity. Extracts from condensed tannin sorghum bran show a clear protective effect against oxidative DNA damage. However, this assay is not suitable for the evaluation of extracts from marama bean seed coats as the prodelphinidins in the extracts bind to DNA causing poor mobility of the bands in the agarose gel resulting in inconclusive results. The extracts have a potential to lower the risk of cancer by inhibiting free radical oxidative DNA damage in cells. Furthermore, these extracts can potentially prevent atherosclerosis associated with cardiovascular disease by inhibiting LDL oxidation. Aqueous extracts from marama bean may be more effective at lower concentration than those from condensed tannin sorghum bran because of their higher concentration of phenolic compounds. In conclusion, both marama bean seed coat and condensed tannin sorghum bran extracts have the potential to protect biomolecules against oxidative damage.