

## Bibliography

- Aben, I., Verhey, F., Lousberg, R., Lodder, J. & Honig, A. (2002). Validity of the Beck Depression Inventory, Hospital Anxiety and Depression Scale, SCL-90, and Hamilton Depression Rating Scale as screening instruments for depression in stroke patients. *Psychosomatics*, 43(5):386-393.
- Ada, L., Dean, C.M., Lindley, R. & Lloyd, G. (2009). Improving community ambulation after stroke: the AMBULATE trial. *BioMedCentral Neurology*, 9:8.
- Agrell, B. & Dehlin, O. (2000). Mini-Mental State Examination in geriatric stroke patients. Validity, differences between subgroups of patients, and relationships to somatic and mental variables. *Aging (Milano)*, 12(6):439-444.
- Antonucci, G., Guariglia, C., Judica, A., Magnotti, L., et al. (1995). Effectiveness of neglect rehabilitation in a randomized group study. *American Psychological Association*, 17(3):383-389.
- Aström, M. (1996). Generalized anxiety disorder in stroke patients. A 3-year longitudinal Study. *Stroke*, 27:270-275.
- Bailey, M.J., Riddoch, M.J., Crome, P. (2000). Evaluation of a test battery for hemineglect in elderly stroke patients for use by therapists in clinical practice. *NeuroRehabilitation*, 14:139-150.

- Bailey, M.J., Riddoch, M.J. & Crome, P. (2002). Treatment of visual neglect in elderly patients with stroke: A single-subject series using either a scanning or cueing strategy or a left-limb activation strategy. *Physical Therapy*, 82(8):782-797.
- Bailey, M.J., Riddoch, M.J., Crome, P. (2004). Test-retest stability of three tests for unilateral visual neglect in patients with stroke: Star Cancellation, Line Bisection, and the Baking Tray Task. *Neuropsychological Rehabilitation*, 14(4):403-419.
- Ball, K.K., Beard, B.L., Roenker, D.L., Miller, R.L. & Griggs, D.S. (1988). Age and visual search: expanding the useful field of view. *The Journal of the Optical Society of America A*, 5(12):2210-2219.
- Barrett, B. T. (2009). A critical evaluation of the evidence supporting the practice of behavioural vision-therapy. *Ophthalmic and Physiological Optics : The Journal of the British College of Ophthalmic Opticians (Optometrists)*, 29:4-25
- Blanton, S., Morris, D.M., Prettyman, M.G. McCulloch, K., Redmond, S., Light, K.E & Wolf, S.L. (2006). Lessons learned in participant recruitment and retention: The EXCITE Trial. *Physical Therapy*, 86 (11):1520-1533.
- Ble, A., Volpato, S., Zuliani, G., et al. (2005). Executive function correlates with walking speed in older persons: The InCHIANTI study. *Journal of the American Geriatrics Society*, 53:410-415.

- Bolognini, N., Rasi, F., Coccia, F. & Làdavas, E. (2005). Visual search improvement in hemianopic patients after audio-visual stimulation. *Brain*, 128:2830-2842.
- Bouwmeester, L., Heutnik, J. & Lucas, C. (2007). The effect of visual training for patients with visual field defects due to brain damage: a systematic review. *Journal of Neurology, Neurosurgery and Psychiatry*, 78:555-564.
- Bowen, A. & Lincoln, N.B. (2007). Cognitive rehabilitation for spatial neglect following stroke (Review). *The Cochrane Library*, 4:1-43.
- Brown, M.M., Brown, G.C., Sharma, S. & Busbee, B. (2003). Quality of life associated with visual loss. A time tradeoff utility analysis comparison with medical health states. *Ophthalmology*, 110:1076-1081.
- Brunila, T., Lincoln, N., Lindell, A., Tenovuo, O. & Hämäläinen, H. (2002). Experiences of combined visual training and arm activation in the rehabilitation of unilateral visual neglect: A clinical study. *Neuropsychological Rehabilitation*, 12(1):27-40.
- Burvill, P.W., Johnson, G.A., Jamrozik, K.D., Anderson, C.S., Stewart-Wynne, E.G. & Chakera, T.M. (1995). Prevalence of depression after stroke: The Perth community stroke study. *British Journal of Psychiatry*, 166:320-327.

- Cahn-Weiner, D.A., Boyle, P.A. & Malloy, P.F. (2002). Tests of executive function predict instrumental activities of daily living in community-dwelling older individuals. *Applied Neuropsychology*, 9(3):187-191.
- Cappa, S.F., Benke, T., Clarke, S., Rossi, B., Stemmer, B. & van Heugten, C.M. (2003). EFNS guidelines on cognitive rehabilitation: Report of an EFNS task force. *European Journal of Neurology*, 10:11-23.
- Carter, L.T., Howard, B.E. & O'Neil, W.A. (1983). Effectiveness of cognitive skill remediation in acute stroke patients (visual scanning, visual spatial, time judgement, retraining). *American Journal of Occupational Therapy*, 37(5):320-326.
- Carter, L.T., Oliveira, D.O., Duponte, J. & Lynch, S.V. (1988). The relationship of cognitive skills performance to activities of daily living in stroke patients. *American Journal of Occupational Therapy*, 42(7):449-455.
- Cassidy, T.P., Lewis, S. & Gray, C.S. (1998). Recovery from visuospatial neglect in stroke patients. *Journal of Neurology, Neurosurgery and Psychiatry*, 64(4):555-557.
- Chaiken, L. R. (2007). Chapter 30. Disorders of vision and visual-perceptual dysfunction. (In Humphred, D. A., ed. *Neurological Rehabilitation*. 5th ed. Maryland Heights, MO : Mosby. p. 973-1004.)

- Chan, D.Y.L., Chan, C.C.H. & Au, D.K.S. (2006). Motor relearning programme for stroke patients: a randomised controlled trial. *Clinical Rehabilitation*, 20:191-200.
- Cherney, L.R., Halper, A.S., Kwasnica, C.N., Harvey, R.L. & Zhang, M. (2001). Recovery of functional status after right hemisphere stroke: Relationship with unilateral neglect. *Archives of Physical Medicine and Rehabilitation*, 82:322-328.
- Cicerone, K.D., Dahlberg, C., Kalmar, K., Langenbahn, D.M., Malec, J.F., Bergquist, T.F., et al. (2000). Evidence-based cognitive rehabilitation: Recommendations for clinical practice. *Archives of Physical Medicine and Rehabilitation*, 81:1596-1615.
- Cicerone, K.D., Dahlberg, C., Malec, J.F., Langenbahn, D.M., Felicetti, T., Kneipp, S., Ellmo, W., Kalmar, K., et al. (2005). Evidence-based cognitive rehabilitation: Updated review of the literature from 1998 through 2002. *Archives of Physical Medicine and Rehabilitation*, 86:1681-1692.
- Ciuffreda, K.J. (2002). The scientific basis for efficacy of optometric vision therapy in nonstrabismic accommodative and vergence disorders. *Optometry*, 73(12):735.
- Ciuffreda, K.J., Kapoor, N., Rutner, D., Suchoff, I.B., Han, M.E. & Craig, S. (2007). Occurrence of oculomotor dysfunctions in acquired brain injury: A retrospective analysis. *Optometry*, 78:155-161.

- Crum, R.M., Anthony, J.C., Bassett, S.S., Folstein, M.F. (1993). Population-based norms for the Mini-Mental State Examination by age and educational level. *Journal of the American Medical Association*, 18:2386-2391.
- Dam, H., Pedersen, H.E. & Ahlgren, P. (1989). Depression among patients with stroke. *Acta Psychiatrica Scandinavica*, 80:118-124.
- Das, A. & Huxlin, K. R. (2010). New approaches to visual rehabilitation for cortical blindness: Outcomes and putative mechanisms. *The Neuroscientist*, 16(4):374-387.
- de Seze, M., Wiart, L., Bon-Saint-Come, A., Debelleix, X., de Seze, M., Joseph, P.A., et al. (2001). Rehabilitation of postural disturbances of hemiplegic patients by using trunk control retraining during exploratory exercises. *Archives of Physical Medicine and Rehabilitation*, 82(6):793-800.
- Duncan, P.W., Wallace, D., Lai, S.M., Johnson, D., Embretson, S. & Laster, L.J. (1999). The Stroke Impact Scale Version 2.0. Evaluation of reliability, validity and sensitivity to change. *Stroke*, 30:2131–2140.
- Duncan, P.W., Lai, S.M., Tyler, D., Perera, S., Reker, D.M. & Studenski, S. (2002a). Evaluation of proxy responses to the Stroke Impact Scale. *Stroke*, 33:2593-2599.

- Duncan, P.W., Bode, R.K., Lai, S.M., Perera, S. (2003b). Rasch analysis of a new stroke-specific outcome scale: The Stroke Impact Scale. *Archives of Physical Medicine and Rehabilitation*, 84:950-963.
- Edwards, B. & O'Connell, B. (2003). Internal consistency and validity of the Stroke Impact Scale 2.0 (SIS 2.0) and SIS-16 in an Australian sample. *Quality of Life Research*, 12:1135.
- Egelko, S., Simon, D., Riley, E., et al. (1989). First year after stroke: Tracking cognitive and affective deficits. *Archives of Physical Medicine and Rehabilitation*, 70:297-302.
- Evers, S., Nabavi, D., Rahmann, A., Heese, C., Reichelt, D. & Husstedt, I.W. (2003). Ischaemic cerebrovascular events in HIV infection: A cohort study. *Cerebrovascular Diseases*. 15(3):199-205.
- Fanthome, Y., Lincoln, N.B., Drummond, A.E. R. & Walker, M.F. (1995). The treatment of visual neglect using feedback of eye movements: A pilot study. *Disability and Rehabilitation*, 17(8):413-417.
- Flansbjerg, U., Holmback, A.M., Downham, D., Patten, C. & Lexell, J. (2005). Reliability of gait performance tests in men and women with hemiparesis after stroke. *Journal of Rehabilitation Medicine*, 37:75-82.

Folstein, M.F., Folstein, S.E., McHugh, P.R. (1975). "Mini-Mental State". A practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research*, 12(3):189-198.

Folstein, M.F., Folstein, S.E., McHugh, P.R. & Fanjiang, G. (2001). *Mini-Mental State Examination User's Guide*. Odessa, FL : Psychological Assessment Resources.

Galetta, K.M., Barrett, J., Allen, M., et al. (2011). The King-Devick Test as a determinant of head trauma and concussion in boxers and MMA fighters. *Neurology*, 76:1-7.

Gilhotra, J.S., Mitchell, P., Healey, P.R., Cumming, R.G. & Currie, J. (2002). Homonymous visual field defects and stroke in an older population. *Stroke*, 33:2417-2420.

Goh, K. (2007). Vision restoration therapy: New hope for stroke for patients with visual field loss. *ANNALS Academy of Medicine Singapore*, 36(Suppl):72-76.

Gordan, W.A., Hibbard, M.R., Egelko, S., Diller, L., Shaver, M.S., Lieberman, A. & Ragnarsson, K. (1985). Perceptual remediation in patients with right brain damage: A comprehensive program. *Archives of Physical Medicine and Rehabilitation*, 66:353-359.

Gorman, S.L. (2007). Chapter 3. Contemporary issues and theories of motor control, motor learning and neuroplasticity: Assessment of movement and



posture. (*In* Humphred, D. A., ed. *Neurological Rehabilitation*. 5th ed. Maryland Heights, MO : Mosby. p. 56-70.)

Gur, S. & Ron, S. (1992). Training in oculomotor training: Occupational health aspects. *Israel Journal of Medical Sciences*, 28(8-9):622-628.

Hafsteinsdóttir, T.B., Algra, A., Kappelle, L.J., & Grypdonck, M.H.F. (2005). Neurodevelopmental treatment after stroke: A comparative study. *Journal of Neurology, Neurosurgery and Psychiatry*, 76:788-792.

Halligan, P.W., Fink, G.R., Marshall, J.C. & Vallar, G. (2003). Spatial cognition: Evidence from visual neglect. *Trends in Cognitive Sciences*, 7(3):125-133.

Hammell, K.W. & Carpenter, C. (2004). *Qualitative research in evidence-based rehabilitation*. Philadelphia, PA. 148 p.

Heller, R.F., Langhorne, P., & James, E. (2000). Improving stroke outcome: The benefits of increasing availability of technology. *Bulletin of the World Health Organization*, 78(11):1337-1343.

Hinojosa, M.S. & Rittman, M. (2009). Association between health education needs and stroke caregiver injury. *Journal of Aging and Health*, 21:1040-1058.

Holtzer, R., Verghese, J., Xue, X. & Lipton, R.B. (2006). Cognitive processes related to gait velocity: Results from the Einstein aging study. *Neuropsychology*, 20(2):215-223.

- Horak, F.B., Henry, S.M. & Shumway-Cook, A. (1997). Postural perturbations: New insights for treatment of balance disorders. *Physical Therapy*, 77:517-533.
- Horgan, N.F., Cunningham, C.J., Coakley, D., Walsh, J.B., O'Regan, M & Finn, A.M. (2006). The Stroke Activity Scale: Results of a validity study. *Disability and Rehabilitation*, 28(15):937-941.
- Hsueh, I.P., Lee, M.M & Hsieh, C.L. (2001). Psychometric characteristics of the Barthel activities of daily living index in stroke patients. *Journal of the Formosan Medical Association*, 100(8):526–532.
- Hung, K.G. & Ciuffreda, K.J. (2002). Models of the visual system. New York: Springer. 781 p.
- International Bobath Instructors Training Association (IBITA) – An international association for adult neurological rehabilitation. Theoretical Assumptions and Clinical Practice (IBITA Annual General Meeting, September 2006), January 2007.
- Jaillard, A., Grand, S., Le Bas, J.F. & Hommel, M. (2010). Predicting cognitive dysfunctioning in nondemented patients early after stroke. *Cerebrovascular Diseases*, 29(5):415-423.
- Jobke, S., Kasten, E. and Sabel, B.A. (2009). Vision restoration through extrastriate stimulation in patients with visual field defects – a double-blind and randomized experimental study. *Neural Rehabilitation and Neural Repair*, 23:246-55.

- Jones, R.N. & Gallo, J.J. (2000). Dimensions of the Mini-Mental State Examination among community dwelling older adults. *Psychological Medicine*, 30:605-618.
- Jones, S.A. & Shinton, R.A. (2006). Improving outcome in stroke patients with visual problems. *Age and Ageing*, 35(6):560–565.
- Kalaria, R.N. & Ballard, C. (2001). Stroke and cognition. *Current Atherosclerosis Reports*, 3(4):334-339.
- Kalra, L., Perez, I., Gupta, S. & Wittink, M. (1997). The influence of visual neglect on stroke rehabilitation. *Stroke*. 28:1386-1391.
- Kerkhoff, G., MünBinger, U., Eberle-strauss, G & Stögerer, E. (1992). Rehabilitation of hemianopic alexia in patients with postgeniculate visual field disorders. *Neuropsychological Rehabilitation*, 2(1):21-42.
- Kerkhoff, G., MünBinger, U. & Meier, E. K. (1994). Neurovisual rehabilitation in cerebral blindness. *Archives of Neurology*, 51(5):474-481.
- Kerkhoff, G. (2000). Multiple perceptual distortions and their modulation in left sided visual neglect. *Neuropsychologia*, 38(7):1073-1086.
- Kerkhoff, G. (2000). Neurovisual rehabilitation: Recent developments and future directions. *Journal of Neurology, Neurosurgery and Psychiatry*, 68:691-706.
- Làdavas, E, Menghini, G. & Umiltà, C. (1994). A rehabilitation study of hemispatial neglect. *Cognitive Neuropsychology*, 11(1):75-95.

- Land, M., Mennie, N. & Rusted, J. (1999). The role of vision and eye movements in the control of activities of daily living. *Perception*, 28:1311-1328.
- Land, M.F. (2009). Vision, eye movements, and natural behaviour. *Visual Neuroscience*, 26:51-62.
- Langhorne, P., Coupar, F. & Pollock, A. (2009). Motor recovery after stroke: A systematic review. *The Lancet Neurology*, 8(8):741-754.
- Lannin, N.A., & Herbert, R.D. (2003). Is hand splinting effective for adults following stroke? A systematic review and methodological critique of published research. *Clinical Rehabilitation*, 17(8):807-816.
- Leigh, R.J. & Kennard, C. (2004). Using saccades as a research tool in the clinical neurosciences. *Brain*, 127:460-477.
- Lennon, S., Ashburn, A., & Baxter, D. (2006). Gait outcome following outpatient physiotherapy based on the Bobath concept in people post stroke. *Disability and Rehabilitation*, 28(13-14):873-881.
- Linden, T., Samuelsson, H., Skoog, I. & Blomstrand, C. (2005). Visual neglect and cognitive impairment in elderly patients late after stroke. *Acta Neurologica Scandinavica*, 111(3):163-168.

- Liu-Ambrose, T., Pang, M.Y. & Eng, J.J. (2007). Executive function is independently associated with performances of balance and mobility in community-dwelling older adults after mild stroke: implications for falls prevention. *Cerebrovascular Diseases*, 23:203–210.
- Luauté, J., Halligan, P., Rode, G., Rossetti, Y. & Boisson, D. (2006). Visuo-spatial neglect: A systematic review of current interventions and their effectiveness. *Neuroscience and Biobehavioral Reviews*, 30(7):961-982.
- Maddock, R.J., Millodot, M., Leat, S. & Johnson, C.A. (1981). Accommodation responses and refractive error. *Investigative Ophthalmology and Visual Sciences*, 20(3):387-388.
- Magill, R.A. (2004). Motor learning and control: Concepts and applications. 7<sup>th</sup> ed  
New York : McGraw-Hill College. 385 p.
- Mahoney, F.I. & Barthel, D.W. (1965). Functional evaluation: The Barthel Index. *Maryland State Medical Journal*, 14:61-65.
- Markowitz, S.N. (2006). Principles of modern low vision rehabilitation. *Canadian Journal of Ophthalmology*, 41(3):289–312.
- Martin T. & Huxlin K.R. (2010). Spontaneous and training-induced visual learning in cortical blindness: Characteristics and neural substrates. *Topics in Cognitive Science*, 2(2):306-319, May.

- McCullagh, E., Brigstocke, G., Donaldson, N. & Kalra L. (2005). Determinants of caregiving burden and quality of life in caregivers of stroke patients. *Stroke*, 36(10):2181-2186.
- Menon, A. & Korner-Bitensky, N. (2004). Evaluating unilateral spatial neglect post-stroke. Working your way through the maze of assessment choices. *Topics in Stroke Rehabilitation*, 11(3):41-66.
- Mesulam, M. (1999). Spatial attention and neglect: Parietal, frontal and cingulate contributions to the mental representation and attentional targeting of salient extrapersonal events. *Philosophical Transactions of the Royal Society of London*, 354:1325-1346.
- Mochan, A., Modi, M. & Modi, G. (2003). Stroke in black South African HIV positive patients: A prospective analysis. *Stroke*, 34:10-15.
- Montgomery, S.A. & Asberg, M. (1979). A new depression scale designed to be sensitive to change. *British Journal of Psychiatry*, 134:382-389.
- Mueller, I., Mast, H. & Sabel, B.A. (2007). Recovery of visual field defects: A large clinical observational study using vision restoration therapy. *Restorative Neurology and Neuroscience*, 25(5-6):563-72.
- Nelles, G., Esser, J., Eckstein, A., Tiede, A., Gerhard, H. & Diener, H.C. (2001). Compensatory visual field training for patients with hemianopia after stroke. *Neuroscience Letters*, 306(3):189-192.

Nelles, G., Pscherer, A., de Greiff, A., Forsting, M., Gerhard, H., Esser, J & Diener, H.C. (2009). Eye-movement training-induced plasticity in patients with post-stroke hemianopia. *Journal of Neurology*, 256(5):726-733.

Ng, S.S. & Hui-Chan, C.W. (2005). The Timed Up and Go Test: Its reliability and association with lower – limb impairments and locomotor capacities in people with chronic stroke. *Archives of Physical Medicine and Rehabilitation*, 86(8):1641-1647.

Niemeier, J.P. (1998). The Lighthouse Strategy: Use of a visual imagery technique to treat visual inattention in stroke patients. *Brain Injury*, 12(5):399-406.

Paci, M. (2003). Physiotherapy based on the Bobath concept for adults with post-stroke hemiplegia: A review of effectiveness studies. *Journal of Rehabilitation Medicine*, 35:2-7.

Pambakian, A.L.M. & Kennard, C. (1997). Can visual function be restored in patients with homonymous hemianopia? *British Journal of Ophthalmology*, 81(4):324-328.

Pambakian, A.L.M., Mannan, S.K. & Hodgson, T.L., et al. (2004). Saccadic visual search training: a treatment for patients with homonymous hemianopia. *Journal of Neurology, Neurosurgery and Psychiatry*, 75:1443-1448.

- Paolucci, S., Antonucci, G., Guariglia, C., Magnotti, L., Pizzamiglio, L. & Zoccolotti, P. (1996). Facilitatory effect of neglect rehabilitation on the recovery of left hemiplegic stroke patients: A cross-over study. *Journal of Neurology*, 243:308-314.
- Paolucci, S., Antonucci, G., Grasso, M. G. & Pizzamiglio, L. (2001). The role of unilateral spatial neglect in rehabilitation of right brain-damaged ischemic stroke patients: A matched comparison. *Archives of Physical Medicine and Rehabilitation*, 82(6):743-749.
- Patterson, K.K., Gage, W.H., Brooks, D., Black, S.E. & McIlroy, W.E. (2010). Changes in gait symmetry and velocity after stroke: A cross-sectional study from weeks to years after stroke. *Neurorehabilitation and Neural Repair*, 24(9):783-790.
- Perry, J., Garrett, M., Gronley, J.K. & Mulroy, S.J. (1995). Classification of walking handicap in the stroke population. *Stroke*, 26: 982-989.
- Pierce, S. & Buxbaum, L.J. (2002). Treatments of unilateral neglect: A review. *Archives of Physical Medicine and Rehabilitation*, 83:256-268.
- Pierce, L.L., Steiner, V., Govoni, A., Thompson, T.C. & Friedemann, M.L. (2007). Two sides to the caregiving story. *Topics in Stroke Rehabilitation*, 14(2):13-20.
- Pizzamiglio, L., Antonucci, G., Judica, A., Joseph, P.A., Montenero, P., Razzano, C., et al. (1992). Cognitive rehabilitation of hemineglect disorder in chronic patients



with unilateral right brain damage. *Neuropsychology, Development and Cognition. Section A, Journal of Clinical and Experimental Neuropsychology*, 14(6):901-923.

Pizzamiglio, L., Fasotti, L., Jehkonen, M., Antonucci, G., Magnotti, L., Boelen, D. & Asa, S. (2004). The use of optokinetic stimulation in the rehabilitation of the hemineglect disorder. *Cortex*, 40:441-450.

Podsiadlo, D. & Richardson, S. (1991). The Timed "Up & Go": A test of basic functional mobility for frail elderly persons. *Journal of the American Geriatrics Society*, 39:142-148.

Poggel, D.A., Kasten, E. & Sabel, B.A. (2004). Attentional cueing improves vision restoration therapy in patients with visual field defects. *Neurology*, 63:2069-2076.

Pollock, A., Baer, G., Pomeroy, V. & Langhorne, P. (2007). Physiotherapy treatment approaches for the recovery of postural control and lower limb function following stroke (Review). *The Cochrane Library (Issue 4)*. John Wiley & Sons, Ltd.

Rawstron, J.A., Burley, C.D. & Elder, M.J. (2005). A systematic review of the applicability and efficacy of eye exercises. *Journal of Pediatric Ophthalmology and strabismus*, 42(2):82-88.

- Reinhard, J., Schreiber, A., Schiefer, A., Kasten, E., Sabel, B.A., Kenkel, S., Vonthein, R. & Trauzettel-Klosinski, S. (2005). Does visual restitution training change absolute homonymous visual field defects? A fundus controlled study. *British Journal of Ophthalmology*, 89:30-35.
- Robertson, I.H., McMillan, T.M., MacLeod, E., Edgeworth, J. & Brock, D. (2002). Rehabilitation by limb activation training reduces left-sided motor impairment in unilateral neglect patients: A single-blind randomised control trial. *Neuropsychological Rehabilitation*, 12 (5):439-454.
- Roth, T., Sokolov, A.N., Messias, A., Roth, P., Weller, M. & Trauzettel-Klosinski S. (2009). Comparing explorative saccade and flicker training in hemianopia: A randomised controlled study. *Neurology*, 72:324-331.
- Sabel, B.A. & Kasten, E. (2000). Restoration of vision by training of residual functions. *Current Opinion in Ophthalmology*, 11:430-436.
- Sabel, B.A., Kenkel, S. & Kasten, E. (2004). Vision restoration therapy (VRT) efficacy as assessed by comparative perimetric analysis and subjective questionnaires. *Restorative Neurology and Neuroscience*, 22:399-420.
- Salter, K., Jutai, J., Zettler, L., Foley, N. & Teasell, R. (2006). *Evidence-based review of stroke rehabilitation: Outcome measures in stroke rehabilitation. Chapter 21*. Retrieved December, 2011, from <http://www.ebrsr.com/>

- Salter, K., Teasell, R., Bitensky, J., Foley, N. & Bhogal, S.K. (2007). *Evidence-based review of stroke rehabilitation: Cognitive disorders and apraxia. Chapter 12*. Retrieved December, 2011, from <http://www.ebrsr.com/>
- Schuett, S., Kentridge, R.W., Zihl, J. & Heywood, C.A. (2009). Adaptation of eye-movements to simulated hemianopia in reading and visual exploration: Transfer or specificity? *Neuropsychologia*, 47(7):1712-1720.
- Schuett, S., Kentridge, R.W., Zihl, J. & Heywood, C.A. (2009). Are hemianopic reading and visual exploration impairments visually elicited? New insights from eye movements in simulated hemianopia. *Neuropsychologia*, 47:733-746.
- Schulmann, D.L., Godfrey, B. & Fisher, A.G. (1987). Effect of eye movements on dynamic equilibrium. *Physical Therapy*, 67(7):1054-1057.
- Shah, S., Vanclay, F. & Cooper, B. (1989). Improving the sensitivity of the Barthel Index for stroke rehabilitation. *Journal of Clinical Epidemiology*, 42:703-709.
- Shimoda, K. & Robinson, R.G. (1998). Effect of anxiety disorder on impairment and recovery after stroke. *The Journal of Neuropsychiatry and Clinical Neurosciences*, 10:34-40.
- Shumway-Cook, A., Brauer, S. & Woollacott, M. (2000). Predicting the probability for falls in community-dwelling older adults using the Timed Up & Go Test. *Physical Therapy*, 80:896-903.

- Shumway-Cook, A., & Woollacot, M. H. (2007). *Motor control: Theory and practical applications*. (Third ed.). Philadelphia: Lippincott Williams & Wilkins.
- Siu, K.C. & Woollacott, M.H. (2006). Attentional demands of postural control: The ability to selectively allocate information-processing resources. *Gait & Posture*, 25(1):121-126.
- Spering, M. & Gegenfurtner, K.R. (2008). Contextual effects on motion perception and smooth pursuit eye movements. *Brain Research*, 1225:76-85.
- Swan, L. (2001). Unilateral spatial neglect. *Physical Therapy*, 81(9):1572-1580.
- Talelli, P., Ellul, J., Terzis, G., et al. (2004). Common carotid artery intima media thickness and poststroke cognitive impairment. *Journal of the Neurological Sciences*, 223:129-134.
- Teasell, R., Bayona, N. & Heitzner, J. (2011). *Evidence-based review of stroke rehabilitation: Clinical consequences of stroke. Chapter 2*. Retrieved December, 2011, from <http://www.ebrsr.com/>
- Teasell, R., Bayona, N. & Bitensky, J. (2011). *Evidence-based review of stroke rehabilitation: Background concepts in stroke rehabilitation. Chapter 3*. Retrieved December, 2011, from <http://www.ebrsr.com/>
- Teasell, R., Salter, K., Bitensky, J., Bhogal, S.K. & Foley, N. (2011). *Evidence-based review of stroke rehabilitation: Perceptual disorders. Chapter 13*. Retrieved December, 2011, from <http://www.ebrsr.com/>

- Tipping, B., De Villiers, L., Wainwright, H., Candy, S. & Bryer, A. (2007). Stroke in patients with human immunodeficiency virus infection. *Journal of Neurology Neurosurgery and Psychiatry*, 78(12):1320-1324.
- Ustun, T.B., Chatterji, S., Bickenbach, J., Kostanjsek, N. & Schneider, M. (2003). The International Classification of Functioning, Disability and Health: A new tool for understanding disability and health. *Disability and Rehabilitation*, 25 (11-12): 565-571.
- Van Vliet, P.M., Lincoln, N.B. & Foxall, A. (2005). Comparison of Bobath based and movement science based treatment for stroke: A randomized controlled trial. *Journal of Neurology, Neurosurgery and Psychiatry*, 76:503-508.
- Wagenaar, R.C., van Wieringen, P.C.W., Netelenbos, J.B., Meijer, O.G. & Kuik, D.J. (1992). The transfer of scanning training effects in visual inattention after stroke: Five single-case studies. *Disability and Rehabilitation*, 14(1):51-60.
- Webster, J.S., Jones, S., Blanton, P., Gross, R., Beissel, G.F. & Wofford, J.D. (1984). Visual scanning training with stroke patients. *Behavior Therapy*, 15(2):129-143.
- Weinberg, J., Diller, L., Gordon, W.A., Gerstman, L.J., Lieberman, A., Lakin, P., Hodges, G. & Ezrachi, O. (1977). Visual scanning training effect on reading – related tasks in acquired right brain damage. *Archives of Physical Medicine and Rehabilitation*, 58:479.

- Weinberg, J., Diller, L., Gordon, W.A., Gerstman, L.J., Lieberman, A., Lakin, P., Hodges, G. & Ezrachi, O. (1979). Training sensory awareness and spatial organisation in people with right brain damage. *Archives of Physical Medicine and Rehabilitation*, 60:491.
- Weinberg, J., Piasetsky, E., Diller, L. & Wayne, A. (1982). Treating perception organization deficits in non-neglecting patients. *Journal of Clinical Neuropsychology*, 4(1):59-75.
- West, C., Bowen, A., Hesketh, A. & Vail, A. (2009). Interventions for motor apraxia following stroke (Review). *The Cochrane Library*, 3:1-14.
- Whitney, S.L., Hudak, M.T. & Marchetti, G.F. (2000). The dynamic gait index relates to self-reported fall history in individuals with vestibular dysfunction. *Journal of Vestibular Research*, 10(2):99-105.
- Wiat, L., Bon Saint Côme, A., Debelleix, X., Petit, H., Joseph, P.A., Mazaux, J.M. & Barat, M. (1997). Unilateral neglect syndrome rehabilitation by trunk rotation and scanning training. *Archives of Physical Medicine and Rehabilitation*, 78:424-429.
- Wilson, B., Cockburn, J. & Halligan, P. (1987). Development of a behavioral test of visuospatial neglect. *Archives of Physical Medicine and Rehabilitation*, 68:98-101.

World Health Organization / World Federation of Neurology (2004). *Atlas: Country Resources for Neurologic Diseases*, Geneva: WHO, 2004.

World Health Organization, (2007). *World Health Organization*. Retrieved December, 2011, from <http://www.who.int/entity/classifications/icf/en/>

Yogev, G., Hausdorff, J.M. & Giladi, N. (2008). The role of executive function and attention in gait. *Movement Disorders*, 23(3):329-472.

Young, G.C., Collins, D. & Hren, M. (1983). Effect of pairing scanning training with block design training in the remediation of perceptual problems in left hemiplegics. *Journal of Clinical Neuropsychology*, 5(3):201-212.

Zigmond, A.S., & Snaith, R.P. (1983). Hospital Anxiety and Depression Scale. *Acta Psychiatrica Scandinavica*, 67:361-370.

Zihl, J. (1995). Eye movement patterns in hemianopic dyslexia. *Brain*, 118:891-912.

Zihl, J. (1995). Visual scanning behaviour in patients with homonymous hemianopia. *Neuropsychologia*, 33(3):287-303.

Zoltan, B. (1996). *Vision, perception and cognition. A manual for the evaluation and treatment of the neurologically impaired adult*. (Third ed., pp. 27-51). USA: SLACK Incorporated.

Zung, W.W.K. (1965). A self-rating depression scale. *Archives of General Psychiatry*, 12:63-70.