

Chapter 1 INTRODUCTION

Filariosis is the infection of vertebrate hosts with nematodes of the superfamily Filarioidea. Earlier known as filariasis, the term is derived from the generic name *Filaria* which according to Stiles (1907) has been used by zoologists and physicians 'as a generic catch-all for slender roundworms which could not be definitely determined'. The predilection sites of the preadult and adult stages of the filarial worms are the body cavities, blood or lymph vessels and connective tissue. The life cycles are indirect with mammals, amphibians, reptiles and birds acting as definitive hosts and haematophagous arthropods as intermediate hosts. Being viviparous nematodes, females produce incompletely differentiated first stage larvae, known as microfilariae, which are found in the blood and/or lymph. Several species are important pathogens of domestic animals and humans in mostly tropical and subtropical areas of the world. In the veterinary field, filariosis of domestic carnivores is of particular importance. Due to its severe pathogenic effects as well as profound financial implications for owners, *Dirofilaria immitis*, colloquially known as 'heartworm' or 'canine heartworm', constitutes the most important species for both dog and cat. Apart from the genus *Dirofilaria*, the spectrum of filarial helminths encountered in domestic carnivores belongs to the genera *Acanthocheilonema*, *Cercopithifilaria* and *Brugia*. Although *Acanthocheilonema*, *Brugia*, *Cercopithifilaria* and *Dirofilaria* species, other than *D. immitis*, were considered as largely non-pathogenic, there is growing evidence that infections are not so innocuous as generally assumed (Schwan, Miller, De Kock & Van Heerden 2000; Tarello 2003, 2004; Schwan & Schröter 2006). With the introduction of macrocyclic lactone-based dewormers for dogs and cats, filariosis of any aetiology has gained significance.

Similarly to the previously widely used diethylcarbamazine, macrocyclic lactones have microfilaricidal activity which can result in a potentially fatal shock-like syndrome and other adverse reactions as demonstrated in *D. immitis* and *Dirofilaria repens* infected dogs and cats (McGaughey 1952; Sasaki, Kitagawa, Ishihara & Shibata 1989; Euzéby 1990; Schrey 1996; Ware 2003; Plumb 2008; V. Schwan, unpublished data 2008).

Apart from the direct effects on the dog and cat population there are also zoonotic implications as humans can act as accidental hosts for some filarial helminths. The ever-increasing movement of people with their pets and climatic changes are regarded as important factors for the continuous spreading of filariosis (Russell 1985; Poglayen 1996; Rossi, Pollono, Meneguz, Gribaudo & Balbo 1996; Bucklar Scheu, Mossi & Deplazes 1998; Irwin 2002; Tarello 2003; Genchi, Rinaldi, Cascone, Mortarino & Cringoli 2005).

In contrast to most other continents, there is a lack of published information on the occurrence and distribution of filarial helminths of dogs and cats in Africa and its islands, which argues for a strong effort to conduct systematic field studies (Lok 1988). In a first attempt to remediate this situation, the objectives of this study are:

- a) to determine the occurrence and prevalence of *Dirofilaria immitis* and other filarial helminths in dogs in Gauteng, KwaZulu-Natal and Mpumalanga provinces, South Africa and Maputo province, Mozambique,
- b) to determine the occurrence and prevalence of filarial helminths of cats in KwaZulu-Natal province,



- c) to evaluate the results of routine examinations for filarial infections of dogs and cats from South Africa conducted between 1994 and 2008,
- d) to evaluate the results of routine examinations for filarial infections of dogs and cats imported from African countries conducted between 1992 and 2008, and
- e) to conduct a literature review on filariasis of dogs and cats in Africa.

Chapter 2

LITERATURE REVIEW OF THE DIFFERENT AETIOLOGIES OF FILARIOSIS IN DOMESTIC CARNIVORES IN AFRICA

A total of ten confirmed filarial species has been reported worldwide in domestic carnivores (Table 2.1). As regards Africa and its islands, there are published reports of autochthonous cases of filarial infections in dogs and cats involving six of these species (Nelson, Heisch & Furlong 1962; Laub 1988).

Of these the species *D. immitis*, *Dirofilaria repens*, *Acanthocheilonema reconditum*, *Acanthocheilonema dracunculoides* and *Brugia patei*, that all belong to the family Onchocercidae of the superfamily Filarioidea, form the subjects of this study.

2.1 *Dirofilaria immitis*

2.1.1 Taxonomy

Dirofilaria immitis (Leidy, 1856), commonly known as heartworm or canine heartworm, belongs to the subfamily Dirofilarinae and was first described as *Filaria canis cordis* in 1850 by Leidy in Philadelphia. In 1856 the worm was renamed *Filaria immitis* by Leidy. Railliet & Henry (1911a) erected the genus *Dirofilaria* and designated *Filaria immitis* as its type species. As a result of subsequent descriptions by various authors the helminth figures in the literature under the following synonyms: *Filaria canis cordis* (Leidy, 1850), *Filaria papillosa haematica canis-domestica* (Gruly & Delafond, 1852), *Filaria immitis* (Leidy, 1856), *Filaria papillosa haematica* (Schneider, 1866), *Filaria spirocauda* (Leidy, 1858), *Filaria cordis phocae* (Joly, 1858), *Filaria haematica* (Leuckart, 1867), *Filaria sanguinis* (Cobbold, 1869), *Filaria hebetata* (Cobbold, 1873), *Filaria spirocauda* (Cobbold, 1879), *Filaria* sp. (Horst, 1889); *Microfilaria immitis* (Neumann & Mayer,

1914), *Dirofilaria nasuae* (Mazza, 1926), *Dirofilaria pongoi* (Vogel & Vogelsang, 1930), *Dirofilaria indica* (Chakravarty, 1936), *Filaria magalhaesi* (Blanchard, 1895), *Dirofilaria magalhaesi* (Blanchard, 1895), *Dirofilaria fausti* (Skrjabin & Schikhobalova, 1948) and *Dirofilaria louisianensis* (Faust, Thomas & Jones, 1941) (Anderson 1952; Sonin 1985).

Faust (1937) proposed that the genus *Dirofilaria* be split into the subgenera *Dirofilaria* and *Nochtiella*. Species whose predilection site is the cardiovascular system were allocated to the subgenus *Dirofilaria*, whereas the subgenus *Nochtiella* contains those whose predilection site is the subcutaneous connective tissue.

2.1.2 Morphology

The morphological features that characterize the genus have been described by Railliet & Henry (1911a), Vogel (1927), Lent & Freitas (1937) and Sonin (1985). A detailed description of the adult stages of *D. immitis* is given by Fülleborn (1912) and Vogel (1927). According to these authors, females are 21-31 cm long and 1-1.3 mm wide whereas males are 12-20 cm long and 0.6-0.9 mm wide. The left spicule is 300-355 µm long, and the right spicule 175- 226 µm.

The microfilariae are unsheathed, and a detailed description was given by Fülleborn (1912) and Taylor (1960a). The cephalic end is conical and the posterior end is acute with the nuclear column (i.e. the cells that constitute the body of the microfilaria) not extending to the end of the body. The tail in unfixed and unstained microfilariae is straight (Marconcini, Magi, Macchioni & Sasseti 1996). Minimum and maximum measurements regarding length and width range from 180-340 µm and 5-7 µm respectively (Table 2.2).

The infective filarial larva in mosquitoes has been originally described by Nelson (1959) with additional information being provided by Taylor (1960b), Orihel (1961), Lichtenfels, Pilitt, Kotani & Powers (1985) and Bain & Chabaud (1986).

2.1.3 Life cycle

Dirofilaria immitis females produce microfilariae which are found in the blood of the definitive host. They are also capable of passing through the placenta and infect foetuses *in utero* (Mantovani & Jackson 1966; Atwell 1981; Todd & Howland 1983), and microfilariae have been found in urine (Kaewthamasorn, Assarasakorn & Niwetpathomwat 2008) and synovial fluid (Hodges & Rishniw 2008). The appearance of microfilariae in dogs in the peripheral blood is nocturnal subperiodic, with maximum levels of microfilaraemia being attained during late afternoon and at night with some geographical variation (Kosuge 1924; Schnelle & Young 1944; Euzéby & Lainé 1951; Webber & Hawking 1955; Newton & Wright 1956; Tongson & Romero 1962). Apart from the daily periodicity, there is also a seasonal periodicity with microfilariae being more abundant in the peripheral blood during spring and summer (Newton 1968; Kume 1975; Sawyer 1975). There exists a coincidence between the time microfilariae are most abundant in the peripheral blood and the time mosquito vectors obtain blood meals, a circumstance which is regarded as an evolutionary adaptation (Abraham 1988). Microfilariaemia in cats is only seen in less than 20 % of cases, and is inconsistent and transient when present (Cusick, Todd, Blake & Daly 1976). In cats the microfilaraemia is also nocturnal subperiodic (Nogami, Marasugi, Shimazaki, Maeda, Harasawa & Nakagaki 2000).

Mosquitoes act as intermediate hosts. Microfilariae develop into 3rd stage infective larvae in the mosquito and their development has been described by Taylor (1960b), Christensen (1977) and Bradley, Sauerman & Nayar (1984). The incubation period in mosquito vectors is largely temperature-dependent and may take as little as 14-17 days in *Aedes aegypti* (Taylor 1960b). While feeding, infective larvae emerge from the tips of the labella together with a drop of haemolymph onto the surface of the host's skin (Lavoipierre 1958). The haemolymph pool provides a medium in which the larvae can maintain their motility to search for and penetrate the puncture wound remaining after the withdrawal of the mosquito fascicle (Zielke 1973). The developing larvae are also pathogenic for the vector itself, which results in an increased mortality (Kartmann 1953; Galliard 1957; Christensen 1977; Hamilton & Bradley 1979).

In the definitive host the infective larvae undergo an extensive somatic migration to so-called intermediate locations, which are the submuscular membrane, subcutaneous tissue, adipose tissue, subserosa and muscles of the upper abdomen, thorax, head, neck and forelimb regions (Kume & Itagaki 1955). During this migration they moult to the L4-stage and then into young adults which finally enter veins to reach their predilection sites, the right ventricle, right auricle and pulmonary artery (Nelson 1966; Kotani & Powers 1982; Orihel 1961). Worms are found in the heart as early as day 67 after infection (Kume & Itagaki 1955) and the migration is always completed by day 90 (Orihel 1961). There are many reports of *D. immitis* found in aberrant sites (Otto 1975) which is more common in cats than in dogs (Dillon 1988). The prepatent period is 6-9 months in dogs (Bancroft 1904; Webber & Hawking 1955; Orihel 1961; Newton 1968; Kotani & Powers 1982) and 8 months in cats (Donahue 1975). The patent period is up to 7½ years in dogs (Newton 1968) and only about 2 years in cats (Donahue 1975;

Wong, Pedersen & Cullen 1983). The maximum life expectancy of microfilariae in the blood of dogs is 2½ years (Underwood & Harwood 1939).

2.1.4 Host range

Domestic dogs act as the preferential and principal definitive host (Abraham 1988). Cats are less prone to develop a patent infection and thus regarded as insignificant reservoirs of infection (Donahue 1975; Dillon 1988; Wong *et al.* 1983). Apart from these hosts, some wild canids, felids, other mammals, including man, and the Humboldt penguin (*Spheniscus humboldti*) have been found to be infected with adult *D. immitis* (Campbell & Blair 1978; Abraham 1988; Starr & Mulley 1988; Vellayan, Omar, Oothuman, Jefferey, Zahedi, Mathew & Krishnasamy 1989; Canestri Trotti, Pampiglione & Rivasi 1997; Sano, Aoki, Takahashi, Miura, Komatsu, Abe, Kakino & Itagaki 2005). In the majority of these hosts the adult worms are found in aberrant locations and do not produce microfilariae (Abraham 1988).

2.1.5 Vectors

About 70 anopheline and culicine mosquitoes throughout the world, that belong to the genera *Aedes*, *Anopheles*, *Coquillettidia*, *Culex*, *Culiseta*, *Mansonia* and *Psorophora* have been identified as potential intermediate hosts (Bemrick & Sandholm 1966; Ludlam, Jachowski & Otto 1970; Lok 1988). However, innate susceptibility is only a component of true vector competence, which is determined by the demonstration of infective larvae in field-captured mosquitoes (Lok 1988). As regards Africa there are very few references to natural infections of mosquitoes with animal filariae. Because infective larvae of *D. immitis* are practically indistinguishable on morphological criteria from those of *D. repens*, available records from Africa that specifically refer to *D. immitis* (Table 2.3) are therefore of only limited value (Nelson *et al.* 1962).

Geographical strains of mosquito species from Africa that have been found to be susceptible after experimental infections and might therefore act as natural vectors, are *Anopheles pambaensis* from Kenya (Nelson *et al.* 1962), *Aedes aegypti* from Kenya (Nelson *et al.* 1962) and Tanzania (Roubaud 1937) as well as *Culex pipiens fatigans* from Kenya (Heisch, Nelson & Furlong 1959; Nelson *et al.* 1962).

2.1.6 Laboratory diagnosis in live animals

The laboratory diagnosis in live animals can be attained by the demonstration and identification of microfilariae, by serology and by molecular techniques. Various methods have been described for the detection of microfilariae in the blood of animals and humans. The preparation of wet blood films, thin and thick blood films stained with Romanovsky-type stains as well as the capillary haematocrit tube method are appropriate if high levels of microfilaraemia prevail (Schalm & Lain 1966; Collins 1971; Kelly 1973; Bailey 1987). Standardized concentration techniques allow detection of low microfilaraemia levels and make it possible to quantify microfilaria densities. In the classical modified Knott's technique haemolysed blood is centrifuged and the sediment screened microscopically for microfilariae (Knott 1939; Newton & Wright 1956). Variations of this technique have been reviewed by Ho Thi Sang & Petithory (1963). In the membrane filtration technique 1 ml of blood treated with an anticoagulant is forced through a 3.0 μm polycarbonate membrane filter which is stained with Giemsa and examined microscopically (Bell 1967; Chularerk & Desowitz 1970; Dennis & Kean 1971; Chlebowski & Zielke 1977). The membrane filtration technique is more sensitive than the Knott's technique in cases where the microfilaria density is low (100-50 microfilariae/ml blood) (Bell 1967; Watson, Testoni & Porges 1973; Southgate 1974; Feldmeier, Bienzle, Schuh, Geister & Guggenmoos-Holzmann 1986; Beugnet, Bima-Blum & Chardonnet 1993a; Martini, Capelli, Poglayen, Bertotti & Turilli 1996). Since

there are several filarial species in both dog and cat that produce microfilariae which eventually appear in the blood, the mere demonstration of microfilariae remains meaningless unless they are identified (Valcárcel, Ferre, Gómez-Bautista & Rojo-Vázquez 1990).

2.1.6.1 MORPHOMETRICAL IDENTIFICATION OF MICROFILARIAE

Depending on the technique of processing, storage of blood samples and geographical origin, the morphometrical data given for heartworm microfilariae in the literature vary considerably, ranging from 180-340 μm and 5-7 μm respectively (Table 2.2) (Sawyer, Weinstein & Bloch 1963; Acevedo, Theis, Kraus & Longhurst 1981). As the size ranges of microfilariae of other species that concurrently occur on the African continent overlap with those of *D. immitis*, it is unreliable to establish a diagnosis on this criterion (Valcárcel *et al.* 1990). Described by Fülleborn (1924), the relative positions of somatic structures of the microfilarial body, which is constant in each species, is still one of the most accurate methods. The method is, however, sufficiently laborious to be impractical except for taxonomic purposes (Kelly 1973).

2.1.6.2 MORPHOLOGICAL IDENTIFICATION OF MICROFILARIAE

The morphology of the tail and the shape of the anterior extremity of the microfilariae have been described as features for identification (Sonin 1985; Marconcini *et al.* 1996). Tail morphology should only be considered in either unstained and unfixed microfilariae or in those isolated with the modified Knott's technique (Marconcini *et al.* 1996). In *D. immitis*, these morphological criteria are of limited value on the African continent, since *Dirofilaria repens* also has a straight tail and a conical anterior extremity (Valcárcel *et al.* 1990).

2.1.6.3 PHYSIOLOGICAL CHARACTERISTICS OF MICROFILARIAE

Motility in wet blood films is a criterion emphasized especially in the North American literature (Thrasher 1963). However, Valcárcel *et al.* (1990) could not confirm any difference in the motility of microfilariae of *D. immitis*, *D. repens*, *A. dracunculoides* and *A. reconditum*. Similarly, the claim that high microfilarial counts are indicative of *D. immitis* infection in contrast as opposed to low counts in *A. reconditum* infections (Wallenstein & Tibola 1960), has proved questionable, as high microfilarial counts have also been observed in dogs infected with *A. reconditum* (Herd 1978; Bobade, Ojebuoboh & Akinboade 1981).

2.1.6.4 HISTOCHEMICAL IDENTIFICATION OF MICROFILARIAE

In studies of filarial infections in monkeys, Chalifoux & Hunt (1971) observed differences in acid phosphatase activity of microfilariae in blood films using the method of Barka (1960). Subsequently, the method was applied to microfilariae of dogs and cats and has, with modifications, since been proven to be the most reliable, consistent and practical differential technique (Balbo & Abate 1972; Kelly 1973; Whitlock, Porter & Kelly 1978; Valcárcel *et al.* 1990; Beugnet, Costa & Lambert 1993b; Ducos de Lahitte, Ducos de Lahitte & Davoust 1993; Peribáñez, Lucientes, Arce, Morales, Castillo & Garcia 2001). In the microfilariae of *D. immitis*, acid phosphatase activity is uniquely restricted to the excretory pore and the anal pore (Chalifoux & Hunt 1971; Balbo & Abate 1972; Valcárcel *et al.* 1990). Histochemistry is ideally combined with the modified Knott's technique or the membrane filtration technique (Williams, Williams, Signs & Hokama 1977; Whitlock *et al.* 1978; Acevedo *et al.* 1981).

2.1.6.5 SEROLOGY

There are two groups of serological tests to detect circulating female heartworm antigen or circulating heartworm antibody. For the detection of antigen, enzyme-linked immunosorbent assay (ELISA) and immunochromatographic test systems are available. The main advantages of serological testing for antigen lie in the identification of 'occult' (amicrofilaremic) infections, the monitoring of adulticide treatment and the fast execution of the test by the veterinarian in the presence of the owner (Beugnet *et al.* 1993a). Disadvantages are the lack of sensitivity in only male worm infections, in prepatent infections and if low numbers of female worms are present (Beugnet *et al.* 1993b; Hoover, Campbell, Fox, Claypool & Mullins 1996). Lack of sensitivity is particularly a problem in low-endemic or newly colonized areas, where concentration tests provide more accurate results (Frank, Grieve, Mok, Smart & Salman 1992; Tarello 2001). False sero-positive results, due to cross-reactions with *D. repens* have to be considered (Valcárcel *et al.* 1990; Beugnet *et al.* 1993b; Schrey 1996; Schwan *et al.* 2000). The abundant North American literature currently regards antigen testing as the most sensitive diagnostic method (Datz 2003). However, data on the accuracy of the various commercial tests available cannot be extrapolated to other geographical areas since, with a single exception, all commercial test kits are manufactured in the USA where cross-reactions with *Dipetalonema reconditum* as the only other filarial species of dogs can be excluded (Schrey 1996). Heartworm antibody tests are used in cats, where infections are usually amicrofilaraemic and antigen is difficult to detect because of the low worm burdens (Datz 2003).

2.1.6.6 MOLECULAR DIAGNOSIS

The polymerase chain reaction (PCR) and DNA probes as tools in the molecular diagnosis have been used to differentiate *D. immitis* from other filarial helminths (WHO 1992; Favia, Lanfrancotti, Della Torre, Cancrini & Coluzzi 1996; Favia, Lanfrancotti, Della Torre, Cancrini & Coluzzi 1997; Bredal, Gjerde, Eberhard, Aleksandersen, Wilhelmsen & Mansfield 1998; Mar, Yan, Chang & Fei 2002; Rishniw, Barr, Simpson, Frongillo, Franz & Dominguez Alpizar 2006). However, due to technical inadequacies, a lack of practical trials to validate the techniques appropriately combined with exhaustive technical requirements, the routine application of molecular-biological techniques remains limited (WHO 1992; Pampiglione, Rivasi & Canestri Trotti 2000; Shaw & Day 2005; Olga & Éva 2006).

2.1.7 *Veterinary and medical importance*

Cardiovascular dirofilariosis is caused by the preadult and adult worms which exert a mechanical and phlogistic effect that, depending on the worm burden, duration of infection and host-parasite interaction, ultimately develops into a multisystemic disorder with the lungs, heart, liver and kidneys mainly affected (Pampiglione & Rivasi 2001; Ware 2003). The pulmonary arterial system is the prime site of pathology and the effects are reflected by alterations in the pulmonary vasculature and interstitial lung tissue (Sutton 1988). Due to the development of pulmonary hypertension, the heart and liver become affected which leads to right-sided circulatory failure with the kidneys also partially becoming involved in the cascade of events (Ducos de Lahitte 1990). The pathogenesis, pathology and clinical manifestations have been subject of many reviews (Knight 1977, 1987; Ducos de Lahitte 1990; Ducos de Lahitte *et al.* 1993; Ware 1998, 2003).

A microfilaria-associated cutaneous syndrome characterized by erythematous, papulo-nodular and/or ulcerative pruritic lesions and a membranous glomerulonephritis have been described on very few occasions (Casey & Splitter 1975; Mozos, Ginel, López, Carrasco, Martín de las Mulas & Molleda 1992; Hargis, Lewis, Duclos, Loeffler & Rausch 1999). However, with the introduction of macrocyclic lactone-based dewormers for dogs and cats, microfilariae of any filarial species have gained significance. Similarly to the previously widely used diethylcarbamazine, macrocyclic lactones have microfilaricidal activity which can result in a potentially fatal shock-like syndrome and other adverse reactions as demonstrated in *D. immitis* and *Dirofilaria repens* infected dogs and cats (Sasaki *et al.* 1989; Euzéby 1990; Schrey 1996; Klotins, Martin, Bonnett & Peregrine 2000; Ware 2003; Plumb 2008; V. Schwan, unpublished data 2008). This is presumably due to immunological reactions against the substances released from dying microfilariae (Sasaki *et al.* 1989; Plumb 2008).

Clinically, cardiovascular dirofilariosis can present as a mild asymptomatic form, which is mostly detected incidentally, as a moderate form with exercise intolerance, chronic cough, dyspnoea and weight loss or as a severe form with right-sided congestive heart failure, syncope, acute or chronic *vena cava* syndrome and sudden death (Atwell 1988; Moraillon 1990).

About 230 human cases have been reported worldwide (Muller 2002). In almost all instances immature worms or unfertilized females have been isolated from the lungs (Pampiglione & Rivasi 2001). Most infections are asymptomatic, showing typical 'coin lesions' on chest radiography which are often mistakenly removed as neoplasms (Ciferri 1982).

The treatment of cardiovascular dirofilariosis consists of chemotherapy directed against preadult and adult worms by means of macrofilaricides (adulticides) and subsequently against microfilariae in the blood by means of 'microfilaricides' (McCall, Guerrero, Genchi & Kramer 2004). Possible systemic side effects such as pulmonary thromboembolism caused by the reaction of the body to the disintegrating adult worms and circulatory collapse following the rapid death of large numbers of microfilariae may require additional treatment with anti-inflammatories and parenteral fluids (Ware 2003).

Melarsomine (Immiticide[®], Merial) is currently the drug of choice for the treatment of preadult and adult stages in dogs (Raynaud 1992). Macrofilaricidal therapy in cats should only be considered as a last resort, as severe complications are very likely to occur (Ware 2003). In dogs, microfilaricidal therapy is recommended to be started 3-4 weeks after macrofilaricidal therapy in dogs (McCall *et al.* 2004). Ivermectin administered as a single dose of 50 µg/kg has to be regarded as the drug of choice for this purpose as it causes fewer side effects than other microfilaricides (Beugnet *et al.* 1993b; Ware 2003). At this dose, the drug is also safe in ivermectin-sensitive dog breeds (Pulliam, Seward, Henry & Steinberg 1985; Paul, Tranquilli, Seward, Todd & Di Pietro 1987).

For preventative use, specific formulations of the macrocyclic lactones ivermectin, milbemycin oxime, moxidectin and selamectin administered once a month and the piperazine diethylcarbamazine (DEC) given daily are available (Ware 2003; McCall *et al.* 2004; Plumb 2008).

2.1.8 *Distribution on the African continent and its islands*

A survey for parasitism in animals conducted by FAO, WHO and OIE (1984) indicates that filariasis of dogs is widespread in Africa. According to Nelson (1966) it was common in veterinary practice to assume that dogs with microfilariae in their blood were infected with *D. immitis* and that this has resulted in a great deal of confusion with other harmless species. This is supported by Levine (1980), who maintains that *D. immitis* is rather rare in Africa.

2.1.8.1 NORTHERN AFRICA

Dirofilaria immitis has been reported from Algeria, Egypt, Morocco, Tunisia and the offshore Canary Islands.

Algeria: The earliest reports come from Beni-Ounif de Figuig on the Algerian-Moroccan border where several small scale surveys were conducted between 1913 and 1923 (Foley 1921; Foley, Catanei & Vialatte 1926) with prevalence rates ranging from 10-22 % based on the demonstration and identification of microfilariae. According to the authors, the microfilariae were 175-253 μm long, 6-5 μm wide and had a cephalic hook. This description is contrary to published information for *D. immitis*, and only the microfilariae of *A. reconditum* possess a cephalic hook (Sawyer, Rubin & Jackson 1965). The misdiagnosis is also supported by the fact that the authors were unable to demonstrate adult worms in the heart of the microfilaraemic animals and that none of them showed any clinical signs attributable to heartworm infection. Choquette, Gayot & Poul (1952) report a case from a dog in the Alger region based on the finding of microfilariae. The authors have not provided any information on what criteria the microfilariae were identified. In a later survey involving 190 dogs from the Alger region, eight were found positive for microfilariae on blood examination with a description given

that is in accordance with published information (Rioche 1960). In the most recent survey conducted in Alger, Montaron (1975) found one out of 215 dogs positive for *D. immitis* on examination of blood. The identification was based on microfilarial motility and the appearance of the tail.

Egypt: *Dirofilaria immitis* was incidentally found during postmortem examination in the pulmonary arteries in 5 out of 50 dogs in Assiut (Mahmoud & Ibrahim 1989). In a later report eight out of 19 police dogs in Assiut were found positive for microfilariae on blood examination (Abd El Rahim 1998). The microfilariae were identified morphologically by their tapered anterior end and their straight tail with no further information provided.

Morocco: Heartworm in Morocco was first reported by Bouin (1921), who conducted a necropsy survey in the south of the country with no exact locality given. The survey involved 109 dogs of which one was found positive with a single female specimen isolated from the right heart. Santucci, Haag & Sendral (1953) reported on a clinical case of dirofilariosis in a 2-year-old male Boxer. The diagnosis was based on the morphometrical identification of microfilariae in Giemsa-stained blood films using the data given by Neveu-Lemaire (1936) as a reference. In a more recent necropsy survey in the Rabat region 7 out of 57 stray dogs were reported to be infected (Pandey, Dakkak & Elmamoune 1987).

Tunisia: In a survey involving 207 dogs from Tunis, one dog was found positive for microfilariae which occurred in large numbers (Yakimoff & Kohl-Yakimoff 1911). The microfilariae are described as sheathless, 214-227 μm long and 4.2-5.6 μm wide with no further details provided. Juminer & Durand (1960) report on a dog with severe polyparasitism which was subsequently euthanased and necropsied. Although the

authors failed to demonstrate adult parasites in the heart during necropsy, microfilariae found in stained bloodfilms were identified as those of *D. immitis* with no details provided on the criteria used for identification. In a survey conducted in Tunis, 25 out of 70 dogs were found to be microfilaraemic (Perrot 1985). The identification of the microfilariae as those of *D. immitis* was based on motility, morphometrical and morphological criteria and is in accordance with published information.

Canary Islands: In a prevalence survey on the Canary Island of Tenerife 130 out of 310 dogs were microfilaraemic (Valladares, Gijon & Lopez-Roman 1987). No details are given on what criteria the microfilariae were identified. However, *D. immitis* infection was confirmed by demonstration of adult parasites in the heart of 14 selected microfilaraemic animals during necropsy. In a more recent survey, heartworm seroprevalence in dogs from Tenerife Island was 21 % (172/823) (Montoya, Morales, Juste, Bañares, Simon & Genchi 2006). Seroprevalence surveys conducted during 1994 to 1996 involving 2034 dogs on Gran Canaria Island showed a mean prevalence of 58.89 % (Montoya, Morales, Ferrer, Molina & Corbera 1998).

2.1.8.2 WESTERN AFRICA

Dirofilaria immitis has been reported from Guinea-Bissau, Nigeria, Senegal and Sierra-Leone. However, most of the reports are based on the demonstration of microfilariae with negative necropsy results, a circumstance which is unlikely as is highlighted by Schillhorn van Veen & Blotkamp (1975), who also maintain that *D. immitis* is rare in Western Africa.

Guinea-Bissau: The existing reports are based on the demonstration of microfilariae only (Tendeiro 1948, 1949). Although Tendeiro (1949) claims that *D. immitis* is very

common in dogs, in what was then Portuguese Guinea, the author also mentions that adult parasites have never been found and that no clinical signs were ever observed in infected animals. Tendeiro (1949) gives a detailed morphological and morphometrical description of the microfilariae and compares the obtained data with, amongst others, those of Fülleborn (1912) and Foley (1921). While Fülleborn's data are quoted incorrectly, Foley's description of the microfilariae having a cephalic hook and the circumstance that no clinical signs were observed in infected dogs suggests the aetiology of *A. reconditum*.

Nigeria: Schillhorn van Veen (1974) maintains that *D. immitis* has not been demonstrated in dogs in Nigeria, although veterinary field officers often assume that dogs with microfilariae in their peripheral blood are infected (Idowu, Okon & Dipeolu 1977). This is the conclusion after routine necropsies of 400 dogs in Zaria failed to demonstrate heartworm.

Senegal: As the only reference available, Pangui & Kaboret (1993) report on a survey conducted in Dakar. Between 1984 and 1992, 72 stray dogs were caught and necropsied of which six were found to be infected with adult worms at the predilection site.

Sierra Leone: The two existing reports from Sierra Leone are based on the finding of microfilariae only (Kamara 1977; Hassan 1984) with no information provided on the criteria used for identification. Necropsies conducted on microfilaraemic dogs in the earlier study failed to demonstrate worms at the predilection site (Kamara 1977).



2.1.8.3 CENTRAL AFRICA

Dirofilaria immitis has been reported from Angola, Cameroon and Gabon.

Angola: Serrano (1962) reports that microfilariae of *D. immitis* have been found in the blood of dogs in Luanda and Nova Lisboa and that the adult stage was recovered in the heart of a dog in Luanda.

Cameroon: Thys, Sawa & Guissart (1982) published a case history of a 7-year-old male Boxer in Maroua who was brought to Cameroon from Yugoslavia 7 months before being presented for treatment. The animal was found to be infected with *D. immitis* based on demonstration and identification of microfilariae in stained bloodfilms. The diagnosis was confirmed at necropsy. However, the authors suggest that the dog was already infected before arriving in Maroua where *D. immitis* appears not to be endemic due to the unfavourable climatic conditions. Heartworm is endemic in the territory of the previously known Yugoslavia where it was first reported by Dzunkovski (1934) in a dog in Belgrade.

Gabon: In a survey involving 48 dogs from Libreville, 50 % tested positive for *D. immitis* based on the identification of microfilariae, positive necropsy results and heartworm antigen testing (Beugnet & Edderai 1998).

2.1.8.4 EASTERN AFRICA

Reports come from Ethiopia, Kenya, Malawi, Mozambique and Tanzania and the islands of Madagascar, Mauritius and Réunion.

Ethiopia: Chiodi (1936) reports that *D. immitis* is common in dogs in Abyssinia. In his collation of data on helminth infection of domestic and wild animals, Graber (1975) cites *D. immitis* which was recovered from the right ventricle and the pulmonary artery from a dog of unknown origin in Ethiopia by Chiodi in 1936.

Kenya: The earliest report comes from the Island of Pate (Heisch *et al.* 1959) where nine out of 12 dogs were infected with *D. immitis*. The diagnosis was based on the identification of microfilariae and demonstration of adults during necropsy. The identity of the microfilariae was confirmed by comparing them with those taken from the gravid uteri of adult worms found at autopsy. In a later survey on filarial infections in man, animals and mosquitoes on the Kenya coast from Somalia to Tanzania, 22 out of 252 dogs were found infected (Nelson *et al.* 1962). A necropsy survey involving 286 dogs from the Nairobi area yielded two cases (Murray 1968). There are two further reports where infection of dogs was demonstrated during necropsy from various unspecified localities in the country (Bwangamoi & Frank 1970; Bwangamoi, Frank, Moulton, Mugeru & Wandera 1971).

Malawi: *Dirofilaria immitis* is mentioned in a check list of helminth parasites from domestic dogs (Fitzsimmons 1964).

Mozambique: In the earliest report, Dias (1954) states that *D. immitis* appears to be rare. The author reports on a single case from the Region of Maputo which was identified in a laboratory with, however, no details provided on what criteria the diagnosis was based. Cruz e Silva (1971) refers to adult specimens (several male and female worms) collected by Travassos Dias in 1969 in Quelimane as well as specimens from Beira collected in 1966 from dogs at necropsy. In a survey conducted in Maputo

Province between 1981 and 1984, Jurášek (1986) found five out of 86 dogs to be microfilaraemic. The author claims that the microfilariae were those of *D. immitis* with, however, no information provided on the criteria used for identification. In a small-scale survey in Quelimane 4 out of 13 indigenous dogs were found positive for microfilariae which were identified by acid phosphatase staining (Schwan & Durand 2002) as those of *D. immitis*.

Tanzania: Infection in a dog was first recorded by the Veterinary Department of Tanganyika (1934). Alley (1950) gives a clinical report on six cases of *D. immitis* infected dogs on the island of Zanzibar. In the annual report of the Veterinary Department, Roe (1958) lists *D. immitis* as having been diagnosed in dogs.

Madagascar: *Dirofilaria immitis* in dogs is mentioned in a host-parasite list of helminths in domestic animals (Daynes 1964). The author claims that *D. immitis* is known for a long time on the island.

Mauritius: Heartworm is mentioned for the first time for Mauritius as part of a helminth collection from domestic animals on the island (Ware 1925). The specimens were recovered from the dog's heart. In a necropsy survey, Webb & Nadeau (1958) found three out of 50 stray dogs to be infected. A more recent survey indicated that 30 out of 184 dogs were microfilaraemic (Sibartie, Beeharry & Jaumally 1983). The microfilariae were identified as those of *D. immitis* according to published criteria and infection was further confirmed by necropsy in four animals.

Réunion: Prunaux & Guignard (1991) report on the end result of a 4-year investigation of the Veterinary Departmental Laboratory of Réunion Island, in which 16 out of 96 dogs were found positive on necropsy.

2.1.8.5 SOUTHERN AFRICA

Dirofilaria immitis has only been reported in imported animals (Van Heerden, Verster & Gouws 1980; Verster, Cilliers & Schroeder 1991; Schwan & Durand 2002).

2.2 *Dirofilaria repens*

2.2.1 Taxonomy

Dirofilaria repens (Railliet & Henry, 1911) has no vernacular name and has been known in the literature by the following names: *Filaria acutiuscula* (Molin, 1858); *Dirofilaria acutiuscula* (Molin, 1858); *Filaria palpebralis* (Pace 1867); *Filaria peritonaei hominis* (Babes, 1880); *Filaria conjunctivae* (Addario, 1885); *Dirofilaria conjunctivae* (Addario, 1885); *Filaria repens* (Braun, 1915) and *Loa extraocularia* (Skrjabin, 1917) (Anderson 1952; Sonin 1985; Chauve 1990).

2.2.2 Morphology

A detailed description of the adult stages of *D. repens* is given by Railliet & Henry (1911b), Vogel (1927) and Le-Van-Hoa & Le Thi-Ty (1971). According to these authors females are 8.4-17 cm long and 380-650 µm wide whereas males are 3.9-7 cm long and 270-450 µm wide. The left spicule is 338-590 µm long, and the right spicule 123-206 µm long.

The microfilariae are unsheathed and a detailed description is given by Gunewardene (1956) and Taylor (1960a). According to these authors the cephalic end is conical with 2-3 nuclei in the head space and the posterior end is acute with the nuclear column not extending to the end of the body. The tail in unfixed and unstained microfilariae is like the handle of an umbrella (Marconcini *et al.* 1996). Influenced by the technique of processing, geographical origin and host, the morphometrical data given in the literature vary considerably. Minimum and maximum measurements regarding length and width range from 207-385 μm and 5-9 μm (Table 2.4).

The infective filarial larva in mosquitoes has been described in detail by Nelson (1959, 1960) with additional information provided by Bain & Chabaud (1986).

2.2.3 Life cycle

Dirofilaria repens females produce microfilariae which are found in the blood of the definitive host. They are also capable of extravascular migration as is evidenced by passing through the placenta and infecting puppies (Mantovani 1966) and the demonstration of microfilariae in urine (Mantovani 1965). The appearance of microfilariae in the peripheral blood of dogs and cats is nocturnal subperiodic, with maximum levels of microfilaraemia between 20:00 and 03:00 (Webber & Hawking 1955; Mantovani & Restani 1965; Kamalu 1986).

Mosquitoes act as intermediate hosts. The incubation period in susceptible mosquito vectors is temperature- and species-dependent and may take as little as 10 days in *Anopheles stephensi* (Webber & Hawking 1955). Development within the vector and subsequent transmission is similar as for *D. immitis* and has been described by

Fülleborn (1908a), Bernard & Bauche (1913), Gunewardene (1956) and Mantovani (1965).

Whether infective larvae follow a complex migration in the definitive host is unknown (Webber & Hawking 1955). The predilection sites are the subcutaneous tissue in most parts of the body (Canestri Trotti *et al.* 1997) and the fascial sheaths overlying the muscles of the hind legs (Heisch *et al.* 1959) The prepatent period is 6-8 months in dogs (Webber & Hawking 1955) and 6 months in cats (Cancrini, Mantovani & Coluzzi 1979; Cancrini & Iori 1981). According to Webber & Hawking (1955) the patent period in dogs is at least 2-3 years. However, based on data obtained from experimentally infected dogs in Italy who remained microfilaraemic for 8-9 years, the patent period appears to be much longer (Cancrini & Iori 1981). The same authors state that experimentally infected cats remained microfilaraemic for about 2 years.

2.2.4 *Host range*

The domestic dog and cat act as the preferential and principal definitive hosts and both appear equally susceptible to infection in Africa (Heisch *et al.* 1959). Apart from these hosts some wild canids, felids and the large-spotted genet (*Genetta tigrina*) were found infected with adult *D. repens* (Canestri Trotti *et al.* 1997).

2.2.5 *Vectors*

Anopheline and culicine mosquito species belonging to the genera *Aedes*, *Anopheles*, *Culex*, *Mansonia* and *Taeniorhynchus* are considered intermediate hosts (Pampiglione, Canestri Trotti & Rivasi 1995).

There are no records on natural infections of mosquitoes with *D. repens* in Africa. Geographical strains of mosquito species from Africa that have been found to be susceptible following experimental infections are *Aedes pambaensis*, *Aedes aegypti*, *Mansonia uniformis* and *Mansonia africanus* from Kenya (Nelson *et al.* 1962) as well as *Aedes aegypti* from Nigeria (Anyanwu, Agbede, Ajanusi, Umoh & Ibrahim 2000).

2.2.6 Laboratory diagnosis in live animals

The laboratory diagnosis is based on the demonstration and identification of microfilariae in blood samples utilizing the same methodologies as outlined for *D. immitis*. Because of the unreliability of most characteristics, histochemical staining for acid phosphatase activity has proved to be the most reliable, consistent and practical differential technique to diagnose infection in dogs and cats (Kelly 1973; Valcárcel *et al.* 1990). In the microfilariae of *D. repens*, acid phosphatase activity is uniquely restricted to the anal pore or to the anal pore and innerbody (Balbo & Abate 1972; Yen & Mak 1978; Valcárcel *et al.* 1990). The utilization of the polymerase chain reaction (PCR) as a tool to differentiate *D. repens* microfilariae and immature adult stages removed from biptic material from those of other filarial species of dogs and cats has been reported (Favia *et al.* 1996, 1997; Vakalis, Spanakos, Patsoula & Vamvakopoulos 1999; Rishniw *et al.* 2006).

2.2.7 Veterinary and medical importance

In the past *D. repens* has been regarded as apathogenic in natural and experimental infections (Webber & Hawking 1955; Heisch *et al.* 1959; Nelson 1966). However, since the 1960s it has been recognized that *D. repens* is not an innocuous parasite in at least a subgroup of infected dogs and cats. Pruritic dermatitis characterized by the presence of erythema, papules, focal or multifocal alopecia, crusting and subcutaneous nodules

containing adult worms is the most commonly observed clinical manifestation of infection in both dogs and cats (Euzéby 1961; Kamalu 1986; Bredal *et al.* 1998; Tarello 2000a, 2002, 2003; Ananda & D'Souza 2006). The part of the body most affected are the lumbosacral and perineal areas, and the hind legs in dogs (Tarello 2002), and the flanks, back, neck, legs and paws in cats (Tarello 2000a). In dogs, the flanks, back and the hind limbs are commonly considered the preferential sites of dwelling for both the larvae and adults which may concentrate in large numbers in a single area (Mandelli & Mantovani 1966). The embolization of microfilariae, the movement of adults in the subcutaneous tissue and the immunological response to the L3, L4, adults and/or microfilariae are thought to cause these cutaneous lesions (Mozos *et al.* 1992; Chauve 1997; Pampiglione *et al.* 1995; Tarello 2002). Circular cutaneous ulcers, subcutaneous tumefactions, subcutaneous oedema and ascitis were reported from infected dogs which resolved following adulticidal and microfilaricidal treatment (Restani, Rossi & Semproni 1963). Cutaneous ulcers have also been reported from a cat (Tarello 2000b). Acute liver failure was reported from a microfilaraemic cat in South Africa which resolved after treatment with ivermectin (Schwan *et al.* 2000). The macropathological and histopathological changes of the spleen, liver, lungs, heart and kidneys, which have been described from some suspected clinical cases, are similar to those observed in cardiovascular dirofilariosis and comprise hyperplastic splenomegaly, plasmocytosis, erythrophagocytosis, haemosiderosis, chronic stasis of the liver with centrolobular steatosis and portal fibrosis, lung atelectasis and chronic bronchitis, glomerular fibrosis of kidneys, myocardosis, vascular tumours and other vascular alterations in the vessels of the nervous tissues (Mantovani 1965; Mandelli & Mantovani 1966; Kamalu 1991; Schwan *et al.* 2000; Martano, Veneziano, Santaniello, Carbone, Paciello, Cataldi, Russo & Maiolino 2004). There is increasing evidence that the pathogenicity may be

influenced by concurrent infections, such as babesiosis, monocytic ehrlichiosis, leishmaniosis and haemobartonellosis (Tarello 2002).

Treatment of *D. repens* infection in dogs and cats is indicated if they are clinically affected and to decrease the risk of human infection in endemic areas (Baneth, Volansky, Anug, Favia, Bain, Goldstein & Harrus 2002). However, reports on treatment are scarce. The adulticide thiacetarsamide (no dosage provided) followed by the microfilaricide diethylcarbamazine (100 mg/kg *per os* daily for 30 days) were used in Italy (Restani *et al.* 1963) effectively. Diethylcarbamazine at 5.5 mg/kg *per os* daily for 1 month was used in Nigeria with no apparent clinical effect (Kamalu 1991). This is less than the recommended daily chemoprophylactic dosage of 6.6 mg/kg for heartworm (Roberson 1988). The adulticide melarsomine (2 x 2.5 mg/kg 24 hours apart intramuscular) subsequently followed by the microfilaricide ivermectin (50 µg/kg) 10 and 30 days later was used for the treatment of dogs and a cat in Italy with resolution of the cutaneous lesions (Tarello 1999, 2000a, b, 2002, 2003). Similarly, melarsomine followed by doramectin (0.4 mg/kg) has been reported from Israel to be effective in clearing infections (Baneth *et al.* 2002).

Three macrocyclic lactones are reported to be used successfully in monthly dosing regimens to prevent infection in dogs, two by oral administration, ivermectin (Marconcini, Magi & Hecht Contin 1993; Pollono, Pollmeier & Rossi 1998) and moxidectin (Rossi, Ferroglio & Agostini 2002), and selamectin by pour-on application (Genchi, Poglayen & Kramer 2002). An injectable, sustained-release formulation of moxidectin shown to confer six-month protection against the related *D. immitis*, has proved to be similarly effective as a prophylactic for *D. repens* (Lok, Knight, Wang, Doscher, Nolan, Hendrick, Steber & Heaney 2001; Rossi, Ferroglio & Agostini 2004).

Dirofilaria repens accidentally affects humans and has been reported about 400 times from 30 countries but mostly from Italy with a more common superficial manifestation and a visceral form which is often confused with neoplastic tumours (Pampiglione *et al.* 1995; Muro, Genchi, Cordero & Simón 1999). There is only a single report worldwide of a patent infection with a microfilariaemia in a human case from Corsica (Nozais, Bain & Gentilini 1994).

2.2.8 Distribution on the African continent

2.2.8.1 NORTHERN AFRICA

Dirofilaria repens has been reported from Egypt, Sudan and Tunisia.

Egypt: In a check list of nematodes collected during 1948-1955, *D. repens* is listed for the golden jackal (*Canis aureus*) (Myers, Kuntz & Wells 1962).

Sudan: Adult *D. repens* adult specimens were recovered from the *Gluteus superficialis* and *Biceps femoris* muscles of 2 lions from Bahr-el-Ghazal Province (Kellas & Webber 1955).

Tunisia: Chatton (1918) reports on a survey conducted in Médine and Gabès in the south of the country, where 2 out of 26 cats were found microfilaraemic. The microfilariae were described as 240-350 µm long and 7-9 µm wide. In a later necropsy survey involving 348 dogs in Tunis, 1 dog was found infected (Bernard, Ben Osman & Juminer 1967). Adult worms were isolated from the supracostal connective tissue. The identification was based on a detailed morphometrical study with results being in accordance with published information.

2.2.8.2 WESTERN AFRICA

Dirofilaria repens has only been reported from Nigeria. Schillhorn van Veen (1974) maintains that *D. repens* is the most common filarial worm in Nigerian dogs and cats. According to this author it is found mainly in the subcutaneous connective tissue without causing any marked pathological changes. The author reports on a 9.4 % prevalence in dogs in the Zaria area based on the demonstration of microfilariae in blood. The size range given is 300-369 μm . There are several other reports of *D. repens* in dogs from Nigeria with information provided on the criteria used for identification of microfilariae and adults which confirm its endemic status (Schillhorn van Veen & Blotkamp 1975; Schillhorn van Veen, Shonekan & Fabiyi 1975; Kamalu 1986, 1991; Anyanwu, Umoh, Ogbogu, Essien, Galadima, Adawa & Hassan 1996).

2.2.8.3 CENTRAL AFRICA

The only report comes from the Kapa River in the northeast of the Central African Republic, where the adult specimens of *D. repens* were recovered from the subcutaneous connective tissue of a lion (Graber, Euzéby, Gevrey, Troncy & Thal 1972). The authors provide morphometrical data which are in accordance with published information.

2.2.8.4 EASTERN AFRICA

Dirofilaria repens has been reported from Kenya, Uganda, Zambia and Zimbabwe.

Kenya: In a survey on the Island of Pate, 2 out of 12 dogs, 27 out of 29 cats and 8 out of 9 large-spotted genet cats were found to be infected with *D. repens* (Heisch *et al.* 1959). Adult worms were found under the skin and were particularly common in the fascial

sheaths overlying the muscles of the hind legs with no obvious pathological lesions. The identity of microfilariae in the blood of animals was cross-checked by comparing them with those taken from the gravid uteri of known adult worms found at necropsy. The authors report that the parasite is not confined to the Island of Pate but that it is fairly common in cats and dogs on the Kenya coast. This was confirmed in a subsequent survey involving the entire Kenya coast from Somalia to Tanzania in which 6 out of 252 dogs and 43 out of 240 cats were found to be infected on day blood films (Nelson *et al.* 1962).

Uganda: Bwangamoi (1973) recovered two female specimens of *D. repens* from the subcutis of a dog from Kampala during necropsy. Except of their length (12.7 and 16.2 cm) no further criteria are presented for their identification. In a microfilarial survey involving 836 dogs from various parts of the country, 8.6 % were found positive (Bwangamoi & Isyagi 1973). The authors present a confusing morphometrical study with no references provided for identification and record, amongst other filarial species, the presence of *D. repens*.

Zambia: Le Roux (1958) reports that a lion on Mbesuma Ranch in the Chinsali District was found heavily infected with amongst other helminths *D. repens*. No details are given on what criteria the identification was based.

Zimbabwe: *Dirofilaria repens* is listed for cats in a checklist of helminth parasites of domestic and wild mammals of Zimbabwe (Jooste 1990).



2.2.8.5 SOUTHERN AFRICA

Dirofilaria repens has only been reported from South Africa in a cat from Pretoria (Schwan *et al.* 2000). The diagnosis was based on the demonstration of microfilariae and their identification by acid phosphatase staining.

2.3 *Acanthocheilonema reconditum*

2.3.1 Taxonomy

Acanthocheilonema reconditum (Grassi, 1889) has no vernacular name and has been known in the literature by the name *Filaria recondita* Grassi, 1889 and the widely used previous name, *Dipetalonema reconditum* (Grassi, 1889) (Sonin 1985). This latest allocation is based on an attempt to rearrange the complex genus *Dipetalonema* on evolutionary lines (Chabaud & Bain 1976; Bain, Baker & Chabaud 1982a). As a result of this the revived genus *Acanthocheilonema* includes those dipetalonematids which were similar to *Dipetalonema s.s.* of New World primates (Muller 1987).

2.3.2 Morphology

The genus *Acanthocheilonema* accommodates those dipetalonematids which have a well-chitinised buccal capsule, a sturdy, divided oesophagus and a right spicule that is provided with a well-developed sheath (Bain *et al.* 1982a).

A detailed description of the adult stages is given by Noè (1907), Nelson (1962), Korkejian & Edeson (1978) and Laub (1988). According to these authors, females are 21-36 mm long and 70-205 µm wide whereas males are 9-17 mm long and 70-133 µm wide. The left spicule is 220-300 µm long, and the right spicule 92-104 µm long.

The microfilariae are unsheathed. A detailed description is given by Laub (1988). They have a long clear head space (Nelson 1962) and a cephalic hook which was first described by Fülleborn (1913) and redescribed by Sawyer *et al.* (1965). The tail is attenuated and free of nuclei (Nelson 1962). The tail in unfixed and unstained microfilariae assumes the shape of a hook (Marconcini *et al.* 1996). Influenced by the technique of processing and geographical origin, the morphometrical data in the literature vary considerably. Minimum and maximum measurements regarding length and width range from 168-292 μm and 4-6.7 μm respectively (Table 2.5).

The infective filarial larva in *Heterodoxus spiniger* and *Ctenocephalides felis* has been described in great detail by Nelson (1962), Bain & Beaucournu (1974) and Laub (1988).

2.3.3 Life cycle

Information in the literature concerning the periodicity of microfilariae in the peripheral blood is controversial. Newton & Wright (1956) in the United States reported a diurnal subperiodicity. Gubler (1966) in Hawaii illustrated a periodic cycle with a diurnal and nocturnal peak in naturally infected dogs. Bobade *et al.* (1981) report on a marked nocturnal subperiodic appearance of microfilariae in an infected dog in Nigeria. However, the aetiology of the filarial infection in the latter report remains questionable since the identification was only based on the length and width of microfilariae, which according to Laub (1988) is inappropriate for species identification. Korkejian & Edeson (1978) report a nocturnal subperiodicity in naturally infected dogs in Lebanon. The results obtained from studies on naturally infected dogs in Brazil and Okinawa are inconclusive (Pennington & Phelps 1969; Lima & Costa 1972). A study conducted on naturally infected dogs in Liberia showed no periodicity of microfilariae (Laub 1988).

The fleas *Ctenocephalides canis*, *Ctenocephalides felis* and *Pulex irritans*, as well as the chewing lice *Heterodoxus spiniger* and *Linognathus setosus* have been identified as intermediate hosts (Grassi & Calandruccio 1890; Newton & Wright 1956; Nelson 1962; Pennington & Phelps 1969). The incubation period in *C. canis* is 20-23 days at 28 °C and 80 % humidity (Laub 1988) and 7 days in *C. felis* (Farnell & Faulkner 1978).

According to Nelson (1962) the predilection sites of the adult worms are the subcutaneous fascial spaces of the limbs and back. Korkejian & Edeson (1978) recovered adult worms from the trunk and hindlegs only. Grassi & Calandruccio (1890) found adults near the kidney.

The prepatent period is 61-101 days in experimentally infected dogs (Farnell & Faulkner 1978; Lindemann & McCall 1984). No information is available on the patent period of the species.

2.3.4 *Host range*

Apart from the domestic dog, which is the preferred and principal definitive host, *A. reconditum* has also been isolated from some wild canids as well as the spotted hyaena (*Crocuta crocuta*) and the brown hyaena (*Hyaena brunnea*) (Sonin 1985).

2.3.5 *Laboratory diagnosis in live animals*

The laboratory diagnosis is based on the demonstration and identification of microfilariae in blood samples by means of acid phosphatase staining (Kelly 1973; Valcárcel *et al.* 1990). In the microfilariae of *A. reconditum* acid phosphatase activity is either uniform with slighter lighter area from the cephalic end to the excretory vesicle or in some instances with diffuse denser staining in the area of the excretory pore,

innerbody and anal pore (Chalifoux *et al.* 1971; Acevedo *et al.* 1981). The presence of a cephalic hook as a differentiating feature is only visible in dehaemoglobinized, undried blood films that are stained with brilliant cresol blue (Sawyer *et al.* 1965). Nelson *et al.* (1962) discovered that by using *Ctenocephalides felis* for xenodiagnosis, infections in dogs were detected with microfilarial densities of less than 10/ml. The utilization of the polymerase chain reaction (PCR) as a tool to differentiate *A. reconditum* microfilariae from other filarial species of dogs has been reported (Mar *et al.* 2002; Rishniw *et al.* 2006).

2.3.6 Veterinary and medical importance

Acanthocheilonema reconditum is widely regarded as apathogenic (Grassi & Calandruccio 1890; Newton *et al.* 1956; Nelson 1962). However, cases of pruritic dermatosis and focal alopecia have been attributed to the action of microfilariae if present in large numbers (Bobade *et al.* 1981; Hubert 1985; Chauve 1990).

2.3.7 Distribution on the African continent

2.3.7.1 WESTERN AFRICA

Acanthocheilonema reconditum has been reported from Liberia and Nigeria.

Liberia: In a survey conducted in Bong County, Montserrado County, Cape Mount County and Grand Bassa County, 56 out of 137 dogs were found positive for microfilariae of *A. reconditum* (Laub 1988). The identification is based on a detailed morphological and morphometrical analysis.

Nigeria: Schillhorn van Veen & Blotkamp (1975) report on 'short-type microfilariae' found in 9.2 % of 369 dogs in the Zaria area which based on morphometrical analysis

were similar to those of *A. reconditum* and *A. dracunculoides*. Idowu *et al.* (1977) report on a survey in Ibadan in which two out of 488 dogs were found positive for microfilariae of *A. reconditum*. The authors do not provide any information on the microfilariae and the criteria used for identification. Bobade *et al.* 1981 report on a dog from Ibadan with microfilariae identified as *A. reconditum* by measuring length and width and tail shape.

2.3.7.2 EASTERN AFRICA

Acanthocheilonema reconditum has been reported from Kenya, Mozambique and Uganda.

Kenya: Nelson (1962) gives the first description of the parasite from Africa from material collected from dogs in Mombasa and Nairobi. The author reports that *A. reconditum* is the most widely distributed and common filarial species of dogs in Kenya. It is particularly common in dogs on the hot coastal strip and in the cooler highlands. It also occurs in jackals (*Canis adustus*, *Canis aureus*, *Canis mesomelas*) and hyaenas (*C. crocuta*, *Hyaena hyaena*). In a survey on filarial infections in man, animals and mosquitoes on the Kenya coast from Somalia to Tanganyika 40 out of 252 dogs were found positive for *A. reconditum* microfilariae on day blood films (Nelson *et al.* 1962).

Mozambique: In a small scale survey conducted in the Quelimane area 1 out of 13 dogs was found positive for *A. reconditum*, as identified by acid phosphatase staining of microfilariae (Schwan & Durand 2002).

Uganda: Bwangamoi (1973) reports on a dog from Kampala with microfilariae which were not identified. At necropsy 16 adult and one immature filarial worms were recovered and identified as *A. reconditum*. Although the overall body measurements fall

in the range of *A. reconditum*, the measurements given for the spicules (left spicule 161 µm, right spicule 48 µm) are not in accordance with published information. In a microfilarial survey involving 836 dogs from various parts of the country, 8.6 % were found positive (Bwangamoi & Isyagi 1973). The authors present a confusing morphometrical study with no references for identification, and conclude, amongst other filarial species, the occurrence of *A. reconditum*.

2.3.7.3 SOUTHERN AFRICA

The only report comes from South Africa. Van Heerden (1986) reports on microfilariae of *A. reconditum* in 6 out of 13 blood samples obtained from wild dogs (*Lycaon pictus*) with no data and details provided on what criteria the diagnosis was based.

2.4 *Acanthocheilonema dracunculoides*

2.4.1 Taxonomy

Acanthocheilonema dracunculoides (Cobbold, 1870) has no vernacular name and has been known in the literature by the names *Acanthocheilonema dagestanica* (Yarulin, 1962), *Microfilaria sp.* (Leger, 1911); *Microfilaria lewisi* (Korke, 1924); *Haematozoon lewisi* (Rao, 1923) and the widely used previous name *Dipetalonema dracunculoides* (Cobbold, 1870) (Sonin 1985).

2.4.2 Morphology

A detailed description of the adult stages is given by Leger (1911), Railliet, Henry & Langeron (1912), Rao (1938), Fraga de Azevedo (1943) and Nelson (1963) with additional information provided by Rioche (1960) and Chauve (1990). According to these authors females are 30-60 mm long and 200-370 µm wide whereas males are 15-

31 mm long and 100-310 μm wide. The left spicule is 320-402 μm long, and the right spicule 120-186 μm long.

Microfilariae are unsheathed and a detailed description was given by Rioche (1960) and Ortega-Mora, Gomez-Bautista & Rojo-Vázquez (1989). They have a clear head space and a short attenuated tail free of nuclei that ends bluntly. There is no cephalic hook (Leger 1911). The tail in unfixed and unstained microfilariae is straight (Marconcini *et al.* 1996). Minimum and maximum measurements regarding length and width range from 121-277 μm and 3.1-7.4 μm respectively (Table 2.6).

The infective larva in *Hippobosca longipennis* has been described by Nelson (1963) and Bain (1971).

2.4.3 Life cycle

Acanthocheilonema dracunculoides females produce microfilariae which are found in the blood of the definitive host. The information in the literature concerning periodicity of microfilariae is controversial. While Bouin (1921) and Montaron (1975) could not observe any periodicity in naturally infected dogs in Algeria and Morocco, Wolfe, Aslamkhan, Sharif & Pervez (1971) reported a diurnal subperiodicity in an infected dog in Pakistan.

So far, the louse fly *Hippobosca longipennis* and the hard tick *Rhipicephalus sanguineus* have been identified as intermediate hosts (Nelson 1963; Olmeda-García, Rodríguez-Rodríguez & Rojo-Vázquez 1993; Olmeda-García & Rodríguez-Rodríguez 1994).

The predilection sites of the adult worms are the abdominal and thoracic cavities (Chauve 1990).

2.4.4 Host range

Apart from the domestic dog, *A. dracunculoides* has been isolated from the aardwolf (*Proteles cristatus*), the spotted hyaena and the red fox (*Vulpes vulpes*) (Sonin 1985).

2.4.5 Laboratory diagnosis in live animals

Similar to other filarial infections of dogs and cats, histochemical staining for acid phosphatase activity of microfilariae has proved to be the most reliable, consistent and practical differential technique to diagnose infection (Valcárcel *et al.* 1990). Acid phosphatase activity is restricted to the cephalic space, the excretory pore, inner body and the anal pore (Ortega-Mora *et al.* 1989; Chauve 1990; Peribáñez *et al.* 2001). The utilization of the polymerase chain reaction (PCR) as a tool to differentiate *A. dracunculoides* microfilariae from other filarial species of dogs has been reported (Rishniw *et al.* 2006).

2.4.6 Veterinary and medical importance

Although *A. dracunculoides* is regarded as non-pathogenic in the dog (Nelson 1966; Montaron 1975), there is some evidence reported from Spain, Kenya, Uganda and Namibia that suggests that the parasite may not be as innocuous as generally assumed. Infection occasionally presents with dermal clinical signs and lesions ranging from pruritus, alopecia, erythema to skin ulcers as well as other clinical signs such as ataxy, incoordination, cachexia, cyanosis, ascitis and pleural effusion (Piercy 1951; Ortega-Mora & Rojo Vázquez 1988; Chauve 1990; Bolio, Montes, Gutierrez, Alonso, Bernal, Sauri & Rodríguez-Vivas 2002; Schwan & Schröter 2006). Dermal clinical signs

in a dog attributed to *A. dracunculoides* infection improved after treatment with ivermectin at a dose rate of 50 µg/kg (Rodríguez 1990). There are no reports of human infections (Chauve 1990).

2.4.7 Distribution on the African continent

Although Nelson (1963) claims that *A. dracunculoides* is widespread in the drier areas of Africa extending from the Mediterranean to South Africa, the parasite has only been reported from some countries. According to Nelson (1963) the distribution of the parasite coincides with the distribution of *Hippobosca longipennis*.

2.4.7.1 NORTHERN AFRICA

Acanthocheilonema dracunculoides has been reported from Algeria, Morocco, Sudan and Tunisia.

Algeria: The first report comes from a dog in the Alger region (Rioche 1960), who found numerous adult worms at necropsy in the peritoneal cavity. The author provides a detailed description of the adult male and female worms and the microfilaria. Montaron (1975) reports on a survey in Alger in which 49 (22.8 %) out of 215 dogs were found positive for microfilariae on examination of blood. In 48 dogs microfilariae were identified as those of *A. drancunculoides* based on motility and appearance of the tail. The diagnosis was confirmed at necropsy were adult worms were found in the abdominal and thoracic cavities.

Morocco: Bouin (1921) reports on a necropsy survey conducted in southern Morocco with no exact locality given. The survey involved 109 dogs of which 19 tested positive for microfilariae prior to necropsy. The microfilariae were identified morphometrically as

those of *A. dracunculoides*. However, adult filarial worms were found in the thoracic, peritoneal and pelvic cavities of 1 dog only. In a later necropsy survey, *A. dracunculoides* was reported in 6 out of 57 stray dogs from Rabat and the nearby towns of Temara, Sidi Yahya des Zaers and Ain Aouda (Pandey *et al.* 1987). Giemsa-stained blood films were prepared prior to euthanasia. The authors do not provide any information on what specimens (microfilariae or adult worms) and what criteria the diagnosis was based.

Sudan: Specimens were obtained from the peritoneal cavity of a dog from an undisclosed locality in the country (Baylis 1929).

Tunisia: Railliet *et al.* (1912) report on a microfilaraemic dog from Tunis. At necropsy adult worms were recovered from the peritoneal cavity. The microfilariae found in the blood were identical to those isolated from the posterior parts of the female uterus. The authors, however, do not provide any description of the microfilariae. In a later survey involving 348 dogs in Tunis, 17.54 % were found to be infected at necropsy, and adult worms were recovered from the peritoneal and pleural cavities (Bernard *et al.* 1967).

2.4.7.2 WESTERN AFRICA

Reports come from Mali and Nigeria.

Mali: Leger (1911) isolated adult worms from the peritoneal cavity of a spotted hyaena in the outskirts of Bamako. Railliet & Henry (1911b) suspected the relation of Leger's filarial species to *A. dracunculoides*. Railliet *et al.* (1912) examined Leger's specimens and confirmed that they were *A. dracunculoides*. In a collation of data on helminth infection of the former French West Africa, the specimens collected by Leger (1911) are

listed and the subsequent identification by Railliet *et al.* (1912) is mentioned (Joyeux, Gendre & Baer 1928).

Nigeria: *Acanthocheilonema dracunculoides* was first reported from dogs in the Zaria area (Schillhorn van Veen 1974). The diagnosis was based on the demonstration and morphometrical analysis of microfilariae, with, however, no details being provided other than the length (234-264 µm). Subsequently, Schillhorn van Veen & Blotkamp (1975) report on 'short-type microfilariae' found in dogs during a survey in the Zaria area, which, based on morphometrical analysis, were similar to those of *A. reconditum* and *A. dracunculoides*. In a host-parasite checklist of helminth parasites of domestic animals in Northern Nigeria, *A. dracunculoides* is listed as recorded in the Zaria area in dogs but is said to be rare (Schillhorn van Veen, Shonekan & Fabiyi 1975).

2.4.7.3 CENTRAL AFRICA

The only report comes from the Democratic Republic of the Congo where it was recovered from the abdominal cavity of a dog and from the pleural cavity of another dog from Katanga by Rodhain (Gedoelst 1916).

2.4.7.4 EASTERN AFRICA

Reports on the occurrence of the parasite come from Kenya, Tanzania, Uganda and Zimbabwe.

Kenya: *Acanthocheilonema dracunculoides* was first discovered in hyaenas (species not specified) near Nairobi and in dogs from the Northern Province (Nelson *et al.* 1962). The authors also report that the parasite had not been seen in dogs at the coast. Another report comes from Lokitaung in northern Turkana where adult specimens were

recovered from the peritoneal cavity of a dog (Nelson 1963). Another report on a survey involving 63 dogs of which 79 % were found to be infected on blood examination and necropsy came from the same locality (Lightner & Reardon 1983). According to the authors, the morphology and dimensions of the microfilariae corresponded with descriptions given by Nelson (1963) and Wolfe *et al.* (1971).

Tanzania: Sachs (1976) reports on *A. dracunculoides* from the abdominal cavity of spotted haenas in the Serengeti.

Uganda: Carmichael & Bell (1943) report on a 6-year-old Alsation which presented with multiple non-pruritic skin lesions. Unsheathed microfilariae, 240 µm long and 4.5 µm wide, were found in the blood but no further information was provided. Necropsy did not reveal any adult worms. The authors claim that the microfilariae were close to *A. dracunculoides*.

Zimbabwe: *Acanthocheilonema dracunculoides* in the spotted hyaena is listed in a check list of helminth parasites of domestic and wild mammals of Zimbabwe (Jooste 1990).

2.4.7.5 SOUTHERN AFRICA

Reports come from Namibia and South Africa.

Namibia: Microfilariae found in two dogs from Windhoek were identified by acid phosphatase staining as those of *A. dracunculoides* (Schwan & Schröter 2006).

South Africa: The country, where the species was first discovered and described from specimens recovered from an aardwolf caught at an undisclosed locality (Cobbold 1870).

2.5 *Cercopithifilaria grassii*

2.5.1 Taxonomy

Cercopithifilaria grassii (Noé, 1907) has no vernacular name and has been known in the literature by the names *Filaria grassii* (Noé, 1907), *Acanthocheilonema grassii* (Baylis, 1929) and the still widely used previous name *Dipetalonema grassii* (Noé, 1907) (Sonin 1985).

2.5.2 Morphology

The genus *Cercopithifilaria* is used to accommodate filarial species considered as specialized *Acanthocheilonema* (Bain *et al.* 1982a). Differentiating features are a short and undivided oesophagus, a very small buccal capsule, a stumpy right spicule without a distinct sheath, and the caudal papillae reduced in number and situated close to the cloaca (Bain *et al.* 1982a).

The morphology of the adult stages of *C. grassii* is described by Noè (1907); Costa & Freitas (1962) and Balasubramaniam, Anandan & Alwar (1975) According to these authors females are 24-27.2 mm long and 56-114 µm wide whereas males are 10-11.2 mm long and 40-45 µm wide. The left spicule is 200-215 µm and the right spicule 50-65 µm long.

Microfilariae are sheathed. According to the only detailed descriptions by Noè (1907, 1908) the microfilariae have a squat, blunt tail that terminates in three papillae. The

cephalic end is slightly distended and there are transversal cuticular ridges along the rest of the body. Compared to the other canine and feline filarial species, the microfilariae are huge, measuring 567 μm in length and 12.25 μm in width (Noè 1907).

The infective larva in *Rhipicephalus sanguineus* has been described by Noè (1908), Bain, Aeschlimann & Chatelanat (1982b) and Bain & Chabaud (1986).

2.5.3 Life cycle

Cercopithifilaria grassii microfilariae are found mainly in the skin (Noè 1907), the lymph (Pampiglione & Canestri Trotti 1990) and occasionally in the blood (Casarosa 1985). *Rhipicephalus sanguineus* is the only vector identified (Noè 1908; Bain *et al.* 1982b).

2.5.4 Host range

Domestic dogs and cats are the only definitive hosts reported for *C. grassii* (Tarello 2004). Adult worms are found in the subcutaneous and intermuscular connective tissue as well as in the abdominal cavity (Costa & Freitas 1962; Chauve 1990).

2.5.5 Laboratory diagnosis in live animals

The laboratory diagnosis is based on the demonstration of sheathed microfilariae in skin snips or blood, taking the unique dimensions of the microfilaria into consideration (Tarello 2004).

2.5.6 Veterinary and medical importance

Although, infection with *C. grassii* is rarely reported and considered to be harmless (Nelson 1966; Bain *et al.* 1982b; Chauve 1990), there is a report by Tarello (2004) of an infected cat that presented with pruritic dermatitis and multifocal alopecia. Following

treatment with melarsomine and ivermectin the lesions and clinical signs resolved. There are no reports of human infections (Chauve 1990).

2.5.7 *Distribution on the African continent*

In the only documentation from Africa, Heisch *et al.* (1959) and Nelson *et al.* (1962) report on finding a *Dipetalonema* species in a dog in Faza, Kenya and a dog in Dar-es-Salaam, with adults recovered from the subcutaneous tissues and under the abdominal muscles. The microfilariae, which are only described as 300 µm long on average, were confined to the skin. The authors suggest that this might be *C. grassii*. The authors do not provide any information on the adults.

2.6 *Brugia patei*

2.6.1 *Taxonomy*

Brugia patei (Buckley, Nelson & Heisch, 1958) is referred to in the literature also as *Wuchereria patei* Buckley, Nelson & Heisch, 1958 (Sonin 1985). The genus name *Brugia*, chosen in honour of the discoverer of *Microfilaria malayi*, Dr SL Brug, was proposed by Buckley in 1960 for a distinct group of filarial helminths which is parasitic in the lymphatic system of primates, carnivores and insectivores. The species was named after Pate Island where it was first discovered by Nelson & Heisch (1957).

2.6.2 *Morphology*

The genus *Brugia* accommodates a small group of morphologically very similar filarial helminths with affinities to *Wuchereria bancrofti* (Sonin 1975). The subtle morphological features that characterize the genus and differentiate it from *Wuchereria* have been described by Buckley (1960).

A description of the adult stages is given by Buckley, Nelson & Heisch (1958). According to these authors females are 34.5-50.7 mm long and 135-190 μm wide whereas males are 14-25.4 mm long and 75-100 μm wide. The left spicule is 255-295 μm and the right spicule 110-130 μm long.

The microfilariae are sheathed and very similar to those of *Brugia malayi* with the only constant difference in the length of the nuclei-free cephalic space which averages 4.8 μm as compared with 6.9 μm in *B. malayi* (Feng 1933; Buckley *et al.* 1958). Measurements regarding width and length range from 165-260 and 5-6 μm respectively and there are characteristic terminal tail nuclei (Feng 1933; Laurence & Simpson 1971).

The infective larva in mosquitoes has been described by Nelson (1959) and Bain & Chabaud (1986).

2.6.3 Life cycle

Brugia patei microfilariae are found in the blood of the definitive host. A non-periodic and nocturnal subperiodic strain have been identified in Kenya (Nelson *et al.* 1962).

Mosquitoes act as intermediate hosts. The incubation period in *Mansonia uniformis* takes 6-9 days at 28 °C (Laurence & Pester 1960; Nelson *et al.* 1962). There is no information on the prepatent and patent period in definitive hosts.

2.6.4 Host range

Apart from the domestic dog and cat, the large-spotted genet and the greater bushbaby (*Galago crassicaudatus*) have been identified as natural hosts (Nelson 1959). Infection

rates in cats are reported to be higher than in dogs and large-spotted genets and adult worms are most commonly found in the lymphatics of the hind legs (Heisch *et al.* 1959).

2.6.5 Vectors

Brugia patei infective larvae were found in *Aedes pembaensis*, *Mansonia uniformis* and *Mansonia africanus* (Nelson *et al.* 1962). The vector *A. pembaensis* has an obligatory phoretic association with salt-water crabs (Goiny, Van Someren & Heisch 1957).

2.6.6 Laboratory diagnosis in live animals

The diagnosis is based on the demonstration of sheathed microfilariae in the blood, taking into consideration the characteristic tail nuclei as a differentiating morphological feature (Buckley 1960).

2.6.7 Veterinary and medical importance

Although *B. patei* is regarded as largely non-pathogenic (Nelson 1966), there is a report from Kenya where two cats with ascitis had adult worms in the abdominal lymphatics and numerous sheathed microfilariae in the ascitic fluid (Heisch *et al.* 1959).

Microfilariae of *B. patei* have never been seen in the blood of man (Heisch *et al.* 1959). However, they may be an important aetiology of tropical pulmonary eosinophilia which is not uncommon on the Kenya coast (Nelson *et al.* 1962). This assumption is based on observations of Buckley (1958) who provided experimental proof by inoculating himself with infective larvae of *Brugia* spp. from animals.

2.6.8 *Distribution on the African continent and its islands*

Brugia patei was first discovered in domestic dogs and cats on Pate Island in Kenya and has not been reported from any other country in Africa (Nelson & Heisch 1957). The discovery was based on finding a sheathed microfilaria of the *Brugia malayi*-type. Buckley *et al.* (1958) report on a 56 % prevalence (14 out of 24 animals) of microfilarial infection in cats from Pate Island and also mention infection in 2 out of 5 dogs and three out of seven large-spotted genets. In a subsequent survey on the island, infection was reported again in dogs, cats and large-spotted genets with a similar higher prevalence in cats (21 out of 29) than in dogs (3 out of 12) (Heisch *et al.* 1959). The diagnosis was made by demonstration of adults at necropsy and microfilariae in day bloods. The microfilariae found in the blood were identical to those isolated from the gravid uteri of adult worms. The authors also found the parasite in cats on the Island of Lamu and in villages on the Tana River. However, *B. patei* could not be demonstrated in day bloods from cats and dogs at Mombasa, Pemba, other unidentified villages along the southern Kenya coast, Zanzibar and Dar-es-Salaam. In another survey on filarial infections in man, animals and mosquitoes on the Kenya coast from Somalia to the former Tanganyika, 15 out of 252 dogs, 38 out of 240 cats, 3 out of 9 large-spotted genets and 1 out of 10 greater bushbabies were found infected on day blood films (Nelson *et al.* 1962).

2.7 **Other reported species**

2.7.1 *Microfilaria auquieri*

Foley (1921) reports on a new microfilaria found in the blood of dogs in Beni-Ounif de Figuig on the Algerian-Moroccan border which was named in memory of Dr Auquier, a former physician in Figuig. The microfilaria was described as unsheathed, without a cephalic hook and characterized by its remarkable shortness (58-102 μm) and relative

large width (6-8 μm). The only other report comes from the Région de Palestro in Algeria, where the microfilaria was found in one dog and is described as 89 μm long and 7 μm wide on average (Rioche 1960). Although a systematic necropsy was conducted, adult worms could not be found.

2.7.2 *Filaria ochmanni*

A new microfilaria described as sheathed and 320 μm long on average in haematoxylin-stained thin blood films was found in a dog from Dar-es-Salaam in Tanzania (Fülleborn 1908b). The microfilaria was named after the state veterinarian Ochmann in Dar-es-Salaam who supplied the material. Buckley *et al.* (1958) suggest that this microfilaria belongs to the '*malayi*' group of microfilariae.



Table 2.1: Filarial helminths described from dogs and cats and their geographical distribution

Species	Host	Geographical distribution
<i>Acanthocheilonema dracunculoides</i>	Dog	Africa, Asia, Europe
<i>Acanthocheilonema reconditum</i>	Dog	Africa, America, Asia, Europe, Australia
<i>Brugia ceylonensis</i>	Dog, cat	Asia
<i>Brugia malayi</i>	Dog, cat	Asia
<i>Brugia pahangi</i>	Dog, cat	Asia
<i>Brugia patei</i>	Dog, cat	Africa
<i>Cercopithifilaria baineae</i>	Dog	South America
<i>Cercopithifilaria grassii</i>	Dog, cat	Europe
<i>Dirofilaria immitis</i>	Dog, cat	Africa, America, Asia, Europe, Australia
<i>Dirofilaria repens</i>	Dog, cat	Africa, Asia, Europe

Table 2.2: Length and width of *Dirofilaria immitis* microfilariae from dogs according to geographical origin and technique of processing

Length [μ m] Range \pm SD or Mean	Width [μ m] Range or Mean	Geographical origin/ Technique of processing	Source
Africa			
264 \pm 5.8	5 \pm 0.2	Algeria Technique not specified	Rioche (1960)
250		Kenya Technique not specified	Heisch <i>et al.</i> (1959)
232-260	4-6	Mozambique Methanol fixation and Giemsa staining	Schwan & Durand 2002
America			
327.46 \pm 2.36		USA Knott's technique	Acevedo <i>et al.</i> (1981)
281.32 \pm 4.24		USA Membrane filtration	Acevedo <i>et al.</i> (1981)
326.15 \pm 3.27		USA Membrane filtration with subsequent methanol fixation and Giemsa staining	Acevedo <i>et al.</i> (1981)
269.5 \pm 3.72		USA Membrane filtration with subsequent formalin fixation and methylene blue staining	Acevedo <i>et al.</i> (1981)
233-270 (253)	5.4-6.5 (5.96)	USA Microfilariae isolated from formalinized female worms	Fülleborn (1912)
285.6 – 339.8	6.1-7.2	USA Knott's technique	Lindsey (1961)
307-322	-	USA Knott's technique	Newton & Wright (1956)
Asia			
260-280	5.7-7.5	Vietnam Giemsa staining	Mathis & Léger (1911)
180-285	5	Vietnam Technique not specified	Railliet & Henry (1911b)
210-253 (233.7)		China Microfilariae isolated from female worms preserved in 70% ethanol	Fülleborn (1912)
245-333	7-8.5	Taschkent Technique not specified	Yakimoff (1917)
260 \pm 5	4.5	China Bouin's fixative and Giemsa staining	Taylor (1960a)
Australia			
256.7 \pm 16.6	-	Membrane filtration	Watson <i>et al.</i> (1973)
301.3 \pm 22.6	-	Knott's test	Watson <i>et al.</i> (1973)
Europe			
220-340	5-6.5	France Technique not specified	Ducos de Lahitte <i>et al.</i> (1993)
290-330	6-6.5	Italy Unfixed and unstained microfilariae	Marconcini <i>et al.</i> (1996)
306.83 \pm 22.41	5.9 \pm 0.69	Spain Knott's technique	Valcárcel <i>et al.</i> (1990)



Table 2.3: Natural culicine vectors of *Dirofilaria immitis* in Africa

Species	Locality	Reference
<i>Aedes aegypti</i>	Kenya	Nelson <i>et al.</i> (1962)
<i>Aedes pembaensis</i>	Kenya Tanzania	Nelson <i>et al.</i> (1962) Mosha & Magayuka (1979)
<i>Anopheles pharoensis</i>	West Africa	Brengues & Nelson (1975)
<i>Anopheles tenebrosus</i>	Tanzania	Gillies (1964); Magayuka (1973); Mosha & Magayuka (1979)
<i>Culex quinquefasciatus</i>	Mauritius	Halcrow (1954)
<i>Mansonia africana</i>	Tanzania	Magayuka (1973); Mosha & Magayuka (1979)
<i>Mansonia uniformis</i>	Madagascar Tanzania West Africa	Brunhes, Rajaonarivelo & Nelson (1972) Magayuka (1973), Mosha & Magayuka (1979) Brengues & Nelson (1975)

Table 2.4: Length and width of *Dirofilaria repens* microfilariae from dogs, cats and other carnivores according to geographical origin and technique of processing

Length [μm] Range \pm SD or Mean	Width [μm] Range or Mean	Geographical origin/ Technique of processing	Source
Africa			
328	-	Kenya (dog/cat/large-spotted genet)	Heisch <i>et al.</i> (1959)
300-369	-	Nigeria (dog) Brilliant cresylblue staining	Schillhorn van Veen (1974)
340-353	7.5-8	Nigeria (dog) Live/Stubbs technique	Kamalu (1986)
320-360	7.5-8	Nigeria (dog) Live/Stubbs technique	Kamalu (1991)
320-350	6.25-7.5	South Africa (cat) Membrane filtration and Giemsa staining	Schwan <i>et al.</i> (2000)
240-350	7-9	Tunisia (cat) Technique not specified	Chatton (1918)
Asia			
300-360	6.5-8	Vietnam (dog) Technique not specified	Railliet & Henry (1911b)
300-375	6.5-7	Vietnam (dog) Technique not specified	Bernard & Bauche (1913)
290 \pm 15	6-8	Sri Lanka (dog) Giemsa staining	Gunewardene (1956)
337	8	Vietnam (fishing cat) Giemsa staining	Le-Van-Hoa & Le Thi-Ty (1971)
Europe			
315 \pm 22	6 \pm 0.5	Sardinia (dog) Unfixed and unstained	Webber & Hawking (1955)
290 \pm 10	6	Italy (dog) Bouin's fixative + Giemsa staining	Taylor (1960a)
340	-	Italy (cat)	Cancrini & Iori (1981)
377	-	Italy (dog)	
207-360	5-8	France Technique not specified	Chauve (1990)
290-360	6-8	France Technique not specified	Ducos de Lahitte & Ducos de Lahitte (1990)
345-385	6.5-7	Italy (red fox) Unfixed and unstained	Marconcini <i>et al.</i> (1996)
345.27 \pm 19.3	6.4 \pm 0.78	Spain (dog) Knott's technique	Valcárcel <i>et al.</i> (1990)

Table 2.5: Length and width of *Acanthocheilonema reconditum* microfilariae from dogs according to geographical origin and technique of processing

Length [μm] Range \pm SD or Mean	Width [μm] Range or Mean	Geographical origin/ Technique of processing	Source
Africa			
270	4.5	Kenya Giemsa-stained blood films	Nelson (1962)
227.6-259.7 (239.3)	4.4-6.7	Liberia Haematoxylin/Ethanol fixation	Laub (1988)
239.8-273.1 (257.1)	4.4-6.7	Liberia Methylene blue/Formaldehyde fixation	Laub (1988)
225-282 (263)	4.93-5.86 (5.02)	Nigeria Knott's technique	Bobade <i>et al.</i> (1981)
200-204	4	Mozambique Giemsa-stained blood films	Schwan <i>et al.</i> (2002)
America			
262.09 \pm 3.36	-	USA Knott's technique	Acevedo <i>et al.</i> (1981)
241.06 \pm 2.34	-	USA Membrane filtration	Acevedo <i>et al.</i> (1981)
246.4-291.6	4.7-5.8	USA Technique not specified	Lindsey (1961)
276	-	USA Technique not specified	Newton & Wright (1956)
Asia			
230-290 (263.6)	4-5	Lebanon Knott's technique	Korkejian & Edeson (1978)
167.8-228.5 (204.34)	2.9-5	Brilliant cresyl blue staining	
230-285 (262)	3.2-5.8 (4.7)	Okinawa Knott's technique	Pennington & Phelps (1969)
Australia			
226.6 \pm 13.4	-	Membrane filtration	Watson <i>et al.</i> (1973)
255.2 \pm 24.8	-	Knott's technique	Watson <i>et al.</i> (1973)
Europe			
269-283	4	France Technique not specified	Chauve (1990)
200-230	4-5	France Technique not specified	Ducos de Lahitte (1990)
210-215	-	France Technique not specified	Euzéby (1961)
270-280	5-5.5	Italy in <i>Vulpes vulpes</i> Unfixed and unstained	Marconcini <i>et al.</i> (1996)
261.96 \pm 14.28	-	Spain Knott's technique	Valcárcel <i>et al.</i> (1990)

Table 2.6: Length and width of *Acanthocheilonema dracunculoides* microfilariae from dogs and other carnivores according to geographical origin and technique of processing

Length [μm] Range \pm SD or Mean	Width [μm] Range or Mean	Geographical origin/ Technique of processing	Source
Africa			
121-218	4.5-5.2	Algeria (dog) May-Grünwald-Giemsa staining	Rioche (1960)
255	4.5	Kenya (dog) Knott's technique	Lightner & Reardon (1983)
195-230	5-5.5	Mali (spotted hyaena) Technique not specified	Railliet <i>et al.</i> (1912)
240-260	5-6	Morocco (dog)	Bouin (1921)
Asia			
212-265	3.1-5.7	Pakistan (dog)	Wolfe <i>et al.</i> (1971)
213	4.4	Knott's test Haematoxylin-stained blood films	
Europe			
185-230	5-6	France Technique not specified	Chauve (1990)
199-230	5-6	France Technique not specified	Ducos de Lahitte & Ducos de Lahitte (1990)
237-247	4.2-4.4	Italy (red fox) Unfixed and unstained	Marconcini <i>et al.</i> (1996)
145-233	5.3-7.4	Portugal (dog) Giemsa-stained blood films	Fraga de Azevedo (1943)
233-277	4.5-6	Spain (dog) Knott's technique	Ortega-Mora <i>et al.</i> (1989)
263.51	5.04	Spain (dog) Knott's technique	Valcárcel <i>et al.</i> (1990)