

## REFERENCES

- Abe H, Urao T, Ito T, Seki M, Shinozaki K, Yamaguchi-Shinozaki K. 2003. *Arabidopsis* AtMYC2 (bHLH) and AtMYB2 (MYB) function as transcriptional activators in abscisic acid signaling. *Plant Cell* 15: 63-78.
- Abe K, Emori Y, Kondo H, Suzuki K, Arai S. 1987. Molecular cloning of a cysteine proteinase inhibitor of rice (oryzacystatin) –Homology with animal cystatins and transient expression in the ripening process of rice seeds. *J. Biol. Chem.* 262: 16793-16797.
- Abe K, Kondo H, Watanabe H, Emori Y, Arai S. 1991. Oryzacystatins as the first well-defined cystatins of plant origin and their target proteinases in rice seeds. *Biomed. Biochim. Acta* 50: 637-641.
- Abe M, Abe K, Domoto C, Arai S. 1995. Two distinct species of corn cystatin in corn kernels. *Biosci. Biotech. Biochem.* 59: 756-758.
- Abe M, Abe K, Kuroda M, Arai S. 1992. Corn kernel cysteine proteinase inhibitor as a novel cystatin superfamily member of plant origin molecular cloning and expression studies. *Eur. J. Biochem.* 209: 933-937.
- Abe Y, Shirane K, Yokosawa H, Matsushita H, Mitta M, Kato I, Ishii S. 1993. Asparaginyl endopeptidase of jack bean seeds. Purification, characterization, and high utility in protein sequence analysis. *J. Biol. Chem.* 15: 3525-3529.
- Abrahamson M. 1994. Cystatins. *Methods Enzymol.* 244: 685-700.
- Akasofu H, Yamauchi D, Mitsuhashi W, Minamikawa T. 1989. Nucleotide sequence of cDNA for sulfhydryl-endopeptidase (SH-EP) from cotyledons of germinating *Vigna mungo* seeds. *Nucl. Acids Res.* 17: 6733.

- Alfonso M, Yruela I, Almarcegui S, Torrado E, Perez MA, Picorel R. 2001. Unusual tolerance to high temperatures in a new herbicide-resistant D1 mutant from *Glycine max* (L.) Merr. cell cultures deficient in fatty acid desaturation. *Planta* 212: 573-582.
- Alia HH, Mustardy L, Deshniem P, Ida M, Murata N. 1997. Transformation of *Arabidopsis thaliana* with the *codA* gene for choline oxidase; accumulation of glycinebetaine and enhanced tolerance to salt and cold stress. *Plant J.* 12: 133-142.
- Alia HH, Sakamoto A, Murata N. 1998. Enhancement of the tolerance of *Arabidopsis* to high temperatures by genetic engineering of the synthesis of glycinebetaine. *Plant J.* 16: 155-161.
- Alonso JM, Granell A. 1995. A putative vacuolar processing protease is regulated by ethylene and also during fruit ripening in *Citrus* fruit. *Plant Physiol.* 109: 541-547.
- Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. 1997. GAPPED BLAST and PSI-BLAST: a new generation of protein database search program. *Nucl. Acids Res.* 25: 3389-3402.
- Apse MP, Aharon GS, Snedden WA, Blumwald E. 1999. Salt tolerance conferred by overexpression of a vacuolar Na<sup>+</sup>/H<sup>+</sup> antiport in *Arabidopsis*. *Science* 285: 1256-1258.
- Araus JL, Slafer GA, Reynolds MP, Royo C. 2002. Plant breeding and drought in C3 cereals: What should we breed for? *Ann. Bot. (Lond)* 89: 925-940.
- Artus NN, Uemura M, Steponkus PL, Gilmour SJ, Lin C, Thomashow MF. 1996. Constitutive expression of the cold-regulated *Arabidopsis thaliana* COR15a gene affects both chloroplast and protoplast freezing tolerance. *Proc. Natl. Acad. Sci. USA* 93: 13404-13409.

- Ashouri A, Michaud D, Cloutier C. 2001. Recombinant and classically selected factors of potato plant resistance to the Colorado potato beetle, *Leptinotarsa decemlineata*, variously affect the potato aphid parasitoid *Aphidius nigripes*. *BioControl* 46: 401-418.
- Atkinson HJ, Johnston KA, Robbins M. 2004. Prima facie evidence that a phytocystatin for transgenic plant resistance to nematodes is not a toxic risk in the human diet. *J. Nutr.* 134: 431-434.
- Atkinson HJ, Urwin PE, McPherson MJ. 2003. Engineering plants for nematode resistance. *Ann. Rev. Phytopathol.* 41: 615-639.
- Bajaj S, Targolli J, Liu L-F, Ho T-HD, Wu R. 1999. Transgenic approaches to increase dehydration-stress tolerance in plants. *Mol. Breed.* 5: 493-503.
- Barrett AJ. 1994. Classification of peptidases. In: Barrett AJ, ed. *Methods Enzymol.* New York, Academic Press, 244: 1-15.
- Beers EP, Jones AM, Dickerman AW. 2004. The S8 serine, C1A cysteine and A1 aspartic protease families in *Arabidopsis*. *Phytochem.* 65: 43-58.
- Belenghi B, Acconcia F, Trovato M, Perazzolli M, Bocedi A, Polticelli F, Ascenzi P, Delledonne M. 2003. AtCYS1, a cystatin from *Arabidopsis thaliana*, suppresses hypersensitive cell death. *Eur. J. Biochem.* 270: 2593-2604.
- Benfey NP, Ren L, Chua NH. 1990. Tissue-specific expression from CaMV 35S enhancer subdomains in early stages of plant development. *EMBO J.* 9: 1677-1684.
- Bhalerao R, Keskitalo J, Sterky F, Erlandsson R, Bjorkbacka H, Birve SJ, Karlsson J, Gardstrom P, Lundeberg J, Jansson S. 2003. Gene expression in autumn leaves. *Plant Physiol.* 131: 430-442.

- Bohnert HJ, Nelson DE, Jensen RG. 1995. Adaptations to environmental stresses. *Plant Cell* 7: 1099-1111.
- Bohnert HJ, Sheveleva E. 1998. Plant stress adaptations— making metabolism move. *Curr. Opin. Plant Biol.* 1: 267-274.
- Boominathan P, Shukla R, Kumar A, Manna D, Negi D, Verma PK, Chattopadhyay D. 2004. Long term transcript accumulation during the development of dehydration adaptation in *Cicer arietinum*. *Plant Physiol.* 135: 1608-1620.
- Bordas M, Montesinos C, Debausa M, Salvador A, Roig LA, Serrano R, Moreno V. 1997. Transfer of the yeast salt tolerance gene HAL1 to *Cucumis melo* L. cultivars and *in vitro* evaluation of salt tolerance. *Transgenic Res.* 6: 41-50.
- Boston RS, Viitanen PV, Vierling E. 1996. Molecular chaperones and protein folding in plants. *Plant Mol. Biol.* 32: 191-222.
- Botella MA, Xu Y, Prabha TN, Zhao Y, Narasimhan ML, Wilson KA. 1996. Differential expression of soybean cysteine proteinase inhibitor genes during development and in response to wounding and methyl jasmonate. *Plant Physiol.* 112: 1201-1210.
- Bottari A, Capocchi A, Galleschi L, Jopova A, Saviozzi F. 1996. Asparaginyl endopeptidases during maturation and germination of durum wheat. *Physiol plant.* 97: 475-480.
- Bowler LD, Hubank M, Spratt BG. 1999. Representational difference analysis of cDNA for the detection of differential gene expression in bacteria: development using a model of iron-regulated gene expression in *Neisseria meningitides*. *Microbiology* 145: 3529-3537.
- Boyer JS. 1982. Plant productivity and environment. *Science* 218: 443-448.
- Bradford MM. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of dye-binding. *Anal. Biochem.* 72: 248-254.

- Bray EA, Bailey-Seres J, Weretilnyk E. 2000. Responses to abiotic stresses. *In*: BB Buchanan, W Gruissem, RL Jones, eds, *Biochemistry & Molecular Biology of Plants*, Split edition. American Society of Plant Physiologists, Rockville, MD, pp 1158-1203.
- Bray EA. 1997. Plant responses to water deficit. *Trends Plant Sci.* 2: 48-54.
- Bray EA. 2002. Classification of genes differentially expressed during water-deficit stress in *Arabidopsis thaliana*: an analysis using microarray and differential expression data. *Ann. Bot.* 89: 803-811.
- Broderson P, Petersen M, Pike HM, Olszak B, Skov S, Odum N, Jorgensen LB, Brown RE, Mundy J. 2002. Knockout of *Arabidopsis* ACCELERATED-CELL-DEATH11 encoding a sphingosine transfer protein causes activation of programmed cell death and defense. *Genes Develop.* 16: 490-502.
- Brown WM, Dziegielewska KM. 1997. Friends and relations of the cystatin superfamily new members and their evolution. *Protein Sci.* 6: 5-12.
- Buchanan-Wollaston V, Ainsworth C. 1997. Leaf senescence in *Brassica napus*: cloning of senescence related genes by subtractive hybridisation. *Plant Mol. Biol.* 33: 821-834.
- Callis J. 1995. Regulation of protein degradation. *Plant Cell* 7: 845-857.
- Cannon RJC. 2000. Bt transgenic crops: risks and benefits. *Integr. Pest Manag. Rev.* 5: 151-173.
- Cercos M, Santamaria S, Carbonell J. 1999. Cloning and characterization of TPE4A, a thiol-protease gene induced during ovary senescence and seed germination in pea. *Plant Physiol.* 119: 1341-1348.

- Chartzoulakis K, Patakas A, Kofidis G, Bosabalidis A, Nastou A. 2002. Water stress affects leaf anatomy, gas exchange, water relations and growth of two avocado cultivars. *Scientia Hort.* 1778: 1-13.
- Chaves MM, Pereira JS, Maroco J, Rodrigues ML, Ricardo CP, Osorio ML, Carvalho I, Faria T, Pinheiro C. 2002. How plants cope with water stress in the field. Photosynthesis and growth. *Ann. Bot.* 89: 907-916.
- Chen GH, Huang LT, Yap MN, Lee RH, Huang YJ, Cheng MC, Chen SCG. 2002. Molecular characterization of a senescence associated gene encoding cysteine proteinase and its gene expression during leaf senescence in sweet potato. *Plant Cell Physiol.* 43: 984-991.
- Chen THH, Murata N. 2002. Enhancement of tolerance of abiotic stress by metabolic engineering of betaines and other compatible solutes. *Curr. Opin. Plant Biol.* 5: 250-257.
- Cheong YH, Chang HS, Gupta R, Wang X, Zhu T, Luan S. 2002. Transcriptional profiling reveals novel interactions between wounding, pathogen, abiotic stress, and hormonal responses in *Arabidopsis*. *Plant Physiol.* 129: 661-677.
- Cheong YH, Kim KN, Pandey GK, Gupta R, Grant JJ, Luan S. 2003. CBL1, a calcium sensor that differentially regulates salt, drought, and cold responses in *Arabidopsis*. *Plant Cell* 15: 1833-1845.
- Choi HI, Hong JH, Ha J, Kang JY, Kim SY. 2000. ABFs, a family of ABA-responsive element binding factors. *J. Biol. Chem.* 275: 1723-1730.
- Christeller JT, Burgess EPJ, Mett V, Gatehouse HS, Markwick NP, Murray C, Malone LA, Wright MA, Philip BA, Watt D, Gatehouse LN, Lovei GL, Shannon AL, Phung MM, Watson LM, Laing WA. 2002. The expression of a mammalian proteinase inhibitor,

- bovine spleen trypsin inhibitor in tobacco and its effects on *Helicoverpa armigera* larvae. *Transgenic Res.* 11: 161-173.
- Cipriani G, Fuentes S, Bello V, Salazar LF, Ghislain M, Zhang D. 2000. Transgene expression of rice cysteine proteinase inhibitors for the development of resistance against Sweetpotato Feathery Mottle virus. CIP Program Report 1999-2000, pp. 267-271.
- Contento AL, Kim SJ, Bassham DC. 2004. Transcriptome profiling of the response of *Arabidopsis* suspension culture cells to sucrose starvation. *Plant Physiol.* 135: 2330-2347.
- Corre-Menguy F, Cejudo FJ, Mazubert C, Vidal J, Lelandais-Briere C, Torres G, Rode A, Hartmann C. 2002. Characterization of the expression of a wheat cystatin gene during caryopsis development. *Plant Mol. Biol.* 50: 687-698.
- Coupe SA, Sinclair BK, Watson LM, Heyes JA, Eason JR. 2003. Identification of dehydration-responsive cysteine proteases during post-harvest senescence of broccoli florets. *J. Exp. Bot.* 54: 1045-1056.
- Cowgill SE, Bardgett RD, Kiezebrink DT, Atkinson HJ. 2002b. The effect of transgenic nematode resistance on non-target organisms in the potato rhizosphere. *J. Appl. Ecol.* 39: 915-923.
- Cowgill SE, Wright C, Atkinson HJ. 2002a. Transgenic potatoes with enhanced levels of nematode resistance do not have altered susceptibility to non target aphids. *Mol. Ecol.* 11: 821-827.
- Crafts-Brandner SJ, Salvucci ME. 2000. Rubisco activase constrains the photosynthetic potential of leaves at high temperature and CO<sub>2</sub>. *Proc. Natl. Acad. Sci. USA* 97: 13430-13435.

- Craufurd PQ, Peacock JM. 1993. Effect of heat and drought stress on sorghum. *Exp. Agr.* 29: 77-86.
- Creelman RA, Mason HS, Bensen RJ, Boyer JS, Mullet JE. 1990. Water deficit and abscisic acid cause differential inhibition of shoot versus root growth in soybean seedlings. Analysis of growth, sugar accumulation, and gene expression. *Plant Physiol.* 92: 205-214.
- Cullis CA. 2004. *Plant Genomics and Proteomics*. John Wiley and Sons Inc. Hoboken New Jersey.
- Cushman JC, Bohnert HJ. 2000. Genomic approaches to plant stress tolerance. *Curr. Opin. Plant Biol.* 3: 117-124.
- Dash S, Mohanty N. 2002. Response of seedlings to heat-stress in cultivars of wheat: Growth temperature-dependent differential modulation of photosystem 1 and 2 activity, and foliar antioxidant defense capacity. *J. Plant Physiol.* 159: 49-59.
- Dat JF, Lopez-Delgado H, Foyer CH, Scott IM. 1998. Parallel changes in H<sub>2</sub>O<sub>2</sub> and catalase during thermotolerance induced by salicylic acid or heat acclimation in mustard seedlings. *Plant Physiol.* 116: 1351-1357.
- Davison PA, Hunter CN, Horton P. 2002. Overexpression of beta-carotene hydroxylase enhances stress tolerance in *Arabidopsis*. *Nature* 418: 203-206.
- De Barros EG, Larkins BA. 1994. Cloning of a cDNA encoding a putative cysteine protease gliadin during wheat germination. *Phytochemistry* 43Z: 39-44
- De Leo F, Volpicella M, Licciulli F, Liuni S, Gallerani R, Ceci LR. 2002. PLANT-PIs: A database for plant protease inhibitors and their genes. *Nucl. Acids Res.* 30: 347-348.



- De Rocher EJ, Vargo-Gogola TC, Diehn SH, Green PJ. 1998. Direct evidence for rapid degradation of *Bacillus thuringiensis* toxin mRNA as a cause of poor expression in plants. *Plant Physiol.* 117: 1445-1461.
- Delledonne M, Allegro G, Belenghi B, Balestrazzi A, Picco F, Levine A, Zelasco S, Calligari P, Confalonieri M. 2001. Transformation of white poplar (*Populus alba* L.) with a novel *Arabidopsis thaliana* cysteine proteinase inhibitor and analysis of insect pest resistance. *Mol. Breed.* 7: 35-42.
- Diop NN, Kidric M, Repellin A, Gareil M, d'Arcy-Lameta A, Pham Thi AT, Zuily-Fodil Y. 2004. A multicystatin is induced by drought-stress in cowpea (*Vigna unguiculata* (L.) Walp) leaves. *FEBS Lett.* 577: 545-550.
- Dolja VV, Peremyslov VV, Keller KE, Martin RR, Hong J. 1998. Isolation and stability of histidine-tagged proteins produced in plants via potyvirus gene vectors. *Virology* 252: 269-274.
- Dombrowski JE. 2003. Salt stress activation of wound-related genes in tomato plants. *Plant Physiol.* 132: 2098-2107.
- Down RE, Ford L, Bedford SJ, Gatehouse LN, Newell C, Gatehouse JA, Gatehouse AMR. 2001. Influence of plant development and environment on transgene expression in potato and consequences for insect resistance. *Transgenic Res.* 10: 223-236.
- Downing WL, Mauxion F, Fauvarque MO, de Vienne D, Vartanian N, Giraudat J. 1992. A *Barassica napus* transcript encoding a protein related to the Kunitz protease inhibitor family accumulates upon water stress in leaves, not in seeds. *Plant J.* 5: 685-693.

- Drake R, John I, Farrell A, Cooper W, Schuch W, Grierson D. 1996. Isolation and analysis of DNAs encoding tomato cysteine proteinases expressed during leaf senescence. *Plant Mol. Biol.* 30: 755-767.
- Eason JR, Ryan DJ, Pinkney TT, O'Donoghue EM. 2002. Programmed cell death during flower senescence: isolation and characterization of cysteine proteinases from *Sandersonia aurantiaca*. *Funct. Plant Biol.* 29: 1-10.
- Eason JR, Ryan DJ, Watson LM, Hedderley D, Christey MC, Braun RH, Coupe SA. 2005. Suppression of the cysteine protease, aleurain, delays floret senescence in *Brassica oleracea*. *Plant Mol. Biol.* 57: 645-657.
- Eckes P, Vajtewaal B, Donn G. 1989. Synthetic gene confers resistance to the broad spectrum herbicide L- phosphinothricin in plants (supplement 13 D). *J. Cell Biochem.*
- Elhag GA, Thomas FJ, McCreery TP, Bourque DP. 1992. Nuclear-encoded chloroplast ribosomal protein L12 of *Nicotiana tabacum*: characterization of mature protein and isolation and sequence analysis of cDNA clones encoding its cytoplasmic precursor. *Nucl. Acids Res.* 20: 689-697.
- Fischer J, Becker C, Hillmer S, Horstmann C, Neubohn B, Schlereth A, Senyuk V, Shutov A, Muntz K. 2000. The families of papain- and legumain-like cysteine proteinases from embryonic axes and cotyledons of *Vicia* seeds: developmental patterns, intracellular localization and functions in globulin proteolysis. *Plant Mol. Biol.* 43: 83-101.
- Fowler S, Thomashow MF. 2002. Arabidopsis transcriptome profiling indicates that multiple regulatory pathways are activated during cold acclimation in addition to the CBF cold response pathway. *Plant Cell* 14: 1675-1690.

- Foyer CH, Descourviere P, Kunert KJ. 1994. Protection against oxygen radicals: an important defence mechanism studied in transgenic plants. *Plant Cell Environ.* 17: 507-523.
- Fujita Y, Fujita M, Satoh R, Maruyama K, Parvez M, Seki M, Hiratsu K, Ohme-Takagi M, Shinozaki K, Yamaguchi-Shinozaki K. 2005. AREB1 Is a transcription activator of novel ABRE-dependent ABA signaling that enhances drought stress tolerance in *Arabidopsis*. *Plant Cell* 17: 3470-3488.
- Funk V, Kositsup B, Zhao C, Beers EP. 2002. The *Arabidopsis* xylem peptidase XCP1 is a tracheary element vacuolar protein that may be a papain ortholog. *Plant Physiol.* 128: 84-94.
- Gaddour K, Vicente-Carbajosa J, Lara P, Isabel-Lamoneda I, Diaz I, Carbonero P. 2001. A constitutive cystatin-encoding gene from barley (*Icy*) responds differentially to abiotic stimuli. *Plant Mol. Biol.* 45: 599-608.
- Gan S, Amasino RM. 1997. Making sense of senescence. Molecular genetic regulation and manipulation of leaf senescence. *Plant Physiol.* 113: 313-319.
- Gasteiger E, Gattiker A, Hoogland C, Ivanyi I, Appel RD, Bairoch A. 2003. ExPASy: the proteomics server for in-depth protein knowledge and analysis. *Nucl. Acids Res.* 31: 3784-3788.
- Gaxiola RA, Li J, Undurraga S, Dang LM, Allen GJ, Alper SL, Fink GR. 2001. Drought- and salt-tolerant plants result from overexpression of the AVP1 H<sup>+</sup>-pump. *Proc. Natl. Acad. Sci. USA* 98: 1444-11449.
- Gepstein S, Sabehi G, Carp M-J, Hajouj T, Neshar MFO, Yariv I, Dor C, Bassani M. 2003. Large-scale identification of leaf senescence-associated genes. *Plant J.* 36: 629-642.

- Gietl C, Schmid M. 2001. Ricinosomes: an organelle for developmentally regulated programmed cell death in senescing plant tissue. *Naturwissenschaften* 88: 49-58.
- Gilmour SJ, Sebolt AM, Salazar MP, Everard JD, Thomashow MF. 2000. Overexpression of the arabidopsis CBF3 transcriptional activator mimics multiple biochemical changes associated with cold acclimation. *Plant Physiol.* 124: 1854-1865.
- Girard C, Bonadé-Bottino M, Pham-Delegue MH, Jouanin L. 1998. Two strains of cabbage seed weevil (*Coleoptera: Curculionidae*) exhibit differential susceptibility to a transgenic oilseed rape expressing oryzacystatin I. *J. Insect Physiol.* 44: 569-577.
- Godde D. 1999. Adaptations of the photosynthetic apparatus to stress conditions. In: Lerner HR, ed. Plant responses to environmental stresses. New York, NY: Marcel Dekker, 449-474.
- Görlach J, Volrath S, Knauf-Beiter G, Hengy G, Beckhove U, Kogel KH, Oostendorp M, Staub T, Ward E, Kessmann H, Ryals J. 1996. Benzothiadiazole, a novel class of inducers of systemic acquired resistance, activates gene expression and disease resistance in wheat. *Plant Cell* 8: 629-643.
- Granell A, Cercos M, Carbonell J. 1998. Plant cysteine proteinases in germination and senescence. In: AJ Barrett, ND Rawlings, JF Woessner (Eds.), The Handbook of Proteolytic Enzymes, Academic Press, San Diego, pp. 578-583.
- Greene JK, Turnipseed SG, Sullivan MJ, Herzog GA. 1999. Boll damage by southern greenstink bug (*Hemiptera: Pentatomidae*) and tarnished plant bug (*Hemiptera: Miridae*) caged on transgenic *Bacillus thuringiensis* cotton. *J. Econ. Entomol.* 92: 941-944.

- Griffiths CM, Hosken SE, Oliver D, Chojeki J, Thomas H. 1997. Sequencing, expression pattern and RFLP mapping of a senescence-enhanced cDNA from *Zea mays* with high homology to oryzain gamma and aleurain. *Plant Mol. Biol.* 34: 815-821.
- Grudkowska M, Zagdańska B. 2004. Multifunctional role of plant cysteine proteinases. *Acta Biochem. Polon.* 51: 609-624.
- Gruis DF, Selinger DA, Curran JM, Jung R. 2002. Redundant proteolytic mechanisms process seed storage proteins in the absence of seed-type members of the vacuolar processing enzyme family of cysteine proteases. *Plant Cell* 14: 2863-2882.
- Guerrero C, Calle M, Reid MS, Valpuesta V. 1998. Analysis of the expression of two thiolprotease genes from daylily (*Heemerocallis* spp.) during flower senescence. *Plant Mol. Biol.* 36: 565-571.
- Guo Y, Cai Z, Gan S. 2004. Transcriptome of Arabidopsis leaf senescence. *Plant Cell Environ.* 27: 521-549.
- Gupta AS, Heinen JL, Holaday AS, Burke JJ, Allen RD. 1993a. Increased resistance to oxidative stress in transgenic plants that overexpress chloroplastic Cu/Zn superoxide dismutase. *Proc. Natl. Acad. Sci. USA* 90: 1629-1633.
- Gupta AS, Webb RP, Holaday AS, Allen RD. 1993b. Overexpression of superoxide dismutase protects plants from oxidative stress. *Plant Physiol.* 103: 1067-1073.
- Gutierrez-Campos R, Torres-Acosta J, Saucedo-Arias LJ, Gomez-Lim MA. 1999. The use of cysteine proteinase inhibitors to engineer resistance against potyviruses in transgenic tobacco plants. *Nature Biotech.* 17:1223-1226.

- Guy C. 1999. The influence of temperature extremes on gene expression, genomic structure, and the evolution of induced tolerance in plants. *In*: Lerner HR, ed. Plant responses to environmental stresses. New York, NY: Marcel Dekker, 497-548.
- Haake V, Cook D, Riechmann JL, Pineda O, Thomashow MF, Zhang JZ. 2002. Transcription factor CBF4 is a regulator of drought adaptation in *Arabidopsis*. *Plant Physiol.* 130: 639-648.
- Hall AE. 2001. Crop Responses to the Environment. CRC Press, Boca Raton, FL.
- Hammond JP, Bennett MJ, Bowen HC, Broadley MR, Eastwood DC, May ST, Rahn C, Swarup R, Woolaway KE, White PJ. 2003. Changes in gene expression in arabidopsis shoots during phosphate starvation and the potential for developing smart plants. *Plant Physiol.* 132: 578-596.
- Haq SK, Atif SM, Khan RH. 2004. Protein proteinase inhibitor genes in combat against insects, pests, and pathogens: natural and engineered phytoprotection. *Arch. Biochem. Biophys.* 431: 145-159.
- Hara-Nishimura I, Hatsugai N, Nakaune S, Kuroyanagi M, Nishimura M. 2005. Vacuolar processing enzyme: an executor of plant cell death. *Curr. Opin. Plant Biol.* 8: 404-408.
- Hara-Nishimura I, Maeshima M. 2000. Vacuolar processing enzymes and aquaporins. *In*: Vacuolar Compartments in Plants, ADG Robinson, JC Rogers, eds (London: Sheffield Academic Press) pp. 20-42.
- Härndahl U, Hall RB, Osteryoung KW, Vierling E, Bornman JF, Sundby C. 1999. The chloroplast small heat shock protein undergoes oxidation-dependent conformational changes and may protect plants from oxidative stress. *Cell Stress Chaperones* 4: 129-138.

- Harrak H, Azelmat S, Baker EN, Tabaeizadeh Z. 2001. Isolation and characterization of gene encoding a drought-induced cysteine protease in tomato (*Lycopersicon esculentum*). *Genome* 44: 368-374.
- Hasegawa K, Yukawa Y, Sugita M, Sugiura M. 2002. Organization and transcription of the gene family encoding chlorophyll a/b-binding proteins in *Nicotiana sylvestris*. *Gene* 289: 161-168.
- Hatsugai N, Kuroyanagi M, Yamada K, Meshi T, Tsuda S, Kondo M, Nishimura M, Hara-Nishimura I. 2004. A plant vacuolar protease, VPE, mediates virus-induced hypersensitive cell death. *Science* 305: 855-858.
- He C, Yan J, Shen G, Fu L, Holaday A S, Auld D, Blumwald E, Zhang H. 2005. Expression of an Arabidopsis vacuolar sodium/proton antiporter gene in cotton improves photosynthetic performance under salt conditions and increases fiber yield in the field. *Plant Cell Physiol.* 46: 1848-1854.
- He X, Kermode AR. 2003. Proteases associated with programmed cell death of megagametophyte cells after germination of white spruce (*Picea glauca*) seeds. *Plant Mol. Biol.* 52: 729-744.
- Hensel LL, Grbic V, Baumgarten DA, Bleeker AB. 1993. Developmental and age- related processes that influence the longevity and senescence of photosynthetic tissues in Arabidopsis. *Plant Cell* 5: 553-564.
- Herde O, Atzorn R, Fisahn J, Wasternack C, Willmitzer L, Peña-Cortés H. 1996. Localized wounding by heat initiates the accumulation of proteinase inhibitor II in abscisic acid-deficient plants by triggering jasmonic acid biosynthesis. *Plant Physiol.* 112: 853-860.

- Hermesmeier D, Schittko U, Baldwin IT. 2001. Molecular interactions between the specialist herbivore *Manduca sexta* (Lepidoptera, Sphingidae) and its natural host *Nicotiana attenuata*: large-scale changes in the accumulation of growth- and defense-related plant mRNAs. *Plant Physiol.* 125: 683-700.
- Hilder VA, Boulter D. 1999. Genetic engineering of crop plants for insect resistance - a critical review. *Crop Protection* 18: 177-191.
- Hildmann T, Ebner M, Peña-Cortés H, Sánchez-Serrano JJ, Willmitzer L, Prat S. 1992. General roles of abscisic and jasmonic acid in gene activation as a result of mechanical wounding. *Plant Cell* 4: 1157-1170.
- Himelblau E, Amasino RM. 2001. Nutrients mobilized from leaves of *Arabidopsis thaliana* during leaf senescence. *J. Plant Physiol.* 158: 1317-1323.
- Hines ME, Osuala CI, Nielsen SS. 1992. Screening for cysteine proteinase inhibitor activity in legume seeds. *J. Sci. Food Agric.* 59: 4555-4557.
- Hirai MY, Fujiwara T, Awazuhara M, Kimura T, Noji M, Saito K. 2003. Global expression profiling of sulfur-starved *Arabidopsis* by DNA microarray reveals the role of O-acetyl-L-serine as a general regulator of gene expression in response to sulfur nutrition. *Plant J.* 33: 651-663.
- Hong SW, Vierling E. 2000. Mutants of *Arabidopsis thaliana* defective in the acquisition of tolerance to high temperature stress. *Proc. Natl. Acad. Sci. USA* 97: 4392-4397.
- Hong SW, Vierling E. 2001. Hsp101 is necessary for heat tolerance but dispensable for development and germination in the absence of stress. *Plant J.* 27: 25-35.



- Hong Z, Lakkineni K, Zhang K, Verma DPS. 2000. Removal of feedback inhibition of <sup>1</sup>-pyrroline-5-carboxylate synthetase results in increased proline accumulation and protection of plants from osmotic stress. *Plant Physiol.* 122: 1129-1136.
- Hortensteiner S, Feller U. 2002. Nitrogen metabolism and remobilisation during senescence. *J. Exp. Bot.* 53: 927-937.
- Hoth S, Morgante M, Sanchez JP, Hanafey MK, Tingey SV, Chua NH. 2002. Genome-wide gene expression profiling in *Arabidopsis thaliana* reveals new targets of abscisic acid and largely impaired gene regulation in the *abi1-1* mutant. *J. Cell Sci.* 115: 4891-4900.
- Hsieh TH, Lee JT, Yang PT, Chiu LH, Charng YY, Wang YC, Chan MT. 2002. Heterology expression of the *Arabidopsis* C-repeat/dehydration response element binding factor 1 gene confers elevated tolerance to chilling and oxidative stresses in transgenic tomato. *Plant Physiol.* 129: 1086-1094.
- Hubank M, Schatz DG. 1994. Identifying differences in mRNA expression by representational difference analysis of cDNA. *Nucl. Acids Res.* 22: 5640-5648.
- Hubank M, Schatz DG. 1999. cDNA Representational Difference Analysis: a sensitive and flexible method for identification of differentially expressed genes. *Methods Enzymol.* 303: 325-349.
- Humbeck K, Quast S, Krupinska K. 1996. Functional and molecular changes in the photosynthetic apparatus during senescence of flag leaves from field-grown barley plants. *Plant Cell Environ.* 19: 337-344.
- Iba K. 2002. Acclimative response to temperature stress in higher plants: Approaches of gene engineering for temperature tolerance. *Annu. Rev. Plant Biol.* 53: 225-245.

- Ingram J, Bartels D. 1996. The molecular basis of dehydration tolerance in plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 47: 377-403.
- Irie K, Hosoyama H, Takeuchi T, Iwabuchi K, Watanabe H, Abe M, Abe K, Arai S. 1996. Transgenic rice established to express corn cystatin exhibits strong inhibitory activity against insect gut proteinases. *Plant Mol. Biol.* 30: 149-157.
- Ito Y, Katsura K, Maruyama K, Tajiri T, Kobayashi M, Seki S, Shinozaki K, Yamaguchi-Shinozaki K. 2006. Functional analysis of Rice DREB1/CBF-type transcription factors involved in cold-responsive gene expression in transgenic rice. *Plant Cell Physiol.* 47: 141-153.
- Jaglo KR, Kleff S, Amundsen KL, Zhang X, Haake V, Zhang JZ, Deits T, Thomashow MF. 2001. Components of the *Arabidopsis* C-repeat/dehydration-responsive element binding factor cold-response pathway are conserved in *Brassica napus* and other plant species. *Plant Physiol.* 127: 910-917.
- Jaglo-Ottosen KR, Gilmour SJ, Zarka DG, Schabenberger O, Thomashow MF. 1998. *Arabidopsis* CBF1 overexpression induces COR genes and enhances freezing tolerance. *Science* 280: 104-106.
- Jagtap V, Bhargava S, Streb P, Feierabend J. 1998. Comparative effect of water, heat and light stresses on photosynthetic reactions in *Sorghum bicolor* (L.) Moench. *J. Exp. Bot.* 49: 1715-1721.
- Jefferson RA, Kavanagh TA, Bevan MW. 1987. Gus fusion:  $\beta$ -glucuronidase as a sensitive and versatile gene marker in higher plants. *EMBO J.* 6: 3901-3907.
- Jiang Y, Huang B. 2001. Drought and heat stress injury to two cool season turf grasses in relation to antioxidant metabolism and lipid peroxidation. *Crop Sci.* 41: 436-442.

- Jones JT, Mullet JE. 1995. A salt and dehydration inducible pea gene *Cyp15a*, encodes a cell-wall protein with sequence similarity to cysteine proteinase. *Plant Mol. Biol.* 28: 1055-1065.
- Jongsma MA, Bolter C. 1997. The adaptation of insects to plant protease inhibitors. *J. insect physiol.* 43: 885-895.
- Kamel A, Loser DM. 1995. Contribution of carbohydrates and other solutes to osmotic adjustment in wheat leaves under water stress. *J. Plant Physiol.* 145: 363-366.
- Kampinga HH, Brunsting JF, Stege GJ, Burgman PW, Konings AW. 1995. Thermal protein denaturation and protein aggregation in cells made thermotolerant by various chemicals: role of heat shock proteins. *Exp. Cell Res.* 219: 536-546.
- Kang JY, Choi HI, Im MY, Kim SY. 2002. *Arabidopsis* basic leucine zipper proteins that mediate stress-responsive abscisic acid signaling. *Plant Cell* 14: 343-357.
- Kardailsky IV, Brewin NJ. 1996. Expression of cysteine protease genes in pea nodules development and senescence. *Mol. Plant-Microbe Interact.* 9: 689-695.
- Karrer KM, Peiffer SL, DiTomas ME. 1993. Two distinct gene subfamilies of cysteine proteinase gene. *Proc. Natl. Acad. Sci. USA* 90: 3063-3067.
- Kasuga M, Liu Q, Miura S, Yamaguchi-Shinozaki K, Shinozaki K. 1999. Improving plant drought, salt, and freezing tolerance by gene transfer of a single stress-inducible transcription factor. *Nature Biotech.* 17: 287-291.
- Kasuga M, Miura S, Shinozaki K, Yamaguchi-Shinozaki K. 2004. A combination of the *Arabidopsis* DREB1A gene and stress-inducible rd29A promoter improved drought- and low-temperature stress tolerance in tobacco by gene transfer. *Plant Cell Physiol.* 45: 346-350.

- Katiyar-Agarwal S, Agarwal M, Grover A. 2003. Heat-tolerant basmati rice engineered by overexpression of *hsp101*. *Plant Mol. Biol.* 51: 677-686.
- Kawaguchi R, Girke T, Bray EA, Bailey-Serres JN. 2004. Differential mRNA translation contributes to gene regulation under non-stress and dehydration stress conditions in *Arabidopsis thaliana*. *Plant J.* 38: 823-839.
- Kim S, Hong YN, An CS, Lee KW. 2001. Expression characteristics of serine proteinase inhibitor II under variable environmental stresses in hot pepper (*Capsicum annum* L). *Plant Sci.* 161: 27-33.
- Kingston-Smith AH, Bollard AL, Minchin FR. 2005. Stress-induced changes in protease composition are determined by nitrogen supply in non-nodulating white clover. *J. Exp. Bot.* 56: 745-753.
- Kinoshita T, Yamada K, Hiraiwa N, Nishimura M, Hara-Nishimura I. 1999. Vacuolar processing enzyme is up-regulated in the lytic vacuoles of vegetative tissues during senescence and under various stressed conditions. *Plant J.* 19: 43-53.
- Kishor KPB, Hong Z, Miao GH, Hu CAA, Verma DPS. 1995. Overexpression of  $\delta^1$ -pyrroline-5-carboxylate synthetase increase proline production and confers osmotolerance in transgenic plants. *Plant Physiol.* 108: 1387-1394.
- Koiwa H, Bressan RA, Hasegawa PM. 1997. Regulation of protease inhibitors and plant defense. *Trends Plant Sci.* 2: 379-384.
- Koizumi M, Yanaguchi-Shinozaki K, Tsuji H, Shinozaki K. 1993. Structure and expression of two genes that encode distinct drought-inducible cysteine proteinases in *Arabidopsis thaliana*. *Gene* 129: 175-182.

- Kondo H, Abe K, Arai S. 1989. Immunoassay of oryzacystatin occurring in rice seeds during maturation and germination. *Agric. Biol. Chem.* 53: 2949-2954.
- Konno K, Hirayama C, Nakamura M, Tateishi K, Tamura Y, Hattori M, Kohno K. 2003. Papain protects papaya trees from herbivorous insects: role of cysteine proteinases in latex. *Plant J.* 37: 370-378.
- Konstantinova T, Parvanova D, Atanassov A, Djilianov D. 2002. Freezing tolerant tobacco, transformed to accumulate osmoprotectants. *Plant Sci.* 163:157-164.
- Kramer PJ. 1980. Drought, stress, and the origin of adaptation. *In: Adaptation of plants to water and high temperature stress* (NC Turner and PJ Kramer eds.). John Wiley, New York, pp. 7-20
- Kreps JA, Wu Y, Chang H-S, Zhu T, Wang X, Harper JF. 2002. Transcriptome changes for *Arabidopsis* in response to salt, osmotic, and cold stress. *Plant Physiol.* 130: 2129-2141.
- Krüger J, Thomas CM, Golstein C, Dixon MS, Smoker M, Tang S, Mulder L, Jones JD. 2002. A tomato cysteine protease required for Cf-2-dependent disease resistance and suppression of autonecrosis. *Science* 296: 744-747.
- Kuroda M, Ishimoto M, Suzuki K, Kondo H, Abe K, Kitamura K, Arai S. 1996. Oryzacystatins exhibit growth-inhibitory and lethal effects on different species of bean insect pests *Callosobruchus chinensis* (Coleoptera) and *Riptortus clavatus* (Hemiptera). *Biosci. Biotechnol. Biochem.* 60: 209-212.
- Kuroda M, Kiyosaki T, Matsumoto I, Misaka T, Arai S, Abe K. 2001. Molecular cloning, characterization and expression of wheat cystatins. *Biosci. Biotech. Biochem.* 65: 22-28.
- Laemmli UK. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227: 680-685.

- Larkindale J, Hall JD, Knight MR, Vierling E. 2005. Heat stress phenotypes of *Arabidopsis* mutants implicate multiple signaling pathways in the acquisition of thermotolerance. *Plant Physiol.* 138: 882-897.
- Larkindale J, Knight MR. 2002. Protection against heat stress induced oxidative damage in *Arabidopsis* involves calcium, abscisic acid, ethylene, and salicylic acid. *Plant Physiol.* 128: 682-695.
- Law R, Crafts-Brandner SJ. 1999. Inhibition and acclimation of photosynthesis to heat stress is closely correlated with activation of ribulose-1,5-bisphosphate carboxylase/oxygenase. *Plant Physiol.* 120: 173-182.
- Lawlor DW, Cornic G. 2002. Photosynthetic carbon assimilation and associated metabolism in relation to water deficits in higher plants. *Plant Cell Environ.* 25: 275-294.
- Lawlor DW. 2002. Limitation to photosynthesis in water-stressed leaves: stomata vs metabolism and the role of ATP. *Ann. Bot.* 89: 871-885.
- Lecardonnell A, Chauvin L, Jouanin L, Beaujean A, Prevost G, Sangwan-Norreel X. 1999. Effects of rice cystatin I expression in transgenic potato on Colorado potato beetle larvae. *Plant Sci.* 140: 71-79.
- Lee JH, Hübel A, Schöffl F. 1995. Derepression of the activity of genetically engineered heat shock factor causes constitutive synthesis of heat shock proteins and increased thermotolerance in transgenic *Arabidopsis*. *Plant J.* 8: 603-612.
- Lepilé JC, Bonadé-Bottino M, Augustin S, Pilate G, Letan VD, Delplanque A. 1995. Toxicity to *Chrysomela tremulae* (Coleoptera: chrysomelidae) of transgenic poplars expressing a cysteine proteinase inhibitor. *Mol. Breed.* 1: 319-328.

- Levitt, J. 1980. Responses of plants to environmental stresses. *In: Physiological Ecology* (Kozlowski, T.T., Ed.) New York: Academic Press, pp. 347-448.
- Li Y, Lu YY. 2002. Applying a highly specific and reproducible cDNA RDA method to clone garlic up-regulated genes in human gastric cancer cells. *World J. Gastroent.* 8: 213-216.
- Liang C, Brookhart G, Feng GH, Reeck GR, Kramer KJ. 1991. Inhibition of digestive proteinases of stored grain Coleoptera by oryzacystatin, a cysteine proteinase inhibitor from rice seed. *FEBS Lett.* 278: 139-142.
- Lichtenthaler HK. 1998. The stress concept in plants: an introduction. *Ann. N.Y. Acad. Sci.* 851: 187-198.
- Lidgett AJ, Moran M, Wong KAL, Furze J, Rhodes MJC, Hamill JD. 1995. Isolation and expression pattern of a cDNA encoding a cathepsin B-like protease from *Nicotiana rustica*. *Plant Mol. Biol.* 29: 379-384.
- Lilley CJ, Urwin PE, Katherine KA, Howard J, Atkinson HJ. 2004. Preferential expression of a plant cystatin at nematode feeding sites confers resistance to *Meloidogyne incognita* and *Globodera pallida*. *Plant Biotech. J.* 2: 3-12.
- Lim CO, Lee ST, Chung WS, Park SH, Hwang I, Cho MJ. 1996. Characterization of a cDNA encoding cysteine proteinase inhibitor from Chinese cabbage (*Brassica campestris* L. spp. *Pekinensis*) flower buds. *Plant Mol. Biol.* 30: 373-379.
- Lin JF, Wu SH. 2004. Molecular events in senescing *Arabidopsis* leaves. *Plant J.* 39: 612-628.
- Ling J, Kojima T, Shiraiwa M, Takahara H. 2003. Cloning of two cysteine proteinases genes, *CysP1* and *CysP2*, from soybean cotyledons by cDNA representational difference analysis. *Biochim. Biophys. Acta* 1627: 129-139.

- Linthorst HJM, Van der Does C, Brederode FT, Bol JF. 1993. Circadian expression and induction by wounding of tobacco genes for cysteine proteinase. *Plant Mol. Biol.* 21: 685-694.
- Lisitsyn N, Wigler M. 1993. Cloning the differences between two complex genomes. *Science* 259: 946-951.
- Liu X, Kasuga M, Sakuma Y, Abe H, Miura S, Yamaguchi- Shinozaki K, Shinozaki K. 1998. The transcription factors, DREB1 and DREB2, with an EREBP/AP2 DNA binding domain separate two cellular signal transduction pathways in drought and low-temperature-responsive gene expression, respectively, in *Arabidopsis*. *Plant Cell* 10: 1391-1406.
- Lobell DB, Asner GP. 2003. Climate and management contributions to recent trends in U.S. agricultural yields. *Science* 299: 1032.
- Lohman KN, Gan S, John MC, Amasino RM. 1994. Molecular analysis of natural leaf senescence in *Arabidopsis thaliana*. *Physiol. Plant.* 92: 322-328.
- MacKinney G. 1941. Absorption of light by chlorophyll solutions. *J. Biol. Chem.* 140: 315-322.
- Majoul T, Bancel E, Triboï E, Ben Hamida J, Branlard G. 2003. Proteomic analysis of the effect of heat stress on hexaploid wheat grain: Characterization of heat-responsive proteins from total endosperm. *Proteomics* 3: 175-183.
- Malik MK, Slovin JP, Hwang CH, Zimmerman JL. 1999. Modified expression of a carrot small heat shock protein gene, Hsp17.7, results in increased or decreased thermotolerance. *Plant J.* 20: 89-99.



- Margis R, Reis EM, Villeret V. 1998. Structural and phylogenetic relationships among plant and animal cystatins. *Arch. Biochem. Biophys.* 359: 24-30.
- Maroco JP, Edwards GE, Ku MSB. 1999. Photosynthetic acclimation of maize to growth under elevated levels of carbon dioxide. *Planta* 121: 115-125.
- Marschner H. 1995. Mineral nutrition of higher plants. Academic press London.
- Martiez M, Abraham Z, Carbonero P, Diaz I. 2005. Comparative phylogenetic analysis of cystatin gene families from arabidopsis, rice and barley. *Mol. Gen. Genom.* 273: 423-432.
- Martinez M, Rubio-Somoza I, Carbonero P, Diaz I. 2003. A cathepsin B-like cysteine protease gene from *Hordeum vulgare* (gene *CatB*) induced by GA in aleurone cells is under circadian control in leaves. *J. Exp. Bot.* 54: 951-959.
- Maruyama K, Sakuma Y, Kasuga M, Ito Y, Seki M, Goda H, Shimada Y, Yoshida Y, Shinozaki K, Yamaguchi-Shinozaki K. 2004. Identification of cold-inducible downstream genes of the *Arabidopsis* DREB1A/CBF3 transcriptional factor using two microarray systems. *Plant J.* 38: 982-993.
- Massonneau A, Condamine P, Wisniewski JP, Zivy M, Rogowsky PM. 2005. Maize cystatins respond to developmental cues, cold stress and drought. *Bioch. Biophys. Acta* 1729: 186-199.
- McKersie BD, Bowley SR, Harjanto E, Leprince O. 1996. Water-deficit tolerance and field performance of transgenic alfalfa overexpressing superoxide dismutase. *Plant Physiol.* 111: 1177-1181.
- McKersie BD, Bowley SR, Jones KS. 1999. Winter survival of transgenic alfalfa overexpressing superoxide dismutase. *Plant Physiol.* 119: 839-848.

- McKersie BD, Murnaghan J, Jones KS, Bowley SR. 2000. Iron-superoxide dismutase expression in transgenic alfalfa increases winter survival without a detectable increase in photosynthetic oxidative stress tolerance. *Plant Physiol.* 122: 1427–1438.
- Meyer P, Saedler H. 1996. Homology-dependent gene silencing in plants. *Ann. Rev. Plant Physiol. Plant Mol. Biol.* 47: 23-48.
- Michaud D, Faye L, Yelle S. 1993b. Electrophoretic analysis of plant cysteine and serine proteinases using gelatin-containing polyacrylamide gels and class specific proteinase inhibitors. *Electrophoresis* 14: 94-98.
- Michaud D, Nguyen-Quoc B, Bernier-Vadnais N, Faye L, Yelle S. 1994. Cysteine proteinase forms in sprouting potato tuber. *Physiol. Plant.* 90: 497-503.
- Michaud D, Nguyen-Quoc B, Yelle S. 1993a. Selective inhibition of colorado potato beetle cathepsin H by oryzacystatins I and II. *FEBS Lett.* 331: 173-176.
- Minami A, Fukuda H. 1995. Transient and specific expression of a cysteine endopeptidase associated with autolysis during differentiation of *Zinnia* mesophyll cells into tracheary elements. *Plant Cell Physiol.* 36: 1599-1606.
- MINITAB for Windows Release 12.21, Minitab Inc. 1995.
- Misaka T, Kuroda M, Iwabuchi K, Abe K, Arai S. 1996. Soyacystatin, a novel cysteine proteinase inhibitor in soybean, is distinct in protein structure and gene organization from other cystatins of animal and plant origin. *Eur. J. Biochem.* 240: 609-614.
- Mishra SK, Tripp J, Winkelhaus S, Tschiersch B, Theres K, Nover L, Scharf KD. 2002. In the complex family of heat stress transcription factors, HsfA1 has a unique role as master regulator of thermotolerance in tomato. *Genet. Develop.* 16: 1555-1567.

- Mitsuhashi W, Yamashita T, Toyomasu T, Kashiwagi Y, Konnai T. 2004. Sequential development of cysteine proteinase activities and gene expression during somatic embryogenesis in carrot. *Biosci. Biotech. Biochem.* 68: 705-712.
- Mittler R. 2002. Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci.* 9: 405-410.
- Moura DS, Ryan CA. 2001. Wound-inducible proteinase inhibitors in pepper: differential regulation upon wounding, systemin, and methyl jasmonate. *Plant Physiol.* 126: 289-298.
- Murdock LL, Brookhart G, Dunn PE, Foard DE, Kelley S, Kitch L, Shade RE, Shukle RH, Wolfson JL. 1987. Cysteine digestive proteinases in Coleoptera. *Comparative Biochem. Physiol.* 87: 783-787.
- Nagata K, Kudo N, Abe K, Arai S, Tanokura M. 2000. Three-dimensional solution of oryzacystatin-I, a cysteine proteinase inhibitor of rice, *Oryza sativa* L. japonica. *Biochemistry* 39:14753-14760.
- Naito Y, Fujie M, Usami S, Murooka Y, Yamada T. 2000. The involvement of cysteine proteinase in the nodule development in Chinese milk vetch infected with *Mesorhizobium huakuii* subsp. *rengei*. *Plant Physiol.* 124: 1087-1096.
- Nakaune S, Yamada K, Kondo M, Kato T, Tabata S, Nishimura M, Hara-Nishimura I. 2005. A vacuolar processing enzyme,  $\delta$ vpe, is involved in seed coat formation at the early stage of seed development. *Plant Cell* 17: 876-887.
- Nanjo T, Kobayashia M, Yoshibab Y, Kakubaric Y, Yamaguchi-Shinozaki K, Shinozaki K. 1999. Antisense suppression of proline degradation improves tolerance to freezing and salinity in *Arabidopsis thaliana*. *FEBS Lett.* 461: 205-210.

- Ndong C, Danyluk J, Wilson KE, Pocock T, Huner NPA, Sarhan F. 2002. Cold-regulated cereal chloroplast late embryogenesis abundant-like proteins. Molecular characterization and functional analyses. *Plant Physiol.* 129: 1368-1381.
- Neumann K, Droge-Laser W, Kohne S, Broer I. 1997. Heat treatment results in a loss of transgene-encoded activities in several tobacco lines. *Plant Physiol* 115: 939- 947.
- Nieto-Sotelo J, Kannan KB, Martinez LM, Segal C. 1999. Characterization of a maize heat-shock protein 101 gene, HSP101, encoding a ClpB/Hsp100 protein homologue. *Gene* 230: 187-195.
- Nikiforova V, Freitag J, Kempa S, Adamik M, Hesse H, Hoefgen R. 2003. Transcriptome analysis of sulfur depletion in *Arabidopsis thaliana*: interlacing of biosynthetic pathways provides response specificity. *Plant J.* 33: 633-650.
- Nishio N, Satoh H. 1997. A water-soluble chlorophyll protein in cauliflower may be identical to BnD22, a drought-induced 22 kilodalton protein in rapeseed. *Plant Physiol.* 115: 841-846.
- Noctor G, Foyer C. 1998. Ascorbate and glutathione: keeping active oxygen under control. *Annu. Rev. Plant Physiol. Mol. Biol.* 49: 249-279.
- Noctor G, Veljovic-Jovanovic S, Foyer CH. 2000. Peroxide processing in photosynthesis: antioxidant coupling and redox signalling. *Phil. Trans. R. Soc. Lond B* 355: 1465-1475.
- Noh YS, Amasino RM. 1999. Regulation of developmental senescence is conserved between *Arabidopsis* and *Brassica napus*. *Plant Mol. Biol.* 41: 195-206.
- Noodén LD. 1988. The phenomena of senescence and aging. *In*: LD Noodén, AC Leopold, eds, *Senescence and Aging in Plants*. Academic Press, Inc., San Diego, CA, pp 1-50.

- Oberschall A, Deák M, Török K, Sass L, Vass I, Kovács I, Fehér A, Dudits D, Horváth GV. 2000. A novel aldose/aldehyde reductase protects transgenic plants against lipid peroxidation under chemical and drought stresses. *Plant J.* 24: 437-446.
- Oh SJ, Song SI, Kim YS, Jang HJ, Kim SY, Kim M, Kim YK, Nahm BH, Kim JK. 2005. *Arabidopsis* CBF3/DREB1A and ABF3 in transgenic rice increased tolerance to abiotic stress without stunting growth. *Plant Physiol.* 138: 341-351.
- Ohtsubo S, Kobayashi H, Noro W, Taniguchi M, Saitoh E. 2005. Molecular cloning and characterization of oryzacystatin-III, a novel member of phytocystatin in rice (*Oryza sativa* L. japonica). *J. Agric. Food Chem.* 53: 5218-5224.
- Ojima A, Shiota H, Higashi K, Kamada H. 1997. An extracellular insoluble inhibitor of cysteine proteinases in cell cultures and seeds of carrot. *Plant Mol. Biol.* 34: 99-109.
- Okamoto T, Minamikawa T. 1998. A vacuolar cysteine endopeptidase (SH-EP) that digests seed storage globulin: characterization, regulation of gene expression and posttranslational processing. *J. Plant Physiol.* 152: 675-682.
- Okamoto T, Shimada T, Hara-Nishimura I, Nishimura M, Minamikawa T. 2003. C-terminal KDEL sequence of a KDEL-tailed cysteine proteinase (sulfhydryl-endopeptidase) is involved in formation of KDEL vesicle and in efficient vacuolar transport of sulfhydryl-endopeptidase. *Plant Physiol.* 132: 1892-1900.
- Olsen K, Daly J, Finnegan J, Holt H, Mahon R. 2001. The effect of two environmental factors, temperature and insect damage, on the efficacy of presquare Bt cotton plants. In: Gupta VVSR, ed. *Proceedings of 4th Pacific rim conference on the biotechnology of Bacillus thuringiensis and its environmental impact*, 11-15 November 2001, Australian National University, Australia, 37.

- Orr GL, Strickland JA, Walsh TA. 1994. Inhibition of *Diabrotica* larval growth by a multicystatin from potato tubers. *J. Insect Physiol.* 40: 893-900.
- Otegui MS, Noh YS, Martinez DE, Vila Petroff MG, Staehelin LA, Amasino RM, Guamet JJ. 2005. Senescence-associated vacuoles with intense proteolytic activity develop in leaves of *Arabidopsis* and soybean. *Plant J.* 41: 831-844.
- Outchkourov NS, de Kogel WJ, Wieggers GL, Abrahamson M, Jongsma MA. 2004a. Engineered multidomain cysteine protease inhibitors yield resistance against western flower thrips (*Frankliniella occidentalis*) in greenhouse trials. *Plant Biotech. J.* 2: 449-458.
- Outchkourov NS, de Kogel, WJ, Schuurman-de Bruin A, Abrahamson M, Jongsma MA. 2004b. Specific cysteine protease inhibitors act as deterrents of Western flower thrips *Frankliniella occidentalis* (Pergande) in transgenic potato. *Plant Biotech. J.* 2: 439-448.
- Outchkourov NS, Rogelj B, Strukelj B, Jongsma MA. 2003. Expression of sea anemone equistatin in potato: effects of plant proteases on heterologous protein production. *Plant Physiol.* 133: 379-390.
- Panchuk I, Volkov R, Schoffl F. 2002. Heat stress and HSF dependent expression of ascorbate peroxidase in *Arabidopsis*. *Plant Physiol.* 129: 838-853.
- Park S, Li J, Pittman JK, Berkowitz GA, Yang H, Undurraga S, Morris J, Hirschi KD, Gaxiola RA. 2005. Up-regulation of a H<sup>+</sup>-pyrophosphatase (H<sup>+</sup>-PPase) as a strategy to engineer drought-resistant crop plants. *Proc. Natl. Acad. Sci. USA* 102: 18830-18835.

- Pechan T, Chang YL, Mitra A, Lin L, Davis FM, Williams WP, Luthe DS. 2000. A unique 33-kd cysteine proteinase accumulates in response to larval feeding in maize genotypes resistant to fall armyworm and other lepidoptera. *Plant Cell* 12: 1031-1040.
- Pechan T, Cohen A, Williams WP, Luthe DS. 2002. Insect feeding mobilizes a unique plant defense protease that disrupts the peritrophic matrix of caterpillars. *Proc. Natl. Acad. Sci. USA* 99: 13319-13323.
- Pechan T, Jiang B, Steckler D, Ye L, Lin L, Luthe DS, Williams WP. 1999. Characterization of three distinct cDNA clones encoding cysteine proteinases from maize (*Zea mays* L.) callus. *Plant Mol. Biol.* 40: 111-119.
- Peferoen M. 1997. Progress and prospects for field use of Bt genes in crops. *Trends Biotech.* 15: 173-177.
- Pellegrineschi A, Reynolds M, Pacheco M, Brito RM, Almeraya R, Yamaguchi-Shinozaki K, Hoisington D. 2004. Stress-induced expression in wheat of the *Arabidopsis thaliana* DREB1A gene delays water stress symptoms under greenhouse conditions. *Genome* 47: 493-500.
- Pelleschi S, Rocher JP, Prioul JL. 1997. Effects of water restriction on carbohydrate metabolism and photosynthesis in mature maize leaves. *Plant Cell Environ.* 20: 493-503.
- Pernas M, Sánchez-Monge R, Salcedo G. 2000. Biotic and abiotic stress can induce cystatin expression in chestnut. *FEBS Lett.* 467: 206-210.
- Pic E, de la Serve B, Tardieu F, Turc O. 2002. Leaf senescence induced by mild water deficit follows the same sequence of macroscopic, biochemical and molecular events as monocarpic senescence in pea. *Plant Physiol.* 128: 136-246.

- Pinheiro C, Kehr J, Ricardo CP. 2005. Effect of water stress on lupin stem protein analysed by two-dimensional gel electrophoresis. *Planta* 221: 716-728.
- Pinheiro C, Chaves MM, Ricardo CP. 2001. Alterations in carbon and nitrogen metabolism induced by water deficit in the stems and leaves of *Lupinus albus* L. *J. Exp. Bot.* 52: 1063-1070.
- Pitcher LH, Zilinskas BA. 1996. Overexpression of copper/zinc superoxide dismutase in the cytosol of transgenic tobacco confers partial resistance to ozone-induced foliar necrosis. *Plant Physiol.* 110: 583-588.
- Pnueli L, Hongjian L, Mittler R. 2003. Growth suppression, altered stomatal responses, and augmented induction of heat shock proteins in cytosolic ascorbate peroxidase (Apx1)-deficient Arabidopsis plants. *Plant J.* 34: 187-203.
- Poorter H, Nagel O. 2000. The role of biomass allocation in the growth response of plants to different levels of light, CO<sub>2</sub>, nutrients and water: a quantitative review. *Aust. J. Plant Physiol.* 27: 595-607.
- Prändl R, Hinderhofer K, Eggers-Schumacher G, Schöffl F. 1998. HSF3, a new heat shock factor from *Arabidopsis thaliana*, derepresses the heat shock response and confers thermotolerance when overexpressed in transgenic plants. *Mol. Gen. Genet.* 258: 269-278.
- Prasad PV, Boote KJ, Allen LH, Jr., Thomas JMG. 2002. Effects of elevated temperature and carbon on seed-set yield of kidney bean (*Phaseolus vulgaris* L.). *Global Change Biol.* 8: 710-721.
- Quan R, Shang M, Zhang H, Zhao Y, Zhang J. 2004. Engineering of enhanced glycine betaine synthesis improves drought tolerance in maize. *Plant Biotech. J.* 2: 477-486.



- Quirino BF, Normanly J, Amasino RM. 1999. Diverse range of gene activity during *Arabidopsis thaliana* leaf senescence includes pathogen-independent induction of defense-related genes. *Plant Mol. Biol.* 40: 267-278.
- Rabbani MS, Maruyama K, Abe H, Khan MA, Katsura K, Ito Y, Yoshiwara K, Seki M, Shinozaki K, Yamaguchi-Shinozaki K. 2003. Monitoring expression profiles of rice genes under cold, drought, and high-salinity stresses and abscisic acid application using cDNA microarray and RNA gel-blot analyses. *Plant Physiol.* 133: 1755-1767.
- Rahbé Y, Deraison C, Bonadé-Bottino M, Girard C, Nardon C, Jouanin L. 2003. Effects of the cysteine protease inhibitor oryzacystatin (OC-I) on different aphids and reduced performance of *Myzus persicae* on OC-I expressing transgenic oilseed rape. *Plant Sci.* 164: 441-450.
- Rainwater DT, Gossett DR, Millhollon EP, Hanna HY, Banks SW, Lucas MC. 1996. The relationship between yield and the antioxidant defense system in tomatoes grown under heat stress. *Free Radic. Res.* 25: 421-435.
- Rakawal R, Agrawal KG, Kubo A, Yonekura M, Tamogami S, Saji H, Iwahashi H. 2002. Defense/stress responses elicited in rice seedlings exposed to the gaseous air pollutant sulfur dioxide. *Environ. Exper. Bot.* 00: 1-13.
- Ramanjulu S, Bartels D. 2002. Drought- and desiccation-induced modulation of gene expression in plants. *Plant Cell Environ.* 25: 141-151.
- Rawlings ND, Tolle DP, Barrett AJ. 2004. *MEROPS*: the peptidase database. *Nucl. Acids Res.* 32 (database issue): D160-D164.
- Reis EM, Margis R. 2001. Sugarcane phytoalexins: Identification, classification and expression pattern analysis. *Genetics and Mol. Biol.* 24: 291-296.

- Reymond P, Bodenhausen N, Van Poecke RMP, Krishnamurthy V, Dicke M, Farmer EE. 2004. A conserved transcript pattern in response to a specialist and a generalist herbivore. *Plant Cell* 16: 3132-3147.
- Reymond P, Weber H, Damond M, Farmer EE. 2000. Differential gene expression in response to mechanical wounding and insect feeding in *Arabidopsis*. *Plant Cell* 12: 707-719.
- Riccardi F, Gazeau P, Jacquemot MP, Vincent D, Zivy M. 2004. Deciphering genetic variations of proteome responses to water deficit in maize leaves. *Plant Physiol. Biochem.* 42: 1003-1011.
- Rizhsky L, Liang H, Mittler R. 2002. The combined effect of drought stress and heat shock on gene expression in tobacco. *Plant Physiol.* 130: 1143-1151.
- Rizhsky L, Liang H, Shuman J, Shulaev V, Davletova S, Mittler R. 2004. When defense pathways collide. The response of *Arabidopsis* to a combination of drought and heat stress. *Plant Physiol.* 134: 1683-1696.
- Rose TM, Henikoff JK, Henikoff S. 2003. CODEHOP (COnsensus-DEgenerate Hybrid Oligonucleotide Primer) PCR primer design. *Nucl. Acids Res.* 31: 3763-3766.
- Rotari VI, Dando PM, Barrett AJ. 2001. Legumain forms from plants and animals differ in their specificity. *Biol. Chem.* 382: 953-959.
- Roxas VP, Smith RK Jr, Allen ER, Allen RD. 1997. Overexpression of glutathione S-transferase/glutathione peroxidase enhances the growth of transgenic tobacco seedlings during stress. *Nature Biotech.* 15: 988-991.
- Rus AM, Estaño MT, Gisbert C, Garcia-Sogo B, Serrano R, Caro M, Moreno V, Bolarín MC. 2001. Expressing the yeast HAL1 gene in tomato increases fruit yield and enhances K<sup>+</sup>/Na<sup>+</sup> selectivity under salt stress. *Plant Cell Environ.* 24: 875-880.

- Ryan CA. 1990. Protease inhibitors in plants: genes for improving defenses against insects and pathogens. *Annu. Rev. Phytopathol.* 28: 425-449.
- Ryan CA. 2000. The systemin signaling pathway: differential activation of plant defensive genes. *Biochim. Biophys. Acta* 1477: 112-121.
- Saijo Y, Hata S, Kyojuka J, Shimamoto K, Izui K. 2000. Over-expression of a single Ca<sup>2+</sup>-dependent protein kinase confers both cold and salt/drought tolerance on rice plants. *Plant J.* 23: 319-327.
- Salekdeh GH, Siopongco J, Wade LJ, Ghareyazie B, Bennett J. 2002. A proteomic approach to analyzing drought- and salt-responsiveness in rice. *Field Crops Res.* 76: 199-219.
- Samac DA, Smigocki AC. 2003. Expression of oryzacystatin I and II in alfalfa increases resistance to the root-lesion nematode. *Phytopathology* 93: 799-804.
- Sambrook J, Fritsch EF, Maniatis T. 1989. *Molecular cloning: A laboratory manual*. Cold Spring Harbor, NY.
- Sambrook J, Russell DW. 2001. *Molecular cloning: A laboratory manual*. 3<sup>rd</sup> ed. Cold Spring Harbor, NY.
- Sangwan V, Orvar BL, Beyerly J, Hirt H, Dhindsa RS. 2002. Opposite changes in membrane fluidity mimic cold and heat stress activation of distinct plant MAP kinase pathways. *Plant J.* 31: 629-638.
- Savin R, Nicolas ME. 1996. Effects of short periods of drought and high temperature on grain growth and starch accumulation of two malting barley cultivars. *Aust. J. Plant Physiol.* 23: 201-210.
- Schaffer MA, Fischer RL. 1988. Analysis of mRNAs that accumulates in response to low temperature identifies a thiol protease gene in tomato. *Plant Physiol.* 87: 431-436.

- Schlereth A, Standhardt D, Mock HP, Muntz K. 2001. Stored cysteine proteinases start globulin mobilization in protein bodies of embryonic axes and cotyledons during vetch (*Vicia sativa* L.) seed germination. *Planta* 212: 718-727.
- Schmid M, Simpson D, Gietl C. 1999. Programmed cell death in castor bean endosperm is associated with the accumulation and release of a cysteine endopeptidase from ricinosomes. *Proc. Natl. Acad. Sci. USA* 96: 14159-14164.
- Schmid M, Simpson D, Kalousek F, Gietl C. 1998. A cysteine endopeptidase with a C-terminal KDEL motif isolated from castor bean endosperm is a marker enzyme for the ricinosome, a putative lytic compartment. *Planta* 206: 466-475.
- Schmid M, Simpson D, Sarioglu H, Lottspeich F, Gietl C. 2001. The ricinosomes of senescing plant tissue bud from the endoplasmic reticulum. *Proc. Natl. Acad. Sci. USA* 98: 5353-8535.
- Schouten A, Roosien J, Engelen FA, van Jong GAM, de Borst Vrenssen AWM, Zilverentant JF, Bosch D, Stiekema WJ, Gommers FJ, Schots A. 1996. The C-terminal KDEL sequence increases the expression level of a single-chain antibody design to be targeted to both the cytosol and the secretory pathway in transgenic tobacco. *Plant Mol. Biol.* 30: 781-793.
- Seki M, Narusaka M, Ishida J, Nanjo T, Fujita M, Oono Y, Kamiya A, Nakajima M, Enju A, Sakurai T, Satou M, Akiyama K, Taji T, Yamaguchi-Shinozaki K, Carninci P, Kawai J, Hayashizaki Y, Shinozaki K. 2002a. Monitoring the expression profiles of 7000 Arabidopsis genes under drought, cold and high-salinity stresses using a full-length cDNA microarray. *Plant J.* 31: 279-292.

- Seki M, Ishida J, Narusaka M, Fujita M, Nanjo T, Umezawa T, Akiyama K, Yamaguchi-Shinozaki K, Caeninici P, Kawai J, Hayashizaki Y, Shinozaki K. 2002b. Monitoring expression pattern of about 7000 Arabidopsis genes under ABA treatment using full length cDNA microarray. *Funct. Integr. Genomics* 2: 282-291.
- Seki M, Narusaka M, Abe H, Kasuga M, Yamaguchi-Shinozaki K, Carninci P, Hayashizaki Y, Shinozaki K. 2001. Monitoring the expression pattern of 1300 Arabidopsis genes under drought and cold stresses by using a full-length cDNA microarray. *Plant Cell* 13: 62-72.
- Selote DS, Bharti S, Khanna-Chopra R. 2004. Drought acclimation reduces  $O_2^-$  accumulation and lipid peroxidation in wheat seedlings. *Biochem. Biophys. Res. Commun.* 314: 724-729.
- Sharma HC, Sharma KK, Seetharama SN, Ortiz R. 2000. Prospects for using transgenic resistance to insects in crop improvement. *Electronic J. Biotech.* 3: 76-95.
- Sheveleva E, Chmara W, Bohnert HJ, Jensen RG. 1997. Increased salt and drought tolerance by D-ononitol production in transgenic *Nicotiana tabacum* L. *Plant Physiol.* 115: 1211-1219.
- Sheveleva EV, Marquez S, Chmara W, Zegeer A, Jensen RG, Bohnert HJ. 1998. Sorbitol-6-phosphate dehydrogenase expression in transgenic tobacco. *Plant Physiol.* 117: 831-839.
- Shi H, Lee BH, Wu SJ, Zhu JK. 2003. Overexpression of a plasma membrane  $Na^+/H^+$  antiporter gene improves salt tolerance in *Arabidopsis thaliana*. *Nature Biotech.* 21: 81-85.
- Shimada T, Hiraiwa N, Nishimura M, Hara-Nishimura I. 1994. Vacuolar processing enzyme of soybean that converts proproteins to the corresponding mature forms. *Plant Cell Physiol.* 35: 713-718.

- Shimada T, Yamada K, Kataoka M, Nakaune S, Koumoto Y, Kuroyanagi M, Tabata S, Kato T, Shinozaki K, Seki M, Kobayashi M, Kondo M, Nishimura M, Hara-Nishimura I. 2003. Vacuolar processing enzymes are essential for proper processing of seed storage proteins in *Arabidopsis thaliana*. *J. Biol. Chem.* 278: 32292-32299.
- Shinozaki K, Yamaguchi-Shinozaki K, Seki M. 2003. Regulatory network of gene expression in the drought and cold stress responses. *Curr. Opin. Plant Biol.* 6: 410-417.
- Shinozaki K, Yamaguchi-Shinozaki K. 1996. Molecular responses to drought and cold stress. *Curr. Opin. Biotechnol.* 7: 161-167.
- Shinozaki K, Yamaguchi-Shinozaki K. 1997. Gene expression and signal transduction in water-stress response. *Plant Physiol.* 115: 327-334.
- Shinozaki K, Yamaguchi-Shinozaki K. 1999. Molecular responses to drought stress. *In* Molecular Responses to Cold, Drought, Heat and Salt Stress in Higher Plants (Shinozaki K, Yamaguchi-Shinozaki K., eds). Austin, TX: RG Landes, pp. 11-28.
- Shinozaki K, Yamaguchi-Shinozaki K. 2000. Molecular responses to dehydration and low temperature: differences and cross-talk between two stress signaling pathways. *Curr. Opin. Plant Biol.* 3: 217-223.
- Shou H, Bordallo P, Wang K. 2004. Expression of the Nicotiana protein kinase (NPK1) enhanced drought tolerance in transgenic maize. *J. Exp. Bot.* 55: 1013-1019.
- Shyu DJ, Chou WM, Yiu TJ, Lin CP, Tzen JT. 2004. Cloning, functional expression, and characterization of cystatin in sesame seed. *J. Agric. Food Chem.* 52: 1350-1356.
- Sin SF, Chye ML. 2004. Expression of proteinase inhibitor II proteins during floral development in *Solanum americanum*. *Planta* 219: 1010-1022.

- Siqueira-Junior CL, Fernandes KVS, Machado OLT, Cunha M, Gomes VM, Moura D, Jacinto T. 2002. 87-kDa tomato cystatin exhibits properties of a defence protein and forms crystals in prosystemin over-expressing transgenic plants. *Plant Physiol. Biochem.* 40: 247-254.
- Sivamani E, Bahieldin A, Wraithc JM, Al-Niemia T, Dyera WE, Hod THD, Qu R. 2000. Improved biomass productivity and water use efficiency under water deficit conditions in transgenic wheat constitutively expressing the barley HVA1 gene. *Plant Sci.* 155: 1-9.
- Smart CM, Hosken SE, Thomas H, Greaves JA, Blair BG, Schuch W. 1995. The timing of maize leaf senescence and characterization of senescence-related cDNAs. *Physiol Plant.* 93: 673-682.
- Smirnoff N. 1993. The role of active oxygen in the response of plants to water deficit and desiccation. *New Phytol.* 125: 27-58.
- Solomon KF, Labuschagne MT. 2003. Variation in water use and transpiration efficiency among durum wheat genotypes grown under moisture stress and non-stress conditions. *J. Agri. Sci.* 141: 31-41.
- Solomon M, Belenghi B, Delledonne M, Menachem E, Levine A. 1999. The involvement of cysteine proteases and protease inhibitor genes in the regulation of programmed cell death in plants. *Plant Cell* 11: 431-444.
- Sotiropoulou G, Anisowicz A, Sager R. 1997. Identification, cloning, and characterization of cystatin M, a novel cysteine proteinase inhibitor, down-regulated in breast cancer. *J. Biol. Chem.* 272: 903-910.
- Sousa-Majer MJ, Turner NC, Hardie DC, Morton RL, Lamont B, Higgins TJV. 2004. Response to water deficit and high temperature of transgenic peas (*Pisum sativum* L.)

- containing a seed specific  $\alpha$ -amylase inhibitor and the subsequent effects on pea weevil (*Bruchus pisorum* L.) survival. *J. Exp. Bot.* 55: 497-505.
- Steponkus PL, Uemura M, Joseph RA, Gilmour SJ, Thomashow MF. 1998. Mode of action of the COR15a gene on the freezing tolerance of *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* 95: 14570-14575.
- Stevens LH, Stoopern GM, Elbers IJW, Molthoff JW, Bakker HAC, Lommen A, Bosch D, Jordi W. 2000. Effect of climate conditions and plant developmental stage on the stability of antibodies expressed in transgenic tobacco. *Plant Physiol.* 124: 173-182.
- Stoll M, Loveys B, Dry P. 2000. Hormonal changes induced by partial root zone drying of irrigated grapevine. *J. Exp. Bot.* 51: 1627-1634.
- Stroehrer VL, Maclanag JL, Good AG. 1997. Molecular cloning of a *Brassica napus* cysteine protease gene inducible by drought and low temperature stress. *Physiol. Plant* 101: 389-397.
- Sugino M, Hibino T, Tanaka Y, Nii N, Takabe T. 1999. Overexpression of DnaK from a halotolerant cyanobacterium *Aphanothece halophytice* acquires resistance to salt stress in transgenic tobacco plants. *Plant Sci.* 146: 81-88.
- Sun W, Bernard C, van de Cotte B, Van Montagu M, Verbruggen N. 2001. At-HSP17.6A, encoding a small heat-shock protein in *Arabidopsis*, can enhance osmotolerance upon overexpression. *Plant J.* 27: 407-415.
- Suzuki N, Rizhsky L, Liang H, Shuman J, Shulaev V, Mittler R. 2005. Enhanced tolerance to environmental stress in transgenic plants expressing the transcriptional coactivator multiprotein bridging factor. *Plant Physiol.* 139: 1313-1322.



- Swidzinski JA, Sweetlove LJ, Leaver CJ. 2002. A custom microarray analysis of gene expression during programmed cell death in *Arabidopsis thaliana*. *Plant J.* 30: 431-446.
- Takahashi S, Seki M, Ishida J, Satou M, Sakurai T, Narusaka M, Kamiya A, Nakajima M, Enju A, Akiyama K, Yamaguchi-Shinozaki K, Shinozaki K. 2004. Monitoring the expression profiles of genes induced by hyperosmotic, high salinity, and oxidative stress and abscisic acid treatment in *Arabidopsis* cell culture using a full-length cDNA microarray. *Plant Mol. Biol.* 56:29-55.
- Tanaka T, Yamauchi D, Minamikawa T. 1991. Nucleotide sequence for an endopeptidase (EP-C1) from pods of maturing *Phaseolus vulgaris* fruits. *Plant Mol. Biol.* 16: 1083-1084.
- Tardieu F, Katerji N, Bethenod J, Zhang J, Davis WJ. 1991. Maize stomatal conductance in the field: its relationship with soil and plant water potentials, mechanical constraints and ABA concentration in the xylem sap. *Plant Cell Environ.* 14: 121-126.
- Tewari AK, Tripathy BC. 1998. Temperature-stress-induced impairment of chlorophyll biosynthetic reactions in cucumber and wheat. *Plant Physiol.* 117: 851-858.
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG. 1997. The ClustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucl. Acids Res.* 25: 4876-4882.
- Tiedemann J, Schlereth A, Muntz K. 2001. Differential tissue-specific expression of cysteine proteinases forms the basis for the fine-tuned mobilization of storage globulin during and after germination in legume seeds. *Planta* 212: 728-238.
- Toyooka K, Okamoto T, Minamikawa T. 2000. Mass transport of a proform of a KDEL-tailed cysteine proteinase (SH-EP) to protein storage vacuoles by endoplasmic reticulum-

- derived vesicle is involved in protein mobilization in germinating seeds. *J. Cell Biol.* 148: 453-563.
- Traore SB, Carlson RE, Pilcher CD, Rice ME. 2000. Bt and non-Bt maize growth and development as affected by temperature and drought stress. *Agron. J.* 92: 1027-1035.
- Turk V, Bode W. 1991. The cystatins: protein inhibitors of cysteine proteinases. *FEBS Lett.* 285: 213-219.
- Turner NC, Wright GC, Siddique KHM. 2001. Adaptation of grain legumes (pulses) to water-limited environments. *Advan. Agron.* 71: 193-231.
- Ueda T, Seo S, Ohashi Y, Hashimoto J. 2000. Circadian and senescence-enhanced expression of a tobacco cysteine protease gene. *Plant Mol. Biol.* 44: 649-657.
- Umezawa T, Yoshida R, Maruyama K, Yamaguchi-Shinozaki K, Shinozaki K. 2004. SRK2C, a SNF1-related protein kinase 2, improves drought tolerance by controlling stress-responsive gene expression in *Arabidopsis thaliana*. *Proc. Natl Acad. Sci. USA* 101: 17306-17311.
- Uno Y, Furihata T, Abe H, Yoshida R, Shinozaki K, Yamaguchi-Shinozaki K. 2000. Arabidopsis basic leucine zipper transcription factors involved in an abscisic acid-dependent signal transduction pathway under drought and high-salinity conditions. *Proc. Natl. Acad. Sci. USA* 97: 11632-11637.
- Urwin PE, Atkinson HJ, Waller DA, McPherson MJ. 1995. Engineered oryzacystatin-I expressed in transgenic hairy roots confers resistance to *Globodera-pallida*. *Plant J.* 8: 121-131.

- Urwin PE, Green J, Atkinson HJ. 2003. Expression of a plant cystatin confers partial resistance to *Globozadara*, full resistance is achieved by pyramiding a cystatin with natural resistance. *Mol. Breed.* 12: 263-269.
- Urwin PE, Levesley A, McPherson MJ, Atkinson HJ. 2000. Transgenic resistance to the nematode *Rotylenchulus reniformis* conferred by *Arabidopsis thaliana* plants expressing proteinase inhibitors. *Mol. Breed.* 6: 257-264.
- Urwin PE, Lilley CJ, McPherson MJ, Atkinson HJ. 1997. Resistance to both cyst and root-knot nematodes conferred by transgenic *Arabidopsis* expressing a modified plant cystatin. *Plant J.* 12: 455-461.
- Vacca RA, de Pinto MC, Valenti D, Passarella S, Marra E, DeGara L. 2004. Production of reactive oxygen species, alteration of cytosolic ascorbate peroxidase, and impairment of mitochondrial metabolism are early events in heatshock-induced programmed cell death in tobacco Bright-Yellow 2 cells. *Plant Physiol.* 134: 1100-1112.
- Vain P, Worland B, Clarke MC, Richard G, Beavis M, Liu H, Kohli A, Leech M, Snape J, Christou P. 1998. Expression of an engineered cysteine proteinase inhibitor (Oryzacystatin-I delta D86) for nematode resistance in transgenic rice plants. *Theor. Appl. Genet.* 96: 266-271.
- Valpuesta V, Lange NE, Guerrero C, Reid MS. 1995. Up-regulation of a cysteine protease accompanies the ethylene-insensitive senescence of daylily (*Hemerocallis* spp.) flowers. *Plant Mol. Biol.* 28: 575-582.
- Van Camp W, Capiou K, Van Montagu M, Inze D, Sloaten L. 1996. Enhancement of oxidative stress tolerance in transgenic tobacco plants overproducing Fe-superoxide dismutase in chloroplasts. *Plant Physiol.* 112: 1703-1714.

- Van der Vyver C, Schneidereit J, Driscoll S, Turner J, Kunert K, Foyer CH. 2003. Oryzacystatin-I expression in transformed tobacco produces a conditional growth phenotype and enhances chilling tolerance. *Plant Biotech. J.* 1: 101-112.
- Van Doorn WG, Woltering EJ. 2005. Many ways to exit? Cell death categories in plants. *Trends Plant Sci.* 10: 117-122.
- Vierling E. 1991. The roles of heat shock proteins in plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 42: 579-620.
- Vierstra RD. 1993. Protein degradation in plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 44: 385-410.
- Vierstra RD. 1996. Proteolysis in plants: mechanisms and functions. *Plant Mol. Biol.* 32: 275-302.
- Visal S, Taylor MAJ, Michaud D. 1998. The proregion of papaya proteinase IV inhibits Colorado potato beetle digestive cysteine proteinases. *FEBS Lett.* 434: 401-405.
- Vogel JT, Zarka DG, Van Buskirk HA, Fowler SG, Thomashow MF. 2005. Roles of the CBF2 and ZAT12 transcription factors in configuring the low temperature transcriptome of *Arabidopsis*. *Plant J.* 41: 195-211.
- Wagstaff C, Leverentz MK, Griffiths G, Thomas B, Chanasut U, Stead AD, Rogers HJ. 2002. Cysteine proteinase gene expression and proteolytic activity during senescence of *Alstroemeria* petals. *J. Exp. Bot.* 53: 233-240.
- Waldron C, Wegrich LM, Merlo PAO, Walsh TA. 1993. Characterization of a genomic sequence coding for potato multicystatin, an eight-domain cysteine proteinase inhibitor. *Plant Mol. Biol.* 23: 801-812.

- Walker AJ, Urwin PE, Atkinson HJ, Brain P, Glen DM, Shewry PR. 1999. Transgenic *Arabidopsis* leaf tissue expressing a modified oryzacystatin shows resistance to the field slug *Deroceras reticulatum* (Müller). *Transgenic Res.* 8: 95-103.
- Walsh TA, Strickland JA. 1993. Proteolysis of the 85-kilodalton crystalline cysteine proteinase inhibitor from potato releases functional cystatin domains. *Plant Physiol.* 103: 1227-1234.
- Walter C, Broer I, Hillemann D, Pühler A. 1992. High frequency, heat treatment-induced inactivation of the phosphothricin resistance gene in transgenic single cell suspension cultures of *Medicago sativa*. *Mol. Gen. Genet.* 235: 189-196.
- Wan L, Xia Q, Qiu X, Selvaraj G. 2002. Early stages of seed development in *Brassica napus*: a seed coat-specific cysteine proteinase associated with programmed cell death of the inner integument. *Plant J.* 30: 1-10.
- Wang HY, Huang YC, Chen SF, Yeh KW. 2003. Molecular cloning, characterization and gene expression of a water deficiency and chilling induced proteinase inhibitor I gene family from sweet potato (*Ipomoea batatas* Lam.) leaves. *Plant Sci.* 165: 191-203.
- Wang R, Guegler K, LaBrie ST, Crawford NM. 2000. Genomic analysis of a nutrient response in *Arabidopsis* reveals diverse expression patterns and novel metabolic and potential regulatory genes that are induced by nitrate. *Plant Cell* 12: 1491-1510.
- Wang W, Barak T, Vinocur B, Shoseyov O, Altman A. 2003. Abiotic resistance and chaperones: possible physiological role of SP1, a stable and stabilizing protein from *Populus*. In: Vasil IK (ed) *Plant biotechnology 2000 and beyond*. Kluwer, Dordrecht, pp 439-443.

- Wang W, Vinocur B, Altman A. 2003. Plant responses to drought, salinity and extreme temperatures: towards genetic engineering for stress tolerance. *Planta* 218: 1-14.
- Wawrzynska A, Lewandowska M, Hawkesford MJ, Sirko A. 2005. Using a suppression subtractive library-based approach to identify tobacco genes regulated in response to short-term sulphur deficit. *J. Exp. Bot.* 56: 1575-1590.
- Weaver LM, Gan S, Quirino BF, Amasino RM. 1998. A comparison of the expression patterns of several senescence-associated genes in response to stress and hormone treatment. *Plant Mol. Biol.* 37: 455-469.
- Williams J, Bulman M, Huttly A, Phillips A, Neill S. 1994. Characterization of a cDNA *Arabidopsis thaliana* encoding a potential thiol protease whose expression is induced independently by wilting and abscisic acid. *Plant Mol. Biol.* 25: 259-270.
- Wiśniewski K, Zagdańska B. 2001. Genotype-dependent proteolytic response of spring wheat to water deficiency. *J. Exp. Bot.* 52: 691-713.
- Wolfson JL, Murdock LL. 1990. Diversity in digestive proteinase activity among insects. *J. Chem. Eco.* 16: 1089-1102.
- Wong EY, Hironaka CM, Fischhoff DA. 1992. *Arabidopsis thaliana* small subunit leader and transit peptide enhance the expression of *Bacillus thuringiensis* proteins in transgenic plants. *Plant Mol. Biol.* 20: 81-93.
- Wu JW, Haard NF. 2000. Purification and characterization of a cystatin from the leaves of methyl jasmonate treated tomato plants. *Comp. Biochem. Physiol.* 3: 209-220.
- Xu FX, Chye ML. 1999. Expression of cysteine proteinase during developmental events associated with programmed cell death in brinjal. *Plant J.* 17: 321-327.

- Xu Q, Huang B. 2001. Morphological and physiological characteristics associated with heat tolerance in creeping bentgrass. *Crop Sci.* 41:127–133.
- Yamada K, Shimada T, Nishimura M, Hara-Nishimura I. 2005. A VPE family supporting various vacuolar functions in plants. *Physiol. Plant.* 123: 369-375.
- Yamaguchi-Shinozaki K, Kasuga M, Liu Q, Nakashima K, Sakuma Y, Abe H, Shinwari ZK, Seki M, Shinozaki K. 2002. Biological mechanisms of drought stress response. JIRCAS Working Report, pp. 1-8.
- Yamaguchi-Shinozaki K, Shinozaki K. 2005. Organization of cis-acting regulatory elements in osmotic- and cold-stress responsive promoters. *Trends Plant Sci.* 10: 88-94.
- Yamaguchi-Shinozaki K, Shinozaki K. 1994. A novel cis-acting element in an Arabidopsis gene is involved in responsiveness to drought, low temperature, or high-salt stress. *Plant Cell* 6: 251-264.
- Yamanouchi U, Yano M, Lin H, Ashikari M, Yamada K. 2002. A rice spotted leaf gene, Spl7, encodes a heat stress transcription factor protein. *Proc. Natl. Acad. Sci. USA* 99: 7530-7535.
- Yan J, He C, Wang J, Mao Z, Holaday SA, Allen RD and Zhang H. 2004. Overexpression of the *Arabidopsis* 14-3-3 protein GF14 $\lambda$  in cotton leads to a “Stay-Green” phenotype and improves stress tolerance under moderate drought conditions. *Plant Cell Physiol.* 45: 1007-1014.
- Younis ME, El-Shahaby OA, Abo-Hamed SA, Ibrahim AH. 2000. Effects of water stress on growth, pigments and <sup>14</sup>CO<sub>2</sub> assimilation in three sorghum cultivars. *J. Agron. Crop Sci.* 185: 73-82.

- Zagdańska B, Wisniewski K. 1996. Endoproteinase activities in wheat leaves upon water deficit. *Acta Biochim. Polo.* 43: 515-519.
- Zhang HX, Blumwald E. 2001. Transgenic salt-tolerant tomato plants accumulate salt in foliage but not in fruit. *Nature Biotech.* 19: 765-768.
- Zhang HX, Hodson JN, Williams JP, Blumwald E. 2001. Engineering salt-tolerant *Brassica plants*: characterization of yield and seed oil quality in transgenic plants with increased vacuolar sodium accumulation. *Proc. Natl Acad. Sci. USA* 98: 12832-12836.
- Zhang J, Klueva NY, Zhang Z, Wu R, Ho T-H D, Nguyen HT. 2000. Genetic engineering for abiotic stress resistance in crop plants. *InVitro Cellular Develop. Biol -Plant* 36: 108-114.
- Zhang N, Jones BL. 1995. Characterization of germinated barely endoproteolytic enzymes by two dimensional gel electrophoresis. *J. Cereal Sci.* 21: 145-153.
- Zhang Y, Rouf Mian MAR, Chekhovskiy K, So S, Kupfer D, Lai H, Roe BA. 2005. Differential gene expression in *Festuca* under heat stress conditions. *J. Exp. Bot.* 56: 897-907.
- Zhao Y, Botella MA, Subramanian L, Niu X, Nielsen SS, Bressan RA, Hasegawa PM. 1996. Two wound inducible soybean cysteine proteinase inhibitors have greater insect digestive proteinase inhibitory activities than a constitutive homolog. *Plant Physiol.* 111: 1299-1306.
- Zhu JK. 2002. Salt and drought stress signal transduction in plants. *Annu. Rev. Plant Biol.* 53: 247-273.
- Zinselmeier C, Sun Y, Helentjaris T, Beatty M, Yang S, Smith H, Habben J. 2002. The use of gene expression profiling to dissect the stress sensitivity of reproductive development in maize. *Field Crops Res.* 75: 111-121.