

Literature cited

Alexander, S.A.; Horner, W.E. and Lewis, K.J. (1988). *Leptographium procerum* as a pathogen of pines. In: *Leptographium* root diseases on conifers. (ed. T.C. Harrington & F.W. Cobb Jr.), American Phytopathological Society, St. Paul, Minnesota, pp. 97-112.

Anderson, R.L. and Alexander, S.A. (1979). How to identify and control white pine root decline. *Forestry Bulletin SA-FR/P6*.

Aoshima, K. (1965). Studies of wood-staining fungi of Japan, Doctoral thesis, University of Tokyo. (In Japanese with English summary), pp. 285.

Arx, J.A., von. (1952). Über die Ascomycetengattungen *Ceratostomella* Sacc., *Ophiostoma* Syd. und *Rostrella* Zimmermann. *Antonie van Leeuwenhoek* **18**, 201-213.

Ayer, W.A., Browne, L.M. and Lovell, S.H. (1983). Biologically active phenolic metabolites of a *Verticicladiella* species. *Phytochemistry* **22**, 2267-2271.

Ayer, W.A., Browne, L.M. and Lin, G. (1989). Metabolites of *Leptographium wagneri*, the causative agent of black stain root disease of conifers. *Journal of Natural Products* **52**, 119-129.

Bakshi, B. K. (1950). Fungi associated with ambrosia beetles in Great Britain. *Transactions of the British Mycological Society* **33**, 111-120.

Bakshi, B.K. (1951). Studies on four species of *Ceratocystis*, with a discussion on fungi causing sapstain in Britain. *Mycological paper* **35**, 1-16.

Barnard, E.L., Blakeslee, G.M., English, J.T., Oak, S.W. and Anderson, R.L. (1985). Pathogenic fungi associated with sand pine root disease in Florida. *Plant Disease* **69**, 196-199.

Barras, S.T. (1973). Reduction of progeny and development in the Southern pine

beetle following removal of symbiotic fungi. *Canadian Entomologist* **105**, 1295-1299.

Barras, S.J. and Perry, T. (1971a). *Leptographium terebrantis* sp. nov. associated with *Dendroctonus terebrantis* in loblolly pine. *Mycopatologia et Mycologia applicata* **43**, 1-10.

Barras, S.J. and Perry, T. (1971b). Gland cells and fungi associated with prothoacic mycangium of *Dendroctonus adjunctus* (Coleoptera: Scolytidae). *Annals of the Entomological Society of America* **64**, 123-126.

Batra, L.R. (1963). Ecology of ambrosia fungi and their dissemination by beetles. *Transactions of the Kansas Academy of Science* **66**, 213-236.

Bedard, W. and Ferrell, G. (1990). Trapping evaluation of beetle vectors of black-stain root disease in Douglas fir. *Canadian Entomologist* **122**, 459-468.

Bennet, E.M. and Tattar, T.A. (1988). Bluestain fungi and insect vector interaction in Japanese black and scots pine mortality. *Arboricultural Journal* **12**, 237-247.

Berryman, A.A. (1972). Resistance of conifers to invasion by bark beetle-fungus associations. *BioScience* **22**, 598-602.

Bertagnole, C.L., Woo, J.Y. and Partridge, A.D. (1983). Pathogenicity of five *Verticicladiella* species to lodgepole pine. *Canadian Journal of Botany*, **61**, 1861-1867.

Bertagnole, C.L., Partridge, A.D. and LeTourneau, D.J. (1987). Interacting root pathogens associated with black stain root disease of *Pinus ponderosa* Laws. *Phytopathology* **77**, 1237.

Bevan, B. (1962). Pine shoot beetle. Forestry Commission Leaflet 3. HMSO. London. Pp.8

Blackwell, M., Bridges, J.R., Moser, J.C. and Perry, T.J. (1986). Hyperphoretic dispersal of *Pyxidiophora* anamorph. *Science* **232**, 993-995.

Bramble, W.C. and Holst, E.C. (1935). Microorganisms infecting pine attacked by *Dendroctonus frontalis*. *Phytopathology* **25**, 7.

Bramble, W.C. and Holst, E.C. (1940) Fungi associated with *Dendroctonus frontalis* in killing shortleaf pines and their effect on conduction. *Phytopathology* **30**, 881-899.

Brand, J.M., Bracke, L.N., Makovetz, A.J. and Barras, S.J. (1976). Bark beetle pheromones: production of verbenone by a mycangial fungus of *Dendroctonus frontalis*. *Journal of Chemical Ecology* **2**, 195-199.

Britz, H. (1997). *Fusarium subglutinans* f.sp. *pini*: a taxonomic, molecular and population study. MSc thesis, University of the Orange Free State. pp. 95.

Butin, H. and Aquilar, A.M. (1984). Blue-stain fungi on *Nothofagus* from Chile - including two new species of *Ceratocystis* Ellis and Halst. *Phytopathologische Zeitschrift* **109**, 80-89.

Byler, J.W., Cobb, F.W. (jr) and Rowney, D.L. (1979). An evaluation of black stain root disease on the Georgetown divide Eldorado County. *Forest Insect and Disease Management Report* **79**, 1-15.

Byler, J.W., Harrington, T.C., James, R.L. and Haglund, S. (1983). Black stain root disease in Douglas-Fir in Western Montana. *Plant Disease* **67**, 1037 - 1038.

Cobb, F.W. (jr.) (1988). *Leptographium wageneri*, cause of black-stain root disease: a review of its discovery, occurrence and biology with emphasis on pinyon and ponderosa pine. In: *Leptographium* root diseases on conifers. American Phytopathological Society, St Paul, Minnesota, pp. 41-62.

Cobb, F.W. and Platt, W.D. (1967). Pathogenicity of *Verticicladiella wageneri* to Douglas-fir. *Phytopathology* **57**, 998-999.

Cobb, F.W. (Jr.), Parmeter, J.R. (jr.), Wood, D.L. and Stark, R.W. (1974). Root pathogens as agents predisposing ponderosa pine and white fir to bark beetles. In: Proceedings of the fourth international conference on *Fomes annosus* (ed. E.G.

Kuhlman), pp. 8-15.

Cobb, F.W. (Jr.), Slaughter, G.W., Rowney, D.L. and DeMars, C.J. (1982). Rate of spread of *Ceratocystis wageneri* to ponderosa pine stands in Central Sierra Nevada. *Phytopathology*, **72**, 1359-1362.

Cobb, F.W. jr., Goheen, D.J. and Harrington, T.C. (1984). Black-stain root disease caused by *Verticicladiella wageneri*. (Abstract). Fourth International Congress for Plant Pathology, Abstract 333, p84.

Cobb, F.W. (Jr.), Lawson, T.T and Popenuck, T.L. (1987). Interactions among the three variants of *Verticicladiella wageneri* and the three host types. *Phytopathology* **77**, 1717-1718.

Coetsee, C. (1999). Characterisation of selected Ophiostomatoid fungi. M.Sc. Thesis, Department of Microbiology and Biochemistry, University of the Orange Free State, Bloemfontein, South Africa. pp. 104.

Correll, J.C., Klittich, C.J.R. and Leslie, J.F. (1987). Nitrate nonutilising mutants of *Fusarium oxysporum* and their use in vegetative compatibility tests. *Phytopathology* **77**, 1640-1646.

Correll, J.C., Klittich, C.J.R. and Leslie, J.F. (1989). Heterokaryon self-incompatibility in *Gibberella fujikuroi* (*Fusarium moniliforme*). *Mycological Research* **93**, 21-27.

Craighead, F.C. (1928). Interrelation of tree-killing bark beetles (*Dendroctonus*) and blue-stain. *Journal of Forestry* **26**, 886-887.

Crane, J.L. (1971). Illinois fungi II. A new species of *Phialocephala*. *Transactions of the British Mycological Society* **56**, 160-163.

Davidson, R.W. (1935). Fungi causing stain in logs and lumber in the southern states, including five new species. *Journal of Agricultural Research* **50**, 789-806.

Davidson, R.W. (1942). Some additional species of *Ceratostomella* in the United

States. *Mycologia* **34**, 650-662.

Davidson, R.W. (1955). Wood-staining fungi associated with bark beetles in Engelmann spruce in Colorado. *Mycologia* **47**, 58-67.

Davidson, R.W. (1958). Additional species of Ophiostomataceae from Colorado. *Mycologia* **50**, 661-670.

Davidson, R.W. (1966). New species of *Ceratocystis* from conifers. *Mycopathologia et Mycologia applicata* **28**, 273-286.

Davidson, R.W. (1971). New species of *Ceratocystis*. *Mycologia* **63**, 5-15.

Davidson, R.W. (1976). Sapwood staining fungi from two tree species. *Memoirs of the New York Botanical Garden* **28**, 45-49.

Davidson, R.W. (1978). Staining fungi associated with *Dendroctonus adjunctus* in pines. *Mycologia* **70**, 35-40.

Davidson, R.W., Francke-Grosmann, H. and Käärik, A. (1967). A restudy of *Ceratocystis penicillata* and report of two American species of this genus from Europe. *Mycologia* **59**, 928-931.

Davidson, R.W. and Robinson-Jeffrey, R.C. (1965). New records of *Ceratocystis europhioides* and *C. huntii* with *Verticicladiella* imperfect stages from conifers. *Mycologia* **57**, 488-490.

De Hoog, G.S. (1974). The genera *Blastobotrys*, *Sporothrix*, *Calcarisporium* and *Calcarisporiella* gen. nov. *Studies in Mycology*, 1-84.

De Hoog, G.S. (1993). *Sporothrix*-like anamorphs of *Ophiostoma* species and other fungi. In: *Ceratocystis and Ophiostoma: Taxonomy, Ecology and Pathogenicity* (eds. M.J. Wingfield, K.A. Seifert and J.F. Webber), American Phytopathological Society, St. Paul, Minnesota, pp. 53-60.

- De Hoog, G.S. and Scheffer, R.J. (1984). *Ceratocystis* vs. *Ophiostoma*: a reappraisal. *Mycologia* **76**, 292-299.
- Diamandis, S., Epstein, L. and Cobb, F.W. jr. (1987). Germination of conidia of *Verticicladiella wagneri* on root surface. *Phytopathology* **77**, 1238.
- Dochinger, L.S. (1967). *Leptographium* root decline of eastern white pine. (Abstract). *Phytopathology* **57**, 809.
- Farrell, R.L., Hadar, E., Kay, S.J., Blanchette, R.A. and Harrington, T.C. (1997). Survey of sapstain organisms in New Zealand and albino anti-sapstain fungi. Conference proceedings on the Biology and Prevention of Sapstain. Canada.
- Fenn, M.E., Dunn, P.H. and Wilborn, R. (1990). Black-stain root disease in ozone stressed ponderosa pine. *Plant Disease* **74**, 426-430.
- Fergus, C.L. (1956). The influence of actidione on wood-staining fungi. *Mycologia* **48**, 468-472.
- Francke-Grosmann, H. (1965). Über symbiosen von xylo-mycetophagen und ploeoophagen Scolytoidea mit holzbewohnenden pilzen. Supplement to Materials and Organisms. International symposium Berlin-Dahlem, 1965. 503-522.
- Francke-Grosmann, H. (1967). Ectosymbiosis in Wood-Inhabiting Insects. In: Symbiosis. Volume 2. (ed. S.M. Henry). Accademic Press, New York and London, pp 141-205.
- French, D.W. (1949). Pole blight- This is how to recognized it. Pest Leaflet, Division of Forest Pathology, Forest Service. Pp4.
- Furniss, M.M., Solheim, H. and Christiansen, E. (1990). Transmission of blue-stain fungi by *Ips typographus* (Coleoptera: Scolytidae) in Norway spruce. *Annals of the Entomological Society of America* **83**, 712-716.
- Foster, R.E. (1957). Pole blight of Western white pine. Timber of Canada,

contribution no 275. 4pp.

Gams, W. (1971). *Cephalosporium* -artig Schimmelpilze (Hyphomycetes). G. Fisher: Stuttgart.

Gambogi, P. and Lorenzini, G. (1977). Conidiophore morphology in *Verticicladiella serpens*. *Transactions of the British Mycological Society* **69**, 217-223.

Gibbs, J.N. (1993). The biology of ophiostomatoid fungi causing sapstain in trees and freshly cut logs. In: *Ceratocystis and Ophiostoma: Taxonomy, Ecology and Pathogenicity*. (ed. M.J. Wingfield, K.A. Seifert and J.F. Webber), pp. 153-160. American Phytopathological Society Press: St Paul, Minnesota, pp. 153-160.

Gibbs, J.N. and Inman, A. (1991). The pine shoot beetle *Tomicus piniperda* as a vector of blue-stain fungi to windblown pine. *Forestry* **64**, 239-249.

Gill, L.S. and Andrews, S.R. (1949). Note on a *Scopularia* attacking western white pine. *Plant Disease Reporter* **33**, 227.

Gill, L.S., Leaphart, C.D. and Andrews, S.R. (1951). Preliminary results of inoculations with a species of *Leptographium* on Western white pine. *Forest Pathology Special Release* **35**, 1-14.

Goheen, D.J. (1976). *Verticicladiella wageneri* on *Pinus ponderosa*: Epidemiology and interrelationships with insects. Ph. D. thesis. University of California, Berkeley. 118 pp.

Goheen, D.J. and Cobb, F.W.(Jr.) (1978). Occurrence of *Verticicladiella wageneri* and its perfect state, *Ceratocystis wageneri* sp. nov., in insect galleries. *Phytopathology* **68**, 1192-1195.

Goheen, D.J. and Cobb, F.W. (Jr.) (1980). Infestation of *Ceratocystis wageneri* -infected ponderosa pines by bark beetles (Coleoptera:Scolytidae) in the Central Sierra Nevada. *The Canadian Entomologist* **112**, 725-730.

Goheen, D.J., Cobb, F.W. (Jr.), Wood, D.L. and Rowney, D.L. (1985). Visitation frequencies of some insect species on *Ceratocystis wageneri* infected and apparently healthy ponderosa pines. *The Canadian Entomologist* **117**, 1535-1543.

Goheen, D.J., Cobb, F.W. (Jr.) and McKibbin, G.N. (1978). Influence of soil moisture on infection of Ponderosa pine by *Verticicladiella wageneri*. *Phytopathology* **68**, 913-916.

Goheen, D.J. and Hansen, E.M. (1978). Black stain root disease in Oregon and Washington. *Plant Disease Reporter* **62**, 1098-1102.

Goidanich, G. (1936). Il genera di Ascomiceti "Grosman" G. Goid. *Bollettino della Roma Stazione di Patologia Vegetale-Roma N.S.* **16**, 26-60.

Griffin, H.D. (1968). The genus *Ceratocystis* in Ontario. *Canadian Journal of Botany* **46**, 689-718.

Grosman, H. (1931). Beiträge zur Kenntnis der Lebensgemeinschaft zwischen Borkenkäfern und Pilzen. (Contributions to the knowledge concerning the life partnership between bark beetles and fungi) *Zeitschrift für Parasitenkunde* **3**, 56-102.

Grosman, H. (1932). Über die systematischen Beziehungen der Gattung *Leptographium* Lagerberg and Melin zur Gattung *Ceratostomella* Sacc. *Hedwigia* **72**, 183-193.

Halambek, M. (1976). Dieback of eastern white pine (*Pinus strobus* L.) in cultures. *Agriculturae Conspectus Scientificus* **39**, 495-498.

Halambek, M. (1981). *Verticicladiella procera* Kendrick causal organism of Eastern white pine wilting in conifers culture. (English summary). *Plant Protection* **32**, 313-323.

Hallaksela, A.M. (1977). Microbial flora isolated from Norway spruce stumps. *Acta Forestalia Fennica* **158**, 5-41.

- Halsted, B.D. (1890). Some fungous diseases of the sweet potato. N.J. Agr. Exp. Sta. **76**, 1-32.
- Hansen, E.M. (1978). Incidence of *Verticicladiella wagneri* and *Phellinus weirii* in Douglas-fir adjacent to and away from roads in Western Oregon. *Plant Disease Reporter* **62**, 179-181.
- Hansen, E.M. and Goheen, D.J. (1988). Rate of increase of black-stain root disease in Douglas-fir plantations in Oregon and Washington. *Canadian Journal of Forest Research* **18**, 942-946.
- Hansen, E.M., Goheen, D.J., Hessburg, P.F., Witcosky, J.J. and Schowalter, T.D. (1988). Biology and management of black-stain root disease in Douglas-fir. In: *Leptographium* root diseases on conifers. (ed. T.C. Harrington & F.W. Cobb Jr.), American Phytopathological Society, St Paul, Minnesota, pp. 63-80.
- Harding, S. (1995). Fungal associated of *Ips typographus* L. in Denmark-occurrence, frequency and pathogenicity. (ed. Christiansen, E), Proceedings from a symposium held at the Norwegian Forest Research Institute (NISK), Ås, Norway, 31 July 1995 – 2 August, 1995. *Aktuelt fra Skogforsk* 6/95, pp. 36.
- Harrington, T.C. (1981). Cycloheximide sensitivity as a taxonomic character in *Ceratocystis*. *Mycologia* **73**, 1123-1129.
- Harrington, T.C. (1982). *Verticicladiella wagneri*: Taxonomy and vector relations. Ph.D. thesis. University of California, Berkeley. 113 pp.
- Harrington, T.C. (1987). New combinations in *Ophiostoma* of *Ceratocystis* species with *Leptographium* anamorphs. *Mycotaxon* **28**, 39-43.
- Harrington, T. C. (1988). *Leptographium* species, their distributions, hosts and insect vectors. In: *Leptographium* root diseases on conifers. (ed. T.C. Harrington & F.W. Cobb Jr.), American Phytopathological Society, St. Paul, Minnesota, pp. 1-39.
- Harrington, T.C. (1992). *Leptographium*. In: Methods for research on soilborne

phytopathogenic fungi. (eds. L.L. Singleton, J.D. Mihail and C.M. Rush). American Phytopathological Society Press, St. Paul, Minnesota, pp129-133.

Harrington, T.C. (1993), Diseases of conifers caused by species of *Ophiostoma* and *Leptographium* In: *Ceratocystis and Ophiostoma: Taxonomy, Ecology and Pathogenicity* (eds. M.J. Wingfield, K.A. Seifert and J.F. Webber), American Phytopathological Society, St. Paul, Minnesota, pp. 161-172.

Harrington, T.C. and Cobb, F.W. (Jr.). (1983). Pathogenicity of *Leptographium* and *Verticicladiella* spp. isolated from roots of Western North American conifers. *Phytopathology* **73**, 596-599.

Harrington, T.C., Reinhart, C., Thornburgh, D.A. and Cobb, F.W. (jr.) (1983). Association of black-stain root disease with precommercial thinning of douglas-fir. *Forest Science* **29**, 12-14.

Harrington, T.C. and Cobb, F.W. (jr.) (1984). Host specialization of three morphological variants of *Verticicladiella wageneri*. *Phytopathology* **74**, 286-290.

Harrington, T.C., Cobb, F.W. (jr.) and Lownsbery, J.W. (1985). Activity of *Hylastes nigrinus*, a vector of *Verticicladiella wageneri*, in thinned stands of Douglas-fir. *Canadian Journal of Forest Research* **15**, 519-523.

Harrington, T.C. and Cobb, F.W. (Jr.) (1986). Varieties of *Verticicladiella wageneri*. *Mycologia* **78**, 562-567.

Harrington, T.C. and Cobb, F.W. (Jr.) (1987). *Leptographium wageneri* var. *pseudotsugae*, var. nov., cause of black stain root disease on Douglas-fir. *Mycotaxon* **30**, 501-507.

Harrington, T.C. and Cobb, F.W. (Jr.) (1988). *Leptographium* root diseases on conifers. American Phytopathological Society Press, St. Paul Minnesota, pp. 150.

Hausner, G., Reid, J. and Klassen, G.R. (1993a). On the subdivision of *Ceratocystis* s.l., based on partial ribosomal DNA sequences. *Canadian Journal of Botany* **71**, 52-

63.

Hausner, G., Reid, J. and Klassen, G.R. (1993b). *Ceratocystiopsis*: a reappraisal based on molecular criteria. *Mycological Research* **97**, 625-633.

Hawthorne, B.T. and Rees-George, J. (1996). Use of nitrate non-utilizing mutants to study vegetative incompatibility in *Fusarium solani* (*Nectria haematococca*), especially members of mating populations I, V and VI. *Mycological Research* **100**, 1075-1081.

Helms, J.A., Cobb, F.W. (Jr.) and Whitney, H.S. (1971). Effect of infection by *Verticicladiella wageneri* on the physiology of *Pinus ponderosa*. *Phytopathology* **61**, 920-925.

Hessburg, P.F. and Hansen, E.M. (1982). Histopathology of *Verticicladiella wageneri* in Douglas-fir. *Phytopathology* **72**, 930.

Hessburg, P.F. and Hansen, E.M. (1983). The effect of soil temperature on the *in vivo* growth and development of *Verticicladiella wageneri* in Douglas-fir roots. Fourth International Congress of Plant Pathology, abstract 287.

Hessburg, P.F. and Hansen, E.M. (1986a). Mechanisms of intertree transmission of *Ceratocystis wageneri* in young Douglas-fir. *Canadian Journal of Forest Research* **16**, 1250-1254.

Hessburg, P.F. and Hansen, E.M. (1986b). Soil temperature and rate of colonization of *Ceratocystis wageneri* in Douglas-fir. *Phytopathology* **76**, 627-631.

Hessburg, P.F. and Hansen, E.M. (1987). Pathological anatomy of black stain root disease in Douglas-fir. *Canadian Journal of Botany* **65**, 962-971.

Highley, L. and Tattar, T.A. (1985). *Leptographium terebrantis* and black turpentine beetles associated with blue stain and mortality of black and scots pines on Cape Cod, Massachusetts. *Plant Disease* **69**, 528-530.

Highley, L. and Tattar, T.A. (1987). Patterns of bluestain discoloration and associated organisms in Japanese Black and Scots pines on Cape Cod, Massachusetts. *Arboricultural Journal* **11**, 105-113.

Hicks, B.R. (1973). Growth of *Verticicladiella wageneri* through soil and infection of *Pinus ponderosa* as related to selected soil properties. M.Sc. thesis. University of California, Berkeley. pp. 135.

Hicks, B.R., Cobb, F.W. (jr.) and Gersper, P.L. (1980). Isolation of *Ceratocystis wageneri* from forest soil with a selective medium. *Phytopathology* **70**, 880-883.

Hinds, T.E. (1972). Insect transmission of *Ceratocystis* species associated with aspen canker. *Phytopathology* **62**, 221-225.

Hinds, T.E and Davidson, R.W. (1972). *Ceratocystis* species associated with the aspen ambrosia beetle. *Mycologia* **64**, 405-409.

Hobson, K.R., Parmeter, J.R. and Wood, D.L. (1991). Bluestain fungi and the xylem occlusion enigma. Proceedings: North American Forest Insect Work Conference. (ed: D.C. Allen & L.P. Abrahamson), pp156-157.

Horner, W.E. and Alexander, S.A. (1983a). *Verticicladiella procera* in pine seed orchards in the south. (Abstract) *Phytopathology* **73**, 835.

Horner, W.E. and Alexander, S.A. (1983b). *Verticicladiella procera* on *Pinus sylvestris* christmas trees. *Phytopathology* **73**, 949.

Horrner, W.E., Alexander, S.A. and Julian, M.M. (1986). Qualitative determination of cellulose in the cell walls of *Verticicladiella procera*. *Mycologia* **78**, 300-303.

Horner, W.E., Alexander, S.A. and Lewis, K.J. (1987). Colonization patterns of *Verticicladiella procera* in Scots and Eastern Pine and associated resin - soaking, reduced sapwood moisture content, and reduced needle water potential. *Phytopathology* **77**, 557-560.

- Hornvedt, R., Christiansen, E., Solheim, H. and Wang, S. (1983). Artificial inoculation with *Ips typographus* -associated blue-stain fungi can kill healthy Norway spruce trees. *Reports of the Norwegian Forest Research Institute* **38**, 5-20.
- Houston, D.R. (1969). Basal canker of white pine. *Forest Science* **15**, 66-83.
- Hubert, E.E. (1953). Studies of *Leptographium* isolated from Western white pine. *Phytopathology* **43**, 637-641.
- Hughes, S.J. (1953). Conidiophores, conidia and classification. *Canadian Journal of Botany* **31**, 577-659.
- Hunt, J. (1956). Taxonomy of the genus *Ceratocystis*. *Lloydia* **19**, 1-58.
- Hunt, R.S and Morrison, D.J. (1980). Black stain root disease in British Columbia. *Pest Leaflet* **67**, 1-4.
- Hunt, R.S. and Morrison, D.J. (1986). Black-stain root disease on lodgepole pine in British Columbia. *Canadian Journal for Forest Research* **16**, 996-999.
- Hutchison, L.J. and Reid, J. (1988). Taxonomy of some potential wood-staining fungi from New Zealand 1. Ophiostomataceae. *New Zealand Journal of Botany*, **26**, 63-81.
- Isaev A.S., Rozhkov A.S., Kiselev V.V.(1988) Fir sawyer beetle *Monochamus urussovi* (Fisch.)-Novosibirsk:"Nauka". -270p. (in Russian)
- Ivory, M.H and Andre, W. (1995). A preliminary report of *Verticillium* wilt of Takamaka (*Calophyllum inophyllum* L.) in Seychelles. *The African Journal of Mycology and Biotechnology* **3**, 169-170.
- Jacobi, W.R. (1992). Potential insect vectors of the black stain root disease pathogen on Southern Vancouver Island. *Journal of the Entomological Society of British Columbia* **89**, 54-56.

- Jacobs, M. (1966). The compact key. *Flora Malesiana Bulletin* **21**, 1428-1431.
- Jacobs, K; Wingfield, M.J. and Bergdahl, D. (1997). A new species of *Ophiostoma* from North America, similar to *Ophiostoma penicillatum*. *Canadian Journal of Botany* **75**, 1315-1322.
- Jacobs, K., Wingfield, M.J, Wingfield, B.D. and Yamaoka, Y. (1998). Comparison of *Ophiostoma huntii* and *O. europioides* and description of *O. aenigmaticum* sp. nov. *Mycological Research* **102**, 289-294.
- Jacobs, K., Wingfield, M.J., Crous, P.W. and Harrington, T.C. (1998) *Leptographium engelmannii*, a synonym of *Leptographium abietinum*, and description of *Leptographium hughesii* sp. nov. *Canadian Journal of Botany* **76**, 1660 - 1667.
- Jacobs, K., Wingfield, M.J. Coetsee, C., Kirisits, T. and Wingfield, B.D. (1999). *Leptographium guttulatum* sp. nov., a new species from spruce and pine in Europe. *Mycologia* (in press).
- Jacobs, K., Wingfield, M.J. and Roux, J. (1999). *Leptographium eucalyptophilum*, a new species from *Eucalyptus* in the Congo. *South African Journal of Botany* (in press).
- Jacobs, K, Wingfield, M.J., Uzunovic, A. and Frisullo, S. (1999). Three new species of *Leptographium* from pine, similar to *L. procerum*. *Mycological Research* (submitted)
- Jacobs, K.; Wingfield, M.J and Bergdahl, B.D. (1999). New *Leptographium* species from Indonesia and Eastern North America. *Mycoscience* (in press).
- Jacobs, K., Wingfield, M.J., Pashenova, N.V. and Vetrova, V.P. (1999). A new *Leptographium* species from Russia. *Mycological Research* (submitted).
- Jewell, T.R. (1974). A qualitative study of cellulose distribution in *Ceratocystis* and *Europhium*. *Mycologia* **66**, 139-146.

Jong, S.C. and Davis, E.E. (1972). *Phialocephala humicola*, a new hyphomycete. *Mycologia* **64**, 1351-1356.

Jooste, W.J. (1978). *Leptographium reconditum* sp. nov. and observations on conidiogenesis in *Verticicladiella*. *Transactions of the British Mycological Society* **70**, 152-155.

Joubert, W.A. and Britz, T.J. (1987). A simple and inexpensive method for the long term preservation of microbial cultures. *Journal of Microbiological Methods* **7**, 73-76.

Kaneko, S. and Harrington, T.C. (1990). *Leptographium truncatum* isolated from Japanese red and black pines. *Report of the Tottori Mycological Institute* **28**, 171-174.

Kendrick, W.B. (1961). The *Leptographium* complex. *Phialocephala* gen. nov. *Canadian Journal of Botany* **39**, 1080-1085.

Kendrick, W.B. (1962). The *Leptographium* complex. *Verticicladiella* S.Hughes. *Canadian Journal of Botany* **40**, 771-797.

Kendrick, W.B. (1963a). The *Leptographium* complex. *Penicillium repens* C. and E. *Canadian Journal of Botany* **41**, 574-577.

Kendrick, W.B. (1963b). The *Leptographium* complex: Two new species of *Phialocephala*. *Canadian Journal of Botany* **41**, 1015-1023.

Kendrick, W.B. (1964a). The *Leptographium* complex. *Hantzschia* Auersw. *Canadian Journal of Botany* **42**, 1291-1295.

Kendrick, W.B. (1964b). The *Leptographium* complex. *Scopularia venusta* Preuss. *Canadian Journal of Botany* **42**, 1119-1122.

Kendrick, W.B. and Molnar, A.C. (1965). A new *Ceratocystis* and its *Verticicladiella* imperfect state associated with the bark beetle *Dryocoetus confusus* on *Abies lasiocarpa*. *Canadian Journal of Botany* **43**, 39-43.

Kilbertus, G., Mangenot, F and Radtke, D. (1980). L'alteration des bios d'oeuvre de hetre (*Fagus silvatica* L.) par *Fusarium solani* (Mart.) Sacc. et *Verticicladiella procera* Kendrick. *Cryptogamie, Mycologie* 1, 97-101.

Klepzig, K. D., Raffa, K. F. and Smalley, E. B. (1991). Association of an insect-fungal complex with red pine decline in Wisconsin. *For. Sci.* 37, 1119-1139.

Klittich, C.J.R. and Leslie, J.F. (1988). Nitrate reduction mutants of *Fusarium moniliforme* (*Gibberella fujikuroi*). *Genetics* 118, 417-423.

Korf, R.P. (1972). Synoptic key to the genera of the Pezizales. *Mycologia* 64, 937 - 994.

Korf, R.P. and Zhuang, W.Y. (1985). A synoptic key to the species of *Lambertella* (Sclerotiniaceae), with comments on a version prepared for taxadat, Anderegg's computer program. *Mycotaxon* 24, 361-386.

Krokene, P. (1996). The role of blue-stain fungi in tree-killing by bark beetles. PhD. thesis. University of Oslo, Norway. pp115.

Krokene, P. and Solheim, H. (1996). Fungal associates of five bark beetle species colonizing Norway spruce. *Canadian Journal of Forest Research* 26, 2115-2122.

Kulhavy, D.L., Chacko, R.J. and Partridge, A.D. (1978). Some decay and disease fungi isolated from Western White pine in Northern Idaho. *Plant Disease Reporter* 62, 332-336.

Kulhavy, D.L., Partridge, A.D. and Stark, R.W. (1984). Root diseases and blister rust associated with bark beetles (Coleoptera: Scolytidae) in western white pine in Idaho. *Environmental Entomologist* 13, 813-817.

Lackner, A.L. (1981). Incidence and pathogenicity of *Verticicladiella procera* Kendrick on pines in Virginia. M.Sc. thesis. Virginia Polytechnic Institute. pp57.

Lackner, A.L. and Alexander, S.A. (1982). Occurrence and pathogenicity of

Verticicladiella procera in Christmas tree plantations in Virginia. *Plant Disease* **66**, 211-212.

Lackner, A.L. and Alexander, S.A. (1983). Root disease and insect infestations on air-pollution-sensitive *Pinus strobus* and studies of pathogenicity of *Verticicladiella procera*. *Plant Disease* **67**, 679-681.

Lackner A.L. and Alexander, S.A. (1984). Incidence and development of *Verticicladiella procera* in Virginia Christmas tree plantations. *Plant Disease* **68**, 210-212.

Lagerberg, T., Lundberg, G and Melin, E. (1927). Biological and practical researches into blueing in pine and spruce. *Svenska Skogsvårdsföreningens Tidskrift* **25**, 145-272.

Landis, T.D. and Helburg, L.B. (1976). Black stain root disease of Pinyon pine in Colorado. *Plant Disease Reporter* **60**, 713-717.

Lane, B.B and Goheen, D.J. (1979). Incidence of root disease in bark beetle-infested eastern Oregon and Washington true firs. *Plant Disease Reporter* **63**, 262-266.

Långström, B. and Hellqvist, C. (1993). Induced and spontaneous attacks by pine shoot beetles on young Scots pine trees : tree mortality and beetle. *J. Appl. Entomol.* **115**, 25-36.

Långström, B., Solheim, H., Hellqvist, C. and Gref, R. (1993). Effects of pruning young Scots pines on host vigor and susceptibility to *Leptographium wingfieldii* and *Ophiostoma minus*, two blue-stain fungi associated with *Tomicus piniperda*. *Eur. J. For. Path.* **25**, 400-415.

Lawson, T.T. and Cobb, F.W. (1987a). Growth reductions in young Douglas-fir infected with *Verticicladiella wagneri*. *Phytopathology* **77**, 1727.

Lawson, T.T. and Cobb, F.W. (1987b). Pathological heartwood formation in pole-

- sized Douglas-fir infected with *Verticicladiella wageneri*. *Phytopathology* **77**, 1727.
- Leach, J.G., Orr, L.W. and Christensen, C. (1934). The interrelationships of bark beetles and blue-staining fungi in felled Norway pine timber. *Journal of Agricultural Research* **49**, 315-341.
- Leaphart, C.D. (1956). Physiological studies of some fungi associated with pole blight of western white pine. *Mycologia* **48**, 25-40.
- Leaphart, C.D. (1958). Pole blight - How it may influence western white pine management in light of current knowledge. *Journal of Forestry* **56**, 746-751.
- Leaphart, C.D. (1960). A root stain disease of Eastern White Pine. *Plant Disease Reporter* **44**, 704 - 706.
- Leaphart, C.D. and Gill, L.S. (1959). Effects of inoculations with *Leptographium* spp. on western white pine. *Phytopathology* **49**, 350-353.
- Leaphart, C.D., Copeland, O.T. (jr.) and Graham, D.P. (1957). Pole blight of Western White pine. Forest Pest leaflet 16, 1-4.
- Lewinsohn, Lewinsohn, Bertagnolli and Partridge (1994). Blue-stain and their transport structures on the Douglas-fir beetle. *Canadian Journal of Botany* **24**, 2275-2283.
- Leslie, J.F. (1993). Fungal vegetative compatibility. *Annual Review of Phytopathology* **31**, 127-150.
- Leslie, J.F. and Klein, K.K. (1996). Female fertility and mating type effects on effective population size and evolution in filamentous fungi. *Genetics* **144**, 557-567.
- Lévieux, J., Piou, D., Cassier, P., André, M. and Guillaumin, D. (1994). Association of phytopathogenic fungi for the scots pine (*Pinus sylvestris* L.) with the European pine weevil *Hylobius abietis* (L.) (Col. Curculionidae). *The Canadian Entomologist* **126**, 929-936.

- Lewis, K.J. (1985). Studies on the spread of *Verticicladiella procera* by soil-borne and insect-borne propagules. MSc thesis. Faculty of the Virginia Polytechnic Institute and State University, Virginia, U.S.A., pp. 119.
- Lewis, K.J. and Alexander, S.A. (1985). Germinability over time of *Verticicladiella procera* propagules in artificially infested soil and their inability to cause disease. *Phytopathology* **75**, 1337.
- Lewis, K.J. and Alexander, S.A. (1986). Insects associated with the transmission of *Verticicladiella procera*. *Canadian Journal of Forest Research* **16**, 1330-1333.
- Lewis, K.J., Alexander, S.A and Horner, W.E. (1987). Distribution and efficacy of propagules of *Verticicladiella procera* in soil. *Phytopathology* **77**, 552-556.
- Lieutier, F., Yart, A. Garcia, J., Poupinel, B. and Leveux, J. (1988). Do fungi influence the establishment of bark beetles in Scots pine? In: Mechanisms of woody plant defenses against insects, search for pattern. (ed. W.J. Mattson, J. Leveux & C. Bernard-Dagan), Springer-Verlag, New York, 416pp.
- Lieutier, F. and Yart, A. (1989). Temperature preference of the fungi associated with *Ips sexdentatus* Boern. and *Tomicus piniperda* L. (Coleoptera: Scolytidae). (French with english summary). *Ann. For. Sci.* **46**, 411-415.
- Lieutier, F., Cheniclet, C. and Garcia, J. (1989a). Comparison of the defense reactions of *Pinus pinaster* and *Pinus sylvestris* to attacks by two bark beetles (Coleoptera: Scolytidae) and their associated fungi. *Environmental Entomology* **18**, 228-234.
- Lieutier, F., Yart, A, Garcia, J., Ham, M.C, Morelet, M. and Leveux, J. (1989b). (French with english summary) Phytopathogenic fungi associated with two bark beetles of scots pine (*Pinus sylvestris* L.) and preliminary studies of their aggressiveness for the host. *Ann. Sci. For.* **46**, 201-216.
- Livingston, W.H. and Wingfield, M.J. (1982). First report of *Verticicladiella procera*

on pines in Minnesota. *Plant Disease* **66**, 260-261.

Livingston, W.H., Mangini, A.C., Kinzer, H.G. and Mielke, M.E. (1983). Association of root diseases and bark beetles (Coleoptera: Scolytidae) with *Pinus ponderosa* in New Mexico. *Plant Disease* **67**, 674-676.

Lorenzini, G. and Gambogi, P. (1976). Decline of *Pinus pinea* associated with the presence of *Verticicladiella* sp. (preliminary note) (Italian). *Informatore Fitopatologico* **5**, 5-8

Mackenzie, M. and Dick, M. (1984). *Verticicladiella* root disease. *Forest Pathology in New Zealand* **8**, 1-4.

Malloch, D. and Blackwell, M. (1993). Dispersal biology of the ophiostomatoid fungi. In: *Ceratocystis and Ophiostoma. Taxonomy, ecology and pathogenicity.* (eds. M.J. Wingfield, K.A. Seifert and J.F. Webber) American Phytopathological Society Press. pp. 195-206.

Marais, G.J. (1996). Fungi associated with infructescences of *Protea* species with special reference to the Ophiostomatales. Ph.D. thesis. University of the Orange Free State. pp.133.

Marais, G.J., Wingfield, M.J., Viljoen, C.D. and Wingfield, B.D. (1998). A new ophiostomatoid genus from *Protea* infructescences. *Mycologia* **90**, 136-141.

Marais, G.J. and Wingfield, M.J. (1999a). Differentiation between species in *Ceratocystis sensu lato* based on substrate utilisation. *South African Journal of Botany*. (in press).

Marais, G.J. and Wingfield, M.J. (1999b). Cycloheximide sensitivity and cell saccharides in *Ceratocystis sensu lato* with special emphasis on species from Proteaceae. *South African Journal of Botany*. (in press).

Masuya, H., Kaneko, S., and Yamaoka, Y. (1998). Blue stain fungi associated with *Tomicus piniperda* (Coleoptera: Scolytidae) on Japanese red pine. *Journal of*

Forestry Research **3**, 213-219.

Masuya, H., Kaneko, S., Yamaoka, Y., Osawa, M. (1999). Comparisons of ophiostomatoid fungi associated with *Tomicus piniperda* and *T. minor* in Japanese red pine. *Journal of Forestry Research* **4**, 131-135.

Masuya, H., Wingfield, M.J., Kaneko, S. and Yamaoka, Y. (1999). *Leptographium pinidensiflorae* sp. nov. from Japanese red pine. *Mycological Research* (submitted).

Mathiesen, A. (1950). Über einige mit bokenkäfern assoziierte bläuepilze in Schweden. *Oikos* **2**, 275-308.

Mathiesen, A. (1951). Einige neue *Ophiostoma*-arten in Schweden. *Svensk Botanisk Tidskrift* **45**, 203-232.

Mathiesen-Käärik, A. (1953). Eine Übersicht über die gewöhnlichsten mit Borkenkäfern assoziierten Bläuepilze in Schweden und einige für Schweden neue Bläuepilze. *Meddelanden Från Statens Skogforskningsinstitut* **43**, 3-74

Mathiesen-Käärik, A. (1960). Studies on the ecology, taxonomy and physiology of Swedish insect-associated blue-stain fungi, especially the genus *Ceratocystis*. *Oikos* **II**, 1-25.

Mathre, D.E. (1964). Survey of *Ceratocystis* spp. associated with bark beetles in California. *Contributions from Boyce Thompson Institute* **22**, 353-362.

Meyer, G.J., Hindal, D.F. and Quinn, D.O. (1983). Occurrence of white pine root decline in western Virginia. *Phytopathology* **73**, 967.

Mielke, M.E. (1979). *Verticicladiella* species and associated root stain and decay fungi isolated from symptomatic Northern Rocky Mountain conifers. M.Sc. thesis, University of Idaho, Moscow.

Mielke, M.E. (1981). Pathogenicity of *Verticicladiella penicillata* (Grosm.) Kendrick to Northern Idaho conifers. *Forest Science* **27**, 103-110.

- Miller, D.R. and Veirs, E.I. (1968). *Verticicladiella wageneri* found on Douglas-fir in El Dorado county, California. *Plant Disease Reporter* **52**, 393.
- Minter, D.W., Kirk, P.M. and Sutton, B.C. (1982). Holoblastic phialides. *Transactions of the British Mycological Society* **79**, 75-93.
- Minter, D.W., Kirk, P.M. and Sutton, B.C. (1983). Thallic phialides. *Transactions of the British Mycological Society* **80**, 39-66.
- Mouton, M, Wingfield, M.J. and Van Wyk, P.S. (1992). The anamorph of *Ophiostoma francke-grosmanniae* is a *Leptographium*. *Mycologia* **84**, 857-862.
- Mouton, M, Wingfield, M.J. and Van Wyk, P.S. (1994). Conidium development in anamorphs of *Ceratocystis sensu lato*: a review. *South African Journal of Science* **90**, 293-298.
- Molnar, A.C. (1965). Pathogenic fungi associated with a bark beetle on alpine fir. *Canadian Journal of Botany* **43**, 563-570.
- Moreau, C. (1952). Coexistence des formes *Thielaviopsis* et *Graphium* chez une souche de *Ceratocystis major* (Van Beyma) nov. comb. *Revue de Mycologie, Suppl. Colonial* **17**, 17-25.
- Morelet, M. (1986). Les *Verticicladiella* des pins en liaison avec les phénomènes de dépérissement. *EPPO Bulletin* **16**, 473-477.
- Morelet, M. (1988). Observations sur trios Deutéromycètes inféodés aux pins. *Annales de la Société des Sciences Naturelles et d'Archéologie de Toulon et du Var*. **40**, 41-43.
- Morrison, D.J. and Hunt, R.S. (1988). *Leptographium* species associated with root disease of conifers in British Columbia. In: *Leptographium* root diseases on conifers. (ed. T.C. Harrington & F.W. Cobb Jr.), American Phytopathological Society, St. Paul, Minnesota, pp. 97-112.

- Moser, J.C. (1985). Use of the sporothecae by phoretic *Tarsonemus* mites to transport ascospores of coniferous bluestain fungi. *Transactions of the British Mycological Society* **84**, 750-753.
- Moser, J.C., Perry, T.J and Solheim, H. (1989). Ascospores hyperphoretic on mites associated with *Ips typographus*. *Mycological Research* **93**, 513-517.
- Münch, E. (1907). Die Blaufäule des Nadelholzes. *Naturwissenschaftliche Zeitschrift für forest und Landw.* **5**, 531-573.
- Nelson, R.M. (1934). Effect of bluestain fungi on southern pines attacked by bark beetles. *Phytopath. Z. Bd.* **7**, 327-353.
- Nelson, R.M. and Beal, J.A. (1929). Experiments with bluestain fungi on southern pines. *Phytopathology* **19**, 1101-1106.
- Nevill, R.J. and Alexander, S.A. (1992a). Transmission of *Leptographium procerum* to Eastern White pine by *Hylobius pales* and *Pissodes nemorensis* (Coleoptera: Curculionidae). *Plant Disease* **76**, 307-310.
- Nevill, R.J. and Alexander, S.A. (1992b). Distribution of *Hylobius pales* and *Pissodes nemorensis* (Coleoptera: Curculionidae) within Christmas Tree Plantations with Procerum root disease. *Environmental Entomology*, **21**, 1077 - 1085.
- Nevill, R.J., Kelley, W.D., Hess, N.J. and Perry, T.J. (1995). Pathogenicity to loblolly pines of fungi recovered from trees attacked by southern pine beetles. *Southern Journal of Applied Forestry* **19**, 78-83.
- Olchowecki, A. and Reid, J. (1974). The genus *Ceratocystis* in Manitoba. *Canadian Journal of Botany* **52**, 1675-1711.
- Onofri, S. and Zucconi, L. (1984). Two new species of the genus *Phialocephala*. *Mycotaxon* **20**, 185-195.
- Orlic, S., Harapin, M., Halambek, M. and Mayer, B. (1973). Dieback of eastern

white pine (*Pinus strobus* L.) in cultures. *Preštampani iz šumarkog lista* br. 9-10.

Otrosina, W.J. (1986). Electrophoretic characteristics of three *Verticicladiella wageneri* variants. (Abstract). *Phytopathology*, **76**, 845.

Otrosina, W.J. and Cobb, F.W. (Jr.) (1987). Analysis of allozymes of three distinct variants of *Verticicladiella wageneri* isolated from conifers in Western North America. *Phytopathology* **77**, 1360-1363.

Otrosina, W.J., Cobb, F.W. (Jr.) and Popenuck, T. (1987). Variation in virulence within the host specific variants of *Verticicladiella wageneri*. (Abstract). *Phytopathology* **77**, 1757.

Otrosina, W.J., Hess, N.J., Zarnoch, S.J., Perry, T.J. and Jones, J.P. (1997). Blue-stain fungi associated with roots of Southern pine trees attacked by the Southern Pine beetle, *Dendroctonus frontalis*. *Plant Disease* **81**, 942-945.

Paine, T.D., Raffa, K.F. and Harrington, T.C. (1990). Interactions among Scolytid bark beetles, their associated fungi, and live host conifers. *Annual Review of Entomology* **42**, 179-206.

Parker, A.K. (1957a). *Europhium*, a new genus of the ascomycetes with a *Leptographium* imperfect state. *Canadian Journal of Botany* **35**, 173-179.

Parker, A.K. (1957b). The nature of the association of *Europhium trinacriforme* with pole blight lesions. *Canadian Journal of Botany* **35**, 845-856.

Partridge, A.D. and Bertagnole, C.L. (1980). *Verticicladiella* species and associated organisms causing black stain root disease of inland Northwest pines. Proceedings of the twenty-eight annual Western international forest disease work conference. pp. 45-53.

Peck, C.H. (1879). Report of the Botanist. *New York State Museum Report* **31**, 19-60.

Perry, T.J. (1991). A synopsis of the taxonomic revisions in the genus *Ceratocystis* including a review of blue-staining species associated with *Dendroctonus* bark beetles. General Technical Report SO-86, New Orleans, L.A.: U.S. Department of Agriculture, Forst Service, Southern Forest Experiment Station 16 pp.

Pest Alert (1977). White pine root decline. US Department of Agriculture leaflet 1977-751-868.

Preuss, G.T. (1851). Uebersicht untersuchter Pilze, besonders aus der Umgegend von Hoyerswerda.

Prey, A.J. (1975). Forest Pest conditions in Wisconsin. Annual Report p. 21, Department of Natural Resources, Madison, Wisconsin.

Puhalla, J.E. (1985). Classification of strains of *Fusarium oxysporum* on the basis of vegetative compatibility. *Canadian Journal of Botany* **63**, 179-183.

Raffa, K.F. (1995). Bark beetles, fungi, trees and humans: Four perspectives, four agendas. (ed. Christiansen, E), Proceedings from a symposium held at the Norwegian Forest Research Institute (NISK), Ås, Norway, 31 July 1995 – 2 August, 1995. *Aktuelt fra Skogforsk* 6/95, pp. 7-9.

Raffa, K.F. and Smalley, E.B. (1995). Interaction of pre-attack and induced monoterpane concentrations in host conifer defense against bark beetle-fungal complexes. *Oecologia* **102**, 285-295.

Ram, C. and Ram, A. (1972). Timber-attacking fungi from the state of Mara-Nhão, Brazil. Some new or interesting wood staining fungi. *Broteria* **41**, 89-112.

Rayner, R.W. (1970). A Mycological color chart. Commonwealth Mycological Institute and British Mycological Society, Kew, Surrey and British Mycological Society.

Redfern, D.B. (1989). The roles of the bark beetles *Ips cembrae*, the woodwasp *Urocerus gigas* and associated fungi in dieback and death of larches. In: Insect-

fungus interactions. (ed: N. Wilding, N.M. Collins, P.M. Hammond and J.F. Webber), pp. 195-204, Academic press, London.

Rennerfelt, E. (1950). Über den zusammenhang zwishchen dem verblauen des holzes und den insekten. *Oikos* **2**, 120-137.

Reynolds, K.M. (1992). Relations between activity of *Dendroctonus rufipennis* Kirby on Lutz spruce and blue-stain associated with *Leptographium abietinum* (Peck) Wingfield. *Forest Ecology and Management* **47**, 71-86

Robinson, R.C. (1962). Blue stain fungi in lodgepole pine (*Pinus contorta* Dougl. var. *latifolia* Engelm.) infested by the mountain pine beetle (*Dendroctonus monticolae* Hopk.). *Canadian Journal of Botany* **40**, 609-614.

Robinson-Jeffrey, R.C. and Grinchenko, A.H.H. (1964). A new fungus in the genus *Ceratocystis* occurring on blue-stained lodgepole pine attacked by bark beetles. *Canadian Journal of Botany* **42**, 527-532.

Robinson-Jeffrey, and Davidson, R.W. (1968). Three new *Europhium* species with *Verticicladiella* imperfect states on blue-stained pine. *Canadian Journal of Botany* **46**, 1523-1527.

Rosinski, M.A. and Campana, R.J. (1964). Chemical analysis of the cell wall of *Ceratocystis ulmi*. *Mycologia* **56**, 738-744.

Ross, D.W., Fenn, P. and Stephen, F.M. (1992). Growth of southern pine beetle associated fungi in relation to induced wound response in loblolly pine. *Canadian Journal of Forest Research* **22**, 1851-1859.

Ross, D.W. and Solheim, H. (1995). Pathogenicity of the Douglas-fir beetle associated fungi, *Ophiostoma pseudotsugae* and *Leptographium abietinum*, to Douglas-fir. (ed. Christiansen, E), Proceedings from a symposium held at the Norwegian Forest Research Institute (NISK), Ås, Norway, 31 July 1995 – 2 August, 1995. *Aktuelt fra Skogforsk* 6/95, pp. 17.

Ross, D.W. and Solheim, H. (1996). Douglas-fir and western larch defensive reactions to *Leptographium abietinum* and *Ophiostoma pseudotsugae*. In: Dynamics of forest herbivore: Quest for pattern and principles. (eds. Matson, W.J., Niemelä, P. & Rousi, M.), USDA For. Serv. Gen. Tech. Rep. N.C-183, For.Exp.Sta. St. Paul, MN, pp. 224-227.

Ross, D.W. and Solheim, H. (1997). Pathogenicity to Douglas-fir of *Ophiostoma pseudotsugae* and *Leptographium abietinum*, fungi associated with Douglas-fir beetle. *Canadian Journal of Forest Research* **27**, 39-43.

Rumbold, C.T. (1931). Two blue-staining fungi associated with bark-beetle infestation of pines. *Journal of Agricultural Research* **43**, 847-873.

Rumbold, C.T. (1936). Three blue-staining fungi, including two new species, associated with bark beetles. *Journal of Agricultural Research* **52**, 419-437.

Saccardo, P.A. (1886). *Sylloge Fungorum* **4**, 270-275.

Samuels, G.J. (1993). The case for distinguishing *Ceratocystis* and *Ophiostoma*. In: *Ceratocystis* and *Ophiostoma*. Taxonomy, Ecology and Pathogenicity. (ed. M.J. Wingfield, K.A. Seifert and J.F. Webber). American Phytopathological Press, St. Paul, Minnesota, pp. 15-20.

Samuels, G.J. and Müller, E. (1978). Life-history studies of Brazilian Ascomycetes 5. Two new species of *Ophiostoma* and their *Sporothrix* anamorphs. *Sydowia* **31**, 169-179.

Samuels, G.J. and Seifert, K.A. (1995). The impact of molecular characteristics on systematics of filamentous ascomycetes. *Annual Review of Phytopathology* **33**, 37-67.

Seifert, K.A. and Okada, G. (1993). *Graphium* anamorphs of *Ophiostoma* species and similar anamorphs of other Ascomycetes. In: *Ceratocystis* and *Ophiostoma*. Taxonomy, Ecology and Pathogenicity. (ed. M.J. Wingfield, K.A. Seifert and J.F.

- Webber). American Phytopathological Press, St. Paul, Minnesota, pp. 27-42.
- Seifert, K.A., Webber, J.F. and Wingfield, M.J. (1993). Methods of studying species of *Ophiostoma* and *Ceratocystis*. In: *Ceratocystis* and *Ophiostoma*. Taxonomy, Ecology and Pathogenicity. (ed. M.J. Wingfield, K.A. Seifert and J.F. Webber). American Phytopathological Press, St. Paul, Minnesota, pp. 255-259.
- Shaw, C.G. and Hubert, E.E. (1952). A review of the *Leptographium-Scopularia-Hantszchia* nomenclature. *Mycologia* **44**, 693-704.
- Shaw, C.G III and Dick, M. (1980). *Verticicladiella* root disease of *Pinus strobus* in New Zealand. *Plant Disease* **64**, 96-98.
- Siegfried, A.L., Seifert, K.A. and Bilmer, B.C. (1992). A new species of *Phialocephala* (Hyphomycetes). *Canadian Journal of Botany* **70**, 2484-2489.
- Siemaszko, W. (1939). Fungi associated with bark-beetles in Poland. *Planta Polonica* **7**, 1-54.
- Sinclair, W.A. and Hudler, G.W. (1980). Tree and shrub pathogens new or noteworthy in New York state. *Plant Disease* **64**, 590-592.
- Singleton, L.L, Mihail, J.D. and Rush, C.D. (1992). Methods for research on soilborne phytopathogenic fungi. (eds). American Phytopathological Society Press, St, Paul, Minnesota, pp 265.
- Sivasithamparam, K. (1975). Two dematiaceous hyphomycetes with a similar mode of conidiogenesis. *Transactions of the British Mycological Society* **64**, 335-338.
- Six, D.L. and Paine, T.D. (1995). Evolutionary association among bark beetles, mycangial fungi, and host conifers. (ed. Christiansen, E), Proceedings from a symposium held at the Norwegian Forest Research Institute (NISK), Ås, Norway, 31 July 1995 – 2 August, 1995. *Aktuelt fra Skogforsk* 6/95, pp. 13-14.
- Six, D.L. and Paine, T.D. (1996). *Leptographium pyrinum* is a mycangial fungus of

Dendroctonus adjunctus. *Mycologia* **88**, 739-744.

Smalley, E.B., Raffa, K.F., Proctor, R.H. and Klepzig, K.D. (1993). Tree responses to infection by species of *Ophiostoma* and *Ceratocystis*. In: *Ceratocystis* and *Ophiostoma*. Taxonomy, Ecology and Pathogenicity. (ed. M.J. Wingfield, K.A. Seifert and J.F. Webber). American Phytopathological Press, St. Paul, Minnesota, pp. 207-217.

Smith, M.J., Patik, C.M. and Rosinski, M.A. (1967). A comparison of cellulose production in the genus *Ceratocystis*. *Mycologia* **59**, 965-969.

Smith, R.S., Jr. (1967). *Verticicladiella* root disease of pines. *Phytopathology* **57**, 935-938.

Smith, R.S. (Jr.) (1969). The inability of *Verticicladiella wageneri* to break down cellulose. *Phytopathology* **58**, 1050.

Smith, R.S. (Jr.). (1974). Prospects for reducing forest damages from the black stain root disease caused by *Verticicladiella wageneri*. *California Plant Pathology* **19**, 1-2.

Smith, R.S. (Jr.) and Graham, D. (1975). Black stain root disease of conifers. *Forest Pest Leaflet* **145**, 1-4.

Smith, V.L. (1991). Occurrence of Procerum root disease caused by *Leptographium procerum* on White pine in Connecticut. *Plant Disease* **75**, 431.

Snyder, W.C. and Hansen, H.N. (1946). Control of culture mites by cigarette paper barriers. *Mycologia* **38**, 455-462.

Solheim, H. (1986). Species of Ophiostomataceae isolated from *Picea abies* infested by the bark beetle *Ips typographus*. *Nordic Journal of Botany* **6**, 199-207.

Solheim, H. (1992a). Fungal succession in sapwood of Norway spruce infested by the bark beetle *Ips typographus*. *European Journal of Forest Pathology* **22**, 136-148.

Solheim, H. (1992b). The early stages of fungal invasion in Norway spruce infested by the bark beetle *Ips typographus*. *Canadian Journal of Botany* **70**, 1-5.

Solheim, H. (1993a). Ecological aspects of fungi associated with the spruce bark beetle *Ips typographus* in Norway spruce. In: *Ceratocystis and Ophiostoma. Taxonomy, Ecology and Pathogenicity*. (ed. M.J. Wingfield, K.A. Seifert and J.F. Webber). American Phytopathological Press, St. Paul, Minnesota, pp. 21-25.

Solheim, H. (1993b). Fungi associated with the spruce bark beetle *Ips typographus* in an endemic area in Norway. *Scandinavian Journal of Forest Research* **8**, 118-122.

Solheim, H. (1995a). Blue-stain fungi associated with the spruce beetles *Dendroctonus rufipennis*. (ed. Christiansen, E), Proceedings from a symposium held at the Norwegian Forest Research Institute (NISK), Ås, Norway, 31 July 1995 – 2 August, 1995. *Aktuelt fra Skogforsk* 6/95, pp. 43.

Solheim, H. (1995b). A comparison of blue-stain fungi associated with the North American spruce beetle *Dendroctonus rufipennis* and the Eurasian spruce bark beetle *Ips typographus*. In: *Forest pathology research in the Nordic countries 1994*. (ed: D. Aamlid) *Aktuelt fra Skogforsk* **4**, 61-67.

Solheim, H. (1995c). Early stages of blue-stain fungus invasion of lodgepole pine sapwood following mountain pine beetle attack. *Canadian Journal of Botany* **73**, 70-74.

Solheim, H. and Långström, B. (1991). Blue-stain fungi associated with *Tomicus piniperda* in Sweden and preliminary observations on their pathogenicity. *Ann. Sci. For.* **48**, 149-156.

Solheim, H., Långström, B. and Hellqvist, C. (1993). Pathogenicity of the blue-stain fungi *Leptographium wingfieldii* and *Ophiostoma minus* to Scots pine: effect of tree pruning and inoculum density. *Canadian Journal of Forest Research* **23**, 1438-1443.

Solheim, H. and Safranyik, L. (1997). Pathogenicity to Sitka spruce of *Ceratocystis rufipenni* and *Leptographium abietinum*, blue-stain fungi associated with the spruce beetle. *Canadian Journal of Forest Research* **27**, 1336-1341.

Solheim, H. and Krokene, P. (1998). Growth and virulence of *Ceratocystis rufipenni* and three blue-stain fungi isolated from the Douglas-fir beetle. *Canadian Journal of Botany* **76**, 1763-1769.

Strydom, R.C., Wingfield, B.D. and Wingfield, M.J. (1997). Ribosomal DNA sequence comparison of *Leptographium lundbergii* and *L. truncatum* and neotypification of *L. lundbergii*. *Systematic and Applied Microbiology* **20**, 295-300.

Spatafora, J.W and Blackwell, M. (1994). The polyphyletic origins of ophiostomatoid fungi. *Mycological Research* **98**, 1-9.

Spencer, J.F.T. and Gorin, P.A.J. (1971). Systematics of the genera *Ceratocystis* and *Graphium*. Proton magnetic resonance spectra of the mannose containing polysaccharides as an aid in classification. *Mycologia* **63**, 387-402.

Swai, I.S. and Hindal, D.F. (1981). Selective medium for recovering *Verticicladiella procera* from soils and symptomatic white pines. *Plant Disease* **65**, 963-965.

Sydow, H and Sydow, P. (1919). Mykologische Mitteilungen. *Annales Mycologici* **17**, 3-47.

Towers, B. (1977). The occurrence of *Verticicladiella procera* in Pennsylvania: 1976. *Plant Disease Reporter* **61**, 477.

Upadhyay, H.P. (1981). A monograph of *Ceratocystis* and *Ceratocystiopsis*. University of Georgia Press, Athens, Ga.

Upadhyay, H.P. and Kendrick, W.B. (1975). Prodrum for a revision of *Ceratocystis* (Microascales, Ascomycetes) and its conidial states. *Mycologia* **67**, 797-805.

Valkama, H. (1995). Does *Ips duplicatus* transport sapwood staining fungi ? (ed.

Christiansen, E), Proceedings from a symposium held at the Norwegian Forest Research Institute (NISK), Ås, Norway, 31 July 1995 – 2 August, 1995. *Aktuelt fra Skogforsk* 6/95, pp. 44-45.

Van der Westhuizen, K., Wingfield, M.J., Yamaoka, Y., Kemp, G.H.J. and Crous, P.W. (1995). A new species of *Ophiostoma* with a *Leptographium* anamorph from Larch in Japan. *Mycological Research* **99**, 1334-1338.

Van Wyk, P.S. and Wingfield, M.J. (1987). Does sympodial development occur in *Leptographium* spp.? (Abstract). *Phytophylactica* **19**, 127.

Van Wyk, P., Wingfield, M.J. and Marasas, W.F.O. (1987). Differences in synchronisation of stages of conidial development in *Leptographium* species. *Transactions of the British Mycological Society* **90**, 451-456.

Van Wyk, P.W.J. and Wingfield, M.J. (1990). Ascospore development in *Ceratocystis sensu lato* (Fungi): a review. *Bothalia* **20**, 141-145.

Van Wyk, P.W.J., Wingfield, M.J. and Van Wyk, P.S. (1991). Ascospore development in *Ceratocystis moniliformis*. *Mycological Research* **95**, 96-103.

Van Wyk, P.W.J., Wingfield, M.J. and Van Wyk, P.S. (1993). Ultrastructure of centrum and ascospore development in selected *Ceratocystis* and *Ophiostoma* species. In: *Ceratocystis and Ophiostoma. Taxonomy, Ecology and Pathogenicity.* (ed. M.J. Wingfield, K.A. Seifert and J.F. Webber). American Phytopathological Press, St. Paul, Minnesota, pp. 133-138.

Viiri, H. (1997). Fungal associates of the spruce bark beetle *Ips typographus* L. (Coleoptera:Scolytidae) in relation to different trapping methods. *Journal of Applied Entomology* **121**, 529-533.

Viljoen, C.D., Wingfield, M.J., Jacobs, K. and Wingfield, B.D. (1999). *Comuvesica*, a new genus to accommodate *Ceratocystiopsis falcata*. *Mycological Research* (in press).

- Wainhouse, D., Murphy, S., Grieg, B., Webber, J. and Vielle, M. (1998). The role of the bark beetle *Cryphalus trypanus* in the transmission of the vascular wilt pathogen of takamaka (*Calophyllum inophyllum*) in the Seychelles. *Forest Ecology and Management* **108**, 193-199.
- Wagener, W.W. and Mielke, J.L. (1961). A staining fungus root disease of ponderosa, jeffrey and pinyon pines. *Plant Disease Reporter* **45**, 831-835.
- Wagner, R.E. (1977). *Verticicladiella* species associated with *Hylurgops porosus* LeConte (Coleoptera: Scolytidae) in lodgepole pine. M.Sc. thesis. University of Idaho, Moscow.
- Walters, J.W. and Walters, N.R. (1977). *Verticicladiella wagneri* in the Southwest. *Plant Disease Reporter* **61**, 419.
- Weber, G., Spaaij, F. and Wingfield, M.J. (1996). *Leptographium costaricense* sp. nov., a new species from roots of *Talauma sambuensis*. *Mycological Research* **100**, 732-736.
- Webber, J.F. and Hansen, E.M. (1990). Susceptibility of European and north-west American conifers to the North American vascular pathogen *Leptographium wagneri*. *European Journal of Forest Research* **20**, 347-354.
- Webber, J.F., Jacobs, K and Wingfield M.J. (1999) A re-examination of the vascular wilt pathogen of takamaka (*Calophyllum inophyllum*). *Mycological Research* **103**, 1588-1592.
- Weijman, A.C.M. and De Hoog, G.S. (1975). On the subdivision of the genus *Ceratocystis*. *Antonie van Leeuwenhoek* **41**, 353-360.
- Werner, R.A. (1995). Effects of fertilization on induced host resistance to attack by spruce bark beetles in Lutz spruce in Alaska. (ed. Christiansen, E), Proceedings from a symposium held at the Norwegian Forest Research Institute (NISK), Ås, Norway, 31 July 1995 – 2 August, 1995. *Aktuelt fra Skogforsk* 6/95, pp. 22-23.

Wiehe, P.O. (1949) Wilt of *Calophyllum inophyllum* L. var. *tacamaca* (Willd.) R.E.V. caused by *Haploglyphium calophylli* sp. nov. in Mauritius. *Mycological Papers* **29**, 1-12.

Wilks, D.S., Gersper, P.L. and Cobb, F.W. (jr). (1983). Relation of soil redox potential to infection of Ponderosa pine by *Ceratocystis wageneri*. *Phytopathology* **73**, 1120 - 1125.

Wilks, D.S., Gersper, P.L. and Cobb, F.W. (jr). (1985). Association of soil moisture with spread of *Ceratocystis wageneri* in ponderosa pine. *Plant disease* **69**, 206 - 208.

Wingfield, B.D., Grant, W.S., Wolfaardt, J.F. and Wingfield, M.J. (1994). Ribosomal RNA sequence phylogeny is not congruent with ascospore morphology among species in *Ceratocystis* sensu stricto. *Molecular Biology and Evolution* **11**, 376-383.

Wingfield, B.D., Viljoen C.D. and Wingfield, M.J. (1999). Phylogenetic relationships of ophiostomatoid fungi associated with *Protea* infructescences in South Africa. *Mycological Research* (in press).

Wingfield, M.J. (1982). *Verticicladiella procera* associated with root weevil damage. *Phytopathology* **72**, 141.

Wingfield, M.J. (1983). Association of *Verticicladiella procera* and *Leptographium terebrantis* with insects in the Lake states. *Canadian Journal of Forest Research* **13**, 1238-1245.

Wingfield, M.J. (1985). Reclassification of *Verticicladiella* based on conidial development. *Transactions of the British Mycological Society* **85**, 81-93.

Wingfield, M.J. (1986). Pathogenicity of *Leptographium procerum* and *L. terebrantis* on *Pinus strobus* seedlings and established trees. *European Journal of Forest Pathology* **16**, 299-308.

Wingfield, M.J. (1988). A case for revising the genus *Ceratocystiopsis*. (Abstract)

Phytophylactica **20**, 97.

Wingfield, M.J. (1993a). *Leptographium* species as anamorphs of *Ophiostoma*: progress in establishing acceptable generic and species concepts. In: *Ceratocystis* and *Ophiostoma*. Taxonomy, Ecology and Pathogenicity. (ed. M.J. Wingfield, K.A. Seifert and J.F. Webber). American Phytopathological Press, St. Paul, Minnesota, pp. 43-51.

Wingfield, M.J. (1993b). Problems in delineating the genus *Ceratocystiopsis*. In: *Ceratocystis* and *Ophiostoma*. Taxonomy, Ecology and Pathogenicity. (ed. M.J. Wingfield, K.A. Seifert and J.F. Webber). American Phytopathological Press, St. Paul, Minnesota, pp. 21-25.

Wingfield, M.J. and Knox-Davies, P.S. (1980a). Root-disease, associated with *Verticicladiella alacris*, of pines in South Africa. *Plant Disease* **64**, 569-571.

Wingfield, M.J. and Knox-Davies, P.S. (1980b). Observations on diseases in pine and eucalyptus plantations in South Africa. *Phytophylactica* **12**, 57-63.

Wingfield, M.J. and Marasas, W.F.O. (1980). *Verticicladiella alacris* sp. nov., associated with root disease of pines in South Africa. *Transactions of the British Mycological Society* **75**, 21-28.

Wingfield, M.J. and Marasas, W.F.O. (1981). *Verticicladiella alacris*, a synonym of *V. serpens*. *Transactions of the British Mycological Society* **76**, 508-510.

Wingfield, M.J. and Marasas, W.F.O. (1983). Some *Verticicladiella* species, including *V. truncata* sp. nov., associated with root diseases of pine in New Zealand and South Africa. *Transactions of the British Mycological Society* **80**, 231-236.

Wingfield, M.J., Van Wyk, P.S. and Wingfield, B.D. (1987). Reclassification of *Phialocephala* based on conidial development. *Transactions of the British Mycological Society* **89**, 509-520.

Wingfield, M.J., Capretti, P. and Mackenzie, M. (1988). *Leptographium* spp. as root

pathogens on conifers. An international perspective. In: *Leptographium* root diseases on conifers. (eds: T.C. Harrington & F.W. Cobb. Jr.), American Phytopathological Society, St. Paul, Minnesota, pp. 113-128.

Wingfield, M.J. and Gibbs, J.N. (1991). *Leptographium* and *Graphium* species associated with pine-infesting bark beetles in England. *Mycological Research* **95**, 1257-1260.

Wingfield, M.J., Seifert, K.A. and Webber, J.F. (1993). *Ceratocystis* and *Ophiostoma*. Taxonomy, Ecology and Pathogenicity. (ed. M.J. Wingfield. K.A, Seifert and J.F. Webber). American Phytopathological Press, St. Paul, Minnesota, pp. 293.

Wingfield, M. J., Crous, P.W. and Tzean, S.S. (1994). *Leptographium elegans*: a new species from Japan. *Mycological Research* **98**, 781-785.

Wingfield, M.J., Harrington, T.C. and Crous, P.W. (1994). Three new *Leptographium* species associated with conifer roots in the United States. *Canadian Journal of Botany* **72**, 227-238.

Wingfield, M.J., Harrington, T.C. and Solheim, H. (1995). Do conifer bark beetles require fungi to kill trees ? (ed. Christiansen, E), Proceedings from a symposium held at the Norwegian Forest Research Institute (NISK), Ås, Norway, 31 July 1995 – 2 August, 1995. *Aktuelt fra Skogforsk* 6/95, pp. 6

Wingfield, M.J., De Beer, C., Visser, C. and Wingfield, B.D. (1996). A new *Ceratocystis* species defined using morphological and ribosomal DNA sequence comparisons. *Systematics and Applied Microbiology* **19**, 191-202.

Witcosky, J.J. (1981). Insects associated with black-stain root disease of Douglas-fir in Western Oregon. M. Sc. thesis. Oregon State University. 51 pp.

Witcosky, J.J. (1989). Root beetles, stand disturbance, and management of black-stain root disease in plantations of Douglas-fir. In: *Insects affecting reforestation: Biology and Damage*. (eds. R.I. Alfaro & S.G. Glover). Forestry Canada, pp. 256.

Witcosky, J.J. and Hansen, E.M. (1985). Root-colonizing insects recovered from Douglas-fir in various stages of decline due to Black-stain root disease. *Phytopathology* **75**, 399-402.

Witcosky, J.J., Schowalter, T.D. and Hansen, E.M. (1986). *Hylastes nigrinus* (Coleoptera: Scolytidae), *Pissodes fasciatus*, and *Steremnius carinatus* (Coleoptera: Curculionidae) as vectors of black-stain root disease of Douglas-fir. *Environmental Entomology* **15**, 1090-1095.

Witthuhn, R.C., Wingfield, B.D., Wingfield, M.J. and Harrington, T.C. (1997). Comparison of three varieties of *Leptographium wageneri* using Random Amplified Polymorphic DNA. *South African Journal of Botany* **63**, 198-200.

Whitney, H.S. and Farris, S.H. (1970). Maxillary mycangium in the mountain pine beetle. *Science* **167**, 54-55.

Wolfaardt, J.F., Wingfield, M.J. and Kendrick, W.B. (1992). Synoptic key and computer database for identification of species of *Ceratocystis sensu lato*. *South African Journal of Botany* **58**, 277-285.

Wright, E.F. and Cain, R.F. (1961). New species of the genus *Ceratocystis*. *Canadian Journal of Botany* **39**, 1215-1230.

Yamaoka, Y., Wingfield, M.J., Takahashi, I. and Solheim, H. (1997). Ophiostomatoid fungi associated with the spruce bark beetle *Ips typographus japonicus* in Japan. *Mycological Research* **101**, 1215-1227.

Yamaoka, Y., Wingfield, M.J., Ohsawa, M. and Kuroda, Y. (1998). Ophiostomatoid fungi associated with *Ips cembrae* in Japan and pathogenicity to Japanese larch. *Mycoscience* **39**, 367-378.

Zambino, P.J. and Harrington, T.C. (1987). Geographic distributions of genotypes of the conifer root pathogen, *Verticicladiella wageneri*. *Annual Meetings of the Mycological Society of America and the Canadian Phytopathological Society*, p. 145.

Zambino, P.J. and Harrington, T.C. (1989). Isozyme variation within and among host-specialized varieties of *Leptographium wageneri*. *Mycologia* **81**, 122-133.

Zambino, P.J. and Harrington, T.C. (1990). Heterokaryosis and vegetative compatibility in *Leptographium wageneri*. *Phytopathology* **80**, 1460-1469.

Zambino, P.J. and Harrington, T.C. (1992). Correspondence of isozyme characterization with morphology in the asexual genus *Leptographium* and taxonomic implications. *Mycologia* **84**, 12-25.

Zambino, E., Harrington, T.C. and O'Malley, D. (1987). Isozyme analysis of three variants of the conifer root pathogen *Verticicladiella wageneri*. *Phytopathology* **77**, 124.

Zhou, X.D., Jacobs, K., Wingfield, M.J and Morelet, M. (1999). A new *Leptographium* species associated with *Tomicus piniperda* in South Western China. *Mycoscience* (in press).

Ye, H. (1991). On the bionomy of *Tomicus piniperda* (L.) (Col. Scolytidae) in the Kunming region of China. *Journal of Applied Entomology* **112**, 366-369.

Ye, H. and Dang, C. L. (1986) (in Chinese). Study on the feature of the pine shoot beetle injuring Yunnan pine. *J. Yunnan Univ.* **8**, 218-222.