

RESEARCH ARTICLE

Open Access

Evaluation of the inhibition of carbohydrate hydrolysing enzymes, antioxidant activity and polyphenolic content of extracts of ten African *Ficus* species (Moraceae) used traditionally to treat diabetes

Oyinlola O Olaokun^{1,3}, Lyndy J McGaw¹, Jacobus N Eloff¹ and Vinny Naidoo^{1,2*}

Abstract

Background: Some *Ficus* species have been used in traditional African medicine in the treatment of diabetes. The antidiabetic potential of certain species has been confirmed *in vivo* but the mechanism of activity remains uncertain. The aim of this study was to determine the activity and to investigate the mechanism of antidiabetic activity of ten selected *Ficus* species through inhibition of α -amylase and α -glucosidase activity, and the possible relationship between these activities, the total polyphenolic content and the antioxidant activity.

Methods: Dried acetone leaf extracts were reconstituted with appropriate solvents and used to determine total polyphenolic content antioxidant activity, α -amylase and α -glucosidase inhibitory activity.

Results: The crude acetone extract of *F. lutea* had the highest polyphenolic content (56.85 ± 1.82 mg GAE/g of dry material) and the strongest antioxidant activity with a TEAC value of 4.80 ± 0.90 . The antioxidant activity of the acetone extracts of the *Ficus* species may not be ascribed to total polyphenolic content alone. The crude extract at a concentration of 0.5 mg/ml of *F. lutea* ($64.3 \pm 3.6\%$) had the best α -glucosidase (sucrase) inhibitory activity. The EC₅₀ of *F. lutea* (290 ± 111 μ g/ml) was not significantly different from that of *F. sycomorus* (217 ± 69 μ g/ml). The α -amylase inhibitory activity of *F. lutea* ($95.4 \pm 1.2\%$) at a concentration of 1 mg/ml was the highest among the *Ficus* species screened. The EC₅₀ for *F. lutea* (9.42 ± 2.01 μ g/ml), though the highest, was not significantly different ($p < 0.05$) from that of *F. craterostoma* and *F. natalensis*. It was apparent that the crude acetone extract of *F. lutea* is a partially non-competitive inhibitor of α -amylase and α -glucosidase. Based on correlation coefficients polyphenolics may be responsible for α -glucosidase activity but probably not for α -amylase activity.

Conclusion: Antidiabetic activity potential via inhibition of α -amylase and α -glucosidase was discovered in *Ficus lutea* which has not been previously reported. The acetone extract of the leaves was high in total polyphenolic content and antioxidant activity, and was a potent inhibitor of α -amylase activity. Research is underway to isolate the active compound(s) responsible for the antidiabetic activity and to confirm the *in vitro* antidiabetic activity and to investigate *in vitro* toxicity.

Keywords: African *Ficus* species, *Ficus lutea*, Diabetes, α -amylase inhibitory activity, α -glucosidase inhibitory activity, Antioxidant activity, Polyphenol content

* Correspondence: vinny.naidoo@up.ac.za

¹Phytomedicine Programme, Department of Paraclinical Sciences, University of Pretoria, Private Bag X04, Onderstepoort 0110, South Africa

²Biomedical Research Centre, Faculty of Veterinary Sciences, University of Pretoria, Onderstepoort 0110, South Africa

Full list of author information is available at the end of the article

Background

Diabetes mellitus is an endocrine disease characterised by chronic hyperglycaemia with the disturbance of carbohydrate, fat and protein metabolism resulting from defects in insulin secretion, insulin action or both [1], and is typically associated with the failure of pancreatic β -cells. There are two major types of diabetes mellitus - type 1 and type 2. In type 1 diabetes, or insulin dependent diabetes mellitus, the body has little or no insulin secretory capacity and depends on exogenous insulin to prevent metabolic disorders and death. In type 2 diabetes, a non-insulin dependent diabetes mellitus, the body retains some endogenous insulin secretory capability; however, their insulin levels are low relative to their blood glucose levels and/or there is a measure of insulin resistance. Type 2 diabetes is the most prevalent form of the disease, accounting for 90-95% of cases [2,3].

Obesity, a sedentary lifestyle and a diet high in carbohydrate and fats are known risk factors for type 2 diabetes [4]. Globally, 171 million people were diagnosed with the disease in 2000, and it has been predicted that the prevalence will grow to 366 million by 2030 if no intervention action is taken [5]. In sub-Saharan Africa the disease is also considered to be an important emerging disease partly due to changes in diet and lifestyle of the people (a highly refined diet coupled to reduced physical activity). This is brought about by westernization, urbanization and mechanization, i.e. the disease which was once associated with only the affluent has increased in prevalence from 1% to 20% while in South Africa it is 4% to 6% [3,6,7].

The prolonged hyperglycaemia that results in diabetic patients has been speculated to induce oxidative stress through the excessive generation of free radicals which may impair endogenous antioxidant defence [8]. The stabilization of blood glucose in diabetic patients is therefore important to prevent these hyperglycaemic complications [9]. The main therapeutic approach used to achieve this objective in the diabetic patient is to stimulate insulin release, increase the number of glucose transporters, inhibit gluconeogenesis or reduce the absorption of glucose [10]. Another therapeutic approach is to decrease the post-prandial hyperglycaemia [11], which is achieved with the enzyme inhibitors such as acarbose, voglibose and miglitol [12], which function by retarding the action of the gastrointestinal carbohydrate hydrolysing enzymes α -amylase and α -glucosidase. As a result, these substances delay carbohydrate digestion thereby decreasing the rate of glucose absorption, i.e. they blunt the post-prandial plasma glucose rise [13]. While effective, the gastrointestinal side effects of acarbose, which includes bloating, abdominal discomfort, diarrhoea, and flatulence make them less attractive as therapeutic agents. New anti-diabetic compounds which

function by this mechanism but devoid of side effect are therefore desirable [12,14].

Ficus is a genus of about 800 species of woody trees, shrubs and vines in the family Moraceae. They are found in all tropical habitat types, with about 100 species occurring in Africa and the surrounding islands [15]. Several species of the genus *Ficus* are used traditionally in a wide variety of ethnomedical remedies all over the world [16,17]. They have long been used in folk medicine as antidiabetic, anthelmintic, hypotensive, mild laxative, anti-rheumatic, digestive and anti-dysentery drugs [18,19]. From previous studies, these plants are known to have chemical constituents such as triterpenes, sterols, polyphenols, flavonoids, coumarins, alkaloids and other metabolites [20]. Polyphenolics are among the naturally occurring antidiabetic agents [21], which may function via various biological effects, of which one is the inhibition of hydrolysing enzymes [22]. Polyphenolic compounds are also one of the major constituents of medicinal compounds which act as free radical scavengers and antioxidants. Free radicals can react with biological molecules, leading to cell and tissue injuries and pathological events. Therefore the discovery of polyphenolic compounds with a potential to inhibit the activity of digestive enzymes and having excellent antioxidant activity with low adverse effects is important for the treatment of diabetes.

Twelve *Ficus* species have antidiabetic activity with glucose lowering activity in alloxan or streptozotocin induced diabetic laboratory animals. These are *Ficus benghalensis* L. [23], *Ficus carica* L. [24], *Ficus racemosa* L. [25], *Ficus hispida* L.f. [26], *Ficus microcarpa* L.f. [27], *Ficus religiosa* L. [28], *Ficus thonningii* Blume [29], *Ficus glumosa* Delile. [30] *Ficus arnottiana* (Miq.)Miq. [31], *Ficus glomerata* Roxb. [32], *F. sycomorus* L. [33] and *F. deltoidea* Jack [34]. Several *Ficus* species are also used traditionally to treat diabetes and other ailments (Table 1). Four of the *Ficus* species used in the current study have *in vivo* blood glucose lowering potential (Table 2). With the mechanism(s) of action being unknown, we investigate the α -amylase and α -glucosidase enzymes inhibitory activity of the acetone extracts of the leaves of the ten *Ficus* species (Table 1). Acetone was selected as extractant because it dissolves many hydrophilic and lipophilic components from plants [48], it is volatile and has low toxicity for use in bioassays [49]. Furthermore acetone does not extract sugars, which if present (as would be the case with water and alcoholic extracts) would complicate α -amylase and α -glucosidase inhibitory assays [50].

Methods

Materials and chemicals

Dimethyl sulphoxide (DMSO), acetone, methanol and Whatman No. 1 filter paper were purchased from Merck

Table 1 Previous reports on the traditional uses of the selected *Ficus* species **F. religiosa* is a naturalized exotic from China

| Species | Section | Sub-section | Parts used | Uses | Reference |
|---|-------------|--------------|----------------------------|--|------------|
| <i>F. capreifolia</i> Delile (sandpaper fig) | sycidium | - | Leaf, root | Infections, vulnerary schistosomiasis | [35] |
| <i>F. cordata</i> Thunb. (Namaqua rock fig) | urostigma | urostigma | Bark stem | Treatment of diarrhoea, tuberculosis | [36] |
| <i>F. craterostoma</i> Mildbr. & Burret. (forest fig) | galoglychia | chlamydodora | leaves | Treatment of stomach ache | [37] |
| <i>F. glumosa</i> Delile (hair rock fig) | galoglychia | platyphyllae | Leaves, stem/bark | Treatment of skin diseases, diabetes, | [30,38] |
| <i>F. lutea</i> Vahl. (giant-leaves fig) | galoglychia | galoglychia | Leaves, bark | Treatment of sores, boils, madness, rabies | [39,40] |
| <i>F. natalensis</i> Hochst. (coastal strangler fig) | galoglychia | chlamydodora | Leaf, fruits, bark | Treatment of skin infections, colic, venereal diseases | [18,38,41] |
| <i>F. polita</i> Vahl. (heart-leaved fig) | galoglychia | caulocarpae | Leaves, fruits, bark, root | Dyspepsia, infectious diseases, abdominal pain, diarrhoea | [42] |
| <i>F. religiosa</i> L. (bo tree or sacred fig) | *- | *- | Leaves, bark/stem | Treatment of ulcers, bacterial infections, asthma, diabetes | [43,44] |
| <i>F. sycomorus</i> L. (sycamore fig) | sycomorus | sycomorus | Stem/bark, leaves, root, | Treatment of tumours, inflammation, mental madness, pain, bacterial infections, diabetes | [45-47] |
| <i>F. thonningii</i> Blume (bark-cloth fig) | galoglychia | chlamydodora | Leaves, fruits | Bronchitis, urinary tract infection, epilepsy, diabetes | [41,47] |

(South Africa). Folin-Ciocalteu reagent, gallic acid, sodium carbonate (Na₂CO₃), potassium persulphate, ABTS [2, 2-azinobis-(3- ethylbenzothiazoline-6-sulfonic acid)], potato starch, porcine pancreatic α- amylase enzyme (type VI), sodium chloride, sodium phosphate, 3, 5-dinitrosalicylic acid (DNS), sodium potassium tartrate, sodium hydroxide, acarbose, rat intestinal acetone powder, potassium phosphate, sucrose, TRIS (hydroxymethyl) aminomethane, hydrogen chloride (HCl), the glucose oxidase kit (GAGO 20) and Trolox (a synthetic water soluble vitamin E analogue) were sourced from Sigma (South Africa). The absorbance measurements were read using a microtitre plate reader (Versamax, Molecular Devices).

Plant material

The leaves of ten *Ficus* species were collected at the Manie van der Schijff Botanical Garden (University of Pretoria), South Africa in February 2009, and voucher specimens were conserved in the HGWJ Schweikerdt Herbarium of the University of Pretoria. The names of

plant species and voucher numbers are as follow: *Ficus capreifolia* Delile - PRU 33124; *Ficus cordata* Thunb - PRU 35501; *Ficus craterostoma* Warb. ex Mildbr. & Burret.- PRU 48293; *Ficus glumosa* Delile - PRU 38554, *Ficus lutea* Vahl.- PRU 074568; *Ficus natalensis* Hochst - PRU 078082; *Ficus polita* Vahl - PRU 35945; *Ficus religiosa* L. - PRU 115415; *Ficus sycomorus* L. - PRU 066173; and *Ficus thonningii* Blume - PRU 57036. The leaves were separated from the stems and dried at room temperature. The dried plant materials were milled to a fine powder in a Macsalab mill (Model 200 LAB, Eriez® Bramley) and stored at room temperature in closed glass bottles in the dark until extracted.

Preparation of plant extracts

The powdered plant material (2 g) was extracted with acetone (20 ml) in polyester centrifuge tubes using a shaker (Labotec model 20.2) at room temperature for 30 min. The extracts were centrifuged and filtered through Whatman No. 1 filter paper. This procedure was repeated three times on the same plant material to ensure that all possible compounds were extracted. The filtered extracts of each species were transferred into pre-weighed glass vials and solvent was left to evaporate at room temperature under a stream of cold air. The percentage yield was expressed on the air-dried mass of the leaves. The dried crude acetone extracts obtained were dissolved in appropriate solvent and used for analyses of total polyphenolics, antioxidant activity and *in vitro* α-amylase and α-glucosidase inhibition.

Table 2 *Ficus* species demonstrated to have blood glucose lowering potential *in vivo*

| Species | Part used | Solvent for extraction | Effective dose (mg/kg) <i>in vivo</i> | References |
|-------------------------|-----------|------------------------|---------------------------------------|------------|
| <i>Ficus glumosa</i> | stem/bark | methanol | 65 | [30] |
| <i>Ficus religiosa</i> | bark | water | 50 and 100 | [28] |
| <i>Ficus sycomorus</i> | Stem/bark | methanol | 250 | [33] |
| <i>Ficus thonningii</i> | stem/bark | ethanol | 120 and 240 | [29] |

Total polyphenolic content

The amount of total polyphenolics was determined using the method described by Djeridane et al. [51]. This method depends on the reduction of Folin-Ciocalteu reagent (a 2 N phosphomolybdic-phosphotungstic reagent) by phenols to a mixture of blue oxides. The commercial reagent was diluted 10-fold with double-distilled water. A standard curve was developed using gallic acid. Different concentrations of gallic acid were prepared in 80% methanol in a 96-well microtitre plate, and their absorbance immediately recorded at 760 nm. The crude acetone extracts were dissolved in 80% methanol to a final concentration of 1 mg/ml. The plant extract (100 μ l of the 1 mg/ml) was dissolved in 500 μ l (1/10 dilution) of the Folin-Ciocalteu reagent. After the addition of 1000 μ l of distilled water the mixture was allowed to stand for 1 min at room temperature, where after 1500 μ l of a 20% Na₂CO₃ solution was added. The final mixture was shaken, followed by an incubation period of 1 h in the dark at room temperature. The absorbance of all samples was measured at 760 nm. All analyses were performed in triplicate and repeated three times. The results are expressed in mg of gallic acid equivalent (GAE) per gram of dry weight of plant material.

Trolox equivalent anti-oxidant capacity (TEAC)

The total antioxidant activity was estimated using the Trolox equivalent capacity (TEAC) test [52]. In this assay, the relative capacity of antioxidant to scavenge the ABTS⁺ [2, 2-azinobis- (3-ethylbenzothiazoline-6-sulfonic acid)] radical compared to the antioxidant potency of Trolox (6-hydroxyl-2,5,7,8-tetramethylchromane-2-carboxylic acid), a synthetic water soluble vitamin E analogue, was measured.

The ABTS⁺ radical was produced by reacting 7 mM ABTS aqueous solution with 2.4 mM potassium persulphate in the dark for 12–16 h at room temperature. Prior to assay, the pre-formed ABTS⁺ radical was diluted in methanol (about 1:89 v/v) and equilibrated to give an initial absorbance (A_i) of 0.700 (\pm 0.02) at 734 nm. The crude acetone extracts were made up in methanol to a concentration of 0.0625 mg/ml to 1 mg/ml and the Trolox was made up to a concentration of 0.0625 mg/ml to 0.5 mg/ml in methanol. The crude acetone extract or Trolox standard (20 μ l) was added to the ABTS⁺ radical (180 μ l) and the mixture allowed to react for 6 minutes, and the absorbance (A) read at 734 nm in a spectrophotometer at one-minute intervals. Appropriate solvent blanks were included in each assay. The percentage change in absorbance was calculated for each concentration of extracts and Trolox. To calculate the TEAC, the gradient of the plot of the percentage inhibition of absorbance versus concentration plot for the extract under investigation is divided by the gradient of the plot for Trolox [52]. As a result, the TEAC of a substance is a ratio value

and has no unit. All assays were carried out in triplicate and repeated three times. The percentage change in absorbance (A) was measured as follows;

$$\% \text{ change in A} = \left(\frac{A_i \text{ ABTS}^+(734 \text{ nm}) - \text{New mean A ABTS}^+}{A_i \text{ ABTS}^+(734 \text{ nm})} \right) \times 100$$

α -Amylase inhibition assay

An adapted α -amylase inhibition assay as described by Ali et al. [50] was utilised. The dried crude acetone extracts were re-dissolved in DMSO to a concentration of 20 mg/ml and used for the α -amylase inhibition assay. Ice cold porcine pancreatic α -amylase solution (200 μ l) at 4 unit/ml (type VI) was pre-incubated with 40 μ l of crude acetone extracts and 160 μ l of distilled water, and mixed in a screw-top plastic tube. The reaction was started by the addition of 400 μ l of potato starch (0.5%, w/v) in 20 mM phosphate buffer (pH 6.9) containing 6.7 mM sodium chloride, and thereafter incubated at 25°C for 3 min. Final concentrations in the incubation mixture were plant extract (1 mg/ml), 0.25% (w/v) starch and 1 unit/ml enzyme. An aliquot of the mixture (200 μ l) was removed and placed into a separate tube containing 100 μ l DNS colour reagent solution (96 mM 3, 5-dinitrosalicylic acid, 5.31 M sodium potassium tartrate in 2 M NaOH) and placed into an 85°C water bath. After 15 min, this mixture was removed from the water bath, cooled and diluted with 900 μ l distilled water. α -Amylase activity was determined by measuring the remaining starch content by measuring the absorbance of the mixture at 540 nm. Control incubations, representing 100% enzyme activity were conducted by replacing the plant extract with DMSO (40 μ l). For the blanks (negative controls) the enzyme solution was replaced with distilled water and the same procedure was carried out as above. Acarbose was used as positive control (acarbose 1 mg/ml; α -amylase 1 unit/ml). The α -amylase inhibition activity was expressed as;

$$\% \text{ Inhibition} = 100 \times \left(\frac{\Delta A_{\text{Control}} - \Delta A_{\text{Sample}}}{\Delta A_{\text{Control}}} \right)$$
$$\Delta A_{\text{Control}} = A_{\text{Test}} - A_{\text{Blank}}$$
$$\Delta A_{\text{Sample}} = A_{\text{Test}} - A_{\text{Blank}}$$

α -Glucosidase inhibition assay

A commercial rat intestine acetone powder was purchased and used to prepare enzyme solution according to the method of Oki et al. [53] with slight modifications. In brief, rat intestinal acetone powder (25 mg/ml) was homogenized in ice-cold 50 mM phosphate buffer. After centrifugation at 6 000 \times g for 15 min, the clear supernatant obtained was used as the enzyme solution for determining α -glucosidase activity.

The rat intestinal sucrase inhibitory activity was determined using the method of Bhandari et al. [54], with a slight modification. Sucrose (200 µl of a 56 mM solution) in 0.1 M potassium phosphate buffer (pH 7) was mixed with 100 µl of crude acetone extracts (2.5 mg/ml) in 50% aqueous dimethyl sulphoxide (DMSO) in a test tube. After pre-incubation at 37°C for 5 min, 200 µl of rat intestinal α-glucosidase solution prepared as described above was added. Whereas 100 µl DMSO (50%) was used in place of the plant extract to represent 100% enzyme activity, Acarbose (0.1 mg/ml) was used as the positive control and for the blanks (negative controls) the enzyme solution was replaced with distilled water and the same procedure was carried out as above. Final concentrations in the incubation mixture for plant extracts were 0.5 mg/ml and 0.02 mg/ml for acarbose.

After thoroughly mixing, the test sample, solvent and positive control test tubes were incubated at 37°C for 20 min and then the reaction was stopped by adding 750 µl of 2 M Tris-HCl buffer (pH 6.9). The amount of liberated glucose was determined by the glucose oxidase method using a commercial test kit (Sigma GAGO 20) according to the manufacturer's instructions. The absorbance of the wells was measured at 540 nm and the inhibitory activity was calculated using the following formula:

$$\% \text{ Inhibition} = 100 \times \left(\frac{\Delta A_{\text{Control}} - \Delta A_{\text{Sample}}}{\Delta A_{\text{Control}}} \right)$$

$$\Delta A_{\text{Control}} = A_{\text{Test}} - A_{\text{Blank}}$$

$$\Delta A_{\text{Sample}} = A_{\text{Test}} - A_{\text{Blank}}$$

Kinetics of inhibition against α- amylase and α- glucosidase activities

The kinetics of inhibition of the crude acetone extract of *F. lutea* against α- amylase and α- glucosidase activities were measured by increasing substrate concentrations of starch (0.016% - 0.5%) and sucrose (7 mM - 56 mM) respectively in the absence and presence of extract of *F. lutea* at concentrations of 25-100 µg/ml for α-amylase inhibitory assay and 62.5 - 250 µg/ml for α- glucosidase inhibitory assay. The type of inhibition was determined by Lineweaver-Burk double reciprocal plot analysis of the data, which was calculated from the result according to Michaelis-Menten kinetics [11].

Calculation of EC50

A dose response effect was conducted on α- amylase and α-glucosidase for the extracts. The concentrations of extracts used ranged from 16 to 500 µg/ml. The concentration of extract that inhibited 50% of the enzyme activity under assay conditions was defined as the effective concentration (EC50) and this was established using

the programme Kinetica 5 (Thermo). The concentration response relationship was plotted and fitted to a Hill's model. The model was described by the following equation; $y = \frac{y_{\text{max}}x^n}{c^n + x^n}$. The EC50 concentrations attained were used for comparison of the potency of the various crude acetone plant extracts.

Statistical analysis

All experiments were performed in triplicate and repeated three times to yield nine dose- response curves. Data are presented as the mean ± standard error of mean (S.E.M.). Statistical significances were evaluated by one-way analysis of variance (ANOVA) and student's *t*-test, with *post hoc* analysis using the Bonferroni corrected *t*-test. *P*-values less than 0.05 were considered to be significant.

Results

Extraction of plant material

Ten *Ficus* species were selected on the basis of availability and were extracted with acetone. In Table 3, the percentage yields of crude acetone extracts of the ten *Ficus* species are reported. The percentage yield ranged from 2.32% to 3.70% among the leaf extracts where the extract of *F. lutea* had the highest yield (3.70%), followed by *F. polita* (3.15%). The extracts of *F. natalensis* (2.35%) and *F. capreifolia* had the lowest yields (2.32%).

Total polyphenol content and antioxidant activity

The total polyphenol content of the crude acetone extracts of the ten *Ficus* species has highly variable (Table 3), with a range of 4.64 to 56.85 mg GAE/g dry weight of plant. When the total polyphenol content of

Table 3 Percentage yield, total polyphenol content and antioxidant activity of crude acetone extracts of leaves of ten *Ficus* species

| Species | Yield (%) | ^a Total polyphenol (mg GAE/g dry weight) | ^{ab} Antioxidant activity TEAC |
|---------------------------|-----------|---|---|
| <i>Ficus capreifolia</i> | 2.32 | 4.73 ± 0.26 ^c | 0.34 ± 0.05 ^c |
| <i>Ficus cordata</i> | 2.65 | 8.23 ± 1.00 ^d | 0.27 ± 0.03 ^c |
| <i>Ficus craterostoma</i> | 2.60 | 9.80 ± 0.93 ^d | 0.66 ± 0.06 ^d |
| <i>Ficus glumosa</i> | 2.60 | 19.24 ± 0.79 ^e | 1.29 ± 0.30 ^e |
| <i>Ficus lutea</i> | 3.70 | 56.85 ± 1.82 ^f | 4.80 ± 0.90 ^f |
| <i>Ficus natalensis</i> | 2.35 | 4.75 ± 0.92 ^c | 0.69 ± 0.08 ^d |
| <i>Ficus polita</i> | 3.15 | 8.04 ± 0.52 ^d | 0.31 ± 0.06 ^c |
| <i>Ficus religiosa</i> | 2.45 | 5.40 ± 0.35 ^c | 0.59 ± 0.18 ^c |
| <i>Ficus sycamoros</i> | 2.60 | 12.33 ± 0.26 ^e | 1.91 ± 0.19 ^e |
| <i>Ficus thonningii</i> | 2.40 | 4.64 ± 0.48 ^c | 0.77 ± 0.06 ^d |

^aValues are means of triplicate determinations done three times (n = 9) ± standard error;

^bAntioxidant activity (Trolox equivalent antioxidant capacity).

^{c,d,e,f}No significant difference between extracts with same value, but significant difference *p*<0.05 between different values.

each extract was compared, the extract of *F. lutea* (56.85 ± 1.82 mg/g) was found to have a remarkably higher content ($p < 0.001$) followed in decreasing order by extracts of *F. glumosa* and *F. sycomorus* with total polyphenol content of 19.24 ± 0.79 and 12.33 ± 0.26 mg GEA/g dry weight of plant respectively. The extracts with the lowest values in decreasing order were *F. natalensis*, *F. capreifolia* and *F. thonningii* with total phenolic content of 4.75 ± 0.92, 4.73 ± 0.26 and 4.64 ± 0.28 mg GAE/g dry weight of plant respectively.

The antioxidant activities of the extracts, expressed as Trolox equivalent antioxidant capacity (TEAC) are listed in Table 3. The crude acetone extracts of the ten *Ficus* species had different antioxidant capacities (Table 3). The total antioxidant activity for the ten *Ficus* species varied widely from 4.80 ± 0.90 to 0.27 ± 0.03 TEAC under the assay conditions. The antioxidant activity of the extract of *F. lutea* (4.80 ± 0.90 TEAC) was significantly different ($p < 0.001$) when compared with that of the other extracts, followed by *F. sycomorus* and *F. glumosa* in decreasing order with antioxidant activity of 1.91 ± 0.18 and 1.29 ± 0.30 TEAC respectively. The extracts with the lowest TEAC in decreasing order were *F. capreifolia*, *F. polita* and *F. cordata* with 0.34 ± 0.05, 0.31 ± 0.06 and 0.27 ± 0.03 TEAC respectively. The correlation coefficient between total polyphenolic content and TEAC of the crude acetone extracts of the ten *Ficus* species R^2 was 0.50.

α-Amylase inhibitory activity of extracts of *Ficus* species

The inhibitory activity of the crude acetone extracts of the ten *Ficus* species are presented in Table 4. The extract of *F. lutea* had the highest inhibitory potential with activity of 95.4 ± 1.2% inhibition at 1 mg/ml. This was followed by the extract of *F. glumosa* with 65.1 ± 3.0% inhibition at the same concentration while Acarbose, the positive control, had inhibitory activity of 96.7 ± 0.3% at 0.04 mg/ml. The extracts of the other *Ficus* species moderately inhibited α-amylase with activity between 40% and 45%, with the exception of the extracts of *F. religiosa* and *F. thonningii* which had poor inhibitory activities. The EC50s are presented in Table 4. The extract of *F. lutea* was most potent (EC50 of 9.42 ± 2.01 µg/ml) followed by *F. craterostoma* (EC50 = 11.41 ± 4.68 µg/ml) and *F. natalensis* (EC50 = 17.85 ± 4.42 µg/ml) with no significant difference present between them ($p < 0.05$). Acarbose was characterised by an EC50 of 0.04 ± 0.03 µg/ml. The correlation coefficient between total polyphenolic content and inhibition of α-amylase activity by the extracts of the ten *Ficus* species R^2 was 0.65. Conversely, the correlation coefficient between antioxidant activity (TEAC) and inhibition of α-amylase activity by the crude acetone extracts of the ten *Ficus* species R^2 was 0.41.

Table 4 The percentage inhibition of α-amylase activity (1 mg/ml) and concentration leading to 50% inhibition (EC50) of crude acetone extracts of the ten selected *Ficus* species

| Species | (%) α- amylase inhibition ^a | EC50 (µg/ml) |
|---------------------------|--|---------------------------|
| <i>Ficus capreifolia</i> | 43.8 ± 3.3 ^b | >100 |
| <i>Ficus cordata</i> | 45.9 ± 5.2 ^b | > 100 |
| <i>Ficus craterostoma</i> | 48.3 ± 1.6 ^b | 11.41 ± 4.68 ^b |
| <i>Ficus glumosa</i> | 65.1 ± 3.0 ^c | > 100 |
| <i>Ficus lutea</i> | 95.4 ± 1.2 ^d | 9.42 ± 2.01 ^b |
| <i>Ficus natalensis</i> | 43.7 ± 1.8 ^b | 17.85 ± 4.42 ^b |
| <i>Ficus polita</i> | 40.2 ± 2.6 ^b | > 100 |
| <i>Ficus religiosa</i> | 35.3 ± 2.8 ^b | > 100 |
| <i>Ficus sycomorus</i> | 40.0 ± 2.8 ^b | > 100 |
| <i>Ficus thonningii</i> | 37.6 ± 2.7 ^b | > 100 |
| Acarbose | 96.7 ± 0.3 | 0.04 ± 0.03 |

^a% α-amylase inhibitory activity of crude acetone extracts of ten South African *Ficus* species (1 mg/ml) and control (acarbose) (0.04 mg/ml). Results expressed as % mean ± SEM (n = 9). ^{b,c,d}No significant difference between extracts with same value, but significant difference $p < 0.05$ between different values.

α-Glucosidase inhibitory activity of extracts of *Ficus* species

The crude acetone extracts of the ten *Ficus* species were generally weak inhibitors of α- glucosidase activity (Table 5), with only *F. lutea* demonstrating inhibitory activity of 64.3 ± 3.6% at a concentration of 0.5 mg/ml (Significantly different ($p < 0.05$) to the other extracts). Acarbose (positive control) was a potent inhibitor of α-glucosidase (84.8 ± 1.7%) at a concentration of 0.02 mg/ml. The EC50s for the inhibitory activity is presented in

Table 5 The percentage inhibition of α-glucosidase activity (0.5 mg/ml) and concentration leading to 50% inhibition (EC50) of crude acetone extracts of ten *Ficus* species

| Species | (%) α-glucosidase inhibition ^a | EC50 (µg/ml) |
|---------------------------|---|------------------------|
| <i>Ficus capreifolia</i> | 24.3 ± 1.7 ^b | > 1000 |
| <i>Ficus cordata</i> | 22.0 ± 3.6 ^b | > 1000 |
| <i>Ficus craterostoma</i> | 28.2 ± 7.0 ^b | > 1000 |
| <i>Ficus glumosa</i> | 38.7 ± 6.5 ^b | > 1000 |
| <i>Ficus lutea</i> | 64.3 ± 3.6 ^c | 290 ± 111 ^b |
| <i>Ficus natalensis</i> | 23.6 ± 8.1 ^b | > 1000 |
| <i>Ficus polita</i> | 29.2 ± 5.8 ^b | > 1000 |
| <i>Ficus religiosa</i> | 17.6 ± 8.0 ^b | > 1000 |
| <i>Ficus sycomorus</i> | 35.7 ± 5.4 ^b | 217 ± 69 ^b |
| <i>Ficus thonningii</i> | 25.3 ± 5.0 ^b | > 1000 |
| Acarbose | 84.8 ± 1.7 | 3.4 ± 0.5 |


^a% α-glucosidase inhibitory activity of crude acetone extracts of ten South African *Ficus* species (0.5 mg/ml) and control (acarbose) (0.02 mg/ml). Results expressed as % mean ± SEM (n = 9).

^{b,c}No significant difference between extracts with same value, but significant difference $p < 0.05$ between different values.

Table 5. The extract of *F. sycomorus* was most potent with an EC₅₀ of 217 ± 69 µg/ml, followed closely by the extract of *F. lutea* (290 ± 111 µg/ml), with no significant difference between them ($p > 0.05$). The EC₅₀ of Acarbose was 3.4 ± 0.5 µg/ml. The correlation coefficient between total polyphenolic content and inhibition of α-glucosidase activity by the crude acetone extracts of the ten *Ficus* species R² was 0.74. In addition, the correlation coefficient between antioxidant activity (TEAC) and inhibition of α-glucosidase activity by the crude acetone extracts of the ten *Ficus* species R² was 0.67.

The enzyme kinetics of α- amylase and α- glucosidase inhibition by extract of *F. lutea*

Analysis of the α-amylase and α-glucosidase kinetics by the crude acetone extract of *F. lutea* is shown in Figure 1. For the α-amylase (Figure 1A) and α-glucosidase (Figure 1B) inhibition by *F. lutea*, the intersection of the double reciprocal plot is seated at a point above the +1/[s] axis, indicating that *F. lutea* acts as a partially non-competitive-type inhibitor of α-amylase and α- glucosidase.



Discussion

The most challenging goal in the management of type 2 diabetes mellitus is to achieve blood glucose levels as close to normal as physiologically possible [55]. With postprandial hyperglycaemia being the earliest metabolic abnormality detectable, modulation of this rise is an important tool in the management of the diabetic patient [56,57]. α-Amylases are endoglucanases, which hydrolyse the internal α-1,4 glucosidic linkages in starch, and α-glucosidase (sucrase), one of the glucosidases located in the brush border surface membrane of intestinal cells, is a key enzyme for carbohydrate digestion and absorption. These enzymes have been recognized as therapeutic targets for modulation of postprandial hyperglycaemia. In addition, prolonged hyperglycaemia is an independent risk factor for the development of microvascular and macrovascular complications of diabetes mellitus which is mediated through oxidative stress [55,58].

In this study the inhibitory effectiveness of the acetone extract of the leaves of ten *Ficus* species against the α-amylase and α-glucosidase enzymes were evaluated. As a secondary objective, the relationship between total polyphenolic content and antioxidant activity was also studied. For this study, a positive correlation between total polyphenolic content and inhibition of α-amylase and α-glucosidase activity was present with R² values of 0.65 and 0.74 respectively. This therefore indicates the likelihood of the polyphenols in the acetone extracts being partly responsible for the inhibition of the activity of the enzymes. The latter was not an unexpected finding as the polyphenolic extracts from a number of plants have been previously shown to be effective inhibitors of the intestinal α-glucosidase and α-amylase enzyme systems [59].

The extracts of ten *Ficus* species were evaluated for their potential to specifically inhibit activity of porcine pancreatic α-amylase and rat small intestinal α-glucosidase. The ten *Ficus* species were generally potent inhibitors of activity of the porcine pancreatic α-amylase but weak inhibitors of activity of the rat small intestinal α-glucosidase. Three of the four *Ficus* species with *in vivo* blood glucose lowering potential in previous studies (Table 2) were all weak inhibitors of the activity of α-amylase and α-glucosidase *in vitro*. While the reason for the difference could be due to the different compounds being extracted as a result of different solvents used in these studies, the more likely reason was that the glucose lowering potential of these plants may be due to non-α-amylase or α-glucosidase inhibitory activities, such as the stimulation of glucose uptake, the stimulation of insulin release or the inhibition of gluconeogenesis.

The fourth *Ficus* species (*F. sycomorus*) which is used traditionally by the Hausa and Fulani tribes of northern

Nigeria to treat diabetes [33] together with known glucose lowering potential *in vivo* (Table 2) was a weak inhibitor of α -amylase activity but a potent inhibitor of α -glucosidase (sucrase) activity with an EC₅₀ of 217 ± 69 μ g/ml. We therefore believe that the glucose lowering potential demonstrated *in vivo* by the extract of *F. sycomorus* may be due to the inhibition of α -glucosidase activity (sucrase). Some authors have shown that the extracts of other *Ficus* species such as *F. racemosa* [60], *F. benghalensis* [61] and *F. deltoidea* [62] inhibit the activity of sucrase with an EC₅₀ of 367 ± 15.2 μ g/ml and 239 ± 14.3 μ g/ml respectively for cold and hot water extract of *F. racemosa*, and 193 ± 21.6 μ g/ml and 141 ± 22.1 μ g/ml respectively for cold and hot water extract of *F. benghalensis*. This was in agreement with our results. Furthermore, studies also showed the activity of α -amylase is inhibited with an EC₅₀ of $0.94 \pm 0.15\%$ and $0.58 \pm 0.15\%$ respectively for the cold and hot water extracts of *F. racemosa*, [60] and, 4.4 and 125 μ g/ml respectively for the cold and hot water extracts of *F. benghalensis* [63], this was also in agreement with our results.

Some studies have observed the potency with which polyphenols inhibited the activity of porcine pancreatic α -amylase and rat small intestinal α -glucosidase to be different [59,64]. Strawberry and raspberry extracts were the most effective inhibitors of α -amylase followed by blueberry and blackcurrant. Although these extracts also inhibited rat intestinal α -glucosidase activity, the order of effectiveness was different than with α -amylase. Blueberry and blackcurrant were the most effective followed by strawberry and raspberry [59]. Rat intestinal α -glucosidase is generally weakly inhibited by many flavonoids, while flavonoids are often potent inhibitors of porcine pancreatic α -amylase [64]. It is possible that the extracts of the *Ficus* species used in this study demonstrated potent α -amylase inhibitory activity because they contain more of the flavonoid groups with the extract of *F. lutea* being the most potent.

Polyphenolic compounds may also be indirectly beneficial in disease by chelating metal ions [65] or activating the expression of antioxidant enzymes [66] or directly acting as antioxidants [67]. High total phenolic contents may not always translate to a high antioxidant activity. Pulido et al. [68] reported that antioxidant efficiency of the polyphenols seemed to depend on the position and extent of hydroxylation and conjugation. This was also observed in this study where the correlation between total polyphenolic content and antioxidant activity had an R² value of 0.5. We found that extracts of *F. glumosa*, *F. sycomorus* and *F. lutea* had good antioxidant activities and high total phenolic contents. Some authors have shown that *F. glumosa* [30], *F. sycomorus* [69] and *F. lutea* [39] have very good DPPH[•] scavenging activities with reported IC₅₀ of 79.5 ± 1.77 μ g/ml and 11.9 ± 0.3 μ g/ml

respectively for *F. sycomorus* and *F. lutea*, and this is in agreement with our results. Among these three *Ficus* species, the extract of *F. lutea* displayed the highest total polyphenolic content and the highest antioxidant activity.

In addition, the qualitative phytochemical analysis conducted (results not included) showed that tannin was one of the prominent phytochemicals contained in the crude acetone extract of *F. lutea*. Hagerman et al. [70] reported that the high molecular weight polyphenols (tannins) have the ability to quench free radicals (ABTS^{•+}) and their effectiveness depends on the molecular weight, the number of aromatic rings and the nature of the hydroxyl group substitutions. It is also possible that not all polyphenolic compounds possess ABTS^{•+} radical scavenging activities [71]. Free radical (ABTS^{•+}) scavenging of *Ficus* species may be due to the presence of high molecular weight polyphenols, such as catechins, pelargonidins and leucopelargonidin derivatives in addition to the flavonoids [72]. A study of the correlation between antioxidant activity and inhibition of α -amylase and α -glucosidase activity by the ten *Ficus* species had an R² value of 0.41 and 0.67 respectively indicating that the polyphenolic compounds responsible for antioxidant activity may likely be responsible for the inhibition of α -glucosidase activity but may not be responsible for the inhibition of α -amylase activity. Some authors have demonstrated that compounds with potent antioxidant activity are also strong inhibitors of α -glucosidase activity.

When studying the kinetics of inhibition, the extract of *F. lutea* showed partial non-competitive inhibition against porcine pancreatic α -amylase and α -glucosidase. A fully non-competitive inhibitor binds the enzyme substrate [ES] complex, not the free enzyme and affects the breakdown of the [ES] to form a product. Likewise, a partial non-competitive inhibitor binds to the enzyme-substrate complex but it is assumed that the inhibitor is released from the enzyme at the same time [ES] is broken down to the product [73]. This type of inhibition does occur where there are multiple inhibitors as will be the case for a crude extract of *F. lutea* [73]. Another study demonstrated that pure compounds such as D-xylose and (+)-catechin are un-competitive inhibitors of rat intestinal α -glucosidase activity [53].

The results of this study revealed that the crude acetone extract of *F. lutea* is high in total polyphenolic content and antioxidant activity, and is a potent inhibitor of α -amylase activity. This *in vitro* study has demonstrated that polyphenolic compounds present in *F. lutea* may likely be responsible for the inhibitory activity against the carbohydrate hydrolysing enzymes.

Conclusion

This study investigated the potential antidiabetic activity of ten South African *Ficus* species, focussing on effects on

digestive enzymes, polyphenolic content and antioxidant activity. All acetone extracts of the ten *Ficus* species had α -amylase and α -glucosidase enzyme inhibitory activity. The crude acetone extract of *F. lutea* was the most active in all the assays. This is the first report of the antidiabetic potential of *F. lutea* in potentially lowering blood glucose levels *in vitro*. The crude acetone extract of *F. lutea* was more active *in vitro* than the crude acetone extracts of the three *Ficus* species that are reported to be used traditionally to treat diabetes. The activity that is responsible for the antidiabetic activity appears to reside in the polar fraction of the extract of *F. lutea*, so rural people should be able to extract the active compound(s) with water or alcohol. Research is underway to isolate the active compound responsible for the antidiabetic activity, to confirm the *in vitro* antidiabetic activity, and to investigate toxicity and efficacy *in vivo*.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

OOO carried out the study and wrote the manuscript; VN and JNE contributed to conception, design, analysis and interpretation of data, and LJM, JNE and VN assisted with and supervised the manuscript writing. All authors have read and approved the final manuscript.

Acknowledgement

The National Research Foundation (NRF) of South Africa provided funding and leaves were collected from the Manie van der Schijff Botanical Garden at the University of Pretoria.

Author details

¹Phytomedicine Programme, Department of Paraclinical Sciences, University of Pretoria, Private Bag X04, Onderstepoort 0110, South Africa. ²Biomedical Research Centre, Faculty of Veterinary Sciences, University of Pretoria, Onderstepoort 0110, South Africa. ³Permanent address: Federal Institute of Industrial Research Oshodi (FIRO), Ikeja, Nigeria.

Received: 30 October 2012 Accepted: 18 April 2013

Published: 4 May 2013

References

1. Anonymous: Definition, diagnosis and classification of diabetes mellitus and its complications. In *Part 1: diagnosis and classification of diabetes mellitus*. Geneva: Report of a WHO Consultation; 1999.
2. Kaneto H, Katakami N, Kawamori D, Miyatsuka T, Sakamoto K, Matsuoka T-A, Matsuhisa M, Yamasaki Y: Involvement of oxidative stress in the pathogenesis of diabetes. *Antioxid Redox Signal* 2007, **9**:355–366.
3. Jung M, Park M, Lee HC, Kang YH, Kang ES, Kim SK: Antidiabetic agents from medicinal plants. *Curr Med Chem* 2006, **13**:1203–1218.
4. Jain S, Saraf S: Review on Type 2 diabetes mellitus—its global prevalence and therapeutic strategies. *Diabetes and Metabolic Syndrome: Clinical Research and Reviews* 2010, **4**:48–56.
5. Wild S, Roglic G, Green A, Sicree R, King H: Global prevalence of diabetes Estimates for the year 2000 and projections for 2030. *Diabetes Care* 2004, **27**:1047–1053.
6. Ogunbodede EO, Fatusi OA, Akintomide A, Kolawole K, Ajayi A: Oral health status in a population of Nigerian diabetics. *J Contemp Dent Pract* 2005, **6**:75–84.
7. Deuschländer MS, van de Venter M, Roux S, Louw J, Lall N: Hypoglycaemic activity of four plant extracts traditionally used in South Africa for diabetes. *J Ethnopharmacol* 2009, **124**:619–624.
8. Johansen JS, Harris AK, Rychly DJ, Ergul A: Oxidative stress and the use of antioxidants in diabetes: Linking basic science to clinical practice. *Cardiovasc Diabetol* 2005, **4**(1):5. doi:10.1186/1475-2840-4-5.
9. Mai TT, Chuyen NV: Anti-hyperglycemic activity of an aqueous extract from flower buds of *Cleistocalyx operculatus* (Roxb.) Merr and Perry. *Biosci Biotechnol Biochem* 2007, **71**:69–76.
10. Youn JY, Park HY, Cho KH: Anti-hyperglycemic activity of *Commelina communis* L: inhibition of alpha-glucosidase. *Diabetes Res Clin Pract* 2004, **66S**:S149–S155.
11. Kim YM, Jeong YK, Wang MH, Lee WY, Rhee HI: Inhibitory effect of pine extract on alpha-glucosidase activity and postprandial hyperglycemia. *Nutrition* 2005, **21**:756–761.
12. De Fronzo RA: Pharmacologic therapy for type 2 diabetes mellitus. *Ann Intern Med* 1999, **131**:281–303.
13. Rhabasa-Lhoret R, Chiasson JL: α -Glucosidase inhibitors. In *International Textbook of Diabetes Mellitus, vol. 1*. 3rd edition. Edited by DeFronzo RA, Ferrannini E, Keen H, Zimmet P. UK: John Wiley & Sons Ltd; 2004:901–914.
14. Moller DE: New drug targets for type 2 diabetes and the metabolic syndrome. *Nature* 2001, **414**:821–827.
15. Serrato A, Ibarra-Manriquez G, Oyama K: Biogeography and conservation of the genus *Ficus* (Moraceae) in Mexico. *J Biogeogr* 2004, **31**:475–485.
16. Hansson A, Zelada JC, Noriega HP: Re-evaluation of risks with the use of *Ficus insipida* latex as a traditional anthelmintic remedy in the Amazon. *J Ethnopharmacol* 2005, **98**:251–257.
17. Koné WM, Atindehou KK, Terreaux C, Hostettmann K, Traoré D, Dosso M: Traditional medicine in North Côte-D'Ivoire: Screening of 50 medicinal plants for antibacterial activity. *J Ethnopharmacol* 2004, **93**:43–49.
18. Watt JM, Breyer-Brandwijk MG: *Medicinal and poisonous plants of Southern and Eastern Africa*. London: Livingstone Ltd; 1962:773–780.
19. Trivedi P, Hind S, Sharma RC: Preliminary phytochemical and pharmacological studies on *Ficus racemosa*. *J Med Res* 1969, **56**:1070–1074.
20. Ramadan MA, Ahmad AS, Nafady AM, Mansour AI: Chemical composition of the stem bark and leaves of *Ficus pandurata* Hance. *Nat Prod Res* 2009, **23**:1218–1230.
21. Andrade-Cetto A, Becerra-Jiménez J, Cárdenas-Vázquez R: Alpha-glucosidase inhibiting activity of some Mexican plants used in the treatment of type 2 diabetes. *J Ethnopharmacol* 2008, **116**:27–32.
22. Mai TT, Thu NN, Tien PG, Van Chuyen N: Alpha-glucosidase inhibitory and antioxidant activities of Vietnamese edible plants and their relationships with polyphenol contents. *J Nutr Sci Vitaminol* 2007, **53**:267–276.
23. Singh RK, Mehta S, Jaiswal D, Rai PK, Watal G: Antidiabetic effect of *Ficus bengalensis* aerial roots in experimental animals. *J Ethnopharmacol* 2009, **123**:110–114.
24. Perez C, Dominguez E, Ramiro JM, Romero A, Campillo JE, Torres MD: A study on the glycaemic balance in streptozotocin-diabetic rats treated with an aqueous extract of *Ficus carica* (fig tree) leaves. *Phytother Res* 1996, **10**:82–83.
25. Rao RB, Murugesan T, Sinha S, Saha BP, Pal M, Mandal SC: Glucose lowering efficacy of *Ficus racemosa* bark extract in normal and alloxan diabetic rats. *Phytother Res* 2002, **16**:590–592.
26. Ghosh R, Sharatchandra KH, Rita S, Thokchom IS: Hypoglycaemic activity of *Ficus hispida* (bark) in normal and diabetic albino rats. *Indian J Pharmacol* 2004, **36**:222–225.
27. Kumar KA, Maheshwari MU, Shivashanmugam AT, Devi VS, Prasanth NV, Ravi TK: Hypoglycemic effect of *Ficus microcarpa* leaves (Chinese banyan) on alloxan-induced diabetic rats. *J Biol Sci* 2007, **7**(2):321–326.
28. Pandit R, Phahke A, Jagtap A: Antidiabetic effects of *Ficus religiosa* extract in streptozotocin-induced diabetic rats. *J Ethnopharmacol* 2010, **128**:462–466.
29. Musabayane CT, Gondwe M, Kamadyaapa DR, Chuturgoon AA, Ojewole JA: Effects of *Ficus thonningii* (Blume) [Moraceae] stem-bark ethanolic extract on blood glucose, cardiovascular and kidney functions of rats, and on kidney cell lines of the proximal (LLC-PK1) and distal tubules (MDBK). *Ren Fail* 2007, **29**:389–397.
30. Madubunyi II, Onoja SO, Asuzu IU: *In vitro* antioxidant and *in vivo* antidiabetic potential of the methanolic extract of *Ficus glumosa* Del (Moraceae) stem bark in alloxan-induced diabetic mice. *Comp Clin Pathol* 2012, **21**:389–394.
31. Mazumder PM, Farswan M, Parcha V: Hypoglycaemic effect of *Ficus arnottiana* Miq. bark extracts on streptozotocin induced diabetes in rats. *Natural Product Radiance* 2009, **8**:478–482.
32. Kar A, Choudhary BK, Bandyopadhyay NG: Comparative evaluation of hypoglycaemic activity of some Indian medicinal plants in alloxan diabetic rats. *J Ethnopharmacol* 2003, **84**:105–108.

33. Adoum OA, Michael BO, Mohammad IS: **Phytochemicals and hypoglycaemic effect of methanol stem-bark extract of *Ficus sycamoros* Linn (Moraceae) on alloxan induced diabetic Wistar albino rats.** *Afr J Biotechnol* 2012, **11**(17):4095–4097.
34. Adam Z, Khamis S, Ismail A, Hamid M: ***Ficus deltoidea*: a potential alternative medicine for diabetes mellitus.** *Evidence-Based Complementary and Alternative Medicine* 2012. doi:10.1155/2012/632763.
35. Brink M: ***Ficus capreifolia* Delile.** In *[Internet] Record from Protabase*. Edited by Brink M, Achigan-Dako EG. Wageningen, Netherlands: PROTA (Plant Resources of Tropical Africa / Ressources végétales de l'Afrique tropicale); 2010.
36. Kuete V, Ngameni B, Fotso Simo CC, Tankeu RK, Ngadjui BT, Meyer JJM, Lall N, Kuate JR: **Antimicrobial activity of the crude extract and compounds from *Ficus chlamydoparp* and *Ficus cordata* (Moraceae).** *J Ethnopharmacol* 2008, **120**:17–24.
37. Bhat RB, Jacobs TY: **Traditional herbal medicine in Transkei.** *J Ethnopharmacol* 1995, **48**:7–12.
38. Hutchings A, Scott AH, Lewis G, Cunningham AB: ***Zulu medicinal plants. An inventory.*** Pietermaritzburg: University of Natal Press; 1996:352.
39. Marwah RG, Fatope MO, Al Mahrooqi R, Varma GB, Al Abadi H, Al-Burtamani SKS: **Antioxidant capacity of some edible and wound healing plants in Oman.** *Food Chem* 2006, **101**:465–470.
40. Chifundera K: **Livestock diseases and the traditional medicine in the Bushi area, Kive province, Democratic Republic of Congo.** *African Study Monographs* 1998, **19**:13–33.
41. Cousins D, Huffman MA: **Medicinal properties in the diet of gorillas: an ethno- pharmacological evaluation.** *African Studies Monographs* 2002, **23**:65–89.
42. Kuete V, Kamga J, Sandjo LP, Ngameni B, Poumale HMP, Ambassa P, Ngadjui BT: **Antimicrobial activities of the methanol extract, fractions and compounds from *Ficus polita* Vahl. (Moraceae).** *BMC Complement Altern Med* 2011, **11**:10. 1186/1472-6882-11-6.
43. Kirtikar J, Basu B: ***Ficus religiosa*.** In *Indian Medicinal Plants, vol. 10, 2nd edition*. Edited by Blatter E, Caius J, Mhaskar K. New Delhi: Oriental Enterprises; 2001:3200–3202.
44. Simmonds MSJ, Howes MJR: **Plants used in the treatment of diabetes.** In *Traditional medicine for modern times - Antidiabetic Plants, 6th volume*. Edited by Soumyanath A. UK: CRC Press/Taylor and Frances Group; 2006:19–82.
45. Lansky EP, Paavilainen HM, Pawlus AD, Newman RA: ***Ficus* spp. (fig): ethnobotany and potential as anticancer and anti-inflammatory agents.** *J Ethnopharmacol* 2008, **119**:195–213.
46. Sandabe UK, Onyeyili PA, Chibuzo GA: **Phytochemical screening and effect of aqueous extract of *Ficus sycamoros* L. (Moraceae) stem bark on muscular activity in laboratory animals.** *J Ethnopharmacol* 2006, **104**:283–285.
47. Etuk EU, Bello SO, Isezuo SA, Mohammed BJ: **Ethnobotanical survey of medicinal plants used for the treatment of diabetes mellitus in the north western region of Nigeria.** *Asian J Exp Biol Sci* 2010, **1**:55–59.
48. Kotze M, Eloff JN: **Extraction of antibacterial compounds from *Combretum microphyllum* (Combretaceae).** *South African J Botany* 2002, **66**:62–70.
49. Eloff JN: **Which extractant should be used for the screening and isolation of antimicrobial components from plants?** *J Ethnopharmacol* 1998, **60**:1–8.
50. Ali H, Houghton PJ, Soumyanath A: **Alpha-amylase inhibitory activity of some Malaysian plants used to treat diabetes; with particular reference to *Phyllanthus amarus*.** *J Ethnopharmacol* 2006, **107**:449–455.
51. Djeridane A, Yousfi M, Nadjemi B, Boutassouna D, Stocher P, Vidal N: **Antioxidant activity of some Algerian medicinal plants extracts containing phenolic compounds.** *Food Chem* 2006, **97**:654–660.
52. Re-Roberta P, Proteggente N, Pannala A, Yang A, Rice-Evans C: **Antioxidant activity applying an improved ABTS radical cation decolorization assay.** *Free Radic Biol Med* 1999, **26**:1231–1237.
53. Oki T, Matsui T, Osajima Y: **Inhibitory effect of alpha-glucosidase inhibitor varies according to its origin.** *Journal of Agriculture and Food Chemistry* 1999, **47**:550–553.
54. Bhandari MR, Jong-Anurakkun N, Hong G, Kawabata J: **Alpha-glucosidase and α - amylase inhibitory activities of Nepalese medicinal herb Pakhanbhed (*Bergenia ciliata*, Haw.).** *Food Chem* 2008, **106**:247–252.
55. Tiwari AK, Rao JM: **Diabetes mellitus and multiple therapeutic approaches of phytochemicals: Present status and future prospects.** *Curr Sci* 2002, **83**:30–38.
56. Lebovitz HE: **α -Glucosidase inhibitors as agents in the treatment of diabetes.** *Diabetes Review* 1998, **6**:132–145.
57. Baron AD: **Postprandial hyperglycaemia and alpha-glucosidase inhibitors.** *Diabetes Res Clin Pract* 1998, **40**:51–55.
58. Kim J-S, Kwon C-S, Son K-H: **Inhibition of alpha glucosidase and amylase by luteolin, a flavonoid.** *Biosci Biotechnol Biochem* 2000, **64**:2458–2461.
59. McDougall GJ, Shpiro F, Dobson P, Smith P, Blake A, Stewart D: **Different polyphenolic components of soft fruits inhibit α -amylase and α -glucosidase.** *Journal of Agriculture and Food Chemistry* 2005, **53**:2760–2766.
60. Ahmed F, Urooj A: **Effect of *Ficus racemosa* stem bark on the activities of carbohydrate hydrolysing enzymes: an *in vitro* study.** *Pharm Biol* 2010, **48**(5):518–523.
61. Ahmed F, Chavan S, Satish A, Punith Kumar R: **Inhibitory activities of *Ficus benghalensis* bark against carbohydrate hydrolysing enzymes- An *in vitro* study.** *Pharmacognosy Journal* 2011, **3**(20):33–37.
62. Farsi E, Shafaei A, Hor SY, Khadeer Ahamed MB, Yam MF, Attitalla IH, Asmawi MZ, Ismail Z: **Correlation between enzymes inhibitory effects and antioxidant activities of standardised fractions of methanolic extract obtained from *Ficus deltoidea* leaves.** *Afr J Biotechnol* 2011, **10**(67):15184–15194.
63. Ponnusamy S, Ravindran R, Zinjarde S, Bhargava S, Kumar AR: **Evaluation of traditional Indian antidiabetic medicinal plants for human pancreatic amylase inhibitory effect *in vitro*.** *Evidence-Based Complementary and Alternative Medicine* 2011. doi:10.1155/2011515647.
64. Tadera K, Minami Y, Takamatsu K, Matsuoka T: **Inhibition of α -glucosidase and α - amylase by flavonoids.** *J Nutr Sci Vitaminol* 2006, **52**:149–152.
65. Brown JE, Khodr H, Hider RC, Rice-Evans CA: **Structural dependence of flavonoid interactions with Cu²⁺ ions: implications for their antioxidant properties.** *Biochem J* 1998, **330**:1173–1178.
66. Carlsen H, Myhrstad MCW, Thoreson M, Moskaug JO, Blomhoff R: **Berry intake increases the activity of the gamma-glutamyl cysteine synthetase promoter in transgenic reporter mice.** *J Nutr* 2003, **133**:2137–2140.
67. Rice-Evans CA, Miller NJ, Paganga G: **Structure-antioxidant activity relationships of flavonoids and phenolic acids.** *Free Radic Biol Med* 1996, **20**:933–956.
68. Pulido R, Bravo L, Saura-Calixto F: **Antioxidant activity of dietary polyphenols as determined by a modified ferric reducing/antioxidant power assay.** *Journal of Agriculture and Food Chemistry* 2000, **48**:3396–3402.
69. Abdel-Hameed E-SS: **Total phenolic contents and free radical scavenging activity of certain Egyptian *Ficus* species leaf samples.** *Food Chem* 2009, **114**:1271–1277.
70. Hagerman AE, Riedl KM, Jones GA, Sovik KN, Ritchard NT, Hartzfeld PW, Riechel TL: **High molecular weight plant polyphenolics (tannins) as biological antioxidants.** *Journal of Agriculture and Food Chemistry* 1998, **46**:1887–1892.
71. Ivanova D, Gerova D, Chervenkov T, Yankova T: **Polyphenols and antioxidant capacity of Bulgarian medicinal plants.** *J Ethnopharmacol* 2005, **96**:145–150.
72. Manian R, Anusuya N, Siddhuraju P, Manian S: **The antioxidant activity and free radical scavenging potential of two different solvent extracts of *Camellia sinensis* (L.) O. Kuntz, *Ficus bengalensis* L. and *Ficus racemosa* L.** *Food Chem* 2008, **107**:1000–1007.
73. Dixon M, Webb EC: **Enzyme inhibition and activation.** In *Enzymes*. 3rd edition. New York: Academic Press Inc; 1999:332–380. Longman Group Ltd. London.

doi:10.1186/1472-6882-13-94

Cite this article as: Olaokun et al.: Evaluation of the inhibition of carbohydrate hydrolysing enzymes, antioxidant activity and polyphenolic content of extracts of ten African *Ficus* species (Moraceae) used traditionally to treat diabetes. *BMC Complementary and Alternative Medicine* 2013 **13**:94.