

Understanding diversity patterns in bacterioplankton communities from a Sub-Antarctic peatland

María Victoria Quiroga^{1*}, Angel Valverde^{2*}, Gabriela Mataloni¹ and Don Cowan²

¹Instituto de Investigación e Ingeniería Ambiental (3iA), Universidad Nacional de San Martín, Buenos Aires, Argentina

²Department of Genetics, Centre for Microbial Ecology and Genomics (CMEG), Genomics Research Institute (GRI), University of Pretoria, Pretoria, South Africa

*These authors contributed equally to this work

Correspondence:

Angel Valverde, Department of Genetics, Centre for Microbial Ecology and Genomics, University of Pretoria, Lynnwood Road, Pretoria 0028, South Africa

Phone: (+27) 012 420 6944

Fax: (+27) 012 420 6870

Email: angel.valverde@up.ac.za

Running title: Bacterioplankton diversity in a Sub-Antarctic peatland

Keywords: Bacterioplankton, bog pools, diversity, phylogenetic clustering, pyrosequencing

Abstract

Bacterioplankton communities inhabiting peatlands have the potential to influence local ecosystem functions. However, most microbial ecology research in such wetlands has been done in ecosystems (mostly peat soils) of the Northern Hemisphere, and very little is known of the factors that drive bacterial community assembly in other regions of the world. In this study we used high-throughput sequencing to analyse the structure of the bacterial communities in five pools located in a sub-Antarctic peat bog (Tierra del Fuego, Argentina), and tested for relationships between bacterial communities and environmental conditions. Bacterioplankton communities in peat bog pools were diverse and dominated by members of the *Proteobacteria*, *Actinobacteria*, *Bacteroidetes* and *Verrucomicrobia*. Community structure was largely explained by differences in hydrological connectivity, pH and nutrient status (ombrotrophic vs minerotrophic pools). Bacterioplankton communities in ombrotrophic pools showed phylogenetic clustering, suggesting a dominant role of deterministic processes in shaping these assemblages. These correlations between habitat characteristics and bacterial diversity patterns provide new insights into the factors regulating microbial populations in peatland ecosystems.

INTRODUCTION

Peatlands are ecosystems characterised by high acidity, low temperatures, and low concentrations of mineral nutrients, in which the amount of carbon sequestered in net primary production (NPP), mainly by *Sphagnum* mosses, exceeds the amount of carbon lost by decomposition of organic matter by microorganisms (Dedysh, 2011). Thus, peatlands are acknowledged as an important sink for atmospheric CO₂, with carbon accumulation rates of 10-30 g C m⁻² year⁻¹ and a global carbon pool of 200-450 Pg of carbon, which constitutes about 30% of the global soil C pool (Gorham, 1991). Decomposition of organic matter in deep anoxic peat layers generates methane (CH₄), which diffuses to the surface and is then partially emitted to the atmosphere, making peatlands a globally important source of methane (Basiliko *et al.*, 2013). Furthermore, in addition to their importance in the terrestrial carbon cycle, these ecosystems hold a key role in the global water balance, regulating the hydrological regime of rivers and represent one of the largest reservoirs of fresh water (Andersen *et al.*, 2013).

Microbial diversity research in peatlands has shown these habitats containing highly specific bacterial communities being dominated by members of the phyla *Acidobacteria*, *Proteobacteria*, *Bacteroidetes* and *Verrucomicrobia* (reviewed in Dedysh 2011). In these ecosystems bacterial communities have been found to vary along small-scale hydrological and chemical gradients. For instance, within a given peatland type, dissolved organic carbon (DOC) content (Lin *et al.*, 2012), as well as substrate quality and site wetness (Jaatinen *et al.*, 2007), have been proven to affect the composition of bacterial communities. Vegetation type also plays an important role in shaping bacterial communities in peatlands (Bragina *et al.*, 2012). However, an important limitation is that most of these researches have been done in a limited number of peatland ecosystems (i.e. peat soils), mainly situated in the Northern Hemisphere (see Andersen *et al.*, 2013). Therefore, more geographically and ecologically diverse samples (e.g. bog pools) are needed to elucidate the biogeographic patterns and environmental factors that shape the structure of bacterial communities in peatland ecosystems at different spatial scales.

Here we have investigated, for the first time, the bacterial diversity in five freshwater pools located along a 500 m transect in Rancho Hambre peat bog (Tierra del Fuego, Argentina). Pools are critical habitats for biodiversity in natural peatlands (Mazerolle *et al.*, 2006). Taking advantage of differences in hydrological connectivity, pool morphometry, and minero- vs ombro-trophic status (González Garraza *et al.*, 2012), we

addressed the following questions: (i) Does bacterial community composition vary within and between pools, and differ from that of northern peatlands? (ii) Do variations in bacterial diversity reflect the underlying environmental conditions (Table S1)? And, if this is observed, (iii) Are bacterial communities from ombrotrophic (more acidic and hydrologically isolated) pools more phylogenetically clustered than those inhabiting minerotrophic pools (less acidic and interconnected)? Analysis of the degree of phylogenetic relatedness of taxa found within and between communities should provide insights into the processes that organize these communities (Webb *et al.*, 2002). To answer these questions we used high-throughput barcode amplicon pyrosequencing and a set of multivariate statistic tools.

RESULTS AND DISCUSSION

We have investigated bacterial diversity patterns in five pools from a sub-Antarctic peat bog located in Tierra del Fuego, Argentina (see *Supporting information* for materials and methods). A total of 897 bacterial operational taxonomic units (OTUs, sequences binned at a 97% similarity cutoff) were identified in 30,225 sequences (2015 per sample, n=15); an average of 109 ± 5 OTUs (mean \pm SE) per sample (Fig. S1). A collector's curve of the number of OTUs per sample revealed that additional bacterial taxa are likely to appear with every additional sample (Fig. S2). The total size of the bacterial OTU pool was estimated to be $1,764 \pm 111$ OTUs (mean \pm SE of Chao1 estimator of total OTU pool richness). Minerotrophic (less acidic) pools (RH1, RH4) typically had higher bacterial diversity than ombrotrophic (more acidic) pools (Fig. S1), although differences were statistically significant only for Shannon and inverse Simpson indices (paired t-test, $P < 0.05$); which agrees with previous results from other peatland environments (Opelt *et al.*, 2007; Peltoniemi *et al.*, 2009; Lin *et al.*, 2012).

Thirteen phyla were detected across all samples (Fig. S3), of which *Proteobacteria* (51% of all sequences), *Actinobacteria* (22%), *Bacteroidetes* (15%) and *Verrucomicrobia* (7%) were the most abundant, constituting 95% of the sequences. These four phyla have been shown to be widely distributed in *Sphagnum*-dominated wetlands (Juottonen *et al.*, 2005; Dedysh *et al.*, 2006; Morales *et al.*, 2006; Hartman *et al.*, 2008; Ausec *et al.*, 2009; Kanokratana *et al.*, 2011; Pankratov *et al.*, 2011; Bragina *et al.*, 2013), although their relative abundances appear to vary across ecosystems.

In contrast to those studies, although detected, we did not find members of the


Acidobacteria to be important components of the freshwater bacterial communities, more likely because large differences in habitat characteristics (peat soil vs. water material). Indeed, bacterial communities were found to shift in composition along the landscape from peat communities dominated by *Acidobacteria* to freshwater communities dominated by *Actinobacteria* (Kulichevskaya et al., 2011). In spite of this striking difference, the overlap between peat soil and freshwater bacterial communities support previous findings obtained across soil, sediment, stream and lake habitats; which suggest that aquatic environments are strongly coupled to terrestrial ecosystems through hydrological networks (Crump et al., 2012, and references therein).

The vast majority of the taxa found were rare, with 72% of OTUs occurring in a single sample (Fig. S4). Nevertheless, common and abundant bacterial phylotypes were also detected. Ten bacterial OTUs belonged to two phyla that were present in 80% or more of the samples, representing 1.1% of the bacterial taxonomic diversity but 53% of sequences (Fig. 1). These phyla were *Proteobacteria* (38% of all sequences, mostly *Betaproteobacteria*) and *Actinobacteria* (12%). The most abundant individual OTUs that could be classified in the “core microbiome” at the genus level were *Polynucleobacter* (11%, *Betaproteobacteria*) and *Novosphingobium* (3%, *Alphaproteobacteria*). Members of the genera *Mucilaginibacter* (*Bacteroidetes*), *Flavobacterium* (*Bacteroidetes*) and *Limnohabitans* (*Betaproteobacteria*) were also abundant (representing 7%, 3% and 1% of all sequences, respectively), but were found in a lower number of samples (Fig. 2).

Knowledge is scarce about the specific ecology of these microorganisms in peatland ecosystems. Many of the bacterial taxa identified are not available in pure culture, which hampers determination of their physiologies and consequently assessment of their functional roles in these environments. Nevertheless, bacterial strains affiliated to *Polynucleobacter*, namely *P. neccessarius*, have been reported to occur as obligate endosymbionts of ciliates (subsp. *neccessarius*) [commonly found in the Rancho Hambre water bodies (Quiroga et al., 2013)], and also as free-living strains (subsp. *asymbioticus*). The complete genomic sequence of both subspecies has shown the inability of either living forms to exploit sugars as carbon or energy sources or to perform nitrification, denitrification or nitrogen fixation (Boscaro et al., 2013). Conversely, the free-living form can perform the assimilatory reduction of nitrate and assimilate sulphur and sulphate.

Members of the genus *Limnohabitans*, highly widespread in freshwater food webs, have

Fig. 1. Phylogenetic assignment and sequence distribution of ‘core’ OTUs (97% cutoff). Taxonomic assignments are the finest level that passed the Ribosomal Database Project classifier’s (80% confidence threshold).



been ascribed a prominent role in the transfer of DOC from both autochthonous and terrestrial sources to the top trophic levels (Kasalický *et al.*, 2013; Šimek *et al.*, 2013). Most Bacteroidetes (i.e. *Flavobacterium* and *Mucilaginibacter*) have the ability to use a broad range of biopolymers, particularly polysaccharides and proteins (Thomas *et al.*, 2011), and it has been proposed that a primary role for Flavobacteria is the conversion of high molecular weight (HMW) compounds into low molecular weight (LMW) compounds (Teeling *et al.*, 2012). Indeed, analysis of DOC quality in Rancho Hambro pools revealed the predominance of coloured, aromatic, HMW compounds (Quiroga *et al.*, 2013), as found in other bog ecosystems (Hodgkins *et al.*, 2014). Several *Novosphingobium* strains seem to have the capacity to degrade phenolic compounds from *Sphagnum*-derived litter (DeAngelis *et al.*, 2013).

Interestingly, all OTUs belonging to *Limnohabitans* and *Flavobacterium* were found exclusively in minerotrophic pools (Fig. 2), which agrees with the fact that these genera strongly prefer non-acidic habitats (Šimek *et al.*, 2010). In direct contrast, members of the *Mucilaginibacter*, which are reported to grow in a wide range of pH (Pankratov *et al.*, 2007), were only found in ombrotrophic pools. Whether or not this lack of co-occurrence is due to biological interactions (e.g. competition or predation) or species-specific habitat associations deserves future investigation. All in all, the results suggest that there is a high degree of niche specialization among these taxa and that they play an important role for nutrient cycling in peatland limnetic ecosystems.

Bacterial communities in a given pool were highly similar and significantly different from those in other pools (ANOSIM $R=1$, $P<0.001$). The inverse of Whittaker's β -diversity measure, in which mean within-pool diversity (α -diversity) is divided by overall regional diversity (γ -diversity), was close to 0 (Whittaker's $\beta^{-1} = 0.12$), indicating that between-pool diversity contributed more to total diversity than within-site diversity (Anderson *et al.*, 2011). Variation in bacterial community structure was related to pH, ammonium, phosphate and total nitrogen. Based on Bray-Curtis dissimilarities, a permutational multivariate analysis of variance (PERMANOVA) found each of these four factors to be significant in structuring pool bacterial communities (PERMANOVA: $P < 0.01$; combined coefficient of determination $R^2 = 0.69$). pH explained most of the variation ($R^2 = 0.25$), followed by total N ($R^2 = 0.20$), ammonium ($R^2 = 0.14$) and phosphate ($R^2 = 0.10$). As pH can be considered a 'master variable' for the chemical state of aquatic ecosystems that influences and is influenced by other variables (Blodau, 2006), we cannot determine whether pH has a direct or

Fig. 2. Heatmap displaying the most abundant genera for ombrotrophic and minerotrophic samples. Pools are clustered based on the percent relative abundance of the twenty three genus;phylum level classifications shown as rows in this figure. Each row was scaled so that the mean of each taxonomic group across samples was calculated and coloured by corresponding z-score of each cell.




Fig. 3. nMDS ordination plot (Hellinger-transformed Bray-Curtis dissimilarity matrix). Each point represents the bacterial community of an individual sample and shapes denote different pools (n=5). Environmental vectors were fit to the centroids of each pool.




Fig. 4. Standardised effect sizes of MPD (SES.MPD) and MNTD (SES.MNTD) for bacterioplankton communities. Asterisks indicate significant results ($P < 0.05$).



indirect effect on these communities. Notwithstanding that the amount of community variation explained by each factor was conditional on the distance metric used, these four factors always remained significant (data not shown).

A non-metric multidimensional scaling plot confirmed the strong effect of these factors in the assembly of bacterial communities (Fig. 3), and grouped pools according to their trophic status; that is, minerotrophic (RH1 and RH4) vs. ombrotrophic (RH2, RH3 and RH5), as described in González Garraza *et al.* (2012). When pH, ammonium, phosphate and total N were fitted to linear vectors in the ordination space, these vectors were highly correlated with the placement of water pool centroids (pH: $r^2 = 0.60$, $P < 0.01$; ammonium: $r^2 = 0.79$, $P < 0.01$; phosphate: $r^2 = 0.95$, $P < 0.01$; total N: $r^2 = 0.78$, $P < 0.01$). Strong relationships between pH, nutrient concentrations, and bacterial community composition have been observed in other wetlands (Hartman *et al.*, 2008), and suggest environmental filtering (Wang *et al.*, 2013). Indeed, environmental filtering seems to shape bacterial communities in some ombrotrophic pools (i.e. RH3 and RH5) (Fig. 4), as we were able to detect phylogenetic clustering using ecological null models (Kembel, 2009). The link between bacterial phylogenetic clustering and ombrotrophic status (SES.MPD = -3.1 ± 0.1 , SES.MNTD = -1.6 ± 0.1 ; both $P < 0.05$) indicates that the prevailing conditions of these pools act as a stronger ecological filter compared to those of minerotrophic pools (SES.MPD = 0.5 ± 0.2 , SES.MNTD = -0.6 ± 0.2 ; both $P > 0.05$). Habitat filtering has also been proposed as the major mechanism explaining bacterial community assembly in a peaty acidic soil (Felske *et al.*, 1998).

Ombrotrophic environments are hydrologically isolated from the surrounding landscape and characterised by acidic pH and low nutrient values (Wheeler and Proctor, 2000). Both environmental conditions and spatial isolation can cause phylogenetic clustering in bacterial communities (Horner-Devine and Bohannan, 2006; Bryant *et al.*, 2008; Stegen *et al.*, 2013), and suggest that habitat heterogeneity can contribute to observable biogeographic patterns in microbial communities at limited (within-peatland) spatial scales. It is important to note, however, that habitat filtering (abiotic factors and biotic interactions), together with differential dispersal (connectivity among communities as a result of water run out and physical barriers) and ecological drift (stochastic changes in the relative abundance of taxa), likely act in concert (Vellend, 2010; Stegen *et al.*, 2013) to produce the bacterial communities studied here.

Acknowledgments

Financial support was provided by CONICET (Research grant PIP 11220090100050), the National Research Foundation (South Africa), and the Ministerio de Ciencia, Tecnología e Innovación Productiva (South Africa-Argentina International Science and Technology agreements). We are grateful to the Secretaría de Desarrollo Sustentable y Ambiente, Dirección General de Recursos Hídricos, Provincia de Tierra del Fuego and the Centro Austral de Investigaciones Científicas (CADIC)-CONICET for valuable logistical support. The help of Sergio Camargo and Gabriela González Garraza during sampling was much appreciated.

Conflict of interest

The authors declare no conflict of interest

References

- Andersen R., Chapman S.J., and Artz R.R.E. (2013) Microbial communities in natural and disturbed peatlands: A review. *Soil Biol Biochem* **57**: 979-994.
- Anderson M.J., Crist T.O., Chase J.M., Vellend M., Inouye B.D., Freestone A.L., *et al.* (2011) Navigating the multiple meanings of beta-diversity: A roadmap for the practicing ecologist. *Ecol Lett* **14**: 19-28.
- Ausec L., Kraigher B., and Mandic-Mulec I. (2009) Differences in the activity and bacterial community structure of drained grassland and forest peat soils. *Soil Biol Biochem* **41**: 1874-1881.
- Basiliko N., Henry K., Gupta V., Moore T.R., Driscoll B.T., and Dunfield P.F. (2013) Controls on bacterial and archaeal community structure and greenhouse gas production in natural, mined, and restored Canadian peatlands. *Front Microbiol* **4**: 215
- Blodau C. (2006) A review of acidity generation and consumption in acidic coal mine lakes and their watersheds. *Sci Tot Environ* **369**: 307-332.
- Boscaro V., Felletti M., Vannini C., Ackerman M.S., Chain P.S.G., Malfatti S., *et al.* (2013) Polynucleobacter necessarius, a model for genome reduction in both free-living and symbiotic bacteria. *Proc Natl Acad Sci USA* **110**: 18590-18595.
- Bragina A., Berg C., Cardinale M., Shcherbakov A., Chebotar V., and Berg G. (2012) Sphagnum mosses harbour highly specific bacterial diversity during their whole lifecycle. *ISME J* **6**: 802-813.
- Bragina A., Berg C., Müller H., Moser D., and Berg G. (2013) Insights into functional bacterial diversity and its effects on Alpine bog ecosystem functioning. *Scientific reports* **3**: 1955.
- Bryant J.A., Lamanna C., Morlon H., Kerkhoff A.J., Enquist B.J., and Green J.L. (2008) Microbes on mountainsides: Contrasting elevational patterns of bacterial and plant diversity. *Proc Natl Acad Sci USA* **105**: 11505-11511.

- Crump B.C., Amaral-Zettler L.A., and Kling G.W. (2012) Microbial diversity in arctic freshwaters is structured by inoculation of microbes from soils. *ISME J* **6**: 1629-1639.
- DeAngelis K.M., Chivian D., Fortney J.L., Arkin A.P., Simmons B., Hazen T.C., and Silver W.L. (2013) Changes in microbial dynamics during long-term decomposition in tropical forests. *Soil Biol Biochem* **66**: 60-68.
- Dedysh S.N., Pankratov T.A., Belova S.E., Kulichevskaya I.S., and Liesack W. (2006) Phylogenetic analysis and in situ identification of Bacteria community composition in an acidic Sphagnum peat bog. *Appl Environ Microbiol* **72**: 2110-2117.
- Dedysh S.N. (2011) Cultivating uncultured bacteria from northern wetlands: Knowledge gained and remaining gaps. *Front Microbiol* **2**: 184
- Felske A., Wolterink A., Van Lis R., and Akkermans A.D.L. (1998) Phylogeny of the main bacterial 16S rRNA sequences in drentse A grassland soils (The Netherlands). *Appl Environ Microbiol* **64**: 871-879.
- González Garraza G., Mataloni G., Iturraspe R., Lombardo R., Camargo S., and Quiroga M.V. (2012) The limnological character of bog pools in relation to meteorological and hydrological features. *Mires Peat* **7**: 1-14.
- Gorham E. (1991) Northern peatlands: role in the carbon cycle and probable responses to climatic warming. *Ecol Appl* **1**: 182-195.
- Hartman W.H., Richardson C.J., Vilgalys R., and Bruland G.L. (2008) Environmental and anthropogenic controls over bacterial communities in wetland soils. *Proc Natl Acad Sci USA* **105**: 17842-17847.
- Hodgkins S.B., Tfaily M.M., McCalley C.K., Logan T.A., Crill P.M., Saleska S.R., *et al.* (2014) Changes in peat chemistry associated with permafrost thaw increase greenhouse gas production. *Proc Natl Acad Sci USA* **111**: 5819-5824.
- Horner-Devine M.C., and Bohannon B.J.M. (2006) Phylogenetic clustering and overdispersion in bacterial communities. *Ecology* **87**: S100-S108.
- Jaatinen K., Fritze H., Laine J., and Laiho R. (2007) Effects of short- and long-term water-level drawdown on the populations and activity of aerobic decomposers in a boreal peatland. *Glob Chang Biol* **13**: 491-510.
- Juottonen H., Galand P.E., Tuittila E.S., Laine J., Fritze H., and Yrjälä K. (2005) Methanogen communities and Bacteria along an ecohydrological gradient in a northern raised bog complex. *Environ Microbiol* **7**: 1547-1557.
- Kanokratana P., Uengwetwanit T., Rattanachomsri U., Bunternngsook B., Nimchua T., Tangphatsornruang S., *et al.* (2011) Insights into the Phylogeny and Metabolic Potential of a Primary Tropical Peat Swamp Forest Microbial Community by Metagenomic Analysis. *Microb Ecol* **61**: 518-528.
- Kasalický V., Jezbera J., Hahn M.W., and Šimek K. (2013) The Diversity of the Limnohabitans Genus, an Important Group of Freshwater Bacterioplankton, by Characterization of 35 Isolated Strains. *PLoS ONE* **8**: e58209.
- Kembel S.W. (2009) Disentangling niche and neutral influences on community assembly: assessing the performance of community phylogenetic structure tests. *Ecol Lett* **12**: 949-960.
- Kulichevskaya I.S., Belova S.E., Komov V.T., Dedysh S.N., and Zavarzin G.A. (2011) Phylogenetic composition of bacterial communities in small boreal lakes and ombrotrophic bogs of the upper Volga basin. *Microbiology* **80**: 549-557.
- Lin X., Green S., Tfaily M.M., Prakash O., Konstantinidis K.T., Corbett J.E., *et al.* (2012) Microbial community structure and activity linked to contrasting biogeochemical gradients in bog and fen environments of the glacial lake agassiz peatland. *Appl Environ Microbiol* **78**: 7023-7031.

- Mazerolle M.J., Poulin M., Lavoie C., Rochefort L., Desrochers A., and Drolet B. (2006) Animal and vegetation patterns in natural and man-made bog pools: Implications for restoration. *Freshwater Biol* **51**: 333-350.
- Morales S.E., Mouser P.J., Ward N., Hudman S.P., Gotelli N.J., Ross D.S., and Lewis T.A. (2006) Comparison of bacterial communities in New England Sphagnum bogs using terminal restriction fragment length polymorphism (T-RFLP). *Microb Ecol* **52**: 34-44.
- Opelt K., Berg C., Schönmann S., Eberl L., and Berg G. (2007) High specificity but contrasting biodiversity of Sphagnum-associated bacterial and plant communities in bog ecosystems independent of the geographical region. *ISME J* **1**: 502-516.
- Pankratov T.A., Tindall B.J., Liesack W., and Dedysh S.N. (2007) Mucilaginibacter paludis gen. nov., sp. nov. and Mucilaginibacter gracilis sp. nov., pectin-, xylan and laminarin-degrading members of the family Sphingobacteriaceae from acidic Sphagnum peat bog. *Int J Syst Evol Microbiol* **57**: 2349-2354.
- Pankratov T.A., Ivanova A.O., Dedysh S.N., and Liesack W. (2011) Bacterial populations and environmental factors controlling cellulose degradation in an acidic Sphagnum peat. *Environ Microbiol* **13**: 1800-1814.
- Peltoniemi K., Fritze H., and Laiho R. (2009) Response of fungal and actinobacterial communities to water-level drawdown in boreal peatland sites. *Soil Biol Biochem* **41**: 1902-1914.
- Quiroga M.V., Unrein F., Garraza G.G., Küppers G., Lombardo R., Marinone M.C., *et al.* (2013) The plankton communities from peat bog pools: Structure, temporal variation and environmental factors. *J Plankton Res* **35**: 1234-1253.
- Šimek K., Kasalický V., Jezbera J., Jezberová J., Hejzlar J., and Hahn M.W. (2010) Broad habitat range of the phylogenetically narrow R-BT065 cluster, representing a core group of the betaproteobacterial genus limnohabitans. *Appl Environ Microbiol* **76**: 631-639.
- Šimek K., Kasalický V., Jezbera J., Horňák K., Nedoma J., Hahn M.W., *et al.* (2013) Differential freshwater flagellate community response to bacterial food quality with a focus on Limnohabitans bacteria. *ISME J* **7**: 1519-1530.
- Stegen J.C., Lin X., Fredrickson J.K., Chen X., Kennedy D.W., Murray C.J., *et al.* (2013) Quantifying community assembly processes and identifying features that impose them. *ISME J* **7**: 2069-2079.
- Teeling H., Fuchs B.M., Becher D., Klockow C., Gardebrecht A., Bennke C.M., *et al.* (2012) Substrate-controlled succession of marine bacterioplankton populations induced by a phytoplankton bloom. *Science* **336**: 608-611.
- Thomas F., Hehemann J.H., Rebuffet E., Czjzek M., and Michel G. (2011) Environmental and gut Bacteroidetes: The food connection. *Front Microbiol* **2**: 93
- Vellend M. (2010) Conceptual synthesis in community ecology. *Q Rev Biol* **85**: 183-206.
- Wang J., Shen J., Wu Y., Tu C., Soininen J., Stegen J.C., *et al.* (2013) Phylogenetic beta diversity in bacterial assemblages across ecosystems: deterministic versus stochastic processes. *ISME J* **7**: 1310-1321.
- Webb C.O., Ackerly D.D., McPeck M.A., and Donoghue M.J. (2002) Phylogenies and community ecology. *Annu Rev Ecol Syst* **33**: 475-505.
- Wheeler B.D., and Proctor M.C.F. (2000) Ecological gradients, subdivisions and terminology of north-west European mires. *J Ecol* **88**: 187-203.

Supporting Information

Supplementary materials and methods.

Fig. S1. Distribution and diversity of bacterial OTUs (97% cutoff) after rarefying the data set to 2015 sequences per sample. Error bars depict standard errors. PD, phylogenetic diversity.

Fig. S2. Collector's curve (mean \pm 95% confidence interval) of bacterial OTU (97% cutoff) richness vs. number of samples.

Fig. S3. Taxonomic distribution of bacterial OTUs (97% cutoff). Affiliation was performed using the Ribosomal Database Project Classifier with a confidence threshold of 80%. The sequence data generated in this study were deposited in the NCBI Sequence Read Archive and are available under the project number SRP050278. A supplementary table containing sample metadata conforming to MIMARKS standards is also provided (Table S1).

Fig. S4. Frequency histogram: number of bacterial OTU (97% cutoff) that occurred in each sample (of 15 possible).

Table S1. Metadata, including the chemistry of the samples, conforming to MIMARKS standards.