

RESEARCH ARTICLE

Nosema Tolerant Honeybees (*Apis mellifera*) Escape Parasitic Manipulation of Apoptosis

Christoph Kurze^{1*}, Yves Le Conte², Claudia Dussaubat², Silvio Erler¹, Per Kryger³, Oleg Lewkowski¹, Thomas Müller⁴, Miriam Widder⁴, Robin F. A. Moritz^{1,5,6}

1 Institute for Biology, Martin-Luther-Universität Halle-Wittenberg, Halle (Saale), Germany, **2** UR 406 Abeilles et Environnement, INRA, Avignon, France, **3** Department of Agroecology, Aarhus University, Flakkebjerg, Denmark, **4** Department of Internal Medicine IV, Martin-Luther-Universität Halle-Wittenberg, Halle (Saale), Germany, **5** German Institute for Integrative Biodiversity Research (iDiv), Leipzig, Germany, **6** University of Pretoria, Department of Zoology and Entomology, Pretoria, South Africa

* christoph.kurze@zoologie.uni-halle.de



OPEN ACCESS

Citation: Kurze C, Le Conte Y, Dussaubat C, Erler S, Kryger P, Lewkowski O, et al. (2015) *Nosema* Tolerant Honeybees (*Apis mellifera*) Escape Parasitic Manipulation of Apoptosis. PLoS ONE 10(10): e0140174. doi:10.1371/journal.pone.0140174

Editor: Wolfgang Blenau, University of Cologne, GERMANY

Received: May 20, 2015

Accepted: September 21, 2015

Published: October 7, 2015

Copyright: © 2015 Kurze et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This study was funded by the Deutsche Forschungsgemeinschaft through the priority programme "Host-Parasite Coevolution – Rapid Reciprocal Adaptation and its Genetic Basis" (SPP 1399, Grant number MO373/26-2) to RFAM and a scholarship from The Foundation for the Preservation of Honey Bees, Inc. to CK.

Competing Interests: The authors have declared that no competing interests exist.

Abstract

Apoptosis is not only pivotal for development, but also for pathogen defence in multicellular organisms. Although numerous intracellular pathogens are known to interfere with the host's apoptotic machinery to overcome this defence, its importance for host-parasite coevolution has been neglected. We conducted three inoculation experiments to investigate in the apoptotic respond during infection with the intracellular gut pathogen *Nosema ceranae*, which is considered as potential global threat to the honeybee (*Apis mellifera*) and other bee pollinators, in sensitive and tolerant honeybees. To explore apoptotic processes in the gut epithelium, we visualised apoptotic cells using TUNEL assays and measured the relative expression levels of subset of candidate genes involved in the apoptotic machinery using qPCR. Our results suggest that *N. ceranae* reduces apoptosis in sensitive honeybees by enhancing *inhibitor of apoptosis protein-(iap)-2* gene transcription. Interestingly, this seems not be the case in *Nosema* tolerant honeybees. We propose that these tolerant honeybees are able to escape the manipulation of apoptosis by *N. ceranae*, which may have evolved a mechanism to regulate an anti-apoptotic gene as key adaptation for improved host invasion.

Introduction

In insects, epithelial cells of the intestine are typically the first line of pathogen defence. They produce not only antimicrobial peptides (AMPs) and reactive oxygen species (ROS) but they can also respond with programmed cell death (including apoptosis) of infected cells. The infection may then be simply cleared by defecation. Hence, it is not surprising to see that intracellular pathogens have evolved mechanisms to overcome apoptosis for self-protection and increase of reproductive success within their host cell [1–3]. This is also the case for microsporidia [4–6], a group of highly specialised intracellular fungal parasites, causing diseases in a wide range of animal hosts, including humans and several animal species important for agriculture and aquaculture [7].

Although strategies of numerous pathogens have been studied in some detail [1–3], adaptations by the host to withstand the manipulation by these pathogen are neglected. The inhibition of apoptosis was recently shown to be pivotal for a successful infection of the microsporidian pathogen *Nosema ceranae* [5], which is considered to be a major threat to honeybees and wild bee pollinators [8–10]. Several honeybee transcriptome studies also indicated a link between *Nosema* spp. infections and apoptosis and epithelium renewal [11–13], supporting the idea that apoptosis might play a central role during *Nosema* infection. *Nosema* spores are transmitted via the faecal-oral route and germinate in the host midgut and enter epithelial cells, where they replicate and produce a new generation of spores after 4 days [9, 14].

Because Nosemosis can seriously impact colony health, Danish bee breeders successfully selected for *Nosema* resistant colonies, which appear to be the result of tolerance at the individual level [15, 16]. To examine the importance of the apoptotic defence system in the adaptation of these *Nosema* tolerant honeybees, we compared them with sensitive honeybees in three controlled inoculation experiments and screened for apoptotic processes in the honeybee midgut epithelium.

Materials and Methods

Experimental inoculation

One colony of *Nosema* tolerant honeybees from Aarhus (Denmark) was transported to Avignon (France), where one colony with *Nosema* sensitive honeybees was chosen for positive controls. We replicated three independent inoculation experiments in May 2013 following standard methods [17]. Briefly, newly emerged workers (< 24 h old) were collected from the brood frames. *Nosema* sensitive (SN) and tolerant (TN) honeybees were individually fed with 10^5 freshly extracted and purified *N. ceranae* spores in 2 μ l sucrose solution. Uninfected controls of the sensitive (SC) and tolerant (TC) honeybee strain were only fed with 2 μ l sucrose solution. Individuals that have not consumed the inoculum were discarded from the experiment. Twenty worker bees per honeybee strain were housed in sterile stainless steel cages (10×10×5.5 cm) with a piece of clean wax foundation in an incubator at $34 \pm 1^\circ\text{C}$, 60% relative humidity and provided with 50% (w/v) sucrose solution *ad libitum*. Bees were sacrificed either on one or six days post infection (p.i.). Their midguts (ventriculi without rectum) were dissected and stored accordingly to the analysis. We confirmed the treatment success by estimating the number of *Nosema* spp. spores for a random subset of 5 midgut samples for each treatment group and replicate using a Fuchs–Rosenthal haemocytometer under a phase-contrast microscope ($\times 400$).

Immunohistochemistry

4% buffered formaldehyde (Süsse) was used for fixation of three midguts per replicate for 24 h at 8°C , followed by paraffin embedding according to standard histological methods. The ratio of apoptosis was determined by TUNEL (Terminal deoxynucleotide transferase mediated X-dUTP nick endlabelling) assays (In Situ Cell Death Detection Kit, Roche) on 7 μ m thick longitudinal sections according to the manufacturer's manual. This method allows the detection of apoptotic cells at the early stage, for which selective internucleosomal DNA degradation is characteristic, by directly labelling of single- and double-stranded DNA nicks with the enzyme TdT (Terminal deoxynucleotide transferase) and fluorescein-dUTP. Prior to the TUNEL reactions we blocked endogenous peroxidase activity (Dual Endogenous Enzyme Block, Dako), followed by permeabilisation step using nucleases-free 10 μ g μ l⁻¹ proteinase K in 10 mM Tris/HCl pH 7.5 for 20 min at room temperature and rinsed the samples twice in PBS (phosphate-buffered saline). The TUNEL reaction was stopped after 1 h at 37°C in the dark by rinsing the




Fig 1. Quantification of apoptosis in the midgut epithelium of honeybees infected with *N. ceranae*. (A) The frequency of apoptotic cells was calculated as the numbers of TUNEL+ve relation to all (DAPI+ve) nuclei. For this, DAPI and TUNEL stained images (top) were merged (bottom left); nuclei were binarised and automatically counted using ImageJ (bottom right; red = TUNEL+ve, white = TUNEL-ve). Scale bars = 25 μ m. (B) Comparison of apoptotic TUNEL+ve cells detected in the posterior end of the midgut in *Nosema* infected sensitive and tolerant honeybees on day 6 p.i. Scale bars = 50 μ m. (C) Apoptosis ratio (mean \pm s.e.) during *Nosema ceranae* infection in *Nosema* sensitive (SN, solid circles) and tolerant (TN, solid squares) honeybees, and their uninfected controls (SC, open circles and TC, open squares) at 1 day (green) and 6 days (blue) after inoculation. Sample sizes are given in [S2 Table](#). Significance between treatment groups **, $P < 0.01$.

doi:10.1371/journal.pone.0140174.g001

samples three times in PBS and then counterstaining with 1 μ g ml⁻¹ DAPI (4',6-diamidino-2-phenylindole) (Sigma-Aldrich). We visualised apoptotic cells (TUNEL+ve) relative to the total number of cells (DAPI+ve) in the posterior part of midguts (primary site of the infection on day 6 p.i.) using fluorescence microscopy and acquired images with CCD camera connected to Axio Vision 4.6 (Zeiss). Automatic cell counting and analyses were performed with ImageJ [18] (Fig 1A) screening at the average 325 \pm 16 s.e. cells per sample (see also [S2 Table](#)).

Gene expression

Midguts of nine workers for each treatment group and for each replicated experiment were sampled in pools of three individuals, flash-frozen in liquid nitrogen and stored at -80°C until subsequent qPCR analyses. Briefly, total RNA was isolated using TRIZOL extraction procedure and 1 μ g RNA each were reverse transcribed. Obtained cDNA were purified using QIAquick PCR Purification Kit (Qiagen). We performed a TBLASTN search [19] of *Apis mellifera* (taxid 7460) database using full-length amino acid sequences for key proteins involved in apoptosis known from *Drosophila melanogaster* [20]. Only homologous proteins with at least 20% identity were considered for this study ([S1 Table](#)). Thus some proteins known to be relevant for apoptosis in *D. melanogaster* were not included due to insufficient homology. Gene specific primers spanning one intron of subset of nine potential candidate genes, including the genes *basket* (*bsk*), *tumor protein p53-like* (*p53*), *inhibitor of apoptosis protein 2* (*iap-2*; homologous gene to *Diap-1*), *caspase-2-like* (*casp-2*; possible homologous gene to *Dronc*) and *caspase-10-like* (*casp-10*; homologous gene to *Dredd*), were designed using Primer-BLAST on the *A. mellifera* reference genome (release v.4.5, GenBank, [S1 Table](#)). *Ribosomal protein S5a* (*RpS5a*) and *actin related protein 1* (*arp1*, also known as *actin*) were initially chosen in order to standardize expression levels between pools and treatment groups [21]. For qPCR, we used 20 ng cDNA mixed with 5 μ l SensiMixPlus (Bioline), 0.25 μ M of each primer and DEPC-water in 10 μ l final

volume. Initial denaturation at 95°C for 10 min was followed by 40 amplification cycles (95°C for 15 s, 59°C for 30 s, 72°C for 30 s), ending with melting curve analysis from 50°C to 98°C in 1°C increments. At least two technical replicates were run per sample using Chromo4™ (Bio-Rad) and repeated if necessary to obtain a delta C_t (threshold cycle) value below 0.5 between two replicates (LinRegPCR [22]). Correct amplicon sizes absence of non-specific products were verified using the high-resolution automatic capillary electrophoresis system QIAxcel® (Qiagen). We tested the suitability of *RpS5a* and *arp1* as previously described [23], and found the housekeeping gene *RpS5a* (s.d. = 0.86) to be more suitable for normalisation of gene expression levels among samples, and thus excluded the *arp1* (s.d. = 1.55) from further analyses.

Statistics

All statistical analyses and data plotting (mean ± s.e.) were performed in R (v.3.0.2) [24]. Spore load between infected groups was tested using Welch's two-tailed *t*-test. We used Generalized Linear Models (GLM) based on quasilielihood estimation with a binomial error distribution to test the effects of honeybee strain and treatment and their interactions on the apoptosis ratio for each day p.i. independently. Post-hoc analyses were performed using the Tukey's HSD. The effects of day p.i., honeybee strain and treatment and their interactions on relative gene expression were tested using linear models (LM) for each gene separately, accounting for multiple testing with Bonferroni adjustments. We used the likelihood ratio test to test single parameters and their interactions, comparing the goodness-of-fit between the models [25]. If a model was found to be unstable with all interactions included, we removed non-significant interactions step-wise. Model validity was tested by comparing full models to their null models without any fixed factors included using likelihood ratio test. Tukey's HSD post-hoc contrast analyses were performed using the *glht* function with Bonferroni adjustment (multcomp package, v.1.3–2.).

Results and Discussion

There were generally higher apoptosis rates in all four treatment groups tested on day 1 p.i. than on day 6 p.i. (GLM: day, estimate ± s.e. = -0.904 ± 0.239 , $P < 0.001$, Fig 1B), suggesting that mid-gut epithelial cells in those young bees were still undergoing morphogenetic developments at age of 1–2 day [26]. More interestingly, however, was the interaction between treatment and honeybee strain (GLM: -0.899 ± 0.293 , $P < 0.005$; Fig 1B) whereby *N. ceranae* infection reduced the rate of apoptosis in sensitive honeybees (Tukey's HSD: -0.679 ± 0.210 , $P < 0.007$), confirming the suppression of apoptosis by microsporidia [5], but interestingly this was not the case in the infected tolerant honeybees (Tukey's HSD: 0.220 ± 0.205 , $P = 1$). Hence, these results suggest an adaptive mechanism in the tolerant honeybees which enables them to remove infected cells from the epithelia into the gut lumen and presumably to eventually defecate.

As honeybees usually defecate during short defecation flights and avoid defecation in their nest [27], infected apoptotic cells have presumably been accumulated in the midgut and rectum in *Nosema* tolerant honeybees at this early stage of the infection in our cage experiments. This in fact would be a plausible explanation why we have not counted less numbers of *Nosema* spores between sensitive (SN: $6.0 \pm 1.2 \times 10^6$ spores; $n = 14$) and tolerant honeybees (TN: $8.0 \pm 1.4 \times 10^6$ spores; $n = 15$) on day 6 p.i. (*t*-test: $t = 1.115$, *d.f.* = 26.77, $P < 0.275$).

To explore the molecular mechanisms underlying the inhibition of apoptosis in the sensitive honeybee, we measured relative gene expression levels of candidate genes in the apoptotic cascade predicted from *Drosophila* [20] (S1 Table). We found age-related alterations in gene expression levels for Jun N-terminal kinase (JNK)/*basket* (*bsk*, linear model (LM): day p.i., 0.091 ± 0.026 , $P < 0.007$; Fig 2) and *tumor suppressor protein p53* (*p53*, LM: day p.i., 0.022 ± 0.005 , $P < 0.001$; Fig 2), which were slightly higher expressed on day 6 p.i. than on day




Fig 2. Relative expression (mean \pm s.e.) of candidate genes important for apoptosis in *Nosema* infected honeybees. *Nosema* sensitive (SN, solid circles) and tolerant (TN, solid squares) honeybees infected with 10^5 *N. ceranae* spores, and their controls uninfected (SC, open circles and TC, open squares), were sampled at 1 day (green) and 6 days (blue) after inoculation. The genes JNK/*bsk* (Jun N-terminal kinase/ *basket*), *p53* (*tumor protein p53-like*), *iap-2* (*inhibitor of apoptosis protein 2*; predicted homologous gene to *Diap-1* in *D. melanogaster*), *casp-2* (*caspase-2-like*; homologous gene to *Dcp-1*), *casp-10* (*caspase-10-like*; homologous gene to *Dredd*) were predicted from *Drosophila melanogaster*. Sample sizes are ranging between six and ten pools of three individual honeybee midguts (see also [S3 Table](#)). Significance between treatment groups ***, $P < 0.001$.

doi:10.1371/journal.pone.0140174.g002

1 p.i. over all treatment groups. Although both JNK/*bsk* and *p53* are important proapoptotic factors [28, 29], the core of the apoptotic-machine consists of caspases that destroy essential cell proteins that initiate apoptosis [20, 30, 31]. Surprisingly, *caspase-10-like* (homologous gene to *Dredd* in *D. melanogaster*) was slightly higher expressed in the sensitive honeybees regardless the treatment than in the tolerant honeybees on day 6 p.i. (*casp-10*, LM: honeybee strain, 0.286 ± 0.093 , $P < 0.025$; Fig 2). The *caspase-2-like* gene (possible homologous gene to *Dronc* in *D. melanogaster*), however, was not differentially expressed between treatment groups (*casp-2*, LM: 0.074 ± 0.074 , $P < 0.290$; Fig 2). In *D. melanogaster* the apical cell-death caspase DRONC, mediated by the adapter ARK (homolog to the apoptotic protease-activating factor 1, *Apaf 1*), plays a central role due to its chronic activation in many cells [20]. This may also be the case for *casp-2* in honeybees as we measured relatively high expression levels in all treatment groups. Nevertheless, cells survive due to the DIAP1 expression, which suppresses DRONC and other activated caspases [20, 32]. Interestingly, we found an interaction between honeybee strain and treatment on day 6 p.i. (LM: honeybee strain \times treatment: 0.751 ± 0.260 , $P < 0.038$; Fig 2), whereby *iap-2* (predicted homologous gene of *Diap-1* in *D. melanogaster*) expression appeared to be tenfold increased on average in sensitive honeybees when infected with *N. ceranae* (Tukey's HSD: 0.761 ± 0.198 , $P < 0.001$) but we found no alterations in *iap-2* expression level in tolerant honeybees (Tukey's HSD: 0.010 ± 0.169 , $P < 0.980$).

Although IAPs are also known to play a regulatory role in pathogen-sensing pathways and induction of the innate immune system [20, 32–34], we speculate that the up-regulation of *iap-2* in *Nosema* infected sensitive honeybees might be rather involved in cell survival, because previous studies reported that *N. ceranae* causes immunosuppression in sensitive honeybees [16, 35, 36]. Furthermore, this up-regulation of *iap-2* only in *Nosema* infected sensitive honeybees would also plausibly explain the reduced apoptosis activity in sensitive honeybees in our TUNEL assays

and supports previous findings [5]. Unfortunately, we did not measure the expression levels of potential apoptosis inducing proteins such as Reaper, Hid and Grim (known as RHG proteins), which can negatively regulate DIAP1 activity in *D. melanogaster* [20, 34]. Nevertheless, *in vitro* studies have demonstrated in some more detail that infections with protozoans such as *Toxoplasma gondii* [37], *Cryptosporidium parvum* [38] as well as bacteria *Shigella flexneri* [39] and *Neisseria gonorrhoeae* [40] elicit up-regulation of *iap* genes and result in the inhibition of host cell apoptosis in mammalian host cell cultures. The activation of the key transcription factor NF- κ B was shown to correlate with *iap* transcription [37, 40] and may also be the case in our *in situ* honeybee-*Nosema* system and might be triggered by elevated *Nosema* HSP70 levels [41, 42]. The capability to retain high apoptotic activity in spite of a *Nosema* infection might explain why the tolerant honeybees can overcome the infection and eliminate the disease from the colony [15, 16]. Workers presumably simply clear the infection by removing those apoptotic infected cells on defecation flights. In contrast sensitive honeybees might be likely to retain the infection in the gut epithelium for much longer time. The dynamics of intestine epithelium development [26] may also provide an alternative explanation for the age-dependent *Nosema* susceptibility in *Bombus terrestris* and honeybees previously exclusively attributed to age polyethism [8, 43].

Our results provide a snap shot of host-parasite co-evolution, where artificial selection of the honeybee host has presumably accelerated a counter adaptation towards *Nosema*. Irrespective of the actual molecular mechanisms, this study does not only highlight the central role of apoptosis for host immunity in general but also shows the importance of its manipulation for intracellular pathogens. Understanding the molecular dialogue between infecting pathogen and host cell might not only be interesting for evolutionary biology and parasitology, but may also provide novel perspectives for effective immunological strategies in the treatment of animal and human diseases by interfering into the regulatory machinery of apoptosis.

Supporting Information

S1 Table. Primer sequences used in qPCR.

(DOCX)

S2 Table. Data of the estimation of the apoptosis rate in the posterior end of honeybee mid-guts for days one and six post infection (d.p.i).

(DOCX)

S3 Table. Relative gene expression data of predicated candidate genes (*bsk*, *p53*, *iap-2*, *casp-2* and *casp-10*).

(DOCX)

Acknowledgments

We thank Robin Crewe, Robert Paxton and anonymous referees for comments on a previous version of the manuscript, and Panagiotis Theodorou for statistical advice. This study was funded by the Deutsche Forschungsgemeinschaft through the priority programme "Host-Parasite Coevolution–Rapid Reciprocal Adaptation and its Genetic Basis" (SPP 1399, Grant number MO373/26-2) to RFAM and a scholarship from The Foundation for the Preservation of Honey Bees, Inc. to CK.

Author Contributions

Conceived and designed the experiments: CK YLC PK RFAM. Performed the experiments: CK CD. Analyzed the data: CK SE OL TM MW RFAM. Contributed reagents/materials/analysis tools: RFAM YLC CK PK TM. Wrote the paper: CK RFAM.

References

1. Bruchhaus I, Roeder T, Rennenberg A, Heussler VT. Protozoan parasites: programmed cell death as a mechanism of parasitism. *Trends Parasitol.* 2007; 23(8):376–83. doi: [10.1016/j.pt.2007.06.004](https://doi.org/10.1016/j.pt.2007.06.004) PMID: [17588817](https://pubmed.ncbi.nlm.nih.gov/17588817/)
2. Faherty CS, Maurelli AT. Staying alive: bacterial inhibition of apoptosis during infection. *Trends Microbiol.* 2008; 16(4):173–80. doi: [10.1016/j.tim.2008.02.001](https://doi.org/10.1016/j.tim.2008.02.001) PMID: [18353648](https://pubmed.ncbi.nlm.nih.gov/18353648/)
3. Mocarski ES, Upton JW, Kaiser WJ. Viral infection and the evolution of caspase 8-regulated apoptotic and necrotic death pathways. *Nat Rev Immunol.* 2012; 12(2):79–88. doi: [10.1038/nri3131](https://doi.org/10.1038/nri3131)
4. Del Aguila C, Izquierdo F, Granja AG, Hurtado C, Fenoy S, Fresno M, et al. Encephalitozoon microsporidia modulates p53-mediated apoptosis in infected cells. *Int J Parasitol.* 2006; 36(8):869–76. doi: [10.1016/j.ijpara.2006.04.002](https://doi.org/10.1016/j.ijpara.2006.04.002) PMID: [16753166](https://pubmed.ncbi.nlm.nih.gov/16753166/)
5. Higes M, Juarranz A, Dias-Almeida J, Lucena S, Botias C, Meana A, et al. Apoptosis in the pathogenesis of *Nosema ceranae* (Microsporidia: Nosematidae) in honey bees (*Apis mellifera*). *Environ Microbiol Rep.* 2013; 5(4):530–6. doi: [10.1111/1758-2229.12059](https://doi.org/10.1111/1758-2229.12059) PMID: [23864567](https://pubmed.ncbi.nlm.nih.gov/23864567/)
6. Scanlon M, Leitch GJ, Shaw AP, Moura H, Visvesvara GS. Susceptibility to apoptosis is reduced in the microsporidia-infected host cell. *J Eukaryot Microbiol.* 1999; 46(5):34S–5S. PMID: [10519237](https://pubmed.ncbi.nlm.nih.gov/10519237/)
7. Keeling P. Five Questions about Microsporidia. *PLoS Pathog.* 2009; 5(9). doi: [10.1371/journal.ppat.1000489](https://doi.org/10.1371/journal.ppat.1000489) PMID: [19779558](https://pubmed.ncbi.nlm.nih.gov/19779558/)
8. Fürst MA, McMahon DP, Osborne JL, Paxton RJ, Brown MJF. Disease associations between honeybees and bumblebees as a threat to wild pollinators. *Nature.* 2014; 506(7488):364–6. doi: [10.1038/nature12977](https://doi.org/10.1038/nature12977) PMID: [24553241](https://pubmed.ncbi.nlm.nih.gov/24553241/)
9. Higes M, Meana A, Bartolome C, Botias C, Martin-Hernandez R. *Nosema ceranae* (Microsporidia), a controversial 21st century honey bee pathogen. *Environ Microbiol Rep.* 2013; 5(1):17–29. doi: [10.1111/1758-2229.12024](https://doi.org/10.1111/1758-2229.12024) PMID: [23757127](https://pubmed.ncbi.nlm.nih.gov/23757127/)
10. Goulson D, Nicholls E, Botias C, Rotheray EL. Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science.* 2015; 347(6229):1435–45. doi: [10.1126/science.1255957](https://doi.org/10.1126/science.1255957)
11. Aufauvre J, Misme-Aucouturier B, Vigues B, Texier C, Delbac F, Blot N. Transcriptome analyses of the honeybee response to *Nosema ceranae* and insecticides. *PLoS ONE.* 2014; 9(3). doi: [10.1371/journal.pone.0091686](https://doi.org/10.1371/journal.pone.0091686) PMID: [24646894](https://pubmed.ncbi.nlm.nih.gov/24646894/)
12. Dussaubat C, Brunet JL, Higes M, Colbourne JK, Lopez J, Choi JH, et al. Gut pathology and responses to the microsporidium *Nosema ceranae* in the honey bee *Apis mellifera*. *PLoS ONE.* 2012; 7(5). doi: [10.1371/journal.pone.0037017](https://doi.org/10.1371/journal.pone.0037017) PMID: [22623972](https://pubmed.ncbi.nlm.nih.gov/22623972/)
13. Holt HL, Aronstein KA, Grozinger CM. Chronic parasitization by *Nosema* microsporidia causes global expression changes in core nutritional, metabolic and behavioral pathways in honey bee workers (*Apis mellifera*). *BMC Genomics.* 2013; 14. doi: [10.1186/1471-2164-14-799](https://doi.org/10.1186/1471-2164-14-799) PMID: [24245482](https://pubmed.ncbi.nlm.nih.gov/24245482/)
14. Gisder S, Mockel N, Linde A, Genersch E. A cell culture model for *Nosema ceranae* and *Nosema apis* allows new insights into the life cycle of these important honey bee-pathogenic microsporidia. *Environ Microbiol.* 2011; 13(2):404–13. doi: [10.1111/j.1462-2920.2010.02346.x](https://doi.org/10.1111/j.1462-2920.2010.02346.x) PMID: [20880328](https://pubmed.ncbi.nlm.nih.gov/20880328/)
15. Hatjina F, Bienkowska M, Charistos L, Chlebo R, Costa C, Dražić MM, et al. A review of methods used in some European countries for assessing the quality of honey bee queens through their physical characters and the performance of their colonies. *J Apicult Res.* 2014; 53(3):337–63. doi: [10.3896/IBRA.1.53.3.02](https://doi.org/10.3896/IBRA.1.53.3.02)
16. Huang Q, Kryger P, Le Conte Y, Moritz RFA. Survival and immune response of drones of a Nosemosis tolerant honey bee strain towards *N. ceranae* infections. *J Invertebr Pathol.* 2012; 109(3):297–302. doi: [10.1016/j.jip.2012.01.004](https://doi.org/10.1016/j.jip.2012.01.004) PMID: [22285444](https://pubmed.ncbi.nlm.nih.gov/22285444/)
17. Fries I, Chauzat MP, Chen YP, Doublet V, Genersch E, Gisder S, et al. Standard methods for *Nosema* research. *J Apicult Res.* 2013; 52(1). doi: [10.3896/ibra.1.52.1.14](https://doi.org/10.3896/ibra.1.52.1.14)
18. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. *Nat Methods.* 2012; 9(7):671–5. doi: [10.1038/nmeth.2089](https://doi.org/10.1038/nmeth.2089) PMID: [22930834](https://pubmed.ncbi.nlm.nih.gov/22930834/)
19. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol.* 1990; 215(3):403–10. doi: [10.1006/jmbi.1990.9999](https://doi.org/10.1006/jmbi.1990.9999) PMID: [2231712](https://pubmed.ncbi.nlm.nih.gov/2231712/)
20. Hay BA, Huh JR, Guo M. The genetics of cell death: Approaches, insights and opportunities in *Drosophila*. *Nat Rev Genet.* 2004; 5(12):911–22. doi: [10.1038/nrg1491](https://doi.org/10.1038/nrg1491) PMID: [15573123](https://pubmed.ncbi.nlm.nih.gov/15573123/)
21. Evans JD. Beepath: An ordered quantitative-PCR array for exploring honey bee immunity and disease. *J Invertebr Pathol.* 2006; 93(2):135–9. doi: [10.1016/j.jip.2006.04.004](https://doi.org/10.1016/j.jip.2006.04.004) PMID: [16737710](https://pubmed.ncbi.nlm.nih.gov/16737710/)
22. Ramakers C, Ruijter JM, Deprez RHL, Moorman AFM. Assumption-free analysis of quantitative real-time polymerase chain reaction (PCR) data. *Neurosci Lett.* 2003; 339(1):62–6. doi: [10.1016/s0304-3940\(02\)01423-4](https://doi.org/10.1016/s0304-3940(02)01423-4) PMID: [12618301](https://pubmed.ncbi.nlm.nih.gov/12618301/)

23. Pfaffl MW, Tichopad A, Prgomet C, Neuvians TP. Determination of stable housekeeping genes, differentially regulated target genes and sample integrity: BestKeeper—Excel-based tool using pair-wise correlations. *Biotechnol Lett*. 2004; 26(6):509–15. doi: [10.1023/b:bile.0000019559.84305.47](https://doi.org/10.1023/b:bile.0000019559.84305.47) PMID: [15127793](https://pubmed.ncbi.nlm.nih.gov/15127793/)
24. R Foundation for Statistical Computing R: a language and environment for statistical computing (25 September 2013).
25. Zuur A, Ieno EN, Walker N, Saveliev AA, Smith GM. *Mixed effects models and extensions in ecology with R*: Springer; 2009.
26. Pipan N, Rakovec V. Cell death in the midgut epithelium of the worker honey bee (*Apis mellifera carnica*) during metamorphosis. *Zoomorphologie*. 1980; 94(2):217–24. doi: [10.1007/bf01081936](https://doi.org/10.1007/bf01081936)
27. Seeley TD, Nowicke JW, Meselson M, Guillemin J, Akratanakul P. Yellow Rain. *Sci Am*. 1985; 253(3):122–31.
28. Kanda H, Miura M. Regulatory roles of JNK in programmed cell death. *J Biochem*. 2004; 136(1):1–6. doi: [10.1093/jb/mvh098](https://doi.org/10.1093/jb/mvh098) PMID: [15269233](https://pubmed.ncbi.nlm.nih.gov/15269233/)
29. Steller H. *Drosophila* p53: meeting the Grim Reaper. *Nat Cell Biol*. 2000; 2(6):E100–E2. doi: [10.1038/35014093](https://doi.org/10.1038/35014093) PMID: [10854336](https://pubmed.ncbi.nlm.nih.gov/10854336/)
30. Thornberry NA, Lazebnik Y. Caspases: Enemies within. *Science*. 1998; 281(5381):1312–6. doi: [10.1126/science.281.5381.1312](https://doi.org/10.1126/science.281.5381.1312) PMID: [9721091](https://pubmed.ncbi.nlm.nih.gov/9721091/)
31. Cohen GM. Caspases: the executioners of apoptosis. *Biochem J*. 1997; 326:1–16. PMID: [9337844](https://pubmed.ncbi.nlm.nih.gov/9337844/)
32. Deveraux QL, Reed TC. IAP family proteins—suppressors of apoptosis. *Gene Dev*. 1999; 13(3):239–52. doi: [10.1101/gad.13.3.239](https://doi.org/10.1101/gad.13.3.239) PMID: [9990849](https://pubmed.ncbi.nlm.nih.gov/9990849/)
33. Hay BA, Guo M. Caspase-dependent cell death in *Drosophila*. *Annu Rev Cell Dev Biol*. 2006; 22:623–50. doi: [10.1146/annurev.cellbio.21.012804.093845](https://doi.org/10.1146/annurev.cellbio.21.012804.093845) PMID: [16842034](https://pubmed.ncbi.nlm.nih.gov/16842034/)
34. Huh JR, Foe I, Muro I, Chen CH, Seol JH, Yoo SJ, et al. The *Drosophila* inhibitor of apoptosis (IAP) DIAP2 is dispensable for cell survival, required for the innate immune response to gram-negative bacterial infection, and can be negatively regulated by the Reaper/Hid/Grim family of IAP-binding apoptosis inducers. *J Biol Chem*. 2007; 282(3):2056–68. doi: [10.1074/jbc.M608051200](https://doi.org/10.1074/jbc.M608051200) PMID: [17068333](https://pubmed.ncbi.nlm.nih.gov/17068333/)
35. Antunez K, Martin-Hernandez R, Prieto L, Meana A, Zunino P, Higes M. Immune suppression in the honey bee (*Apis mellifera*) following infection by *Nosema ceranae* (Microsporidia). *Environ Microbiol*. 2009; 11(9):2284–90. doi: [10.1111/j.1462-2920.2009.01953.x](https://doi.org/10.1111/j.1462-2920.2009.01953.x) PMID: [19737304](https://pubmed.ncbi.nlm.nih.gov/19737304/)
36. Chaimanee V, Chantawannakul P, Chen Y, Evans JD, Pettis JS. Differential expression of immune genes of adult honey bee (*Apis mellifera*) after inoculated by *Nosema ceranae*. *J Insect Physiol*. 2012; 58(8):1090–5. doi: [10.1016/j.jinsphys.2012.04.016](https://doi.org/10.1016/j.jinsphys.2012.04.016) PMID: [22609362](https://pubmed.ncbi.nlm.nih.gov/22609362/)
37. Molestina RE, Payne TM, Coppens I, Sinai AP. Activation of NF-kappa B by *Toxoplasma gondii* correlates with increased expression of antiapoptotic genes and localization of phosphorylated I kappa B to the parasitophorous vacuole membrane. *J Cell Sci*. 2003; 116(21):4359–71. doi: [10.1242/jcs.00683](https://doi.org/10.1242/jcs.00683)
38. Liu J, Enomoto S, Lancto CA, Abrahamsen MS, Rutherford MS. Inhibition of apoptosis in *Cryptosporidium parvum*-infected intestinal epithelial cells is dependent on survivin. *Infect Immun*. 2008; 76(8):3784–92. doi: [10.1128/iai.00308-08](https://doi.org/10.1128/iai.00308-08) PMID: [18519556](https://pubmed.ncbi.nlm.nih.gov/18519556/)
39. Pedron T, Thibault C, Sansonetti PJ. The invasive phenotype of *Shigella flexneri* directs a distinct gene expression pattern in the human intestinal epithelial cell line Caco-2. *J Biol Chem*. 2003; 278(36):33878–86. doi: [10.1074/jbc.M303749200](https://doi.org/10.1074/jbc.M303749200) PMID: [12813033](https://pubmed.ncbi.nlm.nih.gov/12813033/)
40. Binnicker MJ, Williams RD, Apicella MA. Infection of human urethral epithelium with *Neisseria gonorrhoeae* elicits an upregulation of host anti-apoptotic factors and protects cells from staurosporine-induced apoptosis. *Cell Microbiol*. 2003; 5(8):549–60. doi: [10.1046/j.1462-5822.2003.00300.x](https://doi.org/10.1046/j.1462-5822.2003.00300.x) PMID: [12864814](https://pubmed.ncbi.nlm.nih.gov/12864814/)
41. Vidau C, Panek J, Texier C, Biron DG, Belzunces LP, Le Gall M, et al. Differential proteomic analysis of midguts from *Nosema ceranae*-infected honeybees reveals manipulation of key host functions. *J Invertebr Pathol*. 2014; 121:89–96. doi: [10.1016/j.jip.2014.07.002](https://doi.org/10.1016/j.jip.2014.07.002) PMID: [25038465](https://pubmed.ncbi.nlm.nih.gov/25038465/)
42. Joly A-L, Wettstein G, Mignot G, Ghiringhelli F, Garrido C. Dual Role of Heat Shock Proteins as Regulators of Apoptosis and Innate Immunity. *J Innat Immun*. 2010; 2(3):238–47. doi: [10.1159/000296508](https://doi.org/10.1159/000296508)
43. Smart MD, Sheppard WS. *Nosema ceranae* in age cohorts of the western honey bee (*Apis mellifera*). *J Invertebr Pathol*. 2012; 109(1):148–51. doi: [10.1016/j.jip.2011.09.009](https://doi.org/10.1016/j.jip.2011.09.009) PMID: [22001631](https://pubmed.ncbi.nlm.nih.gov/22001631/)

Supplementary Material

S1 Table. Primer sequences used in qPCR.

Gene	Predicted in <i>Apis mellifera</i>	Homolog in <i>D. melanogaster</i>	GenBank assacion no.	Primer sequences	References
<i>ark</i>	<i>apoptotic protease-activating factor (Apaf) 1-like</i>	<i>Apaf-1-related-killer</i> , isoform B	XR_120278.1	forward, CACCCACTCTTCCTCCAACAT; reverse, ATATTTCTTCATCAACTGGACGTTT	this study
<i>bsk</i>	<i>JNK MAP kinase basket</i>	<i>basket</i> , isoform B	XM_392806.4	forward, ACAAGATAATCGAGCAACTGGGA; reverse, CCTGTTCTCCACGTAATTCCTGA	this study
<i>bcl-2</i>	<i>bcl-2-related ovarian killer protein</i>	<i>Bcl-2 related protein (buffy)</i>	XM_395083.4	forward, GCATTGCCGATGCCTGAAAA; reverse, TCTGCGATAAGGTTGGCCTG	this study
<i>casp-10</i>	<i>caspase-10-like</i> (LOC724930)	<i>death related ced-3/Nedd2-like protein (Dredd)*</i>	XM_001120830.2	forward, GCGTCATAAAGAAAAAGGATCATGG; reverse, CCCCTACTGCATCAATTGTTTT	this study
<i>casp-1</i>	<i>caspase-1-like</i> (LOC412235)	<i>Dredd*</i> , <i>Nedd2-like caspase (Dronc)*</i>	XM_395697.4	forward, CCGCGAGTGGTATTTCTCCA; reverse, ACTGATGGTCAACCAGCTTCT	this study
<i>casp-2</i>	<i>caspase-2-like</i>	<i>Dronc*</i>	XM_003249125.1	forward, TCACGTTGGAAGACAAATCTCAC; reverse, AATGCAAAAGGTCCCCGTGT	this study
<i>casp</i>	<i>caspase-like</i> (LOC411381)	<i>Drice*</i> ; <i>Dronc*</i>	XM_394855.3	forward, ACAGTATGACGGATGCCGAG; reverse, GGCGAATTTCTTCTGTGCATGT	this study
<i>iap-2</i>	<i>inhibitor of apoptosis protein 2</i>	<i>apoptosis 1 inhibitor (Diap-1)</i>	XM_396819.4	forward, TTCAACGTGATGATGAAGTACAATG; reverse, TGGATCATCACCAAGCTCCC	this study
<i>p53</i>	<i>tumor protein p73-like</i>	<i>tumor protein p53</i>	XM_003249744.1	forward, TGGAAATATGACCACTACGAATGGA; reverse, AGGGTACAAATCCAATATCCTTCAC	this study
<i>arp 1</i>	<i>actin related protein 1</i>		NM_001185146.1	forward, TTGTATGCCAACACTGTCCTTT; reverse, TGGCGCGATGATCTTAATTT	[1]
<i>RPS5</i>	<i>ribosomal protein S5a</i>		XM_624081.3	forward, AATTATTTGGTCGCTGGAATTG; reverse, TAACGTCCAGCAGAATGTGGTA	[1]

* Due to high sequence similarities, a distinction between caspases, as known for *D. melanogaster*, was difficult in *Apis mellifera* based on gene homologies.

Additional References

1. Evans JD (2006) Beepath: An ordered quantitative-PCR array for exploring honey bee immunity and disease. *J Invertebr Pathol* 93: 135-139.

S2 Table. Data of the estimation of the apoptosis rate in the posterior end of honeybee midguts for days one and six post infection (d.p.i). The apoptosis ratio was calculated as TUNEL+ve / DAPI+ve nuclei using ImageJ. For day 1 p.i.: $n = 3_{SC}, 5_{SN}, 5_{TC}, 6_{TN}$ and for day 6 p.i.: $n = 5_{SC}, 7_{SN}, 9_{TC}, 7_{TN}$.

#ID	treatment groups	treatment	group	d.p.i.	replicate	number of nuclei	apoptosis ratio
1	SN	infected	sensitive	six	1	354	0.362
2	SN	infected	sensitive	six	1	457	0.225
3	SN	infected	sensitive	six	1	273	0.348
4	SC	control	sensitive	six	1	328	0.451
5	SC	control	sensitive	six	1	342	0.471
6	SC	control	sensitive	six	1	384	0.477
7	TN	infected	tolerant	six	1	515	0.373
8	TN	infected	tolerant	six	1	624	0.404
9	TN	infected	tolerant	six	1	516	0.341
10	TN	infected	tolerant	six	1	538	0.394
11	TC	control	tolerant	six	1	345	0.304
12	TC	control	tolerant	six	1	293	0.294
13	TC	control	tolerant	six	1	283	0.339
14	SN	infected	sensitive	six	3	483	0.178
15	SN	infected	sensitive	six	3	362	0.224
16	SC	control	sensitive	six	3	376	0.314
17	SC	control	sensitive	six	3	362	0.304
18	SC	control	sensitive	six	3	407	0.405
20	TN	infected	tolerant	six	3	341	0.375
21	TC	control	tolerant	six	3	467	0.272
22	TC	control	tolerant	six	3	342	0.398
24	SN	infected	sensitive	six	2	429	0.054
25	SN	infected	sensitive	six	2	359	0.167
26	SC	control	sensitive	six	2	417	0.223
27	SC	control	sensitive	six	2	267	0.228
28	SC	control	sensitive	six	2	296	0.375
29	TN	infected	tolerant	six	2	297	0.387
30	TN	infected	tolerant	six	2	436	0.374
31	TC	control	tolerant	six	2	371	0.251
32	TC	control	tolerant	six	2	431	0.439
33	SN	infected	sensitive	one	1	70	0.629
34	SN	infected	sensitive	one	1	343	0.615
35	SN	infected	sensitive	one	3	288	0.580
43	TC	control	tolerant	one	3	297	0.401
44	SN	infected	sensitive	one	2	204	0.515
45	SN	infected	sensitive	one	2	306	0.533
46	SC	control	sensitive	one	2	364	0.492
47	SC	control	sensitive	one	2	187	0.561
48	SC	control	sensitive	one	3	194	0.500
49	TN	infected	tolerant	one	2	175	0.617
50	TN	infected	tolerant	one	2	372	0.540
51	TN	infected	tolerant	one	2	91	0.835
52	TC	control	tolerant	one	2	310	0.529

53	TC	control	tolerant	one	2	165	0.727
57	TN	infected	tolerant	one	3	174	0.540
58	TN	infected	tolerant	one	1	253	0.443
59	TN	infected	tolerant	one	1	235	0.447
60	TC	control	tolerant	one	1	186	0.608
62	TC	control	tolerant	one	1	218	0.468

S3 Table. Relative gene expression data of predicated candidate genes (*bsk*, *p53*, *iap-2*, *casp-2* and *casp-10*) involved in apoptosis in *Nosema ceranae* infected *Nosema* sensitive (SN) and tolerant (TN) honeybees, and their uninfected controls (SC, TC) after one and six days post infection (d.p.i.). Sample ID represents pools of midguts from three individual honeybees.

#ID	treatment groups	treatment	group	d.p.i.	replicate	<i>bsk</i>	<i>p53</i>	<i>iap-2</i>	<i>casp-2</i>	<i>casp-10</i>
1	TN	infected	tolerant	one	1	0,009	NA	0,010	0,505	0,003
2	TC	control	tolerant	one	1	0,003	0,016	0,012	0,736	0,009
4	SC	control	sensitive	one	1	0,009	0,001	0,001	0,126	NA
5	TN	infected	tolerant	one	2	0,017	0,001	0,002	0,168	0,002
6	TC	control	tolerant	one	2	0,008	0,016	0,024	0,585	0,010
7	TC	control	tolerant	one	3	0,011	0,002	0,002	0,056	0,003
8	SN	infected	sensitive	one	2	0,004	0,008	0,011	1,193	0,010
9	SC	control	sensitive	one	2	0,009	0,002	0,006	0,415	0,003
10	TN	infected	tolerant	one	3	0,006	0,016	0,015	0,432	0,010
11	SN	infected	sensitive	one	3	0,007	0,021	0,088	3,424	0,013
12	SC	control	sensitive	one	3	0,009	NA	0,002	0,086	0,004
23	TN	infected	tolerant	six	1	0,072	0,009	0,033	0,454	0,028
24	TC	control	tolerant	six	1	0,708	0,027	0,034	NA	0,017
25	SN	infected	sensitive	six	1	0,132	0,020	1,471	0,680	0,035
26	SC	control	sensitive	six	1	0,018	0,010	0,015	0,144	NA
27	TN	infected	tolerant	six	1	0,276	NA	NA	0,829	NA
28	TC	control	tolerant	six	2	NA	0,025	0,028	1,049	0,019
29	TN	infected	tolerant	six	3	0,074	0,032	0,024	0,717	0,006
30	TC	control	tolerant	six	3	0,072	0,020	0,017	0,612	0,005
31	SN	infected	sensitive	six	3	NA	NA	NA	NA	NA
32	SC	control	sensitive	six	3	NA	0,024	0,034	0,219	NA
33	TN	infected	tolerant	six	2	0,065	0,031	0,028	0,617	0,016
34	TC	control	tolerant	six	2	0,015	NA	0,014	NA	NA
35	SN	infected	sensitive	six	2	0,033	0,002	0,307	0,128	0,001
36	SC	control	sensitive	six	2	0,048	0,001	NA	0,446	0,012
37	TN	infected	tolerant	one	1	0,034	0,017	0,017	0,243	0,015
38	TC	control	tolerant	one	1	0,015	0,014	0,014	0,489	0,010
39	SN	infected	sensitive	one	1	0,038	0,010	0,024	0,445	0,014
40	SC	control	sensitive	one	1	0,020	0,014	0,029	0,443	0,017
41	TN	infected	tolerant	six	1	0,022	NA	0,031	0,267	0,018
42	TC	control	tolerant	six	1	0,032	0,019	0,027	0,374	0,009
43	SN	infected	sensitive	six	1	0,034	0,016	0,030	0,411	0,471
44	SC	control	sensitive	six	1	0,171	0,019	0,088	NA	0,056
45	TN	infected	tolerant	six	3	0,078	0,130	0,037	0,353	0,014
46	TC	control	tolerant	six	3	NA	NA	0,032	NA	NA
47	SN	infected	sensitive	six	3	0,172	NA	NA	0,420	0,017
48	SC	control	sensitive	six	3	0,042	NA	0,039	0,083	0,011
49	TC	control	tolerant	six	1	NA	0,002	0,002	0,756	0,020
51	TC	control	tolerant	six	3	0,100	NA	0,021	NA	0,014
52	TN	infected	tolerant	six	2	0,104	0,045	0,033	0,305	0,033
53	TN	infected	tolerant	six	2	NA	NA	NA	NA	NA
54	TN	infected	tolerant	six	1	0,307	0,041	0,033	0,936	NA
55	SC	control	sensitive	six	3	0,152	0,048	0,034	0,805	0,064

56	SC	control	sensitive	six	2	0,234	0,057	0,021	NA	0,017
57	SC	control	sensitive	six	2	NA	NA	NA	NA	NA
58	TC	control	tolerant	six	2	0,054	0,075	0,004	0,685	0,016
59	SN	infected	sensitive	six	2	NA	0,029	2,035	NA	NA
60	SN	infected	sensitive	six	3	NA	NA	NA	NA	NA
61	SN	infected	sensitive	six	2	0,023	0,032	0,030	0,062	0,049
62	TN	infected	tolerant	six	3	0,009	NA	0,043	0,629	0,016
63	TC	control	tolerant	six	3	0,448	NA	0,045	0,291	0,014
64	TC	control	tolerant	six	2	0,020	0,060	NA	0,773	0,032
65	SC	control	sensitive	six	1	0,032	0,022	0,030	0,349	0,022
66	SN	infected	sensitive	six	1	0,042	0,047	0,917	0,059	0,696
84	TN	infected	tolerant	one	2	0,052	0,009	0,017	0,357	0,013
70	TN	infected	tolerant	one	3	0,082	0,014	NA	0,333	0,014
71	SC	control	sensitive	one	3	0,059	0,007	0,019	0,294	0,020
72	SN	infected	sensitive	one	3	0,041	0,012	0,012	0,268	0,013
73	TC	control	tolerant	one	3	0,048	0,019	0,013	0,305	NA
74	SN	infected	sensitive	one	2	0,023	0,015	0,008	0,191	NA
75	SC	control	sensitive	one	2	0,016	0,007	0,010	0,262	0,010
76	TN	infected	tolerant	one	2	0,074	0,008	0,014	0,348	0,009
77	SC	control	sensitive	one	2	0,043	0,009	0,017	0,761	0,020
78	TN	infected	tolerant	one	3	0,085	0,023	0,018	0,553	0,007
79	TN	infected	tolerant	one	2	0,001	0,001	0,000	NA	0,000
80	SN	infected	sensitive	one	1	0,021	0,010	0,020	0,421	0,015
81	TC	control	tolerant	one	2	0,047	0,010	0,014	0,212	0,006
82	SN	infected	sensitive	one	2	0,047	0,010	NA	0,285	0,013
83	TC	control	tolerant	one	3	0,060	0,016	0,044	0,318	0,011
85	TC	control	tolerant	one	2	0,026	0,007	0,016	NA	0,006
86	SN	infected	sensitive	one	3	0,001	0,001	0,000	0,010	0,001
87	SC	control	sensitive	one	3	0,037	0,011	NA	0,443	0,012
88	TN	infected	tolerant	one	1	0,044	0,024	0,029	0,455	0,010
89	SC	control	sensitive	one	3	0,003	0,001	0,001	0,016	0,000