

Seroprevalence of Newcastle disease and other infectious diseases in backyard chickens at markets in Eastern Shewa Zone, Ethiopia

H. Chaka,^{*†1}, F. Goutard,[‡] S. P. R. Bisschop,^{*} and P. N. Thompson^{*}

* Department of Production Animal Studies, Faculty of Veterinary Science, University of Pretoria, South Africa and, † National Animal Health Diagnostic and Investigation Center, Sebata, Ethiopia; and ‡ *Centre de coopération internationale en recherche agronomique pour le développement (CIRAD), Département ES, Unité AGIRs, Montpellier, France*

ABSTRACT This study was conducted to estimate the seroprevalence of Newcastle disease (ND), *Pasteurella multocida* (PM) infection, *Mycoplasma gallisepticum* (MG) infection and infectious bursal disease (IBD), and to assess the level of concurrent seropositivity during the dry and wet seasons of the year 2010. A total of 234 and 216 sera were collected during the dry and wet seasons respectively from unvaccinated backyard chickens at four live poultry markets in two Woredas (Districts) of Eastern Shewa Zone, Ethiopia, and were tested using commercial enzyme-linked immunosorbent assay (ELISA) kits. The overall seroprevalence of ND, PM, MG, and IBD was 5.9%, 66.2%, 57.7% and 91.9% respectively during the dry season, and 6.0%, 63.4%, 78.7% and 96.3% respectively during the wet season. The seroprevalence of MG was higher ($P < 0.001$) during the wet season than during the dry season and higher ($P = 0.002$) in Adami-Tulu-Jido-Kombolcha (ATJK) Woreda (74%) than in Ada'a Woreda (60%). Area and season had no significant effect on the seroprevalence of ND, IBD and PM, indicating the widespread presence of those pathogens throughout the year in the study area. Of all the chickens tested, 85.6% had antibodies concurrently to more than

¹corresponding author: hasscha@yahoo.com

Tel: +251 113 383561 Fax: +251 113 380220

one of the pathogens investigated. Birds were concurrently seropositive to more diseases during the wet season (median = 3) than during the dry season (median = 2) ($P = 0.002$). As serology is not able to distinguish between strains, further studies are warranted to better understand the circulating strains, their interactions and their economic effect on backyard poultry production in Ethiopia.

Key words: Newcastle disease, infectious disease, backyard chickens, Ethiopia

INTRODUCTION

Like elsewhere in the developing world, backyard poultry rearing is a common practice in rural Ethiopia. Village backyard poultry, characterized by traditional production methods and local breeds, represents 98% of the total Ethiopian poultry population of 38 million (CSA, 2009). The sector provides eggs and poultry meat to most rural and many urban consumers (Tadelle, 1996). However, the productivity of backyard chickens is hampered by several factors, including a variety of infectious diseases. In Ethiopia an estimated 40-60% of newly hatched chicks die before reaching maturity, mainly due to disease and predation (Tadelle et al., 2003). An ongoing questionnaire-based survey (unpublished data) has revealed that farmers in Eastern Shewa Zone consider any disease that causes moderate to high mortality to be Newcastle disease (**ND**), known locally as *Fengil or Fenqil*, as also reported elsewhere (Sonaiya and Swan, 2004). It is therefore likely that many other infectious diseases have, for a long time, been described incorrectly as ND. Mortalities of backyard chickens may also result from other viral or bacterial diseases, either individually or concurrently. Several viral and bacterial infectious diseases, including ND, *Mycoplasma gallisepticum* (**MG**) infection, infectious bursal disease (**IBD**) and Marek's disease have been described in commercial

poultry farms in Ethiopia (Alamargot, 1987; Nasser et al., 2000; Lobago and Woldemeskel, 2004; Zeleke et al., 2005a,b; Chanie et al., 2009), but their occurrence in backyard poultry production systems has rarely been documented. The key to increasing profitability of backyard poultry production is to know which diseases are prevalent in an area (Bell, 2009). Workers in other African countries have documented the presence of many viral and bacterial diseases in backyard production systems (Bell et al., 1990; Kelly et al., 1994; Chrysostome et al., 1995; Idi et al., 1999; Orjaka et al., 1999; Muhairwa et al., 2001; Ndanyi, 2003; Mushi et al., 2006; Mbuthia et al., 2008). It is therefore also likely that infectious diseases such as ND, IBD, mycoplasmosis, and pasteurellosis could play a role, individually or concurrently, in backyard poultry health in Ethiopia. In addition, backyard poultry could be a potential reservoir of these pathogens which could jeopardize the development of semi-commercial poultry production in Ethiopia.

Newcastle disease is caused by avian paramyxovirus serotype 1 (**APMV-1**), which, with viruses of the other eight APMV serotypes (APMV-2 to APMV-9), have been placed in the genus *Avulavirus*, belonging to the sub-family *Paramyxovirinae*, family *Paramyxoviridae* (OIE, 2009). It has a worldwide distribution and is regarded as one of the most important constraints to the development of backyard poultry production (Alexander et al., 2004; OIE, 2009). Serological and virological evidence has shown the presence of the disease in backyard poultry in many African countries (Bell et al., 1990; Echeonwu et al., 1993; Alders et al., 1994; Chrysostome et al., 1995; Orjaka et al., 1999; Snoeck et al., 2009; Almeida et al., 2009), and the disease has long been known to be endemic in Ethiopia (NVI, 1974). Limited previous serological surveys of ND based on haemagglutination inhibition (HI) titer antibody detection in backyard chickens in Ethiopia have found that 11-38% of chickens had detectable antibodies (Tadesse et al., 2005; Zeleke et al., 2005b; Regasa et al., 2007).

Infectious bursal disease (IBD) is a highly contagious, immunosuppressive infection of immature chickens with a worldwide distribution (Sharma et al., 2000). Two serotypes of IBD virus strains are described: 1 and 2. Serotype 2 strains classified as apathogenic and serotype 1 strain, pathogenic to chickens, classified into several pathotypes, from mild to hypervirulent, according to their virulence (van den Berg et al., 2004). IBD outbreaks amongst backyard chickens have been reported in China (Fa, 1993), Indonesia (Parede, 1992), Vietnam (Vui et al., 2002) and Ecuador (Sonia et al., 2006). IBD was also reported from several countries in Africa, including Zimbabwe (Kelly et al., 1994), Niger (Idi et al., 1999), Kenya (Ndanyi, 2005), Egypt (Azzam et al., 2004), Mauritania (Bell et al., 1990) and Botswana (Mushi et al., 2006). The disease was first diagnosed in Ethiopia in 2002 in commercial poultry (Zelege et al., 2005b), and thereafter in a government-owned poultry multiplication centre (Woldemariam and Wossene, 2007) and a commercial broiler farm (Chanie et al., 2009). In Ethiopia, IBD in backyard chickens has been serologically documented from the northwest and central part of the country (Mazengia et al., 2009, Degefu et al., 2010).

Mycoplasma gallisepticum (MG), together with *Escherichia coli*, is the cause of chronic respiratory disease (CRD) in chickens and is the most economically important of the avian *Mycoplasma* spp. (Bradbury, 2001). Mycoplasmas are also well known for their interactions with other infectious agents, such as ND virus, and environmental factors in producing clinical disease (Kleven, 1998). The disease has been reported, by serology or isolation of the agent, in backyard poultry in a few African countries including Niger (Idi et al., 1990), Zimbabwe (Kelly et al., 1994), Botswana (Mushi et al., 1999), Benin (Chrysostome et al., 1995) and Kenya (Ndanyi, 2003), as well as from Ecuador (Sonia et al., 2006) and Argentina (Xavier et al., 2011), and in fancy breed poultry flocks in Switzerland (Wunderwald and Hoop, 2002) where the management system is equivalent of the one in backyard poultry

flocks. In Ethiopia, the disease was reported recently and MG was isolated from commercial broiler farms (Chanie et al., 2009). To our knowledge there are no reports of its presence in backyard chickens in Ethiopia.

Fowl cholera, caused by *Pasteurella multocida* (**PM**), is another disease of significant economic importance with a worldwide distribution (Christensen and Bisgaard, 2000). The characteristic signs of the disease are respiratory rales, coughing and nasal discharge. Fowl cholera is considered a leading killer of domestic and wild birds in Asia (Rimler and Glisson, 1997). However, literature on the epidemiology and significance of infections caused by PM in poultry in developing countries is scanty, with reports in backyard chickens in Thailand (Thitisak et al., 1989), Zimbabwe (Kelly et al., 1994), Tanzania (Muhairwa et al., 2001) and Kenya (Mbuthia et al., 2008). There are no published reports of this disease in either commercial or backyard chickens in Ethiopia.

The paucity of information on the presence and prevalence of the above diseases in backyard chickens may reflect a lack of resources for disease surveillance and control in backyard production systems. In addition, the diagnostic coverage of poultry diseases in Ethiopia is limited to the extent that, even from commercial farms, only a few cases are brought to either the National Animal Health Diagnostic and Investigation Center (**NAHDIC**) or the National Veterinary Institute (**NVI**). Most poultry outbreaks, particularly in more remote parts of the country, remain undiagnosed and dead chickens are simply discarded. Therefore, information on the prevalence and significance of infectious poultry diseases can only readily be obtained through indirect serological studies on apparently healthy chickens. It is difficult to design and implement chicken health development programs without an understanding of the diseases present in the backyard poultry production system. Hence this study was implemented to determine the seroprevalence of ND and other major poultry diseases potentially affecting backyard poultry health in Ethiopia, to determine

the level of concurrent seropositivity to multiple pathogens, and to determine any seasonal or geographic patterns of seroprevalence.

MATERIALS AND METHODS

Study area, study design and sample size

Administratively, Ethiopia is subdivided into Regions which are again sub-divided into Zones and then Woredas (Districts). The study was conducted in the Eastern Shewa Zone of the Oromia Region, in the Woredas of Adami-Tulu-Jido-Konbolcha (**ATJK**) and Ada'a. In order to have animals coming from a diversity of sources, four rural markets (two from each Woreda) were targeted. Accordingly, Dire and Tuludimitu markets in Ada'a Woreda and Adami Tulu and Bulbula markets in ATJK Woreda were selected.

A cross-sectional serological survey was carried out twice during 2010: in January, representing the dry season, and in September, representing the wet season.

Since various prevalences had been reported for ND virus antibodies in previous studies, ranging from 11% (Regasa et al., 2007) to 38% (Tadesse et al., 2005) an expected seroprevalence of 20% was assumed for ND. Since little or no information was available for the other diseases, the same expected seroprevalence as for ND was assumed. The sample size was therefore calculated in order to estimate a prevalence of 20% with 95% confidence and 10% absolute error; this gave a required sample size of 62 birds for each market, during each season. On each sampling occasion a market was visited twice, on two consecutive market days. Apparently healthy chickens, greater than 2 months of age, were purchased, individually identified using a leg band and transported to NAHDIC. In addition, individual sellers were interviewed regarding chicken disease occurrence during the previous six months

(with symptoms including mass mortality, respiratory distress, diarrhoea, ocular and nasal discharges, or nervous signs) and vaccination history in their village of origin.

Sampling procedure and sample analysis

Immediately after arrival at the NAHDIC laboratory, blood samples were collected from the brachial vein in 3 mL disposable syringes and left horizontally for 3 hrs and then vertically for the serum to ooze out. Serum was collected in 2 mL cryovial tubes and kept at -20°C until testing.

Serum samples were analyzed using commercial ELISA kits for the presence of antibodies to ND (Svanovir NDV-Ab Elisa, Svanova Veterinary Diagnostics, Sweden), MG (*Mycoplasma gallisepticum* Ab test kit, Svanova Veterinary Diagnostics, Sweden), IBD (LSI IBD ELISA, Laboratoire Service International, France), and PM (Synbiotics ProFLOK[®] *Pasteurella multocida*, Synbiotics Corporation, U.S.A.), according to the manufacturers' instructions.

The ND virus antibody and MG ELISAs work on the principle of blocking ELISA and are developed to detect specific antibodies against PMV-1 and MG, respectively, in serum. The sample and control optical density (**OD**) values were read using an ELISA reader (Immunoscan, BDSL) at 450 nm. From OD values percent inhibition (**PI**) was calculated for control and test samples using the following formula:

$$PI = [(OD \text{ negative control} - OD \text{ sample}) / OD \text{ positive control}] \times 100 / OD \text{ negative control}$$

The LSIVET IBD and ProFLOK[®] PM antibody ELISA are based on the principle of indirect ELISA. The sample and control OD values were read using an ELISA reader (Immunoscan, BDSL) at 405 nm. For each sample the sample-to-positive (S/P) ratios were calculated from OD values by the formula:

S/P ratio= [Sample OD– negative control mean OD]/ [Positive control mean OD– negative control mean OD].

Data analysis

Data from the laboratory analyses were stored in a spreadsheet, and PI or S/P values were computed as above. The seroprevalence of each disease, for each market, Woreda and season, with binomial exact 95% confidence intervals, were calculated. Seroprevalence was compared between Woredas and seasons using Fisher's exact test. The median number of diseases for which birds were seropositive was compared between seasons using the Wilcoxon rank-sum test. A significance level of $\alpha = 0.05$ was used. Analyses were done using STATA 11 (Stata Corp, College Station, Texas, U.S.A.)

RESULTS

Serum samples from a total of 250 and 229 apparently healthy chickens were collected during the dry and wet seasons, respectively. However, only 234 and 216 sera, respectively, were analyzed because of the limited quantity and quality of serum obtained from some of the birds.

From the interviews we learned that none of the sellers had ever vaccinated their chickens. However, 23.4% and 30% of sellers, during the dry and wet seasons respectively, claimed to have had disease amongst their poultry flocks during the previous six months, with a combination of signs including depression, inappetance, diarrhea, respiratory distress and paralysis in some cases.

Table1 shows the prevalence of ND, PM, MG and IBD antibodies in the different markets and Woredas during the dry season. The seroprevalence of ND ranged from 4.3% (Bulbula) to 7.8% (AdamiTulu), but no significant difference was seen between Woredas. For PM the seroprevalence varied from 56.8% (Tuludimitu) to 78.5% (Bulbula) and was higher in ATJK

(72.4%) than in Ada'a (58%) Woreda ($P = 0.025$). The seroprevalence of MG had a wider variation, from 32.4% (Tuludimitu) to 67.2% (AdamiTulu) and was also significantly higher in ATJK (63.4%) than in Ada'a (50%) Woreda ($P = 0.045$). IBD seroprevalence was high in all the markets, and did not differ significantly between Woredas.

TABLE 1. Seroprevalence of Newcastle disease (ND), *Pasteurella multocida* (PM), *Mycoplasma gallisepticum* (MG) and infectious bursal disease (IBD) in backyard chickens at markets in Eastern Shewa Zone, Ethiopia, during January 2010 (dry season).

Woreda/market	n	Percent seropositive (exact 95% C.I.)			
		ND	PM	MG	IBD
ATJK Woreda	134	5.9 (2.6;10.0) ^A	72.4 (64;79.5) ^B	63.4 (54.6;72.0) ^B	94.8 (89.5;98) ^A
AdamiTulu	64	7.8 (2.6;17.4) ^a	65.6 (52.6;77.0) ^a	67.0 (54.3;78.4) ^a	100 (94.3;100) ^b
Bulbula	70	4.3 (0.9;12.0) ^a	78.5 (67.1;87.5) ^a	60.0 (47.6;71.5) ^a	90.0 (80.4;95.9) ^a
Ada'a Woreda	100	6.0 (2.2;12.6) ^A	58.0 (47.7;67.8) ^A	50.0 (39.8;60.2) ^A	88.0 (79.9;93.6) ^A
Dire	63	6.3 (1.8;15.5) ^a	58.0 (45.6;71.0) ^a	60.3 (47.2;72.4) ^b	87.3 (76.5;94.4) ^a
Tuludimitu	37	5.4 (0.6;18.2) ^a	56.8 (39.5;73.0) ^a	32.4 (18.0;49.8) ^a	89.2 (74.6;97.0) ^a
TOTAL	234	5.9 (3.3; 9.8)	66.2 (59.7;72.0)	57.7 (51.1;64.0)	91.9 (87.6;95.0)

^{A,B} Woredas with different superscripts differ significantly ($P < 0.05$)

^{a,b} Within Woreda, markets with different superscripts differ significantly ($P < 0.05$)

Table 2 shows the prevalence of ND, PM, MG and IBD antibodies in the different markets and Woreda as during the wet season. The seroprevalence of ND showed wider variation between markets than during the dry season, ranging from 3.2% (Bulbula) to 10.3% (Tuludimitu). No significant difference was seen between Woredas but a relatively lower proportion of seropositive chickens were recorded in ATJK (4.2%) than in Ada'a (8.0%) Woreda. For PM the prevalence varied from 60.3% (Dire) to 72.4% (Tuludimitu). Fairly similar seroprevalence (63%) was recorded between Woredas. The seroprevalence of MG varied from 66.1% (Dire) to 87.5% (AdamiTulu) and was higher ($P = 0.007$) in ATJK (85.7%) than in Ada'a (70.1%) Woreda. During the wet season IBD seroprevalence was

closely similar between Woredas: 96.6% in ATJK and 95.9 % in Ada'a, and it did not differ significantly between markets.

TABLE 2. Seroprevalence of Newcastle disease (ND), *Pasteurella multocida* (PM), *Mycoplasma gallisepticum* (MG) and infectious bursal disease (IBD) in backyard chickens at markets in Eastern Shewa Zone, Ethiopia, during September 2010 (wet season).

Woreda/market	n	Percent seropositive (exact 95% C.I.)			
		ND	PM	MG	IBD
ATJK Woreda	119	4.2 (1.4;9.5) ^A	63.0 (53.7;71.7) ^A	85.7 (78;91.4) ^B	96.6 (91.6;99) ^A
AdamiTulu	56	5.4 (1.1;15.8) ^a	62.5 (48.5;75.0) ^a	87.5 (76;94.8) ^a	94.6 (85.1;98.8) ^a
Bulbula	63	3.2 (0.4;11.0) ^a	63.4 (50.4;75.3) ^a	84.1 (72.7;92.1) ^a	98.4 (91.5;100) ^a
Ada'a Woreda	97	8.0 (3.6;15.6) ^A	63.9 (53.5;73.4) ^A	70.0 (59.9;78.9) ^A	96.0 (89.7;98.8) ^A
Dire	68	7.4 (2.4;16.3) ^a	60.3 (47.6;72.0) ^a	66.1 (53.6;77.2) ^a	94.1 (85.6;98.4) ^a
Tuludimitu	29	10.3 (2.2;27.4) ^a	72.4 (52.7;87.3) ^a	79.3 (60.3;92) ^b	100 (88.1;100) ^a
Total	216	6.0 (3.2;10.0)	63.4 (56;67.0)	78.7 (72.6;84.0)	96.3 (93;98.4)

^{A,B} Woredas with different superscripts differ significantly ($P < 0.05$)

^{a,b} Within Woreda, markets with different superscripts differ significantly ($P < 0.05$)

During both wet and dry seasons a fairly similar proportion (6%) of chickens had antibodies against NDV. But during both seasons the proportion of chickens which were seropositive for PM, MG and IBD could be considered high, reflecting the widespread prevalence of those diseases. Both overall and within Woredas the seroprevalence of MG was significantly higher during the wet season than during the dry season ($P < 0.001$).

The distribution of the number of diseases under investigation to which an individual bird was concurrently seropositive is shown in Figure 1. It was found that 81.2% and 91.2% of the chickens tested had antibodies to at least two of the diseases under investigation during the dry and wet seasons, respectively. The median number of diseases to which the birds were seropositive during the dry and wet seasons were 2 and 3, respectively, which differed significantly ($P = 0.002$). Overall, less than 1% of the chickens were seronegative for all four

diseases, while 2.67% had antibodies to all four diseases investigated. Most of the concurrent seropositivity (>75%) was due to IBD with either MG or PM, or both. No bird was seropositive for ND alone.




Figure 1. Seasonal distribution of the number of diseases to which an individual bird was seropositive, in backyard chickens at markets in Eastern Shewa Zone, Ethiopia, tested for antibodies to Newcastle disease, *Pasteurella multocida*, *Mycoplasma gallisepticum* and infectious bursal disease

DISCUSSION

In unvaccinated flocks, positive serological results are clear evidence that the birds have been exposed to the infectious agent under investigation, although without identifying the infecting strains. In the present study we confirmed from the sellers during purchase that none of them had vaccinated their chickens for any poultry diseases. Hence, the presence of

antibodies to ND, PM, MG and IBD was considered evidence of exposure to natural infection.

The study revealed that the prevalence of ND antibodies in backyard chickens was generally low, around 6%. This is considerably lower than previous reports by Zeleke et al. (2005b) and Tadesse et al. (2005), who reported prevalences of 19.8% in the southern and Rift Valley districts and 32.2% in central Ethiopia, respectively, but closer to that reported by Regasa et al. (2007) in southern Ethiopia (11%). Our results are also consistent with seroprevalences in backyard poultry of 4.8% in Mauritania (Bell et al., 1990), 2.2% in Mexico (Gutierrez-Ruiz et al., 2000), 4.8% in California (McBride et al., 1991) and 5% in South Africa (Thekiso et al., 2003). When chickens are affected by a velogenic ND virus that results in very high mortality, one is likely to find few or no survivors with antibodies. Up to 30% of market sellers claimed to have observed poultry disease signs (sudden death, diarrhea and nervous signs) resembling Newcastle disease during previous months. Moreover, the fact that high ELISA PI values (as high as 94) were recorded for most of the positive individual animal sera, in the absence of vaccination, suggests that velogenic virus outbreaks might have killed most chickens in the villages and left few survivors with high antibody titer (Chrysostome et al., 1995; Alexander et al., 2004). According to Awan et al. (1994), low ND HI antibody prevalence is suggestive of an interepidemic phase and this could partly explain the high proportion of seronegative chickens in the present study. The seroprevalence was far lower than that reported from Ecuador (97%) (Sonia et al., 2006), Tanzania (46.1%) (Yongol et al., 2001), Zambia (36.9%) (Alders et al., 1994), Zimbabwe (27%) (Kelly et al., 1994) and Bangladesh (88%) (Biswas et al., 2009). This could be explained by differences in study settings, or by exposure to mild virus strains that induced immunity but did not kill many chickens. The presence of lentogenic, or possibly mesogenic ND in backyard chickens in an area may result in a constant cycle of infection which periodically boosts the immunity of all exposed chickens resulting in a higher proportion of

chickens with antibodies (Sagild and Haresnape, 1987; Martin, 1992). Another reason for variation between studies could be subjectivity and variation in HI cut-off values used for the interpretation of the result. For instance some authors considered a HI titer $\geq 1\log_2$ as positive (Alders et al., 1994; Chrysostome et al., 1995, Bouzaria and Mausavi, 2006; Biswas et al., 2009), while others used cut-off titers of $3\log_2$ (Zelege et al. 2005b; Tadesse et al. 2005) or of $4\log_2$ (Gutierrez-Ruiz et al., 2000). Given the periodic outbreaks and probable high mortality amongst birds affected with ND, our serological findings are likely to be a reasonable indication of the level of ND virus antibody in individual backyard chickens. There was no observed seasonal or geographic variation in seroprevalence, in the present study, suggesting that the disease is widespread and occurs throughout the year in the study area.

Our study revealed a high seroprevalence of fowl cholera (65%), and this constitutes the first report of fowl cholera seroprevalence in Ethiopia. Our finding is in close agreement with that of Kelly et al. (1994), who documented a prevalence of 52% in backyard chickens in Zimbabwe. The high prevalence may be due to infection of backyard chickens by less virulent strains, with or without any clinical signs or significant mortality. This has also been described after challenge with a low virulence strain causing signs of chronic fowl cholera in Kenya (Mbuthia et al., 2008). Biswas et al. (2005) reported proportional mortality from PM of only 6.7% from a longitudinal study in free-range scavenging chickens' in Bangladesh. Mortality is not a typical outcome of fowl cholera in backyard chickens but it may decrease feed efficacy (Mbuthia et al., 2008). Hence there is a high probability that infected birds will seroconvert and remain convalescent carriers, explaining the observed high prevalence (Muhairwa et al., 2000). Surviving birds from diseased flocks may therefore be a risk to naïve birds. Investigations indicate that carriers of PM may exist within poultry flocks with or without a history of previous outbreaks of fowl cholera (Christensen and Bisgaard, 2000; Mbuthia et al., 2008). Mbuthia et al., (2008) were able to isolate PM from 6.2% of healthy-looking chickens from free-range family poultry farms and at market slaughter slabs in

Kenya. Sharing drinking water and feed, which is the common practice in backyard poultry systems in Ethiopia, can facilitate transmission of the bacteria between birds.

The MG prevalence observed in this study averaged 67.7%. This high prevalence is in close agreement with reports from Benin (62%) (Chrysostome et al., 1992), Botswana (57.8%) (Mushi et al., 1999), South Africa (63%) (Thekisoe et al., 2003), Ecuador (73%) (Sonia et al., 2006) and Bangladesh (58.9%) (Sarkar et al., 2005). Lower seroprevalence was reported in village chickens in Malaysia (18.6-25.7%) (Faisal et al., 2001, Shah-Majid, 1996) and Zimbabwe (<33%) (Kelly et al., 1994). Seasonal variation of MG seroprevalence was observed in our study, with higher seroprevalence observed during the wet season (78.7%) than during the dry season (57.7%). This could be due to cold and wet conditions that stress the birds and make them more susceptible to respiratory infections. Sarkar et al. (2005) also found that MG had a seasonal pattern in Bangladesh, where the seroprevalence was higher during winter than summer. Seroprevalence of MG was also higher in ATJK (74%) than in Ada'a (60%) Woreda. This could be associated with the more dusty nature of the areas in ATJK than in Ada'a, as well as to differences in hygienic conditions. Lack of cleaning and hygienic practices in chicken houses may result in ammonia build-up in the wet season that predisposes to respiratory system infection such as mycoplasmosis (Johnson, 1983). However, this was not assessed in the present study. The occurrence of a MG outbreak in commercial broiler chickens in Ethiopia was associated with overcrowding, poor housing, poor sanitation and changes in environmental factors (Chanie et al., 2009). Mycoplasmas are also well known for their interactions with other infectious agents and environmental factors in producing clinical disease (Kleven, 1998). Hence, it appears that mycoplasmas, along with other bacterial or viral pathogens, maybe responsible for a considerable proportion of the respiratory signs that were reported in poultry.

The survey also indicated that IBD is widespread among village chickens in the study area, with a seroprevalence of 94%. This agrees closely with reports by Degefu et al. (2010) from Ethiopia, Sonia et al. (2006) from Ecuador, and Idi et al. (1999) and Karunakaran et al., (1993) from India, who reported seroprevalences of 76.6%, 100%, 74% and 73.7% respectively. However, lower IBD seroprevalences were recorded in Mauritania (15.8%) (Bell et al., 1990), Zimbabwe (55%) (Kelly et al., 1994), Kenya (49.3%) (Ndanyi, 2005) and Botswana (66.2%) (Mushi et al., 2006). The higher seroprevalence of the disease in the study area, in the apparent absence of mortality, could be due to an IBD virus of lower pathogenicity, unlike the case reported from the Amhara region of Ethiopia (Mazengia et al., 2009) or the outbreak in a commercial broiler farm with evident mortalities (Zelege et al., 2005a). It is also possible that the birds were infected with IBD virus as adults, at which stage they simply seroconvert without any apparent clinical disease. With such a high seroprevalence and low mortality of infected birds, there is the possibility of genetic resistance amongst indigenous breeds of chickens in Ethiopia, as reported from Egypt (Hassan, et al., 2004). This is difficult to demonstrate using serological studies, but further studies could be undertaken to investigate this.

In general, there was very little variation between seasons in the seroprevalence of PM, IBD and ND in this study, suggesting the endemicity of those diseases throughout the year in backyard chickens. But variation in the seroprevalence of PM, MG, and IBD was observed between areas during the dry season of the year, suggesting that there could be variation in disease incidence that could be explained partly by variation in local conditions and variation in age structure of flocks in different areas. This could be further substantiated with longitudinal studies that should also take into account the various potential confounders.

In this study we observed a high percentage of concurrent seropositivity to multiple infectious agents. Although we could not determine whether there had been concurrent

infection, most of these involved the immunosuppressive IBD. Although the exact role of each disease is not very clear, the fact that the vast majority of chickens tested had antibodies to two or more of the diseases under investigation, suggests that their health was impacted by multiple infectious pathogens. A similar observation was made in a commercial production system in Ethiopia (Chanie et al., 2009). Biswas et al. (2009) in a serological investigation had reported the existence of various viral diseases in smallholdings in Bangladesh. Bell et al. (1990), in an investigation of the disease status of village chickens in Mauritania, confirmed serologically that village chickens were exposed to NDV, IBD and *Salmonella pullorum* to varying degrees. Similarly, a study from Zimbabwe demonstrated that backyard chicken flocks had been exposed to several viral and bacterial diseases (Kelly et al., 1994). Biswas et al. (2005) reported the existence of a combination of different viral and bacterial diseases in small holdings in Bangladesh and concluded that death was the synergic cooperation between different pathogens. Hence it is possible that the mortalities amongst backyard chickens in Ethiopia could be explained not only by periodic occurrences of velogenic NDV, but also by the synergic effect of other disease-causing agents. Synergism has been demonstrated between MG, ND and infectious bronchitis (Bradbury, 1984). Carpenter et al. (1991), in a study of turkey flocks, indicated that flocks that had antibodies to ND virus and/or *Mycoplasma meleagridis* had an increased risk of having an outbreak of fowl cholera. It is generally believed that concurrent infection renders backyard chickens more susceptible to ND infection (Martin, 1992). The fact that mortality was high even in ND vaccinated flocks indicates that mortalities are the synergic effect of concurrent infections (Kyvsgaard et al., 1999). Kleven (1998) concluded that control of the clinical manifestations of mycoplasma infections is simplified when concurrent infections are minimized and optimum environmental conditions are provided. However, a serological study such as ours cannot provide an accurate indication of concurrent infection of ND with other diseases, firstly since most chickens might have died due to virulent strains of ND, and secondly because serology

cannot indicate precisely when the infectious agent was present. Nevertheless, the high prevalence of concurrent seropositivity, and the fact that about 2.5% of the chickens had antibodies to all the four diseases suggests that backyard chickens are under constant pressure due to several infectious diseases.

In conclusion, our study revealed that several infectious poultry diseases are widespread in backyard chickens in Ethiopia. These diseases are considered to have a significant economic implication individually and/or concurrently. It is also likely that other infections occur in addition to the ones investigated (Tadesse et al., 2004). As serology is not able to demonstrate which strains are circulating, further work is recommended to better understand the circulating strains or pathotypes and the epidemiology of these diseases. Secondly, improvement of village chicken production is at least partly dependent on successful control of some or all of those diseases. Further study is necessary to understand the interactions of these infectious poultry diseases and to estimate their impact on the backyard poultry production system.

ACKNOWLEDGMENTS

The study was done under the framework of the FSP project [GRIPAVI 2006-26] funded by the French Ministry of Foreign and European Affairs (MAEE). Technical assistance of Mr Mengistu Namera, Mrs Roman Zewdu and Mrs Birhan Ayalewu from NAHDIC is highly appreciated.

REFERENCES

Alamargot, J. 1987. Avian pathology of industrial poultry farms in Ethiopia. In: Proceedings Institute of Agricultural. *Proceedings of the First National Livestock Improvements Conference, February 11–13, 1987*, Addis Ababa, Ethiopia. Ethiopian Society of Animal Production. Pp. 114-117.

- Alders, R. G, S. Inoue, and J. C. Katongo. 1994. Prevalence of NCD and evaluation of Hitchner B1 and B4 vaccine for the control of Newcastle disease in village chicken in Zambia. *Prev. Vet. Med.* 21: 125-132.
- Alexander, D. J., J. G. Bell, and R. G. Alders. 2004. A technology review, Newcastle Disease. FAO Animal production and health book No.161, FAO, and Rome.
- Almeida, R. S., O. M Maminaiaina, P. Gil, S. Hammoumi, S. Molia, V. Chevalier, M. Okob, H. Harentsoaniain, R. Andriamanivo, A. Traoré, K. Samaké, A. Diarra, C. Grillet, D., Martinez, and E. Albina. 2009. Africa, a reservoir of new virulent strains of Newcastle disease virus? *Vaccine.* 27: 3127–3129.
- Awan, M. A., M.J. Otte, and A. D. James. 1994. The epidemiology of Newcastle disease in rural poultry: a review. *Avian Pathol.* 23: 405-423.
- Azzam, A. H., H. M. Z. Youseif, and E. E. K. Ahmed. 2004. Prevalence of gumboro disease in vaccinated and non vaccinated village chickens in Egypt. Improving farmyard poultry production in Africa. Intervention and their economic assessment. Proceeding of final research coordination meeting organized by FAO/IAEA. Vienna, 24-28.May 2004. PP. 67-74.
- Bell, J. G., M. Kane, and C. Le Jan. 1990. An investigation of the disease status of village poultry in Mauritania. *Prev. Vet. Med.* 8: 291-294.
- Bell, J. G. 2009. Factors limiting production efficiency and profitability from smallholder poultry production. *World's Poult. Sci. J.* 65:207-210 doi: 10.1017/S0043933909000142.
- Biswas, D., S. Ahmed, A. Rahman, and N. C. Debnath. 2005. A longitudinal study of the incidence of major endemic and epidemic diseases affecting semi-scavenging chickens reared under the Participatory Livestock Development Project areas in Bangladesh. *Avian Pathol.* 34: 303-312.

- Biswas, P. K., H. Barua, G. M. N. Uddin, D. Biswas, A. Ahad, and N. C. Debnath. 2009. Serosurvey of five viruses in chickens on smallholdings in Bangladesh. *Prev. Vet. Med.* 88: 67–71.
- Bouzari, M., and R. MousaviMorekani. 2006. Seroepidemiology of Newcastle disease in domestic village chickens of plain areas of Isfahan province, central India. *Ind. J. Vet. Res.* 7: 80-84.
- Bradbury, J. M. 2001. Avian Mycoplasmosis, In: Frank Jordan et al. (eds.) *Poultry Diseases*. 5th ed. W.B. Saunders Company, Iowa. pp: 178-193.
- Bradbury, J. M. 1984. Avian Mycoplasma infections: prototype of mixed infections with Mycoplasmas, bacteria and viruses. *Annal. micro.* 135: 183-189.
- Carpenter, T. E., K. P Snipes, D. W. Hird, and D. C. Hirsh. 1991. Serologic associations between fowl cholera and other diseases. *Avian Dis.* 35: 718-722.
- Chanie, M., T. Negash, and B. T. Samuel. 2009. Occurrence of concurrent infectious diseases in broiler chickens is a threat to commercial poultry farms in Central Ethiopia. *Trop. Anim. Health and Prod.* 41:1309–1317.
- Christensen, H. and M. Bisgaard. 2000. Fowl cholera. *Rev. Sci. Tech.* 19: 626-637.
- Chrysostome, C. A., J. G. Bell, F. Demey, and A. Verhulst. 1995. Seroprevalences to three diseases in village chickens in Benin. *Prev. Vet. Med.* 22: 257-261.
- CSA. 2009. Agricultural Sample Survey (2008/9) Volume II, *Report on Livestock and Livestock characteristics*, Statistical Bulletin, 446.
- Degefu, H., M. Balcha, M. Yohannes, and M. Getachew. 2010. Seroprevalence of infectious bursal disease in backyard chickens of Oromia regional state, Ethiopia. *Vet. Res.* 3: 89-93.
- Echeonmu, G. O. N., G. U. Iroegbu, and A. C. Emeruwa. 1993. Recovery of velogenic Newcastle disease virus from dead and healthy free-roaming birds in Nigeria. *Avian Pathol.* 22: 383-387.

- Fa, W. Q. 1993. IBD outbreak of chickens raised in backyard in the Chinese village. In: Proceedings, 10th Poultry Association Congress, Sydney, Australia, pp. 182.
- Faisal, Z., A. Ideris, M. Hair-Bejo, A. R. Omar, and T. Chinggiap. 2011. The prevalence of *Mycoplasma gallisepticum* infection in chicken from peninsular Malaysia. J. Anim. and Vet. Advan.10:1867-1874.
- Gutierrez-Ruiz, E. J., G. T. Ramirez-Cruz, E. I. CamaraGamboa, D. J. Alexander, and R. E. A. Gough. 2000. Serological survey for avian infectious bronchitis virus and Newcastle disease virus antibodies in backyard (free-range) village chickens in Mexico. Trop. Anim. Health and Prod. 32:381-390.
- Hassan, M. K., M. A. Afify, and M. M. Aliy. 2004. Genetic resistance of Egyptian chicken to infectious bursal disease and Newcastle disease. Trop. Anim. Health and Prod. 36: 1-9.
- Idi, A., I. Maikano, and H. Adamou. 1999. Seroprevalence of Newcastle disease and infectious bursal disease in local chickens commercialized in Niamey, Niger. INFPD Newsletter Vol. 9 No. 1, January - March 1999.
- Johnson, D. C. 1983. Role of management and sanitation in controlling mycoplasma outbreaks. Avian Dis. 27: 342-343.
- Karunakaran, K, M. Thanappapilldi, and N. Raghavan. 1993. Seroprevalence of infectious bursal disease (IBD) in parts of Tamil Nadu, India. Comp. Immun. Micro. Inf. Dis.16:241-244.
- Kelly, P. J, D. Chitauro, C. Rohde, J. Rukwava, A. Majok, F. Davelaar, and P. R. Mason. 1994. Diseases and management of backyard chicken flocks in Chitungwiza, Zimbabwe. Avian Dis. 38: 626-629.
- Kleven, S. H. 1998. Mycoplasmas in the etiology of multifactorial respiratory disease. Poult. Sci. 77: 1146– 1149.
- Kyvsgaard, N. C., L. A. Luna, and P. Nansen. 1999. Analysis of a traditional grain- and scavenge-based poultry system in Nicaragua. Poultry as a tool in poverty eradication and

- promotion of gender equality. Proceedings of a Workshop. March 22 – 26, 1999, Tune Landboskole, Denmark.
- <http://www.smallstock.info/reference/KVLDK/tune99/9-Kyvsgaard.htm>.
- Lobago, F., and M. Woldemeskel. 2004. An outbreak of Marek's disease in chickens in central Ethiopia. *Trop. Anim. Health and Prod.* 36:397-406.
- Martin, P. A. J. 1992. The epidemiology of Newcastle disease in village chickens. In: SPRADBROW, P.B. (Ed.). *Newcastle Disease in Village Chickens, Control with Thermostable Oral Vaccines*. Proceedings, International Workshop held in Kuala Lumpur, Malaysia, 6-10 October 1991, Centre for International Agricultural Research (ACIAR), Canberra, Pp. 40-45.
- Mazengia, H., S. T. Bekele, and T. Negash. 2009. Incidence of infectious bursal disease in village chickens in two districts of Amhara Region, Northwest Ethiopia. *Livestock Research for Rural Development. Volume 21, Article #214*. Retrieved June 9, 2011, from <http://www.lrrd.org/lrrd21/12/maze21214.htm>.
- Mbuthia, P. G., L. W. Njagi, P. N. Nyaga, L. C. Bebora, U. Minga, J. Kamundia, and J. E. Olsen. 2008. *Pasteurella multocida* in scavenging family chickens and ducks: carrier status, age susceptibility and transmission between species. *Avian Pathol.* 37:51-57
- McBride, M. D., D. W. Hird, T. E. Carpenter, K. P. Snipes, C. Danaye-Elmi, and W. W. Utterback. 1991. Health survey of backyard poultry and other avian species located within one mile of commercial California meat-turkey flocks. *Avian Dis.* 35:403-407.
- Muhairwa, A. P., J. P. Christensen, and M. P. Bisgaard. 2000. Investigations on the carrier rate of *Pasteurella multocida* in healthy commercial poultry flocks and flocks affected by fowl cholera. *Avian Pathol.* 29:133-142.
- Muhairwa, A. P., M. M. A. Mtambo, J. P. Christensen, and M. P. Bisgaard. 2001. Occurrence of *Pasteurella multocida* and related species in village free ranging chickens and their animal contacts in Tanzania. *Vet. Micro.* 78:139-153.

- Mushi, E. Z., M. G. Binta, R. G. Chabo, M. Mathaio, and R. T. Ndebele. 1999. Detection of *Mycoplasma gallisepticum* and *M. synoviae* antibodies in the sera of indigenous chickens by Rapid Plate agglutination test at Mnopane, Gaborone, Botswana. *Onderst. J. Vet. Res.* 66:333 - 334.
- Mushi, E. Z., M. G. Binta, R. G. Chabo, and K. Itebeng. 2006. Diseases of indigenous chickens in Bokaa village, Kgatleng district, Botswana. *J. South Afr. Vet. Assoc.* 77: 131-133.
- Nasser, M., J. E. Lohr, G. Y. Mebratu, K. H. Zessin, M. P. O. Baumann, and Z. Ademe. 2000. Oral Newcastle disease vaccination trials in Ethiopia. *Avian Pathol.* 29: 27– 34.
- National Veterinary Institute (NVI). 1974. Annual report. Cited in T. Dessie, and Y. A. Jobre. 2004. Review of the importance and control of Newcastle disease in Ethiopia. *Eth. Vet. J.* 8:71-81.
- Ndanyi, M. R. A. 2005. Study to determine causes of mortality and the effect of infectious bursal disease (IBD) vaccination in village chickens in Taita Taveta district of Kenya. (Unpublished M. Sc Thesis, the Royal veterinary and agricultural University). 68 pp.
- OIE. 2009. (International office for Animal Health). Terrestrial manual. Newcastle disease, Chapter 2.3.14. www.oie.int/fileadmin/Home/eng/.../2.03.14_NEWCASTLE_DIS.pdf , accessed on 30/08/10.
- Orajaka, I. J. E., D. F. Adene., B. M. Anene, and H. A. Onuoha. 1999. Sero-prevalence of Newcastle disease in local chicken from southeast savannah Zone of Nigeria. *Rev. d'Elevageet de Med. Vet. des pays Trop.* 52: 185-188.
- Parede, L. 1992. The treatment of Gumboo in native chickens. In: Proceedings, international seminar on livestock services for smallholders. Bogor, pp. 153.
- Regasa, C., D. Fufa, S. Berihanu, and D. Hunduma. 2007. Sero-prevalence of Newcastle disease in backyard chickens in mid rift valley of Oromia, Ethiopia. Work Poster. In the

- Proceedings of the 12th International conference of the association of institutions of tropical veterinary medicine, Montpellier, France, 20-22 August, 2007. Pp121.
- Rimler, R. B., and J. R. Glisson. 1997. Fowl cholera. In: Calnek, B.W., Barnes, H.J., Beard, C.W., McDougald, L.R., Saif, Y.M. (Eds.), Poultry diseases. Mosby-Wolfe, London, pp. 143-159.
- Sagild, I., and J. M. Haresnape. 1987. The status of Newcastle disease and the use of v4 vaccine in Malawi. *Avian Pathol.* 16:165-176.
- Sarkar, S. K., M. B. Rahman, M. Rahman, K. M. R. Amin, M. F. R. Khan, and M. M Rahman. 2005. Seroprevalence of *Mycoplasma gallisepticum* Infection of chickens in model breeder poultry farms of Bangladesh. *Int. J. Poult. Sci.* 4:32-35.
- Shah-Majid, M. 1996. Detection of *Mycoplasma gallisepticum* antibodies in the sera of village chickens by enzyme linked immunosorbent assay. *Trop. Anim. Health and Prod.* 28:181-182.
- Sharma, J. M., I. J Kim, S. Rautenschlein, and H. Y. Yeh. 2000. Infectious bursal disease virus of chickens: pathogenesis and immune suppression. *Dev. and Comp. Immun.* 24:223-235.
- Snoeck, C. J., M.F. Ducatez, A. A. Owoade, O.O. Faleke, B. R. Alkali, M.C. Tahita, Z. Tarnagda, J. B. Ouedraogo, I. Maikano, P. O. Mbah, J. R. Kremer, and C. P. Muller. 2009. Newcastle disease virus in west Africa: new virulent strains identified in non-commercial farms. *Arch. Virol.* 154:47-54.
- Sonia, M. D. Hernandez, P. Villegas, F. Prieto, J. C. Unda, N. Stedman, B. Ritchie, R. Carroll, and J. Stephen. 2006. A survey of selected avian pathogens of backyard poultry in northwestern Ecuador. *J. Avian Med. and Surg.* 20:147-158
- Sonaiya, E. B., and S. E. J. Swan. 2004. Small-scale poultry technical guide, FAO animal production and health manual, (FAO, Rome, Italy).

- Tadelle, D. 1996. Studies on village poultry production systems in the central highlands of Ethiopia. M.Sc Thesis, Swedish university of agricultural sciences, pp 70.
- Tadelle, D., T. Million, Y. Alemu, and K.J. Peters. 2003. Village chicken production systems in Ethiopia: 1. Flock characteristics and performance. *Livestock Research for Rural Development* 15(1). <http://www.lrrd.org/lrrd15/1/tadea151.htm>, accessed 19 February 2010.
- Tadesse, S., A. Ashenafi, and Z. Aschalew. 2005. Seroprevalence study of Newcastle Disease in local chickens in central Ethiopia. *Int. J. of Appl. Res. Vet. Med.* 3: 25-29.
- Tadesse, S., M. Woldemeskel, B. Molla, M. Tibbo, D. Kidane, G. Medhin, and S. Britton. 2004. Avian Mycobacteriosis in domestic chickens from selected agro-climatic regions in Ethiopia. *The Int. J. Appl. Res. Vet. Med.*: 2(1).
- Thekiso, M. M. O., P. A. Mbat, and S. P. R. Bisschop. 2003. Diseases of free-ranging chickens in the Qwa-Qwa District of the north eastern Free State province of South Africa. *J. South Afr. Vet. Assoc.* 74:14–16.
- Thitisak, W., O. Janvirisayopak, R. S. Morris, R. von Krüder, and S. Srihakim. 1989. Profile study on healthy and productivity of native chickens in Northeast Thailand. In: Riest, U. (Ed.), *Proceedings on small holder poultry production in developed countries, 1989*, Vol. 4. pp. 162-170.
- van den Berg, T.P., D. Morales, N. Etteradossi, G. Rivallan, D. Toquin, R. Raue, K. Zierenberg, M.F., Zhang, Y.P. Zhu, C.Q. Wang, H.J. Zheng, X. Wang, G.C. Chen, B.L. Lim, and H. Müller. 2004. Assessment of genetic, antigenic and pathotypic criteria for the characterization of IBDV strains. *Avian Pathol.* 33, 470–476.
- Vui, T. Q, J. E. Lohr, M. N. Kyule, K. H. Zessin, and M. P. O. Baumann. 2002. Antibody Levels against Newcastle Disease Virus, Infectious Bursal Disease Virus and Avian Influenza Virus in Rural Chickens in Viet Nam. *Int. J. Poult. Sci.* 1:127-132.

- Woldemariam, S., and A. Wossene. 2007. Infectious bursal disease (Gumboro Disease): Case report at Andasa poultry farm, Amhara region. *Eth. Vet. J.* 11:141-150.
- Wunderwald, C., and R. K. Hoop. 2002. Serological monitoring of 40 Swiss fancy breed poultry flocks. *Avian Pathol.* 3:157-162.
- Xavier J, D. Pascal, E. Crespo, H. L. Schell, J. A. Trinidad, and D. J. Bueno. 2011. Seroprevalence of Salmonella and Mycoplasma infection in backyard chickens in the state of Entre Rios in Argentina. *Poult. Sci.* 90:746-51.
- Yongolo, M. G. S., A. D. Maeda-machang'u, and U. M. Minga. 2001. Newcastle disease and infectious bursal disease among free range village chickens in Tanzania. In: *Proceedings of Associations of the Institutes of Tropical Veterinary Medicine (AIMVT) conference proceedings, 20-23 August 2001, Copenhagen, Denmark.*
- Zelege, A., E. Gelaye, S. Sori, G. Ayelet, A. Sirak, and B. Zekarias. 2005a. Investigation on infectious bursal disease outbreak in Debre-Zeit. *Int. J. Poult. Sci.* 7: 504-506.
- Zelege, A., T. Sori, E., Gelaye and G. Ayelet. 2005b. Newcastle disease in village chickens in the southern and rift valley districts in Ethiopia. *Int. J. Poult. Sci.* 7:508-510.